Astrocytic lysosome deficits reduce alpha-synuclein degradation and induce spread of

pathology

Authors: Lindsay M. Roth[1], Olga Morozova[1], Jan Stöhr[1], & Jason Schapansky[1*]

[1]AbbVie, Cambridge Research Center, 200 Sidney Street, Cambridge, MA 02139, USA

*Address correspondence to:

Dr. Jason Schapansky

AbbVie, Cambridge Research Center, 200 Sidney Street, Cambridge, MA 02139, USA.

Email: jason.schapanksy@abbvie.com

Abstract

Parkinson's Disease (PD) is a neurodegenerative disorder caused by the loss of

dopaminergic neurons in the substantia nigra due to Lewy body aggregates, primarily

composed of misfolded alpha-synuclein (αSyn). While PD progression is thought to be driven by

a prion-like spread of αSyn aggregates between neurons, the role of astrocytes remains

unclear. Observations of αSyn pathology in PD patient astrocytes suggest their potential

involvement in processing aggregates. To investigate this, we studied astrocytes' interactions

with αSyn pre-formed fibrils (PFFs) and their effects in astrocyte-neuron co-cultures on the

spread of seed-competent αSyn. Primary astrocytes quickly internalized and degraded αSyn

PFFs. However, degradation was significantly hindered by lysosome-compromising agents like

chloroquine, Leupeptin, or CA-074. Adding astrocytes to neuron cultures reduced endogenous

αSyn aggregation, indicating their role in mitigating αSyn pathology. When lysosome efficiency

in astrocytes was compromised, their anti-seeding effect diminished. Moreover, lysosome-

compromised astrocytes preloaded with αSyn PFFs enhanced αSyn pathology in neurons,

whereas unimpaired astrocytes did not. These findings suggest astrocytes can modulate and

contribute to αSyn pathology spread, playing a significant role in PD pathogenesis.

Keywords: alpha-synuclein/astrocytes/co-culture/lysosomes/Parkinson's Disease

Introduction

Synucleinopathies, such as Parkinson's Disease (PD), Lewy body dementia (LBD), and

multiple system atrophy, are a spectrum of neurodegenerative diseases defined by the shared

pathological hallmark of intracellular accumulation and aggregation of alpha-synuclein (αSyn) in

neurons and oligodendrocytes (Calabresi *et al*, 2023; Koga *et al*, 2021). In post-mortem PD and

LBD brains, intraneuronal inclusions, termed Lewy bodies (LB) or Lewy neurites (LN), are rich in

misfolded αSyn phosphorylated at serine 129 (pS129), as well as, lipid, lysosomal, and

mitochondrial components (Ramalingam *et al*, 2023; Anderson *et al*, 2006; Fujiwara *et al*, 2002).

It is believed that the misfolding of αSyn is the central pathogenic process in initiation and

progression of PD. Once misfolded, αSyn aggregates are thought to propagate among

connected brain regions in a prion-like manner, templating endogenous αSyn and leading to

disease progression (Braak *et al*, 2003; Angot *et al*, 2010; Dehay *et al*, 2015; Jucker & Walker,

2018). PD etiology is still unclear but includes a complex involvement of genetic, environmental,

and age-related factors (Pang *et al*, 2019) that potentially converge on αSyn pathogenesis.

Furthermore, gene mutations in αSyn are often associated with both sporadic and familial PD

such as the autosomal dominant G209A mutant that leads to overexpression of pathological

A53T αSyn, producing more aggregate prone αSyn and thus increasing disease severity

(Ohgita *et al*, 2022; Polymeropoulos *et al*, 1997). Multiple copies of the αSyn gene results in

early pathogenesis, with disease onset directly correlating to αSyn expression (Stefanis, 2012).

Lastly, PD patients in prodromal and clinical stages can be precisely diagnosed by the presence

of aggregated, seeding competent αSyn in CSF and skin samples (Russo *et al*, 2021; Peng *et*

al, 2023).

Under physiological conditions, αSyn is highly expressed in neurons and associates with

neuronal membranes to regulate the recycling of synaptic vesicles (Bendor *et al*, 2013). In PD,

accumulation of αSyn-rich LB and LN induce progressive loss of dopaminergic (DA) neurons

projecting from the substantia nigra (Poewe *et al*, 2017), leading to gradual motor deficits such

as tremors, muscle stiffness, and postural instability (Sveinbjornsdottir, 2016). PD research has

primarily focused on this neuron-centric mechanism of DA loss; however, emerging research

suggests that non-cell autonomous mechanisms play a significant role in PD etiology and

disease progression (Panicker *et al*, 2021; Brash-Arias *et al*, 2024; Harackiewicz & Grembecka,

2024; Kam *et al*, 2020; Booth *et al*, 2017). A larger effort to understand the role of glia cells in

PD pathogenesis might be a key to the development of disease-modifying therapeutics.

Astrocytes, the most abundant glia cell present in the brain, are critical for maintaining

brain homeostasis and are highly involved in neuroinflammation, an emerging feature of PD

pathogenesis (Araújo *et al*, 2022; Bartels & Leenders, 2007; Mena & Yébenes, 2008;

Verkhratsky *et al*, 2017). Astrocytes have a robust endolysosomal system that is critical for

release of gliotransmitters, modulating energy metabolism of astrocytes and neurons, pruning

synapses, and clearing protein aggregates (Wong, 2020; Balta & Zunke, 2021). Even though

αSyn is predominantly expressed by neurons, aggregated αSyn has been detected in

astrocytes in PD brains, suggesting that αSyn secreted by neurons is taken up by astrocytes

(Altay *et al*, 2022; Sorrentino *et al*, 2019; Braak *et al*, 2007; Wakabayashi *et al*, 2000; Hishikawa

et al, 2001). Further, it suggests that astrocytes in PD are also impaired as they cannot clear

these aggregates sufficiently leading to intracellular accumulation. Studies have shown that

astrocytes can take up αSyn via the endolysosome pathway and localize to lysosomes,

suggesting a role for astrocytes in the clearance of misfolded αSyn aggregates (Loria *et al*,

2017; Cavaliere *et al*, 2017; Filippini *et al*, 2019). Additionally, studies have reported that αSyn

can be transferred from neurons to astrocytes *in vitro* which may help in the clearance of αSyn

aggregates (Loria *et al*, 2017; Tsunemi *et al*, 2020; Domenico *et al*, 2019). In fact, many PD

genome-wide association studies genes that encode proteins involved in lysosomal function,

such as TMEM175, LRRK2, GCase, ATP13A2, Cathepsin D (Cat D), and Cathepsin B (Cat B),

are highly enriched in astrocytes and their risk variants may alter critical astrocytic lysosomal

functions in addition to neuronal impairments (Wang *et al*, 2021; Kim *et al*, 2023; Nalls *et al*,

2019; Booth *et al*, 2017). Recently, studies using induced pluripotent stem cells (iPSC)-derived

astrocytes from PD patients harboring mutations in LRRK2, GCase, or ATP13A2 have shown

alterations in clearance and metabolism pathways and inflammatory responses, suggesting a

role for astrocytic lysosome dysfunction contributing to PD progression (Booth *et al*, 2019, 2017;

Qiao *et al*, 2016; Sanyal *et al*, 2020; Tsunemi *et al*, 2020; Aflaki *et al*, 2020; Domenico *et al*,

2019; Jacquet *et al*, 2021; Pons-Espinal *et al*, 2024).

Here, we aimed to examine the consequences of lysosome deficiency in astrocytes on

both the metabolism of αSyn pre-formed fibrils (PFFs) and the ramifications on neuronal αSyn

pathology. Primary wild-type (WT) astrocytes, with little to no expression of endogenous αSyn,

were used to track uptake and degradation of our PFFs templated from isolated PD brain

samples. Primary neurons overexpressing A53T αSyn were utilized to identify endogenous

pS129 αSyn aggregation induced by PFFs. These data demonstrate that αSyn PFFs are readily

taken up and preferentially degraded by astrocytes, which prevents pS129 αSyn pathology in

surrounding neurons. Disrupting lysosome function with a general insult, such as chloroquine

(CQ), or with a more targeted approach, such as the irreversible Cat B inhibitor CA-074, results

in accumulation αSyn PFFs in astrocytes. Subsequently, the ability of astrocytes to mitigate

αSyn pathology in neurons is reduced and astrocytes can spread αSyn PFFs capable of

seeding endogenous αSyn in neurons. This work demonstrates that lysosome-compromised

(LC) astrocytes can contribute to the spread of αSyn pathology, and that enhancement of

lysosomal function in astrocytes could help assuage progression of PD or other

synucleinopathies.

Materials and Methods

116   *Preparation of primary mouse CD-1 astroglia cell cultures.* All experiments were performed in

117   accordance with the guidelines set forth by AbbVie Animal Care and Use Committees.

118   *A*strocytes were isolated from brains of postnatal day 1 CD-1 mice (Charles River Laboratories,

119   Wilmington, MA) and plated in T75 flasks as previously described (Schildge *et al*, 2013). We

purified the astrocytes using the "shake-off" method (McCarthy & Vellis, 1980). Three independent assays calculating the percentage of Vimentin+, Glial fibrillary acidic protein (GFAP)+, Aldehyde dehydrogenase 1 family member L1 (ALDH1L1)+, or Aquaporin-4 (AQP4)+ cells in culture averaged 93%, 80%, 73.5%, and 89%, respectively, suggesting most of the cells are astrocytes (Jurga *et al*, 2021). After purification, astrocytes were plated into 6 well plates or 96 well plates for immunoblotting or immunocytochemistry, respectively.

Preparation of A53T αSyn transgenic mouse primary cortical neuron cultures. All experiments were performed in accordance with the guidelines set forth by AbbVie Animal Care and Use Committees. Mixed neuron and glia cultures were isolated from brains of embryonic day 16 A53T αSyn transgenic mice bred in-house. Brains were isolated and dissected cortices were incubated in 3 mL Hank's Balanced Salt Solution with Neuronal Isolation Enzyme plus papain (Thermo Fisher Scientific, Waltham, MA) for 15 minutes in a shaking water bath at 37°C. Cells were then diluted to 10 mL total volume in plating media (Dulbecco's Modified Eagle Medium, 10% fetal bovine serum and 1% penicillin streptomycin (P/S)) and centrifuged at 1500 rpm for 5 minutes. Aspiration, resuspension, and centrifugation of the cells occurred twice more and then resuspended cells were strained with a 70 um filter and counted. For mixed cultures (neurons and astrocytes), cells were plated at 35,000 cells/well into 96-well plates in plating media for 4 hours. Media was then changed to culture media (Neurobasal plus, B27 plus supplement, 1% P/S, and 1% GlutaMAX supplement). The time of plating is considered day in vitro (DIV) 0.

Glia depletion protocol for A53T αSyn transgenic mouse primary neuron cultures. Following the preparation of a neuron and astrocyte cell suspension as described above, cells were diluted to 20 mL culture media and counted. Mitotic inhibitor, 5-Fluoro-2'deoxyuridine (FUdR, 1 μM) was added and the FUdR-containing cell suspension was incubated in a metal bead bath at 37°C for

1 hour. Following incubation, cells were recounted and centrifuged at 300 x g for 5 minutes.

Cells were resuspended in culture media and plated at 55,000 cells/well into 96-well plates.

Preparation of primary human astrocytes. Fetal human cortical astrocytes (ScienCell Research

Laboratories, Carlsbad, CA) were cultured *in vitro* in astrocyte media (Cat # 1801, ScienCell

Research Laboratories, Carlsbad, CA) in T-75 flasks until they reached 90% confluency. Cells

were then incubated with 0.25% trypsin at 37°C for 3 minutes to lift cells. Following

ultracentrifugation at 300 x g for 5 minutes, cells were counted and plated at 10,000 cells/well

into 96-well plates or 500,000 cells/well into 6-well plates for immunocytochemistry and

immunblotting experiments, respectively.

Preparation of αSyn recombinant PFF aggregates and drug treatments. Recombinant WT α-syn

was expressed in *E. coli* strain BL21(DE3) and purified as previously described (Ghee *et al*,

2005). Disease-relevant PFFs were generated by incubating insoluble αSyn seeds purified from

human PD brains for 7 days, as previously described (Marotta *et al*, 2021). Protein

concentration was determined by UV-Vis Nanodrop measurement after incubation with 4M

Guanidine HCL. PFFs were diluted to 0.1 mg/mL in Dulbecco's PBS (DPBS) and sonicated for 1

min at 30% amplitude with 1 sec on, 1 sec off pulses using a QSonica 500 water bath sonicator.

pHrodo-labeled PFFs were generated by following the manufacturers protocol (Cat #: P36011,

ThermoFisher) with a labeling ratio of 3:1. The unreacted dye was removed from solution by

centrifuging the fibril solution at 20,000xg for 20 min, removing the supernatant and

resuspending the pellet in PBS. The following drugs were used in this study: chloroquine (CQ,

25 μM, Millipore Sigma Cat #: C6628), leupeptin (Leu, 10 μM, Selleckchem Cat #: S7380), CA-

074 (10 μM, Tocris Biosciences Cat #: 4863), or pepstatin A (Pep A, 10 μM, Tocris Biosciences

Cat #: 1190). Cells were treated with αSyn PFFs for the time and concentration specified for

each experiment.

Immunofluorescence. All cell culture experiments for immunofluorescence were performed in

96-well plates. All experiments involving primary neurons were fixed and immunostained on DIV

21. Cultures were fixed with ice cold methanol for 10 minutes, rinsed with PBS, and incubated

with 5% goat serum, 0.2% *Triton-x100* in PBS to permeabilize and block cells for 1 hour. Then

cells were incubated at 4°C degree overnight with primary antibodies. Antibodies used per

experiment are specified in figure legends. A list of all primary antibodies used are as follows:

microtubule-associated protein 2 (MAP2, Abcam, Cat #: ab5392, 1:3000); GFAP (Thermo

Fisher Scientific, Cat #: 14-9892-82, 1:3000); pS129 (Cell Signaling Technologies, Cat #:

23706, 1:2000); Lysosomal associated membrane protein 1 (LAMP1) for mouse astrocytes

(Biolegend, Cat #: 121602 1:200); LAMP1 for human astrocytes (Cell Signaling Technologies,

Cat #: 15665, 1:200); total α-synuclein (MJFR1, Abcam, Cat #: ab138501, 1:2000) and

Galectin-3 (Abcam, Cat #: A3A12, 1:200). The next day, cells were rinsed twice with PBS and

incubated with secondary antibodies (Alexa Fluor 488, 568 or 647 [ThermoFisher] at 1:500

dilution) for 1 hour at room temperature. Following a 2x PBS rinse, cells were incubated with

1:2000 dilution of Hoechst in PBS for 10 minutes to stain nuclei. Cells are left in PBS for

imaging to avoid drying out. Cells were imaged using an Opera Phenix high content confocal

microscope (*Revvity*, Shelton, CT) affixed with 4 excitation lasers (405, 488, 561 and 640 nm)

and emission filters (435-550, 435-480, 500-550, 570-630 and 650-760 nm). Images were

captured at 20x or 40x magnification with at least 9 fields acquired per well. Quantification of

marker areas or cells positive for stained markers was performed with algorithms made on the

*H*armony analysis software (*Revvity*).

Immunoblotting. Primary mouse astrocytes were grown in 6-well plate tissue culture dishes for

immunoblot until they reached approximately 80% confluency. Whole cell extracts of primary

mouse astrocyte cultures were prepared with ice cold RIPA buffer (ThermoFisher Scientific,

Waltham, MA), containing protease and phosphatase inhibitor cocktails (Roche Diagnostics)

followed by probe sonication. Protein concentrations were determined by bicinchoninic acid

(BCA) assay (Thermo Fisher Scientific, Waltham, MA). Protein (15 ug/lane) was loaded onto a

12% Bis-Tris gel for separation. After separation, proteins were transferred to nitrocellulose

membranes using an iBlot system (Thermo Fisher Scientific, Waltham, MA) and blocked in tris-

buffered saline with Tween 20 (TBS-T) with 5% milk for 1 hour at room temperature.

Membranes were incubated overnight at 4°C with primary antibodies in TBS-T with 5% milk.

Primary antibodies to the following antigens were used: total αSyn (Syn1/clone 42, 1:500

dilution, BD Biosciences Cat #: 610787), total αSyn (MJFR1, 1:500 dilution, Abcam Cat #:

ab138501), and β-actin (1:8,000 dilution, Millipore Sigma Cat #: A1978). Following 3x washes

with TBS-T, membranes were incubated with corresponding antigen specific horseradish

peroxidase-conjugated secondary antibodies (1:10,000 dilution) in TBS-T with 5% milk.

Membranes were incubated with chemiluminescent substrate SuperSignal West Dura

(ThermoFisher Scientific, Waltham, MA) for 5 minutes and then blots were visualized using a

ChemiDoc MP Imaging System (Biorad, Hercules, CA). Densitometric analysis of band

intensities were conducted using Fiji (NIH RRID: SCR_002285). All bands were normalized to

the loading control as specified in each experiment.

Statistical analysis. Data were analyzed using GraphPad Prism statistical software (version

10.0) and were graphically presented as a fold change from the average value of the

normalizing group per experiment ± SD. The average value of the normalizing group is

represented as a dotted line. The normalizing group is specified in the figure legends per

experiment. All immunoblots were normalized to a loading control, β-actin. Each experiment

was performed at least 3 times on separate 96-well or 6-well plates.

Results

Primary astrocytes readily take up and preferentially degrade αSyn PFFs via the

endolysosomal pathway. Due to the presence of αSyn pathology in astrocytes in PD patients

(Wakabayashi *et al*, 2000; Altay *et al*, 2022), we hypothesized that astrocytes may play a

significant role in metabolism of αSyn PFFs. Using WT astrocytes allowed for a model in which

we could monitor metabolism of external αSyn aggregates since astrocytes express little to no

endogenous αSyn (Braak *et al*, 2007; Mori *et al*, 2002). To measure rate of uptake, mouse

astrocytes were treated with pHrodo-tagged recombinant αSyn monomer or PFFs and imaged

live by fluorescence every 2 hours for 24 hours. The pHrodo tag allows for fluorescence only in

acidic compartments such as endosomes or lysosomes, therefore only active uptake is

observed. Representative images of αSyn monomer- or PFF-treated astrocytes showed

increased pHrodo+ fluorescence over time (Figure 1A). The number of phRodo+ spots over

time (Figure 1B) or at 24 hours (Figure 1C) was significantly increased in astrocytes treated with

αSyn PFFs compared to monomer, implying that aggregates are more actively internalized

compared to soluble αSyn.

Next, we assessed astrocytic degradation of αSyn monomer or PFFs by treating primary

astrocytes with αSyn monomer or PFFs over time and immunoblotting for total αSyn. Total full

length αSyn protein was initially present in both αSyn monomer- and PFF-treated astrocytes,

but we only observed decreased αSyn abundance over time in αSyn PFF-treated astrocytes,

indicating that aggregated, not monomeric, αSyn is targeted for degradation (Figure 1D, E).

Moreover, only αSyn PFF treatment resulted in the presence of degraded αSyn bands,

represented by bands lower than 14 kDa, that increased after 24 h and then decreased at 48

and 72 h (Figure 1D,F). Using a C-terminal-localized αSyn antibody resulted in a loss of

detection of degraded αSyn bands in the αSyn PFF-treated primary astrocyte lysates (Extended

View Figure 1A,B), suggesting that astrocytes are degrading αSyn PFFs at the c-terminal end.

Given that αSyn PFFs were detected in acidic compartments, we asked whether we

would observe decreased total αSyn abundance over time. Primary astrocytes were treated with

αSyn PFFs over time and immunostained for total αSyn and the lysosome marker LAMP1

(Settembre *et al*, 2013). Subsequent quantification of αSyn PFF-treated astrocytes revealed

fewer total αSyn+ spots over time and fewer total αSyn+ and LAMP1+ spots over time (Figure

1G, H, J), with no observed differences in total LAMP1+ spots or total nuclei over time (Figure

1I, J). These results are consistent with our immunoblotting results, suggesting an active

digestion of αSyn aggregates that occurs quickly following exposure.

To confirm this was not a phenomenon localized to mouse astrocytes, primary human

astrocytes were used to assess the differences in uptake and degradation of αSyn monomers or

PFFs. Like mouse astrocytes, we observed that both monomers and PFFs are taken up, while

only the PFFs were subsequently degraded (Extended View figure 1C-E). Additionally, the

immunofluorescent assay results were consistent with the immunoblotting results, with

decreased total αSyn+ spots and total αSyn+ and LAMP1+ spots in human primary astrocytes

treated with αSyn PFFs (Extended View figure 1F-J). Thus, this was not a species-specific

phenomenon but seems to relate to the human cells as well.

αSyn PFFs accumulate in LC primary astrocytes. In figure 1 and extend view figure 1 we

showed that mouse and human astrocytes take up and degrade αSyn PFFs via the

endolysosomal pathway. We hypothesized that αSyn PFFs could accumulate if this pathway

was perturbed. To examine this, we co-treated primary astrocytes with lysosome compromising

agents and αSyn PFFs, followed by immunostaining for total αSyn after 24 hours. We utilized

the lysosomotropic compound chloroquine (CQ, 25 µM), Cat D enzyme inhibitor pepstatin A

(Pep A, 10 µM) or Cat B inhibitors leupeptin (Leu, 10 µM) and CA-074 (10 µM) to challenge

lysosomal activity (Yoon *et al*, 2022; Gacko *et al*, 2007; Baici & Gyger-Marazzi, 1982; Gallagher

et al, 2017) . Quantification of challenged astrocytes revealed more total αSyn+ spots within

lysosomes after 24 hours treatment compared to vehicle-treated astrocytes (Figure 2A,B).

Interestingly, total αSyn+ spots in Pep A-treated astrocytes were similar to that of vehicle-

treated astrocytes (Figure 2A, B), implying that aspartic acid proteases are not involved in

astrocytic αSyn PFF degradation. Furthermore, astrocytes only demonstrated lysosome

damage marker Galectin-3 (Gal-3) with CQ treatment as shown previously (Gallagher *et al*,

2017) and not with cathepsin inhibition (Extended View figure 2D-H), suggesting that lysosome

damage is not required for aSyn accumulation. Moreover, the amount of αSyn remaining

correlated with the severity of insult, with CQ having a greater effect. Therefore, both lysosomal

damage and/or lysosomal enzymatic insufficiency can result in an accumulation of αSyn

aggregates in astrocytes.

To further elucidate the degradative capacity of lysosome-compromised (LC) astrocytes,

we co-treated astrocytes with increasing concentrations of Leu (0.1, 1 or 10 µM), CA-074 (1 or

10 µM) or Pep A (10 µM) and αSyn PFFs for 24 hours and immunoblotted for total αSyn. Leu-

treated astrocytes showed concentration-dependent accumulation of non-degraded αSyn

compared to vehicle-treated astrocytes (Figure 2D,E). In addition to full length αSyn we also

observed an accumulation of partially degraded αSyn suggesting that inhibition of Cat B

reduces the efficiency of PFF degradation in the lysosome. The distinct banding patterns

indicate a specific protease resistant confirmation of the αSyn fibrils (Figure 2D,F). In contrast

we observed no difference in full length or degraded total αSyn in Pep A-treated astrocytes

compared to vehicle-treated astrocytes suggesting that Cat D is not participating in the

degradation of pathological αSyn (Figure 2D-F). Specific inhibition of cathepsin B with CA-074

also showed accumulation of non-degraded αSyn and distinct degraded αSyn bands compared

to vehicle-treated astrocytes (Figure 2G-I). These results are consistent with the

immunofluorescence results in which only Leu or CA-074-treated astrocytes increased

abundance of total αSyn, suggesting that cysteine proteases and specifically cathepsin B

function is critical for degrading αSyn aggregates in astrocytes.

Since CQ is a known inhibitor of endocytosis (Wang *et al*, 1993; Dutta & Donaldson,

2012), we also utilized a different paradigm to specifically demonstrate the effect of CQ on αSyn

PFF degradation. In figure 2J, astrocytes were first preloaded with αSyn PFFs for 4 hours,

rinsed and treated with vehicle or CQ for 24 or 48 hours. This data shows similar results to

those in figure 2D-I, that both non-degraded αSyn and partially-degraded αSyn accumulated in

CQ-treated astrocytes compared to vehicle-treated astrocytes (Figure K,L).

Healthy astrocytes prevent pS129 pathology in neurons induced by αSyn PFFs. We have

shown that astrocytes take up and preferentially degrade αSyn PFFs through the

endolysosomal pathway, in a distinct manner from αSyn monomer. To model how this astrocytic

ability may impact neuronal αSyn pathology, we developed multiple co-culture models. Figure

3A depicts the first co-culture assay we used to understand the effect of astrocytes on neuronal

αSyn pathology. Briefly, DIV 7 A53T αSyn transgenic mouse neuroglia cultures were exposed to

αSyn PFFs to induce endogenous αSyn aggregation. To ensure some uptake of αSyn PFFs into

neurons, we waited four days after PFF treatment before adding increasing densities (5,000,

10,000, or 20,000) of mouse astrocytes to the neuroglia cultures. On DIV 21, the co-cultures

were fixed and immunostained for astrocytes, neurons and endogenous αSyn pathology using

GFAP, MAP2, or pS129, respectively. Representative images and quantification of astrocyte-

added neuroglia cultures show fewer pS129+ spots compared with neuroglia cultures in the

absence of additional mouse astrocytes (Figure 3B,C). Thus, the addition of exogenous

astrocytes was protective, preventing neuronal-based αSyn aggregation. This assay includes

A53T αSyn transgenic astrocytes, and adding exogenous astrocytes significantly decreases

neuronal αSyn pS129+ pathology compared to control.

Next, we hypothesized that depletion of A53T αSyn transgenic astrocytes from neuroglia

cultures would enhance endogenous αSyn pS129 pathology in primary neurons, subsequently

increasing the efficacy of exogenous astrocytes when added. To test this, we used the mitotic

inhibitor 5-Fluoro-2'deoxyuridine (FUdR) to deplete proliferating glia cells prior to plating

neurons. Endogenous αSyn pS129+ pathology was examined in FUdR-treated pure neuron

cultures compared to untreated cultures induced by αSyn PFFs. Representative images and

quantification show more pS129+ spots in FUdR-treated neuron cultures compared to untreated

cultures (Figure 3G,H). Therefore, depletion of A53T αSyn transgenic astrocytes increased

endogenous αSyn pS129+ pathology in neurons.

Next, we examined whether addition of astrocytes to pure neurons modulated

endogenous αSyn pS129+ pathology to a greater extent than in the neuroglia cultures after

adding additional astrocytes. Following the paradigm depicted in figure 3L, astrocytes were

added four days after pure neurons were treated with αSyn PFFs to allow pure neurons to take

up most of the αSyn PFFs prior to astrocyte addition; Cells were subsequently immunostained

for GFAP, MAP2, or pS129 on DIV 21. Representative images and quantification of cultures

with astrocytes added show fewer pS129+ spots compared to pure neuron control (Figure

3M,N). Astrocyte addition to pure neuron cultures more robustly decreased neuronal αSyn

pS129+ pathology (by ~50%) compared to astrocyte addition to neuronal cultures with A53T

αSyn transgenic astrocytes still present. This data confirmed our hypothesis that astrocytes can

reduce neuronal exposure to propagating αSyn seeds.

LC astrocytes do not mitigate pS129 pathology in neurons induced by αSyn PFFs.

We next hypothesized that LC astrocytes would be unable to prevent endogenous αSyn pS129+ pathology in neurons. We tested this by treating astrocytes with CQ or CA-074 overnight prior to adding to neuroglia cultures already exposed to PFFs (Figure 4A). Representative images and quantification of neuroglia cultures treated with LC astrocytes had more pS129+ spots compared to neuroglia cultures with unimpaired (vehicle-treated) astrocytes added (Figure 4B,C). This suggests that lysosomal function plays a critical role in the ability of astrocytes to reduce endogenous αSyn seeding in neurons.

Coming from cultures with a mixed impaired and unimpaired astrocyte population, we next used a model to assess how solely LC astrocytes would affect endogenous αSyn pS129+ pathology in neurons. In this paradigm, astrocytes were plated first and treated overnight with CQ or CA-074, rinsed and pure neurons were plated on top. Co-cultures were exposed to αSyn PFFs and subsequently immunostained for GFAP, MAP2, or pS129 (Figure 4G). Representative images and quantification of co-cultures with LC astrocytes show increased pS129+ spots compared to co-cultures with vehicle-treated astrocytes (Figure 4H,I). Therefore, astrocytes with insufficient lysosome capacity have a reduced proficiency to mitigate αSyn pathology in neuronal cultures as well as astrocytes with normal lysosome function.

αSyn PFF preloaded LC astrocytes induce endogenous αSyn pS129 pathology in neurons. Since LC astrocytes accumulate αSyn PFFs following exposure to PFFs (Figure 2) and lose the ability to alleviate pS129 pathology in neurons (Figure 4), we examined whether LC astrocytes preloaded with αSyn PFFs could spread seed-competent αSyn to neurons. In this assay, astrocytes are loaded with αSyn PFFs, rinsed, and treated overnight with CQ. αSyn PFF

preloaded astrocytes are added to neuroglia cultures and immunostained on for GFAP, MAP2,

or pS129 (Figure 5A). Quantification reveals the presence of pS129+ spots only in neuroglia

cultures with PFF-preloaded, CQ-treated astrocytes compared to PFF-preloaded, vehicle-

treated astrocytes, highlighting the importance of a functional lysosome to neutralize aSyn

aggregates (Figure 5B,C).

Finally, since the paradigm in figure 5A includes the presence of A53T αSyn transgenic

astrocytes, we decided to use the same assay depicted in figure 4G-L to better assess the

ability of solely LC astrocytes to induce endogenous αSyn pS129 pathology in pure neurons. In

this experiment, we plated astrocytes first, preloaded them with αSyn PFFs, rinsed, and treated

with CQ or CA-074 overnight. Pure neurons were added on top of the astrocytes and the cells

were immunostained for GFAP, MAP2, or pS129 (Figure 5G). Similarly, only neurons cultured

with PFF pre-loaded astrocytes treated with CQ and to a lesser degree with CA-074

demonstrated αSyn pS129 pathology (Figure 5H-I), further suggesting that LC astrocytes can

spread seed-competent αSyn to neurons. Taken together, these data suggest that astrocytic

lysosome efficiency is important for degrading αSyn aggregates and slowing the spread of αSyn

pathology in neurons.

Discussion

Using co-culture models to specifically manipulate astrocytes, we provide evidence that

astrocytic lysosomal deficiency causes the accumulation of αSyn PFFs, which can lead to the

spread of seed-competent αSyn to surrounding neurons for formation of endogenous pS129+

aggregation. We show that healthy WT mouse and human astrocytes can readily take up and

preferentially degrade αSyn PFFs *in vitro* and exogenously added WT astrocytes can mitigate

PFF-induced aggregation of endogenous αSyn in primary neurons. However, either general

astrocytic lysosomal damage and dysfunction or specific minor lysosome enzymatic inhibition

disrupts αSyn PFF degradation, resulting in an elevation of endogenous αSyn aggregation in

neurons and spread of αSyn pathology. This work emphasizes the importance of the astrocytic

lysosomal pathway in the clearance αSyn aggregates and supports a non-cell autonomous

contribution to PD neuropathology.

Firstly, we observed that astrocytes take up more αSyn PFFs over time than αSyn

monomer (Figure 1A-C). Due to the use of the pHrodo tag, we know that both forms of αSyn

were taken up through acidic compartments including the endosome, which agrees with

previous reports of αSyn uptake mechanisms (Lee *et al*, 2008b, 2008a; Andromidas *et al*, 2024;

Koob & Sacchetti, 2018). Interestingly, the observations that monomeric αSyn is taken up less

than aggregates and that it is not degraded in the same manner suggest that the two αSyn

formats are differentially internalized and processed (Figure1D-F). Also, in αSyn PFF-treated

astrocytes, we observed that total αSyn staining localized to LAMP1+ compartments suggesting

that αSyn PFFs are targeted to lysosomes for degradation (Figure 1G-H). Taken together,

astrocytes specifically and preferentially target αSyn PFFs, and not monomeric αSyn, to the

lysosome and degrade them. This specific difference in the astrocytic metabolism of αSyn

aggregates led us to explore the role of lysosome degradation further.

Due to the strong genetic and mechanistic evidence for Cat D and Cat B involvement in

PD, (Lin *et al*, 2024; Nalls *et al*, 2019; Bunk *et al*, 2021; Jones-Tabah *et al*, 2024; McGlinchey &

Lee, 2015; Drobny *et al*, 2022; McGlinchey *et al*, 2019) inhibitors of Cat B (Leu or CA-074) and

an inhibitor of Cat D (Pep A) were utilized to understand cathepsin-specific enzymatic digestion

of αSyn aggregates in astrocytes. The lysosomotropic compound CQ was used to examine how

general lysosome deficiency affects astrocytic αSyn PFF degradation, as it is commonly used to

raise lysosomal pH to limit enzymatic digestion (Zeng *et al*, 2020). In αSyn PFF and CQ-, Leu-

or CA-074-treated astrocytes, we observed unique partially degraded forms of αSyn not

detected in αSyn PFF and vehicle- or Pep A-treated astrocytes (Figure 2). Based on the

antibodies used in this study (Extended View Figure 1A), these degraded forms of αSyn are

likely c-terminally fragmented since they were not seen when αSyn PFF-treated astrocyte

lysates were probed with an αSyn antibody targeting a c-terminal epitope (MJFR1 epitope

aa118-123; Extended View Figure 1B-C). Intriguingly, this accumulation of unique degraded

forms of αSyn was only observed with Cat B inhibition (Leu or CA-074 treatment) and not with

Cat D inhibition (Pep A treatment) (Figure 2). Despite the compelling evidence for the

degradation of αSyn aggregates by Cat D (Bunk *et al*, 2021; Qiao *et al*, 2008; Cullen *et al*,

2009), it's possible that Cat B plays a larger role in the context of astrocytic αSyn degradation.

Moreover, other cathepsins could play a role in astrocytic αSyn degradation and compensate for

Cat B (Matsuki *et al*, 2024). A full analysis of cathepsin activity in astrocytes regarding αSyn

digestion may be warranted.

A significant distinction in our study is that we did not observe overt changes to

lysosome morphology or damage with αSyn PFF treatment alone (Extended View Figure 2).

Many studies of glia cells exposed to various forms of αSyn observed organellar dysfunction or

alterations in the secretome (Lindström *et al*, 2017; Freeman *et al*, 2013; Raj *et al*, 2024;

Chavarría *et al*, 2018). These studies all used different PFFs that were exposed to glia cells at

much higher concentrations than this study, which may account for the differing results.

However, this work adds to a growing number of studies that suggest glia cell-specific

lysosomal dysfunction plays a critical function in the contribution to neurodegenerative disease

pathology and progression (Quick *et al*, 2023; Uddin & Lim, 2022; Balta & Zunke, 2021; Aflaki *et*

al, 2020).

Importantly, only LC astrocytes preloaded with αSyn PFFs induced spread of αSyn

pathology (Figure 5). This observation is aligned with our data in figure 2, clearly showing that

LC astrocytes accumulated αSyn at times where the vehicle-treated astrocytes degraded αSyn

PFFs rapidly and efficiently. It has been suggested that truncated forms of αSyn are more prone

to aggregation (Sorrentino & Giasson, 2020; Liu *et al*, 2005; Zhang *et al*, 2022). Moreover, Cat

B-digested αSyn PFFs form truncated αSyn that has been shown to increase aggregation

propensity in neurons (Tsujimura *et al*, 2015). It is possible that these unique partially degraded

forms of αSyn PFFs, only observed in LC astrocytes, play a significant role in the spread of

αSyn pathology in our model. Furthermore, αSyn PFF preloaded CQ-treated astrocytes induced

more pS129 αSyn pathology in neurons compared to αSyn PFF preloaded CA-074-treated

astrocytes (Figure 5G-L). Since CQ treatment caused overt changes to lysosome size, number,

and lysosome damage (as demonstrated with Gal3 localization) that were absent following CatB

inhibition, it is possible that the mechanism by which these astrocytes induced αSyn pathology

in neurons is different, even though both compounds clearly led to accumulation of non-

degraded and degraded αSyn aggregates (Figure 2). A proposed model for propagation based

on our findings is shown in the synopsis image. However, the potential mechanisms of αSyn

aggregate spread remain to be elucidated.

In PD, αSyn aggregates are hypothesized to spread in a prion-like manner through

synaptically coupled networks of neurons leading to LB and LN pathology and

neurodegeneration (Henrich *et al*, 2020). However, it's suggested that there is no clear

correlation between the strength of the synaptic connections and the magnitude or persistence

of the propagated pathology, which implies that spread of αSyn is not solely through neuronal

synaptic connectivity (Henrich *et al*, 2020). Additionally, there is evidence suggesting that αSyn

is transmitted amongst neurons, astrocytes, and microglia, but the mechanisms by which this

occurs remain intensely debated (Dehay *et al*, 2015; Loria *et al*, 2017; Rostami *et al*, 2017;

Walsh & Selkoe, 2016; Emmanouilidou *et al*, 2010; El-Agnaf *et al*, 2003). Many of these studies

report αSyn is released via exosomes, by lysosome exocytosis following lysosome dysfunction

or through neuroinflammation (Burbidge *et al*, 2022; Alvarez-Erviti *et al*, 2011; Xie *et al*, 2022;

Bae *et al*, 2022; Lee *et al*, 2014). Moreover, two other studies show that neuron to neuron or

astrocyte to astrocyte αSyn propagation can occur through direct contact with tunneling

nanotubes (Abounit *et al*, 2016; Rostami *et al*, 2017). In future studies, we aim to further

investigate these possible mechanisms of astrocytic αSyn PFF spread in our model.

This work supports a growing body of research suggesting a role for glia cells,

specifically astrocytes, in PD etiology and disease progression. We illustrate that overt

lysosome damage or Cat B inhibition in astrocytes can modulate and spread αSyn pathology in

neurons. Astrocytes robustly take up and degrade αSyn aggregates via the endolysosomal

system and that enhancement of this pathway can be harnessed to develop new disease-

modifying therapeutics for PD.

Funding. L.M.R., O.M, J.S, and J.S. are employees of AbbVie. The design, study conduct, and

financial support for this research were provided by AbbVie. AbbVie participated in the

interpretation of data, review, and approval of the publication.

Author contributions. L.M.R., O.M, J.S, and J.S* contributed to the study conception and

design. Material preparation, data collection, and analysis were performed by L.M.R. and O.M.

The first draft of the manuscript was written by L.M.R. and J.S*. All authors commented on

previous versions of the manuscript. All authors read and approved of the final manuscript.

Declaration of Competing Interest. The authors declare no competing financial interests.

Acknowledgements. Jordan Dasilva in the Comparative Medicine East group at AbbVie for her

help with the isolation of primary neuroglia cultures. BioRender was used to create all assay

schematics in the figures and the synopsis image.

References

Abounit S, Bousset L, Loria F, Zhu S, Chaumont F de, Pieri L, Olivo-Marin J, Melki R & Zurzolo
C (2016) Tunneling nanotubes spread fibrillar α-synuclein by intercellular trafficking of
lysosomes. *EMBO J* 35: 2120–2138

Aflaki E, Stubblefield BK, McGlinchey RP, McMahon B, Ory DS & Sidransky E (2020) A
characterization of Gaucher iPS-derived astrocytes: Potential implications for Parkinson's
disease. *Neurobiol Dis* 134: 104647

Altay MF, Liu AKL, Holton JL, Parkkinen L & Lashuel HA (2022) Prominent astrocytic alpha-
synuclein pathology with unique post-translational modification signatures unveiled across
Lewy body disorders. *Acta Neuropathol Commun* 10: 163

Alvarez-Erviti L, Seow Y, Schapira AH, Gardiner C, Sargent IL, Wood MJA & Cooper JM (2011)
Lysosomal dysfunction increases exosome-mediated alpha-synuclein release and
transmission. *Neurobiol Dis* 42: 360–367

Anderson JP, Walker DE, Goldstein JM, Laat R de, Banducci K, Caccavello RJ, Barbour R,
Huang J, Kling K, Lee M, *et al* (2006) Phosphorylation of Ser-129 Is the Dominant
Pathological Modification of α-Synuclein in Familial and Sporadic Lewy Body Disease*. *J Biol
Chem* 281: 29739–29752

Andromidas F, Mackinnon BE, Myers AJ, Shaffer MM, Brahimi A, Atashpanjeh S, Vazquez TL,
Le T, Jellison ER, Staurovsky S, *et al* (2024) Astrocytes initiate autophagic flux and maintain
cell viability after internalizing non-active native extracellular α-synuclein. *Mol Cell Neurosci*
131: 103975

Angot E, Steiner JA, Hansen C, Li J-Y & Brundin P (2010) Are synucleinopathies prion-like
disorders? *Lancet Neurol* 9: 1128–1138

Araújo B, Caridade-Silva R, Soares-Guedes C, Martins-Macedo J, Gomes ED, Monteiro S &
Teixeira FG (2022) Neuroinflammation and Parkinson's Disease—From Neurodegeneration
to Therapeutic Opportunities. *Cells* 11: 2908

Bae E-J, Choi M, Kim JT, Kim D-K, Jung MK, Kim C, Kim T-K, Lee JS, Jung BC, Shin SJ, *et al*
(2022) TNF-α promotes α-synuclein propagation through stimulation of senescence-
associated lysosomal exocytosis. *Exp Mol Med* 54: 788–800

Baici A & Gyger-Marazzi M (1982) The Slow, Tight-Binding Inhibition of Cathepsin B by
Leupeptin. *Eur J Biochem* 129: 33–41

Balta D & Zunke F (2021) The role of lysosomes in alpha-synucleinopathies: a focus on glial
cells. *Neural Regen Res* 17: 1486–1488

Bartels AL & Leenders KL (2007) Neuroinflammation in the pathophysiology of Parkinson's
disease: Evidence from animal models to human in vivo studies with [11C]-PK11195 PET.
Mov Disord 22: 1852–1856

Bendor JT, Logan TP & Edwards RH (2013) The Function of α-Synuclein. *Neuron* 79: 1044–
1066

Booth HDE, Hirst WD & Wade-Martins R (2017) The Role of Astrocyte Dysfunction in
Parkinson's Disease Pathogenesis. *Trends Neurosci* 40: 358–370

Booth HDE, Wessely F, Connor-Robson N, Rinaldi F, Vowles J, Browne C, Evetts SG, Hu MT,
Cowley SA, Webber C, *et al* (2019) RNA sequencing reveals MMP2 and TGFB1
downregulation in LRRK2 G2019S Parkinson's iPSC-derived astrocytes. *Neurobiol Dis* 129:
56–66

Braak H, Sastre M & Tredici KD (2007) Development of α-synuclein immunoreactive astrocytes
in the forebrain parallels stages of intraneuronal pathology in sporadic Parkinson's disease.
Acta Neuropathol 114: 231–241

Braak H, Tredici KD, Rüb U, Vos RAI de, Steur ENHJ & Braak E (2003) Staging of brain
pathology related to sporadic Parkinson's disease. *Neurobiol Aging* 24: 197–211

Brash-Arias D, García LI, Pérez-Estudillo CA, Rojas-Durán F, Aranda-Abreu GE, Herrera-
Covarrubias D & Chi-Castañeda D (2024) The Role of Astrocytes and Alpha-Synuclein in
Parkinson's Disease: A Review. *NeuroSci* 5: 71–86

Bunk J, Huarcaya SP, Drobny A, Dobert JP, Walther L, Rose-John S, Arnold P & Zunke F
(2021) Cathepsin D Variants Associated With Neurodegenerative Diseases Show
Dysregulated Functionality and Modified α-Synuclein Degradation Properties. *Front Cell Dev
Biol* 9: 581805

Burbidge K, Rademacher DJ, Mattick J, Zack S, Grillini A, Bousset L, Kwon O, Kubicki K, Simon
A, Melki R, *et al* (2022) LGALS3 (galectin 3) mediates an unconventional secretion of
SNCA/alpha-synuclein in response to lysosomal membrane damage by the autophagic-
lysosomal pathway in human midbrain dopamine neurons. *Autophagy* 18: 1020–1048

Calabresi P, Mechelli A, Natale G, Volpicelli-Daley L, Lazzaro GD & Ghiglieri V (2023) Alpha-
synuclein in Parkinson's disease and other synucleinopathies: from overt neurodegeneration
back to early synaptic dysfunction. *Cell Death Dis* 14: 176

Cavaliere F, Cerf L, Dehay B, Ramos-Gonzalez P, Giorgi FD, Bourdenx M, Bessede A, Obeso
JA, Matute C, Ichas F, *et al* (2017) In vitro α-synuclein neurotoxicity and spreading among

neurons and astrocytes using Lewy body extracts from Parkinson disease brains. *Neurobiol*
Dis 103: 101–112

Chavarría C, Rodríguez-Bottero S, Quijano C, Cassina P & Souza JM (2018) Impact of
monomeric, oligomeric and fibrillar alpha-synuclein on astrocyte reactivity and toxicity to
neurons. *Biochem J* 475: 3153–3169

Cullen V, Lindfors M, Ng J, Paetau A, Swinton E, Kolodziej P, Boston H, Saftig P, Woulfe J,
Feany MB, *et al* (2009) Cathepsin D expression level affects alpha-synuclein processing,
aggregation, and toxicity in vivo. *Mol Brain* 2: 5

Dehay B, Bourdenx M, Gorry P, Przedborski S, Vila M, Hunot S, Singleton A, Olanow CW,
Merchant KM, Bezard E, *et al* (2015) Targeting α-synuclein for treatment of Parkinson's
disease: mechanistic and therapeutic considerations. *Lancet Neurol* 14: 855–866

Domenico A di, Carola G, Calatayud C, Pons-Espinal M, Muñoz JP, Richaud-Patin Y,
Fernandez-Carasa I, Gut M, Faella A, Parameswaran J, *et al* (2019) Patient-Specific iPSC-
Derived Astrocytes Contribute to Non-Cell-Autonomous Neurodegeneration in Parkinson's
Disease. *Stem Cell Rep* 12: 213–229

Drobny A, Huarcaya SP, Dobert J, Kluge A, Bunk J, Schlothauer T & Zunke F (2022) The role of
lysosomal cathepsins in neurodegeneration: Mechanistic insights, diagnostic potential and
therapeutic approaches. *Biochim Biophys Acta (BBA) - Mol Cell Res* 1869: 119243

Dutta D & Donaldson JG (2012) Search for inhibitors of endocytosis. *Cell Logist* 2: 203–208

El-Agnaf OMA, Salem SA, Paleologou KE, Cooper LJ, Fullwood NJ, Gibson MJ, Curran MD,
Court JA, Mann DMA, Ikeda S-I, *et al* (2003) α-Synuclein implicated in Parkinson's disease is
present in extracellular biological fluids, including human plasma. *FASEB J* 17: 1–16

Emmanouilidou E, Melachroinou K, Roumeliotis T, Garbis SD, Ntzouni M, Margaritis LH,
Stefanis L & Vekrellis K (2010) Cell-Produced α-Synuclein Is Secreted in a Calcium-
Dependent Manner by Exosomes and Impacts Neuronal Survival. *J Neurosci* 30: 6838–6851

Filippini A, Gennarelli M & Russo I (2019) α-Synuclein and Glia in Parkinson's Disease: A
Beneficial or a Detrimental Duet for the Endo-Lysosomal System? *Cell Mol Neurobiol* 39:
161–168

Freeman D, Cedillos R, Choyke S, Lukic Z, Mcguire K, Marvin S, Burrage AM, Sudholt S, Rana
A, O'Connor C, *et al* (2013) Alpha-Synuclein Induces Lysosomal Rupture and Cathepsin
Dependent Reactive Oxygen Species Following Endocytosis. *PLoS ONE* 8: e62143

Fujiwara H, Hasegawa M, Dohmae N, Kawashima A, Masliah E, Goldberg MS, Shen J, Takio K
& Iwatsubo T (2002) α-Synuclein is phosphorylated in synucleinopathy lesions. *Nat Cell Biol*
4: 160–164

Gacko M, Minarowska A, Karwowska A & Minarowski Ł (2007) Cathepsin D inhibitors. *Folia*
Histochem Cytobiol 45: 291–313

Gallagher LE, Radhi OA, Abdullah MO, McCluskey AG, Boyd M & Chan EYW (2017)
Lysosomotropism depends on glucose: a chloroquine resistance mechanism. *Cell Death Dis*
8: e3014

Ghee M, Melki R, Michot N & Mallet J (2005) PA700, the regulatory complex of the 26S
proteasome, interferes with α-synuclein assembly. *FEBS J* 272: 4023–4033

Harackiewicz O & Grembecka B (2024) The Role of Microglia and Astrocytes in the
Pathomechanism of Neuroinflammation in Parkinson's Disease—Focus on Alpha-Synuclein.
J Integr Neurosci 23: 203

Henrich MT, Geibl FF, Lakshminarasimhan H, Stegmann A, Giasson BI, Mao X, Dawson VL,
Dawson TM, Oertel WH & Surmeier DJ (2020) Determinants of seeding and spreading of α-
synuclein pathology in the brain. *Sci Adv* 6: eabc2487

Hishikawa N, Hashizume Y, Yoshida M & Sobue G (2001) Widespread occurrence of
argyrophilic glial inclusions in Parkinson's disease. *Neuropathol Appl Neurobiol* 27: 362–372

Jacquet A de R, Tancredi JL, Lemire AL, DeSantis MC, Li WP & O'Shea EK (2021) The LRRK2
G2019S mutation alters astrocyte-to-neuron communication via extracellular vesicles and
induces neuron atrophy in a human iPSC-derived model of Parkinson's disease. *Elife* 10

Jones-Tabah J, He K, Karpilovsky N, Senkevich K, Deyab G, Pietrantonio I, Goiran T,
Cousineau Y, Nikanorova D, Goldsmith T, *et al* (2024) The Parkinson's disease risk gene
cathepsin B promotes fibrillar alpha-synuclein clearance, lysosomal function and
glucocerebrosidase activity in dopaminergic neurons. *Mol Neurodegener* 19: 88

Jucker M & Walker LC (2018) Propagation and spread of pathogenic protein assemblies in
neurodegenerative diseases. *Nat Neurosci* 21: 1341–1349

Jurga AM, Paleczna M, Kadluczka J & Kuter KZ (2021) Beyond the GFAP-Astrocyte Protein
Markers in the Brain. *Biomolecules* 11: 1361

Kam T-I, Hinkle JT, Dawson TM & Dawson VL (2020) Microglia and astrocyte dysfunction in
parkinson's disease. *Neurobiol Dis* 144: 105028

Kim S, Pajarillo E, Nyarko-Danquah I, Aschner M & Lee E (2023) Role of Astrocytes in
Parkinson's Disease Associated with Genetic Mutations and Neurotoxicants. *Cells* 12: 622

Koga S, Sekiya H, Kondru N, Ross OA & Dickson DW (2021) Neuropathology and molecular
diagnosis of Synucleinopathies. *Mol Neurodegener* 16: 83

Koob AO & Sacchetti P (2018) Pathology, Prevention and Therapeutics of Neurodegenerative
Disease. 81–102

Lee H-J, Bae E-J & Lee S-J (2014) Extracellular α-synuclein—a novel and crucial factor in Lewy
body diseases. *Nat Rev Neurol* 10: 92–98

Lee HJ, Suk JE, Bae EJ, Lee JH, Paik SR & Lee SJ (2008a) Assembly-dependent endocytosis
and clearance of extracellular alpha-synuclein. *Int J Biochem Cell Biol* 40: 1835–49

Lee H-J, Suk J-E, Bae E-J & Lee S-J (2008b) Clearance and deposition of extracellular α-
synuclein aggregates in microglia. *Biochem Biophys Res Commun* 372: 423–428

Lin L, Wu Z, Luo H & Huang Y (2024) Cathepsin-mediated regulation of alpha-synuclein in
Parkinson's disease: a Mendelian randomization study. *Front Aging Neurosci* 16: 1394807

Lindström V, Gustafsson G, Sanders LH, Howlett EH, Sigvardson J, Kasrayan A, Ingelsson M,
Bergström J & Erlandsson A (2017) Extensive uptake of α-synuclein oligomers in astrocytes
results in sustained intracellular deposits and mitochondrial damage. *Mol Cell Neurosci* 82:
143–156

Liu C-W, Giasson BI, Lewis KA, Lee VM, DeMartino GN & Thomas PJ (2005) A Precipitating
Role for Truncated α-Synuclein and the Proteasome in α-Synuclein Aggregation
IMPLICATIONS FOR PATHOGENESIS OF PARKINSON DISEASE*. *J Biol Chem* 280:
22670–22678

Loria F, Vargas JY, Bousset L, Syan S, Salles A, Melki R & Zurzolo C (2017) alpha-Synuclein
transfer between neurons and astrocytes indicates that astrocytes play a role in degradation
rather than in spreading. *Acta Neuropathol* 134: 789–808

Marotta NP, Ara J, Uemura N, Lougee MG, Meymand ES, Zhang B, Petersson EJ, Trojanowski
JQ & Lee VM-Y (2021) Alpha-synuclein from patient Lewy bodies exhibits distinct
pathological activity that can be propagated in vitro. *Acta Neuropathol Commun* 9: 188

Matsuki A, Watanabe Y, Hashimoto S, Hoshino A & Matoba S (2024) Cathepsin L prevents the
accumulation of alpha‐synuclein fibrils in the cell. *Genes Cells* 29: 328–336

McCarthy KD & Vellis J de (1980) Preparation of separate astroglial and oligodendroglial cell
cultures from rat cerebral tissue. *J cell Biol* 85: 890–902

McGlinchey RP, Lacy SM, Huffer KE, Tayebi N, Sidransky E & Lee JC (2019) C-terminal α-
synuclein truncations are linked to cysteine cathepsin activity in Parkinson's disease. *J Biol
Chem* 294: 9973–9984

McGlinchey RP & Lee JC (2015) Cysteine cathepsins are essential in lysosomal degradation of
α-synuclein. *Proc Natl Acad Sci* 112: 9322–9327

Mena MA & Yébenes JG de (2008) Glial Cells as Players in Parkinsonism: The "Good," the
"Bad," and the "Mysterious" Glia. *Neurosci* 14: 544–560

Mori F, Tanji K, Yoshimoto M, Takahashi H & Wakabayashi K (2002) Demonstration of α-
Synuclein Immunoreactivity in Neuronal and Glial Cytoplasm in Normal Human Brain Tissue
Using Proteinase K and Formic Acid Pretreatment. *Exp Neurol* 176: 98–104

Nalls MA, Blauwendraat C, Vallerga CL, Heilbron K, Bandres-Ciga S, Chang D, Tan M, Kia DA,
Noyce AJ, Xue A, *et al* (2019) Identification of novel risk loci, causal insights, and heritable

risk for Parkinson's disease: a meta-analysis of genome-wide association studies. *Lancet*
Neurol 18: 1091–1102

Ohgita T, Namba N, Kono H, Shimanouchi T & Saito H (2022) Mechanisms of enhanced
aggregation and fibril formation of Parkinson's disease-related variants of α-synuclein. *Sci*
Rep 12: 6770

Pang SY-Y, Ho PW-L, Liu H-F, Leung C-T, Li L, Chang EES, Ramsden DB & Ho S-L (2019)
The interplay of aging, genetics and environmental factors in the pathogenesis of
Parkinson's disease. *Transl Neurodegener* 8: 23

Panicker N, Ge P, Dawson VL & Dawson TM (2021) The cell biology of Parkinson's disease. *J*
Cell Biol 220: e202012095

Peng H, Chen S, Wu S, Shi X, Ma J, Yang H & Li X (2023) Alpha-synuclein in skin as a high-
quality biomarker for Parkinson's disease. *J Neurol Sci* 451: 120730

Poewe W, Seppi K, Tanner CM, Halliday GM, Brundin P, Volkmann J, Schrag A-E & Lang AE
(2017) Parkinson disease. *Nat Rev Dis Prim* 3: 17013

Polymeropoulos MH, Lavedan C, Leroy E, Ide SE, Dehejia A, Dutra A, Pike B, Root H,
Rubenstein J, Boyer R, *et al* (1997) Mutation in the α-Synuclein Gene Identified in Families
with Parkinson's Disease. *Science* 276: 2045–2047

Pons-Espinal M, Blasco-Agell L, Fernandez-Carasa I, Andrés-Benito P, Domenico A di,
Richaud-Patin Y, Baruffi V, Marruecos L, Espinosa L, Garrido A, *et al* (2024) Blocking IL-6
signaling prevents astrocyte-induced neurodegeneration in an iPSC-based model of
Parkinson's disease. *JCI Insight* 9: e163359

Qiao C, Yin N, Gu H, Zhu J, Ding J, Lu M & Hu G (2016) Atp13a2 Deficiency Aggravates
Astrocyte-Mediated Neuroinflammation via NLRP3 Inflammasome Activation. *CNS Neurosci*
Ther 22: 451–460

Qiao L, Hamamichi S, Caldwell KA, Caldwell GA, Yacoubian TA, Wilson S, Xie Z-L, Speake LD,
Parks R, Crabtree D, *et al* (2008) Lysosomal enzyme cathepsin D protects against alpha-
synuclein aggregation and toxicity. *Mol Brain* 1: 17

Quick JD, Silva C, Wong JH, Lim KL, Reynolds R, Barron AM, Zeng J & Lo CH (2023)
Lysosomal acidification dysfunction in microglia: an emerging pathogenic mechanism of
neuroinflammation and neurodegeneration. *J Neuroinflammation* 20: 185

Raj A, Banerjee R, Holla V, Kamble N, Yadav R, Pal PK & Datta I (2024) Dysregulation of
protein degradation and alteration of secretome in α-synuclein-exposed astrocytes:
implications for dopaminergic neuronal dysfunction. *Cell Commun Signal* 22: 574

Ramalingam N, Jin S-X, Moors TE, Fonseca-Ornelas L, Shimanaka K, Lei S, Cam HP, Watson
AH, Brontesi L, Ding L, *et al* (2023) Dynamic physiological α-synuclein S129 phosphorylation
is driven by neuronal activity. *npj Park's Dis* 9: 4

Rostami J, Holmqvist S, Lindström V, Sigvardson J, Westermark GT, Ingelsson M, Bergström J,
Roybon L & Erlandsson A (2017) Human Astrocytes Transfer Aggregated Alpha-Synuclein
via Tunneling Nanotubes. *J Neurosci* 37: 11835–11853

Russo MJ, Orru CD, Concha-Marambio L, Giaisi S, Groveman BR, Farris CM, Holguin B,
Hughson AG, LaFontant D-E, Caspell-Garcia C, *et al* (2021) High diagnostic performance of
independent alpha-synuclein seed amplification assays for detection of early Parkinson's
disease. *Acta Neuropathol Commun* 9: 179

Sanyal A, DeAndrade MP, Novis HS, Lin S, Chang J, Lengacher N, Tomlinson JJ, Tansey MG
& LaVoie MJ (2020) Lysosome and Inflammatory Defects in GBA1-Mutant Astrocytes Are
Normalized by LRRK2 Inhibition. *Mov Disord* 35: 760–773

Schildge S, Bohrer C, Beck K & Schachtrup C (2013) Isolation and Culture of Mouse Cortical
Astrocytes. *J Vis Exp*

Settembre C, Fraldi A, Medina DL & Ballabio A (2013) Signals from the lysosome: a control
centre for cellular clearance and energy metabolism. *Nat Rev Mol Cell Biol* 14: 283–296

Sorrentino ZA & Giasson BI (2020) The emerging role of α-synuclein truncation in aggregation
and disease. *J Biol Chem* 295: 10224–10244

Sorrentino ZA, Giasson BI & Chakrabarty P (2019) α-Synuclein and astrocytes: tracing the
pathways from homeostasis to neurodegeneration in Lewy body disease. *Acta Neuropathol*
138: 1–21

Stefanis L (2012) α-Synuclein in Parkinson's Disease. *Cold Spring Harb Perspect Med* 2:
a009399

Sveinbjornsdottir S (2016) The clinical symptoms of Parkinson's disease. *J Neurochem* 139:
318–324

Tsujimura A, Taguchi K, Watanabe Y, Tatebe H, Tokuda T, Mizuno T & Tanaka M (2015)
Lysosomal enzyme cathepsin B enhances the aggregate forming activity of exogenous α-
synuclein fibrils. *Neurobiol Dis* 73: 244–253

Tsunemi T, Ishiguro Y, Yoroisaka A, Valdez C, Miyamoto K, Ishikawa K, Saiki S, Akamatsu W,
Hattori N & Krainc D (2020) Astrocytes Protect Human Dopaminergic Neurons from α-
Synuclein Accumulation and Propagation. *J Neurosci* 40: 8618–8628

Uddin MdS & Lim LW (2022) Glial cells in Alzheimer's disease: From neuropathological
changes to therapeutic implications. *Ageing Res Rev* 78: 101622

Verkhratsky A, Zorec R & Parpura V (2017) Stratification of astrocytes in healthy and diseased
brain. *Brain Pathol* 27: 629–644

Wakabayashi K, Hayashi S, Yoshimoto M, Kudo H & Takahashi H (2000) NACP/alpha-
synuclein-positive filamentous inclusions in astrocytes and oligodendrocytes of Parkinson's
disease brains. *Acta Neuropathol* 99: 14–20

Walsh DM & Selkoe DJ (2016) A critical appraisal of the pathogenic protein spread hypothesis
of neurodegeneration. *Nat Rev Neurosci* 17: 251–260

Wang C, Yang T, Liang M, Xie J & Song N (2021) Astrocyte dysfunction in Parkinson's disease:
from the perspectives of transmitted alpha-synuclein and genetic modulation. *Transl*
Neurodegener 10: 39

Wang LH, Rothberg KG & Anderson RG (1993) Mis-assembly of clathrin lattices on endosomes
reveals a regulatory switch for coated pit formation. *J cell Biol* 123: 1107–1117

Wong CO (2020) Endosomal-Lysosomal Processing of Neurodegeneration-Associated Proteins
in Astrocytes. *Int J Mol Sci* 21

Xie YX, Naseri NN, Fels J, Kharel P, Na Y, Lane D, Burré J & Sharma M (2022) Lysosomal
exocytosis releases pathogenic α-synuclein species from neurons in synucleinopathy
models. *Nat Commun* 13: 4918

Yoon MC, Christy MP, Phan VV, Gerwick WH, Hook G, O'Donoghue AJ & Hook V (2022)
Molecular Features of CA-074 pH-Dependent Inhibition of Cathepsin B. *Biochemistry* 61:
228–238

Zeng J, Shirihai OS & Grinstaff MW (2020) Modulating lysosomal pH: a molecular and
nanoscale materials design perspective. *JoLS, J Life Sci* 2: 25–37

Zhang C, Pei Y, Zhang Z, Xu L, Liu X, Jiang L, Pielak GJ, Zhou X, Liu M & Li C (2022) C-
terminal truncation modulates α-Synuclein's cytotoxicity and aggregation by promoting the
interactions with membrane and chaperone. *Commun Biol* 5: 798

Figure Legends

Figure 1. **Primary astrocytes preferentiality take up and degrade αSyn PFFs via the**

lysosome. A) Bright-field images of astrocytes treated with phRodo-tagged αSyn monomer or

PFFs overtime. Scale bar = 50 µm. Quantification of phRodo+ area per total cell area overtime

(B) and at 24 H (C). Graphs B-C are expressed as a fold change of the 0 h monomer group. N =

6 wells/plate across 3 separate experiments. Unpaired student's t-test, ****p = < 0.0001. D)

Representative immunoblot of astrocyte lysates probed for total αSyn and β actin. Quantification

of band intensities for full length total αSyn (E) and partially degraded αSyn (F). N = 3 separate

experiments. One-way ANOVA with Sidak post hoc, ***$p = < 0.001$, ****$p = < 0.0001$. G)

Representative images of immunocytochemically stained for total αSyn, LAMP1 and nuclei with

Hoechst. Scale bar = 50 μm. Quantification of total αSyn+ spots per nuclei (H), LAMP1+ spots

per nuclei (I), total αSyn+ and LAMP1+ spots per nuclei (J) and nuclei number (K). N = 3-6

wells/plate across 3 separate experiments. Graphs H-K are expressed as a fold change of the 4

H group. One-way ANOVA with Sidak post hoc, **$p = < 0.01$, ****= $p < 0.0001$.

Figure 2. αSyn PFFs accumulate in lysosome-compromised (LC) primary astrocytes. A)

Representative images of astrocytes immunocytochemically stained for total αSyn and nuclei

with Hoechst. Scale bar = 50 μm. Quantification of total αSyn+ spots per nuclei (B) and nuclei

number (C), expressed as a fold change of the vehicle. N = 3-6 wells/plate across 3 separate

experiments. One-way ANOVA with Dunnett's post hoc, *$p = < 0.05$, ****= $p < 0.0001$. D)

Representative immunoblot of astrocyte lysates probed for total αSyn and β actin. Quantification

of band intensities for full length non-degraded total αSyn (E) and partially degraded αSyn (F). N

= 3 separate experiments. One-way ANOVA with Dunnett's post hoc, *$p = < 0.05$, **$p = < 0.01$.

G) Representative immunoblot of astrocyte lysates probed for total αSyn and β actin.

Quantification of band intensities for non-degraded αSyn (H) and partially degraded αSyn (I). N

= 3 separate experiments. One-way ANOVA with Dunnett's post hoc, *$p = < 0.05$. J)

Representative immunoblot of astrocyte lysates probed for total αSyn and β actin. Quantification

of band intensities for non-degraded αSyn (K) and partially degraded αSyn (L). N = 3 separate

experiments. One-way ANOVA with Sidak post hoc, *$p = < 0.05$, **$p = < 0.01$, ****$p = < 0.0001$.

Figure 3. Primary astrocytes prevent pS129 pathology in neurons induced by αSyn PFFs.

A) Schematic of the assay used in B-F. B) Representative images of DIV 21 neuroglia cultures

with increasing densities of primary astrocytes (Asts) added and immunocytochemically stained

for pS129, GFAP, MAP2, and nuclei with Hoechst. Scale bar = 50 μm. Quantification pS129+

spots per MAP2 area (C), MAP2 area (D), GFAP area (E), and nuclei number (F), expressed as

a fold change of the 0k Ast group. N = 3-6 wells/plate across 3 separate experiments. One-way

ANOVA with Sidak post hoc, ****= p < 0.0001. G) Representative images of DIV 21 neuroglia

cultures (-FUdR) and pure neuron cultures (+ FUdR) and immunocytochemically stained for

pS129, GFAP, MAP2, and nuclei with Hoechst. Scale bar = 50 μm. Quantification pS129+

spots per MAP2 area (H), MAP2 area (I), GFAP area (J), and nuclei number (K), expressed as

a fold change of the (-) FUdR) / (+) PFF group.   N = 3-6 wells/plate across 3 separate

experiments. Two-way ANOVA with Sidak post hoc, ***= p < 0.001, ****= p < 0.0001. L)

Schematic of the assay used in M-Q. M) Representative images of DIV 21 pure neuron cultures

-/+ Asts and immunocytochemically stained for pS129, GFAP, MAP2, and nuclei with Hoechst.

Scale bar = 50 μm. Quantification pS129+ spots per MAP2 area (N), MAP2 area (O), GFAP

area (P), and nuclei number (Q), expressed as a fold change of the 15k Asts group. N = 3-10

wells/plate across 3 separate experiments. Unpaired student's t-test, ****= p < 0.0001.

Figure 4. **LC astrocytes do not prevent pS129 pathology in neurons induced by αSyn**

PFFs. A) Schematic of the assay used in B-F. B) Representative images of DIV 21 neuroglia

cultures -/+ healthy or LC Asts and immunocytochemically stained for pS129, GFAP, MAP2,

and nuclei with Hoechst. Scale bar = 50 μm. Quantification pS129+ spots per MAP2 area (C),

MAP2 area (D), GFAP area (E), and nuclei number (F), expressed as a fold change of the 0k

Ast group. N = 3-6 wells/plate across 3 separate experiments. One-way ANOVA with Sidak

post hoc, *= p < 0.05, **= p < 0.01, ****= p < 0.0001. G) Schematic of the assay used in H-L. H)

Representative   images   of   DIV   21   neuron   cultures   +   healthy   or   LC   Asts   and

immunocytochemically stained for pS129, GFAP, MAP2, and nuclei with Hoechst. Scale bar =

50 μm. Quantification pS129+ spots per MAP2 area (I), MAP2 area (J), GFAP area (K), and

nuclei number (L), expressed as a fold change of the vehicle-treated Ast group. N = 3-6

wells/plate across 4 separate experiments. One-way ANOVA with Dunnett's post hoc, *= p <

0.05, **= p < 0.01, ****= p < 0.0001.

Figure 5. αSyn PFF preloaded LC astrocytes induce pS129 pathology in neurons. A)

Schematic of the assay used in B-F. B) Representative images of DIV 21 neuroglia cultures +

αSyn PFF-preloaded Asts and immunocytochemically stained for pS129, GFAP, MAP2, and

nuclei with Hoechst. Scale bar = 50 µm. Quantification pS129+ spots per MAP2 area (C), MAP2

area (D), GFAP area (E), and nuclei number (F), expressed as a fold change of the vehicle-

treated Ast group. N = 3-6 wells/plate across 3 separate experiments. One-way ANOVA with

Dunnett's post hoc, ****= p < 0.0001. G) Schematic of the assay used in H-L. H) Representative

images of DIV 21 neuron cultures + αSyn PFF-preloaded Asts and immunocytochemically

stained for pS129, GFAP, MAP2, and nuclei with Hoechst. Scale bar = 50 µm. Quantification

pS129+ spots per MAP2 area (I), MAP2 area (J), GFAP area (K), and nuclei number (L),

expressed as a fold change of the vehicle-treated Ast group. N = 3-6 wells/plate across 4

separate experiments. One-way ANOVA with Dunnett's post hoc, *= p < 0.05, ****= p < 0.0001.

Extended View Figure 1. Human primary astrocyte take up and degrade αSyn PFFs via

the lysosome. A) Representative immunoblot of mouse primary astrocyte lysates probed for

total αSyn, using MJFR1 antibody, and β actin. B) Quantification of band intensities for full

length total αSyn. N = 3 separate experiments. One-way ANOVA with Sidak post hoc, **p = <

0.01, ****p = < 0.0001. C) Representative immunoblot of human astrocyte lysates probed for

total αSyn, using Syn1 antibody, and β actin. Quantification of band intensities for full length

total αSyn (D) and partially degraded αSyn (E). N = 3 separate experiments. One-way ANOVA

with Sidak post hoc, **p = < 0.01, ***p = < 0.001, ****p = < 0.0001. F) Representative images of

human astrocytes immunocytochemically stained for total αSyn, LAMP1 and nuclei with

Hoechst. Scale bar = 50 µm. Quantification of total αSyn+ spots per nuclei (G), LAMP1+ spots

per nuclei (H), total αSyn+ and LAMP1+ spots per nuclei (I) and nuclei number (J), are

expressed as a fold change of the 4 H group. N = 3-6 wells/plate across 3 separate

experiments. Graphs G-J One-way ANOVA with Sidak post hoc, *p = < 0.05, ***p = < 0.001,

****= p < 0.0001.

Extended View Figure 2. Chloroquine causes decreased lysosome number, increased

size, and damage. A) Representative images of astrocytes immunocytochemically stained for

LAMP1 and nuclei with Hoechst. Scale bar = 50 µm. Quantification of total LAMP1+ spots per

nuclei (B) and Mean LAMP1 spot size (C), expressed as a fold change of the vehicle. N = 3-6

wells/plate across 3 separate experiments. One-way ANOVA with Dunnett's post hoc, ****= p <

0.0001. D) Representative images of astrocytes immunocytochemically stained for Gal3, total

αSyn, and nuclei with Hoechst. Scale bar = 50 µm. Quantification of total αSyn+ spots per nuclei

(E), Gal3+ spots per nuclei (F), total αSyn+ and Gal3+ spots per nuclei, and nuclei number (H),

expressed as a fold change of the vehicle. N = 3-6 wells/plate across 3 separate experiments.

One-way ANOVA with Dunnett's post hoc, ***= p < 0.001, ****= p < 0.0001.

A

	0 H	6 H	12 H	24 H
Monomer				
PFFs				

B

○ αSyn Monomer
● αSyn PFFs

pHrodo+ area per total cell area
Normalized to 0 H Monomer

Time Post pHrodo αSyn Addition (Hours)

C

pHrodo+ area per total cell area
Normalized to 0 H Monomer

Monomer   PFFs   ****

D

UT | 1 μg/mL αSyn Monomer | 1 μg/mL αSyn PFFs
4 H 24 H 48 H 72 H | 4 H 24 H 48 H 72 H

16 kDa → Full Length
Degraded Bands

40 kDa → actin

WB: Syn1 (Total αSyn)

E αSyn Full Length Band

Total Syn1 αSyn/actin Band Density (AU)

UT | 4 H 24 H 48 H 72 H | 4 H 24 H 48 H 72 H
1 μg/mL Monomer | 1 μg/mL PFFs
**** **** ****

F αSyn degraded bands only

Total Syn1 αSyn/actin Band Density (AU)

UT | 4 H 24 H 48 H 72 H | 4 H 24 H 48 H 72 H
1 μg/mL Monomer | 1 μg/mL PFFs
**** **** **** ***

G

| PBS | 1 μg/mL αSyn PFFs |
| | 4 H | 24 H | 48 H | 72 H |

High Mag

αSyn/LAMP1/Hoechst

H

αSyn+ spots per nuclei
Normalized to 4 H

PBS 4 H 24 H 48 H 72 H
1 μg/mL PFFs
**** **** **** ** ns ns

4 H Avg

I

LAMP1+ spots per nuclei
Normalized to 4 H

PBS 4 H 24 H 48 H 72 H
1 μg/mL PFFs
ns ns ns

4 H Avg

J

LAMP1+/αSyn+ spots per nuclei
Normalized to 4 H

PBS 4 H 24 H 48 H 72 H
1 μg/mL PFFs
**** **** ** **** ns ns

4 H Avg

K

Nuclei Number
Normalized to 4 H

PBS 4 H 24 H 48 H 72 H
1 μg/mL PFFs
ns ns ns

4 H Avg

A

LC-Ast αSyn Spreading Model

DIV 0 — αSyn A53T neuroglia cultures 35k/well (96 well plate)

DIV 7 — + PFF preloaded LC-asts (25k/well)

DIV 21 — Fix and ICC: GFAP, MAP2, pS129, Hoechst

= healthy astrocyte
= PFF preloaded astrocyte
= PFF preloaded LC astrocyte
= neuron
= αSyn aggregates

CD1 astrocytes 1M cells/well (6 well plate)

5 H Treatment: 1 μg/mL PFFs

16 H *PFFs washed out Treatment: Vehicle (DMSO), CQ (25 μM)

Lift PFF preloaded LC-asts

B No PFFs Asts / PFFs Veh Asts / PFFs CQ Asts

GFAP / MAP2 / pS129 / High Mag

C pS129+ Spots per MAP2 area Normalized to Vehicle
ns / ****
No PFFs + / PFFs − / Vehicle − / CQ −

D MAP2 area Normalized to Vehicle
ns / ns

E GFAP area Normalized to Vehicle
ns / ns

F Nuclei Number Normalized to Vehicle
ns / ns

G

Pure αSyn A53T Neuron Addition to LC-Ast Spreading Model

DIV -1 — CD1 astrocytes 10k/well (96 well plate)

5 H Treatment: 1 μg/mL PFFs

16 H *PFFs washed out Treatment: Vehicle (DMSO), CQ (25 μM), CA-074 (10 μM)

DIV 0 — + Pure neurons (45k/well) *Asts rinsed of compounds

DIV 21 — Fix and ICC: GFAP, MAP2, pS129, Hoechst

= healthy astrocyte
= PFF preloaded astrocyte
= PFF preloaded LC astrocyte
= neuron
= αSyn aggregates

H No PFFs Asts / PFFs Veh Asts / PFFs CQ Asts / PFFs CA-074 Asts

GFAP / MAP2 / pS129 / High Mag

I pS129+ spots per MAP2 area Normalized to Vehicle
ns / **** / *

J MAP2 area Normalized to Vehicle
ns / ns

K GFAP area Normalized to Vehicle
ns / ns

L Nuclei Number Normalized to Vehicle
ns / ns