The potential role of volatile organic compounds on the colonisation of deadwood by saproxylic beetles

Claudio Sbaraglia[1,2,][*], Simon Thorn [1,3], Lucie Ambrozova[1], Lukas Cizek[1], Petr Kozel[1],

Daniel Sebastián Rodríguez León[4], Thomas Schmitt[4] & Lukas Drag[1]

[1] Biology Centre CAS, Institute of Entomology, České Budějovice, Czechia

[2] University of South Bohemia, Faculty of Science, České Budějovice, Czechia

[3] Department of Biology, Philipps Universität Marburg, Marburg, Germany

[4] Department of Animal Ecology and Tropical Ecology, Biocentre, University of

Würzburg, Würzburg, Germany

* Corresponding author: claudiosbaraglia95@gmail.com

ORCID numbers:

Claudio Sbaraglia 0000-0002-2893-2577, Lukas Drag 0000-0002-6002-9214, Simon

Thorn 0000-0002-3062-3060, Lucie Ambrozova 0000-0002-3258-2661, Petr Kozel 0000-

0001-8779-8024, Lukas Cizek 0000-0002-6447-2002, Thomas Schmitt 0000-0002-6719-

8635, Daniel Sebastián Rodríguez León 0000-0001-9637-1364

Keywords: bark beetles; broadleaf; conifer; decay; monoterpene; sesquiterpene; VOC;
wood-inhabiting fungi

Abstract

1. Volatile organic compounds (VOCs) emitted by deadwood are increasingly

recognised as key olfactory cues used by saproxylic beetles to locate suitable

substrates. However, how beetle communities respond to these cues during the

colonisation of deadwood remains poorly understood. To address this gap, it is

essential to quantify both VOC emissions and beetle assemblages, and to

experimentally disentangle the main ecological drivers of the deadwood volatilome

(i.e., tree species, sun exposure).

2. To explore the potential role of VOCs on saproxylic beetle colonisation, we exposed

1200 freshly cut branches of oak, beech, spruce, and pine across Central Europe. To

mimic natural variation in deadwood and disturbance processes, the bundles were

experimentally manipulated either by sterilisation (to reduce endogenous fungi),

fungal inoculation with a brown rot fungus (*Fomitopsis pinicola*), fungal inoculation

with a white rot fungus (*Fomes fomentarius*), or burning. From each deadwood

bundle, we sampled VOCs and reared emerging beetles.

3. From deadwood bundles, we sampled 448 substances, 89 of which were identified as

VOCs, and reared 134 saproxylic beetle species. We observed distinct VOC profiles

emitted by broadleaf (oak, beech) and conifer (pine, spruce) species, aligning with the

beetles' tree-type preferences. In conifers, bark beetles, longhorn beetles, and jewel

beetles were associated with different chemical cues, whereas beetle taxonomic

separation was not observed in broadleaf species. Although experimental treatments

(sterilisation, fungal inoculation, burning) altered VOC composition, they could not

explain beetle colonisation patterns.

4. Our study highlights that VOCs emitted during the early decay stage of deadwood are

associated with distinct saproxylic beetle assemblages. VOC composition varied

across tree species and treatments, suggesting that chemical variation reflects multiple

ecological factors. These findings indicate that deadwood diversity, particularly in

tree species, contributes to chemical heterogeneity, which is linked to broader beetle

assemblages. Forest conservation efforts may therefore need to consider the role of

chemical variation in deadwood, as it could influence colonisation patterns of

saproxylic beetles and affect the success of biodiversity management strategies.

1. Introduction

Deadwood serves as a crucial habitat for numerous groups of organisms that drive its

decomposition (Grove, 2002), thereby contributing to nutrient cycling and soil development

(Albisetti et al., 2003; Pichler et al., 2012). While fungi and bacteria are the primary

decomposers of deadwood (Fukasawa, 2021; Seibold et al., 2022), insects facilitate

decomposition both by tunnelling through wood, enhancing microbial colonisation, and by

introducing symbiotic fungi and bacteria (Birkemoe et al., 2018). As deadwood decays, there

are various changes in its texture and structure, including an increased occurrence of fungi

(Ulyshen & Hanula, 2010; Seibold et al., 2023). This leads to variation in the emission of

chemical cues, which further influences the composition of deadwood-dependent organisms

(Leather et al., 2014).

65   In deadwood, the composition of Volatile Organic Compounds (VOCs), low-

molecular-weight chemical substances that easily evaporate under natural conditions, changes

over time, and it is shaped by many factors, such as tree species (Mäki et al., 2021), rot type

(Mali et al., 2019), or fungal development stage (Fäldt et al., 1999). Typically, VOCs emitted

by fungi during wood degradation are monoterpenes, sesquiterpenes, and oxygenated VOCs,

i.e. alcohols, ketones, and acetates with chain-length between four and 15 C-atoms (Evans et

al., 2008; El Ariebi et al., 2016). Due to their different wood degradation mechanisms, brown

rot fungi tend to emit higher levels of monoterpenes, while white rot species tend to release

more protein-based volatiles (Mali et al., 2019). VOC emissions also may vary between

fungal structures, with mycelium and sporocarps producing different blends (Orban et al.,

2023), which may attract specific species adapted to various ecological niches (Johansson et

al., 2006). Because fungi are among the first colonisers of deadwood, their early

establishment can lead to priority effects (Chase, 2003). That is, the timing and order of

species arrival shape the assembly of later-arriving organisms (Fukami et al., 2010). Early

fungal colonisers involved in wood degradation alter VOC emissions (Mäki et al., 2021),

potentially influencing the colonisation of saproxylic beetle assemblages attracted to these

VOCs.

Saproxylic beetles are a diverse and ecologically important group of organisms

associated with deadwood. Their colonisation of deadwood is mediated by a mix of visual

and olfactory cues (Jonsell & Nordlander, 1995; Cavaletto et al., 2020). The hypothesis that a

single or a few specific compounds drive host localisation in beetles has limited supporting

evidence (Fraenkel, 1959; Nottingham et al., 1991), whereas the idea that blends or whole

groups of volatiles aid host localisation has received more convincing support in the last

decades (Visser & Avé, 1978; Fraser et al., 2003; Bruce et al., 2005). For instance, the

synergy between terpenoids (e.g., monoterpenes, sesquiterpenes, diterpenes) emitted by

various tree species and the pheromones produced by beetles act as semiochemicals, inducing

attraction or repulsion in several bark and wood-boring beetles (Allison et al., 2004; Miller,

2006; Vuts et al., 2016).

The complex interactions among deadwood, wood-inhabiting fungi, and saproxylic

beetles potentially mediated by VOCs in early-decay stages remain largely unexplored,

despite these stages being widespread and representing key time windows for the colonisation

of diverse beetle communities. While most field studies have focused on individual

saproxylic beetle species and VOCs from live trees (Seybold et al., 2006; Isidorov et al.,

2010; Hulcr et al., 2011), studies focusing on VOCs emitted by deadwood have been mainly

conducted under laboratory conditions (Mali et al., 2019; Mäki et al., 2021). As a result, little

is known about how beetle communities are associated with VOCs released from fresh

deadwood under natural conditions.

To investigate the colonisation process of saproxylic beetles mediated by VOCs, we

first collected VOCs from fresh deadwood bundles exposed in natural conditions, and then

we sampled the beetle assemblages reared from the same bundles. To simulate natural

heterogeneity in deadwood, we created a diverse range of conditions by exposing bundles of

different treatments (fungi-depleted, fungi-inoculated, burned) in combination with four tree

species (oak, beech, spruce, and pine). This experimentally manipulated approach with a

balanced design enabled us to evaluate beetle colonisation choices as a potential response to

chemical cues and to test the following hypotheses:

i)   VOC composition from deadwood bundles differs among tree species, with
        conifer and broadleaf species emitting distinct VOC profiles.

ii)   VOC composition differs among treatments: fungi-inoculated bundles emit
        distinct VOCs due to differences in decay type (brown vs. white rot), while
        burned bundles show unique profiles due to thermal degradation.

iii)   All saproxylic beetles are positively associated with their host tree-specific VOCs;
        and niche specialists, mycetophagous and pyrophilous beetle species, are
        positively associated with VOCs emitted by fungi-inoculated and burned bundles,
        respectively.

2. Methods

122 2.1 Study sites

The 10 study sites span a 650 km longitudinal gradient in Central Europe, ranging from

9.68° E to 17.99° E, with elevations between 344 m and 1069 m (Fig. 1a). These sites include

diverse forest ecosystems, such as mixed mountain forests (spruce, fir, and beech), lowland

oak and pine forests, and upland forests dominated by beech, hornbeam, and maple. (Full site

descriptions, including detailed environmental characteristics, are available in (Appendix S1:

Table S1)

129 2.2 Wood collection and processing

Freshly cut logs were obtained in April 2022 from forest sites within a 20 km radius of Markt

Erlbach, northern Bavaria (Germany). Each log had a similar size with a mean diameter of

7.8 cm (SD ± 1.33) and a mean length of 51.8 cm (SD ± 0.73). Logs consisted of four tree

species: European beech (*Fagus sylvatica*), oak (*Quercus robur/petraea*), pine (*Pinus*

sylvestris), and Norway spruce (*Picea abies*). In total, we prepared 1200 logs (300 logs for

each tree species). To simulate a broad range of conditions, we treated the bundles by light

surface burning, fungal depletion via autoclaving (sterilised), and fungal inoculation with

Fomitopsis pinicola (brown rot) and *Fomes fomentarius* (white rot) (Appendix S1: Section

1).

139 2.3 Experimental design

We established two plots in each site, one in the forest interior shaded by the forest canopy

(hereinafter, as "shaded") and the other in a nearby open landscape exposed to direct sun for

most of the daytime (hereinafter, as "sun-exposed") (Fig. 1b). A total of 1200 logs were

grouped in sets of three, forming 400 bundles, ensuring that logs of the same tree species and

treatments were bundled together. Our experimental design thus consisted of 20 bundle

combinations (4 tree species * 5 treatments, incl. control) exposed in both shaded and sun-

exposed plots, resulting in 40 bundles per site. Two or three bundles were suspended on 2-

metre-high iron poles in a random position to minimise the sampling bias, but at the same

time avoid the presence of the same treatments or tree species on the same pole. Bundles

were exposed from May to October 2022. At the end of the exposure period (October 2022),

all 400 bundles from the experimental sites were placed in plastic tubes and stored at the

rearing station. Each bundle was placed in a plastic tube with a fine mesh-covered ventilation

hole and a collector tube containing propylene glycol as a preservative (Appendix S1: Fig.

S1). Reared beetles were collected from the emergence tubes in September 2023 and

September 2024 and stored in a freezer until identified to species level. The identified beetles

were classified as saproxylic following the criteria in Seibold et al. (2015).

2.4 Collection and identification of VOCs

VOC collection was conducted by active headspace sampling using an adsorbent tube and air

sampling pump for 15 minutes after the deadwood bundle was enclosed in a polyacetate bag

from one side for 45 minutes. The collected samples were analysed using a thermal

desorption unit (TDU; TD100-xr, Markes, Offenbach am Main, Germany) connected to a gas

chromatograph-mass spectrometer (GC/MS; Agilent 7890B GC and 5977 MS, Agilent

Technologies, Palo Alto, USA). We detected 448 peaks in the chromatograms from 400

deadwood bundles across the study sites between July 8–15, 2022, after analysing the volatile

traps using gas chromatography/mass spectrometry coupled to the thermal desorption unit.

Because environmental conditions can alter or dilute VOC emissions (Paiva, 2000), all

samples were collected under comparable weather conditions, including an average site

temperature of ~25 °C, mostly sunny skies, low wind intensity, and dry bundles (Appendix

S1: Table S2), values which suggest a stable and favourable condition for VOCs sampling.

Control samples were collected at each site at about 30 metres from the bundles to account

for background VOCs. Only peaks emitted at least five times more in bundles than in controls

were retained (Appendix S1: Section 2).

2.5 Data analysis

All statistical analyses were performed in R (v4.4.3; R Core Team, 2025) through the

RStudio platform (RStudio Team, 2024). Figures and data manipulation were carried out

using the 'tidyverse' package (Wickham et al., 2019).

2.6 VOCs selection

To assess how tree species, treatment, and exposure influence VOC emissions, we modelled

the relative abundance of each peak (response variable) separately using linear mixed-effect

models (LMM) with a Gaussian error distribution from the 'lme4' package (Bates et al.,

2015). Tree species (oak, beech, spruce, pine), treatment (control, burned, fungi-depleted,

brown rot, white rot), and exposure (sun-exposed, shaded) were included as fixed effects. To

account for the hierarchical structure of the experiment, the site affiliation was included as a

random effect. Peaks with poor model convergence were excluded from further analysis. A

total of 89 peaks were retained in the final matrix and used for further analysis. We

associated individual peaks with treatments and tree species using Tukey's post hoc

comparisons from the LMM models using the 'multcomp' package (Hothorn et al., 2008).

Comparisons were based on model-estimated t-values, with p-values adjusted for multiple

testing. Inoculated logs (brown and white rot) were compared to fungi-depleted logs to isolate

the effect of inoculation, as both were sterilised. Fungi-depleted and burned logs were

compared to untreated controls. For tree species, we tested for species-level differences;

when non-significant (e.g., beech vs oak), compounds were assigned to broader categories

("conifer" or "broadleaf") and tested using the same criteria (Appendix S2).

2.7 Composition analysis

To test whether VOC composition in deadwood bundles differed among tree species and

treatments, we first conducted a preliminary Detrended Correspondence Analysis (DCA),

which revealed a gradient length of 3.39, indicating that a unimodal ordination method was

appropriate (Lepš & Šmilauer, 2003). Based on this result, we performed a Canonical

Correspondence Analysis (CCA) from the 'vegan' package (Oksanen et al., 2001), using the

relative abundance of compounds as the response variable. Tree species, treatment, and their

interaction were included as explanatory variables, and exposure time as a covariate. To

account for differences among study sites, sites were included as a conditioning variable,

effectively removing their influence before testing the effects of the explanatory variables.

Significant differences among explanatory variables were tested through Permutational

Multivariate Analysis of Variance (PERMANOVA) using the 'adonis2' function, with 999

permutations. To account for the hierarchical study design, site affiliations were included as

strata for permutation constraints. Heatmaps were generated through the 'ggplot2' R package

(Wickham, 2016) to visualise potential shifts in VOC composition in response to the applied

treatments and their interaction with tree species.

2.8 Saproxylic beetle preferences

To identify the specific association of saproxylic beetles with tree species and treatments, for

each species we calculated Indicator Values (IndVal) using the 'multipatt' function in the

'indicspecies' package (Cáceres & Legendre, 2009), which quantifies the species' preference

as a percentage. The IndVal is based on two components: specificity (how restricted a species

is to a particular group) and fidelity (how consistently it occurs within that group) (Dufrêne &

Legendre, 1997). The significance of the association was tested through 999 permutations.

Species with significant IndVal ($p < 0.05$) for tree species and/or treatments were retained for

further analysis.

2.9 Association VOC – Beetles

To explore whether beetle colonisation is associated with VOC profiles of specific tree

species and treatments, we performed Non-Metric Multidimensional Scaling (NMDS) via the

'vegan' package (Oksanen et al., 2001), with four dimensions ($k = 4$) based on beetle

abundance, using 40 species selected through Indicator Species Analysis. To assess the

potential association between beetles and VOCs, we fitted the peaks into the NMDS

ordination space using the "envfit" function. A significant association was tested through 999

permutations, accounting for the variation in VOC emissions related to sites by permuting

within each site. Only significant peaks ($p < 0.05$) were plotted.

3. Results

We detected 89 VOCs primarily emitted from deadwood bundles across the different

treatments and tree species. Of these, most peaks were sesquiterpenes (25%) and

monoterpenes (22%), followed by hydrocarbons (14%), alcohols (9%), and unidentified

compounds (9%). Less represented classes included aldehydes (5%), fatty acids (4%), ethers,

ketones, and lactones (each 2%), while disulfides, esters, and furfural were each represented

by a single peak. In total, 41 peaks were tentatively identified at the substance level

(Appendix S1: Table S3).

Within two years, we reared 134 species of saproxylic beetles (24,696 individuals) from the

bundles, with Curculionidae being the most abundant and species-rich family (13,200

individuals across 25 species). The most abundant species were *Xylosandrus germanus* (5091

individuals), *Ernobius mollis* (3397), *Anthaxia quadripunctata* (3231), *Scolytus intricatus*

(2668), and *Taphrorychus bicolor* (1468).

3.1 VOC composition

We found significant effects of tree species (expl. var. = 16.85%, pseudo-F = 27.43, p =

0.001), treatment (expl. var. = 1.59%, pseudo-F = 1.94, p = 0.001), their interaction (expl.

var. = 3.26%, pseudo-F = 1.32, p = 0.003), and exposure (expl. var. = 0.70%, pseudo-F =

3.42, p = 0.001) on the VOC composition (Fig. 2). CCA revealed that VOC collected from

deadwood bundles clustered according to broadleaf and conifer tree species, with a clear

separation between spruce and pine bundles, but not by treatment (Fig. 2). Heatmaps showed

that the relative abundance of VOC composition varied across tree species in response to the

applied treatments (Fig. 3). Among the selected peaks, spruce emitted the fewest VOCs

compared to other tree species. In conifer bundles, we observed a shift in the relative VOC

composition: monoterpenes were more abundant in control bundles, whereas sesquiterpenes

emissions increased in fungi-inoculated bundles. In broadleaf bundles, heatmaps indicated

increased sesquiterpene emissions in fungi-depleted and fungi-inoculated treatments

compared to controls.

3.2 Indicator beetle species

The Indicator Species Analysis identified 45 saproxylic beetle species associated with tree

species and 14 associated with treatments (Appendix S3). In our dataset, most beetle species

exhibited a preference for a specific tree species. Oak and spruce supported the highest

number of beetle species, with 12 and 6 species, respectively. Regarding treatments,

Xylosandrus germanus and *Synchita mediolanensis* mainly occurred in fungi-depleted and

inoculated bundles, while *Leiopus nebulosus* and *Orthoperus atomus* showed a preference for

manipulated bundles (fungi-depleted, burned, or fungi-inoculated). Additionally, *Bitoma*

crenata and *Cryptophagus scanicus* showed a weak but significant association with brown-

rot-inoculated and burned bundles, respectively.

3.3 VOCs – Beetle associations

We displayed indicator beetle species and VOC peaks in NMDS ordination space to explore

their relationship (Fig. 4). Out of the 42 VOC peaks significantly associated with the

ordination axes, 18 showed associations at the species level, seven with pine, five with beech,

four with spruce, and two with oak. In cases where differences between species within a tree

type (e.g., beech or oak and pine or spruce) were not significant, peaks were broadly

categorised by tree type. Based on this grouping, 17 peaks were associated with broadleaf

trees and two with conifers. Five peaks did not show any specific tree association. Beetles

and VOC peaks clustered by tree species (Fig. 4a), with broadleaf-, beech-, and oak-

associated peaks aligning with higher abundances of broadleaf-associated beetles;

additionally, peaks characteristic of conifers were associated with increased abundances of

spruce- and pine-associated beetles. Regarding treatments, eight peaks were linked to fungi-

depleted bundles, two to control, and one each to white rot and burned treatments. None were

associated with brown rot, and 30 had no treatment association. Beetles and peaks did not

cluster clearly by treatment (Fig. 4b). An exception was the ambrosia beetle *X. germanus*,

which was associated with fungi-depleted bundles as well as with some peaks characteristic

of this treatment. The burned-associated peak pointed towards *Cryptophagus scanicus*, an

indicator for burned bundles.

4. Discussion

Our study offers a novel field-based approach combining VOC sampling of deadwood with

insect rearing of saproxylic species to investigate chemically mediated colonisation processes

in deadwood. We show that VOC composition varies across host tree species and, to a lesser

extent, across treatments, and that these chemical differences are associated with distinct

saproxylic beetle assemblages. This highlights the potential ecological relevance of host-

specific volatile cues during the early stages of deadwood colonisation for saproxylic beetle

communities.

4.1 Tree species variation in VOCs composition

As hypothesised, tree species accounted for most of the variation in VOC composition (Fig.

2). The distinct chemical profiles between conifers and broadleaves likely reflect differences

in wood anatomy and stress-response strategies. Conifers store terpene-rich oleoresin in

specialised structures (resin ducts, glands), which release volatiles like monoterpenes and

sesquiterpenes upon damage (Franceschi et al., 2005; Raffa, 2014; Celedon & Bohlmann,

2019). In contrast, broadleaf species lack such resin-containing structures and therefore

exhibit lower terpene emissions upon physical damage (Holighaus & Schulz, 2006; Thakeow

et al., 2007). We observed a distinct VOC composition between spruce and pine bundles (Fig.

2), with pine emitting both a greater number of VOCs and higher levels of monoterpenes in

control bundles. These differences likely reflect their contrasting defence strategies: pine

relies more on constitutive defences, storing large amounts of resin in preformed ducts, while

spruce depends more on induced responses involving de novo synthesis of terpenoids and

phenolics (Franceschi et al., 2005; Zhao et al., 2011; Celedón & Bohlmann, 2019). Since

VOCs were collected from deadwood, where induced responses are inactive, the stronger

emissions from pine likely result from its larger stored resin reserves, whereas spruce's

reliance on inducible pathways may explain its lower VOC emissions.

4.2 Fungal colonisation influences VOC composition

The breakdown of wood tissues during fungal colonisation releases a wide array of VOCs

(Mäki et al., 2021). In our design, we minimised the a priori fungal presence through

sterilisation and used fungi-depleted bundles as controls to isolate the effect of inoculation

from sterilisation. Although the variation explained by our manipulated deadwood

(inoculation, fungi-depletion, and burning) was low, it significantly influenced VOC

composition.

We observed higher relative abundances of monoterpenes in pine control bundles

compared to those that were fungi-depleted, inoculated, or burned (Fig. 3). High monoterpene

levels in freshly cut logs are expected in conifers, due to the passive release of oleoresin after

cutting (Franceschi et al., 2005; Isidorov et al., 2010). Interestingly, this pattern was not

observed in spruce bundles, which aligns with findings by Mäki et al. (2021), who reported

that monoterpene emissions peak earlier in pine sawdust but later in spruce. However, we

found no increase in monoterpene emissions following inoculation in conifers (Fig. 3),

suggesting that fungal colonisation did not enhance oleoresin release, possibly due to limited

release or rapid degradation of monoterpenes. Alternatively, the heat during the sterilisation

and burning process influenced the monoterpenes' emissions.

Contrary to conifers, beech and oak do not store monoterpenes-rich oleoresin, but rely

on tannins and phenolic compounds in their bark as a defence against biotic attacks (Dübeler

et al., 1997; Mämmelä, 2001). Fungal degradation of wood in these species is therefore more

likely to produce derivatives of these compounds rather than increase monoterpene

emissions. In our study, such compounds were indeed detected, including orcinol, thymol,

and cymeneol. Their presence, particularly in fungi-inoculated bundles, suggests that fungal

activity can enhance the emission of phenolic volatiles, with a stronger effect observed in

beech than in oak (Fig. 3).

Sesquiterpene emissions in deadwood are linked to fungal degradation, as observed

under laboratory conditions, in pine (Mäki et al., 2021), spruce (Mali et al., 2019), and beech

(Holighaus & Schütz, 2006; El Ariebi et al., 2016). In our study, fungi-inoculated bundles in

conifers showed higher relative abundances of sesquiterpenes compared to fungi-depleted

ones (Fig. 3). However, this pattern was not observed in broadleaf bundles, where differences

appeared mainly between control and fungi-depleted bundles. This contrast likely reflects

differences in fungal colonisation processes and substrate properties between conifers and

broadleaves. In conifer wood, fungal colonisation is generally slower than in broadleaf due to

higher lignin content, C:N ratio, and resinous structures (Cornwell et al., 2009). This might

explain the relatively low sesquiterpene emissions in fungi-depleted bundles. In contrast,

broadleaf wood is more easily colonised due to lower lignin content, allowing environmental

fungi to rapidly colonise fungi-depleted bundles. This likely led to similar levels of fungal

activity and sesquiterpene emissions in both inoculated and fungi-depleted bundles.

4.3 VOCs and beetle responses to burned wood

The burning process alters wood's structure, texture, and chemical properties, releasing odour

cues that are ecologically relevant to pyrophilous beetles across local and landscape scales

(Allison et al., 2004; Ramberg et al., 2025). Some of these VOCs (e.g., guaiacol, 5-

methylfurfural) act as chemical cues of burned wood for pyrophilous species of longhorn and

woodboring beetles. However, on burned bundles, we detected neither pyrophilous beetles

nor VOCs typically associated with thermal degradation of cellulose and lignin (Oasmaa et

al., 2003; Karagöz et al., 2005), likely due to the two-month delay between the burning event

and VOC sampling. Additionally, the amount of burned wood in our experiment was limited

and highly fragmented, which may have further reduced its detectability by pyrophilous

beetles that rely on long-range chemical cues to locate large-scale fires (Paczkowski et al.,

2013) rather than isolated burned logs. In contrast, we detected *Cryptophagus scanicus*, a

silken fungus beetle, which associated with burned bundles and an unidentified hydrocarbon

(Fig. 4b). This finding suggests a potential case of priority effects, where early fungal

colonisers during the degradation of deadwood may have produced volatiles that attracted

this species.

4.4 VOC beetle association

In the present study, we found that saproxylic beetle communities were associated with VOC

profiles of their host trees. Beetle assemblages from spruce and pine occurred in deadwood

with higher relative abundances of conifer-associated volatiles (i.e., α-pinene, verbenol),

while assemblages from beech and oak were associated with broadleaf-associated volatiles

(i.e., sativene, β-copaene) (Fig. 4a). Moreover, the composition of VOC profiles differed

markedly between coniferous and broadleaf species: broadleaf bundles emitted a greater

diversity of compound classes, predominantly sesquiterpenes, whereas monoterpenes

dominated conifer bundles. The distinct separation of beetle communities based on host tree

volatiles is particularly striking given the complexity of natural environments, where

overlapping odours create a complex mixture of ambient chemical cues. These findings

suggest that saproxylic beetles can detect and respond to either a single volatile or blends to

locate suitable deadwood for feeding or oviposition.

One distinct cluster of beetles emerging from conifer bundles was dominated by bark

beetles and by the predatory species *Nemozoma elongatum*. This cluster was associated with

high relative abundances of conifer volatiles such as α-pinene (P109), β-pinene (P132),

limonene (P158), verbenone (P243), carene (P148), and terpineol (P229) (Fig. 4a). Several of

these monoterpenes are well-known semiochemicals for bark beetles, involved in host

location and aggregation (Byers, 1989; López et al., 2013; Sánchez-Osorio et al., 2021),

acting as cues. The co-occurrence of *N. elongatum* with bark beetles is likely driven by the

presence of its primary prey, *Pityogenes chalcographus*, whose aggregation pheromone,

chalcogran, is used by *N. elongatum* as a kairomonal cue for prey detection (Heuer & Vité,

1984).

The second cluster, comprised of jewel beetles, longhorn beetles, and the bark beetle

Magdalis violacea, linked to cymeneol (P234), longifolene (P318), caryophyllene oxide

(P380), and an unidentified compound. Longifolene has been suggested as a potential

attractant for *Monochamus galloprovincialis* (Szmigielski et al., 2012), especially when

coupled with other common host volatiles (i.e., limonene, α-pinene) (Pajares et al., 2004). In

our study, the few individuals detected in a single bundle did not allow us to include *M.*

galloprovincialis in our analysis.

Nevertheless, our results suggest that longifolene may serve as a host colonisation cue for

other longhorn beetles. For jewel beetles, these associations are reported here for the first

time, and further experimental studies are needed to determine whether and how these species

detect and respond to these compounds.

A third cluster, including p-cymenene (P187), trans-verbenol (P216), and two unidentified

compounds, was not associated with any beetle species in our dataset. Du et al. (2025)

reported that under intense bark beetle pressure, emissions of p-cymenene and other aromatic

compounds increased in spruce logs before beetle emergence, likely as part of an induced

defence response from the logs. This may suggest a deterrent role for these compounds in

shaping local beetle assemblages.

Among the broadleaf-associated beetles, the ambrosia beetle *Xylosandrus germanus* was

associated with a group of VOCs that comprises 1-ethyl-3-methylbenzene (P123) and three

unidentified sesquiterpenes from oak. This ambrosia beetle is known to respond to VOCs

emitted by its symbiotic fungus *Ambrosiella grosmanniae* and other fungal sources under

bioassay experiments (Mayers et al., 2015; Gugliuzzo et al., 2023). This aligns with our

findings, as we recorded thousands of individuals emerging from both fungi-inoculated and

fungi-depleted bundles (colonised by pioneer fungi), suggesting that it relies on fungal decay

cues to locate and colonise weakened or dead wood in the field.

The second cluster, consisting of a broad range of beetles, including two beech bark

beetles, two oak-associated longhorns *Synchita mediolanensis* (Zopheridae) and *Crypturgus*

dentatus (Cryptophagidae), and *Litargus connexus* (Mycetophagidae), was linked to a

monoterpene emitted by both oak and beech and a sesquiterpene associated with beech.

While their host preferences are consistent with our results, the chemical cues involved in

host localisation remain largely unknown. In our study, broadleaf-associated beetles were

generally linked to sesquiterpenes (Fig. 4a), a pattern consistent with previous findings (Fäldt

et al., 1999; McLeod et al., 2005; Leather et al., 2014). Interestingly, the beech bark beetles,

Taphrorychus bicolor and *Ernoporicus fagi*, were associated with an unidentified

sesquiterpene predominantly emitted from fungi-colonised beech bundles. This association

may indicate the use of specific chemical cues to locate early-decay beech substrates.

The third cluster included unidentified sesquiterpenes associated with broadleaf and

beech. In our study, this cluster was not linked to any of the beetle species sampled, although

these compounds may serve as cues for other species that were either absent or

underrepresented in our dataset.

The fourth cluster had a composition similar to the third, consisting of unidentified

sesquiterpenes, but was linked to oak- or broadleaf-associated beetles from several families.

These compounds may function as general olfactory cues for saproxylic beetles across

different families when locating broadleaf deadwood. The only tentatively identified

compounds were sativene (P315) and β-copaene (P326), known to be emitted by the brown-

rot fungus *Fomitopsis pinicola* (Rösecke et al., 2000), which was used to inoculate a subset of

bundles. Although these compounds were more abundant on beech deadwood, we observed a

non-significant trend toward higher abundance in brown-rot inoculated bundles compared to

white-rot. We suggest that small amounts of these volatiles may serve as host colonisation

cues by *Bitoma crenata*, a cylindrical bark beetle associated with brown-rot bundles in our

study, suggesting a potential role of priority effects, where early fungal colonisers shape

subsequent beetle assemblages through the emission of chemical cues linked to deadwood

degradation. Although experimental validation is needed to confirm whether these

compounds are detected and behaviourally relevant to beetles, considering aspects like blend

composition and concentration, our findings highlight the ecological relevance of VOCs in

the colonisation of deadwood by saproxylic beetles.

5. CONCLUSIONS

This study is among the first that highlight the role of VOCs in shaping saproxylic beetle

communities during the early stage of deadwood decay. We found that beetle assemblages

are associated with VOC profiles characteristic of their host tree (conifer or broadleaf), with

different groups of volatiles affecting taxa such as bark beetles (Scolytinae), longhorn beetles (Cerambycidae), and jewel beetles (Buprestidae). For some species, the altered VOC profile of fungus-colonised deadwood may have influenced colonisation, suggesting a VOC-mediated priority effect. By focusing on the early stage of decay, our study primarily captured pioneering colonisers, which are typically highly abundant and host-tree specialists, such as bark beetles (Bussler et al., 2011; Seibold et al., 2023). This aligns with our findings of strong host-tree preferences and a dominance of bark beetles. The limited effect of fungal cues on beetle communities may reflect the early successional stage, where chemical cues from host trees play a greater role than those from fungal colonisation, which becomes increasingly influential in later decay stages through fungal–beetle interactions and priority effects. Although this study focused on beetles, our findings suggest that VOCs may mediate the colonisation processes of other saproxylic organisms (e.g., termites), an aspect worth considering in forest management. Finally, behavioural assays, in the laboratory and the field, are needed to determine whether the identified compounds function as semiochemicals for saproxylic beetles.

Acknowledgements

This study was supported by the Czech Science Foundation (GACR, project no. 22-27166S). We thank Sven Finnberg for allowing access to part of his forest and for invaluable field assistance. We are also grateful to Jonas Hagge, Max Hetzer, and Michal Perlík for their help during fieldwork, and Lena Unterbauer for managing most of the volatile samples. Special thanks to David Hauck, Jiří Procházka, Pavel Průdek, and František Houška for assistance with beetle identification, and to Michaela Helclová for help with sample sorting and for the beetle illustrations. We appreciate the cooperation and support of Michal Plátek. Thanks to

Matthew Sweeney for proofreading the manuscript. We thank the administrations of Podyjí

National Park, Pálava PLA, and the Stará Obora game reserve for permitting our research.

Conflict of interest

The authors declare no conflict of interest.

Data Availability Statement

The data that support the findings of this study are available for peer review in the Figshare
repository at https://figshare.com/s/680e996049ac87919a90 . The dataset will be made
publicly available upon publication.
480

481   **Author Contributions**
482
Conceptualization: Simon Thorn, Lukas Drag, Thomas Schmitt
Methodology: Simon Thorn, Lukas Drag
Formal analysis: Claudio Sbaraglia, Simon Thorn
Investigation: Claudio Sbaraglia, Simon Thorn, Lukas Drag, Lucie Ambrozova, Petr Kozel,
Lukas Cizek
Data curation: Claudio Sbaraglia, Daniel Rodriguez
Writing – original draft: Claudio Sbaraglia
Writing – review & editing: Simon Thorn, Lukas Drag, Lucie Ambrozova, Petr Kozel,
Daniel Rodriguez, Thomas Schmitt, Lukas Cizek
Visualization: Claudio Sbaraglia, Petr Kozel
Supervision: Lukas Drag, Simon Thorn, Thomas Schmitt
Funding acquisition: Simon Thorn, Lukas Drag

References

Allison, J. D., Borden, J. H., & Seybold, S. J. (2004). A review of the chemical ecology of the

Cerambycidae (Coleoptera). *CHEMOECOLOGY*, *14*(3), 123–150.

https://doi.org/10.1007/s00049-004-0277-1

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting Linear Mixed-Effects

Models Using lme4. *Journal of Statistical Software*, *67*, 1–48.

https://doi.org/10.18637/jss.v067.i01

Birkemoe, T., Jacobsen, R. M., Sverdrup-Thygeson, A., & Biedermann, P. H. W. (2018).

Insect-Fungus Interactions in Dead Wood Systems. In M. D. Ulyshen (Ed.),

Saproxylic Insects (Vol. 1, pp. 377–427). Springer International Publishing.

https://doi.org/10.1007/978-3-319-75937-1_12

Bruce, T. J. A., Wadhams, L. J., & Woodcock, C. M. (2005). Insect host location: A volatile

situation. *Trends in Plant Science*, *10*(6), 269–274.

https://doi.org/10.1016/j.tplants.2005.04.003

Bussler, H., Bouget, C., Brustel, H., Brändle, M., Riedinger, V., Brandl, R., & Müller, J.

(2011). Abundance and pest classification of scolytid species (Coleoptera:

Curculionidae, Scolytinae) follow different patterns. *Forest Ecology and*

Management, *262*(9), 1887–1894. https://doi.org/10.1016/j.foreco.2011.08.011

Byers, J. A. (1989). Chemical ecology of bark beetles. *Experientia*, *45*(3), 271–283.

https://doi.org/10.1007/BF01951813

Cáceres, M. D., & Legendre, P. (2009). Associations between species and groups of sites:

Indices and statistical inference. *Ecology*, *90*(12), 3566–3574.

https://doi.org/10.1890/08-1823.1

Cavaletto, G., Faccoli, M., Marini, L., Spaethe, J., Magnani, G., & Rassati, D. (2020). Effect

of Trap Color on Captures of Bark- and Wood-Boring Beetles (Coleoptera;

Buprestidae and Scolytinae) and Associated Predators. *Insects*, *11*(11), Article 11.

https://doi.org/10.3390/insects11110749

Celedon, J. M., & Bohlmann, J. (2019). Oleoresin defenses in conifers: Chemical diversity,

terpene synthases and limitations of oleoresin defense under climate change. *New*

Phytologist, *224*(4), 1444–1463. https://doi.org/10.1111/nph.15984

Cornwell, W. K., Cornelissen, J. H. C., Allison, S. D., Bauhus, J., Eggleton, P., Preston, C.

M., Scarff, F., Weedon, J. T., Wirth, C., & Zanne, A. E. (2009). Plant traits and wood

fates across the globe: Rotted, burned, or consumed? *Global Change Biology*, *15*(10),

2431–2449. https://doi.org/10.1111/j.1365-2486.2009.01916.x

Du, B., Frühbrodt, T., Delb, H., Burzlaff, T., Biedermann, P. H. W., & Kreuzwieser, J.

(2025). Emission patterns of volatile organic compounds from Norway spruce logs

following bark beetle (Ips typographus L.) infestation. *Tree Physiology*, *45*(1),

tpae152. https://doi.org/10.1093/treephys/tpae152

Dübeler, A., Voltmer, G., Gora, V., Lunderstädt, J., & Zeeck, A. (1997). Phenols from *Fagus*

sylvatica and their role in defence against *Cryptococcus fagisuga*. *Phytochemistry*,

45(1), 51–57. https://doi.org/10.1016/S0031-9422(96)00771-6

Dufrêne, M., & Legendre, P. (1997). Species Assemblages and Indicator Species: The Need

for a Flexible Asymmetrical Approach. *Ecological Monographs*, *67*(3), 345–366.

https://doi.org/10.1890/0012-9615(1997)067[0345:SAAIST]2.0.CO;2

El Ariebi, N., Hiscox, J., Scriven, S. A., Müller, C. T., & Boddy, L. (2016). Production and

effects of volatile organic compounds during interspecific interactions. *Fungal*

Ecology, *20*, 144–154. https://doi.org/10.1016/j.funeco.2015.12.013

Evans, J. A., Eyre, C. A., Rogers, H. J., Boddy, L., & Müller, C. T. (2008). Changes in

volatile production during interspecific interactions between four wood rotting fungi

growing in artificial media. *Fungal Ecology*, *1*(2), 57–68.

https://doi.org/10.1016/j.funeco.2008.06.001

Fäldt, J., Jonsell, M., Nordlander, G., & Borg-Karlson, A.-K. (1999). Volatiles of Bracket

Fungi *Fomitopsis pinicola* and *Fomes fomentarius* and Their Functions as Insect

Attractants. *Journal of Chemical Ecology*, *25*(3), 567–590.

https://doi.org/10.1023/A:1020958005023

Fraenkel, G. S. (1959). The Raison d'Être of Secondary Plant Substances. *Science*,

129(3361), 1466–1470. https://doi.org/10.1126/science.129.3361.1466

Franceschi, V. R., Krokene, P., Christiansen, E., & Krekling, T. (2005). Anatomical and

chemical defenses of conifer bark against bark beetles and other pests. *New*

Phytologist, *167*(2), 353–376. https://doi.org/10.1111/j.1469-8137.2005.01436.x

Fraser, A. M., Mechaber, W. L., & Hildebrand, J. G. (2003). Electroantennographic and

Behavioral Responses of the Sphinx Moth *Manduca sexta* to Host Plant Headspace

Volatiles. *Journal of Chemical Ecology*, *29*(8), 1813–1833.

https://doi.org/10.1023/A:1024898127549

Fukasawa, Y. (2021). Ecological impacts of fungal wood decay types: A review of current

knowledge and future research directions. *Ecological Research*, *36*(6), 910–931.

https://doi.org/10.1111/1440-1703.12260

Grove, S. J. (2002). Saproxylic Insect Ecology and the Sustainable Management of Forests.

Annual Review of Ecology, Evolution, and Systematics, *33*(Volume 33, 2002), 1–23.

https://doi.org/10.1146/annurev.ecolsys.33.010802.150507

Gugliuzzo, A., Kreuzwieser, J., Ranger, C. M., Tropea Garzia, G., Biondi, A., & Biedermann,

P. H. W. (2023). Volatiles of fungal cultivars act as cues for host-selection in the

fungus-farming ambrosia beetle *Xylosandrus germanus*. *Frontiers in Microbiology*,

14, 1151078. https://doi.org/10.3389/fmicb.2023.1151078

Hartikainen, A., Yli-Pirilä, P., Tiitta, P., Leskinen, A., Kortelainen, M., Orasche, J., Schnelle-

Kreis, J., Lehtinen, K. E. J., Zimmermann, R., Jokiniemi, J., & Sippula, O. (2018).

Volatile Organic Compounds from Logwood Combustion: Emissions and

Transformation under Dark and Photochemical Aging Conditions in a Smog

Chamber. *Environmental Science & Technology*, *52*(8), 4979–4988.

https://doi.org/10.1021/acs.est.7b06269

Heuer, H. G., & Vité, J. P. (1984). Chalcogran: Unique kairomone-governed predator-prey

relations among ostomid and solytid beetles. *Naturwissenschaften*, *71*(4), 214–215.

https://doi.org/10.1007/BF00490438

Holighaus, G., & Schütz, S. (2006). Odours of wood decay as semiochemicals for

Trypodendron domesticum L. (Col., Scolytidae).

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous Inference in General Parametric

Models. *Biometrical Journal*, *50*(3), 346–363.

https://doi.org/10.1002/bimj.200810425

Hulcr, J., Mann, R., & Stelinski, L. L. (2011). The Scent of a Partner: Ambrosia Beetles Are

Attracted to Volatiles from Their Fungal Symbionts. *Journal of Chemical Ecology*,

37(12), 1374–1377. https://doi.org/10.1007/s10886-011-0046-x

Isidorov, V. A., Smolewska, M., Purzyńska-Pugacewicz, A., & Tyszkiewicz, Z. (2010).

Chemical composition of volatile and extractive compounds of pine and spruce leaf

litter in the initial stages of decomposition. *Biogeosciences*, *7*(9), 2785–2794.

https://doi.org/10.5194/bg-7-2785-2010

Johansson, T., Olsson, J., Hjältén, J., Jonsson, B. G., & Ericson, L. (2006). Beetle attraction

to sporocarps and wood infected with mycelia of decay fungi in old-growth spruce

forests of northern Sweden. *Forest Ecology and Management*, *237*(1–3), 335–341.

https://doi.org/10.1016/j.foreco.2006.09.056

Jonsell, M., & Nordlander, G. (1995). Field attraction of Coleoptera to odours of the wood-

decaying polypores *Fomitopsis pinicola* and *Fomes fomentarius*. *Annales Zoologici*

Fennici, *32*(4), 391–402. https://www.jstor.org/stable/23735545

Karagöz, S., Bhaskar, T., Muto, A., & Sakata, Y. (2005). Comparative studies of oil

compositions produced from sawdust, rice husk, lignin and cellulose by hydrothermal

treatment. *Fuel*, *84*(7), 875–884. https://doi.org/10.1016/j.fuel.2005.01.004

Kupferschmid Albisetti, A. D., Brang, P., Schönenberger, W., & Bugmann, H. (2003). Decay

of *Picea abies* snag stands on steep mountain slopes. *The Forestry Chronicle*, *79*(2),

247–252. https://doi.org/10.5558/tfc79247-2

Leather, S. R., Baumgart, E. A., Evans, H. F., & Quicke, D. L. J. (2014). Seeing the trees for

the wood – beech (*Fagus sylvatica*) decay fungal volatiles influence the structure of

saproxylic beetle communities. *Insect Conservation and Diversity*, *7*(4), 314–326.

https://doi.org/10.1111/icad.12055

López, S., Quero, C., Iturrondobeitia, J. C., Guerrero, Á., & Goldarazena, A. (2013).

Electrophysiological and behavioural responses of *Pityophthorus pubescens*

(Coleoptera: Scolytinae) to ()-α-farnesene, ()-(+)-limonene and ()-(−)-verbenone in

(Pinaceae) stands in northern Spain. *Pest Management Science*, *69*(1), 40–47.

https://doi.org/10.1002/ps.3359

Mäki, M., Mali, T., Hellén, H., Heinonsalo, J., Lundell, T., & Bäck, J. (2021). Deadwood

substrate and species-species interactions determine the release of volatile organic

compounds by wood-decaying fungi. *Fungal Ecology*, *54*, 101106.

https://doi.org/10.1016/j.funeco.2021.101106

Mali, T., Mäki, M., Hellén, H., Heinonsalo, J., Bäck, J., & Lundell, T. (2019). Decomposition

of spruce wood and release of volatile organic compounds depend on decay type,

fungal interactions and enzyme production patterns. *FEMS Microbiology Ecology*,

95(9), fiz135. https://doi.org/10.1093/femsec/fiz135

Mämmelä, P. (2001). Phenolics in selected European hardwood species by liquid

chromatography–electrospray ionisation mass spectrometry. *Analyst*, *126*(9), 1535–

1538. https://doi.org/10.1039/B104584A

Mayers, C. G., McNew, D. L., Harrington, T. C., Roeper, R. A., Fraedrich, S. W.,

Biedermann, P. H. W., Castrillo, L. A., & Reed, S. E. (2015). Three genera in the

Ceratocystidaceae are the respective symbionts of three independent lineages of

ambrosia beetles with large, complex mycangia. *Fungal Biology*, *119*(11), 1075–

1092. https://doi.org/10.1016/j.funbio.2015.08.002

McLeod, G., Gries, R., von Reuß, S. H., Rahe, J. E., McIntosh, R., König, W. A., & Gries, G.

(2005). The pathogen causing Dutch elm disease makes host trees attract insect

vectors. *Proceedings of the Royal Society B: Biological Sciences*, *272*(1580), 2499–

2503. https://doi.org/10.1098/rspb.2005.3202

Miller, D. R. (2006). Ethanol and (−)-α-Pinene: Attractant Kairomones for Some Large

Wood-Boring Beetles in Southeastern USA. *Journal of Chemical Ecology*, *32*(4),

779–794. https://doi.org/10.1007/s10886-006-9037-8

Nottingham, S. F., Hardie, J., Dawson, G. W., Hick, A. J., Pickett, J. A., Wadhams, L. J., &

Woodcock, C. M. (1991). Behavioral and electrophysiological responses of Aphids to

host and nonhost plant volatiles. *Journal of Chemical Ecology*, *17*(6), 1231–1242.

https://doi.org/10.1007/BF01402946

Oasmaa, A., Kuoppala, E., & Solantausta, Y. (2003). Fast Pyrolysis of Forestry Residue. 2.

Physicochemical Composition of Product Liquid. *Energy & Fuels*, *17*(2), 433–443.

https://doi.org/10.1021/ef020206g

Oksanen, J., Simpson, G. L., Blanchet, F. G., Kindt, R., Legendre, P., Minchin, P. R.,

O'Hara, R. B., Solymos, P., Stevens, M. H. H., Szoecs, E., Wagner, H., Barbour, M.,

Bedward, M., Bolker, B., Borcard, D., Carvalho, G., Chirico, M., De Caceres, M.,

Durand, S., … Weedon, J. (2001). *vegan: Community Ecology Package* (p. 2.6-8)

[Dataset]. https://doi.org/10.32614/CRAN.package.vegan

Orban, A., Schwab, L. J., Lugones, L. G., Wösten, H. A. B., & Rühl, M. (2023). Effect of

fruiting-related genes on the formation of volatile organic compounds in the

mushroom *Schizophyllum commune*. *Mycological Progress*, *22*(9), 67.

https://doi.org/10.1007/s11557-023-01914-4

Paczkowski, S., Paczkowska, M., Dippel, S., Schulze, N., Schütz, S., Sauerwald, T., Weiß,

A., Bauer, M., Gottschald, J., & Kohl, C.-D. (2013). The olfaction of a fire beetle

leads to new concepts for early fire warning systems. *Sensors and Actuators B:*

Chemical, *183*, 273–282. https://doi.org/10.1016/j.snb.2013.03.123

Paiva, N. L. (2000). An Introduction to the Biosynthesis of Chemicals Used in Plant-Microbe

Communication. *Journal of Plant Growth Regulation*, *19*(2), 131–143.

https://doi.org/10.1007/s003440000016

Pajares, J. A., Ibeas, F., Díez, J. J., & Gallego, D. (2004). Attractive responses by

Monochamus galloprovincialis (Col., Cerambycidae) to host and bark beetle

semiochemicals. *Journal of Applied Entomology*, *128*(9–10), 633–638.

https://doi.org/10.1111/j.1439-0418.2004.00899.x

Pichler, V., Homolák, M., Skierucha, W., Pichlerová, M., Ramírez, D., Gregor, J., & Jaloviar,

P. (2012). Variability of moisture in coarse woody debris from several ecologically

important tree species of the Temperate Zone of Europe. *Ecohydrology*, *5*(4), 424–

434. https://doi.org/10.1002/eco.235

Raffa, K. F. (2014). Terpenes tell different tales at different scales: Glimpses into the

Chemical Ecology of conifer - bark beetle - microbial interactions. *Journal of*

Chemical Ecology, *40*(1), 1–20. https://doi.org/10.1007/s10886-013-0368-y

Ramberg, E., Strengbom, J., Wikars, L.-O., & Ranius, T. (2025). Surrounding landscape

composition influences saproxylic beetle assemblages after prescribed burning.

Journal of Applied Ecology, *n/a*(n/a). https://doi.org/10.1111/1365-2664.70084

Rösecke, J., Pietsch, M., & König, W. A. (2000). Volatile constituents of wood-rotting

basidiomycetes. *Phytochemistry*, *54*(8), 747–750. https://doi.org/10.1016/S0031-

9422(00)00138-2

Sánchez-Osorio, I., Tapias, R., Domínguez, L., López-Pantoja, G., & González, M. del M.

(2021). Electroantennographic Responses of *Cerambyx welensii* Küster to Host-

Related Volatiles. *Forests*, *12*(9), Article 9. https://doi.org/10.3390/f12091168

Seibold, S., Brandl, R., Buse, J., Hothorn, T., Schmidl, J., Thorn, S., & Müller, J. (2015).

Association of extinction risk of saproxylic beetles with ecological degradation of

forests in Europe. *Conservation Biology*, *29*(2), 382–390.

https://doi.org/10.1111/cobi.12427

Seibold, S., Müller, J., Allner, S., Willner, M., Baldrian, P., Ulyshen, M. D., Brandl, R.,

Bässler, C., Hagge, J., & Mitesser, O. (2022). Quantifying wood decomposition by

insects and fungi using computed tomography scanning and machine learning.

Scientific Reports, *12*(1), 16150. https://doi.org/10.1038/s41598-022-20377-3

Seibold, S., Weisser, W. W., Ambarlı, D., Gossner, M. M., Mori, A. S., Cadotte, M. W.,

Hagge, J., Bässler, C., & Thorn, S. (2023). Drivers of community assembly change

during succession in wood□decomposing beetle communities. *Journal of Animal*

Ecology, *92*(5), 965–978. https://doi.org/10.1111/1365-2656.13843

Seybold, S. J., Huber, D. P. W., Lee, J. C., Graves, A. D., & Bohlmann, J. (2006). Pine

monoterpenes and pine bark beetles: A marriage of convenience for defense and

chemical communication. *Phytochemistry Reviews*, *5*(1), 143–178.

https://doi.org/10.1007/s11101-006-9002-8

Suckling, D. M., Gibb, A. R., Daly, J. M., Chen, X., & Brockerhoff, E. G. (2001). Behavioral

and Electrophysiological Responses of Arhopalus tristis to Burnt Pine and Other

Stimuli. *Journal of Chemical Ecology*, *27*(6), 1091–1104.

https://doi.org/10.1023/A:1010355710509

Szmigielski, R., Cieslak, M., Rudziński, K. J., & Maciejewska, B. (2012). Identification of

volatiles from *Pinus silvestris* attractive for *Monochamus galloprovincialis* using a

SPME-GC/MS platform. *Environmental Science and Pollution Research*

International, *19*(7), 2860–2869. https://doi.org/10.1007/s11356-012-0792-5

Ulyshen, M., & Hanula, J. (2010). Patterns of saproxylic beetle succession in loblolly pine.

Agricultural and Forest Entomology 12: 187-194, *12*, 187–194.

https://research.fs.usda.gov/treesearch/34896

Visser, J. H., & Avé, D. A. (1978). General Green Leaf Volatiles in the Olfactory Orientation

of the Colorado Beetle, *Leptinotarsa Decemlineata*. *Entomologia Experimentalis et*

Applicata, *24*(3), 738–749. https://doi.org/10.1111/j.1570-7458.1978.tb02838.x

Vuts, J., Woodcock, C. M., Sumner, M. E., Caulfield, J. C., Reed, K., Inward, D. J., Leather,

S. R., Pickett, J. A., Birkett, M. A., & Denman, S. (2016). Responses of the two-

spotted oak buprestid, *Agrilus biguttatus* (Coleoptera: Buprestidae), to host tree

volatiles. *Pest Management Science*, *72*(4), 845–851. https://doi.org/10.1002/ps.4208

Wickham, H. (2016). Data Analysis. In H. Wickham, *Ggplot2* (pp. 189–201). Springer

International Publishing. https://doi.org/10.1007/978-3-319-24277-4_9

Figures

Fig. 1 Ten study sites distributed across Germany and the Czech Republic (a). Each site consisted of two plots (shady, sun-exposed), each with 20 bundles (combinations of four tree species and five treatments) (b). VOCs were collected from each bundle during July 2022 after the bundles had been wrapped for 45 minutes to concentrate the volatiles, followed by a 15-minute collection period (c).

Fig. 2 Canonical Correspondence Analysis (CCA) of VOC composition from deadwood bundles. The first two axes explain significantly (*** = $p<0.005$) 39.5% of the variation. Model-selected peaks (n = 89) were used as response variables; predictors included tree species, treatment, their interaction, and exposure. Bar plots show the variance explained by each predictor. Colours indicate tree species; shapes indicate treatments.

Fig. 3 The heatmap showing the average relative VOC composition across tree species and treatment combinations. Peaks are colour-coded by chemical class, with a gradient indicating the level of VOC emissions. Darker colours represent higher emissions, while lighter colours correspond to lower emissions. Grey indicates missing VOCs.

Fig. 4 NMDS plots showing associations between VOC peaks and saproxylic beetles by (a) tree species and (b) treatments. Dots represent indicator beetles (n = 40), colour-coded by substrate preference. Grey dots and arrows stand for no preferences from the LMM. Arrows indicate significant VOC peaks (n = 42); their direction indicates positive associations with beetle species (dots) positioned in front of them. Only VOC peaks identified at the substance level are labelled; the full list is available in Appendix S3. Arrow lengths are standardised to unit vectors, not showing effect size (R^2). The same colour was used for beetles and VOCs associated with the respective tree species (a) or treatment (b). "Manipulated" includes beetles attracted to all treatments except the control. Beetle illustrations by Michaela Helclová.

a)

b) Shady

Sun-exposed

Control
Burned
Sterilized
White rot
Brown rot

c)