Heterotypic intercellular adhesion tunes efficiency of cell-on-cell migration

Chandrashekar Kuyyamudi[1,2]*#, Suhrid Ghosh[1,2]#, & Cassandra G. Extavour[1,2,3]*

1. Department of Organismic & Organismic Biology, Harvard University. Cambridge, MA, USA
2. Howard Hughes Medical Institute. Chevy Chase, MD, USA
3. Department of Molecular & Cellular Biology, Harvard University. Cambridge, MA, USA

*Correspondence: CK   ckuyyamudiashwinikumar@fas.harvard.edu

                     CGE   extavour@oeb.harvard.edu

# These authors contributed equally.

Significance Statement

Cell adhesion is important for cell migration, and when cells migrate on a substrate of other cells (rather than on extracellular matrix), the adhesive properties of both cell types must be considered. However, whether and how dynamic changes in adhesion regulate cell-on-cell migration remains unclear. Here we approach this problem using transepithelial migration of *Drosophila* embryonic germ cells as a case study. We develop an *in silico* model of the migration process that predicts an optimal level of adhesion between migrating cell and substrate cell to achieve efficient migration. *In vivo* live imaging and genetic manipulation experiments uphold the predictions of this model. This suggests that adhesion is not a simple on/off binary parameter regulating migration.

Abstract

Cell migration across epithelial barriers occurs in diverse developmental, immunological, and pathological contexts. Here we investigate the contribution of heterotypic adhesion between migrating cells and epithelial "substrate" cells to transepithelial migration. Using an *in silico* model inspired by the migration of primordial germ cells across the midgut epithelium in the *Drosophila* embryo, we show that heterotypic adhesion modulates migration efficiency in a non-monotonic manner, revealing the existence of an optimal adhesion regime. Consistent with this prediction, *in vivo* overexpression of E-cadherin in germ cells accelerated their exit from the midgut relative to controls. Beyond providing experimentally testable predictions, our model integrates and explains previous observations on the role of heterotypic adhesion in cell-on-cell migration, offering a framework for understanding transepithelial migration across biological contexts.

Keywords:

Germ cell migration, intercellular adhesion, E-cadherin, Cellular Potts model, CompuCell3D, *in vivo* live imaging

Introduction

Cells frequently carry out their functions at locations distinct from their site of origin, making the individual cellular motility that is required for cell migration a critical aspect of their behavior (1, 2). To achieve motility, cells require persistent and directed membrane activity, which is frequently guided by external chemical cues (3–6). This directed membrane activity is enabled by the dynamic nature of cell membranes, which exhibit continuous turnover of cytoskeletal components (7–9)

Cell migration, division, differentiation and death enable the development of multicellular organisms from a single cell (10). Cells in various developmental contexts migrate individually as well as collectively (1, 2, 11). Embryonic development unfolds as a tightly regulated cascade of events, where each step must occur within a specific time window to prevent disruption of subsequent stages (12–15). These temporal constraints also apply to cell migration, making not only the path taken but also the timing of arrival critical to proper development (1, 2, 16, 17).

In the developing embryo of the fruit fly *Drosophila melanogaster*, the primordial germ cells (germ cells) are the first cells to form (18). The germ cells form at the posterior tip of the embryo roughly 2.5 hours after egg laying (18, 19). Around 15 minutes after their formation, the embryo undergoes gastrulation, and the cluster of germ cells enters the primordial midgut upon invagination of the surrounding tissue during gastrulation (20). Here, the germ cell cluster eventually disassembles, and the germ cells exit the gut by individually passing through the monolayer of epithelial cells that form the midgut (20, 21). Recent work has implicated juvenile hormones as chemical cues that guide germ cells toward the posteriorly located gonad (22).

Complementing this, Wunen/Wunen2, strongly expressed on the ventral side of the

midgut, acts as a repellent that drives germ cells to exit the midgut dorsally (23–26).

Once outside the midgut, the germ cells continue their migration to join the

mesodermal cells that form the somatic gonad primordium (27).

The *Drosophila* adhesion protein E-cadherin is encoded by the *shotgun* (*shg*) gene,

and is maternally deposited in the early embryo (28, 29). Germ cells require this

maternally deposited E-cadherin both to form a cohesive cluster and to tether to the

invaginating midgut primordium during gastrulation (30–32). Regulation of germ cell

E-cadherin during internalization into the midgut is orchestrated by the Trapped in

endoderm (Tre1) protein, a G protein-coupled receptor (31, 32). Tre1 enables the germ

cell cluster to form a radial configuration of polarized cells that detach from the cluster

prior to transepithelial migration (31, 32). In embryos laid by females expressing a loss

of function *shotgun* allele (shg^{A9-49}), individual germ cells separate from the cluster

prematurely (31). Furthermore, in 38% of shg^{A9-49} mutant embryos, germ cells exhibit a

delay in crossing the midgut epithelium (31). The posterior midgut, which is the barrier

that the germ cells must breach, is made up of epithelial cells (33, 34). Remodeling of the

midgut epithelium endows the epithelial barrier with increased permeability, allowing

the germ cells to migrate through it (20, 35). High resolution imaging of the germ cells

crossing the midgut epithelium has shown that germ cells maintain contacts with the

epithelial cells with no evident intercellular space around them (20). In addition, FGF

signaling regulates the distribution of zygotic E-cadherin in the posterior midgut to

maintain its epithelial integrity (36). FGF mutants display mislocalized E-cadherin and

collapse of the midgut lumen, trapping the germ cells within (36).

Germ cells expressing defective E-cadherin (shg^{A9-49}) show delayed

transepithelial migration (31), reminiscent of the delay in border cell migration that

occurs when nurse cells either lack or overexpress E-cadherin in the *D. melanogaster* egg

chamber (37). Together, these observations suggest the existence of a non-monotonic

relationship between migration efficiency and adhesion between migrating and

substrate cells. We set out to understand the role of heterotypic adhesion between the

migrating cells and "substrate" cells in transepithelial migration, taking as a case study

the role of intercellular adhesion mediated by E-cadherin in enabling the germ cells to

transmigrate. Leveraging genetic tools and two-photon microscopy, we live imaged the

process of transepithelial migration *in vivo.* We combined the insights gained from live

imaging with previously published observations to formulate an *in silico* model of the

process. This model, formulated in the Cellular Potts Model framework (38–40),

provides a novel mechanistic understanding of this transepithelial migration process.

Our model reveals the existence of optimal heterotypic adhesion between germ cells

and the epithelial cells, which maximizes the efficiency of midgut exit. In addition, our

in silico model yields testable, potentially generalizable predictions about the role of E-

cadherin and intercellular adhesion that may be applicable to other systems of

heterotypic cell adhesion and migration.

Results

I. Germ cells exit the midgut individually and maintain E-cadherin expression throughout exit

To gain insight into the process of germ cell transepithelial migration, we live imaged the process *in vivo* [**Fig. 1b-b″**]. Before migration, the germ cells are contained by the epithelial barrier of the midgut [**Fig. 1b**], and located at a depth of more than $25 \mu m$ from the cortex within the embryo (41). To visualize the germ cells, we used the previously generated fly line *nos-LifeAct-tdTomato* (42) which specifically labels germ cell membrane and cortex through posteriorly localized maternally deposited protein. The cells in the gut epithelium also show some residual labelling since the gut arises from the invaginating posterior blastoderm. The residual labelling generates contrast and helps us estimate the position of the gut during transepithelial migration.

We tracked germ cells throughout transepithelial migration, characterizing their progress at a five minute time resolution [**SI Videos 1 - 3**], and quantified the distances of germ cells from the center to the edge of the midgut lumen as a function of time [**Fig. 1 c**]. **Figure 1d-d′** shows the distance covered relative to their initial position (i.e. t = 0) as a function of time for two different embryos. We found that individual germ cells took 30 minutes on average to go through the midgut epithelial barrier [**Fig. 1d-d′; SI Videos 1 - 3**]. Our live imaging of the transepithelial migration process did not reveal any cell divisions in the midgut epithelial cells that could have generated spaces in the epithelial barrier that would allow for easier exit of germ cells, in contrast to the mechanism that regulates macrophage infiltration in *D. melanogaster* (43).

To examine E-cadherin dynamics during the transepithelial migration, we

generated a fly line expressing endogenously GFP tagged Vasa and mScarlet tagged E-

cadherin [**SI Methods**]. Using multi-photon confocal microscopy, we observed that E-

cadherin in germ cells changed its subcellular localization at the start of the

transepithelial migration [**SI Figure 5 a,a',b, b'; SI Video 6**]. We observed E-cadherin in

the epithelia also move laterally upon germ cell entry [**SI Figure 5 c,c'; SI Video 6**].

Imaging individual germ cells at higher time resolution revealed transient E-cadherin

foci form at the interface of germ cell and lateral gut epithelia [**SI Figure 5 d,d'; SI**

Video 7]. These foci appear to anchor the germ cell to the baso-lateral epithelial

membrane as the germ cells moves forward [**SI Video 7**] and exit the gut. Our

observations from live imaging were further bolstered by our ~~We also~~ observation of E-

cadherin localization on the surface of germ cells during the transepithelial migration,

detected by immunostaining for GFP in endogenously GFP-tagged E-cadherin embryos

[**Fig. 1e-e'**]. This result is consistent with previously published experiments (31) that

detected E-cadherin by immunostaining.

We quantified the intensity of E-cadherin signal at apical and lateral junctions

between midgut epithelial cells, as well as at the epithelial-germ cell junctions [**Fig 1e''**].

Our quantification revealed that the E-cadherin concentration at the germ cell-epithelial

cell interface is not significantly different from that observed between epithelial cells on

the lateral side (p= 0.31) [**Fig. 1e''**]. The presence of E-cadherin in germ cells during

transepithelial migration, where germ cells physically contact epithelial cells (20), along

with the delayed midgut exit previously observed in *shg* loss of function mutants (31),

suggests that the expression of E-cadherin in germ cells is functional rather than

incidental.

II. *Time to exit the midgut has a non-monotonic dependence on germ cell E-cadherin level*

Although inspired by germ cell transepithelial migration, our central question is

broader: what role does heterotypic E-cadherin adhesion play in cell migration? To

generalize our question beyond germ cell transepithelial migration, we turned to *in*

silico modeling, retaining only essential features of the phenomenon. We developed a

two-dimensional model of transepithelial migration, where a single motile cell breaches

a ring of epithelial cells under the influence of an external, radial chemoattractant

gradient [**Fig. 2a**]. Our two-dimensional model, formulated as a Cellular Potts Model

(38, 39), enabled us to consider the cells as finite sized objects capable of deformations in

shape. We primarily explored the potential role of chemoattractant strength (λ_M) and

the concentration of E-cadherin in the germ cells (C_G) in the time taken for the germ cells

to exit the epithelial barrier (τ_e). We ran our simulations for a maximum of 10,000

Monte Carlo steps, and saved the position of the germ cell as a function of time for each

independent iteration for a given choice of system parameters. **Figure 2a** shows a

typical snapshot from our CompuCell3D simulation, and **Figure 2b-b''** shows the

modeled germ cell at three different stages of the transepithelial migration. In the CPM

framework, the temperature parameter (T) controls boundary pixel activity, where

higher temperatures increase the likelihood of energy-raising pixel flips. To investigate

how membrane fluctuations at interepithelial junctions affect migration, we modulated

the temperature governing pixel acceptance rates at these boundaries. This allowed us

to study how epithelial remodeling influences transepithelial migration.

Our simulations [**Fig. 2c-c′**] revealed that elevated epithelial temperature

increased tissue permeability to germ cells, resulting in faster exit times (τ_e) and higher

escape probabilities (p_e). This is consistent with prior experimental findings where

failure in epithelial remodeling of the midgut adversely affected germ cell

transepithelial migration (20, 35, 44).

The strength of the modeled chemoattractant λ_M beyond which germ cells

successfully migrated through the barrier was positively correlated with epithelial cell

E-cadherin concentration C_E [**Fig. 3a–a′′**]. We interpret this to mean that higher C_E

values, indicating stronger adhesion between the epithelial cells, offered greater

resistance to the trans-migrating cell. Regardless of the value of C_E, we observed that

the time taken to exit the barrier decreased with increased germ cell E-cadherin

concentration only up to a point, beyond which there was a sharp increase in the time

taken to breach the epithelial barrier [**Fig. 3a–a′′**]. This non-monotonic dependence of τ_e

on C_G suggested that there is an optimal range of heterotypic adhesion between germ

cell and the epithelial cell. Within this range, adhesion aids the transepithelial

migration, but beyond the range, it hinders germ cell movement. The minima of the

curves for different values of λ_M in **Figure 3a–a′′** coincide, indicating that the optimal

adhesion value for transepithelial migration is independent of the chemoattractant

strength. Interestingly, the coefficient of variation in the time to exit τ_e as a function of

C_G has a trend that is inverse of the one exhibited by τ_e [**Fig. 3b**]. Specifically, the

coefficient of variation is highest when the mean is the lowest, which is not due to

standard deviation being independent of C_G. Instead, the standard deviation increases

with C_G while the mean decreases with C_G [**SI Fig. 1**].

To better understand the potential role of heterotypic adhesion in determining

the speed of exit, we focused on the trajectories of the migrating cells at different values

of C_G. We ran our simulations for a maximum duration of 10,000 Monte Carlo steps (45),

and the cells which had not exited by then were considered to have failed to exit. In

Figure 3c, the trajectories from ~1000 independent iterations are shown for $C_G = 4.89$,

$C_E = 10.0$ and $\lambda_M = 7000$. While there are numerous instances of the migrating cell

failing at early stages of the transepithelial migration, we see many instances of germ

cells stuck to the outer edge of the epithelial barrier and failing to complete midgut exit.

At low values of CG, germ cells are stuck closer to the apical side of the midgut,

whereas at high values germ cells get attached to the basal side, with much of their cell

body outside the barrier [**SI Figure 2**]. We considered all the successful trajectories for

$C_E = 10$ and $\lambda_M = 7000$ and normalized the time for each trajectory, such that $\tau = 0$ and

$\tau = 1$ mark the start and end of the transepithelial migration respectively. The time-

normalized mean trajectories are shown in **Figure 3d**, colored according to their values

of C_G. We observed that germ cells with lower E-cadherin concentrations spent much of

their time getting to the middle of the epithelial barrier, while the second half of the

journey, culminating in complete exit from the midgut, was quite rapid. The fraction of

total time spent in getting to the middle dropped with increasing germ cell–epithelial

cell adhesion, with speed through the barrier becoming constant for high values of C_G

(~7.0). To highlight the difference in the fraction of total travel time spent at different

halves of the epithelial barrier, we show the fraction of time spent to cover the second

half of the epithelial barrier (referred to as τ_2) in **Figure 3d'**. τ_2 showed a nonlinear

increase with C_G approaching 0.5 at lager values of C_G. The probability of success in

exiting the epithelial barrier had an inverse non-linear relation to time to exit τ_e

regardless of the C_E, C_G and λ_M, as evident from the scatter plot in **Figure 3e**. The model

also predicts that there is an optimal concentration of E-cadherin in the substrate [**SI**

Figure 4], consistent with the hypothesis that the efficiency of transepithelial migration

is a function of the heterotypic adhesive interface, rather than of the migrating germ

cells alone.

In accordance with previous genetic studies (25, 46, 47), we hypothesized that

Wunen/Wunen2, which is strongly expressed ventrally, repels germ cells and biases

their exit toward the dorsal side. When we incorporated a dorsoventral chemical

gradient alongside the radial gradient, germ cells exited the epithelial barrier more

dorsally, supporting this hypothesis [**SI Figure 3**].

III. Overexpression of E-cadherin in germ cells leads to faster exit through

the midgut

The results of our model suggested that increasing E-cadherin levels in germ cells

would enable them to complete the most time-consuming part of their journey, namely

traversing from the middle of the midgut lumen through the luminal epithelial surface,

more rapidly, thus resulting in a faster net transepithelial migration time. To test this

prediction *in vivo*, we employed the UAS-GAL4 system (48). Maternally deposited E-

cadherin serves as the primary source of E-cadherin for all early embryonic cells,

including both germ cells and somatic cells (29, 31, 49). Notably, even during

transepithelial migration (~3 hours after egg laying), germ cell E-cadherin remains

predominantly of maternal origin, making RNAi-mediated knockdown in germ cells

impractical without destroying embryonic integrity (29, 31, 49).

Given these constraints, to assess the effects of E-cadherin overexpression on

transepithelial migration, we analyzed germ cell progression in embryos of four

different maternal genotypes, collected within 30 minutes from synchronously

ovipositing mothers and aged for five hours, as follows: (I) **WT**: Outcrossed wild-type

control, expressing endogenous levels of E-cadherin in all cells [*nos-Gal4/+*]; (II) **Pan-**

Ecad-OE-A: with ubiquitous E-cadherin overexpression in all cells [*nos>Ecad*]; (III) **Pan-**

Ecad-OE-B: with ubiquitous GFP-tagged E-cadherin overexpression in all cells

[*nos>Ecad-GFP*]; and (IV) **GC-Ecad-OE**: with germ cell-targeted overexpression of

mClover2-tagged E-cadherin [*nos> Ecad-mClover2-nosTCE-pgc 3'UTR*]. To assess the

extent of overexpression achieved, we quantified germ cell E-cadherin relative to

epithelial apical and lateral expression using anti-DE-cadherin immunostaining [**SI**

Figure 7]. The results of this quantification confirmed that use of the ubiquitous driver

increased E-cadherin levels in both germ cells and the somatic epithelium, and that the

germ cell driver increased E-cadherin levels in germ cells but not somatic cells [**SI**

Figure 7]. The position of germ cells relative to the midgut lumen boundary in

representative embryos from each group is shown in **Figure 4a–a'''**. Quantitative

analysis revealed that germ cells in all overexpression lines (II–IV) were located farther

from the lumen center compared to controls (I) [**Fig. 4b**]. Furthermore, both the mean

germ cell distance from the lumen center [**Fig. 4c**] and the fraction of germ cells that had

exited the midgut [f_{exit} in **Fig. 4d**] were higher in overexpression groups, with the most

pronounced effect in the germ cell-targeted overexpression line (IV).

These results suggest that higher levels of E-cadherin in germ cells accelerates

transepithelial migration, consistent with our theoretical model. We favor the

interpretation that wild-type germ cell E-cadherin expression is associated with an

intermediate time to exit [gray bar, **Fig. 4e**]. Under this interpretation, loss of function

shg mutants like shg^{A9-49} (31) would drive the embryo toward the red region shown in

Figure 4e, associated with longer τ_e, while *shg* overexpression would drive it towards

the green region in **Figure 4e** associated with faster germ cell exit.

Our model predicted that elevated epithelial E-cadherin would hinder migration

by increasing resistance to germ cell exit. Germ cells in our experiments overexpressing

E-cadherin in both somatic and germ cells showed reduced germ cell exit times. This

apparent discrepancy can be explained by the hypothesis that wild-type E-cadherin

levels in somatic epithelial cells are already near saturation. Under this interpretation,

finite adhesion junction availability and membrane transport limitations would prevent

additional E-cadherin from forming new junctions at high concentrations (50–53). Thus,

in embryos with E-cadherin overexpression in both germ cells and somatic cells, like the

ones we examined herein (**Fig. 4,** groups II and III), we hypothesize that E-cadherin

levels higher than wild type levels would have a stronger effect on germ cells than on

epithelial cells, effectively mimicking germ cell-specific overexpression.

To further test this, we drove zygotic expression of GFP-tagged E-cadherin

specifically in the gut using the 48Y-GAL4 endodermal driver (54) [**SI Figure 6a, b**]. We

observed only a small, statistically insignificant decrease in the fraction of germ cells

that had exited the gut relative to controls [**SI Figure 6f**]. Because the 48Y-GAL4 driver

drives detectable expression 30 minutes after transepithelial migration has initiated [**SI**

Figure 6b], it could be the case that our experiment changed E-cadherin levels too late

to severely perturb the process. However, this result suggests that overexpression of E-

cadherin in the already E-cadherin-rich epithelium has little effect on transepithelial

migration, in contrast to germ cell-specific overexpression.

295   We next used the 48Y-GAL4 driver to induce gut-specific knockdown of E-

296  cadherin (shg-RNAi) and observed a statistically significant reduction in the fraction of

297  germ cells exiting the midgut compared to wild type controls [**SI Figure 6f**]. This

298  phenotype mirrors what Parés and Ricardo (36) reported for FGF pathway mutants,

299  where mislocalization of epithelial E-cadherin caused a substantial proportion of germ

300  cells to fail to exit the midgut. These observations are consistent with our hypothesis

301  that epithelial E-cadherin, and not just germ cell E-cadherin, is required for successful

302  germ cell transmigration. We note that we cannot rule out the alternative hypothesis

303  that germ cell transepithelial migration is impeded in this condition because of broad

304  defects in gut architecture that might be present since epithelial E-cadherin is integral to

305  the structural integrity of the gut and its lumen (36). However, if the gut epithelium

306  were indeed weaker because of E-cadherin depletion, we would expect more efficient

307  germ cell exit, which is the opposite of what we observed. We therefore favor the

308  hypothesis that E-cadherin expression in the gut epithelium is an important contributor

309  to the heterotypic interactions that are required for germ cell exit from the gut.

310 **Discussion**

311 Our work sheds light on the potential role of heterotypic E-cadherin adhesion during

312 the transepithelial migration of germ cells through the midgut. Further, the predictions

313 of our *in silico* model could in principle apply to any heterotypic adhesive migration

314 process, as we incorporated only a few features specific to germ cell transepithelial

315 migration. Our model logic consists of three key elements, i) a migrating cell that forms

316 transient heterotypic adhesive contacts with a barrier epithelium/endothelium, ii) a

317 chemoattractant gradient providing directed motility, and iii) an optimal adhesion

318 range, where too little adhesion fails to provide the traction needed to breach the

barrier, while too much adhesion impedes forward movement by making junction

disassembly rate-limiting. We believe this logic is general, while the migratory and

substrate cell type, the molecular identity of the adhesion molecules and the source of

the driving force are specific to the system.

Our results suggest a non-monotonic dependence of migration efficiency on the

heterotypic adhesion between the migrating cell and the substrate cells [**Fig. 4d**]. We

posit that traction is essential for cell migration, and that transmembrane proteins like

integrins facilitate traction by anchoring cells to the extracellular matrix, thereby

enabling forward movement (55). Additionally, cadherin-like molecules mediate

homotypic adhesion, allowing cells of the same type to form cohesive tissues such as

epithelia (49, 56). These molecules also participate in heterotypic adhesion when

migrating cells interact with others expressing cadherins (57–59). A notable example is

the transendothelial migration of melanoma cells, where heterotypic adhesion is clearly

observed (57). Melanoma cells form heterotypic N-cadherin mediated adhesion

junctions with the endothelial cells during their migration (57, 60). Based on our model,

we predict a non-monotonic relationship between transmigration efficiency and the

heterotypic adhesion between melanoma cells and endothelial cells. More broadly,

diapedesis - the process by which cells breach endothelial barriers - relies on heterotypic

interactions between transmembrane adhesion proteins (58, 59). The prevalence of

heterotypic adhesion during transepithelial migration suggests that our work may have

broader implications beyond the specific context of *Drosophila* primordial germ cell

migration.

Many conserved features of cell migration are applicable well beyond a specific

biological context. In a migrating cell, actin filaments polymerize at the leading edge

and undergo retrograde flow, functioning much like the rotation of wheels in a car (61–

63). However, the forward motion of a cell depends on its physical coupling to the

substrate, just as the tires of a car grip the road to provide traction (61, 62, 64). This

mechanism is described by the *molecular clutch model* of cell motility (65). Adhesion

molecules such as integrins and cadherins allow cells to bind to surfaces or to other

cells, engaging the molecular clutch (65, 66). Integrin-mediated traction is typically used

when cells migrate over an extracellular matrix, whereas cadherins facilitate cell-on-cell

migration, as seen in the case of border cell migration (37, 66), and in the transepithelial

migration case that we consider here.

Our *in silico* model predicts that an intermediate adhesion strength between

migrating cells and substrate cells optimizes transmigration. Our finding aligns with

previous observations on *D. melanogaster* border cell migration in the egg chamber (37,

67). In the case of border cell migration, a cluster of six to eight border cells, enclosing a

pair of non-motile polar cells, migrates through a "substrate" formed by nurse cells (67).

Of these three cell types, polar cells express the highest levels of E-cadherin, which they

use to anchor to motile border cells, which have the second highest E-cadherin levels

(37). Meanwhile, border cells form transient junctions with nurse cell E-cadherin (which

is at the lowest levels of the three cell types in the system), pulling on nurse cells as they

migrate (37). Notably, E-cadherin knockdown in nurse cells impedes migration more

severely than knockdown in border cells, whereas E-cadherin overexpression in nurse

cells slows migration (37). These observations further support the concept that

intermediate heterotypic adhesion is optimal for transmigration, specifically E-cadherin

enabling the engagement of a molecular clutch (66).

Since the Cellular Potts model framework is rooted in Steinberg's differential

adhesion hypothesis (38, 39, 68), transepithelial migration can be interpreted through

this lens. In this framework, epithelial cells, by virtue of their high E-cadherin

expression, strongly prefer to bind one another, creating a thermodynamic barrier that

the germ cells need to breach to intercalate into the epithelium. Germ cell expression of

E-cadherin allows them to compete directly with these homotypic epithelial junctions.

While germ cells form fewer adhesive contacts than two epithelial cells can form with

each other, their E-cadherin expression is sufficient to make heterotypic binding

energetically accessible. The chemotactic gradient then provides the directional bias that

tips the overall energy balance in favor of heterotypic adhesion. Migration in the

absence of sufficient intercellular adhesion resembles climbing a slippery ladder—

lacking the traction needed for upward movement. Conversely, excessive adhesion is

akin to ascending a ladder coated in glue, where detachment becomes the limiting

factor. Thus, an optimal level of "stickiness" is crucial for efficient migration. Inherited

maternal content is known to influence germ cells' ability to migrate successfully to the

gonad (26). A component of this maternal content is the E-cadherin used during

transepithelial migration is one component of this maternally supplied content. We

speculate that the level of E-cadherin in each germ cell may contribute to germ cell

fitness during their journey to the gonad.

Methods

Fly stocks and maintenance

We generated an endogenous C-terminal *shotgun*–mScarlet (referred to as E-cadherin-mScarlet) fusion using scarless CRISPR/Cas9-mediated homology-directed repair in *Drosophila melanogaster* (69), using homology arms flanking the *shotgun* stop codon. We identified HDR events by 3xP3-DsRed fluorescence and validated correct targeting by PCR [**Supporting Methods**]. Fly lines *w[1118]; P{w[+mC]=GAL4::VP16-nanos.UTR}CG6325[MVD1]* (abbreviated herein as *nanos-gal4-VP16*; stock #4937), *w[1118]; P{w[+mC]=UASp-shg.GFP}5B* (#58445), *y[1] w[*]; TI{TI}shg[GFP]* (#60584), and *w[*]; P{w[+mC]=UASp-shg.R}5* (#58494) were obtained from the Bloomington Drosophila Stock Center (Indiana, U.S.A.). Fly line *w[*]; TI{TI}vas^{EGFP.KI}* (Vasa-GFP; stock #118616;) was obtained from the Kyoto Stock Center (Kyoto, Japan). Fly lines *w; nos-Lifeact-tdTomato-P2A-tdKatushka2-CAAX (Lifeact-tdTomato* landing site VK00027; abbreviated herein as *nos-LifeAct-tdTomato*) and *w; UASp-DE-cadherin-mClover2-nosTCE-pgc 3'UTR* (landing site attP2) *(Lin et al., 2022)* were a gift from Ruth Lehmann (Whitehead Institute, MIT, USA). All flies and crosses were maintained on standard fly medium (0.8% agar, 2.75% yeast, 5.2% cornmeal, 11% dextrose) in an incubator at 25°C, 65% RH and 12H:12H light-dark cycle.

Fly crosses and embryo staging

To generate over-expression of E-cadherin during transepithelial migration, UASp (over-expression) or Oregon R (wild type control) lines were crossed to *nanos-gal4-VP16* and heterozygous F1 virgin females were collected. 40-50 F1 virgin females were then

408 crossed with 8-10 Oregon R males and transferred into an egg collection cage with a

409 5mm apple juice agar plate as base (70). The *nanos-gal4-VP16* line is highly expressed in

410 the maternal germ line and deposited in the embryo but is not zygotically expressed in

411 the germ cells or somatic cells during TEM. In the case of *nos-LifeAct-tdTomato* and *Vasa-*

412 *GFP, E-cadherin-mScarlet* flies, an approximate 1:1 ratio of homozygous males and

413 females were transferred into an identical cage setup. Egg laying was allowed to

414 proceed in 30-minute windows and embryos were aged for five hours after the

415 midpoint of the egg laying window. After collection, embryos were either fixed

416 (embryonic over-expression) or used for live imaging (*nos-LifeAct-tdTomato* and *Vasa-*

417 *GFP, E-cadherin-mScarlet)*.

419 *Live imaging*

420 Appropriately staged embryos were manually dechorionated with forceps on double-

421 sided scotch tape and attached to a standard 90 mm polystyrene Petri dish using

422 heptane glue. The dish was then filled with sterile distilled water and mounted on a

423 Zeiss LSM 980 multi-photon microscope. Using a 20x water immersion objective, the

424 total volume of the embryonic gut was imaged at a two-photon excitation wavelength

425 of 1000nm (*nos-LifeAct-tdTomato)*, and 900nm/1050nm (*Vasa-GFP, E-cadherin-mScarlet*),

426 every 1, 5 or 10 minutes for the indicated period.

428 *Fixation and immunostaining*

Appropriately staged embryos were collected on a mesh, rinsed with Milli-Q water, and

dechorionated using 70%(v/v) commercial bleach until most of the embryos had lost

the chorion (confirmed visually). Embryos were fixed in a 1:1 mix of heptane and 4%

paraformaldehyde in 1X phosphate buffered saline (1X PBS) at room temperature with

nutation for 20 minutes. After removing the aqueous layer, an equal volume of 100%

methanol was added, and embryos were devitellinized by vigorous manual shaking for

2–3 minutes. Fixed embryos were stored in 100% methanol at –20 °C.

For staining, embryos were rehydrated in 1X PBS, then washed, permeabilized

and blocked twice for 15 minutes in PBTB (1X PBS, 0.2% Triton X-100, 1 mg/mL bovine

serum albumin (Sigma-Aldrich, A9418). Embryos were then incubated with primary

antibodies diluted in PBTB overnight at 4 °C. The next day, embryos were washed (4 ×

15 min, PBTB) and incubated overnight at 4 °C with fluorophore-conjugated secondary

antibodies and DAPI diluted in PBTB containing 4% normal goat serum (Jackson

Immunoresearch, 005-000-121). On the third day, embryos were washed (4 × 15 min in

1X PBS) and mounted in Vectashield (Vector laboratories, H1000) for imaging. Primary

antibodies used were rat anti-E-cadherin (1:20; DSHB, DCAD2), chicken anti-Vasa

(1:800) (Repouliou et al., 2025) and rabbit anti-GFP conjugated to AlexaFluor 488

(1:1000; Invitrogen, A-21311). Secondary antibody used was goat anti-rat (1:200)

conjugated to AlexaFluor 633 (Invitrogen, A-21094), goat anti-chicken (1:200)

conjugated to AlexaFluor 568 (Invitrogen, A-11041). DNA was stained with DAPI

(Sigma-Aldrich, D9542) at 1:2000 dilution of a 10 mg/mL stock.

Image Analysis and Quantification

Live cell tracking was performed using the built-in Mastodon plugin in Fiji/ImageJ (71).

Cell position relative to the gut lumen was determined manually for each time point.

Fluorescence intensities were quantified by drawing manual regions of interest and

measuring mean gray values using basic tools in Fiji/ImageJ (71). Germ cell positions,

gut lumen boundary, and the center of the lumen were marked using the built-in Cell

Counter function in Fiji/ImageJ (71). Germ cells were classified as "Out" if their

distance from the lumen center exceeded that of the lumen boundary; otherwise, they

were classified as "In".

In silico model of transepithelial migration

A cross-sectional slice of the epithelial midgut was modelled as a two-dimensional ring

formed by 20 epithelial cells. The epithelial ring is embedded in a radial chemical field

representing the spatial concentration of the chemoattractant. The magnitude of the

chemical concentration increases linearly with the radius, and the minimum coincides

with the center of the epithelial ring. At t=0, the germ cell begins slightly off-center,

with its initial angle randomly chosen between 0 and 360 degrees, and then migrates

outward toward regions of higher chemoattractant concentration.

Our two-dimensional model of the epithelial ring and the germ cells was

formulated in the Cellular Potts Model (CPM) framework (38–40). The cells in a CPM

are represented as a contiguous collection of pixels. Pairwise interactions between

pixels at the cell boundaries have an associated energy cost, J. Given two pixels a and b,

we use the value of $J_{a,b}$ to reflect the adhesion energy of the cells to which the two

interacting boundary pixels belong. Our model assumes that adhesion energy increases

with the number of adherens junctions formed, and that the number of adherens

junctions is in turn a function of the E-cadherin concentration. The maximum possible

number of junctions between two cells is thus determined by the surface E-cadherin

concentration of the cell that has fewer such molecules. As a result of our assumptions,

if $\sigma(a)$, $\sigma(b)$ are the indices of the cells to which pixels a and b belong, and $C(\sigma(a))$ and

$C(\sigma(b))$ are the E-cadherin concentrations of the two cells, then the interaction energy

term is $J_{a,b} = min(C(\sigma(a)), C(\sigma(b)))$. In our work, we have assumed that all epithelial

cells have E-cadherin concentration equal to C_E, and that all germ cells have E-cadherin

concentration equal to C_G. The energy function of the system of pixels is given by the

following equation:

485
$$
E = -\sum_{a,b} min(C(\sigma(a)), C(\sigma(b))) - \sum_{\sigma} \lambda_A \, (A_\sigma - A_0)^2 - \sum_{\sigma} \lambda_L \, (L_\sigma - L_0)^2
$$

The λ_A and λ_L terms determine the magnitude of the energy cost required for a cell area

and perimeter to deviate from the steady state values A_0 and L_0 respectively. The

system is simulated using the metropolis algorithm (45), where the probability of a pixel

flip is related to the change in the magnitude of the energy of the system as a result of

that specific pixel flip. Under appropriate energy conditions, pixels at the boundary

may flip to the cell state of one of their neighbors that has a different cell state. At each

iteration of the simulation algorithm, one such potential pixel flip is considered, and the

pixel is flipped if the resulting system has lower energy than the current state. If the

pixel flip would not reduce the energy of the system, the pixel flips with probability

$p_{flip} = exp(-\Delta E / T)$, where ΔE is the change in energy due to the pixel flip and T is the

temperature. The temperature term T represents the extent of stochasticity (noise) in the

pixel dynamics, with higher T increasing the probability of pixel flips that increase the

energy of the system. Chemotaxis is implemented within the CPM framework by

biasing pixel flips that aid cell movement in the direction of the gradient of

chemoattractant. In the presence of the attractant whose value at any pixel in 2D space

is $M_{a,b}$, the change in energy due to a pixel flip, where a pixel at (a, b) is copying the

state of a pixel at (a', b') becomes $\Delta E' = \Delta E - \lambda_C (M_{a',b'} - M_{a,b})$ (40). This additional term

favors the pixel flip when the chemical field value at (a', b') is higher than (a, b). To

prevent the epithelial ring from collapsing and forming a cell aggregate without a

lumen, we model the lumen as one large cell with a volume constraint. In addition, we

mechanically couple the epithelial cells by connecting the centroids of the adjacent cells

with springs using the FocalPointPlasticity plugin of CompuCell3D (72). The key

parameters that we explored in our work are C_G, C_E and λ_M. The CompuCell3D

implementation of our model can be found at

https://github.com/boyonpointe/GermCellTransepithelialMigration with commit ID

7ce2534.

Acknowledgements

We thank members of the Extavour Lab for helpful discussions, in particular Dr. Emily

Rivard and Dr. Petra Kovacikova for their comments on the manuscript. We thank Prof.

James Glazier for helpful discussions regarding CompuCell3D, the Drosophila

Genomics Research Center (NIH Grant 2P40OD010949) and Dr. Xiaona Tang for their

help in making the E-cadherin-mScarlet CRISPR flies. We thank Ruth Lehmann for

sharing fly lines.

Data Availability

523 Scripts for Compucell3D simulations and image analysis are available at

524 https://github.com/boyonpointe/GermCellTransepithelialMigration with commit ID

525 7ce2534.

Author Contributions

528 CK conceived of the study, designed and performed all computational experiments,

529 performed data analysis and interpretation, and wrote the first draft of the manuscript.

530 SG conceived of the study, designed and performed wet lab experiments, and reviewed

531 and edited the manuscript. CGE obtained funding for the study, supervised its execution,

532 and reviewed and edited the manuscript.

Funding

535 This study was supported by a postdoctoral fellowship to CK through the NSF-Simons

536 Center for Mathematical and Statistical Analysis of Biology at Harvard (award number

537 DMS-1764269), the Harvard Quantitative Biology Initiative, and by funds from Harvard

538 University and the Howard Hughes Medical Institute (HHMI). CGE is an HHMI

539 Investigator.

Conflicts of Interest

542 The authors declare no conflicts of interest.

Figure Legends

Figure 1: Germ cells exit the midgut individually at comparable speeds. (a-a'') Schematic of *D. melanogaster* embryonic germ line development. The primordial germ cells are formed at the posterior tip at stage 5. Following gastrulation, they are contained in the midgut at stage 7, and eventually exit the midgut through transepithelial migration at stage 10. **(a''')** Schematic showing the three different types of intercellular junction where we quantified E-cadherin intensity. **(b-b'')** Three snapshots from live imaging of the TEM process as the germ cells make their way through the midgut. **(c)** The distribution of the distances of the germ cells from center of the midgut lumen at five different time points are shown. Small numbers under plots indicate sample sizes (number of cells). **(d- d'')** Movement of germ cells in time measured with respect to their initial (t = 0) position for two different live imaged embryos. Black circles: positions within lumen; magenta rhomboids: positions within the midgut epithelium; white squares: positions outside the midgut. **(e)** Higher magnification of the midgut during germ cell transepithelial migration. Magenta: E-cadherin; green: germ cell-specific protein Vasa. **(e')** Germ cells express E-cadherin during the transepithelial migration (boxes). **(e'')** The relative intensity (au: arbitrary units) of the E-cadherin fluorescent signal at the epithelial cell – germ cell junction (EG), lateral epithelial – epithelial junction (E_{lat}) and apical epithelial – epithelial junctions (E_{api}). n = 24 junctions for each of the three categories. Two-sample Mann-Whitney U tests were used to compute the p values that are shown in panel (e'') for pairwise comparisons of distributions.

Figure 2: Greater epithelial junctional remodeling aids germ cell transmigration. (a) Snapshot of Cellular Potts Model simulation of a germ cell (yellow) exiting the two-dimensional epithelial ring (green). Pixels at the boundary of the cells are shown in black. Colored arrows point in the direction of chemoattractant gradient; colors represent the concentration of the chemoattractant. **(b-b'')** Snapshots of the germ cell (b) before entering the epithelial barrier, (b') while inside the barrier and (b'') just before exiting the barrier. Numbers at top right indicate the time in units of Monte Carlo steps (mcs). **(c)** Average time taken (τ_e) for the germ cell to exit the midgut decreases with increase in interepithelial dynamics (T_E). **(c')** Probability of successfully exiting (p_e) increases with T_E. Trends in c and c' are independent of the strength of the chemoattractant (λ_M). The shaded silver line corresponds to

$T_E = 10$, the value at which al simulations described herein are carried out, unless specified

otherwise. Simulation parameters: $\lambda_M = 7000, C_E = 10, C_G = 5, T = 10$.

Figure 3: Time to exit the epithelial barrier has a non-monotonic dependence on germ cell E-

cadherin concentration. (a-a'') The time a simulated germ cell takes to exit the 2D midgut

epithelial barrier as a function of germ cell E-cadherin concentration (C_G). (a), (a') and (a'')

represent increasing epithelial cell E-cadherin concentrations ($C_E = 8$, $C_E = 10$, $C_E = 12$

respectively), offering increasing resistance to the germ cell transepithelial migration. The

curves in each of the three panels are colored according to the strength of the chemical field

(λ_M). Regardless of the value of C_E, for strong enough values of λ_M the time to exit shows a

minimum that is independent of the λ_M value. X-axis label "C_G" underneath **(a'')** applies also to

(a) and **(a')**. **(b)** The coefficient of variation in time to exit as a function of C_G for the case of $C_E =$

10. **(c)** Trajectories of germ cells for a typical parameter set obtained from 1000 independent

stochastic simulations. Silver: successful TEM; brown: germ cell exit failure; black lines: mean

position of the epithelial cells forming the 2D barrier. **(d)** Mean distance covered by the germ

cell within the barrier (l_e) as a function of time . The time τ is normalized by the total time of

travel, where 0 represents time of entry into the barrier and 1 represents time of exit. When the

germ cell E-cadherin concentration is low, the germ cell covers much of the distance in a very

short interval of time at the end of the journey. Higher germ cell E-cadherin levels lead to the

germ cell migrating at roughly uniform speed throughout the journey. **(d')** The time taken to

cover the second half of the journey τ_2 as a function of germ cell E-cadherin concentration. **(e)**

The relationship between τ_{exit} and p_{exit} is shown using data aggregated by simulating the

model over various values of λ_M, C_G and C_E. τ_{exit} decreases monotonically with p_{exit}, implying

that the shorter the time to exit, the higher the chances of exiting.

Figure 4: Higher germ cell E-cadherin promotes quicker midgut exit *in vivo*. (a) (L-R)

Representative micrographs showing the extent of transepithelial migration in the four

experimental groups. I corresponds to the control, II and III represent lines where E-cadherin

is overexpressed in all cells and IV corresponds to embryos where E-cadherin overexpression

is targeted to the germ cells. **(b)** The distribution of distances of germ cells from the center of

the lumen for the four experimental groups. Gray and black symbols represent cells inside and
outside the gut respectively. The distances of the germ cells from the lumen are largest in the
condition with increased germ cell-targeted E-cadherin overexpression (IV). Small numbers
under plots indicate sample sizes (number of germ cells). **(c)** Mean distance of the germ cells
from the center of the lumen in each embryo for the four experimental groups. Small numbers
under plots indicate sample sizes (number of embryos). **(d)** Fractions of germ cells per embryo
that have exited the midgut by 5.25 hours after egg laying in the four experimental groups.
Small numbers under plots indicate sample sizes (number of embryos). Two sample Mann-
Whitney U test was used to compute the p values that are shown in panels **(b)-(d)** for pairwise
distribution comparisons. **(e)** We hypothesize that the WT embryos (group I) have C_G in the
pink shaded region, and that germ cell-targeted E-cadherin expression pushes the system
towards the green shaded region, leading to shortening of τ_e.

References

1. A. Aman, T. Piotrowski, Cell migration during morphogenesis. *Dev. Biol.* **341**, 20–33 (2010).

2. E. Scarpa, R. Mayor, Collective cell migration in development. *J. Cell Biol.* **212**, 143–155 (2016).

3. D. Dormann, C. J. Weijer, Chemotactic cell movement during development. *Curr. Opin. Genet. Dev.* **13**, 358–364 (2003).

4. P. Roca-Cusachs, R. Sunyer, X. Trepat, Mechanical guidance of cell migration: lessons from chemotaxis. *Curr. Opin. Cell Biol.* **25**, 543–549 (2013).

5. E. T. Roussos, J. S. Condeelis, A. Patsialou, Chemotaxis in cancer. *Nat. Rev. Cancer* **11**, 573–587 (2011).

6. S. SenGupta, C. A. Parent, J. E. Bear, The principles of directed cell migration. *Nat. Rev. Mol. Cell Biol.* **22**, 529–547 (2021).

7. M. Bezanilla, A. S. Gladfelter, D. R. Kovar, W.-L. Lee, Cytoskeletal dynamics: A view from the membrane. *J. Cell Biol.* **209**, 329–337 (2015).

8. K. Rottner, T. E. B. Stradal, Actin dynamics and turnover in cell motility. *Curr. Opin. Cell Biol.* **23**, 569–578 (2011).

9. S. Seetharaman, S. Etienne-Manneville, Cytoskeletal Crosstalk in Cell Migration. *Trends Cell Biol.* **30**, 720–735 (2020).

10. S. F. Gilbert, S. F. Gilbert, *Developmental Biology*, 6th Ed. (Sinauer Associates, 2000).

11. P. Friedl, D. Gilmour, Collective cell migration in morphogenesis, regeneration and cancer. *Nat. Rev. Mol. Cell Biol.* **10**, 445–457 (2009).

12. W. T. Boyce, M. B. Sokolowski, G. E. Robinson, Genes and environments, development and time. *Proc. Natl. Acad. Sci.* **117**, 23235–23241 (2020).

13. M. Ebisuya, J. Briscoe, What does time mean in development? *Development* **145**, dev164368 (2018).

14. A. Kicheva, M. Cohen, J. Briscoe, Developmental Pattern Formation: Insights from Physics and Biology. *Science* **338**, 210–212 (2012).

15. J. Negrete, A. C. Oates, Towards a physical understanding of developmental patterning. *Nat. Rev. Genet.* **22**, 518–531 (2021).

16. C. M. Franz, G. E. Jones, A. J. Ridley, Cell Migration in Development and Disease. *Dev. Cell*
2, 153–158 (2002).

17. C. J. Weijer, Collective cell migration in development. *J. Cell Sci.* **122**, 3215–3223 (2009).

18. A. F. Huettner, The origin of the germ cells in Drosophila melanogaster. *J. Morphol.* **37**, 385–
423 (1923).

19. M. Rabinowitz, Studies on the cytology and early embryology of the egg of Drosophila
melanogaster. *J. Morphol.* **69**, 1–49 (1941).

20. M. K. Jaglarz, K. R. Howard, The active migration of Drosophila primordial germ cells.
Development **121**, 3495–3503 (1995).

21. A. C. Santos, R. Lehmann, Germ Cell Specification and Migration in Drosophila and
beyond. *Curr. Biol.* **14**, R578–R589 (2004).

22. L. J. Barton, *et al.*, Juvenile hormones direct primordial germ cell migration to the
embryonic gonad. *Curr. Biol.* **34**, 505-518.e6 (2024).

23. K. Hanyu-Nakamura, S. Kobayashi, A. Nakamura, Germ cell-autonomous Wunen2 is
required for germline development in Drosophila embryos. *Development* **131**, 4545–4553 (2004).

24. A. D. Renault, Y. J. Sigal, A. J. Morris, R. Lehmann, Soma-Germ Line Competition for Lipid
Phosphate Uptake Regulates Germ Cell Migration and Survival. *Science* **305**, 1963–1966 (2004).

25. H. Sano, A. D. Renault, R. Lehmann, Control of lateral migration and germ cell elimination
by the Drosophila melanogaster lipid phosphate phosphatases Wunen and Wunen 2. *J. Cell*
Biol. **171**, 675–683 (2005).

26. M. Slaidina, R. Lehmann, Quantitative Differences in a Single Maternal Factor Determine
Survival Probabilities among Drosophila Germ Cells. *Curr. Biol.* **27**, 291–297 (2017).

27. R. Warrior, Primordial Germ Cell Migration and the Assembly of the Drosophila
Embryonic Gonad. *Dev Biol* **166**, 180–194 (1994).

28. H. Oda, T. Uemura, Y. Harada, Y. Iwai, M. Takeichi, A *Drosophila* Homolog of Cadherin
Associated with Armadillo and Essential for Embryonic Cell-Cell Adhesion. *Dev. Biol.* **165**,
716–726 (1994).

29. U. Tepass, *et al.*, shotgun encodes Drosophila E-cadherin and is preferentially required
during cell rearrangement in the neurectoderm and other morphogenetically active epithelia.
Genes Dev. **10**, 672–685 (1996).

30. M. DeGennaro, *et al.*, Peroxiredoxin Stabilization of DE-Cadherin Promotes Primordial
Germ Cell Adhesion. *Dev. Cell* **20**, 233–243 (2011).

31. P. S. Kunwar, *et al.*, Tre1 GPCR initiates germ cell transepithelial migration by regulating
Drosophila melanogaster E-cadherin. *J. Cell Biol.* **183**, 157–168 (2008).

32. P. S. Kunwar, M. Starz-Gaiano, R. J. Bainton, U. Heberlein, R. Lehmann, Tre1, a G Protein-
Coupled Receptor, Directs Transepithelial Migration of Drosophila Germ Cells. *PLoS Biol.* **1**,
e80 (2003).

33. J. A. Campos-Ortega, V. Hartenstein, *The Embryonic Development of Drosophila melanogaster*
(Springer Science & Business Media, 2013).

34. D. Poulson, D. Waterhouse, Experimental Studies on Pole Cells and Midgut Differentiation
in Diptera. *Aust. J. Biol. Sci.* **13**, 541–567 (1960).

35. J. R. K. Seifert, R. Lehmann, Drosophila primordial germ cell migration requires epithelial
remodeling of the endoderm. *Development* **139**, 2101–2106 (2012).

36. G. Parés, S. Ricardo, FGF control of E-cadherin targeting in the Drosophila midgut impacts
on primordial germ cell motility. *J. Cell Sci.* **129**, 354–366 (2016).

37. D. Cai, *et al.*, Mechanical Feedback through E-Cadherin Promotes Direction Sensing during
Collective Cell Migration. *Cell* **157**, 1146–1159 (2014).

38. J. A. Glazier, F. Graner, Simulation of the differential adhesion driven rearrangement of
biological cells. *Phys. Rev. E* **47**, 2128–2154 (1993).

39. F. Graner, J. A. Glazier, Simulation of biological cell sorting using a two-dimensional
extended Potts model. *Phys. Rev. Lett.* **69**, 2013–2016 (1992).

40. P. Hogeweg, Evolving Mechanisms of Morphogenesis: on the Interplay between
Differential Adhesion and Cell Differentiation. *J. Theor. Biol.* **203**, 317–333 (2000).

41. D. Sweeton, S. Parks, M. Costa, E. Wieschaus, Gastrulation in Drosophila: the formation of
the ventral furrow and posterior midgut invaginations. *Development* **112**, 775–789 (1991).

42. B. Lin, J. Luo, R. Lehmann, An AMPK phosphoregulated RhoGEF feedback loop tunes
cortical flow–driven amoeboid migration in vivo. *Sci. Adv.* **8**, eabo0323 (2022).

43. M. Akhmanova, *et al.*, Cell division in tissues enables macrophage infiltration. *Science* **376**,
394–396 (2022).

44. M. K. Jaglarz, K. R. Howard, Primordial germ cell migration in Drosophila melanogaster is
controlled by somatic tissue. *Development* **120**, 83–89 (1994).

45. N. Metropolis, A. W. Rosenbluth, M. N. Rosenbluth, A. H. Teller, E. Teller, Equation of
State Calculations by Fast Computing Machines. *J. Chem Phys.* **21**, 1087–1092 (1953).

46. M. Starz-Gaiano, N. K. Cho, A. Forbes, R. Lehmann, Spatially restricted activity of a
Drosophila lipid phosphatase guides migrating germ cells. *Development* **128**, 983–991 (2001).

47. N. Zhang, J. Zhang, K. J. Purcell, Y. Cheng, K. Howard, The Drosophila protein Wunen
repels migrating germ cells. *Nature* **385**, 64–67 (1997).

48. A. H. Brand, N. Perrimon, Targeted gene expression as a means of altering cell fates and
generating dominant phenotypes. *Development* **118**, 401–415 (1993).

49. T. J. C. Harris, U. Tepass, Adherens junctions: from molecules to morphogenesis. *Nat. Rev.*
Mol. Cell Biol. **11**, 502–514 (2010).

50. S. Hong, R. B. Troyanovsky, S. M. Troyanovsky, Binding to F-actin guides cadherin cluster
assembly, stability, and movement. *J. Cell Biol.* **201**, 131–143 (2013).

51. R. B. Troyanovsky, E. P. Sokolov, S. M. Troyanovsky, Endocytosis of Cadherin from
Intracellular Junctions Is the Driving Force for Cadherin Adhesive Dimer Disassembly. *MBoC*
17, 3484–3493 (2006).

52. B.-A. Truong Quang, M. Mani, O. Markova, T. Lecuit, P.-F. Lenne, Principles of E-Cadherin
Supramolecular Organization In Vivo. *Curr. Biol.* **23**, 2197–2207 (2013).

53. Y. Wu, P. Kanchanawong, R. Zaidel-Bar, Actin-Delimited Adhesion-Independent
Clustering of E-Cadherin Forms the Nanoscale Building Blocks of Adherens Junctions. *Dev.*
Cell **32**, 139–154 (2015).

54. M. D. Martin-Bermudo, O. M. Dunin-Borkowski, N. H. Brown, Specificity of PS integrin
function during embryogenesis resides in the α subunit extracellular domain. *EMBO J.* **16**,
4184–4193 (1997).

55. A. Huttenlocher, Cell polarization mechanisms during directed cell migration. *Nat. Cell*
Biol. **7**, 336–337 (2005).

56. J. M. Halbleib, W. J. Nelson, Cadherins in development: cell adhesion, sorting, and tissue
morphogenesis. *Genes Dev.* **20**, 3199–3214 (2006).

57. M. Sandig, E. B. Voura, V. I. Kalnins, C.-H. Siu, Role of cadherins in the transendothelial
migration of melanoma cells in culture. *Cell Motil. Cytoskelet.* **38**, 351–364 (1997).

58. D. Vestweber, How leukocytes cross the vascular endothelium. *Nat. Rev. Immunol.* **15**, 692–
704 (2015).

59. Y.-T. Yeh, *et al.*, Three-dimensional forces exerted by leukocytes and vascular endothelial
cells dynamically facilitate diapedesis. *Proc. Natl. Acad. Sci.* **115**, 133–138 (2018).

60. J. Qi, N. Chen, J. Wang, C.-H. Siu, Transendothelial Migration of Melanoma Cells Involves
N-Cadherin-mediated Adhesion and Activation of the β-Catenin Signaling Pathway. *Mol. Biol.*
Cell **16**, 4386–4397 (2005).

61. C. E. Chan, D. J. Odde, Traction Dynamics of Filopodia on Compliant Substrates. *Science*
322, 1687–1691 (2008).

62. P. Maiuri, *et al.*, Actin Flows Mediate a Universal Coupling between Cell Speed and Cell
Persistence. *Cell* **161**, 374–386 (2015).

63. K. M. Yamada, M. Sixt, Mechanisms of 3D cell migration. *Nat. Rev. Mol. Cell Biol.* **20**, 738–
752 (2019).

64. V. Swaminathan, C. M. Waterman, The molecular clutch model for mechanotransduction
evolves. *Nat. Cell Biol.* **18**, 459–461 (2016).

65. L. B. Case, C. M. Waterman, Integration of actin dynamics and cell adhesion by a three-
dimensional, mechanosensitive molecular clutch. *Nat. Cell Biol.* **17**, 955–963 (2015).

66. L. J. Barton, L. R. la Cruz, R. Lehmann, B. Lin, The journey of a generation: advances and
promises in the study of primordial germ cell migration. *Development* **151**, dev201102 (2024).

67. D. J. Montell, P. Rorth, A. C. Spradling, slow border cells, a locus required for a
developmentally regulated cell migration during oogenesis, encodes Drosophila CEBP. *Cell* **71**,
51–62 (1992).

68. R. A. Foty, M. S. Steinberg, The differential adhesion hypothesis: a direct evaluation. *Dev.*
Biol. **278**, 255–263 (2005).

69. S. J. Gratz, *et al.*, Highly Specific and Efficient CRISPR/Cas9-Catalyzed Homology-Directed
Repair in Drosophila. *Genetics* **196**, 961–971 (2014).

70. W. F. Rothwell, W. Sullivan, Drosophila Embryo Collection. *Cold Spring Harb. Protoc.* **2007**,
pdb.prot4825 (2007).

71. J. Schindelin, *et al.*, Fiji: an open-source platform for biological-image analysis. *Nat. Methods*
9, 676–82 (2012).

72. M. H. Swat, *et al.*, Multi-Scale Modeling of Tissues Using CompuCell3D. *Methods Cell Biol.*
110, 325–366 (2012).

(a) (a') (a'') (a''')

E_api
E_lat
EG

(b) 5 min

(b') 90 min

(b'') 190 min

(c)

d_{gc}

t (min)

(d)

Δd_{gc}

● in □ out ◆ transit

(d')

Δd_{gc}

t (min)

(e) E-cad Vasa

(e')

(e'')

EG
E_lat
E_api

I_{Ecad}

(a)

I: WT   II: Pan-Ecad-OE-A   III: Pan-Ecad-OE-B   IV: GC-Ecad-OE

Vasa   DAPI

(b) $+$ Out $\cdot$ In   0.14

4.78×10^{-43}

7.22×10^{-76}

1.03×10^{-23}

d_{gc}

n = 595   n = 513   n = 521   n = 392

I   II   III   IV

(c)   0.95

8.26×10^{-5}

4.75×10^{-6}

1.9×10^{-4}

$\overline{d_{gc}}$

n = 20   n = 17   n = 18   n = 14

I   II   III   IV

(d) 1.24×10^{-6} 0.24

6.18×10^{-6}

1.39×10^{-5}

f_e

1.0

0.5

0.0

n = 20   n = 17   n = 18   n = 14

I   II   III   IV

(e)   WT   More E-cad   Less E-cad

τ_e

C_G