Old growth attributes by chain saw: how between-patch heterogeneity changes the metacommunities of beetles in temperate forests

Oliver Mitesser[1], Marc Cadotte[2], Akira S Mori[3], Fons van der Plas[4], Anne Chao[5], Julia Rothacher[1], Claus Bässler[6], Mirjana Bevanda[7], Peter H. W. Biedermann[8], Pia Bradler[9], Antonio Castaneda-Gomez[7], Orsi Decker[1], Benjamin M. Delory[10], Sebastian Dittrich[11], Andreas Fichtner[9], Alexander Kreis[1], Lisa Köstler-Albert[12], Goddert von Oheimb[13], Kerstin Pierick[14], Simon Thorn[15], Leah Vogelfänger[16], Wolfgang Weisser[17], Martin Wegmann[7], Clara Wild[1], Jörg Müller[1,18]

Corresponding author: Oliver Mitesser, Chair of Conservation Biology and Forest Ecology, Biocenter, University of Würzburg, Glashüttenstr. 5, 96181 Rauhenebrach, oliver.mitesser@uni-wuerzburg.de, +49 931 3186876

Affiliations:

1) Chair of Conservation Biology and Forest Ecology, Biocenter, Julius-Maximilians-Universität Würzburg, Rauhenebrach, Germany

2) Biological Sciences, University of Toronto-Scarborough, Toronto, Ontario, Canada

3) Research Center for Advanced Science and Technology, University of Tokyo, Tokyo, Japan

4) Plant Ecology and Nature Conservation, Wageningen University, Wageningen, The Netherlands

5) Institute of Statistics, National Tsing Hua University, Hsin-Chu, Taiwan

6) University of Bayreuth, Universitätsstraße 30, 95440 Bayreuth, Germany

7) Department of Remote Sensing, Insitute of Geography and Geology, University of

Würzburg, 97074 Würzburg, Germany

8) Forest Entomology and Forest Protection, Albert Ludwig University of Freiburg, Freiburg

im Breisgau, Germany

9) Institute of Ecology, Leuphana University, Lüneburg, Germany

10) Environmental Sciences Group, Copernicus Institute of Sustainable Development, Utrecht

University, Utrecht, The Netherlands

11) Biodiversity and Conservation, Technical University of Dresden, Tharandt, Germany

12) Animal Population Ecology, Bayreuth Center for Ecology and Environmental Research

(BayCEER), University Bayreuth, Bayreuth, Germany

13) Institute of General Ecology and Environmental Protection, Technical University of

Dresden, Tharandt, Germany

14) Spatial Structures and Digitization of Forests/Silviculture and Forest Ecology of the

Temperate Zones, University of Göttingen, Göttingen, Germany

15) Department of Biology, University of Marburg, Marburg, Germany

16) Forest Zoology, Technical University of Dresden, Tharandt, Germany

17) Terrestrial Ecology Research Group, Department of Ecology and Ecosystem

Management, School of Life Sciences Weihenstephan, Technical University of Munich,

Freising, Germany

18) Bavarian Forest National Park, Freyunger Str. 2, 94481 Grafenau, Germany

Emails:

Marc Cadotte: marc.cadotte@utoronto.ca

Fons van der Plas: fons.vanderplas@wur.nl

Akira S Mori: akkym@g.ecc.u-tokyo.ac.jp

Anne Chao: chao@stat.nthu.edu.tw

Julia Rothacher: julia.rothacher@uni-wuerzburg.de

Antonio Castañeda Gomez: antonio.castaneda-gomez@uni-wuerzburg.de

Orsi Decker: Orsolya.Decker@npv-bw.bayern.de

Mirjana Bevanda: mirjana.bevanda@uni-wuerzburg.de

Peter H. W. Biedermann: peter.biedermann@forento.uni-freiburg.de

Pia Marie Bradler: pia_marie.bradler@tu-dresden.de

Sebastian Dittrich: sdittric@msx.tu-dresden.de

Andreas Fichtner: andreas.fichtner@leuphana.de

Lisa Köstler-Albert: Lisa.Albert@uni-bayreuth.de

Goddert von Oheimb: Goddert_v_Oheimb@tu-dresden.de

Simon Thorn: simon.thorn@hlnug.hessen.de

Wolfgang Weisser: wolfgang.weisser@tum.de

Martin Wegmann: martin.wegmann@uni-wuerzburg.de

Clara Mila Wild: clara.wild@uni-wuerzburg.de

Jörg Müller: Joerg.Mueller@npv-bw.bayern.de

Leah Vogelfänger: leah_rebekka.vogelfaenger@tu-dresden.de

Alexander Kreis: alexanderkreis01@gmail.com

Kerstin Pierick: kerstin.pierick@uni-goettingen.de

Statement of authorship: JM, MC, AM, FP, ST, MW, OM and WW designed the study,

JR,CB, MB, PBi, PBa, AC, OD, BD, SD, AF, AK, LK, GO, ST, LV, and CW performed the

research, AC provided the meta-analysis, and JM and OM wrote the first draft of the

manuscript, and all authors contributed substantially to revisions.

Data and code are available at https://figshare.com/s/4b6b4d5a5cfbcf0dbac3. DOI

10.6084/m9.figshare.29994688 is reserved to be published when the manuscript is accepted.

Short running title: Metacommunities in heterogenized forests

Type of article: Letter

Number of words in the abstract: 150

Number of words in the main text (excluding abstract, acknowledgements, references,

table and figure legends): 5000

Number of references: 67

Number of figures and tables: 3 figures, 1 table

Keywords: diversity, metacommunity paradigms, beetles, Coleoptera, temperate forests

Abstract

Metacommunity theory has expanded our understanding of how spatial dynamics and local interactions influence species communities. Different assembly archetypes, reflecting different roles of species differences, habitat differences, and dispersal have been described, but we lack empirical studies specifically in terrestrial habitats testing which archetype is most important. In a replicated design we experimentally enhanced structural between-patch heterogeneity in homogeneous production forests and developed a statistical framework controlling for sample incompleteness to detect different metacommunity processes. Meta-analyses on >100K individuals of >1.3K beetle species showed an increase of ~60 species in heterogenized forests at γ-level promoted by increasing α-diversity consistent with the *mass-effect* and an increase of β-diversity by ~10% supporting *species-sorting*. Additionally, we tested β-deviations from random assembly as a proxy of *neutral processes*. Findings indicate that enhancing structural heterogeneity can shift forests from *patch-dynamics* dominance towards *mass-effect* and *species-sorting*, offering a promising pathway to restore biodiversity in managed landscapes.

Introduction

During the last two decades, metacommunity theory has expanded our understanding of how spatial dynamics and local interactions influence the structure and diversity of species communities (Leibold *et al.* 2004; Cottenie 2005; Gonzalez 2009; Record *et al.* 2021). Within this framework, four archetypes have been developed to characterize metacommunities (Logue *et al.* 2011; Leibold & Chase 2018). Each of them predicts different assembly mechanisms based on dispersal and differences in habitats and species: The *patch-dynamics* archetype assumes homogeneous patches which are occupied by species with a trade-off between dispersal and local dominance. Superior competitors can displace others in occupied patches, whereas weaker competitors may persist at the regional level by rapidly colonizing newly available patches. Variation in species composition is due to processes of colonization and extinction. The *species-sorting* archetype assumes that habitat patches are heterogeneous, and dispersal capacity is just sufficiently high to reach the different patches. Local community composition is primarily shaped by a match between environmental conditions and the ecological niches of the species. A high level of environmental variation across the landscape thus leads to high regional species richness through different species assemblages in different patches. The *mass-effect* archetype assumes that heterogeneous habitat patches are interconnected, and species' dispersal abilities are high. This allows species to reproduce in source patches and frequent dispersal allows them to also exist in sink patches where local conditions prevent long-term survival or reproduction. Massive dispersal over long distances might override signals of species-niche matching in the overall biodiversity response. Last, the *neutral* archetype relies on the assumption that individuals of different species are equivalent in survival, reproduction, and dispersal abilities. Assembly processes entirely depend on stochastic events including immigration and extinction (Hubbell 2001). Ecological drift and rare speciation shape regional diversity patterns over large time scales and large metacommunities serve as persistent sources of species for local communities.

Theoretical studies have shown that the different metacommunity archetypes make different predictions on how biodiversity is distributed across space and which predictors explain the patterns (Thompson *et al.* 2020). While processes of all archetypes have been demonstrated to matter, these relationships can be utilized to identify the most influential mechanisms in real-world landscapes (see hypotheses below). Despite the theoretical progress in metacommunity theory, it is still insufficiently linked to empirical data, particularly in terrestrial habitats (Logue *et al.* 2011), where soft boundaries between patches with gradual transitions of resources, community composition, and processes are the rule rather than the exception. To disentangle the importance of different metacommunity archetypes, Logue *et al.* (2011) proposed developing an integrative framework that combines new experimental approaches and novel statistical methods and is based on three key axes: species differences (functional equivalence vs. differentiation), habitat differences (homogeneity vs. heterogeneity), and dispersal dynamics (Fig. 1a).

We selected homogeneous production forests as a model system to implement a novel large-scale experimental approach (Müller *et al.* 2023) aimed at identifying the dominant processes driving metacommunity assembly and assessing how their relative importance varies with landscape heterogeneity. At the beginning of the experiment, we manipulated the between-forest-patch heterogeneity by creating gaps and deadwood towards old-growth features by chainsaw and selected homogeneous control forests. In parallel, we developed a new meta-analysis approach to disentangle the effects of within-patch α- and between-patch β-diversity on γ-diversity at landscape scale while accounting for sample incompleteness in empirical data for taxonomic (TD), functional (FD), and phylogenetic diversity (PD) along Hill numbers to capture differences in species abundances and ecological attributes (Chao *et al.* 2023). Considering these multiple dimensions of diversity (facet: TD, FD, PD; scale: α, β, γ; Hill number: q=0, 1, 2) enables the identification of underlying processes structuring species

assemblages, depending on whether diversity is shaped by distinct functional traits, local versus

regional patterns, or rare versus dominant species, respectively (Jeliazkov & Chase 2024).

The design allowed us to use the generalized concept of diversity decomposition efficiently in

a replicated experiment at the landscape level. In the same framework, we were able to estimate

pairwise β-deviations (from randomized null models) between-patches of a forest to create null-

model communities (*neutral archetype;* Logue *et al.* 2011) avoiding well documented artefacts

in β-diversity by under-sampling (Tuomisto 2010; Tucker *et al.* 2016). Subtraction of null-

model β-diversity in general controls for effects by demographic and dispersal stochasticity on

the β-diversity pattern expected from the *neutral* archetype (Tucker *et al.* 2016). β-deviation

and spatial and environmental distances offered the option of deciphering the assembly

mechanisms of metacommunities.

We used beetles as one of the taxonomically, phylogenetically, and functionally most

diverse insect orders (Farrell 1998) which are particularly vulnerable under climate change

(Thom & Seidl 2016). We expected a shift from metacommunities dominated by *patch-*

dynamics in homogeneous control forests to metacommunities structured by *species-sorting*

and *mass-effects* stimulating our hypotheses about diversities (α, β, and γ) and β-deviation. We

first test the hypothesis (H1) that enhancement of between-patch heterogeneity increases γ-

diversity of a forest. Along with this hypothesis, we predict an increase of γ-diversity by an

increase of β-diversity according to *species-sorting* or by an increase of α-diversity by the *mass-*

effect promoting source patches by resource enrichment, but at the same time eventually

reducing β-diversity (Mouquet & Loreau 2003) (Fig. 1). Secondly, we hypothesise (H2) that β-

deviation is better explained by spatial distance in homogeneous forests because other

environmental attributes as potential alternative predictors have only low variation. In

heterogeneous forests where *species-sorting* can increase the correlation between communities

and environment we hypothesise (H3) environmental variation to be the better predictor of β-

deviation. As species differences in habitat requirements and in mobility affect *species-sorting*

and dispersal we expect effects of habitat difference in heterogeneous forests specifically on

functional/phylogenetic β-deviation.

Our results demonstrate an increase in α-, β-, and γ-diversity in response to ESBC treatment

and suggest that metacommunities in homogeneous forests are dominated by *patch-dynamics*,

while those in heterogeneous forests by *mass-effect* and *species-sorting*.

[[Figure 1 about here]]

Materials and methods

Study area and experimental design

We established our pairwise design of a district experimentally enhanced in structural beta

complexity (ESBC) and a homogeneous control district at 11 sites throughout Germany

covering a wide range of climate (c. 50 - 1100 m a.s.l.) and soil conditions (acidic to calcareous)

in European beech (*Fagus sylvatica*) mixed forests (Müller *et al.* 2023). In total, the study

included 22 districts, and within each district a set of 9 or 15 experimental patches was

established, with a spatial extent of 50 m × 50 m in a randomized block design. The

interventions aimed to enhance the variation in canopy density and deadwood structures.

Manipulations were carried out in the winter of 2015/16 (Bavarian Forest, Passau), 2016/17

(Saarland, Lübeck, Hunsrück) and 2018/19 (University Forest, i.e. the forest of the University

of Würzburg) to allow sufficient time for the colonization by forest fauna. LiDAR

measurements validate that the ESBC districts remained structurally more heterogeneous than

the control districts during the study period (Pierick *et al.* 2025, *preprint*).

[[Figure 2 about here]]

Beetle data

In each patch we set up two flight interception traps and two pitfall traps to reach sufficient

sample size (number of individuals) to represent the local environmental conditions (see

(Müller & Brandl 2009). Sampling was conducted from May to July 2022 in the University

Forest (3 sites), the Bavarian Forest (4 sites), and the Passau region (1 site), while the remaining

sites were sampled in 2023. These months represent >90% of individuals in beetle sampling

during the vegetation period. Traps were emptied monthly and sorted on order level. Species

identification was conducted by professional taxonomists. As traits we selected 1) body size as

a universal trait for aspects of reproduction, dispersal ability, and general population size

(Chown & Gaston 2010), 2) brightness level as an important trait for species response to climate

conditions (Zeuss *et al.* 2014), 3) the adoption of a life style in deadwood (saproxylic), and 4)

the feeding strategy. All trait information was taken from literature (Freude *et al.* 1964-1983;

Koch 1989-1992; Seibold *et al.* 2015; Hagge *et al.* 2021). As phylogenies of beetles are still

incomplete, we used as a surrogate a phylogeny based on taxonomic classification (Freude *et*

al. 1964-1983; Abrego *et al.* 2024).

Environmental and spatial difference

To quantify habitat heterogeneity, we measured the distance between two patches of a district

first by forest structure. We used canopy densities measured by LiDAR via drone flights using

the penetration ratio at 6 m above ground (Müller & Brandl 2009) and the volume of deadwood

per patch to quantify the level of ESBC. Variation in both variables was predominantly driven

by our experimental ESBC manipulation. However, as natural tree mortality is still increasing

in Europe (Senf *et al.* 2018), both also capture add-on natural dieback processes. There might

be additional more subtle variation or changes between forest patches in environmental

conditions, e.g. soil moisture, not captured by LiDAR and deadwood volume. Therefore, we

additionally calculated unweighted mean Ellenberg-indicator values based on plant relevés

estimated according to a simplified percentage scale (Dittrich et al. 2013) and recorded in all

patches (Ellenberg *et al.* 1991; Ewald 2003), which has been proven to be highly integrative in

describing such subtle environmental conditions (Ewald 2003, Müller et al. 2009). We used the

difference between patches of these values as a proxy for abiotic distance. To avoid overlap

between information in structure measured by LiDAR and these indicator values we removed

the light indicator value. Spatial distance between patches was calculated in meters based on

projected coordinates in the ETRS1989 format.

Statistical methods

Meta-analyses

Statistical analyses were conducted in R (version 4.3.3) (R Core Team, 2023). To allow for

direct comparison, taxonomic (TD), phylogenetic (PD), and functional diversity (FD) were

assessed as the effective number of equally abundant species, divergent lineages, and distinct

functional groups and for Hill numbers $q = 0, 1, 2$ focusing on rare, common, and dominant

species. These facets of beetle diversity and corresponding confidence intervals were calculated

for each of the 22 districts, utilizing the iNEXT.3D package (Chao *et al.* 2022) and the included

bootstrapping approach with 200 replicates. γ-diversity (at the landscape level) was

decomposed into α (patch level) and multiplicative β-diversity using iNEXT.beta3D (version

2.0.8) (Chao et al., 2023). We applied the 1-S transformation, a normalized dissimilarity

measure equivalent to Jaccard-type turnover, mapping β-diversity to the range of 0 indicating

identical species composition among patches to 1 representing completely disjunct

communities. Standardization also controls for variation in patch numbers of sites (9 and 15)

(Chao *et al.* 2019). For comparison between ESBC and control districts, we calculated

differences in diversity metrics (TD, PD, and FD in combination with α-, β-, and γ-diversity,

as well as with q = 0, 1, and 2) for each of the 11 district pairs with positive values indicating

higher diversity in the ESBC district. Differences from the 11 forest pairs were weighted and

combined   by   meta-analysis   (R-package   in   preparation:

https://github.com/AnneChao/iNEXT.meta)   propagating   district-wise   bootstrapping

confidence intervals to the aggregated metric and being presented as error bars in Fig. 3.

Confidence intervals that exclude the value 0 indicate significant differences between ESBC

and control districts. To control for variation in sampling effort eventually correlating with the

treatment, we standardized diversity estimates to a common coverage level of 0.95 at the

maximum of the observed distribution (Chao *et al.* 2020). District-specific results are shown in

the Supplement, figures S1-S27.

β-*deviation*

To compute pairwise β-diversity indices (distance matrices) as a measure of dissimilarity for

all study plots, we utilized the function iNEXTbeta3D_pair3D adapted from the iNEXT.beta3D

package. This step was repeated for a randomized data set generated from the observed data as

a reference null model. For each site, all observed individuals were shuffled between the patches

of the site and thus randomly redistributed within the ESBC and control district of a site (Kraft

et al. 2015; Mori *et al.* 2015b), preserving γ-diversity and the relative abundance of each species

in a site. β-deviation was calculated as the difference of β-diversities between observed and

randomized data. Increasing positive β-deviation indicates a greater difference between the

higher observed dissimilarity and the lower expected dissimilarity of the null model, suggesting

that patches differ more in their species composition than expected under random community

assembly.

Linear mixed model

To model the β-deviation for all nine facets of diversity (taxonomic, phylogenetic, and

functional diversity in combination with Hill numbers q = 0, 1, and 2) we used the pairwise

differences in forest structure, abiotic conditions, and spatial distance as fixed factors. We

estimated the contribution of each predictor treatment specific (ESBC and control district).

Using this in a single model allows a direct comparison of the effect sizes for each treatment.

For testing for differences in the slopes we also present the full interaction model in the

Appendix (Tab. S1). The final models were fitted using the function *lmer* of the `lme4´ package

(Bates et al. 2015). Rather than assuming that pairwise distances are independent observations

(as in conventional fixed-effects models), we employed site as a random-effect to capture both

within-district and among-district dependence (Harrison *et al.* 2018; Chao *et al.* 2024). We

reported z-values as a standardized metrics of effect sizes.

To additionally analyse when functional diversity patterns reveal metacommunity assembly

processes, we examined functional β-diversity independently from the changes in taxonomic

diversity by using another null model approach. For 100 replicates of the null model, we

permuted species labels across the species trait matrix, while the species abundance matrix and

thus its spatial pattern was kept unchanged (Jeliazkov & Chase 2024). We calculated the

standardized effect size (SES) normalizing observed functional β-diversity by mean and

standard deviation of the randomized samples. Positive SES were considered to indicate higher

functional turnover relative to taxonomic turnover, whereas a SES lower than 0 indicated lower

functional turnover. Absolute differences greater than 1.96 indicate significant effects. Finally,

we compared SES levels in control and ESBC districts by a GLMM again with site as a random

factor.

Results

Diversity

Our final data set comprised 1308 beetle species (37 - 143 per patch) in 103,714 specimens (153 - 1888 per patch). The meta-analyses showed the most pronounced effect size in effective species number for γ-diversity in ESBC districts by an increase of about 60 species compared to control districts when focusing on rare species (q=0, Fig. 3). This was driven by both an increase in α-diversity of about 30 species and of β-diversity of about 10%. The effective number of common species (q=1) also increased at the landscape scale, but by 6 species only, again driven by α- and β-diversity. For dominant species (q=2), we found no effect on γ-diversity, but there was a small increase in α-diversity (0.8 species). Except for q=2 β-diversity of functional and phylogenetic diversities (effective numbers of functional groups and lineages) showed lower responses to ESBC forests than taxonomic diversity, which is an indication of functional and phylogenetic redundancy. Here the effects were most pronounced for β-diversity. From rare to dominant species, the increase in functional and phylogenetic β-diversity differences between ESBC and control districts turned into a significantly negative effect. Underlying district-specific results including local diversity estimates are presented in Supplement, Fig. S1-S27.

[[Figure 3 about here]]

β-*deviation*

Modelling of the β-deviation for all Hill numbers and facets of diversity showed strong positive effects of the manipulated differences between patches via forest structure in ESBC districts, for all Hill numbers of taxonomic diversity (Table 1). Space was significant here only for common species (q=1). The z-values of structure versus space were much larger in the ESBC

districts, indicating the dominance of a *species-sorting* effect in the ESBC patches. In the control districts rare and common species were also affected by structural differences. However, in common and dominant species space was the most relevant predictor. Differences in abiotic conditions, using Ellenberg indicator values as a proxy for unmeasured spatial abiotic heterogeneity, did not significantly contribute to explaining taxonomic β-deviation, suggesting that most of the environmental heterogeneity was already captured by the experimentally manipulated and measured variables. Conditional R squared values were highest for taxonomic diversity (Tab. 1).

The β-deviation of functional and phylogenetic diversity revealed similar patterns. In the treatment districts the structural heterogeneity increased functional/phylogenetic diversity of rare species, additionally supporting the *species-sorting* effect due to differences in species' attributes. In homogeneous control districts, however, functional and phylogenetic diversity of common and dominant species increased with increasing spatial distance, which supports the *patch-dynamics* archetype with species showing differences in dispersal and competition. Phylogenetic β-deviation in heterogeneous forests was the only one which increased for rare species with difference in abiotic conditions between patches, which supports the view that the gain in phylogenetic diversity for rare species (Fig. 3) is promoted not only by pure light or deadwood as quantified for the structure, but also for more subtle gradients, e.g. of moisture. Standardized effect sizes of mean pairwise distances in functional β-diversity significantly decreased from control districts to ESBC districts for common and rare species (Supplement, Table S3) indicating a transition towards metacommunities more affected by habitat filtering even when controlling for taxonomic diversity.

[[Table 1 about here]]

Discussion

Our results supported our first hypothesis that heterogenization of a homogeneous temperate production forest increases the diversity of beetles but only with focus on rare and common species, while dominant species remained unaffected or were even disfavoured. The finding that the increase in γ-diversity was determined by an increase in α- and β-diversity confirmed the predictions that both *mass-effects* and *species-sorting* contribute to the higher γ-diversity. This was also supported by the increase in functional or phylogenetic diversity again on the α-scale. Furthermore, the consistent result of larger effects of space on β-deviation in homogeneous forests shifting to reduced spatial effects, but an increase of forest structure effects in heterogeneous forests, supported that ESBC forests are more driven by *species-sorting* processes rather than *patch-dynamics*. This could particularly be observed for functional and phylogenetic diversity which reflect species differences in habitat requirements and mobility and thus are linked to *species-sorting*.

Our study implements the concluding suggestion of Jeliazkov and Chase (2024) to consider scale-explicit approaches depending on disturbance, scales, and decomposition into α-, γ-, and specifically β-components for improving metacommunity analysis. However, a particular problem in β-diversity analyses has been observed for many years. In observational data a fraction of β-response can be regularly attributed to locally varying unobserved species resulting in artifacts of the analysis which distort the results (Tuomisto 2010). This issue is even more pronounced in experimental approaches with substantial, independent variation in predictors where sample coverage has been shown to systematically correlate with treatment (Kortmann *et al.* 2025b; Rothacher *et al.* 2025) which is increasingly relevant in highly diverse taxa with many unobserved species, as in beetles. It does not only matter in species-rich and structurally extremely heterogeneous tropical forests, but also in European temperate zones. Recent analyses showed that many studies with insect samples from temperate zones are

incomplete, in cases using flight and pitfall traps in beetles, light trapping of moths or even for

total insect data from malaise traps (Müller & Brandl 2009; Kortmann *et al.* 2025a; Püls *et al.*

2025). In addition, it especially affects the conclusions from null model approaches which are

evaluating the relative role of different assembly processes in a spatially hierarchical setting

(Mori *et al.* 2015a). Tucker et al. (2016) demonstrated this effect of a higher β-diversity in

undersampled communities with simulated data and suggested developing methods to control

for it. This has been solved recently by standardizing not only diversity by the comprehensive

approach of attribute diversity, but also dissimilarity between communities, by controlling for

sample coverage (Chao *et al.* 2023). In contrast to the widely used Bray-Curtis-dissimilarity the

generalized concept is applicable even to dissimilarities along Hill numbers and enables a much

wider range of β-diversity applications. Moreover, this unified approach for taxonomic,

functional, and phylogenetic diversity, does not only allow direct comparisons of the different

units, but also to generalize the β-diversity concept to the different aspects of species

dissimilarity as previously proposed by Logue *et al.* (2011). Finally, the difference is not only

reflected in traits. Besides species identity, species abundance substantially affects population

dynamics and community assembly. Larger populations of dominant species have, for instance,

higher dispersal potential than smaller populations and are less affected in survival by negative

stochastic effects even if sometimes most affected in general (see for insects in Müller *et al.*

2024; van Klink *et al.* 2024). The use of Hill numbers in our study provides a generalized

framework to investigate metacommunity patterns from different angles without any additional

information about species attributes (see also Guzman *et al.* 2022).

Patch-dynamics archetype

The homogeneous forests are mostly in accordance with metacommunities dominated by *patch*

dynamics (Levins & Culver 1971; Leibold *et al.* 2004). By centuries of human management,

they are rather homogeneous in terms of tree dimension, vertical and horizontal structure. The

patches within such a managed forest are rather similar, and we can expect community differences mainly to be explained by distance (Holyoak *et al.* 2005). Metacommunities in homogeneous environments are determined by differences in species ability to disperse, to colonize and to outcompete other species. Beetles show a wide range in the dispersal ability from flightless species to species dispersing actively over many kilometres (Buse 2012; Komonen & Müller 2018). Even though the maximum distance between two patches within a district was only ~1km, which might be bridged by all species in our data set (Komonen & Müller 2018), we found a clear spatial effect independent from environmental differences in both homogeneous and heterogeneous forests. This was supported even for the direct measures of functional and phylogenetic β-deviation. *Patch dynamics* occurring in both forest types but being more pronounced in homogeneous forests suggests that this mechanism always contributes to metacommunities of beetles in forests but dominates in homogeneous ones. Even if we cannot compare directly, β-deviation analyses of tree species in continuous tropical old-growth forests also revealed a predominant effect of space (Myers *et al.* 2013).

The beetles in our data do not only differ in their abundance and mobility (from flightless to high mobile), but also in their ability to outcompete other species after colonization. One prominent example is Ambrosia beetles. They have evolved highly flexible life history strategies during the colonization of new patches, e.g. by sib-mating without inbreeding depression, changing the level of social behaviour, and adopting a complex mutualism with their ambrosia fungi (Biedermann 2010; Biedermann *et al.* 2011). In contrast, other beetle species are less flexible in traits resulting in a gradient of species' location on the trade-off association between dispersal ability and competitive strength, which is required for *patch-dynamics* (Logue *et al.* 2011).

Species-sorting

MacArthur and MacArthur (1961) laid the cornerstone for the habitat heterogeneity hypothesis using the gradient of vertical heterogeneity of forests as an axis of different niches for bird species. Later this single gradient of habitat heterogeneity has been expanded to a number of biotic and abiotic heterogeneity dimensions in forests with horizontal heterogeneity as one of the overall most prominent ones among taxa and functions (Heidrich *et al.* 2020). In this light, our substantial manipulation of forest structure, including the horizontal heterogeneity and amount of deadwood should end up in species-sorting (Leibold *et al.* 2004; Van der Gucht *et al.* 2007; Hernandez-Ordonez *et al.* 2019). We expected this especially, because we manipulated deadwood, which promotes many obligatory, but also facultative saproxylic arthropod and particularly beetle species (Seibold *et al.* 2016; Graf *et al.* 2022). In fact, the increase in β-diversity and β-deviation with increasing forest structure differences, supports this at least for rare and common species. The question remains, why we could not find an effect on diversity of dominant species. Here we must note that our local interventions did not change the structure of deadwood and light availability of large, forested landscapes. So, the majority of the forest remained even-aged, closed-canopy, and homogeneous making specific species the dominant species with high abundancies resistant to the ESBC treatment. However, dominant species can be affected by habitat heterogeneity under specific conditions. An analysis of the rarity level of the same forest beetles in Europe in response to wilderness areas revealed that rare species in Europe were dominant in the wilderness areas of Mongolia and vice versa (Müller *et al.* 2013). Similarly, we have reports of single species, e.g., the highly endangered *Peltis grossa* (Trogossitidae) formerly very rare due to modern forestry over centuries, turning into a dominant species after landscape-wide pulses of deadwood in large open forests after windstorm and bark beetle disturbances (Busse *et al.* 2022). Together with

our findings, this shows that our patch-wide interventions were insufficient to affect the *species-*

sorting of dominant species in production forests of the temperate European zone.

Mass-effect

The second archetype commonly stressed in metacommunity theory with increasing

heterogeneity is the *mass-effect* stimulating source-sink dynamics (Mouquet & Loreau 2003;

Leibold *et al.* 2004). It has been stated that without information of dispersal rates or frequencies

it cannot be distinguished from the *species-sorting* effect (Logue *et al.* 2011). However, our

comprehensive approach on different spatial scales provides some support that the *mass-effect*

is a critical mechanism providing higher γ-diversity in ESBC forests. We found a clear effect

of the treatment on α-diversity. This indicates that the gain of the resources deadwood and

sunlight in the former homogeneous, dark forest locally increases the diversity of beetles.

Considering that spatial distance additionally affected the β-deviation beyond the pronounced

effects of forest structure suggests that at least the increase in common species is also triggered

by *mass-effects*. In addition, it could be responsible for the slightly negative effect in β-diversity

in dominant species (Fig. 3), when massive dispersal of the most frequent species is promoted

eventually resulting in homogenization of species distribution (Mouquet & Loreau 2002;

Pardini *et al.* 2010). We could not identify such a spatial effect on β-deviation for rare species,

but we found a strong effect on α-diversity of rare species. This might be explained by the fact

that many rare species are promoted by resource pulses at the α-scale, but as they are too rare,

this does not display in β-deviations. Nevertheless, the large contribution of α-diversity to

higher γ-diversity in ESBC districts cannot be explained only by β-diversity suggesting an

overall *mass-effect*.

Facets of diversity

For maximum significance of functional diversity Jeliazkov and Chase (2024) suggested the

selection of traits related to reproduction and mobility. Thus, we combined body size and

feeding strategy with species response to light (brightness level) and to deadwood (saproxylic

guild) which have been proven to be sensitive for environmental gradients as manipulated in

our experiment (e.g. Gossner *et al.* 2013). Like Jeliazkov & Chase (2024), we found that in

general diversity patterns were explained best for taxonomic diversity. We share their sceptical

perspective on the potential explanatory contribution of functional diversity in addition to

taxonomic diversity for rare species as SES do not indicate trait divergence or convergence

when controlled for taxonomic diversity. However, for common and dominant species mostly

positive SES values hint on trait divergence. In contrast to Jeliazkov and Chase (2024) we based

our SES calculation on the multiplicative Hill number approach for diversity decomposition

(Chao *et al.* 2023) which allows to differentiate the perspective between rare and common

species in addition to dominant species. When considering not only the total explained variance

of a model, but also the relevance of individual predictors in comparison to each other, nuanced

insights can be gained from the differences between PD and FD as well as between the Hill

numbers. For FD and PD forest structure affects mainly rare species indicating *species-sorting*,

while the impact of space prevails in common and dominant species where successful dispersal

can be triggered by abundance and constitute a *mass-effect*. Despite the general correlation

between effect sizes in PD and FD, there are also deviating cases, e.g., in rare species and the

impact of space in ESBC habitat, eventually indicating opposing effects of different traits

aggregated in functional diversity.

From mechanisms to management strategies

At the midpoint of the UN decade of restoration, success is still limited. From a forest

management perspective, increasingly frequent natural disturbances are used as guidelines for

new strategies aiming to improve resistance and biodiversity of forests (Aszalós *et al.* 2022).

Many of these strategies aim to locally increase the number of old-growth attributes, such as

trees with microhabitats or deadwood (Bauhus *et al.* 2009) or to mimic natural disturbance

events (Seibold *et al.* 2016). However, natural events provide no standardized design, are most

often not replicated, and spatially not independent. These factors prevent the necessary

comprehensive understanding of the effects on diversity and the essential mechanisms of

metacommunity theory. In addition, they increase the risk of misinterpretation and might even

contribute to establish unproven paradigms (Weisser *et al.* 2023). In contrast to long lasting

expectations two large global assessments recently found that land use intensity does not

consistently form homogeneous communities, and that higher γ-diversity is often driven by

increasing α-diversity, while effects of β-diversity were less consistent (Gonçalves-Souza &

Sanders 2025; Keck *et al.* 2025). While the statistical tools to identify mechanisms along the

axes of different species, habitat difference, and dispersal seem to be available now, more

replicated studies in which homogeneous landscapes are experimentally heterogenized are

necessary to decouple potential drivers and improve the understanding of decline or restoration

success particularly for invertebrates

Outlook

Taking beetles as a highly diverse model group, we derive several important implications for

restoration of homogeneous temperate production forests. First, canopy openings by gap

felling, group wise dieback of trees by insects, summer storms, or droughts and the

enhancement of deadwood do not affect the diversity of dominant beetles in temperate forests.

Second, all local resource pulses by light and deadwood increase not only the α-diversity but

also translates into a higher γ-diversity of rare and common beetle species, probably by *mass-*

effects. Additionally, increasing the between-patch heterogeneity also increases γ-diversity of

rare and common species most probably by *species-sorting* effects. Therefore, both strategies,

an opportunistic increase of resources and a targeted spatial heterogeneity in forests seem very

promising in restoring the beetle diversity of homogeneous production forests, which could be

extended to other taxa.

Acknowledgements and funding

We thank Elisa Stengel, Luisa Pflumm, Thorben Riehe, Michael Junginger, Boris Büche,

Alexander Szallies, Ruth Pickert, Sophia Hochrein, Katharina Kallnik, Marina Wolz, Manuel

Mauermann, Bernd Anell, Jens Geigenmüller, Hans Bahr, Interns of the Bavarian Forest

Nationalpark and all other assistants in the field and laboratory. We also thank the entire BETA-

FOR team for their support. This work was supported by the Deutsche Forschungsgemeinschaft

[BETA-FOR, 459717468]. J.R. received funding from the Bavarian State Ministry for Food,

Agriculture, Forestry and Tourism (grant no. L062).

References

Abrego, N., Furneaux, B., Hardwick, B., Somervuo, P., Palorinne, I., Aguilar-Trigueros, C.A.

et al. (2024). Airborne DNA reveals predictable spatial and seasonal dynamics of fungi.

Nature, 631, 835-842.

Aszalós, R., Thom, D., Aakala, T., Angelstam, P., Brūmelis, G., Gálhidy, L. et al. (2022).

Natural disturbance regimes as a guide for sustainable forest management in Europe.

Ecological Applications, 32, e2596.

Bauhus, J., Puettmann, K. & Messier, C. (2009). Silviculture for old-growth attributes. Forest

Ecology and Management, 258, 525-537.

Biedermann, P.H.W. (2010). Observations on sex ratio and behavior of males in *Xyleborinus*

saxesenii Ratzeburg (Scolytinae, Coleoptera). ZooKeys, 253-267.

Biedermann, P.H.W., Klepzig, K.D. & Taborsky, M. (2011). Costs of delayed dispersal and

alloparental care in the fungus-cultivating ambrosia beetle *Xyleborus affinis* Eichhoff

(Scolytinae: Curculionidae). Behavioral Ecology and Sociobiology, 65, 1753-1761.

Buse, J. (2012). "Ghosts of the past": flightless saproxylic weevils (Coleoptera:

Curculionidae) are relict species in ancient woodlands. J. Insect Conserv., 16, 93-102.

Busse, A., Cizek, L., Pavla Čížková, Drag, L., Dvorak, V., Heurich, M. et al. (2022). Forest

dieback in a protected area triggers the return of the primeval forest specialist *Peltis grossa*

(Coleoptera, Trogossitidae). Conservation Science and Practice, 4, e612.

Chao, A., Chiu, C.-H., Hu, K.-H., van Der Plas, F., Cadotte, M.W., Mitesser, O. et al. (2024).

Hill-Chao numbers allow decomposing gamma multifunctionality into alpha and beta

components. Ecology Letters, 27, e14336.

Chao, A., Henderson, P.A., Chiu, C.H., Moyes, F., Hu, K.H., Dornelas, M. et al. (2021).

Measuring temporal change in alpha diversity: A framework integrating taxonomic,

phylogenetic and functional diversity and the iNEXT.3D standardization (Jul, 10.1111/2041-

210X.13682, 2022). Methods in Ecology and Evolution, 12, 1926-1940.

Chao, A., Kubota, Y., Zeleny, D., Chiu, C.-H., Li, C.-F., Kusumoto, B. et al. (2020).

Quantifying sample completeness and comparing diversities among assemblages. Ecological

Research, 35, 292-314.

Chao, A., Chiu, C.H., Wu, S.H., Huang, C.L. & Lin, Y.C. (2019). Comparing two classes of

alpha diversities and their corresponding beta and (dis)similarity measures, with an

application to the Formosan sika deer Cervus nippon taiouanus reintroduction programme.

Methods in Ecology and Evolution, 10, 1286-1297.

Chao, A., Thorn, S., Chiu, C.H., Moyes, F., Hu, K.H., Chazdon, R.L. et al. (2023).

Rarefaction and extrapolation with beta diversity under a framework of Hill numbers: The

iNEXT.beta3D standardization. Ecological Monographs, 93, e1588.

Chown, S.L. & Gaston, K.J. (2010). Body size variation in insects: a macroecological

perspective. Biological Reviews, 85, 139-169.

Cottenie, K. (2005). Integrating environmental and spatial processes in ecological community

dynamics. Ecology Letters, 8, 1175-1182.

Dittrich, S., Hauck, M., Jacob, M., Rommerskirchen, A. & Leuschner, C. (2013): Response of

ground vegetation and epiphyte diversity to natural age dynamics in a Central European

mountain spruce forest. J. Veg. Sci. 24: 675–687.

Ellenberg, H., Weber, H.E., Düll, R., Wirth, V., Werner, W. & Paulißen, D. (1991).

Zeigerwerte von Pflanzen in Mitteleuropa. Goltze Verlag, Göttingen.

Ewald, J. (2003). The sensitivity of Ellenberg indicator values to the completeness of

vegetation releves. Basic and Applied Ecology, 4, 507-513.

Farrell, B.D. (1998). "Inordinate Fondness" Explained: Why Are There So Many Beetles?

Science, 281, 555–559.

Freude, H., Harde, K. & Lohse, G.A. (1964-1983). Die Käfer Mitteleuropas. Goecke & Evers,

Krefeld.

Gonçalves-Souza, T. & Sanders, N.J. (2025). Species turnover does not rescue biodiversity in

fragmented landscapes. Nature, 640, 702-706.

Gonzalez, A. (2009). Metacommunities: Spatial Community Ecology. In: eLS.

Gossner, M.M., Lachat, T., Brunet, J., Isacsson, G., Bouget, C., Brustel, H. et al. (2013).

Current "near-to-nature" forest management effects on functional trait composition of

saproxylic beetles in beech forests. Conserv. Biol., 27, 605-614.

Graf, M., Seibold, S., Gossner, M.M., Hagge, J., Weiss, I., Baessler, C. et al. (2022).

Coverage based diversity estimates of facultative saproxylic species highlight the importance

of deadwood for biodiversity. Forest Ecology and Management, 517, 120275.

Guzman, L.M., Thompson, P.L., Viana, D.S., Vanschoenwinkel, B., Horvath, Z., Ptacnik, R.

et al. (2022). Accounting for temporal change in multiple biodiversity patterns improves the

inference of metacommunity processes. Ecology, 103, e3683.

Hagge, J., Müller, J., Birkemoe, T., Buse, J., Christensen, R.H.B., Gossner, M.M. et al.

(2021). What does a threatened saproxylic beetle look like? Modelling extinction risk using a

new morphological trait database. Journal of Animal Ecology, 90, 1934-1947.

Harrison, X.A., Donaldson, L., Correa-Cano, M.E., Evans, J., Fisher, D.N., Goodwin, C.E. et

al. (2018). A brief introduction to mixed effects modelling and multi-model inference in

ecology. PeerJ, 6, e4794.

Heidrich, L., Bae, S., Levick, S., Seibold, S., Weisser, W., Krzystek, P. et al. (2020).

Heterogeneity-diversity relationships differ between and within trophic levels in temperate

forests. Nature Ecology & Evolution, 4, 1431-1431.

Hernandez-Ordonez, O., Santos, B.A., Alexander Pyron, R., Arroyo-Rodriguez, V., Nicolas

Urbina-Cardona, J., Martinez-Ramos, M. et al. (2019). Species sorting and mass effect along

forest succession: Evidence from taxonomic, functional, and phylogenetic diversity of

amphibian communities. Ecology and Evolution, 9, 5206-5218.

Holyoak, M., Leibold, M.A., Mouquet, N., Holt, R.D. & Hoopes, M.F. (2005).

Metacommunities - A framework for large-scale community ecology. Metacommunities:

Spatial Dynamics and Ecological Communities, 1-31.

Hubbell, S.P. (2001). The Unified Neutral Theory of Biodiversity and Biogeography.

Princeton University Press, Princeton.

Jeliazkov, A. & Chase, J.M. (2024). When Do Traits Tell More Than Species about a

Metacommunity? A Synthesis across Ecosystems and Scales. American Naturalist, 203, E1-

E18.

Keck, F., Peller, T., Alther, R., Barouillet, C., Blackman, R., Capo, E. et al. (2025). The

global human impact on biodiversity. Nature, 641, 395-400.

Koch, K. (1989-1992). Die Käfer Mitteleuropas. Ökologie, Bd. 2-3. Goecke & Evers, Krefeld.

Komonen, A. & Müller, J. (2018). Dispersal ecology of dead wood organisms: implications

for connectivity conservation. Conserv. Biol., 32, 535–545.

Kortmann, M., Chao, A., Chiu, C.-H., Heibl, C., Mitesser, O., Morinière, J. et al. (2025a). A

short cut to sample coverage standardization in meta-barcoding data provides new insights

into land use effects on insect diversity. Proc R Soc Lond B Biol Sci, 292, 20242927.

Kortmann, M., Chao, A., Schaefer, H.M., Blüthgen, N., Gelis, R., Tremlett, C.J. et al.

(2025b). Sample coverage affects diversity measures of bird communities along a natural

recovery gradient of abandoned agriculture in tropical lowland forests. Journal of Applied

Ecology, 62, 480-491.

Kraft, N.J.B., Adler, P.B., Godoy, O., James, E.C., Fuller, S. & Levine, J.M. (2015).

Community assembly, coexistence and the environmental filtering metaphor. Functional

Ecology, 29, 592-599.

Leibold, M.A. & Chase, J.M. (2018). Metacommunity Ecology. In: Monographs in

Population Biology (eds. Levin, SA & Horn, HS). Princeton University Press Woodstock, p.

491.

Leibold, M.A., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J.M., Hoopes, M.F. et al.

(2004). The metacommunity concept: a framework for multi-scale community ecology.

Ecology Letters, 7, 601-613.

Levins, R. & Culver, D. (1971). Regional coexistence of species and competition between

rare species. Proc. Natl. Acad. Sci. U. S. A., 68, 1246-1248.

Logue, J.B., Mouquet, N., Peter, H., Hillebrand, H. & The Metacommunity Working Group

(2011). Empirical approaches to metacommunities: a review and comparison with theory.

Trends in Ecology & Evolution, 26, 482-491.

MacArthur, R.H. & MacArthur, J. (1961). On bird species diversity. Ecology, 42, 594-598.

Mori, A.S., Fujii, S., Kitagawa, R. & Koide, D. (2015a). Null model approaches to evaluating

the relative role of different assembly processes in shaping ecological communities.

Oecologia, 178, 261-273.

Mori, A.S., Ota, A.T., Fujii, S., Seino, T., Kabeya, D., Okamoto, T. et al. (2015b). Biotic

homogenization and differentiation of soil faunal communities in the production forest

landscape: taxonomic and functional perspectives. Oecologia, 177, 533-544.

Mouquet, N. & Loreau, M. (2002). Coexistence in metacommunities: The regional similarity

hypothesis. American Naturalist, 159, 420-426.

Mouquet, N. & Loreau, M. (2003). Community patterns in source-sink metacommunities.

American Naturalist, 162, 544-557.

Müller, J., Bässler, C., Strätz, C., Klöcking, B. & Brandl, R. (2009). Molluscs and climate

warming in a low mountain range national park. Malacologia, 51, 133-153.

Müller, J. & Brandl, R. (2009). Assessing biodiversity by remote sensing and ground survey

in montainous terrain: the potential of LiDAR to predict forest beetle assemblages. Journal of

Applied Ecology, 46, 897–905.

Müller, J., Hothorn, T., Yuan, Y., Seibold, S., Mitesser, O., Rothacher, J. et al. (2024).

Weather explains the decline and rise of insect biomass over 34 years. Nature, 628, 349-354.

Müller, J., Jarzabek-Müller, A. & Bussler, H. (2013). Some of the rarest European saproxylic

beetles are common in the wilderness of Northern Mongolia. J. Insect Conserv., 17, 989-1001.

Müller, J., Mitesser, O., Cadotte, M.W., van der Plas, F., Mori, A.S., Ammer, C. et al. (2023).

Enhancing the structural diversity between forest patches-A concept and real-world

experiment to study biodiversity, multifunctionality and forest resilience across spatial scales.

Global Change Biology, 29, 1437-1450.

Myers, J.A., Chase, J.M., Jimenez, I., Jorgensen, P.M., Araujo-Murakami, A., Paniagua-

Zambrana, N. et al. (2013). Beta-diversity in temperate and tropical forests reflects dissimilar

mechanisms of community assembly. Ecology Letters, 16, 151-157.

Pardini, R., Bueno, A.d.A., Gardner, T.A., Prado, P.I. & Metzger, J.P. (2010). Beyond the

Fragmentation Threshold Hypothesis: Regime Shifts in Biodiversity Across Fragmented

Landscapes. PLoS One, 5, e13666.

Pierick, K., Link, R. M., Inderwies, H. L., Puggaard Koolen, S., Wu, B. Schuldt, B. et al.

(2025, preprint) Artificial Gap Creation But Not Thinning Interventions Persistently Enhance

Forest Structural Heterogeneity. http://dx.doi.org/10.2139/ssrn.5157921

Püls, M., Achury, R., Heidrich, L., Seibold, S., Hacker, H., Mitesser, O. et al. (2025).

Functional and phylogenetic beta diversity response of nocturnal moth assemblages to land-

use intensity in grasslands and forests. Oikos, e11426.

Record, S., Voelker, N.M., Zarnetske, P.L., Wisnoski, N.I., Tonkin, J.D., Swan, C. et al.

(2021). Novel Insights to Be Gained From Applying Metacommunity Theory to Long-Term,

Spatially Replicated Biodiversity Data. Frontiers in Ecology and Evolution, 8, 612794.

Rothacher, J., Seidl, R., Thom, D., Kortmann, M., Chao, A., Chiu, C.-H. et al. (2025). The

impact of tree mortality and post-disturbance management on insect diversity in temperate

forests: Insights from a replicated experiment. Journal of Applied Ecology, 62 1878-1888.

Seibold, S., Bassler, C., Baldrian, P., Reinhard, L., Thorn, S., Ulyshen, M.D. et al. (2016).

Dead-wood addition promotes non-saproxylic epigeal arthropods but effects are mediated by

canopy openness. Biological Conservation, 204, 181-188.

Seibold, S., Brandl, R., Buse, J., Hothorn, T., Schmidl, J., Thorn, S. et al. (2015). Association

of extinction risk of saproxylic beetles with ecological degradation of forests in Europe.

Conserv. Biol., 29, 382-390.

Senf, C., Pflugmacher, D., Yang, Z.Q., Sebald, J., Knorn, J., Neumann, M. et al. (2018).

Canopy mortality has doubled in Europe's temperate forests over the last three decades.

Nature Communications, 9, 4978.

Thom, D. & Seidl, R. (2016). Natural disturbance impacts on ecosystem services and

biodiversity in temperate and boreal forests. Biological Reviews, 91, 760-781.

Thompson, P.L., Guzman, L.M., De Meester, L., Horvath, Z., Ptacnik, R., Vanschoenwinkel,

B. et al. (2020). A process-based metacommunity framework linking local and regional scale

community ecology. Ecology Letters, 23, 1314-1329.

Tucker, C.M., Shoemaker, L.G., Davies, K.F., Nemergut, D.R. & Melbourne, B.A. (2016).

Differentiating between niche and neutral assembly in metacommunities using null models of

β-diversity. Oikos, 125, 778-789.

Tuomisto, H. (2010). A diversity of beta diversities: straightening up a concept gone awry.

Part 2. Quantifying beta diversity and related phenomena. Ecography, 33, 23-45.

Van der Gucht, K., Cottenie, K., Muylaert, K., Vloemans, N., Cousin, S., Declerck, S. et al.

(2007). The power of species sorting: Local factors drive bacterial community composition

over a wide range of spatial scales. Proc. Natl. Acad. Sci. U. S. A., 104, 20404-20409.

van Klink, R., Bowler, D.E., Gongalsky, K.B., Shen, M., Swengel, S.R. & Chase, J.M.

(2024). Disproportionate declines of formerly abundant species underlie insect loss. Nature,

628, 359–364.

Weisser, W., Blüthgen, N., Staab, M., Achury, R. & Müller, J. (2023). Experiments are

needed to quantify the main causes of insect decline. Biology Letters, 19, 20220500.

Zeuss, D., Brandl, R., Brandle, M., Rahbek, C. & Brunzel, S. (2014). Global warming favours

light-coloured insects in Europe. Nature Communications, 5, 3874.

Figure 1: a) Extension of Logue et al. (2011)'s scheme of metacommunity mechanisms based on the dimensions of heterogeneity, functional equivalence of individuals, and dispersal. Metacommunity archetypes of *species-sorting*, *mass-effect*, *patch-dynamics*, and *neutral processes* occupy typical areas in the diagram. Both meta-analyses (yellow) and β-deviation analysis (blue) allow discrimination between the processes. b) Conceptual figure on archetypes from metacommunity theory explaining diversity and assembly mechanisms in local beetle communities in homogeneous and heterogenous forests. White and green colours indicate habitat differences. In ESBC patches light and dark beetle symbols indicate matching of beetles' attributes to light and dark green habitats, respectively.

Figure 2: Design of replicated manipulation of habitat heterogeneity in 11 homogeneous production

forests and their 11 control forests. A) Interventions were either aggregated in the centre of the patch

(upper row) or evenly distributed over the patch area (lower row). B) Site with treatment (ESBC:

Enhancement of Structural Beta Complexity) district and control district. C) Geographical distribution

of the 11 sites. D) Schematic representation of the ecosystem elements site, district, and patch in

relation to derived diversity levels for an exemplary site.

Figure 3: Results of the meta-analyses based on 11 pairwise comparisons of homogeneous and heterogenized forests on γ-, α-, and β-level with focus on rare (q=0), common (q=1) and dominant species (q=2) as well as on taxonomic (TD), functional (FD) and phylogenetic diversity (PD) of 1308 beetle species in 234 patches. Positive values indicate higher diversity levels in ESBC districts compared to control districts. Error bars denote the 95% confidence intervals calculated from weighted averages of bootstrapped, district-specific diversity estimates for each of the 27 cases.

Table 1: Results of linear mixed interaction models of the β-deviation, i.e. the difference between β-

diversity of observed vs. null-model data (with randomized distribution of individuals across patches in

each site) for taxonomic (TD), functional (FD), and phylogenetic (PD) β-diversity in combination with

Hill numbers q=0, 1, 2 (rare, common, and dominant species). As predictors we used the structural

difference between patches (Structure), the difference in space (Spatial), and the difference in abiotic

conditions (Abiotic) extracted from Ellenberg indicator values for both levels of deadwood and canopy

structure (Control vs. ESBC), as well as Site as random factor. Last row provides the conditional R

squared value of the model.

Diversity facet	TD	TD	TD	FD	FD	FD	PD	PD	PD
Hill number q	**0**	**1**	**2**	**0**	**1**	**2**	**0**	**1**	**2**
Control:Structure	**3.628**	**3.196**	1.579	-0.068	0.352	0.340	**2.002**	0.791	0.644
ESBC:Structure	**4.936**	**11.75**	**7.685**	**3.865**	0.509	0.495	**2.372**	1.722	-0.254
Control:Abiotic	-0.256	-1.155	-0.467	1.258	**-2.196**	-1.673	-0.471	**-2.369**	**-2.347**
ESBC:Abiotic	1.124	1.152	1.807	**-2.007**	0.530	0.790	**2.373**	1.476	1.040
Control:Spatial	0.835	**6.994**	**7.229**	0.878	**5.844**	**6.287**	0.889	**6.909**	**6.987**
ESBC:Spatial	1.538	**3.092**	1.715	0.017	1.230	1.340	**2.345**	1.313	1.220
R2 conditional	0.11	0.35	0.38	0.071	0.15	0.18	0.10	0.17	0.17