1 **Title**: Identifying Priority Stepping Stone Reefs to Maintain Global Networks of Connected
2 Coral Reefs
3

Authors: Ariel Greiner[1,2], Emily S. Darling[3,4], Amelia Wenger[3,5], Martin Krkošek[4], Marie-Josée
Fortin[4]
[1]Department of Biology, University of Oxford, Oxford, UK
[2]Department of Biology, Pennsylvania State University, University Park, PA, USA
[3]Marine Program, Wildlife Conservation Society, Bronx, New York, USA
[4]Department of Ecology and Evolutionary Biology, University of Toronto, Toronto, Ontario,
Canada
[5]Centre for Biodiversity and Conservation Science, School of the Environment, University of
Queensland, St. Lucia, Queensland, Australia
13
14 **Corresponding Author**: Ariel Greiner, greinerariel@gmail.com, Spaulding Hall - Room 201N,
15 University of New Hampshire, 38 Academic Way, Durham, NH 03824, USA

16
17 **# of Words in Full Manuscript**: 3498 words
18

19 **Keywords**: coral reef, conservation planning, marine, connectivity, metapopulations, stepping
20 stone, network, biogeography

21
22 **Article Impact Statement**: We identify 10 coral reef stepping stones in the Indo-Pacific that
23 may ensure that 84,564km^2 of coral reef refugia could remain connected.
24

25 **Acknowledgements**: We thank Marco Andrello for his helpful comments and Cole B. Brookson
26 for a helpful R function for map generation.

27
28 **Artificial Intelligence tools** were not used in the writing, analysis or formulation of this study.
29

Funding Statement: This research was supported by a Natural Sciences and Engineering
Research Council of Canada (NSERC) Postdoctoral Fellowship to A.G. and NSERC Discovery
Grants (no. 4922 to M.K. and no. 6426 to M.-J.F.) and Canada Research Chairs to M.K. and M.-
J.F. M.K. was also supported by an E.W.R. Steacie Memorial Fellowship (566203-2021).
Funding to E.S.D. and A.W. was provided by grants from the Bloomberg Ocean Initiative and
The Tiffany and Co. Foundation.

Title: Identifying Priority Stepping Stone Reefs to Maintain Global Networks of Connected
Coral Reefs
38
Abstract

Conserving coral reef climate refugia is a key conservation strategy to address climate change.
Yet, the accelerating impacts of coral bleaching and other anthropogenic pressures can
jeopardize refugia persistence. The dispersal of coral larvae can increase long-term reef
persistence through metapopulation dynamics if connected reefs can share beneficial adaptations
to bleaching and facilitate demographic recovery. Here, we modelled reef connectivity using a
network approach to identify present-day coral larval dispersal networks of predicted climate
refugia and then determined locations of 'stepping stone' reefs that can be used to connect these
global networks of refugia. We identify 10 key locations of coral reef stepping stones in
Indonesia, Mozambique, Glorioso Islands, and Malaysia that may ensure that $84{,}564\text{km}^2$ of
refugia could remain connected together even if other (non-refugia) reefs become degraded.
Global coral reef conservation efforts should consider prioritizing these stepping stones to build
more redundancy and resilience into future-proofing conservation strategies.

Introduction

Coral reefs are a vital source of sustenance and livelihood for hundreds of millions of people worldwide (Cruz-Trinidad et al., 2014; FAO, 2012; Sing Wong et al., 2022), yet they face extreme threats from numerous anthropogenic pressures. Some of these pressures are global—such as climate change, which increases the frequency and severity of coral bleaching, cyclones and ocean acidification—while others vary locally, depending on the intensity of pressures like invasive species, overfishing, industrial development, and tourism (Andrello et al., 2022; Burke et al., 2011; Smith et al., 2016; Wenger et al., 2016). Although overfishing and water pollution are among the leading local pressures globally, the dominant pressures on reefs vary across regions (Andrello et al., 2022). By 2100, an estimated 70–99% of reefs could face long-term degradation due to coral bleaching (Frieler et al., 2013; van Hooidonk et al., 2016). Under worst-case climate scenarios, the long-term persistence of coral populations may depend on maintaining connectivity among a fragmented mosaic of climate-resilient habitat fragments (a metapopulation or network of reefs), supported by effective management of local pressures.

Predicting potential climate refugia for reefs has been considered in multiple regional studies (DeCarlo, 2020; McClanahan & Azali, 2021; McManus et al., 2020), but only Beyer et al. (2018) have considered a worldwide portfolio. Beyer et al. (2018) used historic, present-day and future projected thermal conditions, cyclone wave damage, as well as emigration of coral larvae simulation data to define 83 bioclimatic units (hereafter, 'Refugia') for coral reefs. Each of the Beyer et al. (2018) Refugium contain ~500km^2 of spatially contiguous coral reef cover that is predicted to be more resilient to thermal stress and cyclones and be good sources of coral larvae for recovery (due to their spatial position) (Beyer et al., 2018).

Since both global and local anthropogenic pressures are spatially variable, coral recovery can be achieved by coral larval dispersal among a network of reefs. Coral larvae dispersing from one reef to another connect coral reefs into distinct networks ('reef networks'). Here, we define reef networks as the complete set of reef cells (18×18km geographic area containing coral cover, as defined based on the Global Distribution of Coral Reefs Dataset (UNEP-WCMC, WorldFish Centre, WRI, & TNC, 2010) and updates from the NOAA Coral Reef Watch Stations as derived from Wood et al. (2014), Beyer et al. (2018)) that send or receive coral larvae to/from each other according to a reference coral larval dispersal connectivity matrix (Greiner et al., 2022a). Using this empirically consistent reference coral larval dispersal connectivity matrix, Greiner et al. (2022a) found that many reef networks are quite large in size and span large regions (e.g. >50% of reef cells worldwide are in 6 reef networks). Thus, motivating coral reef conservation initiatives to consider these large spatial scales when determining which reefs to prioritize. In this study, we aim to identify reef cells that can act as 'stepping stones'. Stepping stones are ecosystem patches (reef cells, in this case) that provide dispersal linkages between prioritized sets of ecosystem patches (in this case, the Refugia, which each have 2-157 reef cells), connecting or enhancing the connections between these prioritized ecosystem patches (Saura et

al., 2014). Prioritizing stepping stones alongside prioritized ecosystem patches is essential for
enhancing connectivity and achieving Target 3 of the Kunming-Montreal Global Biodiversity
Framework (protecting 30% of terrestrial, inland water areas, coastal and marine areas), as well
as for maintaining ecosystem function globally (Convention on Biological Diversity, 2022).
While the stepping stone concept is general, stepping stones have mainly been determined for
terrestrial ecosystems to date (Luo et al., 2021; Saura et al., 2017; Xu et al., 2024; but see
D'Aloia et al., 2019). Identifying stepping stones for coral reefs will enable us to prioritize the
protection or restoration of locations that could enhance connectivity between Refugia and
provide spatial rescue through time (Gotelli, 1991; Mouquet & Loreau, 2003; Saura et al., 2014),
and ideally provide crucial redundancy against the degradation of Refugia reef cells.
103
In this study, we determine which reef cells outside of Refugia (non-refugia reef cells) could act
as stepping stones between Refugia to help ensure the long-term persistence of Refugia reef
cells. We identify these 'stepping stones' through a network modelling approach based on a
worldwide coral larval connectivity model (Greiner et al., 2022a). Specifically, we determine (1)
a set of non-refugia reef cells ('stepping stones') that could, if prioritized for protection, maintain
connections between Refugia reef cells when all other non-refugia reef cells are degraded; (2) the
extent to which the stepping stones maintain connections among Refugia reef cells; and (3) the
present-day anthropogenic pressure and protection status of the stepping stones.
112
Methods
Overview
We identify these stepping stones through a three step process: first, we determined the present-
day reef networks of the 83 Refugia (Fig.1a); second, we simulated the degradation of all non-
refugia reef cells by removing all non-refugia reef cells, and calculating the degraded reef
networks of the 83 Refugia (Fig.1b); third, we used simulations to identify which non-refugia
reef cells, if maintained, would restore connections between Refugia (i.e. act as 'stepping
stones'; Fig.1c). By doing so, we determined how many Refugia the stepping stones could help
maintain connections with and the present-day status of those stepping stones. We simulate the
degradation of all non-refugia reef cells to determine which non-refugia reef cells, if prioritized
for protection now, could help maintain connections among the Refugia if only the Refugia
remain. We envision that if the identified stepping stones are prioritized for protection at present,
the coral reefs within them will have a better chance at withstanding future climatic or
anthropogenic stressors than they would have had otherwise. We calculated all network
designations and the present-day status of those stepping stones using R (v 4.3.1, R Core Team
2019) and QGIS (QGIS Development Team 2021).
129
Reef Networks of the Refugia
To calculate the present-day reef networks of the 83 Refugia, we used the connectivity matrix
generated by Greiner et al. (2022a) that was derived from Wood et al. (2014). First, we used the

Andrello et al. (2022) grid cells (which used data from Beyer et al. (2018)) to determine which of
the Wood et al. (2014) grid cells (hereafter, 'reef cells') were in Refugia (hereafter, 'Refugia reef
cells'). We work at the level of reef cells because that is the spatial resolution for which we have
connectivity information for reefs worldwide from Greiner et al. (2022a). In this study, we will
use the 'reference connectivity matrix' generated in Greiner et al. (2022a) (see SM1 for more
details).
139
First, we calculated which Refugia reef cells are in which reef networks ('weak cluster' in igraph
in R; Csárdi et al. 2023) and then determined which Refugia reef cells were in the same reef
networks as other Refugia reef cells and which were not in the same reef networks as any other
Refugia reef cells (SM2-Fig.1). Note that based on how the Refugia reef cells were identified by
Beyer et al. (2018), they should export more larvae to other reef cells than the average reef cell.
However, this does not necessarily imply that a large proportion of those receiving reef cells are
other Refugia reef cells.
147
Then, we removed all of the non-refugia reef cells and recalculated the reef networks of the 83
Refugia by removing the rows and columns of the connectivity matrix that corresponded with the
non-refugia reef cells. By doing this, we determined which Refugia would remain connected to
other Refugia ('Fully Connected Refugia'), which Refugia would lose some connections with
other Refugia ('Partially Disconnected Refugia'), and which would lose all connections with
other Refugia ('Disconnected Refugia') if all the non-refugia reef cells were degraded (see SM3-
Fig.1).
155
Identifying and Characterizing Stepping Stones
To identify stepping stones, we focused on the Partially Disconnected Refugia and the
Disconnected Refugia (e.g. Refugia 1, 2, and 3 in Fig.1b; reefs in SM3-Fig.1b,c). First, we
examined non-refugia reef cells that were part of the same networks as the Disconnected Refugia
before any reef cells were removed. We then checked whether these non-refugia reef cells had
direct connections—either sending or receiving larvae, or both—with at least one reef cell in a
Disconnected Refugium (i.e. 'Disconnected Refugium *i*'). Finally, we determined whether these
non-refugia reef cells were also directly connected to a reef cell from any other Refugia that were
originally in the same network as Disconnected Refugium *i*. Any reef cell that satisfied both of
these conditions was defined as a stepping stone. We distinguished which of those stepping
stones were 'connector' stepping stones or 'sink' stepping stones, depending on whether the
stepping stones received larvae from one Refugium and sent larvae to at least one other
Refugium (connector) or received larvae from multiple Refugia but did not send larvae to any
Refugia (sink). We performed this analysis for every Disconnected Refugium. We then repeated
the process for non-refugia reef cells connected to Partially Disconnected Refugia to find
additional stepping stones.
172

Once the connector and sink stepping stones had been identified, we evaluated how adding them
back affected the connectivity of the Refugia reef networks. Specifically, we counted how many
Refugia become (1) fully re-connected with the Refugia they were connected to in the present-
day networks (i.e., become 'Fully Connected Refugia'), (2) remain or become Partially
Disconnected Refugia, and (3) remain Disconnected Refugia (SM3-Fig.2). To better indicate the
scale of these changes, we also noted the total number of Refugia reef cells and the total reef cell
area that changed category. Note that 'total reef cell area' corresponds to the total size of the reef
cells themselves and is an overestimate of the area within those cells that contains coral cover
itself. We could not calculate total coral cover itself because precise estimates of the percent
coral cover was not available for all of the Refugia reef cells. Lastly, we used data from Andrello
et al. (2022) to assess which local anthropogenic pressures were affecting each stepping stone
and the protected planet database (UNEP-WCMC and IUCN, 2024) to evaluate the protection
status of each stepping stone. From the analysis of Andrello et al. (2022), we gathered the top
anthropogenic pressure level and the cumulative anthropogenic pressure level of each stepping
stone (see Fig.4).
188
Results
Stepping Stones
We identified ten non-refugia reef cells as stepping stones (Fig.2). These ten stepping stones
were the only non-refugia reef cells that could maintain connections between Refugia following
a scenario when all other non-refugia reef cells were degraded. Six of these stepping stones are
'connector' stepping stones and four of them are 'sink' stepping stones; these ten stepping stones
are located in Indonesia, Mozambique, Glorioso Islands, and Malaysia (Fig.2).
196
Refugia Connectivity with and without Stepping Stones
When all non-refugia reef cells are degraded, the reef networks become more fragmented and
less connected (Fig.3a,b; SM3-Fig.1). In particular, the number of Refugia in multi-Refugia
networks (networks that contain Refugia reef cells from more than one Refugia) decreases from
73 (3,760 reef cells, 1,218,240km^2 total reef cell area) to 66 (3,388 reef cells, 1,097,712km^2) and
the average reef network size declines from 25,364.57km^2 to 11,548.1km^2. (median network size
is unchanged at 648km^2). Nineteen Refugia maintain all connections with other Refugia, 47
Refugia lose some of their connections with other Refugia (Partially Disconnected Refugia), and
seven Refugia lose all connections with other Refugia (Disconnected Refugia) (Fig.3b; SM3-
Fig.1).
207
When the 10 stepping stone reef cells are added back in (3240km^2, Fig.2), the average network
size increases to 12,598.73km^2 (median remains unchanged at 648km^2) and 69 Refugia are now
in multi-Refugia networks (3,501 reef cells, 1,134,324km^2) (Fig.3c; SM3-Fig.2). Specifically,
nine Refugia (294 grid cells, 95,256km^2) become more connected to other Refugia – 8 Refugia
(261 grid cells, 84,564km^2) become Fully Connected Refugia (6 Partially Disconnected to Fully
Connected Refugia, 2 Disconnected to Fully Connected Refugia) and 3 Refugia (113 grid cells,

36,612km^2) become connected to at least one other Refugia again (2 Disconnected Refugia to
Fully Connected Refugia and 1 Disconnected Refugium to a Partially Disconnected Refugium)
(SM3-Fig.2). If only consider the 6 'connector' stepping stone reef cells (1944km^2, Fig.2), the
average network size becomes 12,476.84km^2 (median remains unchanged at 648km^2) and 3
Refugia (113 grid cells, 36,612km^2) become connected to at least one other Refugium again
(Disconnected to Partially Disconnected Refugia), so the number of Refugia in multi-Refugia
networks remains the same (69 Refugia; 3,501 reef cells, 1,134,324km^2). However, with only the
6 'connector' stepping stones, no additional Refugia become Fully Connected Refugia (i.e. none
of the Partially Disconnected Refugia become Fully Connected Refugia, none of the
Disconnected Refugia become Partially Disconnected Refugia) (SM3-Fig.4).

Properties of the Stepping Stones
The stepping stones varied in cumulative anthropogenic pressure level (0-1, 1 indicating high
pressure), with the stepping stones near South Sumatra experiencing the highest cumulative
pressure levels and the stepping stones near Northern Madagascar and the Riau Islands
experiencing the lowest cumulative pressure levels (Fig.4a). Only one stepping stone is within a
protected or managed area identified by the World Database on Protected Areas (UNEP-WCMC
and IUCN, 2024; Pantai Panjang dan P.Baai nature recreation park) (Fig.4b). The stepping
stones vary in the top pressure that they experience (Andrello et al. 2022; Fig.4b).

Discussion
By prioritizing the reefs within the 10 stepping stones for conservation management, we
maintain Refugia-Refugia connections with nine Refugia that would otherwise be disconnected
with the degradation of non-refugia reef cells. In doing so, these 10 stepping stones may
indirectly enhance the resilience of reefs covering an area up to 25 times larger than the
prioritized region itself (3240km^2 vs 84,564km^2, Fig.3c). These stepping stones are especially
important because they are the only individual non-refugia reef cells capable of independently
maintaining connectivity between Refugia. This small number reflects the fact that many
Refugia remain within multi-Refugia networks even when only Refugia reef cells are intact
(Fig.3b; as suggested in Greiner et al. 2022a), indicating that the Refugia alone may already
provide some level of spatial rescue and redundancy to themselves. Note that Beyer et al. (2018)
accounted for connectivity in their scoring by prioritizing reef cells that exported many larvae to
other reef cells. Thus, it is unsurprising that the reef cells within the Refugia are well-connected
to other reef cells, but it was not clear *a priori* that they would be well-connected to reef cells in
the same or other Refugia specifically. Six of these 10 are an even higher priority for
conservation as they are connector stepping stones, and may allow larvae to flow from one
Refugia to another (Fig.2). By only prioritizing these 6, the same number (69) of Refugia are
restored to multi-Refugia networks but none of the Refugia become fully re-connected to other
Refugia (SM3-Fig.3,4). It is hopeful that seven of the 10 stepping stones have a cumulative
anthropogenic pressure score below 0.5 (Fig.4a), indicating that they may have relatively few

non-climate pressures. However, monitoring is needed to confirm the ecological integrity or
capacity of these locations to have functioning coral communities that can promote coral larval
dispersal. Only one of the stepping stone reef cells is currently designated as 'protected' (Fig.5);
notably, this is also the cell with the highest cumulative anthropogenic pressure score (Fig.4),
highlighting a fortunate alignment between protection status and conservation urgency (even if
protection is unlikely to protect against all pressures).
260
These 10 candidate stepping stones were chosen based on present-day knowledge of the location
of reefs, ocean currents and coral larval dispersal. The goal of this study was to use what we
know of global coral larval dispersal connectivity to identify possible reefs that, if individually
prioritized for protection, could help maintain the 83 Refugia (Beyer et al. 2018). We searched
for single reef cell stepping stones because a single reef cell is already quite a large region to
prioritize and we wanted to find the least additional reef area to prioritize (note that pairs of reef
cells did not re-connect any Refugia, see SM5). We focus on these 83 Refugia as they currently
guide how various reef conservation initiatives and funders (e.g., the Global Fund for Coral
Reefs, Bloomberg Ocean Initiative) prioritize reefs for protection around the world, as they are
thought to have the best chance of being resilient towards future climate impacts. That being
said, this study is meant as a starting point for this type of work and more specific analyses
should be performed in the regions containing the 10 stepping stones identified here to determine
which reefs within these reef cells are most relevant and practical to prioritize (as there are
unlikely to be 3,240km^2 of reefs within these stepping stones). In particular, it would be helpful
to collect species-specific coral larval dispersal information and perform genetic analyses on
coral from the reefs found in these 10 stepping stones (and for reefs in general) to ascertain
whether these reefs do send and/or receive larvae from reefs in reef cells of more than one
Refugia–as the data that we are using to find these stepping stones is consistent with empirical
data (see SM1 and Wood et al. 2014 and Greiner et al. 2022a for more information), has been
used to drive coral reef conservation priorities (see Beyer et al. 2018) and is the only global
connectivity matrix for coral that is available (at this time), but is not directly derived from
empirical data.
283
This study presents a methodology for identifying candidate stepping stone reef cells that can be
adapted for other at-risk ecosystems and used to guide prioritization decisions when connectivity
is a management goal. For example, this methodology could be used to identify key stepping-
stone reefs at regional scales. It could also be used for other at-risk marine ecosystems whose
key organisms disperse passively along ocean currents (e.g. rocky reefs). One could also
calculate candidate stepping stone reef cells that focus on reef fish larval dispersal networks; it
would be interesting to determine whether similar reef cells would be prioritized and how well
connected the Refugia are from a reef fish perspective. Lastly, it would be interesting to assess
whether any of these stepping stones may also be likely to send macroalgal gametes between
Refugia, as macroalgal gamete dispersal may counteract the coral larval dispersal and tip reefs in

these Refugia to a degraded, less coral-dominated state (see Greiner et al., 2022b). The risk of
reefs tipping into a degraded, less coral-dominated state may become more relevant as coral reefs
become more disconnected, as dispersal between reefs has been shown to stabilize reefs (see
Greiner et al. 2025). In general macroalgal gametes are not thought to disperse as far as coral
larvae (Deysher and Norton, 1981; Mumby, 2006; Stiger and Payri, 1999), but long distance
dispersal of macroalgal gametes along the same ocean currents that transport coral larvae is
possible and ensuring that the reefs in those reef cells maintain low macroalgal cover might be
crucial for preserving high coral cover in connected reefs. Comparing which reef cells are
identified as crucial stepping stones for all of these different taxa could further help target coral
reef prioritization efforts around the world and allow us to see which reefs may be most efficient
to prioritize to satisfy multiple different conservation priorities.
305

Conclusion
In this study, we identify 10 stepping stones capable of maintaining connections among Refugia
reef cells. The reefs within these stepping stones are exposed to a range of anthropogenic
pressures, and only one currently holds protected status. The reefs within these 10 stepping
stones should be prioritized for conservation and local management before further degradation
occurs, as our study identifies these locations as the only individual reef cells that can maintain
connectivity between Refugia at this spatial scale (according to the analysis in this study).
Preserving healthy reefs in these stepping stones will enhance the resilience of reefs in nine
Refugia. Strengthening the resilience of Refugia reef cells increases the likelihood that these
reefs—predicted to have the greatest potential to maintain coral cover under climate change—
will persist and continue supplying coral larvae to other reefs, thereby supporting protection of a
much larger global reef area. These findings also underscore the need for further research into
larval and gamete connectivity across reef systems to better identify key connectors (across
spatial scales) and thus optimize global coral reef conservation strategies.
320

Data Availability Statement: The climate refugia scores and pressure scores for each Andrello
et al. (2022) reef cell can be found at https://github.com/WCS-Marine/local-reef-pressures. The
code used to generate the connectivity matrix used in this study can be found in
https://github.com/ArielGreiner/GlobalCoralConnectivity_SpatialRescue_Project, while the
original world-wide coral reef connectivity matrix generated for Wood et al. (2014) and the grid
cell coordinates are available at the University of Bristol data repository, data.bris, at:
https://doi.org/10.5523/bris.2s0fn0bc89omq2kj2rol7iolwt (Hendy & Wood, 2022). The scripts
used to modify these datasets and perform the analyses described above and the datasets needed
for doing so are available at: https://figshare.com/s/3d6dc00129a1da24820c.
330

References

Andrello, M., Darling, E. S., Wenger, A., Suárez☐Castro, A. F., Gelfand, S., & Ahmadia, G. N.
(2022). A global map of human pressures on tropical coral reefs. Conservation Letters, 15(1),
e12858.

Beyer, H. L., Kennedy, E. V., Beger, M., Chen, C. A., Cinner, J. E., Darling, E. S., ... &
Hoegh☐Guldberg, O. (2018). Risk☐sensitive planning for conserving coral reefs under rapid
climate change. Conservation Letters, 11(6), e12587.

Burke, L., Reytar, K., Spalding, M., & Perry, A. (2011). Reefs at risk revisited: technical notes
on modeling threats to the world's coral reefs. Washington, DC: World Resources Institute.

Convention on Biological Diversity. (2022). Decision 15/4: Kunming-Montreal Global
Biodiversity Framework. CBD/COP/DEC/15/4

Cruz-Trinidad, A., P. M. Aliño, R. C. Geronimo, and R. B. Cabral. 2014. "Linking food
security with coral reefs and fisheries in the coral triangle." Coastal Management 42:160-182.

Csárdi G., T. Nepusz, V. Traag, S. Horvát, F. Zanini, D. Noom, and K. Müller. 2023. "Igraph:
Network Analysis and Visualization in R." doi:10.5281/zenodo.7682609

D'Aloia, C. C., Naujokaitis-Lewis, I., Blackford, C., Chu, C., Curtis, J. M., Darling, E., ... &
Fortin, M. J. (2019). Coupled networks of permanent protected areas and dynamic conservation
areas for biodiversity conservation under climate change. Frontiers in Ecology and Evolution, 7,
27.

DeCarlo, T. M. (2020). Treating coral bleaching as weather: a framework to validate and
optimize prediction skill. PeerJ, 8, e9449.

Deysher L, Norton TA (1981) Dispersal and colonization in Sargassum muticum (Yendo)
Fensholt. J Exp Mar Biol Ecol 56:179–195. https://doi.org/10.1016/0022-0981(81)90188-X

FAO. 2012. "The State of World Fisheries and Aquaculture." FAO Fisheries and Aquaculture
Department, Food and Agriculture Organization of the United Nations, Rome. - SOFIA 2012

Frieler, K., Meinshausen, M., Golly, A., Mengel, M., Lebek, K., Donner, S. D., & Hoegh-
Guldberg, O. (2013). Limiting global warming to 2°C is unlikely to save most coral reefs. Nature
Climate Change, 3(2), 165–170.

Gotelli, N. J. (1991). Metapopulation models: The rescue effect, the propagule rain, and the core-
satellite hypothesis. The American Naturalist, 138(3), 768–776.

Greiner, A., Andrello, M., Darling, E., Krkošek, M., & Fortin, M. J. (2022a). Limited spatial
rescue potential for coral reefs lost to future climate warming. Global Ecology and
Biogeography, 31(11), 2245-2258.

Greiner, A., S. Darling, E., Fortin, M. J., & Krkošek, M. (2022b). The combined effects of
dispersal and herbivores on stable states in coral reefs. Theoretical Ecology, 15(4), 321-335.

Greiner, A., Andrello, M., Krkošek, M., Fortin, M. J., Nand, Y., Jupiter, S. D., ... & Darling, E.
S. (2025). Dispersal can spread management benefits: Insights from a modeled Fijian coral reef
network. Ecological Applications, 35(8), e70156.

Luo, Y., Wu, J., Wang, X., & Peng, J. (2021). Using stepping-stone theory to evaluate the
maintenance of landscape connectivity under China's ecological control line policy. Journal of
Cleaner Production, 296, 126356.

Manel, S., Loiseau, N., Andrello, M., Fietz, K., Goñi, R., Forcada, A., ... & Mouillot, D. (2019).
Long-distance benefits of marine reserves: myth or reality?. Trends in Ecology & Evolution,
34(4), 342-354.

McClanahan, T. R., & Azali, M. K. (2021). Environmental variability and threshold model's
predictions for coral reefs. Frontiers in Marine Science, 8, 778121.

McManus, L. C., Vasconcelos, V. V., Levin, S. A., Thompson, D. M., Kleypas, J. A., Castruccio,
F. S., ... & Watson, J. R. (2020). Extreme temperature events will drive coral decline in the Coral
Triangle. Global Change Biology, 26(4), 2120-2133.

Mouquet, N., & Loreau, M. (2003). Community patterns in source-sink
metacommunities. The American Naturalist, 162(5), 544–557.

Mumby PJ (2006) The impact of exploiting grazers (Scaridae) on the dynamics of Caribbean
coral reefs. Ecol Appl 16:747–769. https:// doi.org/10.1890/1051-
0761(2006)016[0747:TIOEGS]2.0.CO;2

QGIS Development Team (2021). QGIS Geographic Information System. Open Source
Geospatial Foundation Project.

R Core Team. 2019. "R: A language and environment for statistical computing." R
1145 Foundation for Statistical Computing, Vienna, Austria. https://www.R- project.org/.

Saura, S., Bodin, Ö., & Fortin, M. J. (2014). EDITOR'S CHOICE: Stepping stones are crucial for
species' long☐distance dispersal and range expansion through habitat networks. Journal of
Applied Ecology, 51(1), 171-182.
415
416 Saura, S., Bastin, L., Battistella, L., Mandrici, A., & Dubois, G. (2017). Protected areas in the
417 world's ecoregions: How well connected are they?. Ecological Indicators, 76, 144-158.
418
419 Sing Wong, A., S. Vrontos, and M. L. Taylor. 2022. "An assessment of people living by coral
420 reefs over space and time." Global Change Biology 28:7139-7153.
421
422 Smith, J. E., Brainard, R., Carter, A., Grillo, S., Edwards, C., Harris, J., ... & Sandin, S. (2016).
423 Re-evaluating the health of coral reef communities: baselines and evidence for human impacts
424 across the central Pacific. Proceedings of the Royal Society B: Biological Sciences, 283(1822),
425 20151985.
426
427 Stiger V, Payri CE (1999) Spatial and temporal patterns of settlement of the brown macroalgae
428 Turbinaria ornata and Sargassum mangarevense in a coral reef on Tahiti. Mar Ecol Prog Ser
429 191:91–100. https://doi.org/10.3354/meps191091
430
431 UNEP-WCMC and IUCN (2024), Protected Planet: The World Database on Protected Areas
432 (WDPA) and World Database on Other Effective Area-based Conservation Measures (WD-
433 OECM) [Online], April 2024, Cambridge, UK: UNEP-WCMC and IUCN. Available at:
434 www.protectedplanet.net.
435
436 UNEP-WCMC, WorldFish Centre, WRI, TNC (2010) Global distribution of coral reefs,
437 compiled from multiple sources including the Millennium Coral Reef Mapping Project. Version
438 1.3. Includes contributions from IMaRS-USF and IRD (2005), MaRSUS (2005) and Spalding et
439 al. (2001). Cambridge (UK): UNEP World Conservation Monitoring Centre. URL:
440 http://data.unep-wcmc.org/datasets/1
441
442 Van Hooidonk, R., Maynard, J., Tamelander, J., Gove, J., Ahmadia, G., Raymundo, L.,
443 Williams, G., Heron, S. F., & Planes, S. (2016). Local- scale projections of coral reef futures and
444 implications of the Paris Agreement. Scientific Reports, 6(1), 1–8.
445
446 Wenger, A. S., Williamson, D. H., da Silva, E. T., Ceccarelli, D. M., Browne, N. K., Petus, C., &
447 Devlin, M. J. (2016). Effects of reduced water quality on coral reefs in and out of no☐take
448 marine reserves. Conservation Biology, 30(1), 142-153.
449

450 Wood, S., Paris, C. B., Ridgwell, A., & Hendy, E. J. (2014). Modelling dispersal and
451 connectivity of broadcast spawning corals at the global scale. Global Ecology and Biogeography,
452 23(1), 1-11.
453
454 Xu, D., Peng, J., Jiang, H., Dong, J., Liu, M., Chen, Y., ... & Meersmans, J. (2024).
455 Incorporating barriers restoration and stepping stones establishment to enhance the connectivity
456 of watershed ecological security patterns. Applied Geography, 170, 103347.
457

466
467 **Figure 2**. *Stepping Stones* - (a) Each point represents a different stepping stone reef cell. (b) Black dots
468 represent the spatial extent and precise locations of stepping stone reef cells. The coloured borders around
469 each stepping stone distinguish them from each other. Black borders in (b) indicate the 'connector'

 stepping stones, while the remainder are the 'sink' stepping stones. Land polygons in (b) are from

 <u>naturalearthdata.com</u> (10m scale resolution, accessed August 19, 2025).

 Figure 3. *Re-Connecting 83 Refugia Networks Worldwide* - Bright coloured (i.e. non-light blue) reef cells
indicate Refugia reef cells, coloured according to which reef network they are in (same reef network,
same colour). (a) The light blue reefs indicate reef cells that are not in Refugia. (c) Black stars indicate
stepping-stone reef cells. Note that if only the 'connector' stepping stones are added back in, the 84
Refugia are in 114 networks (Supplementary Materials 3 - Fig. 3).

Figure 4. *Anthropogenic Pressure Scores of Stepping Stones* - The colour of each stepping stone indicates their (a) cumulative anthropogenic pressure score or their (b) top anthropogenic pressure as calculated by Andrello et al. (2022). To calculate these metrics, Andrello et al. (2022) calculated the percentile of each pressure (Fishing, Coastal Population, Industrial Development, Tourism, Sediments, Nitrogen) within a reef cell from the global distribution of each pressure. The top anthropogenic pressure level of each stepping stone is the pressure with the highest percentile for that stepping stone. The cumulative anthropogenic pressure level is the average of the percentiles of the six pressure layers within that reef cell (additional details in Supplementary Materials 1). Black circles around the stepping stones indicate the 'connector' stepping stones, while the remainder are 'sink' stepping stones. Land polygons are from naturalearthdata.com (10m scale resolution, accessed August 19, 2025). (a) The colour of the

stepping stone reef cells indicates their cumulative anthropogenic pressure score. Cumulative anthropogenic pressure scores were derived from Andrello et al. (2022)'s scores and range between 0 and 1. (b) The colour of each stepping stone reef cell indicates the top anthropogenic pressure according to Andrello et al. (2022) experienced by each stepping stone. Five stepping stones are most threatened by high fishing levels, two by high nitrogen levels and three by proximity to large coastal populations. The blue border indicates the stepping stone that is protected according to the World Database on Protected Areas (accessed on April 3, 2024). Note that because of imperfect overlap between the grid cells of Andrello et al. (2022) and this study (those from Wood et al. 2014), some of the stepping stones overlapped with multiple of Andrello et al. (2022)'s grid cells–though this only caused a discrepancy with one of them (blue-bordered stepping stone; 5 fishing grid cells, 1 industrial development grid cell and 1 coastal population grid cell, see Supplementary Materials 4).