Title:

Experimental evolution toward extinction in a molecular host-parasite system

Authors:

Kohtoh Yukawa[1], Tomoaki Yoshiyama[2], Ryo Mizuuchi[3], Norikazu Ichihashi[*1,4,5]

Affiliations:

[1]Department of Life Science, Graduate School of Arts and Science, The University of
Tokyo, Meguro, Tokyo 153-8902, Japan

[2]Graduate School of Information Science and Technology, Osaka University.

[3]Department of Electrical Engineering and Bioscience, Faculty of Science and Engineering, Waseda
University, Shinjuku, Tokyo 162-8480, Japan.

[4]Komaba Institute for Sciences, The University of Tokyo, Meguro, Tokyo 153-8902, Japan

[5]Universal Biology Institute, The University of Tokyo, Meguro, Tokyo 153-8902, Japan

Correspondence:

Norikazu Ichihashi

Department of Life Science, Graduate School of Arts and Science, The University of Tokyo, 3-8-1
Komaba, Meguro-ku, Tokyo 153-8902, Japan

Tel: 81-3-5465-7307; Fax: 81-3-5465-7307; Email: ichihashi@g.ecc.u-tokyo.ac.jp

Keywords:

Self-replicator, parasite, oscillation, coevolution, diversity, extinction

Abstract

Theoretical studies have proposed that coevolution with parasitic replicators plays a critical role in the evolution of primitive life; however, experimental verification of the potential outcomes of such coevolutionary dynamics remains limited. We previously conducted a co-evolutionary experiment using an RNA-protein replication system that resulted in the spontaneous diversification of host and parasitic RNA into five distinct lineages with robust co-replication. Here, we report contrasting evolutionary outcomes from a second long-term co-evolutionary experiment. Using a flow reactor system with increased dilution frequency over 5000 h (15,000 generations), we observed reduced diversity and frequent extinctions in later experimental stages. Co-replication assays of RNA clones revealed that the primary cause of this diversity loss was shortened reaction time due to frequent dilution. Further analysis of RNA clones that emerged during evolution suggested that the frequent extinctions resulted from the dominance of host species that had either lost parasite resistance or exhibited reduced replication ability. These findings demonstrate that co-evolution between host and parasitic replicators can result in diversity loss and frequent extinctions depending on dilution conditions, highlighting the critical role of environmental parameters, such as dilution frequency, in enabling primitive replicators to evolve sustainably toward the emergence of life.

Main

Introduction

Many scenarios for the origins of life propose that self-replicating molecules, such as self-replicating RNA, acquired the ability of Darwinian evolution in the ancient Earth and subsequently evolved through increasing complexity and diversity to approach extant life[1–9]. However, whether and how self-replicating molecules can spontaneously develop diversity and complexity through evolutionary processes remains largely unknown.

One promising approach involves experimental evolution using replicating molecules capable of evolution. The first experimental molecular evolution was reported in 1967 by Spiegelman's group, in which phage genomic RNA was replicated by purified phage RNA replication enzyme (replicase) across multiple generations[10]. Subsequently, other RNA/DNA replication systems were developed and employed for experimental evolution[11,12]. Although these systems demonstrated biologically relevant

phenomena, including niche differentiation[13] and adaptation to inhibitory chemicals[14,15], the typical

evolutionary outcome was dominance by the fastest replicators under given conditions, with no

observed diversification.

Contrasting results were obtained when a translation system was integrated with Spiegelman's RNA

replication system (designated the translation-coupled RNA replication system) in our previous

study[16]. In this system, RNA encoding an RNA replicase is replicated by the RNA replicase translated

from itself in a reconstituted E. coli translation system (Fig. 1a). A notable feature of this system is the

emergence of parasitic RNAs that have lost the replicase-encoding sequence while retaining the

replicase recognition sequence through RNA recombination. These parasitic RNAs replicate by

exploiting "host" RNAs that retain the replicase gene but replicate significantly faster than host RNAs

due to their reduced size. To prevent excessive amplification of parasitic RNAs, the system was

compartmentalized within water-in-oil droplets. Through serial dilution cycles, the RNA sustained

replication and continued evolving. In previous long-term evolutionary experiments more than 500

generations, we observed co-evolution of host and parasitic RNAs through an evolutionary arms race[17],

wherein the ancestral host RNA diversified into at least five distinct lineages—three host lineages and

two parasite lineages—forming an interdependent replication network[18].

One factor contributing to the distinct evolutionary outcomes in the translation-coupled RNA

replication system may be the emergence of parasitic entities. Parasitic molecules have been

considered a critical obstacle for primitive self-replicators to overcome prior to the emergence of life,

as parasites inevitably arise in self-replicating systems[19] and disrupt interdependent or cooperative

replication networks[20,21]. Several solutions to this problem have been proposed, including

compartmentalization, whether permanent or transient[22–28], limited dispersion[29–35], molecular kin

selection[36], and physical linkage evolution[37]. Even when the problem of parasite amplification is

resolved through these mechanisms, how hosts and parasites co-evolve remains an open question.

Theoretical studies have predicted diverse coevolutionary outcomes (reviewed in [38]) and some are

related to primitive replicators: increased host diversity and complexity[39–42], enhanced host variation

with stabilized oscillation dynamics[43], or extinctions of a host species due to reduced population sizes

during oscillations[44]. While many coevolutionary experiments using extant organisms have been

conducted (reviewed in [45]), these studies employed relatively short time scales compared to the

theoretical frameworks for primitive replicators mentioned above, and theoretical predictions

regarding diversification and extinction dynamics remain insufficiently verified. In our

aforementioned long-term coevolutionary experiment using the translation-coupled RNA replication

system, we observed stable coexistence and divergence of host and parasitic RNAs[18,46]. However,

whether this evolutionary outcome consistently occurs in this system or whether alternative outcomes

are possible under different conditions remains unknown.

In this study, we conducted another long-term evolutionary experiment up to 15,000 generations under

increased serial dilution frequency using a flow reactor system and observed a contrasting

coevolutionary outcome: evolution toward reduced diversity and periodic extinction. During the early

phase of evolution, host and parasitic RNA continued replicating with oscillatory dynamics similar to

our previous experiment; however, in later phases, host RNA concentrations frequently declined

substantially and became undetectable (i.e., underwent extinction). We also observed that the

diversified host and parasite lineages present in the initial population were eliminated shortly after

starting this evolutionary experiment, with only one host and one parasite lineage persisting in later

stages. Short-term replication assays using cloned host and parasitic RNAs indicated that diversity

loss resulted from shortened reaction times caused by frequent dilution, while periodic extinctions

were attributed to reduced parasite resistance or reduced host replication ability.

Result

Evolutionary experiment with a flow-reactor system

For another evolutionary experiment utilizing an automated format, we employed a previously

developed droplet flow reactor system[47]. In this system, droplets containing the translation system

were supplied to the reactor at a constant rate. Within the reactor, droplets, which contains host and

parasitic RNAs, were stirred at a specified frequency to induce fusion and division events, thereby

mixing the contents and enabling RNA to continue replication. Details of the flow reactor system are

presented in Fig. S1. The difference between the new flow reactor (FR) condition and the previous

manual serial dilution (SD) condition is illustrated in Fig. 1b. Under SD condition, each serial dilution

cycle consisted of 5-h incubation at 37°C, followed by 5-fold dilution with fresh oil and the translation

system. The diluted droplets were vigorously mixed using a homogenizer to achieve nearly complete

mixing of the contents of droplets, then incubated for the subsequent replication cycle. Under FR

condition, the reaction mixture was continuously diluted to exchange 20% of the tank volume per hour,

while a magnetic stirrer periodically induced droplet mixing at approximately two fusion-division

events per hour.

Using this FR system, we conducted a long-term evolutionary experiment employing the RNA

population from round 155 of our previous evolutionary experiment under SD condition, in which

RNA had already diversified into three host and one parasite lineage. Host RNA concentrations were

quantified by reverse transcription followed by quantitative PCR (RT-qPCR). Figure 1c presents host

RNA concentrations for both the previous evolutionary experiment under SD condition (light green

lines) and the current experiment under FR condition (dark green lines). Under both conditions, host

RNA concentrations exhibited oscillatory dynamics; however, the average RNA concentration under

FR condition was lower than under SD condition. Host RNA continued to be detected for

approximately 2000 h under FR condition, but undetected (i.e., extinct) after 2264 h (top panel in Fig.

1c). We restarted the experiment using preserved droplets from 1920 h (indicated by the red dotted

line). Then, host RNA replication persisted slightly longer, suggesting that extinction was stochastic,

but again became undetectable after 2600 h (second panel from top). In this way, we continued

evolutionary experiment until 5024 h and experienced extinctions eight times in total. Extinctions

occurred frequently in later experimental phases, particularly around 2000, 3000, and 4500 h,

suggesting that RNA populations evolved during these periods may possess characteristics that induce

them to extinction.

The complete trajectory of RNA concentration for the experiment with the longest duration up to 5000

h (corresponding to 15,000 generations, see Method for calculation) is presented in the bottom panel

of Fig. 1c. For this experiment, we also quantified parasitic RNA concentrations (purple marks) based

on band intensity following polyacrylamide-gel electrophoresis (Fig. S2). The detection limit for the

band intensity method is substantially higher than that of RT-qPCR employed for host RNA

quantification; consequently, parasitic RNA was detectable only at elevated concentrations. Parasites

were detected intermittently around host RNA concentration peaks, indicating that parasite RNA

concentrations also oscillated in accordance with host oscillations. In subsequent analyses, we

examined RNA populations from this longest-duration evolutionary experiment.

Sequence analysis

To elucidate the evolutionary dynamics of hosts and parasites during the FR evolutionary experiment,

we analyzed host and parasite RNA sequences at nine time points (red stars in Fig. 1c) using long-read

deep sequencing (PacBio Sequel) after reverse-transcription. To eliminate the effect of sequence error

and quasispecies, we identified dominant mutations occurring at frequencies of 10% or higher at least

one analyzed time point and classified sequences into genotypes based on the presence or absence of

these dominant mutations (Figs. S3 and S4). We constructed a phylogenetic tree incorporating the

three most abundant genotypes at each analyzed time point (solid lines in Fig. 2a), along with the data

before round 155 of the previous SD evolutionary experiment (dotted lines). We also presented the

frequency of each genotype as a heatmap aligned with the phylogenetic tree.

At the starting point (0 h) of the FR experiment (i.e., round 155 of the SD experiment), three host

lineages (HL1, HL2, and HL3) and one parasite lineage (PL2) were present (Fig. 2). However, after

480 h, HL2 and HL3 disappeared, with only one host (HL1) and one parasite (PL2) remaining

detectable, indicating that diversified lineages were not sustained under FR condition. The existence

of each lineage is schematically shown at the top of the heatmap. The heatmap additionally reveals a

relatively homogeneous population consisting of closely related branches until 5000 h, differing by

only 2-3 mutations in the host population shown in Fig. 2. This result indicated that new distinct

lineages did not emerge under FR condition. These results under FR condition represent a stark

contrast to the previous SD experiment (Fig. S5), in which five distinct lineages emerged.

162   Conditions for maintaining diversity

We investigated the differences between FR and SD conditions that influence diversity maintenance

(i.e., the number of coexisting RNA species). The two primary differences between FR and SD

conditions were 1) the continuous reaction time that RNA experienced prior to dilution and 2) the

dilution rate following the reaction. We hypothesized that these differences affect the number of

coexisting RNA species. To test this hypothesis, we conducted short-term replication experiments

employing serial dilution method using multiple RNA species at varying reaction times and dilution

rates, including those corresponding to FR and SD conditions, then evaluating the number of RNA

species present in the final population.

The reaction time and dilution rate under SD condition are corresponding to 5 h and 0.8/round,

respectively because RNA experienced continuous incubation for 5 h before dilution and then diluted

5-fold (Fig. 1b, left). Under FR condition, RNA in the droplets should have experienced continuous

incubation for 0.5 h before dilution on average because droplets underwent two fusion-division events

per hour (Fig. 1b, right). Assuming same sizes for all droplets, the dilution rate at the fusion-division

event should be 0.5. Therefore, we decided the reaction time and dilution rate under FR condition are

close to 0.5 h and 0.5/round, respectively (more detailed rationale for this calculation is described in

Method section).

As RNA species for this experiment, we initially selected four RNA species from round 155 (HL1, 2,

3, and PL2); however, these did not exhibit stable co-replication (Fig. S6), likely because these RNAs

had not yet established stable interdependencies. We therefore employed five RNA clones from a later

round—three host RNAs (HL1, HL2, and HL3) and two parasitic RNAs (PL2 and PL3)—identified

at round 228 in the previous study[18], in which four of these five RNA species (excluding HL3)

demonstrated co-replication under SD condition (5 h reaction time and 0.8 /round dilution rate).

Prior to conducting the experiments, we investigated the effect of reaction time and dilution rate on

the number of co-replicable RNA species using mathematical modeling and simulation. This

mathematical model was based on a previous study[46]. The model comprised three sequential steps: 1)

RNA replication reactions dependent on the number of RNA molecules within each droplet, 2) random

removal and supply of droplets (dilution), and 3) random fusion-division of droplets (see Methods for

details). In these simulations, we varied the reaction time for RNA replication (0.5, 1, and 5 h) and the

dilution rate (0.5, 0.8, 0.9, and 0.99 per round). Other parameters, including the replication rate of each

RNA species, were derived from experimentally determined values[18]. We executed 50 rounds of the

replication-dilution-fusion-division cycle 20 times. Representative RNA concentration dynamics are

presented in Fig. 3a. The average number of remaining RNA species in final populations is displayed

in the upper left triangle of Fig. 3c.

To validate these results, we conducted nine short-term serial dilution experiments at varying reaction

times and dilution rates and quantified the concentration of each RNA species using specific RT-qPCR

(Fig. 3b). The number of maintained RNA species in the final 40-50 rounds is presented in the bottom

right panel of Fig. 3c. The number of remaining RNA species was significantly dependent on both

reaction time and dilution rate, with simulation and experimental results exhibiting similar trends. At

shorter reaction times and higher dilution rates (upper-left region of Fig. 3c), RNA species rarely

persisted, whereas at longer reaction times and lower dilution rates and (lower-right region), greater

numbers of RNA species were maintained. Specifically, under the previous SD evolutionary condition

(5 h reaction and 0.8/round dilution, green dotted square), 3.6 - 4 RNA types were sustained. Under

the condition corresponding to the FR evolutionary experiment (5 h reaction and 0.5/round dilution,

blue dotted square), only 1.6-3 RNA species persisted, consistent with the loss of pre-existing RNA

diversity during the FR evolutionary experiment shown in Fig. 2, where two RNA lineages remained.

These results suggest that reduced diversity under FR condition can be primarily attributed to shorter

reaction times because another difference under FR condition, reduced dilution rate, should increase

the diversity at the reaction time of 0.5 h based on the data of Fig. 3c. Additional discussion is provided

in Supplemental Discussion.

Cause of extinction

We investigated the underlying cause of host RNA disappearance during later stages of the

evolutionary experiment. Such disappearance may be directly attributed to reduced host RNA

concentrations at the bottom of the oscillation dynamics, which likely results in stochastic removal of

all host RNA molecules from the reactor. If this hypothesis is correct, host RNA concentrations at the

bottom should decrease in the later stage of evolution. To test this, we plotted RNA concentrations at

each bottom of the oscillation in Fig. 4a (blue points), which were approximately 10^{-3} nM during 0-

1000 h and decreased to 10^{-4} nM during 1000-2000 h and 3000-4600 h. Beyond 4600 h, the bottom

concentrations decreased further.

Next, we investigated whether evolved host and parasite RNAs in later stages exhibited characteristics

that reduced host concentrations at the bottom of oscillation. We selected the most frequent host

sequences at nine time points (0, 480, 1152, 1520, 2064, 2848, 3376, 3984, and 4616 h) and prepared

individual RNA clones. For parasite, we selected five major sequences that were identical during 1152-

2064 h and 2848-3376 h because similar sequences were sustained for relatively long periods. At 4616

h, parasitic RNA was undetectable; therefore, we employed the sequence from 3984 h. Using these

host-parasite pairs from each time point (1 nM each), we conducted single-round competition

experiments and quantified RNA replication (Fig. 4b).

For the initial host-parasite pair prior to the FR evolutionary experiment, the host replicated selectively without parasite amplification (Fig. 4b, 0 h). At 480-1520 h following initiation of the FR evolutionary experiment, parasites were amplified to levels equivalent to host replication. During 1520-2064 h, replication of both host and parasite declined. At 2848 and 3376 h, host replication increased modestly while parasite replication increased substantially. Parasite replication subsequently decreased at 3984 h, with both host self-replication activity and parasite replication declining at 4616 h.

We identified two notable patterns in host-parasite replication dynamics during the FR evolutionary experiment: 1) increased parasite replication (indicating loss of parasite resistance, e.g., at 480 and 2848 h) and 2) reduced replication for both host and parasite (suggesting parasite resistance coupled with diminished host replication ability, e.g., at 2064 and 4616 h).

The subsequent question was how these characteristics of evolved hosts and parasites resulted in reduced host RNA concentrations at the bottom of oscillation. In the evolutionary experiments, extinctions were frequently observed around 2000, 3000, and 4600 h. Host and parasite RNAs during these periods (2064, 2848, and 4616 h in Fig. 4b) exhibited two distinct characteristics: reduced host replication with parasite resistance (2064 and 4616 h), and elevated host replication with parasite susceptibility (2848 h). To test whether these two contrasting characteristics could result in lower host RNA concentrations at the bottom of oscillation, we conducted short-term replication experiments using host-parasite pairs from each time point (0, 2064, 2848, 3984 h) under the condition approximating FR condition (0.5 h reaction time and 0.5/round dilution rate), and quantified each host and parasite concentrations. These experiments were performed independently in triplicate (Fig. 4c). The average host concentration at the bottom are presented in Fig. 4d. The bottom concentration at 0 h exceeded 10^{-2} nM but decreased to below 10^{-2} nM for 2064, 2848, and 3984 h (Fig. 4d), indicating that all three evolved RNA pairs reduced the host concentrations at the bottom, consistent with frequent extinctions observed around these time periods.

Discussion

Coevolution between hosts and parasites represents an important evolutionary phenomenon relevant to the origin of life and the modern biology; however, experimental evidence regarding potential coevolutionary outcomes remains limited. Here, we conducted and analyzed a long-term evolutionary experiment under altered dilution frequency and demonstrated that the evolutionary outcome differed significantly from our previous evolutionary experiment. Host RNA exhibited a tendency toward extinction in later evolutionary stages (Fig. 1), and lost their lineage diversity (Fig. 2). Assays using RNA clones revealed that extended reaction times and moderate dilution rates were necessary for maintaining diversity (Fig. 3). Competition assay using isolated RNA clones demonstrated that RNAs selected during later stages exhibited two contrasting properties: high replication ability with low parasite resistance and low replication ability with high parasite resistance (Fig. 4b). Both properties were found to decrease the bottom host concentrations during oscillations (Figs. 4c and 4d), thereby increasing extinction probability. These results indicate that host-parasite coevolution does not invariably result in robust co-replication and spontaneous diversification, as observed in our previous evolutionary experiment, but can lead to diversity loss and increased extinction susceptibility depending on dilution conditions. These findings underscore the critical importance of environmental parameters that influence dilution dynamics for primitive replicators to evolve sustainably toward the emergence of life.

Although evolution is generally regarded as a process that either increases or maintains the fitness of populations of organisms or molecules, it can drive population extinction in theory, a phenomenon known as evolutionary suicide, runaway evolution, or Darwinian extinction[48,49]. Such evolution toward extinction has been investigated in several theoretical models[50–54], including predator-prey systems[55–57], bacterial populations[58], and epidemiological models[59,60]. However, experimental demonstration of evolution toward extinction remains limiting[61]. In this study, we provide an example of such evolution toward extinction using a molecular replication system. During the long-term evolutionary experiment under FR condition, we observed frequent extinctions only after 2000 h (Fig. 1c), suggesting that the host RNA evolved to acquire the characteristics that predispose it to extinction.

Our explanation for this counterintuitive evolutionary trajectory is as follows. First, the relatively short

reaction time under FR condition reduced diversity of RNA species, resulting in co-evolution of only

one host and one parasite species (Fig. 3). In ecological studies, reduced diversity is known to enhance

population fluctuations due to the absence of buffering effects from other species[43]. Additionally, host

RNA evolved either to lose parasite resistance (2848 h and 3376 h in Fig. 4b) or to exhibit decreased

self- replication ability (2064 h and 4616 h in Fig. 4b), both of which reduced host RNA concentrations

at the bottom of oscillation (Fig. 4c). This population reduction could cause stochastic elimination of

all RNAs, resulting in extinction. Similar extinction due to reduced population size during oscillations

was observed in a previous theoretical study[44].

A remaining question is why host RNA evolved to lose parasite resistance or decrease its self-

replication ability. Our proposed explanation is as follows. Under FR condition, dilution occurred

more frequently than under SD condition. This frequent dilution is unfavorable for parasite because

parasite replication requires coexistence with host RNA, potentially reducing the selective pressure

for parasite resistance in host RNA. Consequently, parasite-susceptible hosts would dominate, as

observed at 2848 h and 3376 h. However, when parasite-susceptible hosts dominated, parasitic RNA

would be re-amplified despite unfavorable dilution conditions, placing the host under renewed

selection pressure to increase parasite resistance. If host diversity were maintained and parasite-

resistant hosts were present in the population, such parasite-resistant hosts would increase in

frequency; however, in the FR experiment, host diversity was lost, requiring the remaining parasite-

susceptible host to newly acquire parasite resistance. To acquire parasite resistance, host RNA must

encode a replicase with specificity for the host sequence. Generally, there exists a trade-off between

specificity and enzymatic activity in many proteins[62,63]. This trade-off may decrease self-replication

ability together with parasite resistance, as observed at 2064 h and 4616 h. In summary, we propose

that host-parasite coevolution under frequent dilution results in loss of host diversity, thereby selecting

for RNA species that increased susceptibility to extinction.

Method

319 **The translation-coupled RNA replication system**

320 The translation-coupled RNA replication system used in this study originally consist of two

321 components: a host RNA that encodes a core subunit of the Qβ RNA replicase gene with the

322 recognition sequence, and the reconstituted translation system of *E. coli*, which contains all factors

required for translation, such as ribosome, translation proteins, tRNA, NTPs, and amino acids. The components were prepared as previously described[64]. Parasitic RNAs appeared during the replication from the host RNA.

Evolutionary experiment using a flow reactor

The droplet flow reactor system (Fig. S1) was constructed as described in a previous study[47]. This system consisted of two 505 µL tanks set on magnetic stirrers. Oil phases and aqueous phases (the reconstituted translation system) were supplied by syringe pumps into the 1st tank, where water-in-oil droplets were prepared by stirring with a magnetic stirrer at 2 °C. The oil phase consisted of mineral oil (Sigma) and surfactant, 2% Span 80 (Wako), and 3% Tween 80 (Wako), saturated with non-protein components of the translation system, as described previously[18]. The prepared droplets were transferred to the 2nd tank, in which the droplets contained host and parasitic RNAs, and RNA replication occurred in the droplets at 37 °C. The droplets were mixed for 3 s every 60 s by stirring at 4800 rpm to induce fusion and division between the droplets. This corresponds to the low mixing condition in a previous study[47]. The flow rate was 0.2/h (oil phase, 100 µL/h; aqueous phase, 1 µL/h).

At the start of the experiment, 500 µL of the oil phase and 5 µL of the translation system were added to the 1st tank, and the emulsion was prepared by stirring at 2 °C and 7200 rpm for 10 min. The second tank was filled with 505 µL of water-in-oil droplet, which was prepared by mixing droplets from round 155 of the previous SD evolutionary experiment with 4 µL of translation system and 400 µL of oil phase at 7200 rpm for 3 min using a magnetic stirrer at 2 °C. When restarting the reaction in the middle of this experiment, the water-in-oil droplets (505 µL) were added to the second tank at the restarting time point. The translation system and oil phase were replenished every 2–3 days.

RNA measurement

To measure the host RNA concentration, the water-in-oil droplets were diluted 100-fold with 1 mM EDTA (pH8.0) and heated at 95 °C for 5 min, followed by rapid cooling on ice. The heated sample was then used for quantitative PCR after reverse transcription (RT-qPCR) using primers 1-4 and the One-Step SYBR Prime Script PLUS RT-PCR Kit (Takara). To measure the parasitic RNA concentration, the water phase was collected from 300 µL of the water-in-oil emulsion by centrifugation at 15 krpm for 5 min. The precipitate was then washed with 120 µL of diethylether.

After centrifugation at 10 krpm for 1 min and removal of the ether phase, the remaining aqueous phase

was mixed with 50 μL of water, and RNA was purified using the RNeasy Mini Kit (QIAGEN). Purified

RNA (250 ng) was applied to an 8% polyacrylamide gel, as described previously[18]. As a size marker

for parasitic RNA, we used a short parasite (parasite α in a previous study[17] synthesized through *in*

vitro transcription. After electrophoresis at 150 V for 85 min, the gel was stained with SYBR Green II

(Takara). The intensity of each band was quantified using ImageJ software.

Sequence analysis

RNA was purified from the water-in-oil droplets as described above. Then, RNAs were reverse

transcribed with PrimeScript RTase (Takara) and primers 5 and 6, followed by RNase H (Takara)

treatment at 37 °C for 20 min. The cDNA was PCR-amplified using KOD FX (TOYOBO) and primers

5-6 and purified using the PureLink PCR Micro Kit (Life Technologies). The DNA corresponding to

the host and parasite were then separated by gel extraction using an E-gel agarose electrophoresis

system (Life Technologies), and barcode sequences were attached by PCR with KOD FX (TOYOBO)

and primers 5 and 6 attached to each barcode sequence. Then, the DNA mixture was sequenced using

PacBio Sequel. In PacBio Sequel, a single DNA molecule can be read multiple times owing to its

circular format. We used a consensus sequence of Q > 20 to reduce sequencing error. The number of

consensus sequence used was 10000 at any analyzed time point, except for parasites at 2848 h, 3376

h, and 3984 h. These three points were thought to have insufficient RNA concentrations; however,

more than 2,000 sequences were obtained. Following the method described in a previous study[18], 130

dominant mutations that existed at a frequency of 10% or higher in all the analyzed samples were

identified (Figs. S3 and S4). We classified the genotypes of each consensus sequence based only on

the dominant mutations because each consensus sequence contains many unique mutations, probably

because of the error-prone nature of RNA replication and the remaining sequencing errors. The

genotypes that were observed in more than nine reads were used for the following analysis. The

extracted genotypes and their frequencies are shown in Source Data file. Phylogenetic trees were

constructed using the top three most frequent genotypes from each analysis time point using the

neighbor-joining method in MEGA-X ver. 10.2.5[65].

Coexistence experiment in Fig. 3

The five RNA clones (three types of host RNA and two types of parasite RNA) found in the final round (round 228) of the previous evolutionary experiment (HL1-228, HL2-228, HL3-228, PL2-228, and PL3-228) were prepared by in vitro transcription as described previously[18]. The five RNA clones were mixed at 10 nM each in the translation system, and the mixture (10 µL) and the buffer-saturated oil phase (990 µL) were vigorously mixed with a homogenizer (Polytron PT-1300D, KINEMATICA) to prepare water-in-oil droplets. The droplets were incubated at 37 °C for the indicated periods and diluted with water-in-oil droplets, which were prepared by mixing the saturated oil and translation system, at the indicated dilution ratio using the same method without RNA addition. The diluted droplets were further incubated for the next round of replication, for the indicated periods. After repeating this serial dilution process, each RNA sample was measured using RT-qPCR with specific primers 7-16 and the One Step SYBR PrimeScript PLUS RT-PCR Kit (Takara). Although we used primers that selectively recognized only one of the five RNA clones, a low level of misrecognition occurred (usually 1/100 – 1/1000 of the legitimate recognition) (Fig. S7). This misrecognition effect was subtracted when calculating RNA concentration. For the round 155 RNA set, because sufficient accuracy could not be achieved with one-step RT-qPCR, only the plus strand was measured using primers 17-24. Primer sequences are shown in Source Data file.

Coexistence simulation in Fig. 3

The simulation was based on the replication-mixing-dilution model described in a previous study[18]. This model consists of three steps, 1) replication, 2) dilution, and 3) fusion-division, and 300,000 droplets (compartments). 1) In the replication step, i[th] RNA species replicate logistically in each droplet, according to the differential equations described below.

$$\frac{dRNA_i}{dt} = RNA_i \sum_{j=1}^{5} k_{ij} RNA_j \left(1 - \frac{\sum_{h=1}^{5} RNA_h}{C}\right),$$

where C is a carrying capacity in a droplet (300). The replication rate constants k_{ij}, experimentally determined previously[18], were multiplied by 0.1 in this study because the original values were too large to observe the effect of the reaction time. 2) In the next dilution step, the droplets were randomly

removed according to the dilution rate, and an equal number of empty droplets were added. 3) In the
subsequent fusion-division step, RNAs from two randomly selected droplets were combined and
redistributed into two new droplets according to a binomial distribution to mimic the fusion-division
process of droplets. This process was repeated 1.3 times the droplet number according to the previous
study. Starting from an initial state in which RNAs were distributed according to a Poisson distribution
with a mean of 10 molecules per droplet, the replication, dilution, and redistribution processes were
repeated for 50 rounds. All parameter values used in this simulation are shown in Source Data file.

Rationale of the incubation time and dilution rate under the SD and FR conditions

The condition corresponding to the previous evolutionary experiment (SD condition) was a reaction
time of 5 h and a dilution rate of 0.8 /round because one round of serial dilution cycle consisted of
incubation for 5 h, followed by 5-fold dilution.

The condition close to the new evolutionary experiment using the flow reactor (FR condition) is a
reaction time of 0.5 h and a dilution rate of 0.5 /round for the following reasons. Under FR condition,
we stirred the droplets at a 5% duty ratio, which corresponds to a diffusion coefficient of 0.35 /min^{-1},
which means that droplets experience a fusion-division event approximately once every 0.5 h
according to the data in a previous study[66] (based on Fig. 4 of the previous study). Therefore, we
adopted a continuous reaction time before dilution of 0.5 /h. In addition, RNA in a droplet should be
diluted 1/2 in a single fusion-division event, assuming that all droplets are same size and a fusion-
division event of two droplets mixes the contents completely and produces two droplets of the same
size. Therefore, we chose a reaction time of 0.5 h and a dilution rate of 0.5 /round for that close to FR
condition. It should be noted that the above calculation is valid only when the concentration of RNA-
containing droplets is low and fusion between such droplets is negligible. When the concentration of
RNA-containing droplets becomes high, the effective dilution rate approaches that of the reaction tank
itself (20% /h, Fig. 1b), corresponding to 0.1 per round at 0.5 h reaction time. Therefore, the dilution
rate of 0.5 per round used in this study may represent an overestimation.

Competition assay conducted in Fig. 4b

Representative RNAs (the most frequent genotype) at each time point were used in the co-replication
assay. The RNA clones were prepared as follows: The PCR amplified cDNA was ligated to a vector,

pUC19 or pBAD33, using In-Fusion HD Enzyme Premix (Takara), and amplified in *E. coli* DH5α. We checked the sequence using Sanger sequencing, and if there were any unintended mutations, we corrected them using whole plasmid PCR with primers containing corrected mutations, followed by ligation and transformation as described above. The plasmids containing correct RNA sequence were digested with Sma I (Takara) to determine the 3'-end of RNA, transcribed using T7 RNA polymerase (Takara), and purified using the RNeasy Mini Kit (QIAGEN). DNAs was degraded using DNase I (RNase-free) (Takara) treatment, and the RNAs were purified using the RNeasy Mini Kit (QIAGEN). Each host and parasite RNA clones (1 nM) are mixed in the reconstituted translation system and incubated at 37 °C for 1 h. The concentrations of the host and parasite RNA were measured by RT-qPCR with specific primers. Primer sequences are shown in Source Data file.

Assay of the bottom concentration conducted in Fig. 4c

Each indicated host and parasite RNA clones were mixed at 10 nM each in the translation system, and the mixture (10 μL) was homogenized in the buffer-saturated oil phase to prepare water-in-oil droplets as described above in the coexistence experiment section. The serial dilution cycle was repeated 38–40 times under the condition close to FR condition (0.5 h reaction time and 0.5 /round dilution rate) as described in the same section.

Calculation of generations

In the FR evolutionary experiment, the host RNA concentration was maintained for 5000 h while exchanging 20% of the reaction mixture per hour. Consequently, host RNAs underwent 1.25-fold replication per hour, resulting in 1.25^{5000}-fold amplification at 5000 h. We defined one generation for an RNA replicator as the time required to double. Therefore, the total number of generations was calculated as $\log_2(1.25^{5000}) \sim 15{,}000$.

Supplementary Materials

Extended Data Figures, Supplementary Discussion, and Source Data for this article is available.

Acknowledgements:

We thank Mrs. Kayo Aoyama and Ayu Saito for technical support.

Funding:

This work was supported by JST, CREST Grant Number JPMJCR20S1, Japan, and Kakenhi Grant

Numbers 24H00573 and 24H01111.

Author contributions:

TY performed the evolutionary experiments and KY design and performed the other experiments. RM

and NI design experiments. NI and KY wrote the manuscript.

Competing interests:

The authors declare that they have no conflicts of interest.

Data and materials availability:

The data are available upon request from the authors.

Reference

1.   Szathmáry, E. & John Maynard, S. The major evolutionary transitions. *Nature* **374**, 227–
232 (1995).

2.   Joyce, G. F. The antiquity of RNA-based evolution. *Nature* **418**, 214–221 (2002).

3.   Koonin, E. V. & Martin, W. On the origin of genomes and cells within inorganic
compartments. *Trends in Genetics* **21**, 647–654 (2005).

4.   Yarus, M. Getting past the RNA world: The initial Darwinian ancestor. *Cold Spring*

Harb Perspect Biol **3**, 1–8 (2011).

5.   Pross, A. & Pascal, R. The origin of life: What we know, what we can know and what we

will never know. *Open Biol* **3**, 1–5 (2013).

6.   Higgs, P. G. & Lehman, N. The RNA World: molecular cooperation at the origins of life.

Nat Rev Genet **16**, 7–17 (2015).

7.   Goldenfeld, N., Biancalani, T. & Jafarpour, F. Universal biology and the statistical

mechanics of early life. *Philosophical Transactions of the Royal Society A:*

Mathematical, Physical and Engineering Sciences **375**, 1–14 (2017).

8.   Adamski, P. *et al.* From self-replication to replicator systems en route to de novo life.

Nat Rev Chem **4**, 386–403 (2020).

9.   Brunk, C. F. & Marshall, C. R. 'Whole Organism', Systems Biology, and Top-Down

Criteria for Evaluating Scenarios for the Origin of Life. *Life* **11**, 690 (2021).

10.   Mills, D. R., Peterson, R. L. & Spiegelman, S. An extracellular Darwinian experiment

with a self-duplicating nucleic acid molecule. *Proc Natl Acad Sci U S A* **58**, 217–224

(1967).

11.   Breaker, R. R. & Joyce, G. F. Emergence of a replicating species from an in vitro RNA

evolution reaction. *Proceedings of the National Academy of Sciences* **91**, 6093–6097

(1994).

12.   Wright, M. C. & Joyce, G. F. Continuous in vitro evolution of catalytic function. *Science*

(1979) **276**, 614–617 (1997).

13.   Voytek, S. B. & Joyce, G. F. Niche partitioning in the coevolution of 2 distinct RNA

enzymes. *Proc Natl Acad Sci U S A* **106**, 7780–7785 (2009).

14.   Kramer, F. R., Mills, D. R., Cole, P. E., Nishihara, T. & Spiegelman, S. Evolution in

vitro: Sequence and phenotype of a mutant RNA resistant to ethidium bromide. *J Mol*

Biol **89**, 719–736 (1974).

15.   Paegel, B. M. & Joyce, G. F. Microfluidic compartmentalized directed evolution. *Chem*

Biol **17**, 717–724 (2010).

16.   Ichihashi, N. *et al.* Darwinian evolution in a translation-coupled RNA replication system

521   within a cell-like compartment. *Nat Commun* **4**, 2494 (2013).

522   17.   Furubayashi, T. *et al.* Emergence and diversification of a host-parasite RNA ecosystem
523   through Darwinian evolution. *Elife* **9**, (2020).

524   18.   Mizuuchi, Ryo., Furubayashi, Taro. & Ichihashi, Norikazu. Evolutionary transition from
525   a single RNA replicator to a multiple replicator network. *Nat Commun* **13**, 1460 (2022).

526   19.   Iranzo, J., Puigbo, P., Lobkovsky, A. E., Wolf, Y. I. & Koonin, E. V. Inevitability of
527   genetic parasites. *Genome Biol Evol* **8**, 2856–2869 (2016).

528   20.   Eigen, M. & Schuster, P. The Hypercycle: Part C. *Naturwissenschaften* **65**, 341–369
529   (1978).

530   21.   Maynard Smith, J. Hypercycles and the origin of life. *Nature* **280**, 445–446 (1979).

531   22.   Bresch, C., Niesert, U. & Harnasch, D. Hypercycles, parasites and packages. *J Theor*
532   *Biol* **85**, 399–405 (1980).

533   23.   Szathmáry, E. & Demeter, L. Group selection of early replicators and the origin of life. *J*
534   *Theor Biol* **128**, 463–86 (1987).

535   24.   Furubayashi, T. & Ichihashi, N. Sustainability of a compartmentalized host-parasite
536   replicator system under periodic washout-mixing cycles. *Life* **8**, 3 (2018).

537   25.   Laurent, G., Peliti, L. & Lacoste, D. Survival of self-replicatingmolecules under transient
538   compartmentalization with natural selection. *Life* **9**, 78 (2019).

539   26.   Blokhuis, A., Nghe, P., Peliti, L. & Lacoste, D. The generality of transient
540   compartmentalization and its associated error thresholds. *J Theor Biol* **487**, 110110
541   (2020).

542   27.   Matsumura, S. *et al.* Transient compartmentalization of RNA replicators prevents
543   extinction due to parasites. *Science (1979)* **354**, 1293–1296 (2016).

544   28.   Bianconi, G., Zhao, K., Chen, I. A. & Nowak, M. A. Selection for Replicases in
545   Protocells. *PLoS Comput Biol* **9**, e1003051 (2013).

546   29.   Boerlijst, M. C. & Hogeweg, P. Spiral wave structure in pre-biotic evolution:
547   hypercycles stable against   parasites. *Physica D* **48**, 17–28 (1991).

548   30.   Boerlijst, M. C. & Hogeweg, P. Spatial gradients enhance persistence of hypercycles.
549   *Physica D* **88**, 29–39 (1995).

550   31.   Cronhjort, M. B. & Blomberg, C. Cluster compartmentalization may provide resistance
551   to parasites for catalytic networks. *Physica D* **101**, 289–298 (1997).

552   32.   McCaskill, J. S., Füchslin, R. M. & Altmeyer, S. The stochastic evolution of catalysts in
553   spatially resolved molecular systems. *Biol Chem* **382**, 1343–1363 (2001).

554   33.   Szabó, P., Scheuring, I., Czárán, T. & Szathmáry, E. In silico simulations reveal that
555   replicators with limited dispersal evolve towards higher efficiency and fidelity. *Nature*
556   **420**, 340–343 (2002).

557   34.   Sardanyés, J. & Solé, R. V. Spatio-temporal dynamics in simple asymmetric hypercycles
558   under weak parasitic coupling. *Physica D* **231**, 116–129 (2007).

559   35.   Shay, J. A., Huynh, C. & Higgs, P. G. The origin and spread of a cooperative replicase in
560   a prebiotic chemical system. *J Theor Biol* **364**, 249–259 (2015).

561   36.   Levin, S. R. & West, S. A. The evolution of cooperation in simple molecular replicators.
562   *Proceedings of the Royal Society B: Biological Sciences* **284**, 1–8 (2017).

563   37.   Levin, S. R., Gandon, S. & West, S. A. The social coevolution hypothesis for the origin
564   of enzymatic cooperation. *Nat Ecol Evol* **4**, 132–137 (2020).

565   38.   Buckingham, L. J. & Ashby, B. Coevolutionary theory of hosts and parasites. *J Evol Biol*
566   **35**, 205–224 (2022).

567   39.   Takeuchi, N. & Hogeweg, P. Evolution of complexity in RNA-like replicator systems.
568   *Biol Direct* **3**, 1–20 (2008).

569   40.   Seoane, L. F. & Solé, R. How Turing parasites expand the computational landscape of
570   digital life. *Phys Rev E* **108**, 044407 (2023).

571   41.   Zaman, L. *et al.* Coevolution Drives the Emergence of Complex Traits and Promotes
572   Evolvability. *PLoS Biol* **12**, e1002023 (2014).

573   42.   Hickinbotham, S. J., Stepney, S. & Hogeweg, P. Nothing in evolution makes sense
574   except in the light of parasitism: evolution of complex replication strategies. *R Soc Open*
575   *Sci* **8**, 210441 (2021).

43.   Decaestecker, E., De Gersem, H., Michalakis, Y. & Raeymaekers, J. A. M. Damped

long-term host-parasite Red Queen coevolutionary dynamics: A reflection of dilution

effects? *Ecol Lett* **16**, 1455–1462 (2013).

44.   Gokhale, C. S., Papkou, A., Traulsen, A. & Schulenburg, H. Lotka-Volterra dynamics

kills the Red Queen: population size fluctuations and associated stochasticity

dramatically change host-parasite coevolution. *BMC Evol Biol* **13**, 254 (2013).

45.   Brockhurst, M. A. & Koskella, B. Experimental coevolution of species interactions.

Trends Ecol Evol **28**, 367–375 (2013).

46.   Kamiura, R., Mizuuchi, R. & Ichihashi, N. Plausible pathway for a host-parasite

molecular replication network to increase its complexity through Darwinian evolution.

PLoS Comput Biol **18**, e1010709 (2022).

47.   Yoshiyama, T., Ichii, T., Yomo, T. & Ichihashi, N. Automated in vitro evolution of a

translation-coupled RNA replication system in a droplet flow reactor. *Sci Rep* **8**, 11867

(2018).

48.   Webb, C. A Complete Classification of Darwinian Extinction in Ecological Interactions.

Am Nat **161**, 181–205 (2003).

49.   Kalle Parvinen. Evolutionary Suicide. *Acta Biotheor* **53**, 241–264 (2005).

50.   Ferrière, R. Adaptive responses to environmental threats: evolutionary suicide,

insurance, and rescue. in *Options Spring* 12–16 (IIASA, Laxenburg, Austria, 2000).

51.   Gyllenberg, M. & Parvinen, K. Necessary and sufficient conditions for evolutionary

suicide. *Bull Math Biol* **63**, 981–993 (2001).

52.   Parvinen, K. Evolutionary suicide in a discrete-time metapopulation model. *Evol Ecol*

Res **9**, 619–633 (2007).

53.   Parvinen, K. Adaptive dynamics of cooperation may prevent the coexistence of defectors

and cooperators and even cause extinction. *Proceedings of the Royal Society B:*

Biological Sciences **277**, 2493–2501 (2010).

54.   Parvinen, K. & Dieckmann, U. Self-extinction through optimizing selection. *J Theor*

Biol **333**, 1–9 (2013).

55. Matsuda, H. & Abrams, P. A. Runaway evolution to self-extinction under asymmetrical
competition. *Evolution (N Y)* **48**, 1764–1772 (1994).

56. Hiroyuki Matsuda & Peter A. Abrams. Timid Consumers: Self-Extinction Due to
Adaptive Change in Foraging and Anti-predator Effort. *Theor Popul Biol* **45**, 76–91
(1994).

57. Dieckmann, U., Marrow, P. & Law, R. Evolutionary Cycling in Predator-Prey
Interactions: Population Dynamics and the Red Queen. *Jouranal of Theoretical Biology*
176, 91–102 (1995).

58. Himeoka, Y. & Mitarai, N. Dynamics of bacterial populations under the feast-famine
cycles. *Phys Rev Res* **2**, 013372 (2020).

59. Boots, M. & Sasaki, A. Parasite evolution and extinctions. *Ecol Lett* **6**, 176–182 (2003).

60. Boldin, B. & Kisdi, É. Evolutionary suicide through a non-catastrophic bifurcation:
adaptive dynamics of pathogens with frequency-dependent transmission. *J Math Biol* **72**,
1101–1124 (2016).

61. Rankin, D. J. & López-Sepulcre, A. Can adaptation lead to extinction? *Oikos* **111**, 616–
619 (2005).

62. Tawfik, D. S. Accuracy-rate tradeoffs: How do enzymes meet demands of selectivity and
catalytic efficiency? *Curr Opin Chem Biol* **21**, 73–80 (2014).

63. Jana, T., Brodsky, S. & Barkai, N. Speed–Specificity Trade-Offs in the Transcription
Factors Search for Their Genomic Binding Sites. *Trends in Genetics* **37**, 421–432 (2021).

64. Hagino, K., Masuda, K., Shimizu, Y. & Ichihashi, N. Sustainable regeneration of 20
aminoacyl-tRNA synthetases in a reconstituted system toward self-synthesizing artificial
systems. *Sci Adv* **11**, 6269 (2025).

65. Kumar, S., Stecher, G., Li, M., Knyaz, C. & Tamura, K. MEGA X: Molecular
evolutionary genetics analysis across computing platforms. *Mol Biol Evol* **35**, 1547–
1549 (2018).

66. Yoshiyama, T., Motohashi, C., Ichihashi, N., Ichii, T. & Yomo, T. Maintenance of a
translation-coupled RNA replication in dynamic micro-sized compartments under

continuous stirring and flow. *Chemical Engineering Journal* **283**, 896–902 (2016).

Figures

Figure 1: A long-term experiment using a flow reactor.

(a) Scheme of translation-coupled RNA replication. RNA replicase is translated into a host RNA and

replicates both host and parasitic RNA. (b) Differences between previous serial dilution (SD) and new

flow reactor (FR) methods. In the SD method, water-in-oil droplets containing the host and parasite

RNA and the translation system were incubated for 5 h at 37°C for RNA replication. Then, an aliquot

of the droplets was diluted to 5-fold with new oil phage and the translation system, vigorously mixed

to prepare new droplets with completely mixed contents, and incubated for 5 h at 37°C again for the

next round of replication. In FR method, water-in-oil droplets were incubated at 37°C and stirred for

3 s every 60 s, which allows approximately 2 fusion-division events /h. New droplets containing the

translation system were supplied constantly at a rate of exchange of 20% of the total volume per hour.

(c) Trajectories of host RNA concentration during the FR evolutionary experiment until each

extinction. The host RNA concentrations were measured by RT-qPCR using host-specific but lineage-

nonspecific primers. The results of previous evolutionary experiment under SD condition from rounds

0 to 240 are shown in light green, assuming that one round was equivalent to five hours. The new

evolutionary experiment under FR condition was started from the RNA population of round 155 under

SD condition, and the results are shown in deep green. When the host dropped below the detection

limit multiple times and did not recover, the experiment was restarted from the stocked sample at an

earlier time point (indicated by the red dotted arrows). Parasitic RNA concentrations, measured by

quantification from SDS-PAGE images, are shown in the bottom panel (purple makers). The time

points used for sequence analysis are indicated by red stars.

Figure 2: Phylogenetic tree and frequencies in the population

(a) A phylogenetic tree was constructed using the three most frequent genotypes at each sequencing time point. The branches for the previous SD experiment (until round 155) and new RF experiment are shown as dashed and solid lines, respectively. Different types of RNA are depicted in different colors based on the classification in a previous study[18]. The heatmap shows the frequencies of the genotypes in each branch. (b) Schematic representation of existing RNA lineages. Uncertain regions due to the lack of sequence data were shown in dotted lines.

Figure 3: Effect of dilution rate and reaction time on coexistence of multiple host and parasitic RNA

Short-term serial dilution experiment with five types of RNA species (three hosts and two parasites) were performed for 40 – 50 rounds at different dilution rates and reaction times using both computer simulations (a) and experiments (b). (c) Summary of the number of RNA species in the final population in the simulation (top left, average of 20 runs) and the experiment (bottom right).

Figure 4: Characterization of RNA species that appeared in the FR evolutionary experiment.

(a) Trajectory of the host RNA concentrations at the bottom of oscillation. (b) Competition experiments with a host and parasitic RNA clones that dominated at the indicated time points. Each RNA pair (1 nM each) was incubated in the translation system at 37 °C for 1 h, and the host and parasite concentrations were measured using RTqPCR. (c) A short-term serial dilution experiment using the host and parasite RNA pairs (10 nM each) at 0, 2026, 2848, and 4616 h under the condition close to FR condition (0.5 h reaction time and 0.5 /round dilution rate) for 38–40 rounds. The average host RNA concentrations at the bottom are shown in (d) with standard errors (N = 3 or 4).