Lineage domains and cytoskeletal cables organize a cellular square grid in a crustacean

Beatrice L. Steinert[1,2,3], Leo Blondel[4,5], Chandrashekar Kuyyamudi[1,6], Evangelia Stamataki[7,8], Anastasios Pavlopoulos[7,8], Cassandra G. Extavour[1,4,6] *

1 Department of Organismic and Evolutionary Biology, Harvard University
2 Department of the History of Science, Harvard University
Current address: Brown University
Department of Molecular and Cellular Biology, Harvard University
Current address: Learning Planet Institute, France
Howard Hughes Medical Institute
Institute of Molecular Biology and Biotechnology, Foundation for Research and Technology-Hellas, Heraklion, Greece
Janelia Research Campus, Howard Hughes Medical Institute, Ashburn, VA, USA

* Correspondence: extavour@oeb.harvard.edu

Abstract

To build tissues, and ultimately functional bodies, cells in early embryos must arrange into specific patterns. In most animals, epithelial tissues exhibit a predominantly hexagonal space-packing geometry. However, in many species of the largest group of crustaceans, the Malacostraca, the embryonic epithelium takes on the striking form of a grid made up of predominantly square cells, sequential rows of which establish the adult segmented body plan. After square cells emerge, their organization appears to be maintained by specific cell division patterns. However, the mechanisms that initially generate square cells from hexagonal precursors are unknown. Here we address this problem by combining long-term multiview lightsheet microscopy, immunohistochemistry, laser ablation, and pharmacological perturbation. We show that in the emerging model crustacean *Parhyale hawaiensis* this highly unusual grid geometry is first initiated from two perpendicular axes that are established sequentially according to different cellular mechanisms. The first axis arises dorso-ventrally at a tensile lineage compartment boundary, while the second emerges at the anterior-posterior axis along the ventral midline through lineage-independent cell intercalation driven by tensile actomyosin cables. We show that these midline cables are necessary for organizing square-cell packing as well as for proper expression of the segmentation gene *engrailed*. Our findings show that both cell lineage-specific behaviors, as well as lineage-independent supracellular structures, are required to establish square grid epithelial organization and a segmented body plan.

Keywords: amphipod, *Parhyale hawaiensis*, ectoderm, segmentation, epithelium, morphogenesis

Introduction

Introduction

The organization of cells and tissues during development is fundamental for multicellular

organisms. During embryogenesis, cells must be shaped at both the individual and collective tissue

levels to form functional structures at every stage of ontogeny. It has been appreciated since the

late nineteenth century that mechanical forces and geometric properties of cells and tissues play a

central role in morphogenesis, the generation of form during development[1]. Despite the

predominant focus on the role of genes in tissue patterning during the last half century[2–5], recent

decades have witnessed a resurgence of interest in the mechanics of development at the cell and

tissue scales of biological organization[6–8]. Harnessing techniques for live imaging and

computational image processing and annotation, recent work has begun to couple mechanical and

genetic contributions to tissue organization during embryogenesis[9]. However, these investigations

have been largely limited to a small number of model laboratory organisms, giving us a narrow

phylogenetic sampling.

The arthropods, which include insects and crustaceans, are and always have been the most

speciose group of animals on earth[10]. All arthropods have segmented body plans that are

established early in embryonic development[11]. Segmentation is evident in embryonic ectodermal

epithelia[12], tissues in which cells are tightly adhered to one another in a monolayer sheet and apical

cell faces form a confluent network of polygons[13]. Most epithelia that have been studied to date

are those in which constituent cells exhibit predominantly hexagonal packing with three-way

vertices, which is the geometry with minimal surface energy and maximal space-filling

efficiency[14–18]. During segmentation in the fruit fly *Drosophila melanogaster*[19,20] and the

milkweed bug *Oncopeltus fasciatus*[21], a small number of cells at segment boundaries briefly stray

from this geometry to give rise to an increased number of 4-way vertices, suggesting an altered

mechanical state. However, most epithelial cells retain their hexagonal shapes throughout

segmentation and early morphogenesis[22–24]. In *D. melanogaster*, maintaining boundaries between

embryonic segments requires tensile cables of actin and non-muscle myosin (myosin 2), which are

mechanically coupled across the tissue and prevent cell mixing across the boundary[19,24].

In contrast, in the early embryonic ectodermal epithelium of malacostracan crustaceans

(including shrimp, lobsters, isopods and amphipods) cells are packed into a grid of square cells[25,26]

arranged into highly ordered rows and columns[27,28]. Expression of the segment polarity gene

engrailed is initiated precisely every four rows of the square grid[29]. This emergence and

maintenance of square cells across the tissue is particularly striking given that higher surface

energy is predicted to be required for this space-packing geometry, and that 4-way vertices, which

are essential for square-packing, are unstable in vertex models of epithelia[30]. The amphipod

crustacean *Parhyale hawaiensis* is a powerful model for deepening our understanding of how local

lineage relationships and physical parameters of individual cells can result in correct tissue

morphogenesis and body patterning[31]. In contrast with *D. melanogaster*, in which early patterning

is set in motion within a syncytial embryo[32], the early *P. hawaiensis* embryo cleaves

holoblastically in a highly stereotyped manner that defines the cellular progenitors of all three

germ layers and the germ line as individual blastomeres of the 8-cell embryo[27]. From this stage

onwards, the three blastomeres that will give rise to the ectoderm–the descendants of which make

up the square-packing epithelium–can be tracked and their clones spatially mapped[27,33].

Here we uncover the cellular mechanisms underlying morphogenesis of this highly ordered

tissue in *P. hawaiensis*. Previous work in this amphipod suggests that after square cells appear,

their packing into a grid is maintained by cell divisions that occur in bilaterial, palindromic medial-

to-lateral mitotic waves during the phase when grid rows duplicate along the anterior-posterior

axis to establish clonally related parasegments[34]. However, the mechanisms that ensure that 4-way

vertices, square cells, and the tissue-wide arrangement of cells into rows and columns emerge in

the first place, remain unknown.

To address this problem, we used long-term, multiview lightsheet and confocal

microscopy, and imaged *P. hawaiensis* embryos from the 16-cell stage through to the first

emergence of square cells and the square grid (Fig. 1a). We used semi-automated cell tracking to

trace the ectodermal lineages and map their clonal domains onto the quantitative dynamics of cell

shape within the expanding epithelium. We show that square-packing geometry emerges along

two axes, corresponding with the anterior-posterior and dorsal-ventral body axes, through both

lineage-dependent and lineage-independent morphogenetic mechanisms within the tissue. We

further show that supracellular actomyosin cables form at these axes, and that the cables are under

locally-generated tension higher than that of the surrounding tissue. Finally, we provide evidence

that these cables function as a mechanical organizer of this tissue.

Results

Ectodermal grid formation

The first seven cleavages of the early *P. hawaiensis* embryo are holoblastic, and the progenitors

of all three germ layers and the germ line are established at the 8-cell stage[27,29,33,35]. Beginning at

approximately 20 hours post fertilization ("hpf" hereafter), cells that will contribute to the embryo

migrate towards the anterior pole and coalesce into an embryonic rudiment referred to as a "germ

disc," positioned atop yolk and surrounded by extraembryonic tissue (Fig. 1d). By 38 hpf, after

gastrulation (Fig. 1c-e, blue dotted line[33,36]), the ventral surface of this disc (Fig. 1f, h) consists of

a monolayer of ectodermal cells[15]. By 68 hpf, the cells in the germ disc have begun aligning into

rows and columns and transforming their apical faces into squares (Fig. 1i, j). This square grid sets

up the logic of segmentation of the adult body plan (Fig. 1g), with the conserved segment polarity

gene *engrailed* becoming expressed along its rows (Fig. 1i-k)[29,37].

To visualize this epithelial morphogenesis and its relationship to the ectodermal lineages

established at the 16-cell stage (Fig. 1b outside of blue dotted line, Supplementary Fig. 1a), we

generated volumetric, long-term timelapse images of *P. hawaiensis* embryos injected with nuclear

marker *H2B-mCherry* and plasma membrane marker *Lyn-EGFP* mRNA reporter constructs using

multiview lightsheet microscopy (Fig. 1a, Movies 1-2). We used a combination of manual and

semi-automated methods to annotate cell lineage, cell positions and mitotic activity throughout the

entire embryo in frames captured every ten minutes over more than 60 hours of development. We

captured the period starting from the eight cell stage at 9 hpf (Stage 4 as per Browne and

colleagues[29]) through to Stage 11[29] at approximately 70-74 hpf (Fig. 1a). By the end of the capture

period, the ectodermal grid of square cells that is the subject of this study is clearly established

(Fig. 1i, i'). We performed this annotation for three embryos, all of which hatched successfully

following imaging (Supplementary Table 1).

We wished to determine whether and how cell lineage, revealed by the tracking described

above, was related to the cell shape dynamics and rearrangements that establish the square grid

(Movie 3). However, in the later time-points of these datasets, when the germ disc has expanded

over the gradual curve of the embryo's surface, maximum intensity projections of the image

volume obscure apical cell morphologies and challenge accurate quantification of cell shape

dynamics. To solve this problem, we conformally mapped the 3D surface of the embryo onto a 2D

plane using the tissue cartography tool ImSAnE[34,38] (Fig. 1c'). To visualize the grid at even later

stages when the square-cell packing geometry has extended throughout the expanding ectoderm,

we dissected fixed embryos from the underlying yolk and physically flattened the ectoderm into

the 2D plane (Fig. 1h-k). The surface curvature of the unperturbed tissue at these stages is gradual

enough that, unlike in some large vertebrate embryos[39], flattening does not require incisions and

does not deform cell shapes. Using images of these flattened epithelia, we quantified the dynamics

of cell shape change in the ectoderm, as well as the relationship of these dynamics to the lineage

relationships of the constituent ectodermal cells, from the 16-cell stage through to the

establishment of the grid of square cells that serves as the template for the segmented body plan.

Square-packing geometry initiates at a lineage compartment boundary

Previous work has suggested that an ectodermal lineage compartment boundary runs perpendicular

to the anterior-posterior axis of the *P. hawaiensis* embryo[40]. Given the early stereotypic cleavage

patterns of this embryo, we asked whether lineage could contribute to the initiation of square-grid

morphogenesis. We tracked all six of the 16-cell ectodermal lineages (Era, Ela, Erp, Elp, Epr, Epl;

Supplementary Fig. 1a; nomenclature as per Gerberding and colleagues[27]) in the lightsheet

animations, enabling us to delineate the dynamic lineage territories within the developing germ

disc. We observed that a strict boundary emerged immediately after gastrulation between the

anterior (Eva, Ela, grey in Fig. 2a') and posterior (Erp, Elp, Epr, Epl, striped in Fig. 2a') lineages,

and that clones from each domain never crossed this boundary (Supplementary Fig. 1b, c). This

boundary, which we call the initiation line (IL, arrowheads in Fig. 2a), straightened into a line

perpendicular to the anterior-posterior axis by 68 hpf (Fig. 2b). The IL was made up of over 30%

4-way vertices from 38-56±2 hpf, and of over 40% 4-way vertices by 62±2 hpf (Fig. 2c). This

shows that the first emergence of large numbers of 4-way vertices, required for square cell

formation, is coincident with the formation of the IL, which is an ectodermal compartment

boundary.

4-way cellular vertices first emerge in the local vicinity of the IL

163   Given that 4-way vertices are essential for square cell-packing, and that such vertices first

appear in large numbers at the IL, we next characterized the emergence of square cells at the IL.

From 38-68 hpf, the cells near the IL exhibited extensive changes in shape and arrangement (Fig.

2a). The occurrence of cell rearrangement at the IL was evident by the dynamics of the four 16-

cell stage posterior ectodermal lineage domains (Erp, Elp, Epl, Epr; Supplemental Fig. 1a).

Whereas in the 26±2 hpf germ disc the Elp and Erp domains were on opposite sides (left and right)

of the germ disc, separated from each other primarily by descendants of the anterior Era and Ela

clones (Supplementary Fig. 1b, c, top row), they ultimately became separated by a single column

of posterior cells descended from one or both of Epl and Epr (Supplementary Fig. 1b, c, bottom

row). We also observed cell proliferation throughout the germ disc, including directly at the IL

and in the anterior and posterior compartments, throughout this time window (Supplementary Fig.

2, Supplementary Table 2).

175   To quantitatively describe the cell shape dynamics, including the transient changes in edge

numbers that result from a neighboring cell division and the instability of 4-way vertices[30], we

developed a cell shape descriptor metric, defined as the ratio of the area of a cell to the area of the

bounding circle of that cell. For this metric, values range from 0.5 to 1, where higher values are

closer to hexagons (Fig. 1l). We defined "square cells" as those with values in the range of 0.6-

0.66 (darkest color in Fig. 1l). We analyzed the changes in shape in cells along the IL (Fig. 2d, A1

and P1), and in cells one cell diameter away from the IL in both the anterior and posterior

compartments (Fig. 2d, A2 and P2). At 38 hpf, the rows immediately anterior (A1) and posterior

(P1) to the IL had more square cells (> 25%) than other measured regions (Fig. 2d). After 38 hpf,

P1 and the row one cell diameter anterior to the IL (A2) had the highest proportion of square cells

(20-25%) until 62 hpf, but by 68 hpf the proportion of square cells in A2 dropped to less than 20%

(Fig. 2d). The proportion of square cells immediately anterior to the IL (A1), and one cell diameter

away posterior to the IL (P2), was similar in both regions from 44 to 68 hpf (Fig. 2d). The

proportion of square cells in both regions increased by 68 hpf, comprising the highest proportion

of square cells of all measured regions. This gradual and local increase in square cells near the IL,

suggests that this lineage boundary initiates and maintains the transition to square-packing

geometry along the dorsal-ventral axis.

A simulation of cell sorting can explain the emergence of the IL as a straight boundary

Given that the IL is a lineage compartment boundary, we asked whether and how lineage might

contribute to the emergence of this first grid axis, which becomes increasingly straight from 26-

68 hpf (Fig. 2a, Supplementary Fig. 1b, c). To test the hypothesis that minimization of interfacial

free energy at the boundary between cells of distinct ectodermal lineages might be responsible for

boundary formation, we used CompuCell3D[41–43] to simulate cell sorting within the *P. hawaiensis*

ectoderm. To obtain biologically realistic parameters for this simulation, we first segmented

lightsheet images at 26 hpf, the first time point after the completion of gastrulation, when the

surface epithelium is made up entirely of ectoderm cells (Figs. 2a, 3d, Supplementary Fig. 1b). We

then quantified the number of cell divisions within each of the six 16-cell stage ectodermal lineages

(Fig. 3a) up to 20 hours after this time point (46 hpf). The membrane segmentation and lineage

proliferation rates, as well as increased contact energy between the anterior and posterior lineages

(Fig. 3b'), were the initial conditions for the simulation. When run, the model consistently (n = 10

runs) produced a straight anterior-posterior compartment boundary as well as a clonal domain

mapping like those observed in the actual tissue (Fig. 3e; compare with Supplementary Fig. 1b, c,

bottom row). As observed in living embryos (Supplementary Fig. 1b, c), descendants of anterior

(Era, Ela) and posterior (Erp, Elp, Epr and Epl) ectodermal founder cells in our model became

increasingly mixed with each other anterior and posterior to the IL respectively (Fig. 3c, groups I

and II). In contrast, cells on either side of the IL became increasingly segregated over the analyzed

time course (Fig. 3c, group III). The results of the model are consistent with the hypothesis that

the IL represents a stable compartment boundary arising from the principle of energy minimization

between clones of anterior and posterior cells.

A midline of square cells forms the secondary axis of the grid through intercalation

Previous clonal analyses across multiple species of malacostracan crustaceans have shown that a

morphologically distinct column of cells forms along the ventral midline of the embryo[26,27,44–48].

The cells of the midline were previously reported to be entirely comprised of Ep descendants

(Supplementary Fig. 1a) in *P. hawaiensis*[27], but are described as a polyclone descended from

multiple 8-cell stage ectodermal precursors in the related amphipods *Orchestia cavimana*[46,49] and

Gammarus pulex[45]. We asked whether the *P. hawaiensis* midline played a role in square-cell

packing along the anterior-posterior axis of the ectodermal grid, and whether its formation was

lineage-dependent. We observed that *P. hawaiensis* midline formation began at 50-60 hpf through

sequential mediolateral intercalation[50] of a founder cell population (Fig. 4a-b) in a region we call

the intercalation zone (IZ, Movie 4, Fig. 4a'-a''', Supplementary Fig. 3c, d). This region

subsequently split into anterior and posterior intercalation zones (Fig. 4a''-a'''), defined as regions

of intercalation anterior (IZA) and posterior (IZP) to the IL respectively, as cells aligned at the

midline. The IZ founder cell population was identified in the lightsheet datasets by tracking the

lineages of aligned cells at 68-70 hpf backwards to the point when only one midline cell was

aligned. For the next 10 to 12 hours, the midline elongated (Fig. 4b', c', Supplementary Fig. 3e, f)

by a combination of mitotic divisions within the founder population (Fig. 4b', 9-cell midline) and

cell intercalation. Midline intercalation started at the IL (arrowheads in Fig. 4a'-a''',

Supplementary Fig. 3c-f) and progressed both anteriorly and posteriorly, with the accumulation of

midline cells in the anterior zone lagging behind that in the posterior (Fig. 4c). Cells were added

to the midline sequentially (Fig. 4c', Movie 4) and each new cell added took its final shape as a

square only once it was entirely within the midline (Fig. 4b'). Once aligned, midline cells remained

square and were the first square cells formed along the anterior-posterior axis of the grid (Fig. 4a'-

a''', Supplementary Fig. 3e, f). To the left and right of this axis, the remaining ectodermal cells

first aligned and subsequently transitioned to square-packing in a medio-lateral wave initiated from

the midline (Fig. 4a''-a'''). After the 10-12 hour period of midline elongation by intercalation and

founder cell division, by 60-72 hpf the midline contained a mean of 8.75 cells in total (Fig. 3c).

Subsequently, divisions of parasegment precursor rows[29] in the posterior compartment introduced

another mechanism of further midline elongation (starred cells in Fig. 4a''-a''').

In contrast with the initiation line, the midline grid axis did not strictly correspond to clonal

boundaries between 16-cell ectodermal lineage domains (Supplementary Fig. 3). The emergence

of the midline, with the same cell behaviors extending it in both directions along the anterior-

posterior axis, spanned the anterior and posterior lineage domains (Fig. 4b, Supplementary Fig.

3c, d). However, we observed that the posterior midline was not exclusively made up of Ep

descendants (Supplementary Fig. 3c). This is in disagreement with the original lineage tracing

report for *P. hawaiensis*[27], but is consistent with the report that embryos lacking an Ep blastomere

still form a midline[40]. In the anterior portion of the midline, clones from both anterior ectodermal

lineages (Era and Ela) contributed to the midline founder population with no apparent relationship

between lineage domains and the anterior IZ (Supplementary Fig. 3a, b). Taken together, these

results suggest that the midline initiates the emergence of complete square cells and square-cell

packing along the anterior-posterior axis through a highly coordinated morphogenetic process

based on cell recruitment rather than inherited fates, as it is not dependent on the 16-cell stage

ectoderm lineages (Supplementary Fig. 3).

The initiation line and midline column form tensile cytoskeletal cables

Given the emergence of square-cell packing geometry first at the IL (around 30 hpf) and

subsequently at the midline (around 50 hpf), we hypothesized that these boundaries introduce

tension into the tissue to initiate the transition to square-packing. To test this hypothesis, in *P.*

hawaiensis embryos stained with phalloidin (Fig. 5a) and injected with a myosin 2 mRNA reporter

construct (based on the *P. hawaiensis* ortholog of the *D. melanogaster* gene encoding the

regulatory light chain of the non-muscle type 2 myosin, known as *spaghetti-squash* [51] (*Ph-sqh*),

Supplementary Fig. 4), we measured the relative amounts of filamentous actin (F-actin) and

myosin 2 at the IL when it appeared in earlier stages (38-56 hpf, Fig. 5a-a'), and on either side of

the midline when it formed at later stages (around 68-74 hpf, Fig. 5b, b').

In 38-56 hpf embryos, we found significantly higher levels of F-actin (Fig. 5a, a') and

myosin 2 (Supplementary Fig. 4a, a') at the IL relative to the equivalent region of interest (ROI)

one cell anterior (IL/A1) or posterior (IL/P1), compared with one to two cells posterior (P1/P2).

Similarly, we also found significantly higher levels of F-actin on the left (ML) and right (MR)

sides of the midline in embryos with 6-15 cells in the midline (68-74 hpf) (Fig. 5b, b') compared

with the equivalent ROIs one and two cells to the left (ML1, ML2) and right (MR1, MR2). In

contrast, levels of myosin 2 on either side of the midline were not significantly higher than the

ROIs to the left and right of the midline (Supplementary Fig. 4b'b"). We hypothesize that these

accumulations of cytoskeletal proteins constitute supracellular cables running along the two axes

of the ectodermal grid (Supplementary Fig. 5).

In *D. melanogaster*, actomyosin cables segment embryonic and larval tissues by

introducing local tension into the tissue[19,52,53]. We hypothesized that the supracellular cables we

detected in the *P. hawaiensis* germ band were similarly under tension, allowing them to regulate

the formation of the first square cells in the grid. To test this hypothesis, we cut the cables with a

pulsed laser tuned to 927 μm and measured subsequent tissue recoil around the incision site (Fig.

5c, dashed line and arrows). In 38-44 hpf embryos that had formed the IL, recoil speeds were

significantly (1.5 to 2.5-fold) higher at the IL compared with control cuts made four to seven cells

anterior to the IL (Fig. 4c'). Interestingly, this increase is comparable to the mean of 2.37-fold

higher tension reported at the *D. melanogaster* wing disc anterior-posterior compartment boundary

compared with that at cell edges within the anterior or posterior tissue compartments[52]. In embryos

live-imaged following the IL cut, we observed that the 4-way vertices surrounding the incision site

remained intact (black arrowheads in Fig. 5c), suggesting that maintaining a square shape at the

IL is not dependent on tension in neighboring cells.

In later stage embryos in which 3-6 cells had aligned into the midline (58-69 hpf), recoil at

midline cables in the posterior IZ was significantly (2 to 3-fold) higher than at other non-IZ regions

one to four cell diameters anterior or lateral to the midline (Fig. 4c"). In contrast, recoil speeds at

the cables in the anterior region of the midline where intercalation had finished and cells were

square in shape (P1 in Fig. 5b, c") were not significantly different than those at the lateral and

anterior control regions (Fig. 5c"). To determine whether tension was mechanically coupled across

cells contributing to the cable, we measured the recoil of a second cut in the midline after making

a first cut two to three cells away in the cable (Fig. 4c'), and found that P1 and IZ second cut

recoils were not significantly different from those at first cuts (Fig. 5c"). We note that a similar

phenomenon of local, potentially cell-autonomous contribution to tension across a supracellular

actomyosin cable spanning many cell diameters, has also been reported in the morphogenesis of

an extra-embryonic membrane in the beetle *Tribolium castaneum*[54]. Taken together, these results

suggest that there are supracellular cables under tension at the IL and at the midline, and that

tension within midline cables is heterogeneous and locally generated.

Tension in actomyosin cytoskeletal cables results from the phosphorylation of the

regulatory light chain of myosin 2, which slides the attached actin filaments past one another[55].

The results of our laser ablation experiments suggested that while both the IL and the midline

cables possessed supracellular actomyosin cables that were under tension, tension at IL was not

essential for the formation of 4-way vertices. This led us to hypothesize that abrogation of myosin

2 phosphorylation would not hinder square cell-packing geometry formation at the IL at earlier

stages, but might disrupt square cell formation at the midline at later stages.

To test this prediction, we first assessed the effect of sustained elimination of myosin 2

phosphorylation during the later stages. We focused on the period when intercalation is the only

mechanism of midline formation (50-68 hpf, Fig. 4b, b'), and chose morphologically stage-

matched embryos in which just one cell had aligned into the midline (50-56 hpf, Fig. 4b) for

incubation in 50μM of the Rho-kinase inhibitor Rockout for 12 hours (Fig. 5d, d'). This

concentration of Rockout did not impact normal cell proliferation (Supplementary Fig. 6a, b),

thereby allowing the thresholds of contractility needed for cytokinesis[19,53]. Rockout-treated

embryos at this stage had significantly fewer cells aligned into the midline than controls (mean 1.4

vs 6.4, Fig. 5f) but maintained the same total number of cells in the posterior compartment (Fig.

5f'). Notably, a relatively straight boundary at the IL was maintained in the midline Rockout-

treated embryos (Fig. 5d', arrowheads). In contrast, earlier stage embryos (38 hpf) treated for 12

hours in 50µM Rockout had, by 50 hpf, percentages of 4-way vertices that were not significantly

different from controls (Fig. 5e). Taken together, these results suggest that while 4-way vertices

first appear at the IL (Fig. 2), their maintenance is not dependent on the phosphorylation of myosin

2 (Fig. 5e). In contrast, however, subsequent establishment of the first square cells is dependent

on the contractility of myosin 2 within midline cables (Fig. 5g).

The midline cables organize square-cell packing and the emergence of a stripe of Engrailed-

positive cells

Given the presence of differential tension within the germ disc (Fig. 5c, c"), and the midline as the

initiation site of the transition to square-packing along the axis of grid elongation (Fig, 4,

Supplementary Fig. 3), we hypothesized that localized mechanical perturbation of the midline

would disrupt the organization of square-cell packing across the tissue, and that perturbing square-

cell packing would in turn have consequences for the correct establishment of segments. To test

this, we performed two classes of physical ablations, either directly at the midline (Mid), or three

cell diameters lateral to the midline (Lat) in embryos in which 3-6 midline cells had aligned (58-

69 hpf, A: pink rectangles in Fig. 6b, c). The contralateral side of the same embryo, which was not

ablated (NA: pink rectangles in Fig. 6b, c), served as an internal control. In the resulting four

experimental groups (Mid-A, Mid-NA, Lat-A, Lat-NA), we analyzed posterior compartment cell

shapes (Fig. 6d, d'), number of cells in the midline (Supplementary Fig. 7), and expression of the

segment marker Engrailed[37] (En, Fig. 6e, Supplementary Table 3) 9-12 hours after ablation. In the

100 closest cells to P1 posterior to the IL, we used Kernel density estimation[56,57] to estimate the

cell shape probability densities for all groups, including wild type embryos morphologically stage-

matched to Lat embryos based on the number of cells in the midline. We plotted the deviation of

the four experimental groups from the wild type distribution over the range of cell shape values

observed (Fig. 6d').

We found that at 38 hpf, ablations of the IL and control ablations three cell diameters

anterior to the IL did not prevent the formation of square cells nor the establishment of the

ectodermal square grid (Supplementary Fig. 8). Additionally, 38 hpf embryos in which we made

a single cut in the IL cable maintained a straight line at the IL region of the tissue and began

midline formation 12 hours later (Supplementary Fig. 8). This is consistent with the observation

that 4-way vertices still form correctly after inhibition of myosin 2 phosphorylation at IL

establishment stages (Fig. 5e).

In contrast, in embryos in which 3-6 midline cells had aligned (58-69 hpf), ablations lateral

to the midline significantly perturbed the cell shape distribution on the ablated side of the embryo

(Lat-A, Fig. 6d) compared with the unablated side (Lat-NA), overall driving them towards a

rounder, rather than square, shape (Fig. 6b', d'). Relative to wild type, Lat-NA exhibited the

smallest increase in anisotropic cells (defined as those with a cell shape metric value of $\leqslant 0.5$;

circle (i) in Fig. 6d') and the smallest decrease in square cells (circle (ii) in Fig. 6d') of all four

groups.

Unlike the localized impact of lateral ablations, midline ablations significantly disrupted

cell shape distributions both on the ablated side (Mid-A) and on the contralateral control side (Mid-

NA) (Fig. 6c',d'). On both sides of midline ablated embryos, more cells took on more irregular

shapes, characterized by lower values of the cell shape metric (Fig. 6d). In particular, the

proportion of square cells was lower than wild type in both the experimental and the control

(contralateral) group of midline ablations (circle (ii) in Fig. 6d'). Midline ablation destroyed the

organization of the square-packed grid throughout the posterior germ rudiment (Fig. 6c'). In sum,

the effect of the lateral ablation was minimal and localized to the region immediately around the

ablation, while perturbing just one of the midline cables had a tissue-wide impact on cell shapes,

disrupting square cell formation and abolishing the square ectodermal grid.

Midline, but not lateral, ablations also disrupted midline elongation. Laterally-ablated

embryos developed midlines with total numbers of cells comparable to wild type (12-19 cells,

Supplementary Fig. 7) while midlines in midline-ablated embryos were shorter (5-10 cells,

Supplementary Fig. 7). This suggests that midline cable ablations, but not ablations lateral to the

midline, disrupted posterior intercalation at the midline. These data are consistent with the

hypothesis that the alignment of cells into the midline, driven by the contractility of the midline

cables, organizes cell shape across the tissue.

Finally, we asked whether establishment or maintenance of the square cell grid was

important for correct (para)segment patterning. At stage 11-12 of embryogenesis (66-68 hpf[29]),

when midline elongation by intercalation has completed (Fig. 4a"), the protein product of the

segment polarity gene *engrailed* (En) becomes detectable in two cells anterior to the IL, initially

in a few cells within the same row at or near the midline[29]. The En expression domain subsequently

expands in an increasing, left-right symmetrical number of lateral cells within the row[29]. This

expression marks the prospective mandibular segment (Mn, arrowhead in Fig. 1i, j)[29,44].

Subsequent rows of En-expressing ectodermal cells, delineating the remaining segments of the

body plan, arise correlated with the same stereotyped cell divisions of the cells in the grid that

contribute to midline elongation (Fig. 1j, k) [29,44].

Given that En expression in the prospective mandibular segment is detected right after the

midline intercalation phase that appears to create the first square cells in the grid, we asked whether

this expression was dependent on the tensile midline cables in the same way as square cells and

square-packing order. Across the WT and ablated samples, we found that the total number of

Engrailed-positive nuclei in the prospective mandibular segment was positively correlated with

the number of aligned midline cells (Supplementary Fig. 7b, Supplementary Table 3). In other

words, as the midline became longer, more lateral cells within Mn expressed En. Strikingly,

however, midline ablation reduced the number of En-positive Mn nuclei, even on the contralateral

control side of the embryo (Figure 6e, Supplementary Table 3). In contrast, lateral ablations had

little impact on the number of Engrailed-positive Mn nuclei on either the experimental or the

control side of the embryo (Figure 6e, Supplementary Table 3). Midline but not lateral ablations

also decreased the regularity with which Mn cells expressed En (Figure 6e, Supplementary Table

3). Given these results, we hypothesize that the first establishment of En expression across a

complete segment, as seen in Mn, requires midline-cable driven tissue organization.

Discussion

The role of cell lineage in initiating square cell formation

We found that descendants of the 16-cell stage ectodermal founder blastomeres segregate into

anterior and posterior compartments, and that the resulting compartment boundary (IL) forms the

first axis of the square cell grid. This suggests the possibility that cell lineage at least partly

underlies square cell morphogenesis. The results of our simulation suggest that cell sorting

between the cells of the anterior and posterior compartments, a phenomenon observed in multiple

contexts in animal development including amphibian embryos[58] and *Drosophila* imaginal discs[59],

regulates the formation of a straight compartment boundary. The mechanism that regulates this

cell sorting could in principle be one or more of differential cortical stiffness[60–62], differential

adhesion[61–67], or a mechanical barrier imposed by the supracellular actomyosin cable that forms at

the IL. Relevant to the latter hypothesis, we note that although the cable at the IL is under tension,

coupled tension and the phosphorylation of myosin 2 are not required to maintain 4-way vertices

at this boundary. This suggests that local tension at this IL cable is not necessary to maintain the

compartment boundary. Alternatively, it is possible that the interaction between the anterior and

posterior lineages can restore the tensile cable upon ablation.

The results of previous experiments demonstrating that the *P. hawaiensis* early embryonic

ectoderm has regulative capacity are interpretable as consistent with the cell sorting hypothesis[40].

In these experiments, individual 8-cell stage ectodermal blastomeres were photoablated and the

remaining two blastomeres were injected with a lineage tracer. The results showed that El and Er

can replace each other's anterior descendants, while Ep can replace their posterior descendants[40].

We hypothesize that compensation within the ectoderm, and ultimately grid formation, is possible

only when progenitors of both anterior and posterior lineages remain, and that cells from both

compartments are necessary to initiate square cell-packing geometry at the IL. Under this

hypothesis, both anterior and posterior cell types would be needed to form the boundary because

each lineage would contribute a distinct molecular state (e.g. cortical stiffness and/or adhesive

properties), the combination of which would then lead to emergent cell sorting.

In embryos of diverse animal species–including *D. melanogaster*[68–70], the flour beetle

Tribolium casteneum[54], the cnidarian *Hydra vulgaris*[71], the zebrafish *Danio rerio*[72], and the quail

Coturnix japonica[73]–mechanically integrated actomyosin cables spanning multiple cells play

crucial roles in epithelial morphogenesis. We hypothesize that differential genetically-determined

fate between the anterior and posterior lineages also underlies the assembly, and potentially the

shape organizing function, of the cytoskeletal cable we observed at the anterior-posterior boundary

(IL) from 38 hpf onwards, similar to anterior-posterior boundary formation in the *D. melanogaster*

wing disc[52]. In the early *Caenorhabditis elegans* embryo, both endodermal cell fate and the

phosphorylation of myosin 2, which are necessary for gastrulation in endodermal cells, are

dependent on the Wnt-Frizzled signaling pathway, suggesting a direct connection between cell

fate and morphogenesis[74]. A similar relationship could exist within the *P. hawaiensis* ectoderm,

the molecular determinants of which have yet to be identified.

Further ablation experiments in 16-cell stage embryos that eliminate all anterior or

posterior lineages, or elucidation of the transcriptomic, proteomic and mechanical states of the

anterior and posterior compartment clones, would be useful to elucidate the mechanism of cell

sorting at the IL. A hypothesis of one or both of differential adhesion or differential cortical

stiffness would predict that anterior and posterior cells have different levels of cell-cell adhesion

molecules and/or different cortical cytoskeletal behaviors. The fact that the IL marks a clonal

boundary could in principle provide a heritable mechanism of achieving differential molecular

states between clones. For example, anterior cells might inherit a level of adhesion molecules or a

cortical cytoskeletal conformation distinct from that of their posterior neighbors by virtue of

differential inheritance of cytoplasmic factors, or through inheritance of a gene regulatory state.

Lineage dependent and lineage-independent mechanical cues establish square cell packing

Once the IL has straightened into an actomyosin rich boundary axis, we found that a

midline cell founder population begins to intercalate perpendicular to the IL such that square cells

are added sequentially to the midline between two actomyosin cables. This grid axis does not

correlate with 16-cell lineage domain boundaries, such that the midline founder population is a

polyclone containing descendants of multiple ectodermal founders (Supplementary Fig. 1, 2).

These findings suggest that, unlike the IL, the formation of the midline and its cables is not

regulated by inherited differences among these lineages. We hypothesize that the midline grid axis

is instead initiated by mechanical stimuli, similar to how the formation of the primitive streak in

early avian embryos, which involves anterior-posterior oriented actomyosin cables[75], is driven by

tissue flows that regulate the expression of fate-determining genes.[9] Under this hypothesis, the

midline founder population would become defined–and, in turn, potentially genetically

differentiated–as a result of tissue-wide forces. Future work could examine whether fate-

determining genes of this population include components of the planar cell polarity signaling

pathway, which is involved in medio-lateral intercalation of cells in multiple developmental

contexts.[76–80]

We have provided evidence that the mechanical integrity of these midline cables is

necessary for the formation of isometric cells, the alignment of cells into rows, the maintenance

and packing of square cells across the ectodermal grid, and the establishment of segmentally

iterated En expression. We hypothesize that these cables drive tissue-wide mechanics, just as tissue

flows pattern the retina[81] and embryonic germ band extension[82] in *D. melanogaster*, or how

supracellular actin cables organize whole body regeneration in *H. vulgaris*[71]. In this framework,

the tension introduced by the midline cables at the intercalation zones could be the initial condition

necessary to trigger and perhaps subsequently balance the transition to square-packing. The

variance in recoil times in response to ablation at the IL and midline intercalation zones (Fig. 5c',

c"), suggests that tension at these regions could be dynamically modulated in response to global

tissue forces.

The relationship between square-cell packing, gene expression, and segmental patterning

In ascidian and mouse embryos, cell geometry is crucial for the expression of fate-determining

genes[83,84]. Our study suggests that in *P. hawaiensis*, square grid morphogenesis and subsequent

segment formation emerge from axes that are established according to differing degrees of

instruction from tissue mechanics and genes. We also provide evidence that expression of the

segment polarity gene *en* in the *P. hawaiensis* ectoderm is sensitive to mechanical perturbation

and dependent on the formation of the midline. Reminiscent of this finding, in *D. melanogaster*

embryos injected with paramagnetic-beads and subjected to ectopic tissue-wide forces with

magnetic tweezers, expression of En at parasegment boundaries was differentially altered

according to the location and magnitude of the force[85]. The authors of that study observed a change

in nuclear shape and redistribution of myosin 2 in tissue regions experiencing the applied force,

suggesting that expression of *en* is downstream of these mechanical and geometric parameters[85].

This raises the possibility that mechanical regulation of *en* expression could be a conserved feature

in multiple pancrustacean species.

499    In early *D. melanogaster* embryos, membrane fluidization by an ether treatment applied to

500  whole embryos results in larvae with a dramatic range of segmentation defects, from complete loss

501  of segments to the elimination of only a single segment, which resembles segmentation gene loss

502  of function phenotypes[86]. Our *P. hawaiensis* midline ablation experiments, in which cuts were

503  made in the exact same cell positions across samples, similarly resulted in a range of changes in

504  cell shape, midline cell number, and *en* expression, suggesting that segmentation in *P. hawaiensis*

505  could arise from a balance of tuned forces within the ectodermal epithelium. In this mechanical

506  framework, the ultimate effect of midline cable disruption could depend on the mechanical state

507  elsewhere in the tissue. Further experiments determining the longer-term effect of early midline

508  disruption, and the alteration of square-cell packing, on the formation of more posterior segments

509  will be necessary to establish the extent of the midline cables' organizing capabilities.

510    Beyond amphipods, previous comparative studies of embryogenesis in other malacostracan

511  orders (including Cumacea, Tanaidacea, Isopoda, and Mysidacea) have found that they generate

512  some of the cell material for their ectodermal grids through consecutive divisions of ectodermal

513  stem cells (ectoteloblasts) that provisionally align their clonal progeny into rows in the posterior

514  portion of the grid[87]. However, ectodermal grids in these species also contain cells derived from

515  non-ectoteloblast lineages, suggesting that some other mechanism of tissue organization must be

516  present[28]. Amphipod grids like that of *P. hawaiensis* completely lack ectoteloblasts and thus must

517  be organized by what Wolfgang Dohle and Gerhard Scholtz called a "row-forming factor," which

518  they hypothesized to be a "matrix forcing the cells into a grid-like pattern"[87]. Our results suggest

519  that this "matrix" emerges first from tensile cell-sorting at the IL and subsequently from

520  cytoskeletal cables at the midline. *O. cavimana*, another amphipod, exhibits stereotypic early

521  cleavage patterns similar to that of *P. hawaiensis*, suggesting that fate determination within the 8-

16 cell embryo could also play a role in grid morphogenesis of this species[46]. Further lineage

studies across the Amphipoda, the early embryos of which all undergo holoblastic cleavage[88], are

needed to understand the full extent of this mechanism within this group.

The amphipod anterior-posterior compartment boundary corresponds to a boundary

classically called the naupliar-post-naupliar boundary. Under this concept, the naupliar region of

the germ band, is the anterior region that will give rise to the head and mandibular segments,

formed through a combination of relatively uncoordinated mitoses, cell intercalation, and cell

rearrangements, and the post-naupliar region. In contrast, the post-naupliar germ band is formed

through the activity of either ectoteloblasts or segment precursor row duplications[44,45,89], and gives

rise to the trunk segments and telson. Interestingly, our work shows that the IL, positioned one cell

diameter posterior to the first complete segmental stripe of En expression in the prospective

mandibular segment, corresponds to the classically defined naupliar-post-naupliar boundary.

The anterior-posterior compartment boundary is established by the descendants of

ectodermal lineage precursors defined by stereotypical holoblastic cleavages[27,33] and likely also

by asymmetric inheritance of lineage potential to early blastomeres that begins at first

cleavage[35,90,91]. However, holoblastic cleavage is a derived state for the Malacostraca, in which

many extant species undergo syncytial cleavage[46], and the ancestral cleavage state is thought to be

meroblastic (in which cleavage furrows form but cytokinesis is not completed, effectively resulting

in a syncytium but with some partial yolk partitioning by the formation of incomplete furrows)[92,93].

We speculate that the evolution of holoblastic cleavage in amphipods facilitated the emergence of

lineage-based IL formation and the anterior-posterior compartment boundary as a mechanism for

establishing the ectodermal grid and generating the post-naupliar germ band. Under this

hypothesis, the evolution of an anterior-posterior boundary among the ectoderm clones eliminated

the need for patterning introduced by a row of proliferating ectoteloblasts within the germ disc.

With regards to grid organization along the anterior-posterior axis, we hypothesize that midline

formation via intercalation, and the importance of the mechanical organizing properties of the

midline for establishing correct segmental patterning, are likely to be conserved across the

Malacostraca. Further comparative studies should determine whether midline cables are present in

other species.

Methods

Animal culture and transgenesis

P. hawaiensis were cultured at 26 ºC in 16-quart plastic tanks containing 1.022 specific gravity

Instant Ocean artificial sea water (hereafter ASW) made with distilled water and fed a diet of

TetraPro PlecoWafers and fresh carrots. For live imaging of nuclei (Fig. 4c) we used a previously

reported[94] transgenic *P. hawaiensis* line in which the *D. melanogaster* histone H2B fused with a

monomeric Red Fluorescent Protein is under control of a *P. hawaiensis* heat-shock promoter

(hereafter *H2B-mRFPruby*).

Embryo collection and microinjection

Embryos for all experiments were collected and microinjected according to established

protocols[95]. Embryos were collected into and cultured in ASW filtered with a 0.22µm filter

(hereafter FASW). For precise staging and stage-matching among broods, only 1-cell through 8-

cell stage embryos were collected. All microinjected embryos were injected at the late 1-cell stage

once the chorion had hardened. Microinjected embryo cultures were supplemented with 1:100

Penicillin-Streptomycin (10,000 U/mL penicillin, 10,000 µg/ml streptomycin; Gibco 15140-122)

and 1:200 Amphotericin-B (250 µg/ml, Gibco 15290-026). The culture medium was changed out

every 1-2 days. For live imaging of the actomyosin cytoskeleton, embryos were injected with

180pl of 2 µg/µl of mRNA reporter constructs encoding either the actin-binding peptide LifeAct[96]

fused to the monomeric mNeonGreen fluorescent protein[97] (hereafter *LifeAct-mNeonGreen* or *LA-*

mNG) or the *P. hawaiensis* myosin II regulatory light chain gene (*Ph-sqh*) fused to mNeonGreen

(hereafter *Ph-sqh-mNG*). In vitro transcribed capped mRNAs encoding these reporters were

synthesized with the mMessage mMachine T7 kit (Invitrogen AM1344) from plasmid templates

pBlueSK-LifeAct-mNG and pBlueSK-Phsqh-mNG linearized with the restriction enzyme NotI, as

previously described (Kontarakis 2014). Plasmid templates pBlueSK-LifeAct-mNG and

pBlueSK-Phsqh-mNG were generated by PCR amplification of the *LA-mNG* and the Ph-sqh-mNG

open reading frames (with primer combinations pBlue_LA_GBS_F 5'-

agaactctgggggatctgatAatgGGCGTGGCCGATC-3' / pBlue_mNG_GBS_R 5'-

cggttcagaacttataacaTTACTTGTACAGCTCGTCCATG-3' or pBlue_Phsqh_GBS_F 5'-

agaactctgggggatctgatAATGTCGTCCCGCAAAGC-3' / pBlue_mNG_GBS_R) from vectors

pSL-PhHS-LA-mNG and pSL-PhHS-Phsqh-mNG, respectively (Kalogeridi and Pavlopoulous,

personal communication). PCR amplicons were cloned with Gibson assembly into ClaI-digested

pBlueSKMimRNA[98], replacing the Minos transposase ORF with the LA-mNG or the Ph-sqh-

mNG ORFs, respectively. For long-term lightsheet imaging of cell membranes and nuclei,

embryos were co-injected with in vitro transcribed capped mRNAs synthesized with the

mMessage mMachine SP6 kit (Invitrogen AM1340) either from plasmid template pEXPTol2-

H2A-mCherry (encoding the nuclear fluorescent reporter H2A-mCherry) linearized with the

restriction enzyme BglII[99] or from plasmid template PCS2 + Lyn-EGFP (encoding the membrane

fluorescent reporter Lyn-EGFP) linearized with the restriction enzyme NotI[100] .

Immunohistochemistry and drug treatments

Embryos were dissected away from their outer membrane according to established protocols[101]

and fixed for one hour at room temperature in 4% paraformaldehyde made up in FASW. Fixed

tissue was immediately transferred into a blocking solution (1x phosphate buffered saline (1x PBS)

+ 5% normal goat serum + 0.1% Triton X-100) or stored at 4 ºC in 1x PBS + 0.1% Triton X-100

for up to three days prior to staining. Embryos were stained with 1:100 AlexaFluor 488 Phalloidin

(Invitrogen A12379) or Rhodamine Phalloidin (Invitrogen R415) for F-actin, 1:1000 rabbit anti-

Myosin 2 656[102], 1:80 mouse anti-Engrailed Mab 4D9[37], 1:500 mouse anti-phospho-histone H3

(Cell Signaling 9706S), or 1:50 rabbit anti-cleaved caspase-3 conjugated to AlexaFluor 555 (Cell

Signaling 9604S) overnight at 4ºC. We followed these incubations with a solution containing

1:10,000 Hoechst 33342 (Sigma B2261, 10 µg/µl stock solution) as well as secondary antibodies

1:500 AlexaFluor 647-conjugated Goat Anti-Mouse (Invitrogen A31625) or 1:500 AlexaFluor

555-conjugated Goat Anti-Mouse (Invitrogen A31621) in blocking solution for two hours at room

temperature. Prior to fixation, drug-treated embryos were cultured for 12 hours at 26ºC in a

solution of Rockout (Rho Kinase Inhibitor III, Sigma 55555-3) dissolved in anhydrous DMSO and

added to a 30mm petri dish containing 6ml of 50% FASW, as previously described[36]. We found

that a 50µM concentration of Rockout did not disrupt normal cell proliferation and that 100% of

38 hpf-treated embryos survived up to two days in a continuous treatment (Supplementary Fig. 6).

Embryos cultured in 50% FASW alone at 26ºC, or with the equivalent amount of DMSO, hatched

normally (Supplementary Fig. 6).

Microscopy

Embryos for multiview lightsheet imaging were injected at the 1-cell stage with 2 μg/μl *H2A-*

mCherry (nuclei) and *Lyn-EGFP* (membrane) mRNA reporter constructs. We mounted 4-cell

stage embryos inside of a 2mm PTFE tube and 3mm glass capillary suspended in 1% low melt

agar made in filtered ASW and infused with 500 nm fluorescent beads (F-X050, F-Y050 or F-

XC050 microspheres, Estapor Merck). Capillaries were mounted on a SiMView multiview

lightsheet microscope[103] and embryos imaged with a 488 nm (*Lyn-EGFP*) and 594 nm (*H2A-*

mCherry) laser at ten-minute intervals from three or four angles with two objectives positioned

180º apart. The z-interval was set to 0.41μm. The resulting six or eight views per sample were

registered and fused using the MultiView Reconstruction and BigDataViewer plugins for Fiji[104–

107].

625   Live imaging of *H2B-mRFPruby* transgenic embryos and *LA-mNG* or *Ph-sqh-mNG* mRNA

injected embryos was carried out at 26ºC on a Nikon CSU-W1 SoRa Spinning Disc confocal

microscope with a 20x objective. Embryos were mounted individually on glass microscope slides

so that a 1.5 coverslip with clay-feet spacers could be used to roll the desired region of the germ-

disc or grid into the field of imaging. We applied slight pressure to the coverslip to flatten the

tissue and obtain clearer views of apical cell face geometries. We then filled the space between the

coverslip and slide with FASW and sealed the edges with Vaseline to prevent evaporation. Time-

lapse imaging of live embryos was carried out at ten-minute intervals with a z-interval of 0.9μm.

Fixed and stained embryos were cleared in 70% glycerol and mounted individually in the same

manner as live embryos for confocal imaging on either a Nikon CSU-W1 SoRa Spinning Disc

(0.9µm z-interval) or Zeiss LSM 880 (1µm z-interval). Only timelapse datasets taken of embryos

that hatched successfully were used for analysis.

Image analysis and segmentation

Lineage tracing of the fused lightsheet datasets was performed with either semi-automated

(Embryo 1; Supplementary Table 1) or manual tracking (Embryos 2 and 3). The nuclear channel

of Embryo 1 was first pixel classified and then object tracked in ilastik[108] (ver. 1.3.3), followed by

manual correction in MaMuT[94] (ver. 1.53), a plugin for Fiji. Embryos 2 and 3 were tracked entirely

manually using both the nuclear and membrane channels in MaMuT [94] or Mastodon in Fiji. We

created the 2D conformal maps of the embryo's surface from the lightsheet datasets using a version

of ImSAnE[109] customized for *P. hawaiensis*[34]. Nuclear tracks colored according to 16-cell stage

lineages were superimposed onto the maps using a custom MATLAB script or manually

superimposed in Adobe Illustrator (CC 2023 or 2024). Maximum intensity projections of the

lightsheet 2D conformal maps of *Lyn-EGFP* (membrane marker) were segmented in the Fiji plugin

TissueAnalyzer[110] using a watershed algorithm with manual correction.

To visualize and quantitatively describe cell shapes, we first pixel-classified membranes in

10µm maximum intensity projections of Phalloidin-stained or *LA-mNG*-labeled embryos in

ilastik[108] (ver. 1.4.0) to generate a more consistent cell boundary signal. Especially in the later grid

stages (approximately 62 hpf onwards), the embryo must be carefully laid flat against the slide and

slightly compressed with the coverslip. For samples in which flattening out all wrinkles or

depressions in the tissue was not possible, the probability maps containing apical cell faces for

each tissue region were cut out and composited together using the Image Calculator in Fiji. We

then fed the probability maps exported from ilastik[108] (ver. 1.3.3) into Cellpose[111] (ver. 3.0) and

obtained cell ROIs using a custom-trained model with some manual correction of contiguous

masks. Automatic size calibration was performed and non-contiguous or incomplete cell masks

around the edges were removed. All subsequent quantitative descriptions of cell shape and tissue

organization were carried out on these ROIs using custom Python scripts available at

https://github.com/boyonpointe/parhyale_morpho/tree/main with commit ID a19a88d. All other

measurements, including intensities of cytoskeletal components, were made on 10µm maximum

intensity projections in Fiji (ver. 2.1.0) using the ROI section and associated measurement tools.

For cell shape and vertex IL measurements, each 10 µm maximum intensity projection

was cropped to 207 µm wide x 104 µm high. The crop box was centered on the germ disc

parallel to the dorso-ventral axis and centered on the midpoint of the IL (a clear straight or

slightly bent line) in the anterior-posterior axis. Only cells contained fully within the crop box

were counted.

For IL actin and myosin 2 measurements, in un-cropped 10 µm maximum intensity

projections, the lateral extent of the IL region of interest was determined by drawing a line from

the ventral midline of the germ disc towards the left and right sides of the germ disc. The region

was considered terminated when it reached cells whose sides were bent more than approximately

15º from the leading edge of the region. The mean IL length for phalloidin-stained samples (F-

actin) was 146 µm (n = 38) and the mean IL length for *Ph-sqh*-labeled samples (myosin 2) was

151 µm (n = 14).

To compare cell shape distributions in the posterior of the germ disc, we used a bootstrap

approach. For each pair of distributions, we generated 100,000 surrogate pairs of random

resampling and calculated the difference in their means. This gave us a null distribution of

differences, against which we assessed how often the observed empirical difference would occur.

The p-value corresponds to that probability. For stop motion in Movie 3, images were printed on

light-permeable paper and animated over a lightbox using a Sony A7S camera controlled by the

stop-motion animation software DragonFrame (v4).

Cell shape descriptor

To measure cell shape, we used a custom descriptor in which the area of a cell face is divided by

the area of the circle whose diameter is equal to the distance between the farthest two points on

the cell's perimeter. With this descriptor, a perfect square has a value of 0.64, and hexagons and

higher order polygons have values of approximately $\geqslant 0.75$. We found this measurement to more

usefully describe the emergence of square-cell packing than counting the number of cell sides,

because within this dynamic tissue, 4-way vertices frequently dissolve into small edges at square

corners, and cell divisions can introduce temporary kinks into neighboring cell edges.

Cellular Potts Model Implementation

Our model to test the effect of differential adhesion between different cell lineages was formulated

in the Cellular Potts Model (CPM)[41,67,112]. We implemented our model using Compucell3D[43]. In

CPM, the cells are modeled as a spatially contiguous collection of pixels. The pixels that are at the

boundary of cells have an energy cost associated with interacting with pixels belonging to the

neighboring cells. The energy cost between a pair of pixels (a,b) depends on the cell type

$(\sigma(a), \sigma(b))$ of both the cells to which the interacting pair of pixels belong. The area of (A) of a cell

is defined as the number of pixels the cell encompasses while the perimeter (L) is the number of

pixels at the boundary of the cell.

The following equation shows the expression of energy of the system for given pixel

configuration:

705
$$E = -\sum_{a,b} J\big(\sigma(a), \sigma(b)\big) - \sum_{\sigma} \lambda_A \big(A_\sigma - A_0(t)\big)^2 - \sum_{\sigma} \lambda_L \big(L_\sigma - L_0(t)\big)^2$$

The second and third terms on the right hand side of the equation represent the energy cost for cell

area and cell perimeter to deviate away from the steady state values A_0 and L_0 respectively. To

implement growth, A_0 is made a function of time such that the cell grows to reach A_0 and when it

meets the size to undergo division, it divides. The steady state value for the perimeter is also a

function of time and set to ensure that steady state shape is circular i.e. $L_0(t) = 2\pi\sqrt{A_0(t)/\pi}$. The

symmetric matrix J has its number of rows and columns equal to the number of cell types in the

system. Each entry $J_{i,j}$ in the matrix specifies the interaction energy between pixels belonging to

cell type i and j. In our simulation there are six different cell types corresponding to six different

lineages and an additional one that corresponds to the medium.

The segmented cell masks of tissue imaged at 26 hpf is used to generate the initial

configuration of the pixels in the simulation. Each pixel belongs to one among the 6 different cell

types or to the medium. The cells increase in size by 100 pixels before they divide. The amount of

growth cells undergoes before they divide is chosen to reflect the empirically observed increase in

tissue size and cell numbers seen at 46 hpf. The cells divide symmetrically along the axes

orthogonal to the major axis (long axis). Throughout the simulation, we track the number of pixels

in contact with pixels from cells of different lineages.

Ablation experiments

Ablation experiments were carried out on a Zeiss 980 NLO 2-photon microscope with a pulsed

Spectra-Physics InSight X3+ laser set to 927 µm wavelength and an ablation ROI with a 1µm

width that was passed over five times. *LA-mNG* embryos injected at the 1-cell stage were incubated

to the desired stage and mounted individually on slides with a Vaseline-sealed coverslip. Ablated

embryos were imaged with a LD LCI Plan-Apochromat 25x/0.8 objective for two frames prior to

the ablation followed by every second for two minutes immediately afterwards. Some embryos

were further imaged on the same microscope at intervals of one minute to follow the effects of the

ablation. Embryos used for further analysis were then recovered from the slide and transferred to

a 35mm petri dish with fresh FASW with 1:100 Penicillin-Streptomycin (10,000 U/mL penicillin,

10,000 µg/ml streptomycin; Gibco 15140-122) and 1:200 Amphotericin-B (250 µg/ml, Gibco

15290-026) for incubation at 26 ºC until fixation. Recoil length was calculated in Fiji by

subtracting the distance between cell vertices on either side of the ablation ROI in the first frame

immediately post ablation, from the distance between the same two vertices before the incision

was made.

Figure presentation

Line tracing of membranes shown in figures was performed using the Image Trace function

(Silhouettes or Line Art) on exported Cellpose[111] ROIs or on hand segmented micrographs

(Supplementary Fig. 3) in Adobe Illustrator 2024 or 2025. To more clearly display the

relationship between Engrailed expression and grid geometry in Figure 1, we used ilastik[108]

(ver.1.3.3) to pixel-classify images of anti-Engrailed staining and exported the resulting

probability maps. In Movie 4, TissueAnalyzer[113] was used to track and visualize cell membranes

of the midline founder population in a conformally mapped time-series that was manually

aligned. All figures were assembled with Adobe Illustrator.

Acknowledgements

The authors thank William Lemon and Philipp Keller (HHMI Janelia Research Campus) for providing access and training on their SimView lightsheet microscope, Sebastian Streichan and Dillon Cislo (University of California, Santa Barbara) for help with ImSAnE and applying MaMUT tracks to ImSAnE maps, John Rallis and Myrto Ziogas (IMBB-FORTH) for help with plasmid preparations, as well as members of the Extavour and Pavlopoulos laboratories for discussion.

Author Contributions

BS conceived of the study, designed experiments, performed all wet lab experiments, performed data analysis and interpretation, and wrote the first draft of the manuscript. LB designed, performed embryo collection and injection experiments for, and executed capture of lightsheet datasets, and reviewed the manuscript. CK designed and performed all computational modelling experiments, performed data analysis, and reviewed the manuscript. ES and AP conceived of the study, designed and created all fluorescent *P. hawaiensis* constructs, performed embryo collection and injection experiments for capture of lightsheet datasets, and reviewed the manuscript. CGE conceived of the study, performed data interpretation, obtained funding for the study, supervised its execution, and edited the manuscript.

Funding

This work was supported by a National Science Foundation (NSF) Graduate Student Research Fellowship 2018259071 and a Simmons Award from the Harvard Center for Biological Imaging to BLS, NSF award IOS-2220747 to CGE, the NSF-Simons Center for Mathematical and Statistical Analysis of Biology at Harvard (award number DMS-1764269) and the Harvard Quantitative Biology Initiative (CK), Human Frontiers Science Program Grant RGP022/2023 and HHMI Janelia Research Campus Visitor Program to ES and AP, and funds from the Howard Hughes Medical Institute (HHMI) and Harvard University. CGE is an HHMI Investigator.

Conflicts of Interest

The authors declare no conflicts of interest.

Data Availability

Lightsheet datasets described in Supplementary Table 1. MaMuT lineage tracks for these datasets, as well as scripts for cellular potts model simulations and image analyses, are available at https://github.com/boyonpointe/parhyale_morpho/tree/main with commit ID a19a88d.

Statement on Use of Machine Learning, Artificial Intelligence and Large Language Models

ChatGPT 4.0 and 5.0 (open AI) were used to modify R code for altering plot aesthetics. ilastik[108] and Cellpose[111], used for image segmentation in this study, are machine learning-based programs. ML/AI was not used to generate any content, write any text, or perform any other data analysis.

Figure 1 (previous page). Formation of a cellular square grid in the *Parhyale hawaiensis* embryonic
ectoderm. a Developmental timeline from fertilization to the hatching of juveniles with the window of
lightsheet microscope imaging indicated at the bottom. Stages according to Browne and colleagues[29]. **b-f**
Stills selected from a time-lapse of the first 1.5 days of development in a *P. hawaiensis* embryo that was
live-imaged with a multiview lightsheet microscope; cell membranes marked with *Lyn-EGFP*. Micrographs
are maximum intensity projections of a volume constructed by fusing 3D image stacks from all imaging
angles, and are displayed either from a fixed perspective (b-f) or as conformal surface maps (c'). **b** Ventral
(left) and dorsal (right) views of the 16-cell stage. The first cell divisions in *P. hawaiensis* are highly
stereotypic and segregate the germ layers and germ line (gl and gr) by this stage. The eight micromeres and
two macromeres enclosed within the blue dotted line are the mesoderm (ml, mr, Mav), endoderm (en), and
germ-line (g) precursors, the descendants of which delaminate from the surface during gastrulation (c-e)
leaving at the surface a continuous sheet of epithelial ectoderm (f). The remaining six macromeres (Epr,
Epl, Era, Erp, Ela, Elp) will give rise to all ectodermal cells. **c-e** From approximately 16-22 hours post
fertilization (hpf), cells begin reducing in size and migrate to the anterior pole, condensing into a germ disc
(embryonic rudiment) sitting atop larger yolk-filled cells. **c'** Anterior (left) and posterior (right) conformal
maps of the same image volume displayed in c. These 2D surface projections allow for simultaneous
visualization of cell shapes and relationships that make up the embryo's curved surface. The maroon dotted
line in c corresponds to those along map edges in c'. **f** By approximately 38 hpf, the surface of the germ
disc is an epithelium made up entirely of ectoderm (colored cells). **g** An adult female *P. hawaiensis*. White
lines indicate parasegmental boundaries with corresponding morphological segment identities. **h-k** Time
series of fixed embryos in which the ectoderm was dissected away from the yolk and physically flattened
to make surface cell geometries visible. A grid of four-sided cells emerges within the expanding germ disc.
The transition from hexagonal to square cells (**h'-k'**) progresses from anterior to posterior and laterally left
and right (**i**). A row of cells expressing the segmentation gene *engrailed* appears in the prospective
mandibular segment (Mn) around 68 hpf (arrowheads). **j, k** From 80 hpf onwards, progressing from anterior
to posterior, nuclear-localized Engrailed protein is detectable in cells of the anterior-most cell row in each
parasegment as it is generated via row-duplicating cell divisions from the midline to lateral regions. Dotted
lines in j indicate the region of the embryo displayed in k. **l** Cell shape descriptor used to analyze square
grid morphogenesis in this study. While four-sided cells can be either square or rectangular, for simplicity
we describe them as square throughout the text, and our cell shape descriptor gives square cells a metric
value of 0.64. Mx1: first maxillary segment; Mx2: second maxillary segment; T1: first trunk segment; T2:
second trunk segment; T3: third trunk segment.

Figure 2. Square cell geometry initiates along the dorso-ventral axis at the initiation line, which is an ectodermal lineage compartment boundary. **a-b** 38-68±2 hpf manually flattened germ discs segmented to identify cell membranes, with cells colored according to their shape descriptor value. What we call the "initiation line" (IL; arrowheads in the left column), corresponds to the boundary of the anterior (A: grey) and posterior (P: striped) lineage compartments shown in a'. **a'** Stills from a lightsheet time-lapse that were conformally mapped onto 2D, showing descendants of ectodermal precursor macromeres (grey: anterior ectoderm founders Era, Ela; striped: posterior ectoderm founders Erp, Elp, Epr, Epl) The boundary between anterior and posterior clones begins as a jagged curve at 38±2 hpf and straightens out by 68±2 hpf. **c** Top: Schematic shows the analyzed region (box) within the germ disc (dotted line). Bottom: Percentage of 4-way vertices at the IL (gray), the equivalent line one cell anterior (A1, light pink), and one cell posterior (P1, light green). **d** Top: Cell rows analyzed in four different locations relative to the IL: immediately adjacent to the IL to the anterior (A1: light pink), immediately adjacent to the IL to the posterior (P1: light green), one cell diameter anterior to the IL (A2: dark pink) and one cell diameter posterior to the IL (P2: dark green). Bottom: plot showing percentage of square cells (shape descriptor value ⩾ 0.60 or ⩽0.66). Anterior is up in all images. Sample sizes in c and d indicated by n1: number of embryos and n2: number of vertices.

Figure 3. A simulation of cell sorting can explain the emergence of a straight boundary at the initiation line. a Tracing of a 16-cell stage
embryo with the six ectodermal lineages colored. Grey and striped legend is provided to show the correspondence of these blastomere lineages to
the anterior and posterior compartments shown in Figure 2. **b** Simulation contact energy inputs for each combination of lineage precursors (colored
circles show blastomere identities as defined in a) as well as the environment surrounding the tissue (white circles). Higher contact energy results in
decreased affinity between cells. **c** Change in normalized area of overlap plotted against time in Monte Carlo steps (mcs) for n = 10 runs of the
simulation. Group I (circles) = boundaries among the four posterior lineages (Erp, Elp, Epr, Epl); group II (triangles) = boundary between the two
anterior lineages (Era, Ela); group III (diamonds) = boundary between the anterior and posterior lineages. **d** Membrane segmentation of part of the
conformally mapped germ disc of Embryo 1 (Supplementary Table 1) at 26 hpf. At this stage, gastrulation has already taken place, and the surface
of the embryo is a continuous sheet of ectoderm. This segmentation was seeded as the starting point for the simulation. **e** Stills from three time points
of the simulation of cell sorting among the 16-cell stage ectodermal lineages implemented in CompuCell3D with time denoted in arbitrary units.
Cell division rates for each lineage from one of the lightsheet datasets were input into the model. Anterior is up in all embryo image panels.

Figure 4 (previous page). The secondary grid axis emerges as a column of square cells via the intercalation of a polyclonal founder cell population. a At 52-62 hpf, primarily anisotropic ectodermal cells near the ventral midline and spanning the IL (arrowheads in a', a", a'"), in a region we denote the intercalation zone (IZ), intercalate simultaneously in the anterior and posterior directions to form a single column of square cells forming the midline of the embryo (**a'**). **a"** The cells within the midline are the first cells to become square. Cells flanking the midline on the left and right then adopt square shapes posterior to the IL. Cells in a', a", and a'" are colored according to the cell shape descriptor. * in a" and a'" indicates midline cells that have undergone parasegment precursor row duplication. **b** Cells that join the midline derive from a polyclone of founder cells that span the anterior (grey)-posterior (stripes) compartment boundary (arrowhead). **b'** Founder cells intercalate in both anterior (IZA) and posterior (IZP) intercalation zones (as shown in a', a" and a'"), adding or subtracting sides until they form a square. Arrows indicate the direction of a growing or shrinking cell side. Dotted outlines in rightmost panel of top row and second panel of bottom row indicate the outlines of the sister cell resulting from the division in the previous panel. **c** Numbers of total cells in the midline over time (n = 5 embryos). Midline initiation is defined as the time when the first posterior cell (P1) is aligned. By 12 hours after midline initiation, a mean of nine total cells have been added to the midline. **c'** Numbers of cells in the midline anterior (A ML: pink) and posterior (P ML: green) to the IL over time (in = 5 embryos). Anterior is up in a-a'" and b-b''.

Figure 5 (previous page). The initiation line and midline borders are tensile cytoskeletal cables. a
Maximum intensity projection of 20 0.5µm optical sections of the ventral ectoderm of a 44±2 hpf embryo
stained with phalloidin to label F-actin. **a'** Ratios of mean F-actin intensity measured in 38-56 hpf embryos.
The regions quantified (A1, IL, P1, P2) are shown in **a**. Each region begins at the arrowhead on the right of
the micrograph and extends across approximately 10 contiguous cell sides to the left side of the micrograph.
b Maximum intensity projection of six 0.9µm optical sections of a 68-74 hpf embryo with a 15-cell midline;
F-actin is labeled with phalloidin as in a. Arrowheads indicate the six regions (lines running parallel to the
anterior-posterior axis, named MR, MR1, MR2, ML, ML1, ML2) where F-actin intensity was quantified.
Each region quantified begins at the arrowhead shown in **b** and extends towards the posterior along
contiguous cell sides to the posterior IZ (black arrows). **b'** Ratios of mean F-actin intensity measured in 68-
74 hpf embryos with 7-15-cells in the midline. **c** 38-44 hpf embryo subject to laser ablation at the IL. **c** Pink
rectangle indicates the ablation site; white arrows connected by a dashed line indicate the lengths measured
to calculate recoil; blue arrowheads at IL indicate vertices surrounding the incision site. **c'** Ablation sites
(top: thick line indicates the IL) and recoil plot (bottom) in 38-44 hpf embryos in which recoil was measured
following IL ablation. **c"** Ablation sites (top: thick lines indicate the cables demarcating the nascent
midline) and recoil plot (bottom) in 58-69 hpf embryos with 3-7 cells in the midline. Pink numbers on X-
axis categories indicate the ablation site as shown in the schematic above. 7/6 indicates that a first incision
was made at position 6 (next to the midline P1 cell/at the compartment boundary), followed by a second
incision at position 7 (posterior IZ). In 6/7, the same two incisions were made in reverse order. **d** 1-cell
midline embryos (50-56 hpf) were incubated in 50µM DMSO or Rockout (**d'**), and the cells of the posterior
compartment were segmented and their cell shape descriptor was color-mapped 12 hours later. The IL is
indicated by yellow arrowheads. **e** Percentage of 4-way vertices in 38 hpf embryos incubated in DMSO and
50µM Rockout for 12 hours. P-values for comparisons between DMSO and RO-treated samples as follows:
A1: 0.48; IL: 0.18; P1: 0.48. **f** Total number of midline cells is significantly decreased in 62-68 hpf embryos
treated with Rockout for 12 hours (p-value = 0.004). **f'** Total number of cells in the posterior compartment
of 62-68 hpf embryos treated with Rockout for 12 hours is not significantly different from DMSO controls
(p-value = 0.35). **g** Cell shape values for cells aligned into the midline in 12-hour DMSO (n = 35 cells) and
Rockout (n = 12 cells) treated embryos. Grey symbols indicate the single cells at the anterior-most and
posterior-most positions of the midline; black symbols indicate all other midline cells. Samples are the same
as those plotted in f and f' (p-value = 0.002). * p-value < 0.05; ** p-value < 0.01; *** p-value < 0.001; n.s.
not significant (Mann Whitney U-test; all tests one-tailed). Scale bars for a, c', d = 25µm and c = 15µm.
Anterior is up in all micrographs. Sample sizes (number of embryos) in all plots indicated by numbers
underneath category labels.

a

b Lateral ablated (Lat)

3-6 cell
midline

Ablated (Lat-A)   Not ablated (Lat-NA)

c Midline ablated (Mid)

Ablated (Mid-A)   Not ablated (Mid-NA)

Cell
shape

b'   +9-12
hours

c'

d   d'   e

	WT	Lat-NA	Lat-A	Mid-NA	Mid-A
n1	2	3	3	5	5
n2	*1400*	*300*	*300*	*400*	*399*

Figure 6 (previous page). The midline cables organize square-cell packing and the emergence of the first complete stripe of Engrailed-positive nuclei. a The region within the germ disc where ablations were made, delineated by the dotted line. **b, c** Locations of laser ablations performed in different regions of the ventral epithelium in 58-69 hpf embryos with 3 – 6 cells in the midline. **b', c'** Cell shape descriptor values and En expression 9-12 hours after ablation. **d** Cell shape descriptor value distributions across treatments, including unablated embryos (WT) of equivalent chronological age. Yellow circles indicate the mean, dotted line the value for an isometric square. Only significantly different pairwise comparisons (based on the bootstrapping approach described in Methods) are indicated as follows: * $p < 0.05$; ** $p < 0.01$. Sample sizes indicated by n1: number of embryos and n2: number of cells. **d'** Deviation of the four ablated groups from the WT probability density ($\Delta_p = 0$) over the range of cell shape values observed. A positive value indicates shapes gained and a negative value indicates shapes lost. Dotted vertical line indicates the value for an isometric square; colors are the same as in panel d. **e** Proportion of Mn cells (categories on horizontal axis) expressing En in wild type and in embryos subjected to lateral (brown) or midline (green) ablations shown in b and c respectively. Data in Supplemental Table 3.

Movie Legends Download Link

All movies show Embryo 1 (see Supplementary Table 1).

Movie 1. **The first 2.5 days of development of a membrane-labeled *P. hawaiensis* embryo**. Timelapse animation of the *Lyn-EGFP* (membrane) channel of a reconstructed multiview lightsheet dataset beginning at the 8-cell stage and ending during germ band grid elaboration. Each frame is a maximum intensity projection through half of the embryo and was taken ten minutes apart. 319 total frames are animated at a frame-rate of nine frames per second. Sequential frames from this animation were selected for and sequentially displayed in Figure 1 (first frame is Fig. 1b).

Movie 2. **The first 2.5 days of development of a nuclear-labeled *P. hawaiensis* embryo**. Timelapse animation of the *H2A-mCherry* (nuclei) channel of the same embryo shown in Movie 1. Each frame is a maximum intensity projection through half of the embryo and was taken ten minutes apart. 319 total frames are animated at a frame-rate of nine frames per second.

Movie 3. **Dynamics of the 16-cell stage lineage domains visualized by stop-motion animation**. The left side of this animation shows a still from Movie 1 with the *Lyn-EGPF* marked 16-cell stage ectodermal blastomeres false colored (Era = magenta, Ela = pink, Erp = yellow, Elp = purple, Erp = cyan, Elp = orange). The right side shows 2D maps of the same dataset in which the MaMuT nuclear tracks have been superimposed on the membrane channel to show the lineage of each cell. The rectangular cut-out corresponds to the cropped regions of each map shown in Fig. 2c, Fig. 6a, Supplementary Fig. 1 and Supplementary Fig. 8a. The maps shown here begin at 42 hpf and end at 55.67 hpf and are animated at a frame rate of nine frames per second, displaying 1.5 hours of developmental time every second. Given that the mapping of the 3D surface to 2D was not constrained and the germ disc migrates across the yolk, frames were manually aligned to produce animated motion and keep the emerging grid fixed within the cut-out region. The first sweep of the pointing finger across the map (00:07) indicates the IL (Fig. 2) at the anterior-posterior compartment boundary. The second sweep at the end (00:10) indicates the emerging midline (Fig. 4), which separates the Erp and Elp lineages by one cell beginning just posterior to the IL (P1 in Fig. 4b).

Movie 4. Intercalation of midline cells to form a column of square cells. The midline of the germ band grid forms as a single column of cells via intercalation. Cells aligned into the midline and within the intercalation zone at 66±2 hpf were tracked backwards to the onset of midline formation.

Supplementary Figure 1. 16-cell ectodermal lineage domains undergo extensive rearrangement. Stills from long-term multiview lightsheet timelapses shown either as tracings of maximum intensity projections (a) or 2D conformal maps (b-c) in which lineage was determined by tracking each cell back to its 16-cell stage ancestor. **a** 16-cell ectoderm precursor blastomere identities for Embryo 1 (Supplementary Table 1) colored with reference to anterior (grey)-posterior (striped) compartments or 16-cell stage identity (colors). **b-c** Lineage domain comparison between Embryo 1 and Embryo 2 (Supplementary Table 1) showing that in both embryos, anterior lineages (Era, Ela) do not mix with posterior lineages (Erp, Elp Epr, Epl). In the posterior compartment, the Erp and Elp domains begin on opposite (left and right) sides of the germ disc at 26±2 hpf and migrate closer to one another until they are separated by only a single column of Epr or Epl cells by 56±2 hpf, indicating the presence of extensive cell rearrangement.

972
973
974 **Supplementary Figure 2. Cell proliferation occurs throughout the germ disc from 32-30 hpf, with IL-adjacent divisions oriented**
975 **preferentially perpendicular to the IL. a** Cell division angles measured across the germ disc in 32-50 hpf embryos (sample size in parentheses).
976 Angles are plotted with reference to the orientation of the IL (0°) and colored according to the division region (anterior = pink, IL = grey, posterior
977 = green). **b** Segmented germ disc indicating locations of defined tissue regions. IL cell divisions are those that occur in cells directly adjacent to the
978 IL on the anterior or posterior side (grey).
979

980
981
Supplementary Figure 3. Multiple 16-cell stage ectodermal lineages contribute to the midline,
including non-Ep clones in the posterior. a Legend for 16-cell stage ectodermal precursor blastomeres.
Midline descendants shown in c and e. **b** Legend for 16-cell stage Ep and non-Ep ectodermal precursor
blastomeres. **c, e** The midline is composed of descendants of multiple 16-cell blastomeres both anterior and
posterior to the anterior-posterior compartment boundary, which we call the IL at earlier stages. Midline
cells in 12-cell midline embryos were tracked back to the 16-cell stage in the lightsheet datasets. At the 1-
cell midline timepoint, only cells whose descendants make up the midline or intercalation zones shown in
the 12-cell midline timepoints, are shown. **d, f** The posterior midline is not exclusively made up of
descendants of Ep. Arrowheads indicate the location of the compartment boundary. In c through f, Embryos
1 and 2 correspond to the same datasets shown in Supplementary Figure 1 and described in Supplementary
Table 1.

994
995

Supplementary Figure 4. Myosin 2 accumulates at the IL and midline cables. a 10μm maximum intensity projection of a 56±2 hpf embryo injected with *Ph-sqh-mNG* mRNA to label the myosin 2I regulatory light chain. **a'** Ratios of mean intensity of *Ph-sqh-mNG* signal measured from 38-56 hpf embryos in 10μm maximum intensity projections like that shown in a. The regions where *Ph-sqh* signal was quantified (A1, IL, P1, P2) are shown in a; each begins at the arrowhead and extends across contiguous cell sides to the rightmost edge of the region shown. **b** Maximum intensity projection through 5μm of a 74 hpf embryo at the 6-cell midline stage. Arrows indicate regions of myosin-2 accumulation along the MR and ML cables (arrowheads), and at the anterior and posterior intercalation zones (IZ). **b'** Ratios of mean intensity of *Ph-sqh-mNG* signal measured from 5μm maximum intensity projections labeled with *Ph-sqh-mNG* expression . The regions quantified (MR, MR1, MR2, ML, ML1, ML2) are shown in b; each begins at the arrowhead and extends down contiguous cell sides to the bottom of the region shown. p = 0.05 for ML/ML1 compared to ML1/ML2; p = 0.20 for MR/MR1 compared to MR1/MR2. **c** 5μm maximum intensity projections of a 74 hpf embryo with 11 cells at the midline, labeled with an anti-Myosin 2 antibody. Only the anterior portion of the midline is shown here. Scale bars = 25μm and anterior is up in all micrographs. For all statistical comparisons * p-value < 0.05, ** p-value < 0.01, *** p-value < 0.001, and n.s. = not significant (Student's t-test for a; Mann Whitney U-test for b; all tests one-tailed). Sample sizes in a' and b' (numbers of embryos) indicated by numbers under category labels.

Supplementary Figure 5. IL and midline cables in the context of the whole germ disc. a 10μm maximum intensity projections of 38-56±2 hpf
embryos stained with phalloidin (left) and with membranes traced (right). IL (all stages shown) and midline cables (56 hpf only) demarcated with
thick lines. Anterior is up; scale bar = 100μm. **b** 10μm maximum intensity projections of the regions around the IL and midline cables in 62-68±2
hpf embryos. Arrows indicate the midline cables flanking the intercalation zone (IZ). Arrowheads indicate Engrailed expression in the prospective
mandibular segment (Mn). Anterior is up; scale bar = 20μm.

Supplementary Figure 6. 12-hour Rockout treatment does not impact cell proliferation. a 38±2 hpf embryos treated with either 50% FASW, 1:1 FASW:DMSO, or 50μM Rockout in 50% FASW were fixed 12 hours later (at 50±2 hpf), stained with Hoechst 33342 (top row) and anti-phospho-histone H3 (bottom row). Scale bar = 100μm and applies to all panels. **b** Total number of phospho-histone H3-positive cells for each treatment shown in a. n.s. = not significant (p-value = 0.303, two-tailed Mann-Whitney U Test). **c** Continuous treatment hatching assay. 100% of embryos incubated in 50μM Rockout continuously beginning at 38±2 hpf remained alive after 48 hours of treatment (86±2 hpf), and all died by 91 hours of treatment (129±2 hpf) (n = 11 embryos). By contrast, 100% of embryos in the 50% FASW (n = 8) and 1:1 FASW:DMSO (n = 11) treatments remained alive until 204±2 hpf. 87.5% hatched in 50% FASW and 90.1% of embryos hatched in 1:1 FASW:DMSO. Sample sizes in b indicated by numbers underneath category labels.

Supplementary Figure 7. a The bottom tenth percentile of the cell shape value (Fig. 1l) distribution of the whole tissue (i.e. the most
anisotropic/least round cells) correlates positively with the total number of midline cells for that sample, in both wild type and ablated embryos. For
wild type embryos, the roundness value of the bottom tenth percentile of cells is close to 0.6, implying that the shape of most non-round cells in wild
type tissues is a square. However, upon midline ablation, the roundness value for this bottom tenth percentile is well below 0.6, implying that cells
that would have been square in the wild type adopt more asymmetric shapes. The most dramatic impact on cell shape is experienced by the cells in
the midline, which are approximately 10% of the cells in the tissue at this stage (data not shown). **b** The total number of Engrailed-positive (En+)
nuclei correlates positively with the total number of midline cells.

1038

a

b **Anterior ablated (An)**

c **IL ablated (IL)**

38-44 hpf

Cell shape

0.8

0.7

0.6

0.5

+12 hours

b'

c'

1039
1040
1041 **Supplementary Figure 8. Ablated IL embryos maintain a straight line at the IL and form a midline. a** Schematic showing the location of the
1042 region delineated in b and c. **b** Ablation of a cell bond four cells anterior to the IL (n = 2). Ablation sites indicated by the pink rectangles. **b'** Cell
1043 shape descriptor values mapped on to +12 hour ablation tissue segmentations. **c** Ablation of a cell bond at the IL (n = 5). Ablation site indicated by
1044 the pink rectangle. **c'** Cell shape descriptor values mapped as described in b'. Arrowheads: IL.

1045

Lightsheet Sample	Total imaging time (h)	T-interval (min)	16-cell lineages tracking	Midline tracking	Posterior cell numbers	Cell division orientation	Tissue morphology seeds for simulation
Embryo 1	61.00	10	Yes	Yes	Yes	Yes	Yes
Embryo 2	61.00	10	Yes	Yes	Yes	Yes	No
Embryo 3	66.33	10	No	Yes	Yes	Yes	No
Figure			**Fig. 2, Supp. Fig. 1**	**Fig. 4, Supp. Fig. 2**	**Fig. 5**	**Supp. Fig. 3**	**Fig. 3**

1046

1047 **Supplementary Table 1. Lightsheet datasets used for analyses.** h: hours. min: minutes.

1048

| | | | | | Live: 6hr time window | | | | Fixed | | | | | | | |
| | | | | | Cell divisions in germ disc | | | | Cell divisions in germ disc | | | | Total cells in germ disc | | | |
hpf	N (live)	N (fixed)	N (total)		Total	A	IL	P	Total	A	IL	P	Total	A	IL	P
32	3	0	3	Mean	54.33	31.33	9.67	13.33	4.67	3.33	1.33	0.00	154.00	84.00	16.67	53.33
				SD	17.16	14.05	4.04	1.53	1.53	2.31	1.53	0.00	18.19	14.93	1.15	13.65
				SE	9.91	8.11	2.33	0.88	0.88	1.33	0.88	0.00	10.50	8.62	0.67	7.88
38	3	10	13	Mean	69.33	30.33	13.00	26.00	5.42	2.67	1.00	1.92	238.92	155.33	20.62	64.31
				SD	8.50	5.13	4.58	1.73	2.11	1.61	1.00	1.04	23.60	20.72	2.10	7.97
				SE	2.36	1.42	1.27	0.48	0.58	0.45	0.28	0.29	6.55	5.75	0.58	2.21
				A vs P					0.10565				**<0.00001**			
44	2	10	12	Mean	108.50	66.00	16.00	26.50	5.45	2.91	0.75	1.75	311.91	183.64	24.33	103.83
				SD	21.92	7.07	8.49	6.36	2.62	2.07	0.75	1.36	31.23	19.30	2.57	25.56
				SE	6.33	2.04	2.45	1.84	0.76	0.60	0.22	0.39	9.02	5.57	0.74	7.38
				A vs P					0.09342				**0.00006**			
50	2	9	11	Mean	141.00	79.00	15.50	46.50	5.33	3.33	0.36	1.67	356.22	211.11	24.82	120.11
				SD	22.63	18.38	0.71	4.95	2.45	1.87	0.50	1.12	56.18	36.46	2.48	24.14
				SE	6.82	5.54	0.21	1.49	0.74	0.56	0.15	0.34	16.94	10.99	0.75	7.28
				A vs P					**0.03216**				**0.00029**			

1049

1050 **Supplementary Table 2. Mitotic activity in the germ rudiment at the time of intercalation line (IL) formation.** "Live" samples
1051 correspond to lightsheet sample embryos (Supplementary Table 1). Data from samples WT1, WT2 and WT3 were used to generate rose
1052 plots in Supplementary Figure 2a.

Experiment	Sample #	L7	L6	L5	L5	L3	L2	L1	ML	R1	R2	R3	R4	R5	R6	R7	Total # En+	Sample Mean
								Row cell identity										
Wild Type	1	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	9	0.60
	2	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	11	0.73
	3	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	11	0.73
	4	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	9	0.60
	5	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	11	0.73
	6	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	10	0.67
	7	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	9	0.60
	8	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	0.00	7	0.47
	9	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	9	0.60
	10	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	13	0.87
	11	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	9	0.60
	12	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	0.00	8	0.53
	Cell Mean	0.08	0.08	0.33	0.92	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.83	0.42	0.00	0.00	9.67	
Lateral Ablation	1	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	9	0.60
	2	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	8	0.53
	3	0.00	0.00	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	0.00	5	0.33
	Cell Mean	0.00	0.00	0.00	0.33	0.67	0.67	1.00	1.00	1.00	1.00	1.00	0.67	0.00	0.00	0.00	7.33	
Midline Ablation	1	0.00	0.00	0.00	1.00	1.00	0.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	0.00	0.00	6	0.40
	2	0.00	0.00	0.00	0.00	0.00	0.00	0.00	1.00	1.00	1.00	1.00	0.00	0.00	0.00	0.00	4	0.27
	3	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0	0.00
	4	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0	0.00
	Cell Mean	0.00	0.00	0.00	0.25	0.25	0.00	0.25	0.50	0.50	0.50	0.25	0.00	0.00	0.00	0.00	2.50	

054 **Supplementary Table 3.** En expression in cells of the prospective mandibular segment in wild type (WT) embryos, following ablation of cell bonds
055 lateral to the midline (Lat), or following ablation of cell bonds at the midline, as depicted in Fig. 6b, c. These data were used to generate Fig. 6e.

References

1. His, W. *Untersuchungen über die erste Anlage des Wirbelthierleibes: die erste Entwickelung des Hühnchens im Ei*. (F.C.W. Vogel, Leipzig, 1868).

2. Keller, E. F. *The Century of the Gene*. (Harvard University Press, Cambridge, Massachusetts, 2002).

3. Turing, A. M. The Chemical Basis of Morphogenesis. *Philos. Trans. R. Soc. Lond. Ser. B, Biol. Sci.* **237**, 37–72 (1952).

4. Neumann, C. & Cohen, S. Problems and paradigms: Morphogens and pattern formation. *BioEssays* **19**, 721–729 (1997).

5. Müller, G. B. & Newman, S. *Origination of Organismal Form: Beyond the Gene in Developmental and Evolutionary Biology*. vol. 2 (MIT Press, Cambridge, Mass, 2003).

6. Roeder, A. H. K. Arabidopsis sepals: A model system for the emergent process of morphogenesis. *Quant. Plant Biol.* **2**, E14 (2021).

7. Gross, D., Sevrin, A. & Shraiman, B. The Physics of Living Matter: Space, Time and Information. in *Proceedings of the 27th Solvay Conference on Physics* (eds. Gross, D., Sevrin, A. & Shraiman, B.) (2020). doi:10.1142/10964.

8. Ghosh, S., Kuyyamudi, C., Steinert, B. L. & Extavour, C. G. Mechanical mechanisms of morphogenesis as potential substrates for evolutionary change. *Semin. Cell Dev. Biol.* **175**, 103645 (2025).

9. Caldarelli, P. *et al.* Self-organized tissue mechanics underlie embryonic regulation. *Nature* **633**, 887–894 (2024).

10. Stork, N. E. How Many Species of Insects and Other Terrestrial Arthropods Are There on Earth? *Annu. Rev. Entomol.* **63**, 1–15 (2017).

11. Brusca, R. C., Moore, W. & Shuster, S. M. Chapter 23: Phylum Arthropoda. in *The Invertebrates. Third Edition* (Sinauer Associates, Sunderland, MA, USA, 2016).

12. Clark, E., Peel, A. D. & Akam, M. Arthropod segmentation. *Development* **146**, dev170480 (2019).

13. Buckley, C. E. & Johnston, D. S. Apical–basal polarity and the control of epithelial form and function. *Nat. Rev. Mol. Cell Biol.* **23**, 559–577 (2022).

14. Classen, A.-K., Anderson, K. I., Marois, E. & Eaton, S. Hexagonal Packing of *Drosophila* Wing Epithelial Cells by the Planar Cell Polarity Pathway. *Dev. Cell* **9**, 805–817 (2005).

15. Gibson, M. C., Patel, A. B., Nagpal, R. & Perrimon, N. The emergence of geometric order in
proliferating metazoan epithelia. *Nature* **442**, 1038–1041 (2006).

16. Farhadifar, R., Röper, J.-C., Aigouy, B., Eaton, S. & Jülicher, F. The influence of cell
mechanics, cell-cell interactions, and proliferation on epithelial packing. *Curr. Biol.* **17**, 2095–
2104 (2007).

17. Bardet, P.-L. *et al.* PTEN controls junction lengthening and stability during cell
rearrangement in epithelial tissue. *Dev. Cell* **25**, 534–546 (2013).

18. Nagpal, R., Patel, A. & Gibson, M. C. Epithelial topology. *BioEssays* **30**, 260–266 (2008).

19. Monier, B., Pélissier-Monier, A., Brand, A. H. & Sanson, B. An actomyosin-based barrier
inhibits cell mixing at compartmental boundaries in Drosophila embryos. *Nature Cell Biol.* **12**,
60–69 (2010).

20. Simone, R. P. & DiNardo, S. Actomyosin contractility and Discs large contribute to
junctional conversion in guiding cell alignment within the Drosophila embryonic epithelium.
Development **137**, 1385–1394 (2010).

21. Wright, D. A. & Lawrence, P. A. Regeneration of segment boundaries in Oncopeltus: Cell
lineage. *Dev. Biol.* **85**, 328–333 (1981).

22. Zallen, J. A. & Zallen, R. Cell-pattern disordering during convergent extension in
Drosophila. *J. Phys.: Condens. Matter* **16**, S5073–S5080 (2004).

23. Honda, H. Geometrical Models for Cells in Tissues. *Int. Rev. Cyt.* **81**, 191–248 (1983).

24. Tamada, M. & Zallen, J. A. Square Cell Packing in the Drosophila Embryo through
Spatiotemporally Regulated EGF Receptor Signaling. *Dev. Cell* **35**, 151–161 (2015).

25. Dohle, W., Gerberding, M., Hejnol, A. & Scholtz, G. Cell lineage, segment differentiation,
and gene expression in crustaceans. in *Evolutionary Developmental Biology of Crustacea* (ed.
Scholtz, G.) 95–133 (A.A. Balkema, Lisse, 2004).

26. Gerberding, M. & Scholtz, G. Cell lineage of the midline cells in the amphipod crustacean
Orchestia cavimana (Crustacea, Malacostraca) during formation and separation of the germ
band. *Dev. Genes Evol.* **209**, 91–102 (1999).

27. Gerberding, M., Browne, W. E. & Patel, N. H. Cell lineage analysis of the amphipod
crustacean Parhyale hawaiensis reveals an early restriction of cell fates. *Development* **129**, 5789–
5801 (2002).

28. Scholtz, G. & Dohle, W. Cell lineage and cell fate in crustacean embryos--a comparative
approach. *Int. J. Dev. Biol.* **40**, 211–20 (1996).

29. Browne, W. E., Price, A. L., Gerberding, M. & Patel, N. H. Stages of embryonic
development in the amphipod crustacean Parhyale hawaiensis. *genesis* **42**, 124–149 (2005).

30. Spencer, M. A., Jabeen, Z. & Lubensky, D. K. Vertex stability and topological transitions in
vertex models of foams and epithelia. *Eur. Phys. J. E* **40**, 1–17 (2017).

31. Sun, D. A. & Patel, N. H. The amphipod crustacean Parhyale hawaiensis: An emerging
comparative model of arthropod development, evolution, and regeneration. *Wiley Interdiscip.*
Rev. Dev. Biol. **8**, e355 (2019).

32. Lawrence, P. A. *The Making of a Fly: The Genetics of Animal Design*. (Wiley-Blackwell,
New Jersey, USA, 1992).

33. Alwes, F., Hinchen, B. & Extavour, C. G. Patterns of cell lineage, movement, and migration
from germ layer specification to gastrulation in the amphipod crustacean Parhyale hawaiensis.
Dev. Biol. **359**, 110–123 (2011).

34. Cislo, D. J. *et al.* Active cell divisions generate fourfold orientationally ordered phase in
living tissue. *Nat. Phys.* **19**, 1201–1210 (2023).

35. Extavour, C. G. The fate of isolated blastomeres with respect to germ cell formation in the
amphipod crustacean Parhyale hawaiensis. *Dev. Biol.* **277**, 387–402 (2005).

36. Chaw, R. C. & Patel, N. H. Independent migration of cell populations in the early
gastrulation of the amphipod crustacean Parhyale hawaiensis. *Dev. Biol.* **371**, 94–109 (2012).

37. Patel, N. H. *et al.* Expression of *engrailed* proteins in arthropods, annelids, and chordates.
Cell **58**, 955–968 (1989).

38. Heemskerk, I. & Streichan, S. J. Tissue cartography: compressing bio-image data by
dimensional reduction. *Nat. Methods* **12**, 1139–1142 (2015).

39. Keller, R. E. & Trinkaus, J. P. Rearrangement of enveloping layer cells without disruption of
the epithelial permeability barrier as a factor in Fundulus epiboly. *Dev. Biol.* **120**, 12–24 (1987).

40. Price, A. L., Modrell, M. S., Hannibal, R. L. & Patel, N. H. Mesoderm and ectoderm lineages
in the crustacean Parhyale hawaiensis display intra-germ layer compensation. *Dev. Biol.* **341**,
256–266 (2010).

41. Graner, F. & Glazier, J. A. Simulation of biological cell sorting using a two-dimensional
extended Potts model. *Phys. Rev. Lett.* **69**, 2013–2016 (1992).

42. Holtfreter, J. Tissue Affinity, A Means of Embryonic Morphogenesis. in *Foundations of*
Experimental Embryology (eds. Willier, B. H. & Oppenheimer, J. M.) 186–225 (Prentice-Hall,
Englewood Cliffs, N.J., 1939).

43. Swat, M. H. *et al.* Multi-Scale Modeling of Tissues Using CompuCell3D. *Methods Cell Biol.*
110, 325–366 (2012).

44. Scholtz, G., Patel, N. H. & Dohle, W. Serially homologous engrailed stripes are generated
via different cell lineages in the germ band of amphipod crustaceans (Malacostraca, Peracarida).
Int. J. Dev. Biol. **38**, 471–8 (1994).

45. Scholtz, G. The formation, differentiation and segmentation of the post-naupliar germ band
of the amphipod Gammarus pulex L. (Crustacea, Malacostraca, Peracarida). *Proc. R. Soc. Lond.*
B **239**, 163–211 (1990).

46. Wolff, C. & Scholtz, G. Cell Lineage, Axis Formation, and the Origin of Germ Layers in the
Amphipod Crustacean Orchestia cavimana. *Dev. Biol.* **250**, 44–58 (2002).

47. Gerberding, M. & Scholtz, G. Neurons and Glia in the Midline of the Higher Crustacean
Orchestia cavimana Are Generated via an Invariant Cell Lineage That Comprises a Median
Neuroblast and Glial Progenitors. *Dev. Biol.* **235**, 397–409 (2001).

48. Fischer, A. H. L., Pabst, T. & Scholtz, G. Germ band differentiation in the stomatopod
Gonodactylaceus falcatus and the origin of the stereotyped cell division pattern in Malacostraca
(Crustacea). *Arthropod Struct. Dev.* **39**, 411–422 (2010).

49. Scholtz, G. & Wolff, C. Cleavage, gastrulation, and germ disc formation of the amphipod
Orchestia cavimana (Crustacea, Malacostraca, Peracarida). *Contrib. Zool.* **71**, 9–28 (2002).

50. Keller, R. *et al.* Mechanisms of convergence and extension by cell intercalation. *Philos.*
Trans. R. Soc. Lond. **355**, 897–922 (2000).

51. Karess, R. E. *et al.* The regulatory light chain of nonmuscle myosin is encoded by spaghetti-
squash, a gene required for cytokinesis in Drosophila. *Cell* **65**, 1177–1189 (1991).

52. Landsberg, K. P. *et al.* Increased Cell Bond Tension Governs Cell Sorting at the Drosophila
Anteroposterior Compartment Boundary. *Curr. Biol.* **19**, 1950–1955 (2009).

53. Scarpa, E., Finet, C., Blanchard, G. B. & Sanson, B. Actomyosin-Driven Tension at
Compartmental Boundaries Orients Cell Division Independently of Cell Geometry In Vivo. *Dev*
Cell **47**, 727-740.e6 (2018).

54. Jain, A. *et al.* Regionalized tissue fluidization is required for epithelial gap closure during
insect gastrulation. *Nat. Commun.* **11**, 5604 (2020).

55. Martin, A. C. The Physical Mechanisms of Drosophila Gastrulation: Mesoderm and
Endoderm Invagination. *Genetics* **214**, 543–560 (2020).

56. Parzen, E. On Estimation of a Probability Density Function and Mode. *Ann. Math. Stat.* **33**,
1065–1076 (1962).

57. Rosenblatt, M. Remarks on Some Nonparametric Estimates of a Density Function. *Ann.*
Math. Stat. **27**, 832–837 (1956).

58. Townes, P. L. & Holtfreter, J. Directed movements and selective adhesion of embryonic
amphibian cells. *J. Exp. Zool.* **128**, 53–120 (1955).

59. García-Bellido, A. Pattern reconstruction by dissociated imaginal disk cells of *Drosophila*
melanogaster. *Dev. Biol.* **14**, 278–306 (1966).

60. McMillen, P. & Holley, S. A. Integration of cell–cell and cell–ECM adhesion in vertebrate
morphogenesis. *Curr. Opin. Cell Biol.* **36**, 48–53 (2015).

61. Skamrahl, M. *et al.* Cellular segregation in cocultures is driven by differential adhesion and
contractility on distinct timescales. *Proc. Natl. Acad. Sci.* **120**, e2213186120 (2023).

62. Maitre, J.-L. *et al.* Adhesion Functions in Cell Sorting by Mechanically Coupling the
Cortices of Adhering Cells. *Science* **338**, 253–256 (2012).

63. Bao, M. *et al.* Stem cell-derived synthetic embryos self-assemble by exploiting cadherin
codes and cortical tension. *Nat. Cell Biol.* **24**, 1341–1349 (2022).

64. Steinberg, M. S. Adhesion-guided multicellular assembly: a commentary upon the postulates,
real and imagined, of the differential adhesion hypothesis, with special attention to computer
simulations of cell sorting. *J. Theor. Biol.* **55**, 431–443 (1975).

65. Tepass, U., Godt, D. & Winklbauer, R. Cell sorting in animal development: signalling and
adhesive mechanisms in the formation of tissue boundaries. *Curr. Opin. Genet. Dev.* **12**, 572–
582 (2002).

66. Iijima, N., Sato, K., Kuranaga, E. & Umetsu, D. Differential cell adhesion implemented by
Drosophila Toll corrects local distortions of the anterior-posterior compartment boundary. *Nat.*
Commun. **11**, 6320 (2020).

67. Hogeweg, P. Evolving Mechanisms of Morphogenesis: on the Interplay between Differential
Adhesion and Cell Differentiation. *J. Theor. Biol.* **203**, 317–333 (2000).

68. Blankenship, J. T., Backovic, S. T., Sanny, J. S. P., Weitz, O. & Zallen, J. A. Multicellular
rosette formation links planar cell polarity to tissue morphogenesis. *Dev. Cell* **11**, 459–470
(2006).

69. Jacinto, A. *et al.* Dynamic analysis of actin cable function during Drosophila dorsal closure.
Curr. Biol. **12**, 1245–1250 (2002).

70. Martin, P. & Lewis, J. Actin cables and epidermal movement in embryonic wound healing.
Nature **360**, 179–183 (1992).

71. Maroudas-Sacks, Y. *et al.* Topological defects in the nematic order of actin fibres as
organization centres of Hydra morphogenesis. *Nat. Phys.* **17**, 251–259 (2021).

72. Calzolari, S., Terriente, J. & Pujades, C. Cell segregation in the vertebrate hindbrain relies on
actomyosin cables located at the interhombomeric boundaries. *EMBO J.* **33**, 686–701 (2014).

73. Saadaoui, M., Rocancourt, D., Roussel, J., Corson, F. & Gros, J. A tensile ring drives tissue
flows to shape the gastrulating amniote embryo. *Science* **367**, 453–458 (2020).

74. Lee, J.-Y. *et al.* Wnt/Frizzled Signaling Controls C. elegans Gastrulation by Activating
Actomyosin Contractility. *Curr. Biol.* **16**, 1986–1997 (2006).

75. Serra, M. *et al.* A mechanochemical model recapitulates distinct vertebrate gastrulation
modes. *Sci. Adv.* **9**, eadh8152 (2023).

76. Lienkamp, S. S. *et al.* Vertebrate kidney tubules elongate using a planar cell polarity-
dependent, rosette-based mechanism of convergent extension. *Nat. Genet.* **44**, 1382–1387
(2012).

77. Ybot-Gonzalez, P. *et al.* Convergent extension, planar-cell-polarity signalling and initiation
of mouse neural tube closure. *Development* **134**, 789–799 (2007).

78. Darken, R. S. *et al.* The planar polarity gene strabismus regulates convergent extension
movements in Xenopus. *EMBO J.* **21**, 976–985 (2002).

79. Williams, M. L. & Solnica-Krezel, L. Nodal and planar cell polarity signaling cooperate to
regulate zebrafish convergence and extension gastrulation movements. *eLife* **9**, e54445 (2020).

80. Goto, T. & Keller, R. The Planar Cell Polarity Gene Strabismus Regulates Convergence and
Extension and Neural Fold Closure in Xenopus. *Dev. Biol.* **247**, 165–181 (2002).

81. Gallagher, K. D., Mani, M. & Carthew, R. W. Emergence of a geometric pattern of cell fates
from tissue-scale mechanics in the Drosophila eye. *eLife* **11**, e72806 (2022).

82. Dicko, M. *et al.* Geometry can provide long-range mechanical guidance for embryogenesis.
PLoS Comput. Biol. **13**, e1005443 (2017).

83. Guignard, L. *et al.* Contact area–dependent cell communication and the morphological
invariance of ascidian embryogenesis. *Science* **369**, eaar5663 (2020).

84. Royer, C. *et al.* Establishment of a relationship between blastomere geometry and YAP
localisation during compaction. *Development* **147**, dev189449 (2020).

85. Kumar, A. & Shivashankar, G. V. Mechanical Force Alters Morphogenetic Movements and
Segmental Gene Expression Patterns during Drosophila Embryogenesis. *PLoS ONE* **7**, e33089
(2012).

86. Ho, M. W., Matheson, A., Saunders, P. T., Goodwin, B. C. & Smallcombe, A. Ether-induced
segmentation disturbances in Drosophila melanogaster. *Roux's Arch. Dev. Biol.* **196**, 511–521
(1987).

87. Dohle, W. & Scholtz, G. Clonal analysis of the crustacean segment: the discordance between
genealogical and segmental borders. *Development* **104**, 147–160 (1988).

88. Scholtz, G. From a Single Cell to Segmental Structures: Crustacean Embryology. in (Oxford
University Press, 2020).

89. Scholtz, G. Evolution of the nauplius stage in malacostracan crustaceans. *J. Zool. Syst. Evol.*
Res. **38**, 175–187 (2000).

90. Gupta, T. & Extavour, C. G. Identification of a putative germ plasm in the amphipod
Parhyale hawaiensis. *EvoDevo* **4**, 34 (2013).

91. Nestorov, P., Battke, F., Levesque, M. P. & Gerberding, M. The Maternal Transcriptome of
the Crustacean Parhyale hawaiensis Is Inherited Asymmetrically to Invariant Cell Lineages of
the Ectoderm and Mesoderm. *PLoS ONE* **8**, e56049 (2013).

92. Scholtz, G. Chapter 24: Cleavage, germ band formation and head segmentation: the ground
pattern of the Euarthropoda. in *Arthropod Relationships* (eds. Fortey, R. A. & Thomas, R. H.)
317–332 (Chapman & Hall, London, UK, 1997).

93. Scholtz, G. & Wolff, C. Chapter 4: Arthropod Embryology: Cleavage and Germ Band
Development. in *Arthropod Biology and Evolution* (ed. Minelli, A.) 63–89 (2013).
doi:10.1007/978-3-642-36160-9_4.

94. Wolff, C. *et al.* Multi-view light-sheet imaging and tracking with the MaMuT software
reveals the cell lineage of a direct developing arthropod limb. *eLife* **7**, e34410 (2018).

95. Kontarakis, Z. & Pavlopoulos, A. Chapter 10. Transgenesis in Non-model Organisms: The
Case of Parhyale. in *Hox Genes: Methods and Protocols* (eds. Graba, Y. & Rezsohazy, R.) vol.
1196 145–181 (Humana Press, New York, NY, 2014).

96. Riedl, J. *et al.* Lifeact: a versatile marker to visualize F-actin. *Nat. Methods* **5**, 605–607
(2008).

97. Shaner, N. C. *et al.* A bright monomeric green fluorescent protein derived from
Branchiostoma lanceolatum. *Nat. Methods* **10**, 407–409 (2013).

98. Pavlopoulos, A., Berghammer, A. J., Averof, M. & Klingler, M. Efficient Transformation of
the Beetle Tribolium castaneum Using the Minos Transposable Element Quantitative and
Qualitative Analysis of Genomic Integration Events. *Genetics* **167**, 737–746 (2004).

99. Özpolat, B. D., Handberg-Thorsager, M., Vervoort, M. & Balavoine, G. Cell lineage and cell
cycling analyses of the 4d micromere using live imaging in the marine annelid Platynereis
dumerilii. *eLife* **6**, e30463 (2017).

100. Vopalensky, P., Tosches, M. A., Achim, K., Handberg-Thorsager, M. & Arendt, D. From
spiral cleavage to bilateral symmetry: the developmental cell lineage of the annelid brain. *BMC*
Biol. **17**, 81 (2019).

101. Rehm, E. J., Hannibal, R. L., Chaw, R. C., Vargas-Vila, M. & Patel, N. H. Fixation and
dissection of Parhyale hawaiensis embryos. *Cold Spring Harb. Protoc.* **2009**, pdb.prot5127
(2009).

102. Kiehart, D. P. & Feghali, R. Cytoplasmic myosin from Drosophila melanogaster. *J. Cell*
Biol. **103**, 1517–1525 (1986).

103. Tomer, R., Khairy, K., Amat, F. & Keller, P. J. Quantitative high-speed imaging of entire
developing embryos with simultaneous multiview light-sheet microscopy. *Nat. Methods* **9**, 755–
763 (2012).

104. Pietzsch, T., Saalfeld, S., Preibisch, S. & Tomancak, P. BigDataViewer: visualization and
processing for large image data sets. *Nat. Methods* **12**, 481–483 (2015).

105. Preibisch, S., Saalfeld, S., Schindelin, J. & Tomancak, P. Software for bead-based
registration of selective plane illumination microscopy data. *Nat. Methods* **7**, 418–419 (2010).

106. Preibisch, S. *et al.* Efficient Bayesian-based multiview deconvolution. *Nat. Methods* **11**,
645–648 (2014).

107. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat.*
Methods **9**, 676–82 (2012).

108. Berg, S. *et al.* ilastik: interactive machine learning for (bio)image analysis. *Nature Methods*
16, 1226–1232 (2019).

109. Heemskerk, I. & Streichan, S. J. Tissue cartography: compressing bio-image data by
dimensional reductio n. *Nature Methods* **12**, 1139–1142 (2015).

110. Etournay, R. *et al.* TissueMiner: A multiscale analysis toolkit to quantify how cellular
processes create tissue dynamics. *eLife* **5**, e14334 (2016).

111. Stringer, C. & Pachitariu, M. Cellpose3: one-click image restoration for improved cellular
segmentation. *bioRxiv* 2024.02.10.579780 (2024) doi:10.1101/2024.02.10.579780.

112. Glazier, J. A. & Graner, F. Simulation of the differential adhesion driven rearrangement of
biological cells. *Phys. Rev. E* **47**, 2128–2154 (1993).

113. Aigouy, B., Umetsu, D. & Eaton, S. Segmentation and Quantitative Analysis of Epithelial
Tissues. in *Methods in Molecular Biology: Drosophila Methods and Protocols* (ed. Dahlmann,
C.) vol. 1478 227–239 (Springer New York, New York, NY, 2016).

1316