Task Complexity and Limb Dominance Modulate the Effects of Ageing on Neuromuscular Function

Yuxiao Guo[1], Eleanor J. Jones[2], Abdulmajeed Altheyab[2,3], Nishadi N. Gamage[2,4], Bethan E. Phillips[2], Philip J. Atherton[2], Mathew Piasecki[2]

1. Institute of Sports Medicine and Health, Chengdu Sport University, Chengdu, China

2. Centre of Metabolism, Ageing & Physiology (COMAP), MRC-Versus Arthritis Centre for Musculoskeletal Ageing Research and National Institute for Health Research, (NIHR) Nottingham Biomedical Research Centre, School of Medicine, University of Nottingham, Derby, UK

3. Department of Occupational Therapy, College of Applied Medical Sciences, King Saud Bin Abdulaziz University for Health Sciences, Riyadh, Saudi Arabia

4. Neurophysiology of Human Movement Laboratory, Discipline of Physiology, School of Biomedicine, University of Adelaide, Adelaide, Australia

Abstract

Aim: This study aimed to investigate age-related and limb-specific differences in neuromuscular function of vastus lateralis. While muscle strength is critical for functional independence in ageing, asymmetries between the dominant and non-dominant limbs, which can arise from central or peripheral mechanisms, are not well understood. We specifically examined whether bilateral differences exist in neuromuscular function and motor unit (MU) firing behaviours in young and older people, and whether these differences vary with task complexity.

Methods: Twenty-one healthy young adults (22 ± 4 years; 15M, 6F) and seventeen healthy older adults (74 ± 5 years; 12M, 5F) were recruited. High-density surface electromyography signals were collected bilaterally from the vastus lateralis during constant and variable load contractions normalised to maximal voluntary isometric contraction (MVC) and decomposed into individual MU spike trains. Muscle strength and force control as well as MU firing properties were compared bilaterally using

multilevel mixed-effects linear regression models. Statistical significance was accepted at p<0.05.

Results: Older adults showed reduced maximal voluntary force, increased force tracking error, and lower MU firing rates (MUFR), particularly during sinusoidal contractions. Force steadiness and MUFR variability revealed distinct Leg × AgeGroup interactions: older adults had greater variability in non-dominant legs, while younger adults showed the opposite. MU firing properties differed between contraction types, with age-related impairments most evident during dynamic tasks.

Conclusion: These findings highlight that neuromuscular ageing is not uniformly bilateral but involves asymmetric adaptations, especially under dynamic force demands. Despite symmetrical limb use, leg dominance effects become more pronounced with age, potentially reflecting compensatory neuromuscular strategies. Task complexity amplifies these asymmetries, underscoring the need to consider limb-specific neural control when addressing age-related motor decline.

Keywords: limb dominance, motor units, ageing, neuromuscular, vastus lateralis

Introduction

Inter-limb asymmetry refers to imbalances in performance between opposing limbs, arising from bilateral asymmetry of descending motor output between the right and left hemispheres (Kapreli *et al.*, 2006). Other factors may also contribute to the development of such imbalances, including biomechanical disparities, previous injuries (Kuenze *et al.*, 2015) or disease states (Larson *et al.*, 2013), and the effects of ageing (McGrath *et al.*, 2021). Consequently, such imbalances can give rise to a favoured limb when performing tasks that require strength or control (Newton *et al.*, 2006). Importantly, this has clear implications in ageing as grip strength asymmetry in older people is a strong predictor of future falls (McGrath *et al.*, 2021), and older females who experienced falls exhibited significantly greater bilateral strength differences than those who did not (Skelton *et al.*, 2002).

Wider evidence pertaining to limb strength asymmetry is conflicting and varies between muscle groups (Rahnama *et al.*, 2005; Lanshammar & Ribom, 2011; Lathrop-Lambach *et al.*, 2014), with one study demonstrating significant asymmetry

in the knee extensors and flexors, showing 13.5% and 13% greater strength, respectively, during isokinetic contractions in dominant side compared to non-dominant side, and a further showing 9% greater force in the dominant arm flexors of young (Lecce *et al.*, 2025). However, a comprehensive review of studies examining the comparative strength of the knee extensors as well as their influence on performance in jump tasks, revealed no significant disparities between the limbs (McGrath *et al.*, 2016). The presence of such muscle asymmetry can have implications for the interpretation of clinical evaluations of skeletal muscle and for research studies that employ unilateral intervention models, and explanatory mechanisms are not fully established.

The regulation and coordination of neural inputs play a crucial role in achieving graded increases in muscle force. This involves concurrent variations in the number of active motor units (MU) (known as MU recruitment) and modulation of the discharge frequency of action potentials within these units (known as rate coding) (Del Vecchio *et al.*, 2019), probably leading to the uniformly distributed muscle strength and motor control ability (Enoka & Farina, 2021) across limbs. The evidence regarding disparities in MU recruitment and firing related to laterality is also inconclusive. A study investing MU discharge properties in a hand muscle during 30% of maximal voluntary contraction (MVC) revealed dissimilar MU characteristics between limbs with the dominant hand having lower firing rates (FR), FR variability and recruitment threshold when compared with the non-dominant hand (Adam *et al.*, 1998). A more recent study of unimanual movements showed less common synaptic inputs to motoneurons serving intrinsic and extrinsic muscles of the dominant hand, interpreted as enabling a higher flexibility in MU recruitment (Maillet *et al.*, 2022). Another investigation demonstrated higher MUFR and a greater proportion of common synaptic inputs in dominant biceps (Lecce *et al.*, 2025). Yet in lower limbs, there were similar levels of MUFR and MUFR variability bilaterally across submaximal isometric contractions at 10, 20, 40 and 60% of MVC in tibialis anterior (Petrovic *et al.*, 2022).

The existing body of knowledge primarily includes young adults, leaving a lack of information regarding the influence of age on limb dominance, particularly for the quadriceps, which constitute the largest muscle group in the lower body and are the most susceptible to atrophy with ageing (Maden-Wilkinson *et al.*, 2013). Importantly,

many studies of ageing MUs have focused only on unilateral movements, and any implications arising from the presence of limb strength asymmetry, as well as central and peripheral single motor unit properties, as a result of increasing age are yet to be determined.

The purpose of this study was to investigate bilateral differences in the knee extensors with respect to the muscle strength, force steadiness and vastus lateralis MU characteristics in both young and older adults. We hypothesised that no bilateral differences would present in all outcome measures across age groups, but clear age-related differences would be detected in both physical and motor unit function.

Materials and Methods

Participants

A total of 38 healthy individuals were recruited from the local community through advertisement. The cohort included 21 younger adults (15 males and 6 females) and 17 older adults (12 males and 5 females). Leg dominance was determined by asking participants which leg they would use to kick a football, which has been reported to have a strong agreement with observed leg dominance during bilateral mobilizing and unilateral stabilizing tasks (van Melick *et al.*, 2017).

Prior to enrolment, participants completed a comprehensive clinical screening examination at the University of Nottingham, Royal Derby Hospital site and subsequently provided written informed consent. Participants were excluded if they had BMI< 18.5 or > 35 kg/m^2; were competitive in any athletic discipline at county level or above; had a musculoskeletal disorder; respiratory disease; neurological disorder; metabolic disease; active cardiovascular problems; active inflammatory bowel or renal disease; recent steroid treatment within 6 months or hormone replacement therapy.

Participants were instructed to abstain from any vigorous physical exercise for at least 48 hours prior to testing and to avoid caffeine and alcohol consumption for 24 hours before the testing session. All testing sessions were conducted in the morning, with participants arriving at the laboratory at 0900h in a fasted state.

Experimental protocol

Each participant completed a single experimental session that included the assessment of maximal voluntary isometric contraction (MVC) torque of the knee extensors, a familiarisation task, and the evaluation of muscle force control. Throughout these procedures, high-density surface electromyography (HDsEMG) signals were recorded from the vastus lateralis muscle (Figure 1). All measurements were performed bilaterally.

Figure 1. Experimental approach. (A) A participant is seated with the ankle secured to a load cell to enable isometric knee extension. High-density surface electromyography (HDsEMG) with a 64-channel grid was placed over the belly of the vastus lateralis. (B) Representative data from the trapezoidal contraction at 25% of maximal voluntary contraction (MVC). Force steadiness (CoV-Force) and motor unit (MU) firing characteristics are shown, including recruitment and de-recruitment thresholds (purple arrow), MU firing rate (red dashed line), and firing rate variability (green dashed lines). (C) Representative data from the sinusoidal force-tracking task (target force: 25 ± 4% MVC). Tracking performance was evaluated by calculating the area under the error curve and by cross-correlation between the exerted and target paths. MU potential trains were then extracted and divided into ascending and descending segments for further analysis.

Measurement of knee extensor strength

Participants were seated in a custom-built chair, which was adjusted to ensure approximately 90 degrees of flexion at the hips and knees. The lower leg was secured to a force dynamometer using a non-compliant strap (designed with a calibrated strain gauge, RS125 Components Ltd, Corby, UK) positioned above the medial malleolus. To minimize movement of the hips and upper trunk during contractions, the hips and pelvis were stabilised using a seat belt. Following a standardized warm-up consisting of submaximal contractions, participants were instructed to exert maximum effort with the aid of real-time visual feedback and verbal encouragement, aiming for forceful and rapid contractions. During the trial, participants were not allowed to hold onto the sides of the chair and were instructed to cross their arms over their chest. It was further repeated two to three times interspersed with 60 seconds rest intervals. If there was a difference of less than 5% between the two last attempts, the highest value was accepted as the maximal voluntary contraction (MVC).

Muscle force control assessment

Muscle force control was evaluated during steady submaximal isometric contractions at 25% of MVC. Participants completed four trials following a trapezoidal profile (5s ramp-up, 12s plateau, 5s ramp-down), and the third trial was selected for detailed analysis. Muscle force signal was recorded at 2000Hz via a CED Micro 1401-3 (Cambridge Electronic Design, Cambridge, UK), interfaced with a desktop computer, with data exported for complexity analysis in MATLAB (R2024b; The MathWorks, Natick, MA, USA). Force steadiness was quantified using the coefficient of variation (CoV), calculated as the standard deviation of the force during the steady plateau phase divided by its mean, multiplied by 100. The measure reflects the relative magnitude of force fluctuations. To assess the ability to modulate force output in response to dynamic demands, participants performed a sinusoidal force tracking task using OTBioLab software (OT Bioelettronica, Turin, Italy). Prior to the test, a familiarisation trial was completed to ensure task understanding and reduce learning effects. The contraction included a 10s linear ramp-up phase, followed by eight sinusoidal oscillations at a fixed amplitude of $25 \pm 4\%$ MVC over 30s, and concluded with a 10s ramp-down phase. The recorded sinusoidal contractions were exported and

analysed using MATLAB software (R2024b; The MathWorks, Natick, MA, USA).

Force signals were low-pass filtered using a fourth-order Butterworth filter with a

cutoff frequency of 20Hz prior to analysis. To quantify force tracking accuracy, the

absolute error between the exerted path and the target path was calculated over the

entire sinusoidal phase. The area under this error curve (AUC) was then calculated

and normalised to the total duration of the contraction, yielding a normalised AUC

(NormAUC, in N). Larger NormAUC values indicate greater tracking error, thus

reflecting lower tracking precision. The temporal coupling between target and exerted

force paths during sinusoidal contractions was quantified using cross-correlation

analysis, with the maximum Pearson correlation coefficient (r) used as an index of

synchrony.

High-density Surface Electromyography

High-density surface electromyography (HDsEMG) signals were recorded from the

vastus lateralis muscle during the 25% of MVC trapezoidal contraction and sinusoidal

force tracking task. A semi-disposable HDsEMG array (64 electrodes, 13×5, 8mm,

I.E.D., GR08MM1305, OT Bioelettronica, Inc., Turin, Italy) was positioned over the

muscle belly of the vastus lateralis with approximate orientation of the muscle

fascicles (proximal to distal). Prior to placement, the skin was prepared by shaving,

lightly abrading, and cleansing with 70% ethanol. The electrodes were affixed to the

skin using flexible tape, and disposable bi-adhesive foam layers (SpesMedica,

Bettipaglia, Italy) were used to secure the adhesive grids to the muscle surface. The

skin electrode contact was facilitated by filling the cavities of the adhesive layers with

conductive paste (AC Cream, SpesMedica). A strap ground electrode (WS2,

OTBioelettronica, Turin, Italy) moistened with water was placed around the ankle of

the tested leg. The signals were acquired in a monopolar configuration, amplified

(x256), and filtered within the range of 10-500 Hz. The signals were then digitally

converted at a sampling rate of 2000 Hz using a 16-bit wireless amplifier

(Sessantaquattro, OTBioelettronica, Turin, Italy) and stored for further offline

processing.

The recorded HDsEMG signals were converted into MATLAB format, and

monopolar signals were further band-pass filtered (20 to 500 Hz) before

decomposition into motor unit potential trains (MUPTs) using a convolutive kernel compensation (CKC) algorithm (Holobar & Zazula, 2007). All decomposed MUPTs were manually reviewed by a trained investigator to ensure accuracy of discharge patterns. Only MUs with a pulse-to-noise ratio equal to or greater than 28 dB were retained for further analysis.

During the experiment, MUs meeting the analysis criteria were successfully identified from 18 younger and 17 older participants. The detailed distribution of MUs across different contraction types is as follows: During the trapezoidal contractions, a total of 940 MUs were identified from younger and older adults, including 245 from the dominant and 180 from the non-dominant sides of younger adults, and 244 from the dominant and 271 from the non-dominant sides of older adults. During the sinusoidal contractions, a total of 582 MUs were identified from younger and older adults, including 169 from the dominant and 131 from the non-dominant sides of younger adults, and 148 from the dominant and 134 from the non-dominant sides of older adults. For sinusoidal contractions, only the oscillatory section of the signal was analysed, excluding the initial ramp-up and final ramp-down. Each full oscillation was further segmented into ascending and descending phases, resulting in 14 segments, a total of 7,983 MUs were identified from younger and older adults, including 2,321 from the dominant and 1,802 from the non-dominant sides of younger adults, and 2,007 from the dominant and 1,853 from the non-dominant sides of older adults.

Motor unit firing rate (MUFR) was calculated as the mean instantaneous discharge rate during the plateau phase of trapezoidal contractions and the oscillation cycle of sinusoidal contractions, with phase-specific analysis performed separately for ascending and descending segments. Motor unit firing rate variability (FRV) was quantified as the coefficient of variation (CoV) of the inter-spike intervals (ISI) during the plateau. MUFR at recruitment (FRrec) and derecruitment (FRderec) phases was defined as the mean instantaneous discharge rate across the first five discharges. Recruitment and derecruitment thresholds (RecThrsh and DerecThrsh) were defined as the force level corresponding to the first and final detected firing of each motor unit, respectively, and expressed as a percentage of the individual's maximal voluntary contraction (%MVC).

Statistical Analysis

All statistical analysis was conducted using RStudio. Multilevel mixed-effects linear regression models were used through the package lme4 (Version 1.1-27.1) (Bates *et al.*, 2015) with each individual being regarded as an independent cluster. To explore the bilateral differences in young and older individuals, Leg and AgeGroup were included as categorical variables, and their interactions (Leg $\times$ AgeGroup) were explored in the models. Where the interaction term was statistically significant, simple effects analyses were conducted using estimated marginal means (EMMs) (Lenth, 2016) to examine the effect of Leg within each AgeGroup (and vice versa). If the interaction was not significant, main effects of AgeGroup and Leg were interpreted directly based on the model estimates. To assess model fit and estimate the proportion of variance explained, the marginal R^2 (R^2_m) and conditional R^2 (R^2c) are included in each table of model outputs. The R^2_m represents the proportion of variance explained by the fixed effects, and R^2c represents the variance explained by both fixed effects and random effects (Nakagawa & Schielzeth, 2013). Cohen's d effect size was calculated from EMMs derived from the linear mixed-effects models, accounting for the nested data structure. Commonly used interpretations separate effects sizes intro three categories: small ($d = 0.2$), medium ($d = 0.5$) and large ($d = 0.8$). Statistical significance was accepted when $p < 0.05$.

Results

A total of 38 healthy participants were included in the study, consisting of 21 younger adults (15 males, 6 females; age range: 20-36 years) and 17 older males (12 males, 5 females; age range: 67-83 years) (Table 1).

Table 1. Participant characteristics

Measurements	Younger		Older	
	Dominant	Non-Dominant	Dominant	Non-Dominant
Age (years)	22 ± 4		74 ± 5	
Body mass index (kg/m²)	24.27 ± 3.62		25.57 ± 2.95	
Muscle force control				
MVC (N)	510.64 ± 159.32	495.08 ± 178.57	281.68 ± 94.47	276.81 ± 89.88
CoV-Force	2.50 ± 0.92	2.62 ± 1.23	2.56 ± 0.78	2.18 ± 0.87
Sin NormAUC	0.77 ± 0.18	0.85 ± 0.22	1.14 ± 0.41	1.13 ± 0.39
Sin CC	0.94 ± 0.03	0.93 ± 0.03	0.89 ± 0.06	0.88 ± 0.06
Motor unit properties - Trapezoidal Contraction				
MUFR (Hz)	8.67 ± 1.24	8.68 ± 1.48	7.57 ± 1.10	7.64 ± 1.10
MUFR Variability (%)	12.32 ± 3.88	15.70 ± 9.91	11.40 ± 3.01	12.77 ± 3.84
RecThrsh (%MVC)	12.04 ± 2.22	12.39 ± 2.57	11.16 ± 2.35	11.37 ± 1.60
DerecThrsh (%MVC)	13.50 ± 2.73	12.87 ± 3.02	12.57 ± 4.27	12.58 ± 3.71
Motor unit properties - Sinusoidal Contraction				
MUFR (Hz)	8.98 ± 1.37	8.93 ± 1.20	7.66 ± 1.25	7.76 ± 1.17
MUFR Variability (%)	21.57 ± 5.02	18.39 ± 4.23	16.40 ± 3.57	17.95 ± 4.10

271 *Data are reported as mean ± standard deviation. Abbreviations: MVC, maximal*
272 *voluntary isometric contraction; CoV, coefficient of variation; Sin NormAUC,*
273 *normalised area under sinusoidal curve; Sin CC, cross-correlation coefficients*
274 *between sinusoidal curves; MUFR, motor unit firing rate; RecThrsh, recruitment*
275 *threshold; DerecThrsh, derecruitment threshold.*

Muscle force control

For MVC, there was a significant main effect of AgeGroup (β = -1.29, Δ = -228.97 N, 95% CI [-1.81 to -0.78], p < 0.001), with older adults generating lower maximal force compared to younger adults. There was no significant Leg effect (β = -0.09, Δ = -15.56 N, 95% CI [-0.27 to 0.10], p = 0.344) nor a significant Leg × AgeGroup interaction (p = 0.663) (Table 2, Figure 2A).

In the analysis of force steadiness (CoV-Force) during the trapezoidal contractions, no significant Leg × AgeGroup interaction was observed (p=0.097). There was also no

significant main effect of AgeGroup (β = 0.06, Δ = +0.06, 95% CI [-0.59 to 0.71], p = 0.851), nor significant main effect of Leg (β = 0.13, Δ = +0.12, 95% CI [-0.29 to 0.54], p = 0.549) (Table 2, Figure 2B).

In the segmented sinusoidal force-tracking task, no significant Leg × AgeGroup interaction was observed for either normalised area under the curve (NormAUC) (p=0.409) or cross-correlation coefficients (CC) (p=0.592). There were also no significant main effects of Leg (NormAUC: β = 0.11, p = 0.155; CC: β = -0.03, p = 0.743) or Segment Type (NormAUC: β = -0.05, p = 0.374; CC: β = -0.01, p = 0.854) on tracking accuracy. However, significant main effects of AgeGroup were found for both outcome measures, with older adults showing higher normalised AUC values (β = 0.63, Δ = +0.33, 95% CI [0.09 to 0.58], p = 0.008) and lower cross-correlation coefficients (β = -0.35, Δ = -0.04, 95% CI [-0.07 to -0.01], p = 0.007) (Table 2, Figure C-F).

Table 2. Summary of mixed-effects linear model results for muscle force control measures.

Measures	Leg (NonDom v Dom)	AgeGroup (Old v Young)	Leg × AgeGroup	Model Fit (R²m/R²c)
MVC (N)	↔ -15.56 (β=-0.09, p=0.344)	↓ -228.97 (β=-1.29, p<0.001)	p=0.663	0.390 / 0.913
CoV-Force (%)	↔ 0.13 (β=0.12, p=0.549)	↔ -0.06 (β=-0.06, p=0.851)	p=0.097	0.030 / 0.595
Sin NormAUC	↔ 0.06 (β=0.11, p=0.155)	↑ 0.33 (β=0.63, p=0.008)	p=0.409	0.085 / 0.449
Sin CC	↔ -0.00 (β=-0.03, p=0.743)	↓ -0.04 (β=-0.35, p=0.007)	p=0.592	0.024 / 0.082

*Note. Data presented as fixed effect estimates (β) with corresponding p values. Arrows indicate the direction of the effect: ↑increase, ↓decrease, ↔ no difference. The final column shows model fit statistics. Abbreviations: MVC, maximal voluntary contraction; CoV-Force, coefficient of variation of force; Sin NormAUC, normalised area under the sinusoidal curve; Sin CC, correlation coefficient during sinusoidal tracking; R²$_m$, marginal R² representing variance explained by fixed effects only; R²c, conditional R² representing variance explained by the full model (fixed + random effects). *Statistically significant (p < 0.05).*

Figure 2. *Measures of muscle force control in younger and older adults. Panels show maximal voluntary contraction (MVC) (A), coefficient of variance of force (CoV-Force) (B) during trapezoidal contractions, normalised area under the sinusoidal force curve (C-D) and cross-correlation coefficient (E-F) between target and exerted force during ascending and descending segments of sinusoidal contractions. Each data point represents each leg of an individual participant, with dominant and non-dominant legs indicated by purple circles and blue triangles, respectively. Individual motor unit data across segments are plotted as jittered points in panels C-F. Mean values for each leg per participant are connected by lines to visualise within subject differences across legs. Black dots with error bars represent estimated marginal means (EMMs) ± standard errors from the statistical model.*

Motor Unit Properties During 25% MVC Trapezoidal Contractions

For MUFR, no significant Leg × AgeGroup interaction was observed (p = 0.779). A significant main effect of AgeGroup was found, with older adults exhibiting lower MUFR compared to younger adults (β = -0.64, Δ = -1.18 Hz, 95% CI [-1.03 to -0.26], p = 0.001). No significant main effect of Leg was detected (β = 0.01, Δ = 0.01 Hz, 95% CI [-0.12 to 0.14], p = 0.908) (Table 3, Figure 3A). For MUFR variability, there was no significant Leg × AgeGroup interaction (p = 0.102). A significant main effect of Leg was found (β = 0.28, Δ = 2.08%, 95% CI [0.13 to 0.43], p < 0.001), indicating greater variability in the non-dominant legs in both groups, but no significant main effect of AgeGroup (β = -0.01, Δ = -0.08%, 95% CI [-0.32 to 0.30], p = 0.948) (Table 3, Figure 3B). For recruitment and derecruitment thresholds, neither Leg nor AgeGroup showed significant main effects, and there was no significant interaction (all p > 0.20) (Table 3, Figure 3C-D).

Motor Unit Properties During Sinusoidal Contractions

No significant Leg × AgeGroup interaction in MUFR was observed (p = 0.057). There was a significant main effect of Leg, indicating higher MUFR in the non-dominant limb (β = 0.13, Δ = 0.26 Hz, 95% CI [0.08 to 0.19], p < 0.001), and a nonsignificant trend toward lower MUFR in older adults (β = -0.56, Δ = -1.10 Hz, 95% CI [-1.15 to 0.03], p = 0.061) (Table 3, Figure 3E). For MUFR variability, a significant Leg × AgeGroup interaction was observed (p < 0.001). Main effects were significant for both Leg (β = -0.20, Δ = -0.06%, 95% CI [-0.27 to -0.14], p < 0.001) and AgeGroup (β = -0.34, Δ = -0.10%, 95% CI [-0.56 to -0.12], p = 0.002). Post-hoc analysis revealed that within-group comparisons, younger adults had higher MUFR variability in their dominant leg (p < 0.001, d = 0.26), while older adults showed a lower variability in their dominant legs (p = 0.009, d = -0.09) (Table 3, Figure 3F).

Table 3. Summary of multilevel linear regression model results for motor unit firing properties.

Contraction Types	Measures	Leg (NonDom v Dom)	AgeGroup (Old v Young)	AgeGroup × Leg	Model Fit (R2m/R2c)
Trapezoidal Contraction	MUFR (Hz)	↔ 0.01 (β=0.01, p=0.908)	↓ -1.18 (β=-0.64, p=0.001)	p=0.779	0.360 / 0.628
	MUFR Variability (%)	↑ 2.08 (β=0.28, p<0.001)	↔ -0.08 (β=-0.01, p=0.948)	p=0.102	0.290 / 0.450
	RecThrsh (%MVC)	↔ -0.07 (β=-0.01, p=0.899)	↔ -0.86 (β=-0.16, p=0.237)	p=0.651	0.004 / 0.083
	DerecThrsh (%MVC)	↔ -0.58 (β=-0.10, p=0.215)	↔ -0.88 (β=-0.16, p=0.438)	p=0.267	0.004 / 0.313
Sinusoidal Contraction	MUFR (Hz)	↑ 0.26 (β=0.13, p<0.001)	↓ -1.10 (β=-0.56, p=0.061)	p=0.057	0.142 / 0.541
	MUFR Variability (%)	↓ -0.06 (β=-0.20, p<0.001)	↓ -0.10 (β=-0.34, p=0.002)	p<0.001	0.085 / 0.167

Note. Data are presented as fixed effect estimates (β) with corresponding p values. Arrows indicate the direction of the effect: ↑increase, ↓decrease, ↔ no meaningful difference. The final column shows model fit statistics. Abbreviations: MUFR, motor unit firing rate; MUFR variability, coefficient of variation of motor unit firing rate; RecThrsh, recruitment threshold; DerecThrsh, derecruitment threshold; R²m, marginal R² (variance explained by fixed effects only); R²c, conditional R² (variance explained by the full model including both fixed and random effects). Statistically significant (p < 0.05).

Figure 3. *Motor unit firing properties in younger and older adults during trapezoidal and sinusoidal contractions. Panels A and B show motor unit firing rate (MUFR) and MUFR variability during 25% maximal voluntary trapezoidal contractions. Panel C and D illustrate recruitment and derecruitment thresholds, respectively. Panel E and F present MUFR and MUFR variability during sinusoidal force-tracking tasks. Data are presented separately by AgeGroup (Younger vs older adults) and Leg dominance (Dominant vs non-dominant). Purple circles are representing dominant legs and blue triangles represent non-dominant legs. Individual motor unit data are plotted as jittered points. Mean values for each leg per participant are connected by lines to visualise within subject differences across legs. Black dots with error bars represent estimated marginal means (EMMs) ± standard errors from the statistical model.*

In addition to the main effects of Leg and AgeGroup, significant main effects of

segment type were observed in MUFR and MUFR variability during the sinusoidal

contractions. Specifically, MUFR was higher during ascending phases ($\beta = 0.27$, 95%

CI [0.42 to 0.12], $p < 0.001$), and MUFR variability was markedly greater during

descending segments ($\beta = 0.14$, 95% CI [0.11 to 0.17], $p < 0.001$). These consistent

effects underscore the importance of analysing ascending and descending phases

separately when evaluating age- and leg-related differences in MU behaviour.

When phases were explored separately, the ascending phase of sinusoidal contractions

showed no significant Leg $\times$ AgeGroup interaction for MUFR ($p = 0.096$). There was

a significant main effect of Leg ($\beta = 0.22$, $\Delta = 0.43$ Hz, 95% CI [0.15 to 0.29], $p <$

0.001), with higher MUFR in non-dominant legs, whereas there was no significant

main effect of AgeGroup ($\beta = -0.38$, $\Delta = -0.73$ Hz, 95% CI [-1.04 to 0.29], $p = 0.272$)

(Table 4, Figure 4A). For MUFR variability, a significant Leg $\times$ AgeGroup interaction

was found ($p < 0.001$), along with significant main effects of Leg ($\beta = -0.22$, $\Delta = -$

0.02%, 95% CI [-0.31 to -0.13], $p < 0.001$) and AgeGroup ($\beta = -0.62$, $\Delta = -0.07\%$, 95%

CI [-1.04 to -0.20], $p = 0.004$). Post-hoc analysis showed that MUFR variability

during ascending phases was higher in the dominant legs compared to the non-

dominant legs in younger adults ($\Delta = 0.02\%$, $p < 0.001$, $d = 0.24$), while older adults

showed a reversed but non-significant pattern ($\Delta = -0.01\%$, $p = 0.099$, $d = -0.08$).

Age-related differences revealed greater MUFR variability in younger adults than

older adults, with more pronounced differences in the dominant legs ($\Delta = 0.07\%$, $p =$

0.004, $d = 0.66$) than in the non-dominant legs ($\Delta = 0.03\%$, $p = 0.13$, $d = 0.35$) (Table

4, Figure 4C).

During the descending phase of sinusoidal contractions, no significant Leg $\times$

AgeGroup interaction was found in MUFR ($p = 0.311$) nor a main effect of Leg ($\beta =$

0.08, $\Delta = 0.08$ Hz, 95% CI [-0.03 to 0.12], $p = 0.266$). However, there was a

significant main effect of AgeGroup ($\beta = -0.72$, $\Delta = -1.38$ Hz, 95% CI [-1.29 to -0.15],

$p = 0.013$), with lower MUFR in older adults across both legs (Table 4, Figure 4B).

For MUFR variability, a significant Leg $\times$ AgeGroup interaction was found ($p <$

0.001), along with main effects of Leg ($\beta = -0.25$, $\Delta = -0.10\%$, 95% CI [-0.34 to -0.16],

$p < 0.001$) and AgeGroup ($\beta = -0.38$, $\Delta = -0.14\%$, 95% CI [-0.65 to -0.11], $p = 0.006$).

Post-hoc analyses showed that MUFR variability was greater in the dominant leg

compared to the non-dominant leg in younger adults ($\Delta = 0.10\%$, $p < 0.001$, $d = 0.26$),

whereas older adults exhibited the opposite pattern ($\Delta = -0.04\%$, p = 0.025, d = –0.10). Age-related differences were more pronounced in the dominant leg ($\Delta = 0.14\%$, p = 0.006, d = 0.40) than in the non-dominant leg ($\Delta = 0.01\%$, p = 0.84, d = 0.03) (Table 4, Figure 4D).

Table 4. Summary of multilevel linear regression model results for motor unit firing properties during ascending and descending phases of sinusoidal contractions

Contraction Phases	Measures	Leg (NonDom - Dom)	AgeGroup (Old - Young)	Leg × AgeGroup	Model Fit (R2m/R2c)
Sinusoidal	MUFR (Hz)	↑ 0.43 (β=0.22, p<0.001)	↔ -0.73 (β=-0.38, p=0.272)	p=0.096	0.168 / 0.616
Ascending	MUFR Variability (%)	↓ -0.02 (β=-0.22, p<0.001)	↓ -0.07 (β=-0.62, p=0.004)	p<0.001	0.088 / 0.306
Sinusoidal	MUFR (Hz)	↔ 0.08 (β=0.08, p=0.266)	↓ -1.38 (β=-0.72, p=0.013)	p=0.311	0.109 / 0.487
Descending	MUFR Variability (%)	↓ -0.10 (β=-0.25, p<0.001)	↓ -0.14 (β=-0.38, p=0.006)	p<0.001	0.026 / 0.108

Note. Data are presented as fixed effect estimates (β) with corresponding p values. Arrows indicate the direction of the effect: ↑increase, ↓decrease, ↔ no meaningful difference. The final column shows model fit statistics. Abbreviations: MUFR, motor unit firing rate; MUFR variability, coefficient of variation of motor unit firing rate; R^2□, marginal R^2 (variance explained by fixed effects only); R^2c, conditional R^2 (variance explained by the full model including both fixed and random effects). Statistically significant (p < 0.05).

Figure 4. Motor unit firing properties during ascending and descending phases of sinusoidal contractions in younger and older adults. Panels A and C show motor unit firing rate (MUFR) and MUFR variability during the ascending phase of a sinusoidal contraction. Panels B and D display the corresponding measures during the descending phase. Data are presented separately by AgeGroup (Younger vs older adults) and Leg dominance (Dominant vs non-dominant). Purple circles are representing dominant legs and blue triangles represent non-dominant legs. Individual motor unit data are plotted as jittered points. Mean values for each leg per participant are connected by lines to visualise within subject differences across legs. Black dots with error bars represent estimated marginal means (EMMs) ± standard errors from the statistical model.

Discussion

The present study investigated bilateral differences in muscle force control and MU firing behaviours in young and older adults across simple and complex force tracking tasks. Our key findings revealed clear age-related impairments in neuromuscular function, including reduced maximal voluntary force and reduced accuracy during sinusoidal force tracking tasks. No significant leg asymmetries were observed in either age group, suggesting age-related declines in neuromuscular function are

bilateral and relatively symmetric, rather than being selectively expressed in one limb.
Notably, distinct MU firing patterns were observed between contraction types and age
groups, with clear evidence of variable firing during load matched isometric
shortening and lengthening contractions. During trapezoidal contractions, older adults
exhibited significantly lower MUFR compared to younger adults, whereas MUFR
variability was primarily influenced by leg dominance, with both groups showing
greater variability in the non-dominant legs. During sinusoidal contractions, older
adults exhibited lower MUFR, particularly in the descending phases, while MUFR
variability revealed a significant Leg × AgeGroup interaction. Younger adults
consistently demonstrated higher MUFR variability in the dominant legs, whereas
older adults displayed an opposite yet less severe pattern, with greater MUFR
variability in the non-dominant legs. These findings indicate that leg-related
asymmetries in MU firing behaviour are exacerbated with ageing, particularly during
complex dynamic force modulation.

The existence of bilateral asymmetries in skeletal muscle function has yielded
inconsistent findings, with studies showing both significant limb differences
(Portegijs *et al.*, 2005; Newton *et al.*, 2006) and no clear disparities (Ditroilo *et al.*,
2010; Carpes *et al.*, 2010). In the present study, we observed no significant bilateral
differences in knee extensor strength in either age group. This aligns with previous
findings in other lower limb muscles, such as the tibialis anterior (Petrovic *et al.*, 2022)
and may be attributable to the bilateral nature of lower limb involvement during daily
locomotor activities. Although numerous studies have reported reduced force
steadiness in older adults compared to younger counterparts (Oomen & van Dieën,
2017; Pethick *et al.*, 2022), our findings did not show significant age-related
differences in this measure. This finding is consistent with our previous work (Guo *et*
al., 2024) and may be explained by the relatively high health and functional status of
the older adults included.

The observed bilateral symmetry in both groups suggests that isometric force control
relies on generalized mechanisms rather than limb-specific adaptations, or, that limb-
specific adaptations in lower limbs do not occur. However, this symmetry did not
extend to more complex tasks. In contrast to constant load contractions, older adults
exhibited significantly reduced performance in the sinusoidal force-tracking tasks.
This dissociation highlights the task-specific nature of age-related declines, as the

demands of variable force tracking strongly rely on mechanosensitive afferents,
known to decrease with age (McIntyre *et al.*, 2021). While these force-related findings
suggest age- and limb-related differences in control strategies, they provide limited
insight into the underlying neural mechanisms. To address this, we examined MU
firing behaviour across contraction types.

MUFR exhibited task- and age-specific modulation, with more pronounced effects
during variable force contractions. In constant force contractions, MUFR did not
differ between legs in either age group, consistent with prior findings in lower limbs
(e.g., tibialis anterior) (Petrovic *et al.*, 2022), whereas limb-specific differences have
been reported in the upper limbs (e.g., biceps brachii) (Lecce *et al.*, 2025). However,
older adults exhibited significantly lower MUFR, which is a widely reported finding
in both upper and lower limbs (Orssatto *et al.*, 2022). These results suggest that under
steady contractions, MUFR is primarily influenced by age, rather than leg dominance.
In contrast, sinusoidal (variable force) contractions revealed more complex
modulation patterns. A similar trend to that observed in constant contractions, lower
MUFR in older adults, was also present in the sinewave contraction, although it was
not significantly different (p = 0.061). This effect is likely masked by segment-
specific differences.

To better understand these task-dependent modulations, sinusoidal contractions were
further segmented into ascending (concentric) and descending (eccentric) phases.
MUFR was consistently higher during the ascending phase, reflecting the increased
neural drive required for force generation. While the task was specifically isometric,
there is undoubtedly fibre shortening and lengthening across this form of contraction.
These data align with prior findings demonstrating that MUFRs tend to be greater
during concentric than eccentric contractions, even within the same MU, especially
during low-force dynamic task (Del Valle & Thomas, 2005; Duchateau & Baudry,
2013; Basile *et al.*, 2025). The greater MUFR during the ascending phase may be
attributable to enhanced corticospinal excitability and greater neuromuscular
recruitment demands associated with concentric movement, potentially also reflecting
reduced inhibitory control compared to eccentric contractions (Gueugneau *et al.*,
2023).

Importantly, when examining sinusoidal contractions in detail, a distinct modulation

pattern related to leg dominance was observed, which was not present during constant force tasks. Specifically, MUFR during the ascending phase was significantly higher in the non-dominant legs in both age groups. This finding partially aligns with some prior work but contrasts with others. While several studies have reported higher MUFR on the dominant side across different contraction intensities (Lecce *et al.*, 2025), others have found no difference (Petrovic *et al.*, 2022) or even non-dominant advantages (Adam *et al.*, 1998). These discrepancies may arise from differences in muscle selection, contraction type and the inherently dynamic nature of sinusoidal contractions, which engaged greater neuromuscular demands than constant plateaus. The present findings support the presence of lateralised neuromuscular control strategies that influence force generation more than force reduction. The ascending phase, characterised by increasing force output, relies heavily on corticospinal drive (Clos *et al.*, 2022). While the dominant legs are associated with stronger neural drive (Lecce *et al.*, 2025) and more effective proprioceptive feedback (Strong *et al.*, 2023), the higher MUFR observed in the non-dominant leg may indicate a compensatory strategy, possibly involving changes in inhibition control or motoneuron excitability, though the exact mechanisms remain to be fully clarified. These results collectively suggest that leg dominance interacts with task dynamics to shape neuromuscular recruitment strategies, particularly during phases of increasing force.

When examining these phase-specific effects across age groups, we have observed a significant decrease in MUFR during the descending (eccentric) phase in older adults. Previous studies have highlighted that eccentric contractions rely more heavily on cortical involvement and require greater sensorimotor integration than concentric contractions (Duchateau & Baudry, 2013). However, ageing is associated with reduced corticospinal excitability (Oliviero *et al.*, 2006), particularly during eccentric tasks (Duclay *et al.*, 2014; Clos *et al.*, 2022; Ruas *et al.*, 2025), which likely contributes to impaired discharge modulation during muscle lengthening contractions. Additionally, the reduced MUFR during the eccentric descending phase could reflect not only impaired excitatory drive but also enhanced inhibitory processes within the motor system, Specifically, older adults may exhibit increased presynaptic inhibition at the spinal level (Baudry & Duchateau, 2012; Filho *et al.*, 2021) or reduced intracortical facilitation during eccentric movements (Adam *et al.*, 1998), both of which could further limit the ability to modulate MU firing pattern. This suggests that

the age-related decline in neural drive is more pronounced during phases of force
reduction than force generation. This finding has important considerations for
activities of daily living which rely on eccentric contraction, such as stair descent.

While MUFR serves as a fundamental parameter for investigating neuromuscular
recruitment strategies, the variability of MUFR offers additional critical insights into
the stability and adaptability of neuromuscular system, which is also influenced by
task complexity, leg dominance and age-related physiological changes. In the present
study, during constant load contractions, both age groups showed greater MUFR
variability in the non-dominant legs. This aligns with earlier findings suggesting
greater MUFR fluctuations on the non-dominant side (Adam *et al.*, 1998), potentially
reflecting a stronger reliance on afferent sensory feedback and spinal-level regulation
in that limb. However, others report no significant limb-related differences in MUFR
variability, albeit in the upper limb (Lecce *et al.*, 2025). During sinusoidal
contractions, which require continuous force modulation and more dynamic
engagement of higher-order motor control systems, MUFR variability was markedly
greater across all participants. This task-dependent increase in MUFR variability
highlights the increased demands placed on neuromuscular coordination and
sensorimotor integration throughout the contraction.

MUFR variability is generally higher with higher MUFR in young and older VL (Guo
et al., 2022, 2025), but this relationship dissociates when comparing ascending and
descending segments. Despite lower MUFR, the descending (eccentric) phase
exhibited significantly higher MUFR variability than the ascending (concentric) phase.
This dissociation suggests that MUFR variability is not solely MUFR-dependent but
rather reflects distinct phase-specific neural control strategies. Specifically, the
increased MUFR variability during eccentric contraction may reflect stronger
inhibitory regulation, such as presynaptic inhibition or cortical inhibition, which are
known to be upregulated during lengthening eccentric contractions (Doguet *et al.*,
2017; Papitsa *et al.*, 2022). However, in older adults, age-related impairments in
muscle relaxation (Hunter *et al.*, 2016; Jeon *et al.*, 2023) may influence afferent
feedback and further limit the ability to stabilise motor output during the eccentric
phase. These findings suggest that while dynamic force modulation may remain
relatively intact in older adults, specific deficits in descending-phase control reveal a
selective breakdown in neuromuscular adaptability during eccentric tasks.

Age-related differences in MUFR variability became more pronounced during sinusoidal contractions. In younger adults, MUFR variability was higher in the dominant leg. This greater MUFR variability may be indicative of a more variable use of corticospinal resources by the dominant legs, such as that proposed following the finding of fewer common synaptic inputs in dominant hands (Maillet *et al.*, 2022), thereby enabling task-specific adjustments to motor output. Such flexibility may be particularly advantageous in tasks involving variable forces, as it facilitates rapid transition in force. However, this was not apparent in the older adults, where MUFR variability was more similar across legs, suggesting a shift towards more generalised control strategies. This phenomenon may be attributable to age-related corticospinal degradation or diminished inter-hemispheric communication (Delvenne & Malloy, 2025). Notably, older adults demonstrated reduced bilateral modulation across tasks, with smaller inter-limb differences in MUFR variability during variable force contractions. Overall, these findings underscore the progressive loss of neuromuscular adaptability with aging, particularly the diminished capacity to implement differentiated neural strategies tailored to task-specific motor demands.

Strengths and Limitations

This study investigated the effects of leg dominant on neuromuscular function across different age groups, including both static trapezoidal contractions and dynamic sinusoidal contractions, and this study provides valuable function and mechanistic insight from independent motoneuron pool of vastus lateralis, allowing for a comprehensive understanding of how these factors may vary across legs and age. However, this study still has some limitations. (1) Both males and females were included in the study, but the sample size for female MU data was relatively small. (2) MUs sampled in the current study were only from mid-level contraction intensity, so revealed limited information on later recruited MUs for higher force levels. (3) Knee extensor movements are controlled by multiple muscles, so investigating only the MUs of vastus lateralis MUs may limit insight into whole muscle group.

Conclusions

Our findings collectively demonstrate that the aging neuromuscular system undergoes
strategic reorganisation rather than uniform decline, manifesting through complex
interactions between limb dominance and task complexity, as MU firing behaviours
exhibit distinct task- and leg-specific modulation. Despite the symmetrical use of
lower limbs, subtle lateralised differences are evident, particularly during dynamic
tasks requiring continuous modulation of neural drive. Notably, we also report distinct
differences MU behaviour in contraction type, with lower MUFR and higher
variability during eccentric phases. Rather than a uniform decline, ageing appears to
reshape motor control through altered lateralised strategies. These insights may
inform the development of more targeted interventions to support functional
independence in older adults.

Acknowledgements

The authors thank all of the participants for their time and participation.

Conflict of Interest

The authors have no conflict of interest to declare.

Author Contributions

All authors contributed to the conception and design of the study. YG, AA, NNG, EJ
and JP contributed to the data acquisition. YG analysed the data and drafted the
manuscript. BEP, PJA and MP provided comments. All authors have approved the
final version of the submitted manuscript for publication and are accountable for all
aspects of the work. All persons designated as authors qualify for authorship, and all
those who qualify for authorship are listed.

Ethics Approval

This research was approved by the University of Nottingham Faculty of Medicine and
Health Sciences Research Ethics Committee (FMHS-160-0121, 90-0820, 390-1121)

and was conducted between 2022 and 2024 in accordance with the Declaration of
Helsinki.

Funding information

632 This work was supported by the Medical Research Council [grant number
633 MR/P021220/1] as part of the MRC-Versus Arthritis Centre for Musculoskeletal
634 Ageing Research awarded to the Universities of Nottingham and Birmingham and
635 was supported by the NIHR Nottingham Biomedical Research Centre. This work was
636 also supported by a Physiological Society Research Grant awarded to MP.

Data availability statement

639 The datasets generated and analysed during the current study are available from the
640 corresponding author upon reasonable request.

References

Adam A, Luca CJ De & Erim Z (1998). Hand Dominance and Motor Unit Firing
Behavior. *J Neurophysiol* **80,** 1373–1382.

Basile DC, Wira AD, Rice CL & Power KE (2025). Investigating motor unit firing rates
during arm cycling compared with intensity-matched isometric contractions in
humans. *J Neurophysiol* **134,** 162–170.

Bates D, Mächler M, Bolker B & Walker S (2015). Fitting Linear Mixed-Effects Models
Using lme4. *J Stat Softw* **67,** 1–48.

Baudry S & Duchateau J (2012). Age☐related influence of vision and proprioception on
Ia presynaptic inhibition in soleus muscle during upright stance. *J Physiol* **590,**
5541–5554.

Carpes FP, Diefenthaeler F, Bini RR, Stefanyshyn D, Faria IE, Mota CB, Rahnama N,
Lees A & Bambaecichi E (2010). Comparison of muscle strength and flexibility
between the preferred and non-preferred leg in English soccer players.
Ergonomics **48,** 1230–1236.

Clos P, Mater A, Legrand H, Poirier G, Ballay Y, Martin A & Lepers R (2022).
Corticospinal excitability is lower during eccentric than concentric cycling in men.
Front Physiol **13,** 854824.

Delvenne J-F & Malloy E (2025). Functional implications of age-related atrophy of the

661   corpus callosum. *Neurosci Biobehav Rev* **169,** 105982.

662 Ditroilo M, Forte R, Benelli P, Gambarara D & De Vito G (2010). Effects of age and
663   limb dominance on upper and lower limb muscle function in healthy males and
664   females aged 40-80 years. *J Sports Sci* **28,** 667–677.

665 Doguet V, Nosaka K, Guével A, Thickbroom G, Ishimura K & Jubeau M (2017). Muscle
666   length effect on corticospinal excitability during maximal concentric, isometric and
667   eccentric contractions of the knee extensors. *Exp Physiol* **102,** 1513–1523.

668 Duchateau J & Baudry S (2013). Insights into the neural control of eccentric
669   contractions. *J Appl Physiol* **116,** 1418–1425.

670 Duclay J, Pasquet B, Martin A & Duchateau J (2014). Specific modulation of spinal and
671   cortical excitabilities during lengthening and shortening submaximal and maximal
672   contractions in plantar flexor muscles. *J Appl Physiol* **117,** 1440–1450.

673 Enoka RM & Farina D (2021). Force Steadiness: From Motor Units to Voluntary
674   Actions. *Physiology (Bethesda)* **36,** 114–130.

675 Filho SS, Coelho DB, Ugrinowitsch C, de Souza CR, Magalhães FH, de Lima-Pardini
676   AC, de Oliveira ÉMB, Mattos E, Teixeira LA & Silva-Batista C (2021). Age-
677   Related Changes in Presynaptic Inhibition During Gait Initiation. *The Journals of*
678   *Gerontology: Series A* **76,** 568–575.

679 Gueugneau N, Martin A, Gaveau J & Papaxanthis C (2023). Gravity-efficient motor
680   control is associated with contraction-dependent intracortical inhibition. *iScience*
681   **26,** 107150.

682 Guo Y, Jones EJ, Inns TB, Ely IA, Stashuk DW, Wilkinson DJ, Smith K, Piasecki J,
683   Phillips BE, Atherton PJ & Piasecki M (2022). Neuromuscular recruitment
684   strategies of the vastus lateralis according to sex. *Acta Physiologica* **235,** e13803.

685 Guo Y, Jones EJ, Škarabot J, Inns TB, Phillips BE, Atherton PJ & Piasecki M (2024).
686   Common synaptic inputs and persistent inward currents of vastus lateralis motor
687   units are reduced in older male adults. *Geroscience*; DOI: 10.1007/s11357-024-
688   01063-w.

689 Guo Y, Jones EJ, Smart TF, Altheyab A, Gamage N, Stashuk DW, Piasecki J, Phillips
690   BE, Atherton PJ & Piasecki M (2025). Sex disparities of human neuromuscular
691   decline in older humans. *J Physiol* **603,** 151–165.

692 Holobar A & Zazula D (2007). Multichannel Blind Source Separation Using
693   Convolution Kernel Compensation. *IEEE Transactions on Signal Processing* **55,**
694   4487–4496.

695 Hunter SK, Pereira HM & Keenan KG (2016). The aging neuromuscular system and
696   motor performance. *J Appl Physiol* **121,** 982–995.

697 Jeon W, Ramadan A, Whitall J, Alissa N & Westlake K (2023). Age-related differences
698   in lower limb muscle activation patterns and balance control strategies while
699   walking over a compliant surface. *Sci Rep* **13,** 16555.

700 Kapreli E, Athanasopoulos S, Papathanasiou M, Van Hecke P, Strimpakos N, Gouliamos

A, Peeters R & Sunaert S (2006). Lateralization of brain activity during lower limb joints movement. An fMRI study. *Neuroimage* **32,** 1709–1721.

Kuenze CM, Hertel J, Weltman A, Diduch D, Saliba SA & Hart JM (2015). Persistent neuromuscular and corticomotor quadriceps asymmetry after anterior cruciate ligament reconstruction. *J Athl Train* **50,** 303–312.

Lanshammar K & Ribom EL (2011). Differences in muscle strength in dominant and non-dominant leg in females aged 20-39 years--a population-based study. *Phys Ther Sport* **12,** 76–79.

Larson RD, McCully KK, Larson DJ, Pryor WM & White LJ (2013). Bilateral differences in lower-limb performance in individuals with multiple sclerosis. *J Rehabil Res Dev* **50,** 215–222.

Lathrop-Lambach RL, Asay JL, Jamison ST, Pan X, Schmitt LC, Blazek K, Siston RA, Andriacchi TP & Chaudhari AMW (2014). Evidence for joint moment asymmetry in healthy populations during gait. *Gait Posture* **40,** 526–531.

Lecce E, Del Vecchio A, Nuccio S, Felici F & Bazzucchi I (2025). Higher dominant muscle strength is mediated by motor unit discharge rates and proportion of common synaptic inputs. *Sci Rep* **15,** 8269.

Lenth R V (2016). Least-Squares Means: The R Package lsmeans. *J Stat Softw* **69,** 1–33.

Maden-Wilkinson TM, Degens H, Jones DA & McPhee JS (2013). Comparison of MRI and DXA to measure muscle size and age-related atrophy in thigh muscles. *J Musculoskelet Neuronal Interact* **13,** 320–328.

Maillet J, Avrillon S, Nordez A, Rossi J & Hug F (2022). Handedness is associated with less common input to spinal motor neurons innervating different hand muscles. *J Neurophysiol* **128,** 778–789.

McGrath R, Clark BC, Cesari M, Johnson C & Jurivich DA (2021). Handgrip strength asymmetry is associated with future falls in older Americans. *Aging Clin Exp Res* **33,** 2461–2469.

McGrath TM, Waddington G, Scarvell JM, Ball NB, Creer R, Woods K & Smith D (2016). The effect of limb dominance on lower limb functional performance--a systematic review. *J Sports Sci* **34,** 289–302.

McIntyre S, Nagi SS, McGlone F & Olausson H (2021). The Effects of Ageing on Tactile Function in Humans. *Neuroscience* **464,** 53–58.

van Melick N, Meddeler BM, Hoogeboom TJ, Nijhuis-van der Sanden MWG & van Cingel REH (2017). How to determine leg dominance: The agreement between self-reported and observed performance in healthy adults. *PLoS One* **12,** e0189876-.

Nakagawa S & Schielzeth H (2013). A general and simple method for obtaining R2 from generalized linear mixed-effects models. *Methods Ecol Evol* **4,** 133–142.

Newton RU, Gerber A, Nimphius S, Shim JK, Doan BK, Robertson M, Pearson DR, Craig BW, Häkkinen K & Kraemer WJ (2006). Determination of functional

strength imbalance of the lower extremities. *J Strength Cond Res* **20,** 971–977.

Oliviero A, Profice P, Tonali PA, Pilato F, Saturno E, Dileone M, Ranieri F & Di
Lazzaro V (2006). Effects of aging on motor cortex excitability. *Neurosci Res* **55,**
74–77.

Oomen NMCW & van Dieën JH (2017). Effects of age on force steadiness: A literature
review and meta-analysis. *Ageing Res Rev* **35,** 312–321.

Orssatto LBR, Borg DN, Pendrith L, Blazevich AJ, Shield AJ & Trajano GS (2022). Do
motoneuron discharge rates slow with aging? A systematic review and meta-
analysis. *Mech Ageing Dev* **203,** 111647.

Papitsa A, Paizis C, Papaiordanidou M & Martin A (2022). Specific modulation of
presynaptic and recurrent inhibition of the soleus muscle during lengthening and
shortening submaximal and maximal contractions. *J Appl Physiol* **133,** 1327–1340.

Pethick J, Taylor MJD & Harridge SDR (2022). Aging and skeletal muscle force control:
Current perspectives and future directions. *Scand J Med Sci Sports* **32,** 1430–1443.

Petrovic I, Amiridis IG, Holobar A, Trypidakis G, Kellis E & Enoka RM (2022). Leg
Dominance Does Not Influence Maximal Force, Force Steadiness, or Motor
Unit Discharge Characteristics. *Med Sci Sports Exerc* **54,** 1278–1287.

Portegijs E, Sipilä S, Alen M, Kaprio J, Koskenvuo M, Tiainen K & Rantanen T (2005).
Leg extension power asymmetry and mobility limitation in healthy older women.
Arch Phys Med Rehabil **86,** 1838–1842.

Rahnama N, Lees A & Bambaecichi E (2005). Comparison of muscle strength and
flexibility between the preferred and non-preferred leg in English soccer players.
Ergonomics **48,** 1568–1575.

Ruas C V, Taylor JL, Latella C, Haff GG & Nosaka K (2025). Neuromuscular
characteristics of eccentric, concentric and isometric contractions of the knee
extensors. *Eur J Appl Physiol* **125,** 671–686.

Skelton DA, Kennedy J & Rutherford OM (2002). Explosive power and asymmetry in
leg muscle function in frequent fallers and non-fallers aged over 65. *Age Ageing*
31, 119–125.

Strong A, Grip H, Arumugam A, Boraxbekk C-J, Selling J & Häger CK (2023). Right
hemisphere brain lateralization for knee proprioception among right-limb dominant
individuals. *Front Hum Neurosci*. Available at:
https://www.frontiersin.org/journals/human-
neuroscience/articles/10.3389/fnhum.2023.969101.

Del Valle A & Thomas CK (2005). Firing rates of motor units during strong dynamic
contractions. *Muscle Nerve* **32,** 316–325.

Del Vecchio A, Negro F, Holobar A, Casolo A, Folland JP, Felici F & Farina D (2019).
You are as fast as your motor neurons: speed of recruitment and maximal
discharge of motor neurons determine the maximal rate of force development in
humans. *J Physiol* **597,** 2445–2456.