Title: Hologenomic structure of bacterial and fungal community composition in the West Nile virus

vector *Culex tarsalis*

Authors:

Eunho Suh[1], Naomi Huntley[2,3], Travis van Warmerdam[1], Jamie P. Spychalla[4], Najmeh Nejat[1], and Jason L.

Rasgon[1,3,5,6,7]*

Affiliation:

[1]Department of Entomology, The Pennsylvania State University, University Park, PA, USA

[2]Department of Biology, The Pennsylvania State University, University Park, PA, USA

[3]One Health Microbiome Center, The Pennsylvania State University, University Park, PA, USA
12

13 [4]Department of Plant Pathology and Environmental Microbiology, The Pennsylvania State University,
14 University Park, PA, USA
15

16 [5]Department of Biochemistry and Molecular Biology, The Pennsylvania State University, University Park,
17 PA, USA
18

19 [6]The Center for Infectious Disease Dynamics, The Pennsylvania State University, University Park, PA,
20 USA
21

22 [7]The Huck institutes of the Life Sciences, The Pennsylvania State University, University Park, PA, USA
23

* Correspondence to be sent to jlr54@psu.edu

Running title:

Microbiome and population genetics of *Culex tarsalis*

Abstract

Background

Microbiota play a crucial role in determining the ability for arthropod disease vectors to transmit

pathogens. Microbial community structure can be heavily influenced by microbe-microbe interactions,

host genetics and environmental factors. Here, we characterize the host population genetic structure, and

bacterial and fungal communities in natural populations of the West Nile virus mosquito vector *Culex*

tarsalis. Mosquitoes were collected and analyzed across the species range of the mosquito in the United

States, where we used PoolRADSeq to quantify population genetic structure. Microbial community

composition was characterized using bacterial 16S rRNA gene sequencing (V3-V4 region) and fungal ITS

sequencing (ITS1 region).

Results

PoolRADSeq identified four broad genetic clusters of mosquito populations, which corresponded to

previous clusters identified by microsatellite analysis and RADSeq on individual mosquitoes.

Microbiome diversity grouped mosquito populations into three broad clusters, with each cluster

distinctively represented by diagnostic abundant bacteria (*Ralstonia*, *Pseudomonas,* or

Zymobacter/*Providencia*, respectively). Clustering for fungal taxa was less pronounced. Geographic

distance between populations was positively correlated with microbiome community dissimilarity, and

multiple environmental factors were significantly correlated with microbial species richness and diversity.

Conclusions

These results suggest that bacterial and fungal communities are geographically structured in *Cx. tarsalis*,

interact with important environmental factors, and are partially correlated with host genetic structure. As

microbiota can affect the ability for mosquitoes to transmit pathogens, understanding the factors

underpinning microbiome variation across space and time has important implications for the spread of

vector-borne pathogens such as WNV.

Keywords: West Nile virus, microbiome, bacteria, fungi, mosquito, hologenome, RADSeq

Introduction

Culex tarsalis is one of the most important native North American mosquito arbovirus vectors [1]. The species is widely distributed in western North America, feeds primarily on avian hosts which contributes to virus amplification [2], and can move relatively long distances in search of a blood meal or oviposition sites [1,2,3]. Historically *Cx. tarsalis* is an important vector of Western equine encephalitis virus (WEEV) and Saint Louis encephalitis virus (SLEV) [4]. When West Nile virus (WNV) was introduced into the Americas in 1999, *Cx. tarsalis* was demonstrated to be one of the most competent vectors of this novel invasive pathogen [5].

Previously, population genetic data was used to implicate a role for *Cx. tarsalis* in transporting WNV across the western United States. An analysis of microsatellite loci suggested that across the western USA, *Cx. tarsalis* populations are structured into approximately three broad metapopulation clusters ("Midwest", Sonoran [or "Southwest"], and "Pacific"), which closely matched the observed pattern of WNV geographic spread in the early 2000s [1,3]. A more recent population genetic analysis, using RADSeq on approximately 300 individual mosquitoes, identified a broadly similar pattern (with increased resolution and a new identified cluster ["Northwest"] due to the denser marker set) [6].

Knowledge of the factors influencing the spread of arboviruses in mosquito populations is critical for understanding dynamics of disease incidence, development of risk assessment strategies for novel virus introductions and development of virus transmission biomarkers that can be used to efficiently target control efforts [1,7]. In general, there are three main barriers that an arbovirus must overcome to be transmitted by a mosquito vector. The virus particles must 1) establish an infection of the midgut epithelium and replicate, 2) escape the midgut through the basal lamina and infect other body tissues, and 3) infect the salivary glands, escape into the salivary gland lumen and be transmitted during salivation [8,9,10]. These barriers can be modulated by the intrinsic host genetics of the mosquito, components of the endogenous mosquito microbiota, and environmental factors, which has been demonstrated for multiple mosquito species in both the lab and the field [11,12,13,14,15,16,17]. It has become increasingly clear that while pathogen transmission can be affected by host genetics and environmental factors, this is, at best, an incomplete representation of the factors influencing pathogen transmission phenotypes. Recent research on arthropod holobiomes (the host plus all associated microorganisms) has demonstrated the potential impact of mosquito-associated microorganisms on transmission-related phenotypes of a variety of pathogens [15,16,17,18,19,20,21,22, 23,24,25,26,27,28].

Laboratory studies have demonstrated that components of the mosquito holobiome such as bacteria and fungi can modulate pathogen vector competence phenotypes. For example, bacteria in the genera

Chromobacterium, *Enterobacter*, *Serratia*, *Pseudomonas,* and *Delftia* can decrease *Plasmodium* oocyst

levels in *Anopheles* mosquitoes [17,20,21,23], while bacteria in the genera *Serratia* and *Aeromonas* can

increase mosquito susceptibility to arboviruses [22,24]. Less is known about how fungi alter pathogen

vector competence in mosquitoes, but examples exist. Specific isolates of *Penicillium* enhanced

Anopheles infection with *Plasmodium* [25], while *Metarhizium anisopliae* infection rendered *Aedes*

mosquitoes less susceptible to dengue virus [26]. These data suggest that components of the mosquito

microbiome can be potent modulators of pathogen transmission, but despite their importance there is very

little data on what role these microorganisms play in determining vector competence variation in the field.

Little is known about the natural microbiome of *Cx. tarsalis* or its effects on pathogen infection and

transmission. Current studies are limited to characterizing bacterial communities in *Cx. tarsalis*

mosquitoes reared in experimental mesocosms, where communities were dominated by the aquatic

bacteria *Thorsellia* [29, 30]; a common microbe found in freshwater aquatic environments [30]. Nothing

is currently known about fungal communities in this species. To address this knowledge gap, we

examined the bacterial and fungal communities in wild-caught *Cx. tarsalis* mosquitoes from populations

across the species range in the western USA. Combined with host RADSeq data, we found signatures of

population structuring in microbial communities that were partially correlated with mosquito population

genetic structure, where mosquitoes clustered into three broad groups based on their microbial

composition. Microbial structure was more pronounced for bacteria than fungi, possibly reflecting

differences in microbe acquisition mechanisms. Geographic distance between populations was positively

correlated with microbial community dissimilarity. Environmental factors such as longitude, latitude,

temperature, and precipitation were significantly correlated with species richness and/or diversity. These

results suggest that bacterial and fungal communities in this mosquito are geographically structured, are

potentially shaped by host genetics, and interact with or are affected by environmental factors. These

results have critical significance for understanding patterns of microbial ecology in this mosquito species,

as well as having implications for the epidemiology of vector-borne pathogens such as West Nile virus.

Methods

PoolRADSeq

2b-RADseq on pooled mosquitoes was used to characterize *Cx. tarsalis* population structure. Wild *Cx.*

tarsalis mosquitoes were collected from 15 populations in 2021 and 18 populations in 2022 using CDC

traps (Fig 1, Table 1). Mosquitoes were shipped overnight on dry ice to Penn State and stored at -80 °C

until processing. For each population, we created pools of 10 randomly selected mosquitoes. DNA was

extracted from individuals using Qiagen spin columns, quantified with Qubit, and equal amounts of DNA

from each individual were combined to form each pool. After DNA QC, standard 5' –NNN- 3' adaptors

were used to ligate restriction tags for concatenated Isolength restriction site-associated tags sequencing.

Genomic DNA was digested with the type IIB restriction enzyme BsaXI to produce restriction fragments

of uniform length, followed by adaptor ligation, amplification, purification, concatenating and barcoding

to incorporate sample-specific barcodes. Samples were then pooled, QC assessed, and sequenced on the

Illumina Hiseq Xten/NovaSeq platform using paired end 150bp sequencing. Reads were quality filtered.

Reads containing the restriction endonuclease recognition site were extracted from the sequencing data

and clustering was performed with ustacks software (version 1.34) from the Stacks package to construct a

reference sequence. Sequencing data was aligned to the reference sequence using SOAP software (version

2.21), and SNP calling was performed using Maximum Likelihood (ML) using the RADtyping package

[31]. A total of 9,626 quality-filtered SNP markers were obtained across all samples. Phylogenetic trees

were constructed using Neighbor Joining. Library construction, sequencing, and analysis were conducted

by Novogene.

Microbiome mosquito collection and DNA extraction

Cx. tarsalis mosquitoes were collected from 17 locations in five states (CA, CO, UT, TX, and WA) in

2021 (Fig 2, Table 2). Mosquitoes were collected using CO_2 baited EVS (encephalitis vector surveillance)

or CDC traps overnight and morphologically identified at species level by the collecting laboratory.

Mosquito samples were stored in DNA/RNA Shield (Zymo Research, Irvine, USA) at -80C until they

were shipped to Penn State University. Morphological species identification of *Cx. tarsalis* was confirmed,

and individual mosquitoes were surface sterilized in 500 µl 70% ethanol for 5 minutes followed by 3

washes in PBS. All tools used for handling mosquitoes were sterilized using absolute ethanol to minimize

cross contaminations. DNA extractions were performed using ZymoBiomics DNA/RNA miniprep kit

(Zymo Research, Irvine, USA) following the manufacture's protocol with some modifications. In brief,

individual whole mosquitoes were homogenized in 2.0mm Zymo BeadBeating Lysis tubes (Zymo

Research, Irvine, USA) for 30 second at 6m/s in 750ml Zymo DNA/RNA Shield lysis buffer (Zymo

Research, Irvine, USA) using a Fisher brand Beadmill 24 (Fisher Scientific, Waltham, MA) followed by

additional mechanical homogenization in 0.1-0.5mm BeadBeating Lysis tubes for two 30 second cycles at

6m/s with a 45 second dwell. The homogenization protocol was optimized to ensure extraction of both

bacterial and fungal DNA from a single mosquito while minimizing DNA degradation by excessive

shearing. To minimize batch effects, two randomly selected populations were processed per day by two

personnel, with a blank (PBS) extraction added as negative control for each batch (total 16 negative

controls). All negative controls were determined to be negative for 16S PCR and did not pass QC for

library preparation. For positive control, two units of ZymoBIOMICS Microbial Community Standard

(Zymo Research, Irvine, USA) were used to validate our extraction method by confirming detection of
known bacterial and fungal species in the standard mixture. Quantity and quality of DNA were checked
using Qubit 4.0 fluorometer (Fisher Scientific, Waltham, USA) and nanodrop, respectively. DNA
samples were stored at -80C until down-stream processing. Sample sizes and collection locations are
listed in Table 2.
167

Microbiome sequencing and bioinformatics analysis
Bacterial and fungal communities were characterized by splitting the DNA from same individual
mosquito. For bacterial characterization, the V3-V4 hypervariable region of 16S rRNA gene was
amplified using primers 341F (5'- CCTAYGGGRBGCASCAG-3') and 806R (5'-
GGACTACNNGGGTATCTAAT-3') with expected amplicon size of approximately 450-550bp. For
fungal characterization, the Internal Transcribed Spacer (ITS) 1 region was amplified using primers
ITS1F (5'-CTTGGTCATTTAGAGGAAGTAA-3') and ITS2 (5'-GCTGCGTTCTTCATCGATGC-3')
with expected amplicon size of approximately 200-400bp. PCR was carried out in 30 μL reactions with
15 μL of Phusion High-Fidelity PCR Master Mix (New England Biolabs), 0.6 mM of forward and reverse
primers, and 5~10 ng of template DNA (10 μL for negative controls). Thermal cycling for bacterial
characterization consisted of initial denaturation at 98ºC for 1 min, followed by 30 cycles of denaturation
at 98ºC for 10 sec, annealing at 50ºC for 30 sec, and extension at 72ºC for 30 sec, and lastly by 72ºC for 5
min. The same cycle was used for fungal characterization, except for the annealing temperature of 55ºC
for 30 sec. PCR products within the expected size were selected by 2% agarose gel electrophoresis. The
same amount of PCR product from each sample was pooled, end-repaired, A-tailed and further ligated
with Illumina adapters. PCR products were purified using the Qiagen Gel Extraction Kit (Qiagen,
Germany). Sequencing libraries were generated using NEBNext Ultra DNA Library Pre ®Kit for Illumina.
Libraries were commercially sequenced (Novogene) to generate 250 bp paired-end raw reads. The library
was checked by Qubit and real-time PCR for quantification, and bioanalyzer for size distribution
detection. Quantified libraries were pooled and sequenced on the NovaSeq PE250 platform (Illumina, San
Diego, CA, USA).For sequencing and data processing, paired-end reads were assigned to samples and
truncated by cutting off the barcode and primer sequences. After truncation, FLASH (v1.2.11,
ccb.jhu.edu/software/FLASH/) was used to merge reads to get raw tags [2]. Then, fastp software was used
to perform quality control of raw tags, and high-quality clean tags were obtained. Finally, chimeras were
removed using VSEARCH. For denoising and species annotation of ASVs (Amplicon Sequencing
Variants), DADA2 was used to reduce noise and sequences with abundance less than 5 were filtered out
to obtain the final ASV [32]. Representative sequences for each ASV were annotated to obtain the
corresponding species identification using QIIME2. By applying QIIME2's classify-sklearn algorithm

[33,34], a pre-trained Naive Bayes classifier was used for species annotation of each ASV. PCR, library

preparation, sequencing and data processing were performed by Novogene.

Microbiome data and statistical analysis

Data and statistical analyses were conducted using MicrobiomeAnalyst 2.0 [35], GraphPad Prism (v8.0),

and R. For data filtering prior to downstream analyses, features that were constant across all samples and

singletons (those with only one total count) were removed. To eliminate low-quality or uninformative

features, the low-count filter was set to zero to avoid random selection bias for tied low-count data, and

the low-variance filter was set to 10% interquartile range, followed by a data normalization process.

Filtered data were used for all downstream comparative analyses unless otherwise specified. To

understand overall bacterial and fungal community structure, we first constructed relative abundance and

heat trees of representative taxa. Based on visual inspection of these data, we hypothesized a clustering of

mosquito populations and validated this grouping through a series of comparative analyses including

Alpha and Beta diversity, Random Forest classification, Core microbiome analysis, dendrogram

construction, Linear Discriminant Analysis Effect Size (LEfSe), correlation network analysis, and

correlations between environmental factors, community structure, and geographic distance. For Alpha

diversity analysis, non-filtered original data were used. For Random Forest classification, five

consecutive Out-of-Bag (OOB) error tables were generated to estimate the mean OOB error. Correlation

network analysis employed Sparse Estimation of Correlations among Microbiomes (SECOM) with

Pearson1, which efficiently handles both linear and non-linear relationships among microbes while

maintaining sparsity. Correlation analyses between geographical distance and community dissimilarity

indices were performed using the Mantel test (*vegan* package) in R. All other analyses performed in

MicrobiomeAnalyst 2.0 followed the default settings as recommended [35]

Wolbachia **verification**

Wolbachia was detected by 16S amplicon sequencing in a few (12/197) samples. For these samples we

attempted to verify this finding by amplifying and sequencing the *Wolbachia* surface protein (*wsp*) gene

using primers 81F and 691R [36]. We also amplified and sequenced the mosquito mitochondrial ND4

gene [37] to verify mosquito species identification. PCR and Sanger sequencing for these genes was

carried out as described in the respective references.

Results

RADSeq population genetics

A total of 9,626 SNP markers that passed quality control were obtained across all samples, which were

distributed across UTRs, introns, exons, and intergenic regions. Results from both years were combined

for analysis to obtain a more complete spatial sampling across the species range of the mosquito.

Clustering neighbor-joining results from PoolRADSeq were qualitatively similar to previous results from

RADSeq on individual mosquitoes [6] and microsatellites [1,3], resolving the same metapopulation

clusters (Pacific, Sonoran/Southwest, Northwest, and Midwest) that previous analyses [1,3,6] identified

(Fig 3).

Microbiome composition profile

For processing bacterial sequence data, total read count was 27,771,776, and the average count per

sample was 140,973 (Maximum = 502,316; Minimum = 10,132) from 197 samples that passed QC. A

total 107 ASVs were left of 66396 by removing singletons (one total count) and low abundance ASVs

(less than 4 read count). For fungal sequence data, total read count was 16,402,771 and the average count

per sample was 227,816 (Maximum = 792,580; Minimum = 23,624) from 72 samples that passed QC. A

total of 42 ASVs were left of 8,261 by removing singletons and low abundance ASVs. For bacterial and

fungal combined sequence data, total read count was 26,776,978 and the average count per sample was

377,140 (Maximum = 1,001,576; Minimum = 76,642) from 71 samples. 163 ASVs were left of 74,657 by

removing singletons and low abundance ASVs.

We first investigated overall community composition in bacteria and fungi in across all *Cx. tarsalis*

populations. For bacterial communities, *Proteobacteria* were the most abundant at the phylum level

(89.8%) which was mainly composed of *Gammaproteobacteria* (85.7% of total abundance) (Fig 4;

Additional file 1). Within *Gammaproteobacteria*, *Burkhoderiales* was most abundant followed by

Pseudomonadales, *Enterobacterales* and *Oceanospirillales*, representing over 85% of total abundance at

the order level. Within *Gammaproteaobacteria*, *Ralstonia*, *Pseudomonas*, *Zymobacter*, and *Providencia*

were the most abundant taxa representing 75% of total abundance at the genus level. While the top three

abundant taxa were found in 74% to 96% of all 197 samples, *Providencia* was relatively less prevalent

(25%).

We detected *Wolbachia* in approximately 6% (12/197) of samples. In previous surveys, *Cx. tarsalis* was

found to be uninfected with *Wolbachia* [39], so we attempted to further validate this finding. From these

12 samples, we amplified and sequenced the *Wolbachia wsp* gene to strain type the infection. PCR was

successful in only 2 samples, where the infection was identified as *w*Pip (the canonical *Wolbachia* strain

infecting *Cx. pipiens/quinquefasciatus*; [40]); the remaining 10 samples either did not amplify or

amplified random mosquito DNA. We then amplified and sequenced the mosquito mitochondrial ND4

gene to molecularly identify the mosquitoes. The two verified *Wolbachia*-positive samples were

identified as *Cx. quinquefasciatus*. The other 10 samples were identified as *Cx. tarsalis*. These data

indicate that *Wolbachia* infection is not present in *Cx. tarsalis* mosquitoes and the observed rare positive

samples were due to sample misidentification or potentially contamination. The two *Cx. quinquefasciatus*

samples were removed from further downstream analysis.

For fungal communities, *Ascomycota* and *Basidiomycota* were the two most abundant phyla representing

over 79% of total abundance (Fig 5, Additional file 1). While *Basidomycota* was mainly composed of

Agaricomyctetes (21.4%), *Ascomycota* was mostly represented by *Dothideomycetes* (53.1%) at the class

level. Within *Dothidemycetes*, *Capnodiales* and *Pleosporales* were the two most abundant taxa at the

order level representing over 53% of total abundance. *Mycosphaerella* (28.4%) and *Cladosporium* (16%)

were the most abundant within *Dothideomycetes* while *Alternaria* was most abundant (3.8%) within

Pleosporales at the genus level, and these three taxa were detected in 83 to 94% of all 72 samples. While

Ganoderma was the second most abundant taxa (21.4%) belonging to *Basidiomycota*, its prevalence was

relatively lower (36%).

Clustering of mosquito populations by microbial community structure

We identified a pattern where specific bacterial taxa at the genus level were more abundant in certain

clusters of populations. Based on visual inspection and geographic proximity, we hypothesized three

clustering groups of mosquito populations; CA1, CA2, CO1, CO2, CO3, TX1, and TX2 for group CCT

(initials of California, Colorado and Texas), CA3, CA4, CA5, and CA6 for group CA (California), and

UT1, UT2, UT3, WA1, WA2, and WA3 in group UW (initials of Utah and Washington) (Fig. 6A and

Additional file 2). With this grouping, *Ralstonia* and *Pseudomonas* were the most abundant taxa in group

CCT (70.2% relative abundance) and UW (66%), respectively, while *Zymobacter* (35.6%) and

Providencia (31.3%) were most abundant in group CA (Additional file 2, 3).

Clustering of fungal community structure according to this grouping was less pronounced (Fig. 6B;

Additional file 2, 4). For example, while *Ganoderma* was most abundant in UW (36.5% relative

abundance) and *Cladosporium* was relatively more abundant in CA (37.8%) than CCT (17.8%) or UW

(6.3%), *Mycosphaerella* were found in all three groups with similar abundance ranging from 18.3 to

33.1%. Similarly to the case of bacteria, grouping was more pronounced at order or lower taxonomic

levels (Additional file 4). When community structure of bacteria and fungi were combined, clustering was

moderate, similarly to the case of fungal community (Fig 6C; Additional file 2; Additional file 5) as

relative abundance of major taxa identified above (i.e., *Pseudomonas*, *Zymobacter*, *Providencia*,

Mycosphaerella, *Cladosporium* and *Ganoderma*) followed the hypothesized grouping model.

Alpha and Beta diversity, and Random Forest classification

Alpha diversity analyses, Chao1 (species richness) and Shannon (species diversity) indices were

compared for community structure of bacteria, fungi, or bacterial/fungi combined (Fig. 7A). Chao1

indices were significantly different among the three groups ($P < 0.05$, Kruskal-Wallis) with higher index

value in group CCT relative to CA ($P = 0.006$, Bonferroni corrected after pair-wise comparison among

three groups) while Shannon indices were not different among three groups ($P > 0.05$, Kruskal-Wallis).

For fungal communities, Chao1 indices were similar among three groups while a difference was observed

for Shannon indices among the groups ($P < 0.01$, Kruskal-Wallis). When bacteria/fungi were combined,

no difference was observed for Chao1 indices among the groups ($P > 0.05$, Kruskal-Wallis), but Shannon

index was lowest for UW ($P < 0.001$, Kruskal-Wallis; Bonferroni corrected after pairwise comparison).

Rarefaction curve analyses and comparison of sequencing depth among three groups indicated overall

high sequencing depth across samples with minimal heteroskedasticity regardless of data filtering

(Additional file 6). Beta diversity was next explored using Nonmetric Multidimensional Scaling (NMDS),

revealing a significant differences in community structure among the three groups (PERMANOVA, $P <$

0.01) for bacterial, fungal, and bacterial/fungal combined (Fig. 7B).

We then performed machine learning based Random Forest classification analyses to validate our

grouping model (Fig. 8). For bacterial communities, the mean Out-of-bag (OOB) error was less than 0.1

in five consecutive runs (Fig. 8A; Additional file 7) indicating that about 90% of predictions were correct

by our grouping model, and *Ralstonia* was identified to have the greatest influence on the classification

model followed by *Providencia*, *Pseudomonas*, *Collinsella* and *Zymobacter* (Fig. 8B). Similarly,

dendrogram analysis largely supported our clustering model regardless of algorithms applied (Additional

file 8). Classification was less clear for fungal data with slightly higher OOB error (0.228) (Fig. 8A;

Additional file 7, 9). The top five fungal genera that had greatest influence on the classification were

Sporobolomyces, *Sarocladium*, *Ganoderma*, *Cladosporium*, and *Aspergillus*; *Mycosphaerella* (the most

abundant fungus) had relatively little influence (Fig. 8B). For combined bacterial/fungal data,

intermediate OOB error (0.152) was observed (Fig. 8A; Additional file 7, 10), and *Zymobacter* had

greatest influence on classification followed by *Providencia*, *Sporobolomyces*, *Pseudomonas*, and

Saccharibacter (Fig. 8B).

Core microbiome, taxon differentiation, and correlation network

Prevalence (% detected in samples) profiles were characterized using Core microbiome analysis (Fig. 9).

For bacterial data, *Ralstonia* was most prevalent followed by *Pseudomonas*, *Zymobacter*, *Entomospria*

and *Escherichia* in all samples. When each group was analyzed separately, *Ralstonia* and *Pseudomonas*

were most prevalent in group CCT and UW, respectively while *Zymobacter* and *Providencia* were most

prevalent in group CA. For fungal data, *Cladosporium* and *Mycosphaerella* were widely found in all

three groups while *Alternaria* and *Ganoderma* were found in CCT and UW respectively. A similar pattern

was observed when bacteria/fungal data were combined.

The major taxa identified in Core microbiome analyses were consistently observed in Linear Discriminant

Analysis with effect size (LEfSe) for bacterial data (Additional File 11). *Ralstonia* and *Pseudomonas*

were significantly more abundant in CCT and UW, respectively while *Zymobacter* and *Providencia* were

more abundant in group CA. However, for fungal data, *Cladosporium* and *Ganoderma* were more

abundant in group CA and UW, respectively, but *Mycospharella* was not significantly enriched in either

of three groups. A similar pattern was observed for combined bacterial/fungal data.

To explore whether interactions among taxa had influence on our clustering model, we investigated

correlation network for the major taxa identified above. For bacteria, *Ralstonia*, *Pseudomonas*, and

Zymobacter were uncorrelated (Additional file 12; Additional file 13) while for fungi, *Mycosphaerella*

and *Cladosporium* were positively correlated, while *Mycosphaerella* was negatively correlated with

Alternaria. In total, 33 significant correlations were identified between bacteria and fungi, and 85% (28

cases) of them were negative. The two most abundant fungi, *Mycosphaerella* and *Cladosporium,*

accounted for 50% (14 cases) of total negative correlations (Additional file 14A). Consistent with this

result, species richness (Observed ASV/Chao1) was negatively correlated between bacteria and fungi (R

$= -0.3$, $P = 0.0119$) (Additional file 14B). These results indicate a possible exclusive interaction between

bacteria and fungi within the mosquito microbiota.

Relationship between environmental factors and community structure

Because *Cx. tarsalis* population genetic structure was recently observed to be correlated with

environmental factors [6], we explored whether environmental factors were correlated with microbial

community structure or species richness/diversity in this mosquito. We first examined the relationship

between environmental factors (e.g., longitude, latitude, temperature, and precipitation) and Alpha

diversity indices (Fig. 10A). For bacteria, ASV/Chao1 and ACE were positively correlated with longitude

and precipitation while these two indices were negatively correlated with temperature. For fungi,

ASV/Chao1 and ACE were negatively correlated with longitude while these two indices together with

Shannon and Simpson indices were positively correlated with temperature. Latitude was also positively

correlated with Shannon and Simpson indices. For combined bacterial/fungal data, no correlation was

found with longitude for any indices, but all five indices were positively correlated with temperature and

negatively correlated with latitude. These results suggest species richness/diversity is shaped by multiple

environmental factors. Particularly, species richness of bacteria and fungi were negatively correlated

along longitude.

We noted a significant positive correlation between geographic distance and community dissimilarity for

bacterial, fungal and combined bacteria/fungal data (Mantel test, $P < 0.01$; Fig. 8B), indicating that the

community structure is more dissimilar as distance between collection sites increases for both bacteria

and fungi and suggesting possible phylogenomic signal of microbial diversity in *Cx. tarsalis*.

Discussion

Our population genetic analysis of *Cx. tarsalis* across the western USA identified essentially the same

population structure that previous microsatellite and individual RADSeq analyses. In total, these analyses

demonstrate that gene flow is substantial between *Cx. tarsalis* populations across wide geographic

distances, and that the metapopulation structure can be defined in terms of 3-4 large metapopulation

clusters encompassing the Midwest, the Pacific Coast, the Sonoran desert, and potentially the Northwest

[1,3,6]. Although it does not resolve individuals, PoolRADSeq produced similar population-level results

at significantly lower cost.

There are no previous studies on the diversity of fungi colonizing *Cx. tarsalis*, and previous bacterial

microbiome studies in this species were limited to analysis of mosquitoes that were reared in artificial

field mesocosms [29]. These mesocosms were constructed in an attempt to mimic the natural habitat of

this mosquito; however, the resulting microbial composition data differed significantly compared to our

results using true wild-caught *Cx. tarsalis*. The bacterial microbiomes of mesocosm-reared mosquitoes

were dominated by *Thorsellia* [29] a common aquatic bacterial taxon that has been previously associated

with mosquitoes and other aquatic species [30, 41], and likely reflects the unnatural microbial

colonization of the mesocosm habitats. In our study, we observed that distinct microbial taxa were

enriched in wild-reared mosquitoes, and these taxa differed depending on the geographic region the

mosquitoes were collected from. These data suggest that microbiome studies should, if at all possible, be

conducted on individuals that were collected from natural habitats, as artificial habitats may not reflect the

microbial diversity observed in nature and can give misleading results.

Wolbachia is a maternally inherited alphaproteobacterial symbiont that is of interest as an agent for
control of arbovirus transmission in mosquitoes [27,28,29,40,43,44]. Previous studies suggested that *Cx.*
tarsalis is not naturally infected with *Wolbachia* [39]. Artificial transfection with *Wolbachia* in *Cx.*
tarsalis was shown to increase, rather than decrease, their susceptibility to WNV, thus detection of
Wolbachia in this species has significant implication for the epidemiology of WNV transmission (and
potentially other arboviruses) [45]. Therefore, when we detected *Wolbachia* by 16S amplicon sequencing
in a small subset of individuals we attempted to independently validate the finding. Out of 12 samples
where *Wolbachia* was detected by amplicon sequencing, we were able to independently validate the
infection in only two individuals, which were ultimately molecularly identified as *Cx. quinquefasciatus;* a
species which is known to be natively infected with *Wolbachia* [39,40]. In the remaining 10 samples,
Wolbachia infection could not be verified, and these samples were molecularly identified as *Cx. tarsalis*.
In total, these data affirm previous findings that *Cx. tarsalis* is not naturally infected with *Wolbachia* and
highlight the importance of independently verifying low frequency *Wolbachia* infections detected solely
on the basis of 16S amplicon microbiome sequencing.

Recent studies identified correlations between *Cx. tarsalis* population genetics and environmental
variables such as temperature, humidity, and vegetation [6]. We therefore examined potential links
between microbiome composition and environment. We found that bacterial diversity was positively
correlated with longitude and precipitation, and negatively correlated with temperature. Fungal diversity
was negatively correlated with longitude and positively correlated with temperature. The mechanisms
underlying these relationships are at present unclear, but the data suggest that correlations between
environmental parameters and mosquito genetics are confounded by microbial interactions. Future studies
are indicated to tease apart the mechanisms underlying these interactions.

Our data suggests that microbiome structure in *Cx. tarsalis* is, to some extent, correlated with population
genetic structure. While this relationship is not completely one-to-one with the genetic clusters defined by
RADSeq or microsatellites, it is significant and partially overlaps with these clusters. Mantel regression
demonstrates significant isolation by distance in microbial structure. Phylogenomic signal is more
pronounced for bacterial data compared to fungi. Generally, mosquitoes acquire bacterial microbiome
associates from the aquatic larval environment [19,46]. Less is known about how mosquitoes acquire
fungal associates. Differences between bacteria and fungi could reflect different modes of acquisition
between these two groups, differences in retention between bacteria vs. fungi during mosquito
development, or differences in mosquito immune response to bacteria vs. fungi in response to
colonization.

Conclusions

Microbial associates have been demonstrated to affect pathogen infection and transmission in a variety of

mosquito-pathogen systems [15,16,17,18,19,20,21,22, 23,24,25,26,27,28]. This is the first study to

examine diversity of microbial association in natural populations of *Cx. tarsalis* across its species range.

Identification of microbial taxa that can affect WNV or other pathogens in this mosquito has significant

implications for understanding the spread and invasion of the viral pathogens transmitted by this

mosquito across space and time.

Acknowledgements

We thank our collaborators for providing mosquito samples used in this analysis, including Anna Wanek,

Broox Boze, Jason Williams, Nikki Harris and Will Schlatmann (Vector Disease Control International),

Conlin Reis (Fresno Westside MAD), Corey Brelsfoard (Texas Tech University), Gary Hatch (Mosquito

Abatement District-Davis), Jared Larmirante (Cass County Vector Control), Jeff Hamik (Nebraska

Department of Health and Human Services), John Albright (Shasta Mosquito and Vector Control District),

Joshua Blystone (Cascade County Weed & Mosquito Division), Kevin Shoemaker (Benton County

Mosquito Control District), Kim Hung (Coachella Valley Mosquito & Vector Control District), Ryan

Smith (Iowa State University), Sarah Wheeler (Sacramento-Yolo Mosquito & Vector Control District),

Scott Bradshaw (Tooele Valley Mosquito Abatement District), Scott Larson (Metropolitan Mosquito

Control District), and Todd Hanson (Mosquito Control Division, Grand Forks Public Health Department).

Funding

455 This study was supported by NIH/NIAID grant R01AI150251, USDA Hatch funds (4769), and funds

456 from the Dorothy Foehr Huck and J. Lloyd Huck endowment to J.L.R.

Author contributions

459 All authors contributed intellectually to and agreed to this submission. ES and JLR designed the

460 experiments. ES (and others acknowledged above) collected field mosquito samples. ES, TW, and NN

461 conducted experiments and collected the data. ES, NH, and JPS analyzed the data. ES and JLR wrote the

462 manuscript, and NH, TW, JPS, and NN provided feedback. The authors read and approved the final

463 manuscript.

Availability of data and materials

Raw bacterial microbiome data, fungal microbiome data, and PoolRADseq data are available at

zenodo.org/records/14081783, zenodo.org/records/14080098, and zenodo.org/records/14081214.

Code availability

Code for analyses is available at zenodo.org/records/17014950.

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare no competing interests.

References

1. Venkatesan M, Rasgon JL. Population genetic data suggest a role for mosquito-mediated dispersal
of West Nile virus across the western United States. Mol Ecol. 2010; 19:1573-84.

2. Reisen WK. The contrasting bionomics of Culex mosquitoes in western North America. J Am
Mosq Control Assoc. 2012; 28(4 Suppl):82-91.

3. Venkatesan M, Westbrook CJ, Hauer MC, Rasgon JL. Evidence for a population expansion in the
West Nile virus vector Culex tarsalis. Mol Biol Evol. 2007; 24:1208-18.

4. Reisen WK, Lothrop HD, Hardy JL. Bionomics of Culex tarsalis (Diptera: Culicidae) in relation
to arbovirus transmission in southeastern California. J Med Entomol. 1995; 32:316-27.

5. Goddard LB, Roth AE, Reisen WK, Scott TW. Vector competence of California mosquitoes for
West Nile virus. Emerg Infect Dis. 2002; 8:1385-91.

6. Liao Y, Islam T, Noorai R, Streich J, Saski C, Cohnstaedt LW, Cooper EA. Climate Adaptation
and Genetic Differentiation in the Mosquito Species Culex tarsalis. Genome Biol Evol. 2025;
17:evaf143.

7. Hugo LE, Monkman J, Dave KA, Wockner LF, Birrell GW, Norris EL, Kienzle VJ, Sikulu MT,
Ryan PA, Gorman JJ, Kay BH. Proteomic biomarkers for ageing the mosquito Aedes aegypti to
determine risk of pathogen transmission. PLoS One. 2013;8:e58656. doi:
10.1371/journal.pone.0058656.

8. Hardy JL, Apperson G, Asman SM, Reeves WC. Selection of a strain of Culex tarsalis highly
resistant to infection following ingestion of western equine encephalomyelitis virus. Am J Trop
Med Hyg. 1978; 27(2 Pt 1):313-21.

9. Bosio CF, Beaty BJ, Black WC 4th. Quantitative genetics of vector competence for dengue-2
virus in Aedes aegypti. Am J Trop Med Hyg. 1998; 59:965-70.

10. Black WC 4th, Bennett KE, Gorrochótegui-Escalante N, Barillas-Mury CV, Fernández-Salas I, de
Lourdes Muñoz M, Farfán-Alé JA, Olson KE, Beaty BJ. Flavivirus susceptibility in Aedes
aegypti. Arch Med Res. 2002; 33:379-88.

11. Bennett KE, Flick D, Fleming KH, Jochim R, Beaty BJ, Black WC 4th. Quantitative trait loci that
control dengue-2 virus dissemination in the mosquito Aedes aegypti. Genetics. 2005; 170:185-94.

12. Alto BW, Bettinardi D. Temperature and dengue virus infection in mosquitoes: independent
effects on the immature and adult stages. Am J Trop Med Hyg. 2013; 88:497-505.

13. Bosio CF, Fulton RE, Salasek ML, Beaty BJ, Black WC 4th. Quantitative trait loci that control
vector competence for dengue-2 virus in the mosquito Aedes aegypti. Genetics. 2000; 156:687-98.

14. Bennett KE, Olson KE, Muñoz Mde L, Fernandez-Salas I, Farfan-Ale JA, Higgs S, Black WC
4th, Beaty BJ. Variation in vector competence for dengue 2 virus among 24 collections of Aedes
aegypti from Mexico and the United States. Am J Trop Med Hyg. 2002; 67:85-92.

15. Huang W, Rodrigues J, Bilgo E, Tormo JR, Challenger JD, De Cozar-Gallardo C, Pérez-Victoria I,
Reyes F, Castañeda-Casado P, Gnambani EJ, Hien DFS, Konkobo M, Urones B, Coppens I,
Mendoza-Losana A, Ballell L, Diabate A, Churcher TS, Jacobs-Lorena M. Delftia tsuruhatensis
TC1 symbiont suppresses malaria transmission by anopheline mosquitoes. Science. 2023;
381:533-540.
16. Angleró-Rodríguez YI, Blumberg BJ, Dong Y, Sandiford SL, Pike A, Clayton AM, Dimopoulos
G. A natural Anopheles-associated Penicillium chrysogenum enhances mosquito susceptibility to
Plasmodium infection. Sci Rep. 2016; ;6:34084.
17. Bahia AC, Dong Y, Blumberg BJ, Mlambo G, Tripathi A, BenMarzouk-Hidalgo OJ, Chandra R,
Dimopoulos G. Exploring Anopheles gut bacteria for Plasmodium blocking activity. Environ
Microbiol. 2014; 16:2980-94.
18. Hegde S, Rasgon JL, Hughes GL. The microbiome modulates arbovirus transmission in
mosquitoes. Curr Opin Virol. 2015; 15:97-102.
19. Foo A, Brettell LE, Nichols HL; 2022 UW-Madison Capstone in Microbiology Students; Medina
Muñoz M, Lysne JA, Dhokiya V, Hoque AF, Brackney DE, Caragata EP, Hutchinson ML, Jacobs-
Lorena M, Lampe DJ, Martin E, Valiente Moro C, Povelones M, Short SM, Steven B, Xu J,
Paustian TD, Rondon MR, Hughes GL, Coon KL, Heinz E. MosAIC: An annotated collection of
mosquito-associated bacteria with high-quality genome assemblies. PLoS Biol. 2024;
22:e3002897.
20. Ramirez JL, Short SM, Bahia AC, Saraiva RG, Dong Y, Kang S, Tripathi A, Mlambo G,
Dimopoulos G. Chromobacterium Csp_P reduces malaria and dengue infection in vector
mosquitoes and has entomopathogenic and in vitro anti-pathogen activities. PLoS Pathog. 2014;
10:e1004398.
21. Pumpuni CB, Beier MS, Nataro JP, Guers LD, Davis JR. Plasmodium falciparum: inhibition of
sporogonic development in Anopheles stephensi by gram-negative bacteria. Exp Parasitol. 1993;
77:195-9.
22. Mourya DT, Gokhale MD, Pidiyar V, Barde PV, Patole M, Mishra AC, Shouche Y. Study of the
effect of the midgut bacterial flora of Culex quinquefasciatus on the susceptibility of mosquitoes
to Japanese encephalitis virus. Acta Virol. 2002; 46:257-60.
23. Cirimotich CM, Dong Y, Clayton AM, Sandiford SL, Souza-Neto JA, Mulenga M, Dimopoulos G.
Natural microbe-mediated refractoriness to Plasmodium infection in Anopheles gambiae. Science.
2011;332:855-8.
24. Apte-Deshpande A, Paingankar M, Gokhale MD, Deobagkar DN. Serratia odorifera a midgut
inhabitant of Aedes aegypti mosquito enhances its susceptibility to dengue-2 virus. PLoS One.
2012; 7:e40401.
25. Sun X, Wang Y, Yuan F, Zhang Y, Kang X, Sun J, Wang P, Lu T, Sae Wang F, Gu J, Wang J, Xia
Q, Zheng A, Zou Z. Gut symbiont-derived sphingosine modulates vector competence in Aedes
mosquitoes. Nat Commun. 2024; 15:8221.
26. Garza-Hernández JA, Rodríguez-Pérez MA, Salazar MI, Russell TL, Adeleke MA, de Luna-
Santillana Ede J, Reyes-Villanueva F. Vectorial capacity of Aedes aegypti for dengue virus type 2
is reduced with co-infection of Metarhizium anisopliae. PLoS Negl Trop Dis. 2013; 7:e2013.
27. Dutra HL, Rocha MN, Dias FB, Mansur SB, Caragata EP, Moreira LA. Wolbachia Blocks
Currently Circulating Zika Virus Isolates in Brazilian Aedes aegypti Mosquitoes. Cell Host
Microbe. 2016; 19:771-4.

28. Hugo LE, Rašić G, Maynard AJ, Ambrose L, Liddington C, Thomas CJE, Nath NS, Graham M,
Winterford C, Wimalasiri-Yapa BMCR, Xi Z, Beebe NW, Devine GJ. Wolbachia wAlbB inhibit
dengue and Zika infection in the mosquito Aedes aegypti with an Australian background. PLoS
Negl Trop Dis. 2022; 16:e0010786.
29. Duguma D, Hall MW, Rugman-Jones P, Stouthamer R, Terenius O, Neufeld JD, Walton WE.
Developmental succession of the microbiome of Culex mosquitoes. BMC Microbiol. 2015;
24;15:140.
30. Kämpfer P, Lindh JM, Terenius O, Haghdoost S, Falsen E, Busse HJ, Faye I. Thorsellia anophelis
gen. nov., sp. nov., a new member of the Gammaproteobacteria. Int J Syst Evol Microbiol. 2006;
56(Pt 2):335-338.
31. Fu X, Dou J, Mao J, Su H, Jiao W, Zhang L, Hu X, Huang X, Wang S, Bao Z. RADtyping: an
integrated package for accurate de novo codominant and dominant RAD genotyping in mapping
populations. PLoS One. 2013; 8:e79960.
32. Callahan BJ, McMurdie PJ, Rosen MJ, Han AW, Johnson AJ, Holmes SP. DADA2: High-
resolution sample inference from Illumina amplicon data. Nat Methods. 2016; 13:581-3.
33. Bokulich NA, Kaehler BD, Rideout JR, Dillon M, Bolyen E, Knight R, et al. Optimizing
taxonomic classification of marker-gene amplicon sequences with QIIME 2 ' s q2-feature-
classifier plugin. Microbiome. 2018; 6:17.
34. Bolyen E, Rideout JR, Dillon MR, Bokulich N, Abnet CC, Al-Ghalith GA, et al. Reproducible,
interactive, scalable and extensible microbiome data science using QIIME 2. Nat Biotechnol.
2019; 37:852-7.
35. Lu Y, Zhou G, Ewald J, Pang Z, Shiri T, Xia J. MicrobiomeAnalyst 2.0: comprehensive statistical,
functional and integrative analysis of microbiome data. Nucleic Acids Res. 2023; 51:W310-W318.
36. Zhou W, Rousset F, O'Neil S. Phylogeny and PCR-based classification of Wolbachia strains using
wsp gene sequences. Proc Biol Sci. 1998; 265:509-15.
37. Rasgon JL, Cornel AJ, Scott TW. Evolutionary history of a mosquito endosymbiont revealed
through mitochondrial hitchhiking. Proc Biol Sci. 2006; 273:1603-11.
38. Rasgon JL, Scott TW. An initial survey for Wolbachia (Rickettsiales: Rickettsiaceae) infections in
selected California mosquitoes (Diptera: Culicidae). J Med Entomol. 2004; 41:255-7.
39. Rasgon JL, Scott TW. Wolbachia and cytoplasmic incompatibility in the California Culex pipiens
mosquito species complex: parameter estimates and infection dynamics in natural populations.
Genetics. 2003; 165:2029-38.
40. Chavshin AR, Oshaghi MA, Vatandoost H, Pourmand MR, Raeisi A, Terenius O. Isolation and
identification of culturable bacteria from wild Anopheles culicifacies, a first step in a
paratransgenesis approach. Parasit Vectors. 2014; 4;7:419.
41. Taha MME, Abdelwahab SI, Mohamed HY, Jerah A, Alabsi AM, Abdullah SM, Oraibi B, Alfaifi
HA, Babiker YOH, Aziz Ibrahim IA, Alshahrani S, Farasani AM, Alamer AS, Altherwi T.
Comprehensive review of Wolbachia research (1936-2024): Global landscape, mapping progress
and themes. Parasite Epidemiol Control. 2025; 30:e00438.
42. Anders KL, Indriani C, Ahmad RA, Tantowijoyo W, Arguni E, Andari B, Jewell NP, Dufault SM,
Ryan PA, Tanamas SK, Rancès E, O'Neill SL, Simmons CP, Utarini A. Update to the AWED
(Applying Wolbachia to Eliminate Dengue) trial study protocol: a cluster randomised controlled
trial in Yogyakarta, Indonesia. Trials. 2020; 21:429.

43. Ribeiro Dos Santos G, Durovni B, Saraceni V, Souza Riback TI, Pinto SB, Anders KL, Moreira
LA, Salje H. Estimating the effect of the wMel release programme on the incidence of dengue
and chikungunya in Rio de Janeiro, Brazil: a spatiotemporal modelling study. Lancet Infect Dis.
2022; 22:1587-1595.
44. Dodson BL, Hughes GL, Paul O, Matacchiero AC, Kramer LD, Rasgon JL. Wolbachia enhances
West Nile virus (WNV) infection in the mosquito Culex tarsalis. PLoS Negl Trop Dis. 2014;
8:e2965.
45. Coon KL, Vogel KJ, Brown MR, Strand MR. Mosquitoes rely on their gut microbiota for
development. Mol Ecol. 2014; 23:2727-39.

Figure legends

Fig. 1. Map of collection locations for mosquitoes assayed for PoolRADSeq analysis. Red = 2021; Blue = 2022.

Fig. 2. Map of collection locations for mosquitoes assayed for microbiome analysis. Colors indicate States.

Fig. 3. Neighbor-joining tree generated from PoolRADSeq analysis. Clusters previously identified by Venkatesan and colleagues (microsatellites) [1] and Liao and colleagues (individual RADSeq) [6] are highlighted.

Fig. 4. Bacterial communities in *Culex tarsalis* mosquitoes. **A:** Percent relative abundance of the top 10 identified taxa. **B:** Heat tree showing hierarchical structure of representative taxa. Taxa with 50% prevalence (% ASV detected in all samples) or higher are presented (family names are not labeled for better visibility). Genera within the top 10 relative abundance (see Fig. 1A) are in red text. Color and node indicate prevalence and number of ASV, respectively. See Additional file 1 for heat tree with all taxa represented.

Fig. 5. Fungal communities in *Culex tarsalis* mosquitoes. **A:** Percent relative abundance of the top 10 taxa. **B:** Heat tree showing hierarchical structure of representative taxa. Taxa with 50% prevalence or higher are labeled. Genera within the top 10 relative abundance (see Fig 2 A) are in red text. Color and node size indicate prevalence (% sequence read detected in all samples) and number of ASV, respectively. See Additional file 1 for heat tree with all taxa presented.

Fig. 6. Hypothesized clustering pattern of mosquito populations based on community structure of **A:** bacteria, **B:** fungi, and **C:** bacteria/fungi combined at the population (left) or group level (right). Numbers within each bar graph indicate the number of mosquito samples assayed.

Fig. 7. Alpha and Beta diversity analysis. **A:** Chao1 (species richness) and Shannon index (species diversity) were compared for Alpha diversity analysis. Pair-wise Mann-Witney tests were conducted between groups followed by Benjamini-Hochberg correction. **B:** Nonmetric Multidimensional Scaling (NMDS) with Bray-Curtis distance matric was estimated followed by Permutational Multivariate Analysis of Variance (PERMANOVA) for Beta diversity analysis.

Fig. 8. Random Forest classification analysis with **A:** representative tree plots, and **B:** top 15 taxa at genus level that had greatest influence on classification models. Mean OOB (Out Of Bag) errors were calculated using five consecutively generated OOB tables (see Additional file 7) (*,*Burkholderia, Caballeronia, Paraburkholderia*).

Fig. 9. Core microbiome analysis. Taxa with 20% or higher prevalence are presented at genus level. The detection threshold was set at 0.01% relative abundance (*, *Burkholderia, Caballeronia, Paraburkholderia*).

Fig. 10. Correlation analyses between environmental factors and species richness/diversity or geographic structure. **A:** Correlation between abiotic environmental factors and Alpha diversity indices using linear regression. Positive and negative numbers indicate positive (red) and negative (blue) correlation, respectively (***, $P < 0.001$; **, $P < 0.01$; *, $P < 0.05$). **B:** Correlation between geographic distance and Bray-Curtis dissimilarity index using Mantel test. Line plots with shaded areas indicate linear regression with 95% CI.

Table 1. Collection information for PoolRADSeq samples

Collection year	Sample ID	Location	State	Latitude	Longitude
2021	P1_2021	Lake point	UT	40.669406	-112.305263
2021	P2_2021	Stansbury Park	UT	40.651018	-112.299081
2021	P3_2021	Stansbury Park	UT	40.647411	-112.314970
2021	P4_2021	Yakima/Grandview	WA	46.193421	-119.899400
2021	P5_2021	Yakima/Grandview	WA	46.224977	-119.901053
2021	P6_2021	Yakima/Grandview	WA	46.205611	-119.892026
2021	P7_2021	Lubbock	TX	33.462700	-101.891400
2021	P9_2021	Loveland	CO	40.403933	-105.071421
2021	P10_2021	Loveland	CO	40.401394	-105.108170
2021	P12_2021	Cottonwood, Shasta County	CA	40.414286	-122.216945
2021	P13_2021	Anderson, Shasta County	CA	40.445232	-122.243092
2021	P14_2021	Cottonwood, Shasta County	CA	40.396727	-122.281994
2021	P15_2021	Yolo County	CA	38.610336	-121.711942
2021	P16_2021	Sacramento County	CA	38.107311	-121.649758
2021	P17_2021	Yolo County	CA	38.810503	-121.808285
2022	Z1	Riverside county, Mecca	CA	33.53271	-116.03503
2022	Z2	Riverside county, Thermal	CA	33.46238	-116.06125
2022	Z3	Yolo county, Conaway Ranch	CA	38.64849	-121.71102
2022	Z4	Yolo county, Vic Fazio	CA	38.56179	-121.62924
2022	Z5	Adams County	CO	39.90015	-104.91565
2022	Z6	Broomfield	CO	39.89499	-105.13878
2022	Z7	Larimer county, Loveland	CO	40.43432	-105.10429
2022	Z8	Larimer county, Fort Collins	CO	40.48614	-104.97362
2022	Z9	Mapleton	ND	46.89102	-97.04281
2022	Z10	Mapleton	ND	46.88508	-97.06925
2022	Z11	Grand Forks county, Grand Forks	ND	46.94475	-97.06344
2022	Z12	Grand Forks county, Grand Forks	ND	47.88686	-97.09869
2022	Z13	Cascade county, Sunriver	MT	47.51992	-111.77331
2022	Z14	Cascade, Vaughan	MT	47.53991	-111.63364
2022	Z15	Yakima county, Grandview	WA	46.22498	-119.90105
2022	Z16	Yakima county, Mabton	WA	46.20787	-119.99583
2022	Z17	Lubbock, Lubbock	TX	33.58318	-101.89314
2022	Z18	Lubbock, Lubbock	TX	33.58254	-101.82794

694 **Table 2.** Collection information for microbiome samples

Population ID	State	Number of samples		City/County	Latitude	Longitude
		16S	ITS1			
CA1	CA	17	0	Cottonwood, Shasta County	40.41429	-122.21695
CA2		19	2	Anderson, Shasta County	40.44523	-122.24309
CA3		7	1	Cottonwood, Shasta County	40.39673	-122.28199
CA4		14	8	Yolo County	38.61034	-121.71194
CA5		10	7	Sacramento County	38.10731	-121.64976
CA6		10	11	Yolo County	38.8105	-121.80829
CO1	CO	15	3	Loveland	40.40393	-105.07142
CO2		16	4	Loveland	40.40139	-105.10817
CO3		3	0	Fort Collins	40.52993	-105.05145
TX1	TX	8	8	Lubbock	33.4627	-101.8914
TX2		3	0	Lubbock	33.59046	-101.93866
UT1	UT	14	11	Lake point	40.66941	-112.30526
UT2		13	4	Stansbury Park	40.65102	-112.29908
UT3		13	1	Stansbury Park	40.64741	-112.31497
WA1	WA	16	6	Yakima/Grandview	46.19342	-119.8994
WA2		10	2	Yakima/Grandview	46.22498	-119.90105
WA3		9	4	Yakima/Grandview	46.20561	-119.89203

Figure 1

Figure 2

Figure 3

Fig 4

Fig 5

Fig 6

Fig 7

Fig 8

Fig 9

Fig 10