Discrimination of Annonaceae using herbarium leaf reflectance

spectra under limited sample size conditions

Khalil Boughalmi[1], Paola G. Santacruz Endara[2], Lily A. Bennett[3] , Martin Ecarnot[4], Samantha

Bazan[5], Denis Bastianelli[5], Laurent Bonnal[5], Thomas L.P. Couvreur[1]

1 : DIADE, Univ. Montpellier, CIRAD, IRD, Montpellier, France

2: Grupo de investigación Biodiversidad y Sociedad BioS, Jardín Botánico Padre Julio Marrero,

PUCESD, Santo Domingo de los Tsáchilas, Ecuador

3: School of Natural Sciences, University of Tasmania, Hobart, Tasmania, Australia

4: AGAP, INRAE, Univ. Montpellier, CIRAD, Institut Agro, Montpellier, France

5: SELMET, Univ. Montpellier, CIRAD, INRAE, Institut Agro, F-34398 Montpellier, France

Corresponding author : Khalil Boughalmi; boughalmi.khalil@gmail.com

19 ABSTRACT

Premise : Herbarium collections offer an unparalleled archive of plant biodiversity, but their

use for species identification through spectral data remains constrained by uncertain effects of

preservation histories. This study assesses whether barium specimens can reliably predict

species based on its leaf reflectance spectrum, despite variations in age, geographic origin, or

conservation method under limited sample size conditions.

Methods : We scanned herbarium specimens of different ages and geographic distribution of

14 species of the pantropical Annonaceae. In addition, we used a second dataset of 9 species

where some specimens were conserved in alcohol prior to drying and some not. We used five

supervised classification models frequently used for high-dimensional data such as

spectroscopy.

Results : All models achieved high accuracy (>80%) when trained on multiple specimens per

species. However, when using only one specimen per species, accuracy varied substantially

depending on the taxon.

Discussion : Our findings demonstrate that herbarium specimens often retain a strong

taxonomic signal in their spectra, however, inter-individual variability affects accuracy in

some taxa. These findings confirm the usefulness of herbarium spectroscopy as a non-

destructive tool for species identification and offer a promising avenue for digitizing historical

biodiversity data into high-dimensional trait space.

Keywords
Support Vector Machine; Linear Discriminant Analysis; Species identification; Preservation;

Signature spectral signal

INTRODUCTION

Herbaria hold hundreds of millions of preserved plant specimens, forming one of the largest

and oldest repositories of biodiversity worldwide (Davis, 2023). These collections span more

than four centuries of global plant exploration and constitute irreplaceable archives of

taxonomic, geographic, and temporal information. Originally designed for taxonomy and

floristics, herbaria have long been the foundation of plant classification and nomenclature.

Yet, in recent decades, these collections are increasingly being reinterpreted as data-rich

resources for ecology, evolution, and conservation biology (Besnard et al., 2018; Raxworthy

and Smith, 2021). Herbarium specimens collected decades or even centuries ago are now used

in trait-based ecology, functional biogeography, phylogenetics, and global change research

(Nualart et al., 2017; Lang et al., 2019; Rosche et al., 2025), revealing new value in what was

once considered passive documentation.

This shift has been catalyzed by the development of analytical tools capable of extracting

meaningful data from historical plant material (Raxworthy and Smith, 2021; Davis, 2023).

Among these, leaf reflectance, particularly near-infrared spectroscopy (NIRS), has emerged as

a powerful and non-destructive technique to characterize plant functional traits (Cavender-

Bares et al., 2017). Leaf reflectance spectra are a promising approach to species

identification and even predicting functional traits (Kothari and Schweiger, 2022; Kothari et

al., 2023) because they rapidly produce a snapshot of leaf chemical and structural

composition. In such analyses, reflectance measurements typically span wavelengths from

350 to 2500 nm, with 400–700 nm corresponding to the visible range, 700–1100 nm to the

near-infrared (NIR), and 1100–2500 nm to the shortwave infrared (SWIR). Reflectance is

influenced by biochemical constituents such as cellulose, lignin, water, phenolics, and

proteins, as well as by structural characteristics like leaf thickness or the presence of surface

features such as hairs and stomata (Jacquemoud and Ustin, 2019).

Spectra therefore have the potential to serve as a rapid non-desctructive tool for plant species

identification, as they may contain taxonomically informative features, sometimes referred to

as the Species Spectral Signature (SSS) (Durgante et al., 2013). This taxonomic signal has led

to a growing number of studies using reflectance spectra to discriminate between plant

genera, species, and even populations. For example, Durgante et al. (2013) demonstrated that

leaf reflectance spectra could successfully classify tropical tree species, while Meireles et al.

(2020) revealed a phylogenetic component embedded in spectral variation at the angiosperm

level. Other studies have confirmed the potential of NIRS for identifying hybrids, assessing

intraspecific variability, and monitoring trait shifts across environments (Cavender-Bares et

al., 2017; Stasinski et al., 2021).

Based on these observations, leaf reflectance spectroscopy has recently emerged as a

promising tool for botanists, ecologists, and taxonomists alike. Furthermore, its non-invasive

nature makes it particularly attractive for use on valuable or rare specimens, including

historical herbarium material (types) including extinct species.

However, there is a critical caveat to this optimism. Most studies to date have relied on

freshly collected, purpose-pressed leaves, prepared under relatively standardized and

controlled conditions (referred to here as "pressed specimens") (Durgante et al., 2013;

Meireles et al., 2020; Kothari et al., 2023). Pressed specimens are not equivalent to herbarium

specimens as they have not undergone mounting, long-term storage, transportation, or

historical treatment. Herbarium specimens have often been exposed to a wide variety of in-

field drying methods, conservation practices, and chemical treatments, many of which are

undocumented, and all of which can potentially affect the chemical integrity of the leaves.

The effects of these processes on leaf chemistry (and by extension, on spectral properties) are

yet not well understood. This represents a fundamental limitation to the generalizability of

existing findings.

For example, in humid tropical environments, alcohol is frequently used as a temporary

preservative before heat drying, particularly during multi-day field expeditions (Hodge,

1947). While effective at preventing mold and tissue decay, alcohol is a strong solvent and

oxidizer that can also impact DNA extraction (Forrest et al., 2019). Even today, practices such

as applying glue, or drying methods vary considerably between herbaria and collectors. As a

potential source of chemical modification, all these factors can introduce noise or bias into

spectral data collected from herbarium sheets.

Moreover, the geographic origin of specimens introduces an additional layer of variation that

reflects not only environmental conditions but also underlying genetic and phenotypic

diversity within species. Climate, elevation, soil chemistry, and other environmental

parameters influence leaf chemistry (Maire et al., 2015; Asner and Martin, 2016; Cao et al.,

2020), while genetic variation, phenological stage, and ontogenetic differences can further

contribute to spectral heterogeneity among specimens grouped under a single species

definition. As a result, specimens of the same species, preserved differently and collected

across distinct locations or developmental stages, may display substantial spectral divergence.

If spectroscopy is to be used reliably for taxonomic or ecological inference on herbarium

material, these multiple sources of variation must be understood and, where possible,

accounted for within the analytical framework.

Despite these challenges, recent evidence suggests that herbarium specimens still retain

valuable chemical and even strong taxonomic information (Neto-Bradley et al., 2025; White

et al., 2025b). For instance, Foutami et al. (2018) have shown that the age of a specimen does

not drastically affect some part of its chemical profile, even after decades of storage. While

certain surface properties or volatile compounds may degrade over time, the core chemical

constituents that drive spectral variation often remain stable (Mendes Resende et al., 2020).

The key challenge is not whether herbarium spectra contain useful information, but whether

this information is consistent, reproducible, and interpretable enough to enable species-level

identification and classification, while remaining feasible at low sampling scales given the

frequent scarcity of specimens per species and the logistical demands of projects involving

hundreds or thousands of samples.

Here, we investigate several key questions regarding the application of spectroscopy to

herbarium specimens. Specifically: (i) Can spectral data reliably support species-level

identification? (ii) How many specimens per species are required to adequately represent

intra-specific spectral variability? (iii) Are standard machine learning algorithms sufficiently

robust to handle the heterogeneity typical of herbarium collections? and (iv) To what extent

do pre-drying alcohol treatments affect the spectral signal used for classification? By

addressing these questions, we aim to assess both the practical applicability and

methodological limitations of herbarium spectroscopy under realistic working conditions.

To address the above questions, we focused on the pantropical plant family Annonaceae. With

around 2500 species, it is one of the best-known tropical plant families in terms of taxonomy

with global occurrence data and phylogenetic resources available (Couvreur et al., 2012,

2019; Nge et al., 2024). Despite this taxonomic maturity, species identification in Annonaceae

can be challenging, particularly when reproductive structures (flowers or fruits) are absent or

unavailable. This makes Annonaceae an ideal case study for evaluating the robustness of

spectral approaches for species discrimination. Additionally, the family displays a high degree

of ecological and biogeographic amplitude within the tropics, spanning environments from

Amazonia to African rain forests and Southeast Asian regions, thus offering a natural gradient

of variation in both environmental conditions and specimen collection practices (Pennington

et al., 2004; Couvreur et al., 2011; Chatrou et al., 2012).

142
METHODS
143
144
Taxon sampling

We generated two datasets. First, to address the objective of species identification using

herbarium-generated spectra, we scanned leaves from 14 Annonaceae species (**Table 1,**

Supplementary Table 1) all stored in the herbarium of the Muséum national d'Histoire

naturelle in Paris (P) (Le Bras et al., 2017). As is common for historical herbarium material,

detailed information on pre-drying treatments or conservation practices was generally

unavailable for these specimens, although such treatments are likely to have occurred. For

each species we scanned a minimum of 9 specimens. The selected species were: 1) chosen

from recently revised genera, ensuring expert-based identification; 2) represent different

growth habits (liana, understory and canopy trees); 3) have a broad geographical distribution,

including species from South America, Africa, Madagascar, and Southeast Asia; 4) have

diverse geographical representation within species, including specimens from various

countries and ecotypes when possible; and 5) are phylogenetically diverse across Annonaceae

(at least one species for each of the four subfamilies, Nge et al. 2024). We also included two

species pairs belonging to the same genera in *Ambavia* and *Hexalobus* (**Table 1,**

Supplementary Table 1). In general, we prioritized flowering or fruiting specimens as these

are better identified. This dataset is referred to as ID_PARIS.

A second independent dataset was generated to examine the impact of alcohol preservation.

For this we sampled nine species (**Table 2**) within the 50-ha Yasuní Forest Dynamics Plot

(YFDP) located next to the Yasuní Research Station of the Pontificia Universidad Católica del

Ecuador (Valencia et al., 2004). These species are the same as the study of Santacruz Endara

et al. (2025) and cover the phylogenetic diversity of the family. First, specimens were

sampled directly in the field at the Yasuní Research station of the Pontificia Universidad

Católica del Ecuador (PUCE). Leaves were then pressed in newspaper (paper-flattened) and

dried over an electric stove for 24 hours. For one or two specimens per species they were

sprayed in 70-80% alcohol for three days, and then dried in an electric stove for 24 hours.

This dataset is referred to as ID_YASUNÍ.

To place the two datasets into context, **Figure 1** summarizes the overall workflow, from

specimen selection and scanning procedure acquisition to preprocessing, providing a

framework for the methodological details that follow.

Spectral Measurements

Leaf specimens were scanned following a general protocol (Mersni et al., 2025), with dataset-

specific adaptations (**Table 3**). In both datasets (ID_PARIS and ID_YASUNÍ), spectra were

collected in the 350–2500 nm wavelength range using ASD LabSpec® instruments, with

differences in probe configuration and acquisition setup between datasets. ID_PARIS

measurements were acquired using a Rapid Analysis Probe coupled with a fiber optic

illuminator, whereas ID_YASUNÍ spectra were collected using a 6 mm diameter pen probe.

Spectra were generated by averaging multiple readings per scan (20 readings in ID_PARIS

and 30 in ID_YASUNÍ), with 20 scans per specimen in ID_PARIS and 50 scans per specimen

in ID_YASUNÍ. White reference and calibration procedures differed slightly between datasets

(see **Table 3** for details). Measurements were performed by a single operator in ID_PARIS

and by multiple operators in ID_YASUNÍ, but the same Spectral Black™ strip background

was used.

In both datasets, spectra were collected on the adaxial side of mature leaves under controlled

environmental conditions. Young or unfolding leaves were avoided, as well as the midrib

region and areas affected by fungi, bryophytes, or glue. Whenever possible, multiple leaves

per specimen were scanned, and measurements were distributed across the cleanest and least

damaged areas. Care was taken to ensure contact between the probe and the leaf without

applying excessive pressure that could damage the specimen.

Data Processing
All data processing and analyses were conducted using R software (v4.4.1). The processing of

spectral data ensures consistency, removes noise, and enhances relevant signals for further

analysis and is composed of two steps. The first step involved applying a Standard Normal

Variate (SNV) normalization (Barnes et al., 1989). For each spectrum, the mean and standard

deviation of the 20 readings are calculated. Each reflectance intensity value is then

normalized by subtracting the mean and dividing by the standard deviation, resulting in

spectra with a mean of zero and unit variance. This standardization method removes a part of

noise caused by external factors such as instrument settings, distance between the probe and

the sample, or sample thickness. After normalization, the Savitzky-Golay filter (Savitzky and

Golay, 1964) was used to compute the first derivative of the spectra. A polynomial order of 2

and a window size of 7 were used to smooth the data while capturing the rate of change at

each wavelength. The first derivative accounts for variations in the slope of the spectra.

Analytical Approaches

Five classification models were implemented to evaluate taxonomic discrimination: Partial

Least Squares Discriminant Analysis (PLS-DA), Support Vector Machine (SVM), K-Nearest

Neighbors (KNN), Linear Discriminant Analysis (LDA), and Random Forest (RF). Each

model was selected for its already known capacity to handle spectral data sets (Kong et al.,

2013; Trainor et al., 2017; Jombo et al., 2020; Wang et al., 2020). PLS-DA is widely used in

chemometric and botanical applications due to its ability to model multicollinearity and

extract latent components that maximize class separation (Richter et al., 2016; Song et al.,

2018; Meireles et al., 2020; Kothari et al., 2023). LDA, while more limited in its assumptions,

provides a simple and interpretable baseline model. Random Forest is a robust ensemble

method capable of capturing non-linear relationships (Brabant et al., 2019). KNN offers a

non-parametric approach that leverages local similarity in the spectral space and can be

sensitive to subtle differences between specimens (Kong et al., 2013). Lastly, SVM has

demonstrated strong performance in high-dimensional classification problems, particularly

with spectral data, due to its capacity to define optimal decision boundaries in complex

feature spaces, even when class separation is not linearly achievable (Brabant et al., 2019;

Jombo et al., 2020). All models were trained and tested using stratified resampling with a

strict separation by specimen identity to avoid data leakage, defined here as the presence of

training or calibration specimens, or their associated spectra, in the test dataset. Specimen-

level stratified sampling by species was applied in each iteration. For each species, six

specimens were randomly assigned to a tuning subset. From the remaining specimens,

approximately two-thirds were allocated to the calibration subset, and the remaining

specimens formed the independent prediction subset. For PLS-DA, models were fitted using

the plsda() function from the mixOmics package (Rohart et al., 2017). The number of

components was optimized within each iteration by cross-validation on a dedicated tuning

subset of the ID_PARIS dataset, testing values between 2 and 25. Hyperparameter

optimization was conducted using five-fold cross-validation within the tuning subset

(approximately 80% of specimens used for model fitting and 20% for internal validation at

each fold). The optimal number of components was selected as the one minimizing cross-

validated error rates. The SVM model was implemented using the caret interface to the e1071

(Meyer et al., 1999) and kernlab packages (Karatzoglou et al., 2024). Hyperparameter tuning

was performed on the ID_PARIS dataset via cross-validation on a tuning subset, with a grid

of sigma values (0.001, 0.01, 0.1) and cost values (0.1, 1, 10). Five-fold cross-validation was

used within the tuning subset for hyperparameter selection. The best combination of sigma

and C was retained for final calibration and testing. All predictor variables were standardized

based on calibration data. For KNN, the knn3() function from the caret package (Kuhn et al.,

2024) was used. The number of neighbors was tuned on the ID_PARIS dataset using cross-

validation over k values from 1 to 10. Hyperparameter tuning was conducted using five-fold

cross-validation within the tuning subset. Predictors were standardized according to

calibration data, and the same transformation was applied to tuning and test sets. LDA was

performed using the lda() function from the MASS package (Ripley et al., 2025). As LDA

assumes uncorrelated predictors, a principal component analysis (PCA) was applied to the

calibration data. The first set of components explaining at least 95% of the total variance were

retained, and the same transformation was applied to tuning and test sets before fitting the

LDA model. For Random Forest, the randomForest() function from the randomForest

package (Breiman et al., 2024) was used with 500 trees (ntree = 500). All predictors with non-

zero variance were included, and no additional feature selection or scaling was performed.

Classification was based on majority vote across the ensemble of trees.

After hyperparameters were selected using the tuning subset, final models were trained

exclusively on the independent calibration subset using the optimized parameters. The tuning

subset was not reintegrated into the calibration phase, thereby maintaining strict separation

between hyperparameter optimization, model fitting, and final evaluation. Model performance

was then assessed on the independent prediction subset.

The models were applied to two distinct datasets: ID_PARIS, which was used for model

training and testing with herbarium data, and ID_YASUNÍ, which was used to assess the

effect of alcohol preservation on species discrimination. For ID_PARIS, the dataset was split

into tuning, calibration, and prediction subsets, enabling full hyperparameter optimization. In

contrast, analyses on ID_YASUNÍ and additional analyses on ID_PARIS at varying

calibration sizes were conducted in a robustness-testing framework, where fixed

hyperparameters were reused across all conditions. This ensured comparability across datasets

and specimen sampling levels.

To simulate real-world conditions where the preservation method is generally unknown,

alcohol-preserved specimens were randomly included in either the training or test dataset.

This allowed us to evaluate how alcohol preservation might affect the models' ability to

distinguish between species. For each iteration, classification accuracy, or number of correct

predictions divided by the total number of predictions, was calculated, and confusion matrices

were pooled across resamples to evaluate model performance. To avoid bias, species were

equally represented in all calibration datasets, irrespective of specimen availability. This

balanced sampling ensured that species with more specimens did not dominate the models.

Model predictions were compared to expert identifications to assess accuracy. Overall

performance was computed for each model, and confusion matrices were used to highlight

misclassification trends. We tested the models using both all available spectra per specimen

(twenty spectra) and their average. To further explore the limitations of using a single

herbarium specimen to represent a species, we conducted an analysis using the ID_PARIS

dataset. We first evaluated model performance with only one randomly selected specimen per

species, using its twenty spectra for training and the spectra of all remaining specimens for

testing. We then extended the analysis by progressively increasing the number of training

specimens from one to eight, using all their spectra for training and the spectra of the

remaining specimens for testing.

Finally, to ensure robustness and consistency, all analyses were validated using 100 randomly

generated iterations. This repeated validation allowed us to assess the stability of model

performance across different sampling conditions and to ensure reliable results.

RESULTS

Specimen spectrum sampling

A total of fourteen species were scanned in the P herbarium. Using the previously described

workflow (**Fig. 1**), we were able to scan up to 18 specimens for some species. Preparing the

specimen, entering the required parameters, and initiating the first scan takes approximately 8

minutes. Once this setup is completed, acquiring the remaining scans is rapid, so that

collecting a total of 20 spectra requires about 10 minutes per specimen. This enabled us to

scan 160 specimens over 5 days (see **Table** 1 & **Table S1** for details). The scanned specimens

ranged in age from 9 to 201 years, with an overall average of 69 years. Some species include

older specimens (e.g., *Annona senegalensis*: mean 107 years, SD = 23; *Uvaria grandiflora*:

mean 89 years, SD = 53; *Anaxagorea dolichocarpa*: mean 88 years, SD = 64), while others

include younger ones (e.g., *Xylopia humblotiana*: mean 30 years, SD = 22; *Ambavia spp.*:

mean 33 years, SD = 15) (**Fig. S1**). Several issues arose during sampling due to specific

conditions encountered with some specimens, such as glue or highly curled leaves. Among

the 160 specimens, less than five were affected by the scanning process, and the impacts were

never severe and generally consist of leaf tears on curled specimens and leaf detachment. In

the case of the samples from Yasuní, the scanning workflow was much more efficient because

the data had already been collected and the leaves were not fixed in an herbarium voucher.

Despite taking more scans per specimen (50 instead of 20), it took less than five minutes to

measure a whole specimen.

Spectrum discrimination using herbarium specimen

For all five classification models tested, we observed an overall strong capacity to

discriminate herbarium specimens at the species level. Using all spectra from training

specimens species and all spectra (measurement-level) from the remaining specimens for

testing, we obtained the following overall accuracies: 84.74% (SD = 5.23%) for LDA,

81.45% (SD = 4.50%) for SVM, 81.06% (SD = 4.47%) for RF, 78.57% (SD = 4.77%) for

PLS-DA, and 75.63% (SD = 4.08%) for KNN (**Fig. 2A**). When each specimen in the test set

was represented by a single averaged spectrum rather than the full set of 20 individual spectra

(specimen-level), the discrimination capacity were even higher for all models: 87.48% (SD =

5.57%) for LDA, 86.35% (SD = 4.47%) for SVM, 84.96% (SD = 5.60%) for RF, 83.03% (SD

= 5.82%) for PLS-DA, and 79.23% (SD = 4.86%) for KNN(**Fig. 2A**). This remained true

independent of species diversity in terms of geographic origin or collection date (**Fig. 2B**).

However, two species from the same genus, *Hexalobus crispiflorus* and *H. monopetalus*,

appeared to be less accurately classified, with misclassifications from one to the other. In

most models using individual spectra as the test, there was a tendency for spectra to be

classified as the other species, with a slight bias toward misclassification in favor of *H.*

crispiflorus. When using the average spectrum per specimen in the test set, this pattern

persisted, where *H. monopetalus* was more frequently predicted as *H. crispiflorus* than the

reverse (**Fig. 2B**). This trend was especially pronounced in LDA, where nearly all *H.*

crispiflorus specimens were correctly classified (79.50%) but only 66.67% of *H.*

monopetalus, the misclassified ones were all predicted as *H. crispiflorus*. The two *Ambavia*

species showed a better discrimination rate without a distinct four-square misclassification

pattern. However, there was a slight but consistent tendency to misclassify *A.capuronii* as *A.*

gerrardii. *Uvaria grandiflora* also appeared to be less well classified, where its correct

classification rate was among the lowest observed. Overall, despite a general increase in

model accuracy when using the average spectrum, the same species tended to be misclassified

across both approaches and generally in the same predicted species (**Fig. 2B**).

Alcohol treatment effect

All models showed an accuracy above 90% when every spectrum was included in the test set,

and no model accuracy was below 95% when the species average spectrum was used in the

test set. Three models (SVM, PLS-DA and LDA) reached classification rates above 99%,

with very low standard deviation (SVM and PLS-DA <0.005; LDA <0.02 (**Fig. 2A**)) ,

indicating nearly all specimens were correctly classified to their species, regardless of whether

alcohol-treated specimens were included in the training or test datasets. We did not observe a

strong genus-level misclassification pattern, similar to what was seen in *Hexalobus*, among

the two *Guatteria* species included in this dataset especially with the average specimen in the

test dataset (**Fig. 2B**). Additionally, we found that alcohol-preserved specimens do not appear

to be distant in the 3D space defined by the first three PCA components of the average

spectrum for each specimen (**Fig. S2A**). These specimens cluster closely with others from the

same species and are never misclassified into the spectral space of a different species . The

average raw spectra themselves also generally share the same overall shape as those of other

non-alcohol-preserved specimens from the same species, while differences in magnitude can

be observed (**Fig. S2B**).

One specimen to represent them all

We observed substantial variability in classification accuracy depending on the species when

training the models with only a single specimen but using its 20 spectra (**Fig. 3**). For some

species, such as *Anaxagorea dolichocarpa*, *Duguettia calycina* or *Ambavia gerrardii*,

accuracy remained high more than 80% across all models (Mean values *A. dolichocarpa*

85.22; *A. gerrardii* 82.77; *D. calycina* 84.64 (**Fig. 3**)), while for others like *Uvaria*

grandiflora or both *Hexalobus* species, it dropped below 50% (Mean values *H. monopetalus*

49.33; *H. crispiflorus* 43.47; *U. grandiflora* 37.64 (**Fig. 3**)). Despite this variability, model

performance tended to follow similar patterns: species that were generally well classified

remained so across models, and the same held for poorly classified species. Nonetheless,

LDA generally outperformed the other methods, showing the highest classification accuracy

in 12 out of 14 species. In certain cases, this performance gap was particularly marked (e.g.

Monanthotaxis enghiana).

With an average accuracy of 65.4% (**Fig. 4A**), the models showed an important improvement,

reaching 78.2% with only two specimens used for training, and stabilizing near 90% from five

specimens onward, with only slight increases beyond that point (**Fig. 4B**). LDA consistently

outperformed the other models at all specimen counts, achieving over 70% accuracy with just

one specimen and reaching more than 90% from three specimens onward.

DISCUSSION

We demonstrate that herbarium leaf spectra contain sufficient information to enable accurate

species classification, even with small sample sizes and samples that are decades old. Across

most taxa tested, classification performance remained consistently high (**Fig. 2A**), supporting

the idea that herbarium material can preserve relevant traits for classification (Cavender-Bares

et al., 2017; Meireles et al., 2020; Kothari et al., 2023; Neto-Bradley et al., 2025; White et al.,

2025b). Importantly, high accuracy (often >90%) was achieved using as few as five

specimens per species for model training (**Fig. 4**). Even under single-specimen conditions,

performance far exceeded random expectations (**Fig. 3**), indicating that the spectral signature

(Durgante et al., 2013) of many species is both coherent and stable.

Classification success was not universal across taxa, and performance varied among species.

In the two instances of congeneric species included in this study, classification accuracy was

reduced compared to intergeneric discrimination, yet remained well above random

expectations. This was particularly evident for the two *Hexalobus* species, which showed

comparatively lower accuracy across models (**Fig. 2B**) but were still discriminated at rates

substantially higher than chance. These results support the idea that closely related species

present greater challenges for spectral discrimination, even when they are morphologically,

ecologically, and phylogenetically distinct (Botermans et al., 2011; Dagallier et al., 2023). For

example, *Hexalobus crispiflorus* and *H. monopetalus* occupy contrasting environments,

rainforest and savanna, respectively, yet their spectral signatures did not reflect this ecological

divergence. Such cases indicate that while reflectance spectroscopy is generally effective, it

may occasionally fail to capture biologically meaningful differences among certain closely

related species. In contrast, *Greenwayodendron suaveolens*, a widespread generalist tree

species in the Congo Basin (Lissambou et al., 2018), was robustly classified despite spanning

multiple countries and collection periods (**Fig. 3**), highlighting the potential of spectroscopy

to detect stable traits even under broad ecological and geographic variation. It is noteworthy

that misclassification was not restricted to congeneric species. *Uvaria grandiflora*, which was

the sole representative of its genus in the dataset, also showed reduced classification

performance, indicating that factors beyond taxonomic proximity, such as intraspecific

variability, specimen condition, or unobserved preservation artefacts, may contribute to

classification difficulty. An important consideration is that most species included in this study

belong to different genera, validating this approach at the higher taxonomic level. However,

species-level discrimination in more closely related species may require larger training sets.

We evaluated five classification models (PLS-DA, LDA, RF, KNN, and SVM), each selected

for its capacity to handle high-dimensional spectral data (Kong et al., 2013; Trainor et al.,

2017; Jombo et al., 2020; Wang et al., 2020). Although all models achieved strong

performance, they differed in robustness and scalability (**Fig. 2–4**). LDA emerged as the most

accurate method in this study, likely benefiting from the structured nature of the dataset and

prior dimensionality reduction. It is nevertheless worth noting that, despite its strong overall

performance, LDA also exhibited limitations when discriminating among congeneric species,

highlighting the increased difficulty of intra-generic classification. SVM also performed

consistently well, reflecting its strength in handling complex decision boundaries in high-

dimensional space. In contrast, RF and KNN showed greater variability, while PLS-DA,

although effective, is mathematically less suited to classification as the number of classes

increases (Trainor et al., 2017). Accuracy improved with increasing training set size but

plateaued beyond five specimens per species (**Fig. 4**), a promising result for specimen-poor

but species-rich groups.

We deliberately retained the full spectral range for model training, rather than automatically

excluding the first and last wavelength regions as commonly recommended (e.g. Meireles et

al., 2020), in order to explore the complete spectral signal present in herbarium specimens.

We also used spectra that, as shown in **Fig. S2** for ID_YASUNÍ and **Fig. S3** for ID_PARIS,

occasionally display abrupt jumps in reflectance related to instrument characteristics. In our

case, these artefacts did not appear to reduce model performance when using processed

spectra. Consequently, the classification performance reported here may represent a

conservative estimate and could potentially be further improved through targeted wavelength

selection or trimming, even though accuracy was already high.

In addition to overall classification performance, this study represents a first step toward

exploring the effect of alcohol preservation prior to drying on herbarium-based spectral

classification. Alcohol preservation is commonly used during specimen preparation,

particularly in humid tropical environments, where it helps prevent tissue degradation, mold

development, and other forms of contamination. Despite its widespread use, quantitative

information on the proportion of alcohol-preserved specimens in herbarium collections, as

well as on exposure duration and concentration, remains unavailable. It is nevertheless

reasonable to assume that a substantial fraction of tropical herbarium specimens has been

exposed to alcohol at some stage of preparation, in some cases for extended periods.

Alcohol preservation may influence dried herbarium specimens through both physical and

chemical mechanisms. Physically, alcohol can induce tissue shrinkage or structural alteration,

while chemically it is a well-known solvent for chlorophyll and other compounds contributing

to leaf optical properties. In our experimental setup, visual inspection of PCA space based on

averaged spectra indicates that alcohol-treated specimens generally occupy species-specific

regions overlapping with those of untreated specimens (**Fig. S2A**). Although slight shifts in

spectral magnitude were observed (**Fig. S2B**), the overall spectral shape and the main

taxonomic signal appeared to be preserved.

Consistent with these observations, species-level classification accuracy remained high when

alcohol-treated specimens were included in either the training or test datasets. Across all

models, accuracy exceeded 95%, with particularly strong performance for SVM, LDA, and

PLS-DA (**Fig. 2A**), indicating that alcohol-treated specimens retained sufficient spectral

information for accurate species discrimination in this experimental context. However, several

aspects of the design should be considered when interpreting these results. The alcohol dataset

comprised a relatively limited number of specimens and species, which likely facilitated

discrimination. In addition, classification success may partly reflect the behavior of

supervised learning algorithms, which can downweight spectral regions most affected by

chemical alteration and rely on more stable wavelength ranges for discrimination. Taken

together, these results suggest that under controlled, short-term exposure conditions and at a

limited taxonomic scale, alcohol-preserved herbarium specimens can retain a usable

taxonomic signal for spectral classification. Rather than providing a generalized assessment of

alcohol preservation effects, this experiment offers an initial indication that moderate

chemical alteration does not necessarily preclude the use of herbarium spectra in supervised

classification frameworks.

More broadly, pre-drying preservation methods, including those involving alcohol, vary

widely across herbarium collections and are often insufficiently documented. Differences in

exposure duration (i.e. several weeks), concentration, and storage conditions are likely to

produce a wide range of chemical alterations, some exceeding those captured by controlled

experiments. Addressing this variability will require dedicated reference datasets, either based

on specimens with well-documented preparation histories or purpose-designed experiments

explicitly testing preservation treatments. While recent studies have focused on specimen

aging and drying methods (Kühn et al., 2024), chemical treatments, aside from glue-related

effects (White et al., 2025b), remain comparatively understudied, despite their potential

influence on spectral integrity.

While these results are encouraging, our study was conducted under controlled conditions

using specimens from a single herbarium. We did not isolate the effects of specimen age or

geographic origin, reflecting the reality that such metadata are often incomplete or

unavailable. By working under realistic constraints, our goal was to assess model performance

in conditions commonly encountered in herbarium-based research, balancing methodological

rigor with operational feasibility.

From a methodological perspective, these challenges raise important questions regarding

specimen usability, potential exclusion thresholds, and analytical strategies that explicitly

account for preservation-related effects. Future developments may include preservation-aware

preprocessing pipelines, model architectures accommodating chemically induced spectral

shifts, or scan-level approaches that retain information currently lost through spectrum

averaging. A related methodological consideration concerns the decision to use a single

measurement per specimen versus averaging multiple scans. While averaging replicate scans

can reduce instrumental noise and improve classification stability, it may also obscure within-

specimen variability that could be biologically meaningful. The optimal strategy likely

depends on the research objective, whether prioritizing classification accuracy, trait

estimation, or the study of intra-individual variation. Artefacts introduced by glue, fungal

contamination, or leaf structure further complicate interpretation. Although preprocessing

methods such as SNV and Savitzky–Golay filtering help mitigate noise, they do not fully

resolve the boundary between noise and true biological variation. The choice of preprocessing

methods should therefore be preceded by systematic testing to determine which treatments are

most effective and discriminative, ideally by evaluating multiple combinations of

preprocessing steps.

Beyond analytical considerations, the practical implementation of spectral acquisition also

deserves attention. Using our methods, scanning required approximately 10 minutes per

specimen, allowing up to 50 specimens to be processed during a standard 8-hour working day.

This throughput highlights the operational feasibility of herbarium-based spectral acquisition.

However, more detailed protocols are now available (White et al., 2025a), and the time

required may vary depending on the objectives of the research question, the level of metadata

recorded, and the degree of replication implemented.

These practical considerations further emphasize the importance of harmonized

protocols when combining datasets across studies or collections. Differences in

specimen composition, preservation history, sensor characteristics, and measurement

conditions may introduce systematic biases if datasets are pooled uncritically. In

particular, instrument-specific artefacts, such as abrupt reflectance discontinuities or

jumps between sensor regions **(Fig. S2-S3)**, can vary among devices and may affect

spectral continuity in ways that are not immediately evident. Such features should be

carefully evaluated and, where necessary, corrected or accounted for before merging

datasets acquired with different instruments. Careful dataset-specific evaluation and

harmonization are therefore essential prerequisites for meaningful data aggregation.

More broadly, harmonizing protocols across studies remains essential for building

interoperable spectral datasets. Initiatives such as IHerbSpec (Cavender-Bares et al., 2025)

provide a strong foundation by promoting standardized specimen preparation, scanning

environments, and metadata reporting (White et al., 2025a). Adoption of such frameworks

will be key to ensuring that herbarium spectral data are both locally robust and globally

reusable.

Conclusion

The application of near infrared spectroscopy to herbarium specimens is still in its infancy.

Our results show that even old specimens retain enough taxonomic signal in their leaf

reflectance spectra to allow accurate species classification using standard machine learning

approaches. This confirms spectral scanning as a valuable, non-destructive source of data for

studies in taxonomy, ecology, and evolution (Meireles et al., 2020; Stasinski et al., 2021;

Kothari and Schweiger, 2022).

The potential of spectroscopy applied to herbarium-preserved specimens opens exciting

perspectives for plant biodiversity research, adding yet another important dimension to the use

of herbaria (Davis 2023). In addition to complementing traditional taxonomic research,

spectroscopy could also inform leaf functional traits and, in combination with advances in

machine learning (Soltis et al., 2020), enable large-scale, non-destructive trait estimation,

species identification, and even the detection of cryptic diversity across historical collections.

Ensuring data quality and expanding spectral libraries across taxa and regions will be key to

unlocking the full potential of this approach.

532 CRediT Author contributions:
533

KB: Conceptualization, Data curation, Formal analysis, Writing (original draft). **TLPC**:

Conceptualization, Supervision, Methodology, Validation, Writing (review & editing),

Investigation (field sampling), Funding acquisition. **PGSA**: Methodology, Investigation (field

sampling), Writing (review) **LAB**: Methodology, Writing (review & editing). **ME**:

Methodology, Writing (review & editing). **SB**: Methodology (instrumentation), Writing

(review). **DB**: Methodology (instrumentation), Writing (review). **LB**: Methodology

(instrumentation), Writing (review).

542 Acknowledgment
543

The authors thank the Muséum national d'Histoire naturelle (MNHN), Paris, for access to its

herbarium collections, and the Yasuní National Park authorities for permitting field sampling

and supporting research. We are grateful to Dr. Laura Holzmeyer for taxonomy training prior

to specimen selection at P. This research was supported by the European Research Council

(ERC) under the European Union's Horizon 2020 program (grant agreement No. 865787).

Data - Code availability

All code and data used in this study are available from the IRD Forge repository :

https://forge.ird.fr/diade/global_erc/boughalmi_nirs_paris_yasuni_data_2025

REFERENCES

Asner, G. P., and R. E. Martin. 2016. Convergent elevation trends in canopy chemical traits of
tropical forests. *Global Change Biology* 22: 2216–2227.

Barnes, R. J., M. S. Dhanoa, and S. J. Lister. 1989. Standard Normal Variate Transformation
and De-Trending of Near-Infrared Diffuse Reflectance Spectra. *Applied Spectroscopy*
43: 772–777.

Besnard, G., M. Gaudeul , S. Lavergne, S. Muller, G. Rouhan , A. P. Sukhorukov, A.
Vanderpoorten , and F. Jabbour. 2018. Herbarium-based science in the twenty-first
century. *Botany Letters* 165: 323–327.

Botermans, M., M. S. M. Sosef, L. W. Chatrou, and T. L. P. Couvreur. 2011. Revision of the
African Genus *Hexalobus* (Annonaceae). *Systematic Botany* 36: 33–48.

Brabant, C., A.-V. Emilien, A. Laribi, G. Morin, K. Nguyen, A. Thomas, and T. Houet. 2019.
Comparison of Hyperspectral Techniques for Urban Tree Diversity Classification.
Remote Sensing 11: 1269.

Breiman, L., A. Cutler, A. Liaw, and M. Wiener. 2024. randomForest: Breiman and Cutlers
Random Forests for Classification and Regression.

Cao, J., X. Wang, J. F. Adamowski, A. Biswas, C. Liu, Z. Chang, and Q. Feng. 2020.
Response of leaf stoichiometry of *Oxytropis ochrocephala* to elevation and slope
aspect. *CATENA* 194: 104772.

Cavender-Bares, J., J. A. Gamon, S. E. Hobbie, M. D. Madritch, J. E. Meireles, A. K.
Schweiger, and P. A. Townsend. 2017. Harnessing plant spectra to integrate the
biodiversity sciences across biological and spatial scales. *American Journal of Botany*
104: 966–969.

Cavender-Bares, J., D. M. White, N. I. Ahlstrand, M. Austin, D. Bastianelli, S. Bazan, K.
Boughalmi, et al. 2025. NextGeneration specimen digitization: The international
herbarium community goes spectral!

Chatrou, L. W., R. H. J. Erkens, J. E. Richardson, R. M. K. Saunders, and M. F. Fay. 2012.
The natural history of Annonaceae: NATURAL HISTORY OF ANNONACEAE.
Botanical Journal of the Linnean Society 169: 1–4.

Couvreur, T. L. P., A. J. Helmstetter, E. J. M. Koenen, K. Bethune, R. D. Brandão, S. A.
Little, H. Sauquet, and R. H. J. Erkens. 2019. Phylogenomics of the Major Tropical
Plant Family Annonaceae Using Targeted Enrichment of Nuclear Genes. *Frontiers in*
Plant Science 9: 1941.

Couvreur, T. L. P., P. J. M. Maas, S. Meinke, D. M. Johnson, and P. J. A. KEßLER. 2012.
Keys to the genera of Annonaceae: ANNONACEAE GENERIC KEYS. *Botanical*
Journal of the Linnean Society 169: 74–83.

Couvreur, T. L. P., M. D. Pirie, L. W. Chatrou, R. M. K. Saunders, Y. C. F. Su, J. E.
Richardson, and R. H. J. Erkens. 2011. Early evolutionary history of the flowering
plant family Annonaceae: steady diversification and boreotropical geodispersal.

Journal of Biogeography 38: 664–680.

Dagallier, L.-P. M. J., F. M. Mbago, M. Couderc, M. Gaudeul, A. Grall, C. Loup, J. J.
Wieringa, et al. 2023. Phylogenomic inference of the African tribe Monodoreae
(Annonaceae) and taxonomic revision of *Dennettia*, *Uvariodendron* and *Uvariopsis*.
PhytoKeys 233: 1–200.

Davis, C. C. 2023. The herbarium of the future. *Trends in Ecology & Evolution* 38: 412–423.

Durgante, F. M., N. Higuchi, A. Almeida, and A. Vicentini. 2013. Species Spectral Signature:
Discriminating closely related plant species in the Amazon with Near-Infrared Leaf-
Spectroscopy. *Forest Ecology and Management* 291: 240–248.

Forrest, L. L., M. L. Hart, M. Hughes, H. P. Wilson, K.-F. Chung, Y.-H. Tseng, and C. A.
Kidner. 2019. The Limits of Hyb-Seq for Herbarium Specimens: Impact of
Preservation Techniques. *Frontiers in Ecology and Evolution* 7: 439.

Foutami, I. J., T. Mariager, R. Rinnan, C. J. Barnes, and N. Rønsted. 2018. Hundred Fifty
Years of Herbarium Collections Provide a Reliable Resource of Volatile Terpenoid
Profiles Showing Strong Species Effect in Four Medicinal Species of Salvia Across
the Mediterranean. *Frontiers in Plant Science* 9.

Hodge, W. H. 1947. The Use of Alcohol in Plant Collecting. *Rhodora* 49: 207–210.

Jacquemoud, S., and S. Ustin. 2019. Leaf Optical Properties. Cambridge University Press,
Cambridge.

Jombo, S., E. Adam, M. J. Byrne, K. A. Ali, and S. W. Newete. 2020. Exploring the Potential
of Feature Selection Methods in the Classification of Urban Trees Using Field
Spectroscopy Data.

Karatzoglou, A., A. Smola, K. Hornik, N. I. Australia (NICTA), M. A. Maniscalco, and C. H.
Teo. 2024. kernlab: Kernel-Based Machine Learning Lab.

Khondoker, M., R. Dobson, C. Skirrow, A. Simmons, and D. Stahl. 2016. A comparison of
machine learning methods for classification using simulation with multiple real data
examples from mental health studies. *Statistical Methods in Medical Research* 25:
1804–1823.

Kong, W., C. Zhang, F. Liu, P. Nie, and Y. He. 2013. Rice Seed Cultivar Identification Using
Near-Infrared Hyperspectral Imaging and Multivariate Data Analysis. *Sensors* 13:
8916–8927.

Kothari, S., R. Beauchamp-Rioux, E. Laliberté, and J. Cavender-Bares. 2023. Reflectance
spectroscopy allows rapid, accurate and non-destructive estimates of functional traits
from pressed leaves. *Methods in Ecology and Evolution* 14: 385–401.

Kothari, S., and A. K. Schweiger. 2022. Plant spectra as integrative measures of plant
phenotypes. *Journal of Ecology* 110: 2536–2554.

Kuhn, M., J. Wing, S. Weston, A. Williams, C. Keefer, A. Engelhardt, T. Cooper, et al. 2024.

630   caret: Classification and Regression Training.

631 Kühn, P., T. Proß, C. Römermann, K. Wesche, and H. Bruelheide. 2024. Using near-infrared
632   spectroscopy to predict nitrogen and phosphorus concentrations of herbarium
633   specimens under different storage conditions. *Plant Methods* 20: 19.

634 Lang, P. L. M., F. M. Willems, J. F. Scheepens, H. A. Burbano, and O. Bossdorf. 2019. Using
635   herbaria to study global environmental change. *New Phytologist* 221: 110–122.

636 Le Bras, G., M. Pignal, M. L. Jeanson, S. Muller, C. Aupic, B. Carré, G. Flament, et al. 2017.
637   The French Muséum national d'histoire naturelle vascular plant herbarium collection
638   dataset. *Scientific Data* 4: 170016.

639 Lissambou, B.-J., O. J. Hardy, C. Atteke, T. Stevart, G. Dauby, B. Mbatchi, B. Sonke, and T.
640   L. P. Couvreur. 2018. Taxonomic revision of the African genus *Greenwayodendron*
641   (Annonaceae). *PhytoKeys* 114: 55–93.

642 Maire, V., I. J. Wright, I. C. Prentice, N. H. Batjes, R. Bhaskar, P. M. Van Bodegom, W. K.
643   Cornwell, et al. 2015. Global effects of soil and climate on leaf photosynthetic traits
644   and rates. *Global Ecology and Biogeography* 24: 706–717.

645 Meireles, J. E., J. Cavender-Bares, P. A. Townsend, S. Ustin, J. A. Gamon, A. K. Schweiger,
646   M. E. Schaepman, et al. 2020. Leaf reflectance spectra capture the evolutionary
647   history of seed plants. *New Phytologist* 228: 485–493.

648 Mendes Resende, J. V., N. M. D. de Sá, M. T. L. de Oliveira, R. C. Lopes, R. Garrett, and R.
649   Moreira Borges. 2020. Chemical profiling of herbarium samples of *Solanum*
650   (Solanaceae) using mass spectrometry. *Phytochemistry Letters* 36: 99–105.

651 Mersni, J., S. Bazan, E. Baby, L. Bonnal, and D. Bastianelli. 2025. Protocol for NIR spectra
652   acquisition on herbarium specimen. Website https://agritrop.cirad.fr/612773/ [accessed
653   16 February 2026].

654 Meyer, D., E. Dimitriadou, K. Hornik, A. Weingessel, and F. Leisch. 1999. e1071: Misc
655   Functions of the Department of Statistics, Probability Theory Group (Formerly:
656   E1071), TU Wien. 1.7-17.

657 Neto-Bradley, B. M., P. Bonnet, H. Goëau, A. Joly, J. Cavender-Bares, and D. A. Coomes.
658   2025. Using reflectance spectra and Pl@ntnet to identify herbarium specimens: a case
659   study with *Lithocarpus*. *New Phytologist*.

660 Nge, F. J., T. Chaowasku, A. Damthongdee, C. Wiya, V. R. C. Soulé, C. Rodrigues-Vaz, D.
661   Bruy, et al. 2024. Complete genus-level phylogenomics and new subtribal
662   classification of the pantropical plant family Annonaceae. *TAXON* 73: 1341–1369.

663 Nualart, N., N. Ibáñez, I. Soriano, and J. López-Pujol. 2017. Assessing the Relevance of
664   Herbarium Collections as Tools for Conservation Biology. *The Botanical Review* 83:
665   303–325.

666 Pennington, P. T., Q. C. B. Cronk, J. A. Richardson, J. E. Richardson, L. W. Chatrou, J. B.
667   Mols, R. H. J. Erkens, and M. D. Pirie. 2004. Historical biogeography of two

cosmopolitan families of flowering plants: Annonaceae and Rhamnaceae.
Philosophical Transactions of the Royal Society of London. Series B: Biological
Sciences 359: 1495–1508.

Raxworthy, C. J., and B. T. Smith. 2021. Mining museums for historical DNA: advances and
challenges in museomics. *Trends in Ecology & Evolution* 36: 1049–1060.

Richter, R., B. Reu, C. Wirth, D. Doktor, and M. Vohland. 2016. The use of airborne
hyperspectral data for tree species classification in a species-rich Central European
forest area. *International Journal of Applied Earth Observation and Geoinformation*
52: 464–474.

Ripley, B., B. Venables, D. M. Bates, K. H. (partial port ca 1998), A. G. (partial port ca
1998), and D. F. (support functions for polr). 2025. MASS: Support Functions and
Datasets for Venables and Ripley's MASS.

Rohart, F., B. Gautier, A. Singh, and K.-A. L. Cao. 2017. mixOmics: An R package for
'omics feature selection and multiple data integration. *PLOS Computational Biology*
13: e1005752.

Rosche, C., O. Broennimann, A. Novikov, V. Mrázová, G. V. Boiko, J. Danihelka, M. T.
Gastner, et al. 2025. Herbarium specimens reveal a cryptic invasion of polyploid
Centaurea stoebe in Europe. *New Phytologist* 245: 392–405.

Santacruz Endara, P. G., A. Lozinguez, R. Valencia, S. A. Queenborough, M. Endara, B.
Obando-Tello, and T. L. P. Couvreur. 2025. Discreet but diverse and specific:
Determining plant-herbivore interactions across a species-rich plant family in a
tropical rain forest. *Biotropica* 57: e13392.

Savitzky, Abraham., and M. J. E. Golay. 1964. Smoothing and Differentiation of Data by
Simplified Least Squares Procedures. *Analytical Chemistry* 36: 1627–1639.

Soltis, P. S., G. Nelson, A. Zare, and E. K. Meineke. 2020. Plants meet machines: Prospects
in machine learning for plant biology. *Applications in Plant Sciences* 8: e11371.

Song, W., H. Wang, P. Maguire, and O. Nibouche. 2018. Collaborative representation based
classifier with partial least squares regression for the classification of spectral data.
Chemometrics and Intelligent Laboratory Systems 182: 79–86.

Stasinski, L., D. M. White, P. R. Nelson, R. H. Ree, and J. E. Meireles. 2021. Reading light:
leaf spectra capture fine-scale diversity of closely related, hybridizing arctic shrubs.
New Phytologist 232: 2283–2294.

Trainor, P. J., A. P. DeFilippis, and S. N. Rai. 2017. Evaluation of Classifier Performance for
Multiclass Phenotype Discrimination in Untargeted Metabolomics. *Metabolites* 7: 30.

Valencia, R., R. B. Foster, G. Villa, R. Condit, J.-C. Svenning, C. Hernández, K.
Romoleroux, et al. 2004. Tree species distributions and local habitat variation in the
Amazon: large forest plot in eastern Ecuador. *Journal of Ecology* 92: 214–229.

Wang, Q., X. Cui, G. Teng, Y. Zhao, and K. Wei. 2020. Evaluation and improvement of

model robustness for plastics samples classification by laser-induced breakdown
spectroscopy. *Optics & Laser Technology* 125: 106035.

White, D., N. I. Ahlstrand, M. Austin, D. Bastianelli, S. Bazan, K. Boughalmi, W. Cardinal-
McTeague, et al. 2025a. IHerbSpec/IHerbSpec-protocol: IHerbSpec Protocol v1.0.

White, D. M., J. Cavender-Bares, C. C. Davis, J. A. Guzmán Q., S. Kothari, J. M. Robles, and
J. E. Meireles. 2025b. Seeing herbaria in a new light: leaf reflectance spectroscopy
unlocks trait and classification modeling in plant biodiversity collections. *New
Phytologist*.

Zdunek, R., M. Nowak, and E. Plínski. 2016. Statistical classification of soft solder alloys by
laser-induced breakdown spectroscopy: review of methods. *Journal of the European
Optical Society-Rapid Publications* 11: 16006i.

FIGURES/TABLES

TABLE 1. Overview of Paris herbarium specimens included in the study, with number scanned, geographic origin, specimen age span, and subfamily classification – ID_PARIS.

Species	Specimens scanned	Geographic origin	Specimen age span	Subfamily
Ambavia capuronii (Cavaco & Keraudren) Le Thomas	17	Madagascar	1953-2013	Ambavioideae
Ambavia gerrardi (Baill.) Le Thomas	18	Madagascar	1956-2010	Ambavioideae
Anaxagorea dolichocarpa Sprague & Sandwith	10	French Guiana	1845 - 2008	Anaxagoreoideae
Annona senegalensis Pers.	9	Zambia, Tanganyika, Mali, Sierra leone, Guinea	1893 - 1951	Annonoideae
Anonidium mannii (Oliv.) Engl. & Diels	11	Cameroon	1918 - 1995	Annonoideae
Duguetia calycina Benoist	12	French Guiana	1857 - 2003	Annonoideae
Greenwayodendron suaveolens (Engl. & Diels) Verdc.	11	Gabon, DR Congo, Central African Republic, Cameroon	1881 - 1991	Malmeoideae
Hexalobus crispiflorus A.Rich.	10	Congo republic, Guinea, Cameroon, Gabon, Senegal	1838 - 1988	Annonoideae
Hexalobus monopetalus (A.Rich.) Engl. & Diels	11	Mali, Cameroon, Central African Republic, Senegal, Niger, Ivory coast	1929 - 2007	Annonoideae
Monanthotaxis enghiana (Diels) P.H.Hoekstra	9	Cameroon, Gabon, Central African Republic	1948 - 1989	Annonoideae
Unonopsis guatterioides (A.DC.) R.E.Fr.	12	Brasil, Paraguay, Saint Vincent and the Grenadines, Surinam, Bolivia	1886 - 1979	Malmeoideae
Uvaria grandiflora Roxb. ex Hornem.	11	Vietnam, Malaysia, China, Indonesia	1823 - 2015	Annonoideae
Uvariastrum pierreanum Engl.	10	Gabon, Cameroon, Ivory coast, Congo republic, Liberia, Nigeria	1948 - 2013	Annonoideae
Xylopia humblotiana Baill.	9	Madagascar	1924 - 2007	Annonoideae

721　　**TABLE 2**. Yasuní National Park dataset total specimen count and pre-drying alcohol-treated

722　　specimens – ID_YASUNÍ.

Species	Total specimen count	Alcohol-treated specimen count
Anaxagorea brevipes Benth.	7	2
Duguetia hadrantha (Diels) R.E.Fr.	6	1
Guatteria guianensis (Aubl.) R.E.Fr.	6	1
Guatteria scalarinervia D.R.Simpson	5	1
Oxandra riedeliana R.E.Fr.	5	1
Tetrameranthus globuliferus Westra	7	2
Trigynaea triplinervis D.M.Johnson & N.A.Murray	6	1
Unonopsis veneficiorum (Mart..) R.E.Fr.	7	2
Xylopia cuspidata Diels	6	1

Figure 1. Workflow illustrating the classification pipeline used. Specimens from two

datasets were scanned using spectrometers (ASD LabSpec), collecting 20 individual spectra per

specimen. Spectral data was preprocessed, and databases were split into training and test sets

across 100 replicates. Five supervised classification models were trained and then tested on

either individual spectra or on spectra averaged per specimen. Model performance was assessed

using accuracy.

TABLE 3. Comparison of spectral acquisition protocols and instrumentation between the

ID_PARIS and ID_YASUNÍ datasets.

Aspect	ID_PARIS	ID_YASUNÍ
Instrument	ASD LabSpec® Standard-Res Lab Analyzer	ASD LabSpec® Pro spectrophotometer
Probe type	Rapid Analysis Probe (90° angle) + Fiber Optic Illuminator	6 mm diameter pen probe
Software version	Indico Pro v6.5	Indico Pro v3.1
Reference procedure	White baseline references at start of session and every hour (Spectralon® white)	White reference every 15 minutes (Spectralon® Metal SpectralTM)
Black background	Spectral Black™ strip beneath leaves	Spectral Black™ strip beneath leaves (mounted specimens) or larger black sheet (detached leaves)
Sample count/scan	20 readings averaged to generate one spectrum	30 readings averaged to generate one spectrum
Integration time	68 ms^{-1}	68 ms^{-1}
Number of scans	20 scans per specimen	50 scans per specimen
Operator/time	Single operator	Multiple operators

Figure 2. Species-level classification accuracy of leaf spectra. Classification performance for

14 species from ID_PARIS (a-*Ambavia capuronii*, b-*Ambavia gerrardi*, c-*Anaxagorea*

dolichocarpa, d-*Annona senegalensis*, e-*Anonidium mannii*, f-*Duguetia calycina*, g-

Greenwayodendron suaveolens, h-*Hexalobus crispiflorus*, i-*Hexalobus monopetalus*, j-

Monanthotaxis enghiana, k-*Unonopsis guatterioides*, l-*Uvaria grandiflora*, m-*Uvariastrum*

pierranum, n-*Xylopia humblotiana*) and 9 species from ID_YASUNÍ (o-*Anaxagorea brevipes*,

p-*Duguetia hadrantha*, q-*Guatteria guianensis*, r-*Guatteria scalarinervia*, s-*Oxandra*

riedeliana, t-*Tetrameranthus globuliferus*, u-*Trigynaea triplinervis*, v-*Unonopsis veneficiorum*,

w-*Xylopia cuspidata*) using five models: Partial Least Squares Discriminant Analysis (PLS-

DA), k-Nearest Neighbors (KNN), Principal Component Analysis followed by Linear

Discriminant Analysis (PCA-LDA), Support Vector Machines (SVM), and Random Forest

(RF) from 100 independent replicates using 75 % randomly selected specimens in training (6

for ID_PARIS, 3 for ID_YASUNÍ) all the others in the test, Measurement-level column mean

that all specimen spectra where classify and Specimen-level column that only the average

spectrum is classified. **(A)** Overall model accuracy and standard deviation. **(B)** Confusion

matrices represent mean classification results. Values indicate the proportion of spectra

predicted for each species class.

Figure 3. Performance of models trained on a single specimen per species. Classification accuracy (boxplots) across 100 replicates for each species and model, using one randomly selected specimen (20 spectra) as the training dataset. Background tiles indicate the mean classification accuracy across all models for each species. Results are based on the ID_PARIS dataset. Species are ordered from highest (top) to lowest (bottom) mean accuracy .

Figure 4. Effect of the number of training specimens per species on classification accuracy.

(A) Heatmap of mean classification accuracy across 100 resampling iterations for each model and training size (1 to 8 specimens per species) using the ID_PARIS dataset. **(B)** Accuracy trajectories (± standard deviation) across the same iterations, showing model performance as a function of the number of training specimens per species.

767 **SUPPLEMENTARY**

768 **Table S1.** List of herbarium specimens scanned from ID_Paris (barcode in brackets)

Species	Specimens
Ambavia capuronii	Schatz 2549 (P01966283); Razafimandimbison 247 (P01966290); Kotozafy 66 (P01966293); Rabe 6 (P01966295); Ravelonarivo 421 (P01966288); Ravelonarivo 617 (P01966289); Ravelonarivo 611 (P01966285); Turk 90 (P01966281); SF 8751 (P030238); SF 14256 (P02005959); SF 15301 (P01966279); Schatz 2536 (P01966282); Ranirison 812 (P06901328); Rakotondrajaono 107 (P01966280); Aridy 146 (P01966292); Malcomber 1028 (P01966294); Rakotonirina 128 (P01047823)
Ambavia gerrardi	Bosser 18703 (P02032521); Ranaivojaono 622 (P02032556); Rabenantoandro 1160 (P02032555), Randrianaivo 538 (P02032557); Randrianaivo 716 (P02032507); Rakotomalaza 441 (P02032523); SF 15383 (P02032525); Sauquet 94 (P00465816); Rahajasoa 608 (P02032505); Rabehevitra 3 (P06839299); Schatz 3096 (P02032508); Razanatsoa 522 (P06839300); Razakamalala 318 (P00858933); CNRE/PLARM 98 (P02032527); Evrard 11253 (P02032560); McPherson 18940A (P02032558); Rabehevitra 341 (P02133037); Rabevohitra 4893 (P06901451)
Anaxagorea dolichocarpa	Irwin 54545 (P01960236); Sagot 8 (P01960307); Lescure 875 (P01960277); Melinon 21 (P01960281); Melinon 31 (P01960283), Melinon 218 (P01960280); Granville 3706 (P01960643); Granville 5473 (P01960323); Granville 5950 (P01989415); Tostain 1669 (P01960545); Cremers 7680 (P01989410)
Annona senegalensis	s.c. 46 (P02032725); Nsama 1381 (P02032635); Mamboc 1411 (P02032637); Chevalier 2543 (P00517160); Chevalier 44019 (P00517162); Chevalier 13373bis (P02027965); Vuillet 410 (P00517154); Deigton 5442 (P02032682); Paroisse 107 (P02027962)
Anonidium mannii	Hedin 36bis (P02032735); Hedin 1337 (P02032738); Letouzey 194 (P02032741); Letouzey 11202 (P02032587); Kaji 98 (P02032588); de Wilde 7930 (P00353899); Cheek 7896 (P00956185); Cheek 6009 (P00956187); Raynal 9607 (P02032585); Annet 42 (P02032745); Breteler 2454 (P02032747)
Duguetia calycina	s.c. 7982 (P01984825); Melinon 93 (P01963746); Melinon 633 (P01963748); Deward 132 (P01984833); Mori 25703 (P06901455); Granville 5952 (P00124065); Granville 6253 (P01984844); Oldeman T-575 (P00124063); Villiers 3717 (P01748169); Sagot 1095 (P00083612); Poeppig 2772 (P01984856); Barrier 5112 (P01748171);
Greenwayodendron suaveolens	McPherson 15501 (P01965944); de Wilde 101 (P01965978); de Wilde 7940 (P01985243); Le Testu 7936 (P00363322); Gilbert 936 (P01965953); Tisserant 1172 (P01985835); Soyaux 218 (P00363356); Tisserant 2573 (P01084175); Zenker 1306 (P01985239); Leeuwenberg 7322 (P01985867); Letouzey 10456 (P01985871)

Hexalobus crispiflorus	Bouquet 895 (P00486232); Chevalier 13385 (P00486266); Pobéguin 176 (P00486263); Jaeger 9039 (P00486260); Félix 77 (P00486265); Fay 8299 (P00486244); Fleury 26627 (P00486257); Hallé 606 (P00486236); Harris 516 (P00486243); Heudelot 865 (P00486270); Birnbaum 1238 (P06901043)
Hexalobus monopetalus	Satabié 702 (P00486152); Megnikpa 2 (P00486153), Aubréville 415 (P00486158); Griaule N/A (P00486191); Chevalier 33487 (P00486200), Diallo 649 (P00486198), Boudouresque 5981 (P00486167); Baudet 3752 (P00486203); Peyre de Fabrègues 3408 (P00486160); Coussan 1158 (P00486176)
Monanthotaxis enghiana	Breteler 2137 (P01956022); Hallé 491 (P01985815); Hallé 367 (P01985814); Hallé 799 (P01985821); Hallé 3205 (P01985820); Hallé 2452 (P01985822); Tisserant 1257 (P01985786); Tisserant 2062 (P01985781); Harris 1721 (P01985779)
Unonopsis guatterioides	Labroy 107 (P00506137); Bernardy 19436 (P00506139); von Eggers 6882 (P00506141); Argent 6736 (P02132829); Westra 47623 (P02132839); Fróes 24503 (P02132835); Hostmann 1116 (P00506016); Rusby 1253 (P00506011); Fróes 1700 (P02132833); Harley 10597 (P02132825); Harley 10985 (P02132824); Santos 1660 (P02132826)
Uvaria grandiflora	Poilane 35624 (P00262277); Poilane 30328 (P00262274); Schaller KL3878 (P02132845); Poilane 40896 (P00262249); Hiêp 20 (P00262253); Couvreur 838 (P00938648); Robert 16 (P00262297); s.c. 836 (P00262239); Prawiroatmodjo 1387 (P02132755); Poilane 9789 (P00262263); Dilmy 12 (P02132754)
Uvariastrum pierreanum	McPherson 16128 (P01983191); Thomas 8094 (P01983291); Fogiel 1039 (P01983300); Letouzey 10225 (P01983295); Couvreur 590 (P02170712); Bakayoko 99 (P01983224); Koechlin 7758 (P01983223); Adam 26222 (P01983225); Akpabla 1121 (P01983228); Hallé 604 (P01983238)
Xylopia humblotiana	Razafindraibe 101 (P06901345); McPherson 98 (P06839280); Rabevohitra 3793 (P01031514); Rabevohitra 3792 (P01031513); Rogers 1007 (P02005890); Poncy 1597 (P00373125); McPherson 14371 (P01987117); Poncy 1594 (P00373123); Rakotonandrasana 676 (P01986879); Rakotovao 3823 (P01986878); Perrier de la bathie 15998 (P00524382); Lowry II 6030 (P01987062)

A

B

Supp. Figure 1. Age distribution of scanned Annonaceae specimens from the Paris herbarium (ID_Paris dataset)**. (A)** Histogram of specimen ages (calculated as 2025 minus collection year) showing the distribution of 160 scanned Annonaceae specimens from the Paris herbarium. **(B)** Boxplots of specimen ages grouped by genus. ID_PARIS.

Supp. Figure 2. Alcohol preservation effects on leaf spectral data. **(A)** three-dimensional PCA projection of the first three principal components; Find in the Forge repositories a link to the 3D space plot named FS2_3D_plot.Rdata . **(B)** Mean raw reflectance spectra per specimen grouped by species and pre-drying treatment (*Anaxagorea brevipes* and *Duguetia hadrantha*

782   alcohol (_A) vs. directly dried). An averaged spectrum is used to represent the specimen in both

783   analyses. ID_YASUNÍ.

785   **Supp. Figure 3. Reflectance spectra from ID_PARIS.** Mean raw reflectance spectra per

786   specimen for *Greenwayodendron suaveolens* and *Hexalobus monopetalus*. ID_PARIS.