Exploring the roles of trophic mode and microbial prey size in grazing

pathways of tropical oligotrophic waters of the eastern Indian Ocean

Claudia Traboni[a*], Grace F. Cawley[b], Karen E. Selph[c], Michael R. Landry[b], Moira Décima[b]

[a] Integrative Marine Ecology, Stazione Zoologica Anton Dohrn, Naples, 80121, Italy

[b] Scripps Institution of Oceanography, University of California at San Diego, La Jolla, CA

92093, USA

[c] Department of Oceanography, University of Hawai'i at Manoa, Honolulu, HI 96822, USA

*Corresponding author

E-mail address: claudia.traboni@szn.it (C. Traboni)

Keywords: copepod grazing, microzooplankton grazing, top-down control, microbial food

web, mixotrophy, trophic cascade, nutrient channeling

ABSTRACT

Prey removal incubations were conducted in the Argo Basin (eastern Indian Ocean) to

investigate the trophic ecology of the zooplankton community supporting larvae of Southern

Bluefin Tuna. Grazing and selectivity were evaluated considering prey trophic mode and cell

size as key food quality descriptors in copepod diets, and the impact of microzooplankton

grazing was quantified for comparison. Copepod ingestion rate on eukaryotes ranged from

3.4 to 138 ng C ind^{-1} d^{-1}. Diet was mostly derived from mixotrophic (5-89%) and

heterotrophic (0-84%) prey, while autotrophs had a minor contribution (2-17%). Overall,

nanoplankton provided the highest C intake to copepods (62-99%) rather than more

abundant but smaller picoplankton (0.8-38%). Copepod-mediated Chl$a$ removal was not

detected, although gut content revealed ingestion of 0.8 µg Chl$a$ ind^{-1} d^{-1}. Slight selection

emerged in favor of picoplankton versus nanoplankton, suggesting a possible trophic

cascade. Ingestion of ^{15}N-labelled prey indicated active consumption by copepods of

consumers (protistan nano-micrograzers or small metazoans). We found significantly greater

nitrogen consumption (~5-fold) when using 1-2µm ^{15}N-prey (*Synechococcus*), compared to

<1µm sized prey (*Prochlorococcus*). Microzooplankton grazing on eukaryotic (0.07–2.5 d^{-1})

and prokaryote (0.3–2.1 d^{-1}) prey greatly exceeded copepod grazing. Microzooplankton diet

consisted mainly of heterotrophs (25-59%), and mixotrophs (13-41%) with a lower

contribution of autotrophs (12-33%). Nanoplankton constituted the main dietary source for

microzooplankton (95-98%), whereas picoplankton was 2-5%. Most of the daily production

was removed by microzooplankton (111%) leaving copepods a secondary role in

consumption (7%), likely supported by ingestion of larger prey and indirect nutrient

channeling.

1. Introduction

Copepods feed on a wide range of protistan prey varying in taxonomy, size, mobility, trophic mode and biochemical composition (Saiz and Calbet, 2011). The size ratio between prey and predator is a master trait determining capture and ingestion probability (Hansen et al., 1994), but trophic mode can also be an important factor affecting nutritional quality of copepod prey (Traboni et al., 2021). Autotrophic phytoplankton depend on inorganic nutrients and sunlight to sustain their photosynthetic metabolism and accumulate carbon-rich biomass, whereas heterotrophic protozooplankton are obligate consumers of organic matter that are more enriched in nitrogen and phosphorus relative to carbon. Mixotrophic protists (mixoplankton) combine both trophic modes (Flynn et al., 2019; Stoecker et al., 2017) and are characterized by intermediate nutritional composition. Especially under conditions where dissolved concentrations of nutrients are very low, the ability of mixotrophs to acquire nutrients from phagotrophy can improve food quality by reducing prey–predator stoichiometric mismatch (Schenone et al., 2024) and enhancing nutrient transfer efficiency (Ward and Follows, 2016). Laboratory experiments have demonstrated that copepods feed abundantly on mixotrophs (Jeong et al., 2010; Ptacnik et al., 2004) and are not negatively affected by consumption of mixotrophs compared to other trophic modes (Traboni et al., 2020). However, only a few field studies consider this food-web interaction (Castellani et al., 2005; Fileman et al., 2007; Jonsson and Tiselius, 1990), likely due to methodological challenges in distinguishing mixotrophs in natural assemblages.

Mixotrophy is a prevalent adaptive strategy among planktonic protists of oligotrophic regions, overcoming resource limitation of photo- and phagotrophy specialists (Anschütz et al., 2024; Edwards, 2019) by using photosynthesis for growth when nutrients allow and phagotrophy when nutrients/light are scarce (Stoecker et al., 2017). The mixotrophic strategy has been reported worldwide in oligotrophic regions of the North Atlantic (Hartmann et al., 2012; Millette et al., 2016), East China Sea (Li et al., 2024), Sargasso Sea (Arenovski et al., 1995) and Mediterranean Sea (Unrein et al., 2014). Of particular relevance to the current study, Raes et al. (2022) found an 18S-rRNA diversity peak for mixotrophs in high-temperature low-nutrient tropical waters of the eastern Indian Ocean (IO). Elevated food-web transfer efficiency associated with mixotrophy was also invoked to explain especially high zooplankton grazing on chlorophyll-containing protists in this region (Landry et al., 2020a).

In January-March 2022, the BLOOFINZ-IO (Bluefin Larvae in Oligotrophic Ocean Foodwebs, Investigations of Nutrients to Zooplankton – Indian Ocean) expedition investigated the physical, biogeochemical and food web characteristics of waters overlying the Argo Abyssal Plain (hereafter, Argo Basin), a 5000-m deep oceanic basin directly downstream of the Indonesian Throughflow (ITF), where the tropical surface water mass

from the western Pacific first enters the eastern IO (Woo and Pattiaratchi, 2008). This is also

the only known global spawning region for Southern Bluefin Tuna, *Thunnus maccoyii* (Davis

et al., 1991), which underlies the BLOOFINZ goal of understanding details of the food web

supporting feeding, growth and survival of the species through its vulnerable larval stages.

As part of this effort, we conducted experimental incubations to explore the roles of size

structure and trophic mode in grazing pathways to microzooplankton and copepods.

Mixotrophs were identified by acidic vacuole staining and flow cytometry. We hypothesized

that protistan phagotrophy would play a significant role in nutrient channeling in the microbial

food web of this oligotrophic region and that mixotrophs would be a high fraction of

community composition and copepod diet.

2. Materials and methods

2.1. Grazing experiments: study area and timeline of the incubations

Zooplankton grazing experiments were performed aboard R/V *Roger Revelle* during

BLOOFINZ-IO cruise RR2201, with sampling taking place between February 4[th] and March

2[nd] 2022, off NW Australia (Argo Basin) (Fig. 1A,B). During the cruise, daily CTD casts were

conducted to monitor vertical profiles of abiotic (temperature, salinity, irradiance, oxygen)

and biotic features (Chl*a*, microbial abundance and biomass) of the water column, as well as

to collect water for experiments (Table 1). Grazing incubations were mostly conducted

during multi-day experiments (hereafter "cycles") following a satellite-tracked drifter (Landry

et al., 2009; Landry et al. this volume). Nine grazing experiments were conducted in total

(Exps 1-9). The first seven experiments were performed during Cycles 1-4, and two final

grazing incubations were conducted during transits along the return leg of the cruise (Table

1; Fig. 1B). Each experiment consisted of two incubation phases: a 24-h incubation with the

microzooplankton community alone ("Microzoo" incubation, T_0-T_{24}) followed by a 12-h

incubation with added copepods ("Microzoo+copepod" incubation, T_{24}-T_{36}) (Fig. 2, Table 2).

2.2. Seawater and zooplankton sampling

Seawater for grazing incubations was typically collected from Niskin bottles on CTD

casts in early evening (19:00 to 20:30 local; Fig. 2). During cycles (Exps 1-7), low-Chl*a* low-

nutrient water was sampled between 28 and 30 m depth. For the two transit experiments

(Exps 8-9), water was collected from the surface (2 m depth) using trace metal-clean bottles

(Table 1).

Between 20:30 and 23:00, after seawater collection, zooplankton live samples were

collected (Fig. 2) with gentle vertical tows (~10 m min^{-1}) using a 1-m^2 ring net (200-μm

mesh), equipped with a non-filtering 30-L cod end and depth sensor attached to the net

frame. Sampling depths ranged from 38 to 60 m depending on the experiment (Table 1). The

animals were carefully placed in two 20-L buckets, diluting them by 1/3 with fresh filtered

seawater, and allowed to acclimate for ~22 h in a temperature-controlled room (set to the

average sea surface temperature) in the darkness until sorted for experiments.

2.3. Bottle treatments and experimental procedures

Each experiment included: a) three initial "Start" bottles, filled with seawater and

immediately sampled after filling (T_0); b) three "30%" bottles containing 30% seawater and

70% 0.1 µm-Acropack filtered seawater (FSW); c) three "100%" bottles with undiluted

seawater; d) three "CopStart" bottles including undiluted seawater, representing the starting

conditions for copepod incubations; e) three "Cop+Isotopes" bottles containing undiluted

seawater, ^{15}N-labelled prey and ^{13}C bicarbonate inoculated at T_0, and copepods, which were

added ~22-24 h later (at T_{24}; Fig. 2).

Microzoo incubation (T_0-T_{24}). Water from the evening CTD cast was transferred with

acid-washed silicone tubes into 2.7-L acid-washed transparent polycarbonate bottles

following three seawater rinses. The bottles were filled to capacity avoiding bubbles and

ensuring a slow and gentle flow. For Exps 8 and 9, the water was first collected with trace

metal-clean 5-L GO-Flo bottles, then gently homogenized in carboys and added to the 2.7-L

bottles as for other experiments. During these two last experiments, 5 mL water was

removed from all the bottles and replaced with 5 mL of underway surface seawater to ensure

non-limiting trace metal concentration for photosynthesis.

Before the start of the cruise, ^{15}N-labelled *Prochlorococcus* (*Pro*) and ^{15}N-labelled

*Synechococcu*s (*Syn*) were prepared in the laboratory by filtering ^{15}N-labelled cell culture

stocks onto 25-mm diameter 0.2-µm polycarbonate filters (ca. 2-8 x 10^8 cells filter^{-1}). Filters

were flash frozen in liquid nitrogen in 2-mL cryo-eppendorf tubes and long-term stored at -

80°C. Prior to each experiment, the ^{15}N-labelled cells from one filter were resuspended in

13-14 mL of 0.1-µm Acropack FSW, mixed by vortex, and refrigerated for ~1 h before

spiking (2-4 mL) experimental bottles to start (T_0) the Microzoo incubation into the

Cop+Isotope bottles (where copepods will be added on the next day; Fig. 2). Depending on

the experiment, the ^{15}N-labelled cyanobacteria ranged from 6,700 to 102,000 cells mL^{-1}

(Table 2). Additionally, 1 mL of ^{13}C bicarbonate stock was inoculated to reach a final

concentration of 0.25 mM ^{13}C in the same Cop+Isotope bottles. For Exps 1-3, only ^{15}N-*Pro*

was added, whereas Exps 4-9 also received ^{15}N-*Syn,* usually in concentrations that were 5

to 10-fold those of ^{15}N-*Pro* (Table 2). After filling and adding labelled-prey, the bottles were

placed in a deckboard incubator exposed to 25% of incident PAR to match the irradiance at

the depth of seawater collection. The incubators were covered at night to ensure complete

darkness, and three Start bottles were sampled for initial conditions of the different abiotic

and biotic parameters.

Microzoo+copepod incubation (T_{24}-T_{36}). Copepods were sorted out of the cod end

community after ~20-24 h of conditioning in a constant-temperature room, by concentrating

the sample onto 200-µm mesh filters and transferring into small plastic jars containing

seawater. Aliquots of the concentrated zooplankton sample were poured into Petri dishes

and only healthy intact individuals were sorted with a glass Pasteur pipette under a

dissecting microscope. The selected copepod specimens were calanoids (*Candacia* spp.,

Euchaeta spp., *Labidocera* spp., other calanoids) and cyclopoids (*Oncaea* spp., *Sapphirina*

spp., and Corycaeidae). Sorted individuals were temporarily pipetted into 50-mL Falcon

tubes prefilled with 40 mL of FSW and covered with aluminum foil prior to inoculation. The

number of copepods incubated varied by experiment depending on availability (10-70; Table

2). Prior to introducing copepods into the experimental Cop+Isotope bottles, spiked with ^{13}C

and ^{15}N-labelled prey the previous day, the sorted animals were concentrated down to 15-20

mL by 50 µm-reverse filtration, and 15-20 mL of the bottle water was removed with a

serological pipette to allow for the addition of the copepods. The seawater volume containing

the copepods was then gently mixed and transferred to the corresponding bottles. From the

moment copepods were inoculated, the incubations lasted generally 12 h, except during

Exp2 (6 h). Three Cop Start bottles were sampled for initial conditions, and the samples

were processed according to the same protocols for the Microzoo incubation samples. An

aliquot of copepods from the mixed community were stored on 50-µm mesh filters and

frozen at -80ºC until analysis of their elemental contents and natural C and N isotope values

(see Section 2.4). After ~12 h from copepod addition *(T_{36})*, all the bottles were removed from

the deck incubator, quickly transported into the laboratory in the dark and processed for final

sampling following the same order in which they were previously filled and sampled.

2.4. Sample processing

At T_0 and T_{24}, experimental bottles were homogenized, and aliquots were collected for

Chl*a* fluorescence and flow cytometric measurements of cell abundances (Fig. 2). At the end

of copepod incubations (T_{36}), the copepods were removed from the Cop+Isotope bottles by

reverse filtration through 200-µm mesh, counted under a dissecting microscope, and filtered

onto pre-weighed 50-µm sieves. Each filter was stored in a capped petri dish and

immediately frozen (-80°C). The remaining water in the bottles was homogenized and

subsampled as for the Microzoo incubation.

For Chl*a* estimates, 285 mL subsamples were collected in dark amber bottles and

filtered onto 25-mm diameter GF/F Whatman filters. The filters were folded and the Chl*a*

extracted in 5 mL 90% acetone kept in glass screw-cap tubes in a -20°C freezer in total

darkness. After 24 h, the samples were vortexed, filters removed and Chl*a* fluorescence

measured with a Turner Designs model 10AU Field fluorometer calibrated against a pure

Chl*a* standard (Strickland and Parsons, 1972). Flow cytometry (930 µL) samples were

collected to estimate cell abundances of prokaryotes and protistan eukaryotes. Live runs

were done to enumerate pico- and nanoplankton eukaryotes in the categories of

picoautotrophs, nanoautotrophs, picomixotrophs, nanomixotrophs, and nanoheterotrophs.

The live samples were stained with two probes: Hoechst 34580 (DNA-specific; Selph, 2021)

and LysoTracker Green (acidic organelle-specific). The latter was used to detect the

presence of food vacuoles in heterotrophic (Rose et al., 2004) and mixotrophic protists (Sato

and Hashihama, 2019) on living cells since Lysotracker Green is not retained in dead cells'

vacuoles (more details in Selph et al., this volume). After 5-10 min incubation with stains

under dim light, the live samples were analyzed for particle concentration based on detectors

for blue fluorescence from DNA (450±45 nm, excited with a 375 nm laser), green and red

fluorescence of LysoTracker Green (525±40 nm) and Chl*a* (690±50 nm), respectively, from a

488 nm laser, phycoerythrin (575 nm) from a 561 nm laser, and forward and side light

scatter using a portable Beckman Coulter CytoFlex S flow cytometer. For prokaryotes, an

additional aliquot (930 µL) was preserved with 30 µL of 16% paraformaldehyde and

immediately frozen at -80°C for a posteriori enumeration of heterotrophic bacteria,

Prochlorococcus, and *Synechococcus* (referred to as Hbact, *Pro*, and *Syn,* respectively).

These samples were thawed in batches on the ship and abundances determined by Hoechst

34580 DNA after staining 1 h before analysis (Selph, 2021).

Copepods, both initial and those in the grazing experiments, were dried and packaged in

tin cups for elemental analyzer isotope-ratio mass spectrometry. Initials copepods were

measured for natural δ^{13}C and δ^{15}N, while grazing experiment copepods were measured for

enriched δ^{13}C and δ^{15}N. Depending on copepod size, we packaged between 2-8 copepods

per tin cups to ensure sufficient material for analysis. To estimate the isotopic content of

copepod food, particulate organic matter (POM) from the final spiked incubations was filtered

onto pre-combusted GF/F filters (1100-1880 mL), dried at 60°C for 24 h, packed into tin

cups, and sent for analysis. All isotopic samples were processed at the University of

California (Davis).

2.5. Determination of feeding rates and food selection

The net growth rates (Eq. 1a, b) of prey in the 30% diluted samples (k_d) and in the 100%

undiluted samples (k) were calculated as follows:

213
$$k_d = \frac{LN\left(\frac{Prey_{end_{30\%}}}{Prey_{start_{100}}} \; x\frac{1}{D}\right)}{t} \qquad \text{(Eq. 1a)}$$

214
$$k = \frac{LN\left(\frac{Prey_{end_{100\%}}}{Prey_{start_{100}}}\right)}{t} \qquad \text{(Eq. 1b)}$$

where $Prey_{end}$ and $Prey_{start}$ are prey concentrations (cells mL^{-1}) at the end and start of the

incubation respectively, D is the dilution factor, and t is the incubation duration (d^{-1}).

Prey mortality by microzooplankton grazing rates (m, Eq. 2) were determined for Chl$a$

and unicellular prey groups including prokaryotes (Hbact, Pro, Syn), and eukaryotes

(picoautotrophs, nanoautotrophs, picomixotrophs, nanomixotrophs, nanoheterotrophs) using

equations from Landry et al. (2008; this volume) as:

221
$$m = \frac{(k_d - k)}{(1 - D)} \qquad \text{(Eq. 2)}$$

222   To correct copepod grazing estimates for microzooplankton grazing, the adjustment

223 suggested by Nejstgaard et al. (2001) was applied (Eq. 3, Eq. 4). Nanoheterotrophs and

224 nanomixotrophs (total nanophagotrophs) were assumed to exert the main grazing impact

225 within the microzooplankton community analyzed, not only for their abundance but also for

226 their phagotrophic potential over picograzers:

$$k_p = g_{phago,p} \; x \frac{(C_{phago_{control}} - C_{phago_{cop}})}{C_{phago\_control}} \qquad \text{(Eq. 3)}$$

where
$g_{phago,p}$
is the
grazin
g by
nanop

$$g_{corr,p} = g_{cop,p} + k_p \qquad \text{(Eq. 4)}$$

233 hagotrophs on each prey type, C_{phago} is the mean nanophagotroph concentration in the

234 control and in the copepod bottles, and $g_{cop,p}$ is the grazing by copepods on each prey type.

235 Finally corrected clearance rate (F, mL ind^{-1} d^{-1}; Eq. 5) and ingestion rate (I, cells ind^{-1} d^{-1};

236 Eq. 6) were calculated according to Frost (1972) as:

237
$$F = \frac{V \; x \; g_{corr,p}}{N} \qquad \text{(Eq. 5)}$$

238
$$I = (C) \; x \; F \qquad \text{(Eq. 6)}$$

where V is the total volume of the incubation (mL), N is the number of live copepods

retrieved after the incubation and (C) is the average prey concentration (cells mL^{-1})

measured between start and end of the incubation in the copepod grazing bottles.

Food selection by copepods was expressed through the Chesson's α index (Chesson,

1978) to determine preference for the five eukaryotic prey groups and through the Ivlev's E

index (Ivlev, 1961) for trophic and size classes separately. Both α and E and were calculated

according to equations 7 and 8, respectively.

246
$$\alpha = \frac{(r_i/p_i)}{\Sigma(r_i/p_i)} \qquad \text{(Eq. 7)}$$

247   where $\Sigma(r_j/p_j)$ is the sum of all prey types' relative availability. α ranges from 0 to 1, with equal

248   preference being α =1/n (where n is the number of prey types, in our case, α = 1/5 = 0.2).

249   Values of α>0.2 indicate preference for prey type i, whereas α<0.2 represents avoidance of

250   prey type i.

251
$$E = \frac{(r_i - p_i)}{(r_i + p_i)} \qquad \text{(Eq. 8)}$$

where r_i is the proportion of prey type i in the predator's diet, and p_i is the proportion of prey

type i in the environment. E ranges from -1 to +1, being $E>0$ indicative of preference for prey

type i, and $E<0$ indicative of prey i avoidance, whereas $E=0$ indicates neutral ingestion,

which corresponds to equal preference. Both indices give a measure of how much an

ingested prey is preferred or avoided by copepods in proportion to its relative contribution in

the environment. However, Chesson's α index takes rare groups of prey into account,

whereas Ivlev's E index does not. Prokaryotes (Hbact, Pro, Syn) were excluded from the

copepod grazing analyses due to negligible or null consumption.

Prey abundance and copepod ingestion rates were converted to C biomass units from

literature equations. Prokaryote conversions were: Hbact: 5 fg C/cell (Gundersen et al.,

2002), Pro: 32 fg C/cell (Garrison et al., 2000), and Syn: 101 fg C/cell (Garrison et al., 2000).

Eukaryotic conversions used the mean equivalent spherical diameter (ESD, μm) measured

by epifluorescence microscopy to estimate biovolume (Yingling et al., this volume), then

converted to carbon biomass using the Menden-Deuer and Lessard (2000) relationship in

each cycle for the different groups of eukaryotes, namely picoeukaryotes: 0.32 pg C/cell

(mean biovolume of 0.8-2 μm cells), auto- and mixotrophic nanoeukaryotes: between 5 and

9 pg C cell^{-1}, and heterotrophic nanoeukaryotes: 5 pg C cell^{-1}. A carbon conversion of 5 pg C

cell^{-1} was used for heterotrophic nanoeukaryotes based on mean 3.74-μm ESD of these cells

(Yingling et al., this volume). For transit experiments (Exps 8 and 9), no data were available

from epifluorescence samples; hence, cell C contents for nanoauto- and nanomixotrophs

were inferred from FChl$a$:C ratios measured at the surface during cycles and calculated

according in proportion to those measured during transits (6 pg C cell^{-1}).

The percent contributions of microzooplankton to the daily removal of unicellular

planktonic production were calculated as:

$$\text{Microzoo production removal} = \left(\frac{m}{\mu}\right) x\ 100 \qquad \text{(Eq. 9)}$$

where m is the grazing rate (d^{-1}) by microzooplankton on a specific prey item and μ is the

growth rate (d^{-1}) of the prey type (Landry et al., this volume), calculated as:

$$\mu = k + m \qquad \text{(Eq. 10)}$$

The percent contributions of copepods to the daily removal of protist production were

calculated using clearance rate (F, L ind^{-1} d^{-1}), copepod abundance in our incubations (N, ind

L^{-1}), the abundance of each prey (Cp, cells L^{-1}), and total prey ($TotC$, cells L^{-1}):

$$\text{Copepod production removal} = \left(\frac{F\ x\ N\ x\ Cp}{TotC}\right) x\ 100 \qquad \text{(Eq. 11)}$$

To obtain independent estimates of copepod trophic reliance on auto- vs mixo-

heterotrophic diet, we estimated the pathway of C ingestion from primary producers and the

consumption of N grazers of ^{15}N-labelled *Pro* and *Syn*. The copepod biomass (W, µg ind^{-1})

and isotopic signals for δ^{13}C and δ^{15}N of both prey (*food_at*, %) and copepods (*cop_at*, %)

were provided directly from the isotope-ratio mass spectrometry analyses. The values used

for '*food_at$_{T36}$*' were the POM values taken from the copepod incubation bottles at the end of

the experiment.

The daily copepod C and N ingestion rates (IR, µg ind^{-1} d^{-1}, Eq. 12) were calculated as

the difference in copepod δ^{13}C and δ^{15}N isotopic signal between the start (T$_{24}$) and the end

(T$_{36}$) of the incubation (t, hours), and appropriately subtracting the corresponding baseline

food isotopic signal from the copepod isotopic signal (Verschoor et al., 2005). The

percentage of ingested biomass to the total copepod biomass gives the C- and N- specific

daily ration (DR, % d^{-1}, Eq. 13).

$IR_{(C,N)} = (W$ x (cop_at_{T36} / 100) - (cop_at_{T24} / 100) x W) / ($food\_at_{T36}$ / 100) / t x 24 $\qquad$ (Eq. 12)

$$DR_{(C,N)} = IR / W \ x\ 100 \qquad \text{(Eq. 13)}$$

Data were checked for normality (Shapiro-Wilk's test) and homoscedasticity (Bartlett's

test) assumptions. Differences in copepod feeding rates (clearance, ingestion) and

selectivity and microzooplankton grazing rates were tested as a function of food quality

indicators: prey trophic mode and size class. One-Way ANOVA and/or non-parametric

Kruskal-Wallis tests were applied to test a priori differences in variables between trophic

modes (autotrophic-mixotrophic-heterotrophic) while *t*-tests were used to compare size

classes (picoplankton-nanoplankton). For tests yielding significantly different variance

between trophic groups, post-hoc Tukey HSD and/or Dunn's tests were applied for multiple

comparisons. Differences were considered significant at the 0.05 level. Graphical exploration
of results and statistical analyses were performed using RStudio v4.2.

3. Results

3.1. Food landscape: biomass, size and trophic spectra of unicellular plankton

At the start of the microzooplankton incubation (T_0), Chl$a$ biomass ranged from 0.05 to
0.1 µg L^{-1}, with slightly lower concentrations for the shallower depths sampled for transit
experiments (0.05-0.06 µg Chl$a$ L^{-1}) compared to cycle experiments (0.07-0.1 µg Chl$a$ L^{-1})
(Table3). The prey community available for microzooplankton was numerically dominated by
heterotrophic bacteria (Hbact) (482,000 – 625,000 cells mL^{-1}) and *Prochlorococcus* (*Pro*)
(128,000-276,000 cells mL^{-1}; Table 3). *Synechococcus* (*Syn*) (1,260-3,460 cells mL^{-1}) had
similar abundances as nanoheterotrophic eukaryotes (817-2610 cells mL^{-1}). The remaining
eukaryotic groups represented by nanomixotrophs (28-666 cells mL^{-1}), picomixotrophs (19-
282 cells mL^{-1}), picoautotrophs (29-202 cells mL^{-1}), and nanoautotrophs (28-169 cells mL^{-1})
were much less abundant (Table3). On average, Hbact accounted for 70-80% of
microzooplankton prey, followed by *Pro* (20-30%), whereas *Syn* and all the eukaryotic
populations were negligible (<1%) in numerical terms.

In terms of carbon (C), unicellular community biomass was higher in Cycle 2 (21 µg C L^{-1}
) and Cycle 4 (24.5 µg C L^{-1}) compared to Cycle 3 and transit experiments (~17.5 µg C L^{-1})
Fig. 3A). *Pro* and nanoheterotrophs contributed most (30-40% each), nanomixotrophs and
Hbact were slightly lower (20-30% each), and *Syn*, picoautotrophs, picomixotrophs and
nanoautotrophs together comprised <10% of total biomass. Overall, unicellular heterotrophs
accounted for 58% of the C standing stock, whereas autotrophs were 18 to 32% and the
remaining 8 to 22% were mixotrophs (Fig. 3B). Prokaryotes and smaller eukaryotes
(picoplankon) were the most abundant but contributed less to total biomass (25-40%) than
nanoplankton (58-72%) (Fig. 3C).

At the start of the copepod incubation (T_{24}), Chl$a$ biomass varied between 0.03 and 0.13
µg Chl$a$ L^{-1} (Table 3). Eukaryotic groups accounted for a total biomass of 5.6-12 µg C L^{-1},
higher in Cycle 2 (12 µg C L^{-1}) and Cycle 4 (10.6 µg C L^{-1}) compared to Cycle 1, Cycle 3,
and transit experiments (5.8-7.3 µg C L^{-1}), in line with initial biomass (T_0; Fig. 3D).
Heterotrophic protists accounted for 53-79% of eukaryotic biomass, followed by mixotrophs
(12-33%), and autotrophs (3-15%) (Fig. 3E). The prey biomass available to copepods, i.e.
sampled in bottles 24 h later, was mostly nanoplankton (93-99%) rather than picoplankton
(1-7%) (Fig. 3F).

3.2. Copepod feeding on eukaryotes: ingestion, diet composition and selectivity

After the copepod incubation, Chl$a$ biomass remained stable in control bottles and

increased by ~30-60% in copepod grazing bottles (not shown); hence, grazing of copepods

based on Chl$a$ disappearance was not estimated. Total grazing on eukaryotes oscillated

between 0.43 and 1.64 d^{-1} (median= 1.31±0.23 d^{-1}; Fig. 4A), with clearance rates ranging

from 18.5 to 336 mL ind^{-1} d^{-1} (median= 80.7±55.7 mL ind^{-1} d^{-1}; Fig. 4B). Significant

differences in copepod feeding emerged in both grazing (m) and clearance (F) between

hetero-and mixotrophs (m: p= 0.0058; F: p= 0.0045) and between pico- and nano sized prey

(m:, $p<0.0001$; F: p=0.0029). Considering all prey types, between 3.4 and 138 ng C ind^{-1} d^{-1}

(median=31.4±16 ng C ind^{-1} d^{-1}) were ingested by copepods in total including

nanomixotrophs (0-12417 cells ind^{-1} d^{-1}, 0-74.5 ng C ind^{-1} d^{-1}) and nanoheterotrophs (0-5,300

cells ind^{-1} d^{-1}, 0-26.5 ng C ind^{-1} d^{-1}) at higher rates compared to picomixotrophs (356-12,400

cells ind^{-1} d^{-1}, 0.11-3.96 ng C ind^{-1} d^{-1}), picoautotrophs (390-30,200 cells ind^{-1} d^{-1}, 0.12-9.68

ng C ind^{-1} d^{-1}) and nanoautotrophs (0-339 cells ind^{-1} d^{-1}, 0-3.05 ng C ind^{-1} d^{-1}) (Fig. 4C,D).

Cell-specific ingestion increased linearly with prey abundance ($p<0.001$, R^2=0.25, not

shown), and differed significantly as functions of trophic mode (hetero-mixo, p=0.017) and

size class (p=0.006). C ingestion scaled linearly with prey biomass ($p<0.001$, R^2=0.49, not

shown) and according to prey size (p= 0.021).

Copepods consumed a diverse diet, including a considerable proportion of each group in

cell abundance terms (Fig 5A). Numerically, picoplankton cells contributed to a higher

proportion of diet than nanoplankton (Fig. 5E), but nanoplankton dominated C intake (61.7-

99.2%) (Fig. 5F). Among nanoplankton categories, C consumption derived mainly from

nanoheterotrophs (0-84.3%) and nanomixotrophs (0-81.5%), leaving smaller fractions of C

acquired from ingestion of picomixotrophs (0.4-20.8%), picoautotrophs (0.4-17.5%), and

nanoautotrophs (0-9.7%) (Fig 5B). Higher contributions of nanoheterotrophs to copepod diet

were recorded for Cycle 1,2 and 4 experiments, whereas an enhanced contribution of

nanomixotrophs to copepod diet emerged in Cycle 3 and transit experiments. Despite the

slight differences between cell-based and C-based estimates, mixotrophic (5.5-89.9%) and

heterotrophic (0-84.3%) protists accounted for the highest contributions to copepod diets

while strict autotrophs contributed substantially less (2.2-17.5%) (Fig. 5C,D).

Copepods showed neutral selection (α=~0.2) for nanomixotrophs and picoautotrophs,

while nanoheterotrophs, nanoautotrophs and picomixotrophs tended toward negative

selection ($\alpha<0.2$, Fig. 6A). Overall, however, no significant differences were found in

selectivity among prey groups (p=0.894). Autotrophs were ingested in proportion to their

availability (E=+0.01), heterotrophs were neglected (-1) and mixotrophs were slightly

preferred (+0.31; Fig. 6B), but these differences among trophic categories were not

significant due to high data variability (p=0.289). Positive selection was suggested for

picoplankton (+0.3) at the expense of nanoplankton (-0.05), but this was not significant

(p=0.563; Fig 6C).

The daily total C ingestion rates of copepods based on bottle-incubation disappearance

estimates ranged between 0.02 and 0.1 µg C ind^{-1} d^{-1} (equivalent to 20-100 ng C ind^{-1} d^{-1};

Fig. 7A). The daily rations were generally very low, ranging between 0.05 and 0.34% d^{-1}

(Fig. 7B). In contrast, estimating consumption of POM using ^{13}C as a tracer yielded C-

specific ingestion rates (Fig. 7C) and daily rations (Fig. 7D) that were one order of magnitude

higher compared to those obtained using prey abundances analysed by flow cytometry.

These ranged from 0.25 to 0.42 µg C ind^{-1} d^{-1} (250-420 ng C ind^{-1} d^{-1}), which represented

0.8-2.4% d^{-1} of the copepod C biomass. Estimates of N ingestion through ^{15}N-labelled prey

ranged 0.01-0.07 µg N ind^{-1} d^{-1}, (10-70 ng N ind^{-1} d^{-1}) with higher values observed in Cycle

3, 4 and transit experiments (Fig. 7E) when the larger labelled prey (*Syn*) was used. N daily

rations accounted for 0.25-1.0% d^{-1} (Fig. 7F). The daily ration for C (% body carbon) and N

(% body nitrogen) based on the isotope tracers was ~1% for both elements during Cycle 3,

4, and transits (Fig. 7D, F).

3.3. Microzooplankton grazing on prokaryotes and eukaryotes

Microzooplankton grazed variably on Chl*a* (0.34-0.66 d^{-1}) with lower values in Cycles 1

and 4 compared to the other experiments but was not significantly different throughout the

cruise (p=0.195; Fig. 8A). Among prokaryotes, microzooplankton grazed similarly on *Pro*

(0.36-2.14 d^{-1}), *Syn* (0.36-1.26 d^{-1}) and Hbact (0.48-1.46 d^{-1}; Fig. 8B). Among eukaryotes,

nanoheterotrophs were the most grazed prey (0.92-2.48 d^{-1}), especially in Cycle 3 and along

transits, followed by nanomixotrophs (0.48-1.24 d^{-1}), nanoautotrophs (0.18-1.02 d^{-1}; Fig. 8C),

picoautotrophs (0.06-1.51 d^{-1}) and picomixotrophs (0.08-0.93 d^{-1}). Considering only

eukaryotes, microzooplankton grazed on heterotrophs at significantly higher rates compared

to autotrophs (p=0.0002), and mixotrophs (p<0.0001). Similarly, picoeukaryotes were grazed

at significantly lower rates compared to nanoeukaryotes (p<0.0001). Considering both

prokaryotes and eukaryotes, grazed biomass did not differ significantly between trophic

modes (p=0.927), but it did between size classes (p=0.0002). The highest contribution to diet

consisted mostly of nanoeukaryotes, namely nanoheterotrophs (24.7-58.6%),

nanomixotrophs (13.8-40.1%) and nanoautotrophs (7.5-32.9%), whereas the other

prokaryotic and eukaryotic prey accounted for <7% of the food intake (Fig. 8D). Out of the

total grazed biomass, 24.8-58.7% was heterotrophic, 13.9-41.7% was mixotrophic, and a

lower fraction (12.1-33.5%) originated from autotrophic nutrition (Fig. 8E). As the cruise

progressed, increased heterotrophic relative dietary contributions were accompanied by a

relative decreases in autotrophic diet, whereas the contribution of mixotrophs did not show a

clear pattern. Nanoplankton contributed the highest microzooplankton food intake (95.1-

97.9%) as opposed to picoplankton (2.1-4.9%) (Fig. 8F).

3.4. Daily removal of unicellular plankton production

On average, 110% of the daily Chl*a* production was removed by microzooplankton (Fig.

9A), and 104%, 149% and 125% of the daily production of Hbact, *Pro* and *Syn* were

consumed, respectively. Among eukaryotes, microzooplankton contributed to the high

removal of nanoheterotrophs (149%), picoautotrophs (109%), nanoautotrophs (128%).

Between 96% (pico) and 76% (nano) of daily mixotrophic production was removed by

microzooplankton. Among eukaryotes, the highest copepod grazing impact was exerted on

picoautotrophs (13.8%), followed by picomixotrophs (9.6%), nanoheterotrophs (7.6%),

nanomixotrophs (3.9%), while production of nanoautotrophs was the least impacted by

copepod grazing (<0.1%; Fig. 9B).

4. Discussion

4.1. Community biomass, size classes and trophic modes

The Chl*a* and microbial standing stock estimates in our incubations align with values

calculated for the upper euphotic zone during the same cruise (Landry et al., this volume), as

well as those from other oligotrophic regions (Landry and Kirchman, 2002; Paffenhöfer et al.,

2007)*.* The low Chl*a* biomass reflected scarce concentrations of inorganic nutrients in the

upper 50 m (NO_3^{2-}: <0.01-0.03 µM, PO_4^{3-}: ~0.03-0.12 µM; Kranz et al., this volume). In the

eastern Indian Ocean, a moderate fraction of C fixation can, in fact, originate from

diazotrophy (N_2 fixation) (Kranz et al., this volume; Nieblas et al. 2014; Sato et al. 2022), and

lateral advection of nutrients (Kehinde et al., 2023).

Prokaryotes comprised almost 50% of microbial community carbon (C), and *Pro* was

responsible for the Chl*a* biomass change observed across cycles and stations (linear

regression, *p*= 0.0226*,* not shown*).* Heterotrophic and mixotrophic nanoplankton dominated

in abundance and biomass for the eukaryotic component. Given the small size and surface-

to-volume ratio of the dominant cells and the prevalence of heterotrophs *vs* autotrophs, it is

reasonable to think that these groups were selected for in these conditions (Chisholm,

1992), in contrast to productive upwelling areas dominated by larger autotrophs (e.g.

diatoms). Mixotrophs and heterotrophs were key players in the eukaryotic community (Selph

et al., this volume) given the scarcity of inorganic nutrients and the rich prey size spectrum

available (from bacteria to nanoplankton). Nutrient shortage is known to favor mixotrophy in

marine plankton (Christaki et al., 1999; Hartmann et al., 2012; Unrein et al., 2014), enabling

them to acquire organic nutrients through direct predation, symbiosis and kleptoplastidy

under conditions where obligate autotrophs would be limited (Flynn et al., 2019). The

abundant and diversified prey size spectrum could explain the higher abundance of

nanoheterotrophs over mixotrophs, with their mainly higher ingestion and growth rates

compared to non-specialist phagotrophs (Edwards et al., 2023; Jeong et al., 2005).

4.2 Copepod feeding: limitation by food quantity, quality and trophic cascades

Copepods cleared on average 75 mL ind^{-1} d^{-1}, and ingested between 3.4 and 138 ng C

ind^{-1} d^{-1} (median: 32.9 ng C ind^{-1} d^{-1}), in agreement with field studies on copepod feeding on

a natural plankton assemblage in oligotrophic conditions (2-400 mL ind^{-1} d^{-1}, 10-200 ng C

ind^{-1} d^{-1}) (Broglio et al., 2004; Calbet and Saiz, 2005). The entire bottle volumes (108%)

were cleared by the copepods, indicating perhaps that prey availability was not sufficient to

sustain copepod demand. Occasionally, gut Chl*a* was also determined for the copepods

(Exps 2, 8, and 9) and accounted for an equivalent of 0.8±0.4 µg Chl*a* ind^{-1} d^{-1} (for full

description of the methodological procedure, see Décima et al., this volume). Hence,

although our food removal method could not quantify copepod Chl*a* grazing, gut content

data suggested recent ingestion equivalent to 32 ng C ind^{-1} d^{-1}, when applying a C:Chl=58

as done for the integrated mesozooplankton community (Décima et al., this volume).

Interestingly, some pelagic tunicates (appendicularians) were present in the incubation

water. Copepods may have gained indirect access to Chl*a* especially from picoplankton

trapped in the appendicularians' mucous houses, given the ability of certain copepods to

directly feed on them (Ohtsuka et al., 1996). Community grazing was also assessed via Chl*a*

gut content analyses of size-fractionated net collections during the cruise and showed

positive ingestion of Chl*a* by mesozooplankton (>0.2 mm) and up to 8% removal of Chl*a*

production, though the highest Chl*a* grazing impact was exerted by zooplankton size

fractions <1 and <0.5 mm (Décima et al., this volume).

It is possible that copepods fed on different prey that were not explicitly accounted for,

such as large dinoflagellates and ciliates, known to comprise on average 33% and 43% of

the diet in copepods in low nutrient environments (Kleppel, 1993; Saiz and Calbet, 2011).

The highest contributions to dietary C were from nanomixotrophs and nanoheterotrophs,

which are normally dinoflagellates and ciliate groups, preferred prey for copepods given their

nutritional value and detectability (Saiz and Calbet, 2011). Other experiments also report

high grazing and enhanced selection on heterotrophic protists, but only a minimal impact on

total production (2%) (Broglio et al., 2004), consistent with our observed nanoheterotroph

removal rates (7.6%). It may also be true that larger prey might have contributed to high but

undetected C intake. In fact, the dominant copepods sorted for incubation were mostly

carnivores/omnivores and detritivores (*Candacia* spp., *Corycaeus* spp., *Euchaeta* spp.,

Labidocera spp., *Oncaea* spp., *Sapphirina* spp.), which preferentially feed on larger prey

such as microplanktonic dinoflagellates, ciliates, copepod nauplii and marine snow (Atienza

et al., 2006; Landry, 1978; Yen, 1985). Since larger motile cells generate a stronger

hydrodynamic disturbance making them easier to detect by rheotactic predators (Almeda et

al., 2018), it is likely that pico-nanoplankton size might have been suboptimal for efficient

capture by these copepods (~1.5 mm prosome length) (Hansen et al., 1994).

However, bottle incubations are known to provide underestimates of absolute quantities

of C consumed (Zeldis and Décima, 2020) and their best application is to investigate

selectivity. The underestimate is caused at least in part by the inability to replicate the local

concentrations of prey encountered by a copepod. The ingestion calculation includes a

clearance rate term (F, ml ind^{-1} d^{-1}) and a concentration (cells mL^{-1}), which is typically taken

as the average concentration of the incubation or Niskin bottle, but likely does not represent

small scale patchiness or thin layers exploited by copepods in the field to obtain their daily

ration (Mullin and Brooks, 1976). In addition, low prey concentrations can be easily

exhausted over the course of an experiment (Zeldis and Décima, 2020). However, when

investigating selectivity of many types of prey, experiments need to be run long enough to

detect the decrease or depletion of different preferred prey types. Thus, quantifying absolute

ingestion and selectivity within a bottle experiment is usually not possible given the trade-offs

described above, in addition to the inherent limitation of a bottle experiments not replicating

the local prey concentration patchiness.

Copepods exhibit reduced selectivity under food limitation to increase the chance of food

intake (Vincent and Hartmann, 2001). This and the high clearance rate observed might

explain the absence of a clear preference for any of the five specific prey groups analyzed in

our study, presumably in favor of other prey as previously pointed out. Though not

significant, a moderate overall selection for mixotrophs emerged in parallel with avoidance of

auto- and heterotrophs despite their abundance. Laboratory studies report no strong

copepod preference based on trophic guild or nutritional value of their prey (Isari et al., 2013;

Traboni et al., 2020). On the other hand, ingestion of phagotrophs can provide nitrogen,

phosphorus and essential fatty acids, increasing the food quality for copepods (Klein Breteler

et al., 1999) and enhancing the reproductive yield (Kleppel, 1993). Surprisingly, in our

incubations, copepods seem to prefer and impact smaller picoplankton over nanoplankton.

Compensatory feeding mechanisms might have led copepods to feed even on smaller prey

to buffer the limiting food quantity (Burian et al., 2018). In a subtropical ecosystem,

copepods fed natural nanoplankton (>2 μm, 2-5 μm, >5 μm) showed similar patterns of

selection towards the smallest prey size class, ascribed to the dominance of picoplankton

over other prey types, inducing copepods to maximize the intake of the most available prey

(Calbet et al., 2000). Nevertheless, considering the large body size and the hunting behavior

of the copepods in our experiments, active selection of picoplankton seems unlikely and our

observations more likely the result of trophic cascades. This may have been triggered by

microzooplankton preying on nanoplankton and releasing picoplankton from grazing

pressure, which then became more abundant and more detectable and ingested by

copepods. Appendicularians can compete with copepods for food and thus might have

contributed to internal feeding together with protistan and metazoan micrograzers

(Alldredge, 1984; Gorsky and Fenaux, 1998). As copepods ingested microzooplankton, the

grazing pressure on Chl*a* standing stock would be released. Notably, Chl*a* concentration

increased in copepod bottles by 30-60% compared to controls rather than declining due to

grazing, supporting in part this cascade effect. However, one needs also to consider that by

the end of the incubation (T_{36}), the Chl*a* content of *Pro* cells (from flow cytometric

measurements) increased (~35%) significantly in the presence of copepods compared to

bottles without copepods (p=0.037) in several experiments, suggesting that phototrophs

might have benefitted from nutrients excreted by copepods, masking any copepod grazing

impact on Chl*a*.

4.3. Microzooplankton grazing impact exceeds that of copepods

Microzooplankton were the dominant consumers of the microbial community, removing

on average 110% of Chl*a* standing stock d^{-1}, 126% of bacterial production and 111% of

eukaryotic production. Grazing on prokaryotes removed 12% more production as on

eukaryotes, less than the 33% difference reported by Landry et al. (this volume) from in situ

incubations in the upper euphotic zone. Our estimates differ as i) *Syn* and nanoheterotrophs

were included only in this study, ii) our sampling was depth-specific, whereas Landry et al.

(this volume) sampled across a vertical gradient encompassing a wider range of conditions

and determined rates over many more experiments, and iii) these incubations were done on

deckboard incubators as opposed to in situ. Overall, our results align with the 40-100%

range of daily primary production removal by microzooplankton in tropical and subtropical

regions, including the Indian Ocean (Landry, 2009; Landry et al., 2022; Schmoker et al.,

2013).

As expected for an oligotrophic ecosystem, copepods were less important grazers,

removing on average 7% of protistan pico-nanoplankton production within the 2-12 µm size

range. Consequently, the top-down control of eukaryotic production alone depended very

little on copepods and was mostly driven by microzooplankton. Despite differences in

grazing effort, both zooplankton groups consumed a high proportion of nanoheterotrophs,

but copepods relied on nanomixotrophs to a higher extent.

4.4 Mixotrophy and intraguild predation: hidden pathways in oligotrophic food webs

Mixotrophs were important contributors to eukaryotic biomass. FlowCam images from

the cruise detected putatively mixotrophic dinoflagellates (*Prorocentrum* spp., *Tripos* spp.)

and ciliates (*Tontonia* spp.) (J. Goes, pers. comm.). A recent study in the same Indian

Ocean region found rhizarians, including mixotrophic representatives, to be dominant

together with copepods (Davies et al., 2022). Despite the absence of taxonomic detail for the

mixotrophic and heterotrophic community in these experiments, LysoTracker Green coupled

to flow cytometry (Selph et al., this volume) and previous community analysis (Davies et al.,

2022) validate observations that mixotrophy is a widespread strategy spanning different

protistan groups in this region, consistent with their well-known eco-evolutionary advantages

in low-nutrient waters (Stoecker et al., 2017). Molecular analyses of microbial plankton from

the eastern Indian Ocean have previously pointed to the elevated proportions of mixotrophic

taxa, relative to declining contributions of true autotrophs and heterotrophs, in warm tropical

waters that come from our study region (Raes et al., 2022). This also supports the

hypothesis based on mesozooplankton grazing rates that the lower food web of these

tropical waters may allow a more efficient trophic coupling to higher levels due to the high

levels of mixotrophy (Landry et al., 2020a). While nanoheterotrophs contributed equally to

what microzooplankton and copepod consumed in these experiments, nanomixotrophs were

especially important for copepods. Phagotrophic protists provide important essential

elements and can enhance nutrient transfer across trophic levels when ingested.

Copepod C ingestion rates estimated through the disappearance method were

approximately an order of magnitude lower compared to those attained using stable isotope

labelling. We can speculate on a few reasons for this discrepancy. From an ecological

standpoint, we know that flow cytometry does not quantify the larger prey and consequently

the disappearance estimates preclude the inclusion of their contribution to copepod diet.

However, the contribution of microplankton to the overall plankton assemblage was very low

(Yingling et al., this volume), such that it's not likely that high enough abundances were

present in our bottles to account for this difference. From a methodological standpoint, we

used the δ^{13}C values of POM as representative of what copepods were consuming.

However, since POM includes all prey items and detritus (Stukel et al., 2014), it's unlikely

that copepods feed on all these components equally since copepods are highly selective

(Broglio et al., 2004; Isari and Saiz, 2011). If the prey items consumed by copepods were

small primary producers (with higher δ^{13}C values than POM because they are actively

incorporating this tracer), then we have overestimated C-incorporation by using the values of

POM. If the prey items consumed by copepods had lower δ^{13}C values than POM because

they were many trophic steps removed from primary producers, then our calculation

underestimates C-consumption by copepods. The mean δ^{13}C-based estimated C daily

rations were 1.7% d^{-1}, in the range of the estimates calculated for prey <15 μm in calanoid

and cyclopoid copepods (Isari and Saiz, 2011; Schnetzer and Caron, 2005), but not

sufficient to sustain basic respiratory and production rates at these temperatures (Décima et

al. this volume; Landry et al., 2020b). However, these estimates are expected given the

method, because bottles do not contain the prey densities to support the ingestion needed

for respiration and production over 12 hours, as discussed above, but are in line with

previous similar experiments (Isari and Saiz, 2011; Schnetzer and Caron, 2005).

When investigating patterns in the incorporation of N using the ^{15}N-labelled prey, we

cannot compare equally across cycles given the methodological differences associated with

using *Syn* and *Pro*. However, we can compare patterns in ^{15}N incorporation from Cycle 1

and 2 as representative of phagotrophic pathways connecting copepods to *Pro*, and Cycles

3, 4 and the transect, as representative of phagotrophic pathways connecting copepods to

larger-sized prey similar to *Syn* (because *Syn* was not present in high abundances in these

waters dominated by *Pro*). N daily rations were similar to C daily rations when we used *Syn*

(both C and N approximately 1%), but was much lower when using *Pro* (1.8% C compared

to 0.3% N). The higher N daily rations observed in *Syn* experiments point to a more efficient

trophic pathway via the consumption of mixo- and hetero- trophs (nano- and micro- sized),

as confirmed by food removal incubations, that had more likely fed directly on ^{15}N-labelled

Syn due to its larger size (>1 μm). This highlights their presumptive role in nutrient

channeling and trophic upgrading as previously reported for protozooplankton and more

recently for mixoplankton (Klein Breteler et al., 1999; Ptacnik et al., 2004; Traboni et al.,

2021). Selph and colleagues highlighted that mixotrophs were correlated with *Syn* during the

cruise (Selph et al., this volume), contributing to a high N transfer from ^{15}N-*Syn* to copepods.

We know that copepods did not feed directly on ^{15}N-labelled *Syn* cells because the ^{15}N-

labelled prey had already been consumed by the time copepods were inoculated into the

experimental bottles at T_{24}. We cannot rule out that copepods may have acquired $^{1}\square$N by

ingesting clumps of labelled cells that did not resuspend as single cells and/or that had been

concentrated by appendicularians. However, appendicularians need to be collected very

gently in order for them to exhibit natural behavior in containers (Masunaga et al., 2020;

Scheinberg et al., 2005), which we did not do since they would have been transferred to our

bottles directly from the Niskin. And while their abundances where high in our study, the

highest concentrations were 200 individuals m^{-3} (Davies et al., this volume), which means

that on average there was zero or 1 appendicularian per bottle (0.54 individuals in 2.7L),

unlikely to significantly affect these results. Carnivory of copepods on younger stages such

as copepod nauplii and copepodites, which can feed on much smaller prey items than adult

copepods, is potentially more likely given their higher concentrations (Davies et al., this

volume; Swalethorp et al., this volume), and the inclusion of omnivorous/carnivorous taxa in
our incubations. This could also explain the discrepancy in estimated C and N daily rations
based on POM (which would include nauplii caught on the GF/F) and based on flow
cytometry, which only includes direct consumption of unicellular prey.

In oligotrophic regions dominated by small-size cells, the ability of appendicularians to
concentrate bacteria, picoplankton, and detritus into mucilaginous houses that they regularly
abandon provides a major mechanism for transferring seemingly inaccessible nutrition to
copepods. Copepods of the genus *Oncaea* are known to feed on marine snow and
appendicularian houses (Ohtsuka et al., 1996) and were rather common in the Argo Basin
community (Davies et al., this volume; Swalethorp et al., this volume). Appendicularians
were also highly selected by *T. maccoyii* larvae during this cruise and accounted for up to
75% of their prey diet (Swalethorp et al., this volume), highlighting the role of gelatinous
zooplankton in ichthyoplankton nutrition (Laiz-Carrion et al., 2015; Mianzan et al., 2001) and
in nutrient transfer throughout the food web.

5. Summary and conclusions

Copepods grazed a small fraction of daily production, while microzooplankton are the
main players in top-down control of protistan biomass in the Argo Basin. Mixotrophs and
heterotrophs emerged as dominant functional groups in these oligotrophic food webs,
supporting copepod and microzooplankton diets that channeled nutrients to higher trophic
levels. Isotope tracing using labeled ^{15}N-prey confirmed the importance of copepod
consumption of phagotrophs and/or other metazoans, which was much more efficient at
transferring energy from cells 1-2 µm compared to the <1µm-sized dominant
Prochlorococcus. With ocean warming, smaller cell sizes (Morán et al., 2010) and increased
phagotrophy (Wilken et al., 2013) are expected, making mixotrophy a competitive strategy
for protist survival and increasing efficiency of energy transfer across trophic levels (Ward
and Follows, 2016). This may help counteract the declining food quality predicted under
nutrient limitation as a consequence of water-column stratification (Schenone et al., 2024). In
regions dominated by small cells of poor stoichiometric and biochemical quality, copepod
nutrition may strongly rely on larger more nutritional prey, and benefit from the nutrient
channeling and buffering mediated by microzooplankton and mixotrophs. The statistical
limitations of this study highlight the need for further investigations of prey diversity and
selection towards or against specific taxonomic groups that encompass a larger size
spectrum than the one examined herein.

Declaration of competing interest

The authors declare no competing financial nor personal interests.

Author statement

M.D. and C.T. conceived the study. C.T., G.F.C., K.E.S. and M.D. conducted sampling, experiments and sample processing onboard. K.E.S. processed flow cytometry data, G.F.C. processed samples for elemental content estimates. M.D. supported with analysis and interpretation of isotope data. K.E.S and M.R.L. provided cell biomass data, and support for interpretation of the results. C.T. analyzed results, conducted statistical analyses and drafted the manuscript. All authors contributed to provide conceptual feedbacks, comments, edits and thorough review of the manuscript.

Acknowledgements

We are grateful to captain and crew of R/V *Roger Revelle* and to all the RR2201 science party. We thank N. Yingling and M. Stukel for providing stable isotope stocks, R. Swalethorp and C. Davies for assistance during zooplankton sampling, and J. Goes for providing FlowCam images of the protistan community. This study was supported by U. S. National Science Foundation grants OCE-1851558 (M.R.L.) and -1851381 (K.E.S.) and is a contribution to the Second International Indian Ocean Expedition (IIOE-2 endorsed project EP046). Seawater and plankton samples were collected under Australian Government permit AU-COM2021-520 and Australian Marine Parks permit PA2021-00062-2 issued by the Director of National Parks, Australia. Views expressed in this publication do not necessarily represent those of the Director of National Parks or the Australian Government.

References

Alldredge, A.L., 1984. The quantitative significance of gelatinous zooplankton as pelagic consumers, in: Flows of Energy and Materials in Marine Ecosystems. Springer US, Boston, MA, pp. 407–433. https://doi.org/10.1007/978-1-4757-0387-0_16

Almeda, R., van Someren Gréve, H., Kiørboe, T., 2018. Prey perception mechanism determines maximum clearance rates of planktonic copepods. Limnol. Oceanogr. 63, 2695–2707. https://doi.org/10.1002/lno.10969

Anschütz, A., Maselli, M., Traboni, C., Boon, A.R., Stolte, W., 2024. Importance of integrating mixoplankton into marine ecosystem policy and management—Examples from the Marine Strategy Framework Directive. Integr. Environ. Assess. Manag. 20, 1366–1383. https://doi.org/10.1002/ieam.4914

Arenovski, A.L., Lim, E.L., Caron, D.A., 1995. Mixotrophic nanoplankton in oligotrophic
surface waters of the Sargasso Sea may employ phagotrophy to obtain major nutrients.
J. Plankton Res. 17, 801–820. https://doi.org/10.1093/plankt/17.4.801
Atienza, D., Calbet, A., Saiz, E., Alcaraz, M., Trepat, I., 2006. Trophic impact, metabolism,
and biogeochemical role of the marine cladoceran *Penilia avirostris* and the co-
dominant copepod *Oithona nana* in NW Mediterranean coastal waters. Mar. Biol. 150,
221–235. https://doi.org/10.1007/s00227-006-0351-z
Broglio, E., Saiz, E., Calbet, A., Trepat, I., Alcaraz, M., 2004. Trophic impact and prey
selection by crustacean zooplankton on the microbial communities of an oligotrophic
coastal area (NW Mediterranean Sea). Aquat. Microb. Ecol. 35, 65–78.
https://doi.org/10.3354/ame035065
Burian, A., Grosse, J., Winder, M., Boschker, H.T.S., 2018. Nutrient deficiencies and the
restriction of compensatory mechanisms in copepods. Funct. Ecol. 32, 636–647.
https://doi.org/10.1111/1365-2435.13016
Calbet, A., Landry, M.R., Scheinberg, R.D., 2000. Copepod grazing in a subtropical bay:
species-specific responses to a midsummer increase in nanoplankton standing stock.
Mar. Ecol. Prog. Ser. 193, 75–84. https://doi.org/10.3354/meps193075
Calbet, A., Saiz, E., 2005. The ciliate-copepod link in marine ecosystems. Aquat. Microb.
Ecol. 38, 157–167. https://doi.org/10.3354/ame038157
Castellani, C., Irigoien, X., Harris, R.P., Lampitt, R.S., 2005. Feeding and egg production of
Oithona similis in the North Atlantic. Mar. Ecol. Prog. Ser. 288, 173–182.
https://doi.org/10.3354/meps288173
Chesson, J., 1978. Measuring preference in selective predation. Ecology 59, 211–215.
Chisholm, S.W., 1992. Phytoplankton size, in: Primary Productivity and Biogeochemical
Cycles in the Sea. Environmental Science Research. pp. 213–237.
https://doi.org/10.1007/978-1-4899-0762-2_12
Christaki, U., Van Wambeke, F., Dolan, J.R., 1999. Nanoflagellates (mixotrophs,
heterotrophs and autotrophs) in the oligotrophic eastern Mediterranean: standing
stocks, bacterivory and relationships with bacterial production. Mar. Ecol. Prog. Ser.
181, 297–307. https://doi.org/10.3354/meps181297
Davies, C.H., Beckley, L.E., Richardson, A.J., 2022. Copepods and mixotrophic Rhizaria
dominate zooplankton abundances in the oligotrophic Indian Ocean. Deep-Sea. Res. II.
202, 105136. https://doi.org/10.1016/j.dsr2.2022.105136
Davies, C.H., Eriksen, R.S., Jayasinghe, S., Landry, M.R. This issue. Plankton assemblages
across a coastal to oceanic gradient of Northern Australia and responses to a storm
perturbation in the Argo Basin. Deep-Sea Res. II.

Davis, T.L.O., Lyne, V., Jenkins, G.P., 1991. Advection, dispersion and mortality of a patch
of Southern Bluefin Tuna larvae *Thunnus maccoyii* in the East Indian Ocean. Mar. Ecol.
Prog. Ser. 73, 33–45. https://doi.org/10.3354/meps073033

Décima, M., Swalethorpe, R., Cawley, G., Traboni, C., Davies, C., Landry, M.R. This issue.
Zooplankton trophic processes in the Eastern Indian Ocean. Deep-Sea Res. II.

Edwards, K.F., 2019. Mixotrophy in nanoflagellates across environmental gradients in the
ocean. Proc. Natl. Acad. Sci. U. S. A. 116, 6211–6220.
https://doi.org/10.1073/pnas.1814860116

Edwards, K. F., Li, Q., Steward, G. F., 2023. Ingestion kinetics of mixotrophic and
heterotrophic flagellates. Limnol. Oceanogr. 68, 917–927.
https://doi.org/10.1002/lno.12320

Fileman, E.S., Smith, T., Harris, R., 2007. Grazing by *Calanus helgolandicus* and *Para-*
Pseudocalanus spp. on phytoplankton and protozooplankton during the spring bloom in
the Celtic Sea. J. Exp. Mar. Bio. Ecol. 348, 70–84.
https://doi.org/10.1016/j.jembe.2007.04.003

Flynn, K.J., Mitra, A., Anestis, K., Anschütz, A.A., Calbet, A., Ferreira, G.D., Gypens, N.,
Hansen, P.J., John, U., Martin, J.L., Mansour, J.S., Maselli, M., Medić, N., Norlin, A.,
Not, F., Pitta, P., Romano, F., Saiz, E., Schneider, L.K., Stolte, W., Traboni, C., 2019.
Mixotrophic protists and a new paradigm for marine ecology: where does plankton
research go now? J. Plankton Res. 41, 375–391. https://doi.org/10.1093/plankt/fbz026

Frost, B.W., 1972. Effects of size and concentration of food particles on the feeding behavior
of the marine planktonic copepod *Calanus pacificus*. Limnol. Oceanogr. 17, 805–815.
https://doi.org/10.4319/lo.1972.17.6.0805

Garrison, D.L., Gowing, M.M., Hughes, M.P., Campbell, L., Caron, D.A., Dennett, M.R.,
Shalapyonok, A., Olson, R.J., Landry, M.R., Brown, S.L., Liu, H. Bin, Azam, F.,
Steward, G.F., Ducklow, H.W., Smith, D.C., 2000. Microbial food web structure in the
Arabian Sea: A US JGOFS study. Deep-Sea Res. II. 47, 1387–1422.
https://doi.org/10.1016/S0967-0645(99)00148-4

Gorsky, G., Fenaux, R., 1998. The role of Appendicularia in marine food webs, in: The
Biology of Pelagic Tunicates. Oxford University PressOxford, pp. 161–170.
https://doi.org/10.1093/oso/9780198540243.003.0010

Gundersen, K., Heldal, M., Norland, S., Purdie, D.A., Knap, A.H., 2002. Elemental C, N, and
P cell content of individual bacteria collected at the Bermuda Atlantic Time-series Study
(BATS) site. Limnol. Oceanogr. 47, 1525–1530.
https://doi.org/10.4319/lo.2002.47.5.1525

Hansen, B., Bjornsen, P.K., Hansen, P.J., 1994. The size ratio between planktonic predators

and their prey. Limnol. Oceanogr. 39, 395–403.

https://doi.org/10.4319/lo.1994.39.2.0395

Hartmann, M., Grob, C., Tarran, G.A., Martin, A.P., Burkill, P.H., Scanlan, D.J., Zubkov, M.

V., 2012. Mixotrophic basis of Atlantic oligotrophic ecosystems. Proc. Natl. Acad. Sci.

U. S. A. 109, 5756–5760. https://doi.org/10.1073/pnas.1118179109

Isari, S., Saiz, E., 2011. Feeding performance of the copepod *Clausocalanus lividus* (Frost

and Fleminger, 1968). J. Plankton Res. 33, 715–728.

https://doi.org/10.1093/plankt/fbq149

Isari, S., Antó, M., Saiz, E., 2013. Copepod foraging on the basis of food nutritional quality:

can copepods really choose? PLoS One 8, 1–12.

https://doi.org/10.1371/journal.pone.0084742

Ivlev, V.S., 1961. Experimental ecology of the feeding of fishes. Yale University Press, New

Haven.

Jeong, H. J., Yoo, Y. D., Kim, J. S., Seong, K. A., Kang, N. S., Kim, T. H., 2010. Growth,

feeding and ecological roles of the mixotrophic and heterotrophic dinoflagellates in

marine planktonic food webs. Ocean Sci. J. 45, 65–91. https://doi.org/10.1007/s12601-

010-0007-2

Jeong, H.J., Yoo, Y. Du, Kim, J.S., Seong, K.A., Kang, N.S., Kim, T.H., 2010. Growth,

feeding and ecological roles of the mixotrophic and heterotrophic dinoflagellates in

marine planktonic food webs. Ocean Sci. J. 45, 65–91. https://doi.org/10.1007/s12601-

010-0007-2

Jonsson, P., Tiselius, P., 1990. Feeding behaviour, prey detection and capture efficiency of

the copepod *Acartia tonsa* feeding on planktonic ciliates. Mar. Ecol. Prog. Ser. 60, 35–

44. https://doi.org/10.3354/meps060035

Kehinde, O., Bourassa, M., Kranz, S., Landry, M.R., Kelly, T., Stukel, M.R., 2023. Lateral

advection of particulate organic matter in the eastern Indian Ocean. J. Geophys. Res.

Ocean. 128. https://doi.org/10.1029/2023JC019723

Klein Breteler, W.C.M., Schogt, N., Baas, M., Schouten, S., Kraay, G.W., 1999. Trophic

upgrading of food quality by protozoans enhancing copepod growth: role of essential

lipids. Mar. Biol. 135, 191–198. https://doi.org/10.1007/s002270050616

Kleppel, G.S., 1993. On the diets of calanoid copepods. Mar. Ecol. Prog. Ser. 99, 183–195.

https://doi.org/10.3354/meps099183

Kranz S.A., Rose, J., Stukel, M., Selph, K.E., Yingling, N., Landry, M.R. This issue.

Characterization of marine primary productivity and N_2-fixation in tropical waters of the

bluefin tuna spawning region off northwestern Australia. Deep-Sea Res. II.

Laiz-Carrion, R., Gerard, T., Uriarte, A., Malca, E., Quintanilla, J.M., Muhling, B.A., Alemany,

F., Privoznik, S.L., Shiroza, A., Lamkin, J.T., Garcia, A., 2015. Trophic ecology of

Atlantic Bluefin Tuna (*Thunnus thynnus*) larvae from the Gulf of Mexico and NW

Mediterranean spawning grounds: a comparative stable isotope study. PLoS One 10,

1–22. https://doi.org/10.1371/journal.pone.0133406

Landry, M.R., 2009. Grazing processes and secondary production in the Arabian Sea: a

simple food web synthesis with measurement constraints, in: Indian Ocean

Biogeochemical Processes and Ecological Variability. American Geophysical Union, pp.

133–146. https://doi.org/10.1029/2008GM000781

Landry, M.R., 1978. Predatory feeding behavior of a marine copepod, *Labidocera trispinosa.*

Limnol. Oceanogr. 23, 1103–1113. https://doi.org/10.4319/lo.1978.23.6.1103

Landry, M.R., Brown, S.L., Rii, Y.M., Selph, K.E., Bidigare, R.R., Yang E.J., Simmons, M.P.,

2008. Depth-stratified phytoplankton dynamics in Cyclone *Opal,* a subtropical mesoscale

eddy. Deep-Sea Res. II 55, 1348–1359. https://doi.org/10.1016/j.dsr2.2008.02.001

Landry, M.R., Hood, R.R., Davies, C.H. 2020a. Mesozooplankton biomass and temperature-

enhanced grazing along a 110°E transect in the eastern Indian Ocean. Mar. Ecol. Prog.

Ser. 649, 1-19. https://doi.org/10.3354/meps13444

Landry, M., Stukel, M., Décima, M., 2020b. Food-web fluxes support high rates of

mesozooplankton respiration and production in the equatorial Pacific. Mar. Ecol. Prog.

Ser. 652, 15–32. https://doi.org/10.3354/meps13479

Landry, M.R., Hood, R.R., Davies, C.H., Selph, K.E., Antoine, D., Carl, M., Beckley L.E.,

2022. Microbial community biomass, production and grazing along 110°E in the eastern

Indian Ocean. Deep-Sea Res. II 202, 105134.

https://doi.org/10.1016/j.dsr2.2022.105134

Landry, M.R., Kirchman, D.L., 2002. Microbial community structure and variability in the

tropical Pacific. Deep-Sea Res. II. 49, 2669–2693. https://doi.org/10.1016/S0967-

0645(02)00053-X

Landry, M.R., Ohman, M.D., Goericke, R., Stukel, M.R., Tsyrklevich, K., 2009. Lagrangian

studies of phytoplankton growth and grazing relationships in a coastal upwelling

ecosystem off Southern California. Prog. Oceanogr. 83, 208–216.

https://doi.org/10.1016/j.pocean.2009.07.026

Landry, M.R., Stukel, M.R., Yingling, N., Selph, K.E., Kranz, S.A., Fender, C.K., Swalethorp,

R., Bhabu, R.I., This issue. Microbial food web dynamics in tropical waters of the bluefin

tuna spawning region off northwestern Australia. Deep-Sea Res. II.

Li, Q., Dong, K., Wang, Y., Edwards, K.F., 2024. Relative importance of bacterivorous

mixotrophs in an estuary-coast environment. Limnol. Oceanogr. Lett. 9, 81–91.

https://doi.org/10.1002/lol2.10362

Malzahn, A.M., Aberle, N., Clemmesen, C., Boersma, M., 2007. Nutrient limitation of primary

producers affects planktivorous fish condition. Limnol. Oceanogr. 52, 2062–2071.

https://doi.org/10.4319/lo.2007.52.5.2062

Masunaga, A., Liu, A. W., Tan, Y., Scott, A., Luscombe, N. M., 2020. Streamlined sampling

and cultivation of the pelagic cosmopolitan larvacean, *Oikopleura dioica*. J. Vis. Exp.

160: e61279. https://doi.org/10.3791/61279

Menden-Deuer, S., Lessard, E.J., 2000. Carbon to volume relationships for dinoflagellates,

diatoms, and other protist plankton. Limnol. Oceanogr. 45, 569–579.

https://doi.org/10.4319/lo.2000.45.3.0569

Mianzan, H., Pájaro, M., Colombo, G.A., Madirolas, A., 2001. Feeding on survival-food:

gelatinous plankton as a source of food for anchovies. Jellyfish Bloom. Ecol. Soc.

Importance 45–53. https://doi.org/10.1007/978-94-010-0722-1_5

Millette, N.C., Pierson, J.J., Aceves, A., Stoecker, D.K., 2016. Mixotrophy in Heterocapsa

rotundata: a mechanism for dominating the winter phytoplankton. Limnol. Oceanogr.

62, 836–845. https://doi.org/10.1002/lno.10470

Morán, X.A.G., López-Urrutia, Á., Calvo-Díaz, A., LI, W.K.W., 2010. Increasing importance

of small phytoplankton in a warmer ocean. Glob. Chang. Biol. 16, 1137–1144.

https://doi.org/10.1111/j.1365-2486.2009.01960.x

Mullin, M. M., Brooks, E. R., 1976. Some consequences of distributional heterogeneity of

phytoplankton and zooplankton. Limnol. Oceanogr. 21, 784–796.

https://doi.org/10.4319/lo.1976.21.6.0784

Nejstgaard, J.C., Naustvoll, L.J., Sazhin, A., 2001. Correcting for underestimation of

microzooplankton grazing in bottle incubation experiments with mesozooplankton. Mar.

Ecol. Prog. Ser. 221, 59–75. https://doi.org/10.3354/meps221059

Nieblas, A.E., Demarcq, H., Drushka, K., Sloyan, B., Bonhommeau, S., 2014. Front

variability and surface ocean features of the presumed Southern Bluefin Tuna spawning

grounds in the tropical southeast Indian Ocean. Deep-Sea Res. II. 107, 64–76.

https://doi.org/10.1016/j.dsr2.2013.11.007

Ohtsuka, S., Böttger-Schnack, R., Okada, M., Onbé, T., 1996. In situ feeding habits of

Oncaea (Copepoda: Poecilostomatoida) from the upper 250 m of the central Red Sea,

with special reference to consumption of appendicularian houses. Bull. Plankt. Soc.

Japan 43, 89–105.

Paffenhöfer, G.A., Sherr, B.F., Sherr, E.B., 2007. From small scales to the big picture:

persistence mechanisms of planktonic grazers in the oligotrophic ocean. Mar. Ecol.

https://doi.org/10.1111/j.1439-0485.2007.00162.x

Ptacnik, R., Sommer, U., Hansen, T., Martens, V., 2004. Effects of microzooplankton and

mixotrophy in an experimental planktonic food web. Limnol. Oceanogr. 49, 1435–1445.

https://doi.org/10.4319/lo.2004.49.4_part_2.1435

Raes, E.J., Hörstmann, C., Landry, M.R.., Beckley, L.E.. Marin, M., Thompson, P., Antoine,
D., Focardi, A., O'Brien, J., Ostrowski, M., Waite, A.M. 2022. Dynamic change in an
ocean desert: microbial diversity and trophic transfer along the 110° E meridional in the
Indian Ocean. Deep-Sea Res. II. 201, 105097.
https://doi.org/10.1016/j.dsr2.2022.105097
Rose, J.M., Caron, D.A., Sieracki, M.E., Poulton, N., 2004. Counting heterotrophic
nanoplanktonic protists in cultures and aquatic communities by flow cytometry. Aquat.
Microb. Ecol. 34, 263–277. https://doi.org/10.3354/ame034263
Saiz, E., Calbet, A., 2011. Copepod feeding in the ocean: scaling patterns, composition of
their diet and the bias of estimates due to microzooplankton grazing during incubations.
Hydrobiologia 666, 181–196. https://doi.org/10.1007/s10750-010-0421-6
Sato, M., Hashihama, F., 2019. Assessment of potential phagotrophy by pico- and
nanophytoplankton in the North Pacific Ocean using flow cytometry. Aquat. Microb.
Ecol. 82, 275–288. https://doi.org/10.3354/ame01892
Sato, T., Shiozaki, T., Hashihama, F., Sato, M., Murata, A., Sasaoka, K., Umeda, S. ichiro,
Takahashi, K., 2022. Low nitrogen fixation related to shallow nitracline across the
eastern Indian Ocean. J. Geophys. Res. Biogeosciences 127.
https://doi.org/10.1029/2022JG007104
Scheinberg, R., Landry, M., Calbet, A., 2005. Grazing of two common appendicularians on
the natural prey assemblage of a tropical coastal ecosystem. Mar. Ecol. Prog. Ser. 294,
201–212. https://doi.org/10.3354/meps294201
Schenone, L., Aarons, Z.S., García-Martínez, M., Happe, A., Redoglio, A., 2024. Mixotrophic
protists and ecological stoichiometry: connecting homeostasis and nutrient limitation
from organisms to communities. Front. Ecol. Evol. 12, 1–14.
https://doi.org/10.3389/fevo.2024.1505037
Schmoker, C., Hernández-León, S., Calbet, A., 2013. Microzooplankton grazing in the
oceans: impacts, data variability, knowledge gaps and future directions. J. Plankton
Res. 35, 691–706. https://doi.org/10.1093/plankt/fbt023
Schnetzer, A., Caron. D.A., 2005. Copepod grazing impact on the trophic structure of the
microbial assemblage of the San Pedro Channel, California. J. Plankton Res. 27, 959–
971. https://doi.org/10.1093/plankt/fbi049
Selph, K.E., 2021. Enumeration of marine microbial organisms by flow cytometry using near-
UV excitation of Hoechst 34580-stained DNA. Limnol. Oceanogr. Methods 19, 692–
701. https://doi.org/10.1002/lom3.10454
Selph, K.E., Yingling, N., Traboni, C., Landry, M.R., This issue. Acidic vacuole-containing
organisms are a majority of the eukaryotic microbial community in oligotrophic Argo
Basin waters (Eastern Indian Ocean). Deep-Sea Res. II.

Stoecker, D.K., Hansen, P.J., Caron, D.A., Mitra, A., 2017. Mixotrophy in the marine
plankton. Ann. Rev. Mar. Sci. 9, 311–335. https://doi.org/10.1146/annurev-marine-
010816-060617

Strickland, J.D.H., Parsons, T.R., 1972. A practical handbook of seawater analysis, 2nd ed,
Fisheries Research Board of Canada. Bulletin Fisheries Research Board of Canada,
Ottawa, Canada. https://doi.org/http://dx.doi.org/10.25607/OBP-1791

Stukel, M. R., Mislan, K. A. S., Décima, M., Hmelo L., 2014. Detritus in the pelagic ocean, p.
49–76. In P. F. Kemp (Eds), Eco-DAS IX Symposium Proceedings. Association for the
Sciences of Limnology and Oceanography.

Swalethorp, R., Malca, E., Shiroza, A., Kim, L., Décima, M., Quintanilla, J., Borrego-Santos,
R., Davies, C., Die, D., Beckley, L., Traboni, C., Cawley, G., Walsh, K., Landry, M.R.,
Laiz-Carrión, R. This issue. Selective feeding in Southern Bluefin Tuna (*Thunnus*
maccoyii) larvae in the Argo Basin, Eastern Indian Ocean spawning ground.

Traboni, C., Calbet, A., Saiz, E., 2021. Mixotrophy upgrades food quality for marine calanoid
copepods. Limnol. Oceanogr. 66, 4125–4139. https://doi.org/10.1002/lno.11948

Traboni, C., Calbet, A., Saiz, E., 2020. Effects of prey trophic mode on the gross-growth
efficiency of marine copepods: the case of mixoplankton. Sci. Rep. 10, 1–14.
https://doi.org/10.1038/s41598-020-69174-w

Unrein, F., Gasol, J.M., Not, F., Forn, I., Massana, R., 2014. Mixotrophic haptophytes are
key bacterial grazers in oligotrophic coastal waters. ISME J. 8, 164–176.
https://doi.org/10.1038/ismej.2013.132

Verschoor, A.M., Boonstra, H., Mejer, T., 2005. Application of stable isotope tracers to
studies of zooplankton feeding using the rotifer *Brachionus calyciflorus* as an example.
Hydrobiologia. 546, 535–549. https://doi.org/10.1007/s10750-005-4296-x

Vincent, D., Hartmann, H.J., 2001. Contribution of ciliated microprotozoans and
dinoflagellates to the diet of three copepod species in the Bay of Biscay. Hydrobiologia
443, 193–204. https://doi.org/10.1023/A:1017502813154

Ward, B.A., Follows, M.J., 2016. Marine mixotrophy increases trophic transfer efficiency,
mean organism size, and vertical carbon flux. Proc. Natl. Acad. Sci. 113, 2958–2963.
https://doi.org/10.1073/pnas.1517118113

Wilken, S., Huisman, J., Naus-Wiezer, S., Van Donk, E., 2013. Mixotrophic organisms
become more heterotrophic with rising temperature. Ecol. Lett. 16, 225–233.
https://doi.org/10.1111/ele.12033

Woo, M., Pattiaratchi, C. 2008. Hydrography and water masses off the western Australian
coast. Deep-Sea Res. I 55, 1090–1104. https://doi.org/10.1016/j.dsr.2008.05.005

Yen, J., 1985. Selective predation by the carnivorous marine copepod *Euchaeta elongata*:

laboratory measurements of predation rates verified by field observations of temporal

and spatial feeding patterns. Limnol. Oceanogr. 30, 577–597.

https://doi.org/10.4319/lo.1985.30.3.0577

Yingling, N., Selph, K.E., Landry, M.R., Kranz, S.A., Johnson, M., Stukel, M.R., This issue.

Phytoplankton nutrient uptake, abundance, biomass and community composition in the

oligotrophic Argo Basin, Indian Ocean. Deep-Sea Res. II.

Zeldis, J. R., Décima, M., 2020. Mesozooplankton connect the microbial food web to higher

trophic levels and vertical export in the New Zealand Subtropical Convergence Zone.

Deep-Sea Res. I 155, 103146, https://doi.org/doi:10.1016/j.dsr.2019.103146

Table legends

Table 1. General spatio-temporal details of the grazing experiments and main parameters relative to seawater sampled through Niskin bottles during evening CTD casts and zooplankton live net sampling. Abbreviations: Exp: experiment, Lat: latitude, Long: longitude, Temp: temperature, Oxy: dissolved oxygen, FChl$a$: chlorophyll a from fluorescence; Zoo: zooplankton.

Table 2. General details of the grazing experiments. Duration of each incubation phase, mean number of copepods incubated per bottle, copepod weight, abundance of ^{15}N-*Pro* (*Prochlorococcus*) and ^{15}N-*Syn* (*Synechococcus*) inoculated in copepod bottles. Errors are standard deviations (sd).

Table 3. Concentration of Chl$a$ (μg L^{-1}) and abundance of plankton groups (cells mL^{-1}) at the start of Microzoo incubation (T$_0$) and at the start of Microzoo+copepod incubation (T$_{24}$). Samples not analysed: n.a. Data are means ± sd (N=3).

Figure 1. A) BLOOFINZ research cruise study area. B) Geographical position where grazing experiments were conducted, which corresponds to the region enclosed within the square in A). Circles indicate cycle experiments; triangles indicate transit experiments.

Figure 2. Timeline of the experimental phases. Evening sampling of seawater and zooplankton, filling of bottles and isotope spike at the onset of the Microzoo incubation (Start, T_0); 24h later, acclimated zooplankton were then added in the isotope-containing bottles, and start of the Microzoo+copepod incubation (T_{24}). After 12h incubation with copepods, termination of the experiment (End, T_{36}). Red arrows indicate the bottles sampled for analyses. The illustration is merely a representative scheme, since triplicate bottles per treatment were used.

Figure 3. Initial biomass of the unicellular groups of pico- and nanoplankton available as prey for microzooplankton at T_0 (A,B,C) and copepods at T_{24} (D,E,F) and relative contribution of trophic modes and size classes to the biomass in cycle and transit experiments. Data are means (N=3-6). Note that during Cycle1, data on eukaryotes abundance at T_0 were not available.

Figure 4. Copepod feeding. A) Grazing rate, B) clearance rate, C) cell-specific ingestion rates and D) C-specific ingestion rate on the different protist populations in cycle and transits experiments. Absent values indicate null or negative grazing. Data are mean ± standard error (N=3-6).

Figure 5. Copepod diet composition on eukaryotes in the different cycles and transit experiments. Data are mean relative contributions of groups (A, B), trophic modes (C, D), and size classes (E, F) to the copepod diet expressed as cell-specific ingestion (left panels) and C-specific ingestion (right panels).

Figure 6. Copepod food selection. Chesson's selectivity (α) index on A) the prey groups, and Ivlev's electivity index (E) on the prey groups based on B) trophic mode, and C) size. Mean values from all experiments were pooled together per each category. Median values are reported for each prey and/or category. Isolated dots represent outliers. The dashed line represents neutral selection.

Figure 7. Copepod ingestion rate and daily ration. A) C-specific ingestion rate and B)

daily ration estimated by food removal (black), C) C-specific ingestion rate and D) daily

ration inferred through isotope signal (δ^{13}C); E) N-specific ingestion rate, F) and daily

ration inferred through isotope signal (δ^{15}N). The concentration of ^{15}N-*Pro* was

negligible (10-13%) compared to that of ^{15}N-*Syn* (87-90%) in Cycle 3, Cycle 4, and

Transit. Data are mean±standard errors (N=9-18).

1018

1019 **Figure 8.** Grazing rate by microzooplankton on A) Chl*a*, B) prokaryotes, and C)

1020 eukaryotes and contribution to dietary C of the different D) prey groups, E) trophic

1021 modes and F) size classes. Data in the upper panels are mean ± standard error (N=3-

1022 6). Note that during Cycle1, data on eukaryotes abundance at T_0 were not available,

1023 thus grazing rates were not measured.

1024

1025 **Figure 9.** Percentage of prey production removed daily by A) microzooplankton over

1026 24h, and B) copepods over the following 12h. Median values are reported on the top of

1027 each group in each station (N=5, including cycle and transit experiments).

Table 1. General spatio-temporal details of the grazing experiments and main parameters relative to seawater sampled through Niskin bottles during evening CTD casts and zooplankton live net sampling. Abbreviations: Exp: experiment, Lat: latitude, Long: longitude, Temp: temperature, Oxy: dissolved oxygen, FChl*a*: chlorophyll *a* from fluorescence; Zoo: zooplankton.

Date	Cycle / Station	Cycle day	Exp	Lat °N	Long °E	Depth (m)	Salinity	Temp (°C)	Oxy (µmol L^{-1})	FChl*a* (µg L^{-1})	Zoo sampling Depth (m)
04-feb-22	1	1	1	-15.37	114.43	30	34.53	28.61	190.7	0.118	48
09-feb-22	2	1	2	-16.74	115.69	30	34.93	27.93	193.5	0.118	38
12-feb-22	2	3	3	-16.99	116.05	30	34.97	28.09	192.6	0.108	38
15-feb-22	3	1	4	-15.98	115.83	28	34.66	28.63	194.8	0.125	41
17-feb-22	3	3	5	-15.81	115.82	30	34.56	28.01	195.3	0.143	38
20-feb-22	4	1	6	-15.89	118.14	30	34.52	29.18	193.5	0.144	60
22-feb-22	4	3	7	-15.95	118.11	30	34.51	29.22	193.1	0.125	42
25-feb-22	St 37-Argo4	na	8	-15.01	117.44	2	34.59	30.39	186.7	0.050	42
28-feb-22	St 44-Argo11	na	9	-13.53	116.60	2	34.42	29.94	103.3	0.044	50

Table 2. General details of the grazing experiments. Duration of each incubation phase, mean number of copepods incubated per bottle, copepod weight, abundance of ^{15}N-*Pro* (*Prochlorococcus*) and ^{15}N-*Syn* (*Synechococcus*) inoculated in copepod bottles. Errors are standard deviations (sd).

Cycle / Station	Exp	Microzoo incubation T_0-T_{24}	Microzoo + copepod incubation T_{24}-T_{36}	Total incubation T_0-T_{36}	Copepods Abundance	Weight	^{15}N-Enrichment ^{15}N-*Pro*	^{15}N-*Syn*
		(h)			(cop. bottle^{-1})	(µg C ind^{-1})	(cells mL^{-1})	
1	1	23.0	12.2	35.2	48 ± 5	24.1 ± 7.5	13,400	
2	2	21.4	6.0	27.4	38 ± 5	8.4 ± 0.6	6,730	
2	3	21.5	14.6	36.1	70 ± 0	34.1 ± 12.8	6,710	
3	4	22.7	13.5	36.2	61 ± 4	56 ± 19.3	5,680	25,680
3	5	23.8	12.9	36.7	39 ± 4	24.2 ± 8.6	5,660	57,090
4	6	22.4	13.5	35.9	26 ± 2	17.2 ± 1.5	8,540	73,800
4	7	22.5	14.2	36.7	32 ± 5	26.2 ± 5.2	9,480	66,600
37	8	21.1	12.8	33.9	11 ± 1	20.6 ± 9.3	10,970	89,940
44	9	21.7	13.1	34.8	27 ± 9	104 ± 18	10,940	91,160

Table 3. Concentration of Chl$a$ (μg L^{-1}) and abundance of plankton groups (cells mL^{-1}) at the start of Microzoo incubation (T$_0$) and at the start of Microzoo+copepod incubation (T$_{24}$). Samples not analysed: n.a. Data are means ± sd (N=3).

Incubation phase	Cycle/ Station	Exp	Chl$a$	Hbact	Pro	Syn	Picoauto	Nanoauto	Picomixo	Nanomixo	Nanohetero
			(μg L^{-1})	(x10^3 cells mL^{-1})		(cells mL^{-1})					
Microzoo incubation (T$_0$)	1	1	0.07±0.009	482±26	128±13	1,261±111	n.a.	n.a.	n.a.	n.a.	n.a.
	2	2	0.11±0.003	549±28	271±5	1,684±62	29±2.27	134±21	44±35	273±104	1,600±211
	2	3	0.07±0.012	606±15	276±2	3,464±2,673	33±10	169±34	19±15	28±15	1,220±66.8
	3	4	0.1±0.007	592±29	201±21	1,506±123	169±36	41±8	244±25	481±40	1,010±352
	3	5	0.1±0.003	625±19	213±3	1,908±119	134±32	66±22	218±21	666±82	817±280
	4	6	0.09±0.004	552±5	216±1	2,373±302	182±38	72±11	131±40	563±122	2,610±983
	4	7	0.07±0.004	532±14	188±0.4	1,895±33	182±107	28±7	237±43	416±41	1,530±856
	37	8	0.05±0.008	514±5	168±2	1,529±71	165±27	40±10	282±64	561±35	1,720±326
	44	9	0.06±0.004	539±3	166±2	1,740±5	202±17	41±8	247±32	590±52	581±96.2
Microzoo + copepod incubation (T$_{24}$)	1	1	n.a.	n.a.	n.a.	n.a.	446±77	117±36	297±40	328±76	465±89
	2	2	n.a.	n.a.	n.a.	n.a.	126±115	313±116	18.7±15	64±7	803±185
	2	3	0.05±0.032	634±49	283±14	1,876±202	175±40	60±14	199±41	474±44	2,380±1702
	3	4	0.1±0.02	517±4	112±33	1,457±284	180±12	74±6	173±22	355±60	481±208
	3	5	0.061±0.037	585±17	167±96	1,762±313	187±19	83±6	168±41	418±19	839±356
	4	6	0.05±0.036	536±34	94±106	1,506±300	95.9±12	60±16	115±33	441±64	1,620±424
	4	7	0.09±0.015	525±11	181±46	2,411±37	70.1±19	58±8	90±19	472±110	823±341
	37	8	0.03±0.011	483±7	45±23	876±391	144±38	65±3	131±46	424±73	946±359
	44	9	0.13±0.007	593±57	95±74	1,416±966	134±19	88±26	218±75	545±110	558±73

Figure 1. A) BLOOFINZ research cruise study area. B) Geographical position where grazing experiments were conducted, which corresponds to the region enclosed within the square in A). Circles indicate cycle experiments, denoted as C1-2-3-4; triangles indicate transit experiments (Tr) at stations 37 and 44.

Figure 2. Timeline of the experimental phases. Evening sampling of seawater and zooplankton, filling of bottles and isotope spike at the onset of the Microzoo incubation (Start, T_0); 24h later, acclimated zooplankton were then added in the isotope-containing bottles, and start of the Microzoo+copepod incubation (T_{24}). After 12h incubation with copepods, termination of the experiment (End, T_{36}). Red arrows indicate the bottles sampled for analyses. The illustration is merely a representative scheme, since triplicate bottles per treatment were used.

Figure 3. Initial biomass of the unicellular groups of pico- and nanoplankton available as prey for microzooplankton at T_0 (A,B,C) and copepods at T_{24} (D,E,F) and relative contribution of trophic modes and size classes to the biomass in cycle and transit experiments. Data are means (*N*=3-6). Note that during Cycle1, data on eukaryotes abundance at T_0 were not available.

Figure 4. Copepod feeding. A) Grazing rate, B) clearance rate, C) cell-specific ingestion rates and D) C-specific ingestion rate on the different protist populations in cycle and transits experiments. Absent values indicate null or negative grazing. Data are mean ± standard error (*N*=3-6).

Figure 5. Copepod diet composition on eukaryotes in the different cycles and transit experiments. Data are mean relative contributions of groups (A, B), trophic modes (C, D), and size classes (E, F) to the copepod diet expressed as cell-specific ingestion (left panels) and C-specific ingestion (right panels).

Figure 6. Copepod food selection. Chesson's selectivity (α) index on A) the prey groups, and Ivlev's electivity index (*E*) on the prey groups based on B) trophic mode, and C) size. Mean values from all experiments were pooled together per each category. Median values are reported for each prey and/or category. Isolated dots represent outliers. The dashed line represents neutral selection.

Figure 7. Copepod ingestion rate and daily ration. A) C-specific ingestion rate and B) daily ration estimated by food removal, C) C-specific ingestion rate and D) daily ration inferred through isotope signal (δ^{13}C); E) N-specific ingestion rate, F) and daily ration inferred through isotope signal (δ^{15}N). The concentration of ^{15}N-*Pro* was negligible (10-13%) compared to that of ^{15}N-*Syn* (87-90%) in Cycle 3, Cycle 4, and Transit. Data are mean±standard errors (*N*=9-18).

Figure 8. Grazing rate by microzooplankton on A) Chl*a*, B) prokaryotes, and C) eukaryotes and contribution to dietary C of the different D) prey groups, E) trophic modes and F) size classes. Data in the upper panels are mean ± standard error (*N*=3-6). Note that during Cycle1, data on eukaryotes abundance at T_0 were not available, thus grazing rates were not measured.

Figure 9. Percentage of prey production removed daily by A) microzooplankton over 24 h, and B) copepods over the following 12 h. Median values are reported on the top of each group (*N*=5, including cycle and transit experiments).