Questioning the utility of oxidative stress measurements as biomarkers of physiological condition and fitness

Rachel R Reid[1]*, Davide M Dominoni[1] *, Jelle Boonekamp[1]

[1]School of Biodiversity, One Health and Veterinary Medicine, Graham Kerr Building, University of Glasgow, Glasgow, G12 8QQ, UK

*Email: Rachelrosereid141@outlook.com

Keywords: Oxidative stress, intraindividual variation, measurement reliability, ageing physiology

Abstract

Markers of oxidative stress are widely used as biomarkers of health, ageing, and physiological
stress. However, their reliability as biomarkers remains uncertain due to high intraindividual
variation, obscuring associations with environmental conditions, lifestyle, frailty, and physiological
indicators of health such as telomere length. Whilst numerous longitudinal studies exist, individual
repeatability of oxidative stress measurements is rarely reported. This study presents the first
meta-analysis assessing individual repeatability of oxidative stress markers, comprising 123
repeatability estimates from 22 studies. We found that oxidative stress exhibits low individual
repeatability (Intraclass correlation = 0.164), regardless of marker type, taxa, sex, study design,
or environment. We also found that different markers of oxidative stress were often poorly
correlated, which could be due to their low repeatability. This flags serious limitations regarding
the utility of oxidative stress measurements as health biomarkers. To illustrate this point, we
simulated causal effects of oxidative stress on telomere length to reveal statistical power
limitations on the detection of a relationship between oxidative stress and telomere length when
individual repeatability is low. Our simulations reveal that substantially larger samples sizes are
required than those typically used in this field. On a more positive note, we also show that
increasing the number of repeated samples can improve statistical power. Such longitudinal
studies would create valuable opportunities for untangling the causes of intraindividual variation in
oxidative stress, elucidating the functional consequences of oxidative stress for health and
ageing.

Introduction

Oxidative stress is a pervasive feature of life, implicated in cellular senescence, cell proliferation,
inflammation, ageing, and disease (1–4). It arises when prooxidants, exceed the neutralising
capacity of antioxidants (5–7). When oxidative stress becomes deregulated, the imbalance
initiates a cascade of reactions causing DNA damage, alterations to membrane lipids, and
damage to other macro-molecules (8–10). On a systemic level such cellular damage is widely
believed to contribute to organismal senescence (1,11,12) and ageing related disease (13–16).
Understanding the sources of individual variation in oxidative stress may therefore offer crucial
insights into the fundamental mechanisms of ageing and health. For example, elevated levels of
oxidative stress might reveal an unhealthy lifestyle if oxidative stress measurements link lifestyle
factors to health. However, health status beyond obvious disease is difficult to measure.
Therefore, the utility of oxidative stress as biomarkers depends on their associations with other
health markers such as telomere length. Herein lies a fundamental challenge as oxidative stress
is often poorly correlated with other biomarkers. For example, although telomere length is widely
believed to be linked with adult mortality (17) and oxidative stress (8), a recent meta-analysis (18)
found that the correlation between oxidative stress and telomere length is surprisingly weak (r =
0.027). This raises concerns about whether markers of oxidative stress capture meaningful
aspects of health (18). This issue is of particular importance given the widespread use of
oxidative stress biomarkers in clinical and pharmaceutical research (15, 16), ecotoxicology (17)
and evolutionary ecology (5,19,20). The cause of the weak association between oxidative stress
and telomere length remains unclear. One simple explanation could be that oxidative stress
markers show high intraindividual variability such that the observed level of oxidative stress at a
given point in time might poorly reflect the oxidative stress profile of that individual. Such high
intraindividual variability becomes a challenge especially the intraindividual variation is
unpredictable. Hence, the *conditional* repeatability (controlling statistically for potential sources of
variation in the data (21)) fundamentally underpins the utility of oxidative stress measurements as
biomarkers of health and fitness (Fig. 1).

Figure 1. Conceptual depiction of the impact of different scenarios of intraindividual variation in oxidative stress on their utility to predict telomere length in benign and harsh environments. Panels A-C depict repeated individual measurements of oxidative stress across time for (A) high repeatability and predictability, (B) low repeatability and high predictability, (C) low repeatability and low predictability for an individual living in a benign (white bars), and another individual living a harsh environment (grey bars). Panels D-F illustrate the impact of these corresponding scenarios on the population-level associations between oxidative stress, telomere length, and environmental condition. Panel E illustrates that there is little negative impact of low repeatability when one can statistically control for the confounding variation, in this case month. However, oxidative stress is a poor biomarker in the third scenario (panels C and F) because it fails to meaningfully link the environmental effect on oxidative stress to telomere length. Following the logic set out in these simplistic conceptual scenarios, the utility of oxidative stress measurements as health biomarkers hinges on their conditional repeatability, i.e., the repeatability after successfully controlling for confounding variables.

While establishing individual repeatability has been a key research focus for several physiological
traits including metabolic rate (22), glucocorticoid hormones (23,24), cognitive performance (25),
and telomere length (26), the individual repeatability of oxidative stress markers remains unclear.
Where studies do report them, repeatability estimates evidently vary widely. For example,
glutathione peroxidase displayed moderate repeatability (r = 0.47) in great tits (*Parus major*)
across seasons (27). In collared flycatchers' (*Ficedula albicollis*) reactive oxygen metabolites
were not repeatable (r = 0.03), whereas antioxidant defences exhibited moderate repeatability (r
= 0.581) (28). A growing body of evidence suggests that the individual repeatability of oxidative
stress can be affected by a range of environmental and life-history factors. For example, changes
in oxidative stress levels have been observed across life-history stages (5,29,30) and has been
shown to be elevated during mating and reproduction (31,32), and high-energy activities like
migration (33). Oxidative stress is also affected by certain environmental conditions including
predation risk (34), territory quality (35), and seasonal changes (36,37). The plethora of factors
that can influence oxidative stress raises the challenge of how to control for this multitude of
environmental effects on intraindividual variation in oxidative stress. Further adding to this
challenge, oxidative stress has been shown to fluctuate over very short timescales (38,39),
raising concerns about their fine scale temporal stability. From a 'biomarker utility' perspective,
the question remains, to what extent is the observed intraindividual variation predictable? Are
there universal factors that underpin intraindividual variation in oxidative stress? Are there certain
types of oxidative stress markers that might be more stable and therefore more suitable as a
biomarker? Certain antioxidants like vitamin E will largely reflect diet (40,41) whereas more 'pure'
measurements of reactive oxygen species might reflect more dynamic physiological processes
linked to metabolism. There could also be differences in the transient behaviour of enzymatic
versus non-enzymatic, and intra- or extracellular markers of oxidative stress. Indeed, different
oxidative stress markers are often very poorly correlated, even within the same individual
(6,16,42–44). These issues highlight a need to quantify intraindividual variation across a panel of
commonly used oxidative stress markers, to elucidate the potential causes of this variation, and
to determine whether certain markers of oxidative stress might be more suitable as biomarkers
than others.

The increasing availability of longitudinal studies offers the opportunity to investigate the level of
individual repeatability of oxidative stress markers for a wide range of species and identify factors
contributing to intraindividual variation. Here, we present the first formal meta-analysis of
individual repeatability of oxidative stress. By examining biological and methodological factors, we
aim to estimate the overall repeatability and to identify the causes of variation in repeatability. We
also evaluate the correlations among oxidative stress markers to determine if any
systematic/conserved patterns exist. We subsequently explore the consequences of the observed
low repeatability for study design and statistical power using individual based simulations on the
association between oxidative stress and telomere length and provide recommendations
addressing these issues for future studies.

96
97
Results

99
Our meta-analysis synthesises 123 estimates of individual repeatability, estimated with the
intraclass correlation (ICC), derived from 22 published datasets. Across all studies, the overall
repeatability of oxidative stress markers was low (ICC = 0.164, 95% confidence interval (CI) =
[0.093, 0.233], P < 0.001; Table S1, Fig.2), with substantial heterogeneity among effect sizes (I^2 =
57.4). Mammals exhibited slightly higher ICC values (ICC = 0.212, 95% CI = [0.058, 0.356], P =
0.007) compared to birds (ICC = 0.155, 95% CI = [0.062, 0.245], P = 0.012; Table S1, Fig.2), but
the difference between groups was small and not statistically significant (estimate = 0.059, 95%
CI = [-0.123, 0.238], P = 0.526). These findings indicate that oxidative stress markers show low
repeatability overall, regardless of taxonomic group. The high heterogeneity observed among
effect sizes is noteworthy and indicative of a range of repeatability estimates including instances
of moderate to high repeatability. This raises the question whether there could be common
factors underpinning the higher repeatability observed in a minority of cases, and vice versa
whether there are systematic explanations for low repeatability observed in the majority of
studies.

Figure 2: Caterpillar plot displaying individual effect sizes sorted from lowest to highest repeatability and split by taxonomic class. *Yellow points represent ICC values for the "Aves" class while blue points correspond to ICC values for the "Mammal" class. The global ICC estimate for "Aves" is shown at the bottom of the Aves class facet, and the global estimate for "Mammals" is shown similary. These estimates were extracted from the meta-model where taxonomic class was included as a moderator. ICC values are presented along with their 95% confidence intervals. The dashed line represents an ICC value of zero. The red diamond and error bars at the bottom of the plot represents the overall ICC value and its 95% confidence intervals from the intercept only model. The corresponding number of effect sizes is shown on the right side of the plot, with the number of studies indicated in parentheses.*

Moderator analyses investigating the potential causes of low repeatability

We estimated ICC within treatment and control groups for experimental studies, and within sex to
enable testing the influence of these variables as moderators in our meta-analysis (see methods
for details). The control groups of experimental studies were pooled with the effect sizes of
correlational studies to create a 2-level moderator comprising manipulated animals and
unmanipulated animals. Contrary to our prediction, effect sizes obtained from unmanipulated
animals showed slightly lower repeatability (ICC = 0.146, 95% CI = [0.053, 0.230]) than effect
sizes from animals exposed to experimental manipulations (ICC = 0.208, 95% CI = [0.091,
0.319]), but the difference was not significant (estimate = -0.064, 95% CI = [-0.204, 0.077], P =
0.86; Table S1, Fig. 3A). Females also showed similar repeatability (ICC = 0.167, 95% CI =
[0.065, 0.266]) to males (ICC = 0.154, 95% CI = [0.03, 0.273]; estimate = -0.013, 95% CI = [-
0.163, 0.137], P = 0.86; Fig. 3A-B; Table S1, Fig. 3B).

We observed higher repeatability for studies on livestock populations (ICC = 0.391, 95% CI =
[0.178, 0.569], P = 0.005), compared to studies using wild (ICC = 0.119, 95% CI = [-0.003,
0.239], P = 0.057), wild captive (ICC = 0.165, 95% CI = [-0.052, 0.367], P = 0.136), or captive-
bred animals (ICC = 0.158, 95% CI = [-0.07, 0.371], P = 0.173; Table S1, Fig. 3C). Using
livestock as the reference category, model estimates indicated significantly lower repeatability in
wild populations (estimate = -0.285, 95% CI = [-0.506, -0.029], P = 0.029). Repeatability was
lower in wild-captive (estimate = -0.242, 95% CI = [-0.513, 0.072], P = 0.129), and captive
populations (estimate = -0.248, 95% CI = [-0.523, 0.074], P = 0.074) but neither were significantly
different from the livestock reference level. The elevated repeatability in livestock must be
interpreted with caution, as these datasets included a higher number of repeated measures per
individual, which can increase ICC estimates.

We found no evidence for systematic differences in repeatability among markers of oxidative
damage versus antioxidant protection (estimate = -0.016, 95% CI = [-0.140, 0.109], P = 0.803).
Both categories showed low but significant repeatability: oxidative damage (ICC = 0.156, 95% CI
= [0.056, 0.253], P = 0.002); antioxidant defence (ICC = 0.171, 95% CI = [0.082, 0.258], P =
<0.001; Table S1, Fig.3D). Further analysis of oxidative damage marker subtypes revealed that
only lipid peroxidation markers exhibited moderate repeatability (ICC = 0.368, 95% CI = [0.182,
0.529], P <0.001; Table S2, Fig.3E). Other damage types, including DNA damage (ICC = 0.135,
95% CI = [-0.224, 0.461], P = 0.464), protein damage (ICC = -0.068, 95% CI = [-0.346, 0.220], P
= 0.646), and nonspecific damage (ICC = 0.059, 95% CI = [-0.137, 0.252], P = 0.59), were not
statistically significant. Using lipid peroxidation as a reference level, protein damage (estimate = -
0.426, 95% CI = [-0.654, -0.128], P = 0.006) and non-specific damage (estimate = -0.316, 95% CI
= [-0.539, -0.051], P = 0.020) were shown to have significantly lower repeatability. DNA damage
also had lower repeatability than lipid peroxidation, but this difference was not significant
(estimate = -0.245, 95% CI = [-0.578, 0.156], P = 0.228). In a similar way, enzymatic antioxidants
showed slightly higher repeatability (ICC = 0.204, 95% CI = [0.081, 0.320], P = 0.001), compared
to non-enzymatic defence markers (ICC = 0.136 95% CI = [0.042, 0.228], P = 0.005; Table S2,
Fig.3F), but the difference between these two categories was not statistically significant (estimate
= -0.069, 95% CI = [-0.198, 0.061], P = 0.296), and both subtypes showed low repeatability
overall.

Finally, we tested whether the mean time elapsed between repeated oxidative stress
measurements and the number of repeated samples per individual influenced estimates of within-
individual repeatability, as we would expect these to reduce and increase repeatability,
respectively. Surprisingly, the average time interval between repeated measurements had a small
but statistically significant positive effect on ICC (slope estimate = 0.022, 95% CI = [0.006, 0.038],
P = 0.008; Table S1, Fig. 3G). In line with expectations, the number of repeated measurements
per individual had a significant positive effect on ICC (slope estimate = 0.097, 95% CI = [0.064,
0.130], P <0.001; Table S1, Fig. 3H).

Figure 3: Within-individual repeatability of oxidative stress measures for each moderator level.
Panels A-F show repeatability estimates for categorical moderators (A) Study type, (B) Sex, (C) Population source, (D) Oxidative stress biomarker, (E) Type of oxidative damage, (F) Type of antioxidant defence. For each level, the model-estimated repeatability (ICC) is plotted with 95% confidence intervals, back transformed from Fishers Z. The dashed vertical line indicates an ICC of zero. Panels G and H display the relationship between continuous moderators and repeatability: (G) Log-transformed average time between measures and (H) Log-transformed average number of measures per individual. In each panel, grey points represent individual effect sizes, the black line shows the model-predicted relationship, and the shaded ribbon denotes the 95 confidence intervals. The red dashed line indicates an ICC = 0. ICC is plotted on the y-axis in G and H, and on the x axis in A-F.

Conditional repeatability analysis

To assess whether controlling for individual-level covariates enhanced repeatability estimates of oxidative stress biomarkers, we conducted a separate meta-analysis using raw data from eight studies that included information on age, to compute conditional repeatability on age. We explored if a similar approach would have been possible for other variables, but unfortunately there was limited information available from the 22 studies at our disposal. For each study we calculated repeatability estimates using four model types: (1) raw (no covariates), (2) age adjusted, (3) age and sex adjusted (where data on sex was available), (4) age and treatment adjusted (where data on treatment was available). Across all models, estimates of repeatability were broadly similar (Fig. 4). Raw repeatability (R = 0.195, 95% CI = [0.067, 0.316], P = 0.003), and conditional repeatability estimates controlling for age (R = 0.187, 95% CI = [0.058, 0.309], P = 0.004), or age and sex (R = 0.167, 95% CI = [0.025, 0.302], P = 0.021) were all indistinguishable from each other. However, conditional repeatability adjusting for age and treatment (R = 0.356, 95% CI = [-0.035, 0.653], P = 0.073, Fig.4) yielded a higher repeatability estimate although this was only based on two effect sizes. These findings suggest that controlling for common covariates such as age and sex has minimal impact on repeatability estimates. This reinforces the conclusion that oxidative stress markers tend to show low repeatability across contexts and that we still have little understanding of the drivers' causing variation in oxidative stress.

Figure 4: Within-individual repeatability of oxidative stress measures for the type of repeatability calculation. Repeatability type refers to what variables were controlled for in the model when estimating repeatability. Model estimates are shown along with the 95% confidence intervals. The dashed line indicates an R value of zero. The moderator is shown on the y axis, and R values (back transformed from Fishers Z values) are plotted on the x-axis.

Are biomarkers of oxidative stress correlated?

Oxidative stress biomarkers are often assumed to reflect interrelated physiological processes, yet
studies frequently report weak correlations among them (27,28). Our meta-analysis, showing
generally low within-individual repeatability for oxidative stress markers further challenges this
view. From a statistical point of view, given an average repeatability of 20%, the maximum
average correlation between two markers would be $r = 0.04$, underscoring the multiplicative
impact low repeatability has on the expected correlations among different markers of oxidative
stress. Using individual-level raw data available from included studies, we were able to directly
assess the correlations between commonly measured oxidative stress markers, including the
following pairs of markers; protein carbonyls – superoxide dismutase (carbonyl-SOD),
malondialdehyde – antioxidant capacity (MDA-OXY), antioxidant capacity – superoxide
dismutase (OXY-SOD), malondialdehyde – superoxide dismutase (MDA-SOD), and reactive
oxygen metabolites – antioxidant capacity (dROMs-OXY) (Fig. 5). In accordance with their low
repeatability, most pairwise correlations were weak and statistically non-significant (Table S3).
However, we found a significant and modest positive correlation between dROMs and OXY ($r =$
0.250, 95% CI = [0.091, 0.396]; Table S3, Fig.4). This finding is noteworthy because the low ICC
values of dROMs and OXY (0.107 and 0.133, respectively) would predict a much weaker
correlation (expected $r = 0.014$). This could suggest that both markers show synchronous
temporal variation within individuals. Nevertheless, the overall pattern reinforces the idea that
oxidative stress markers are poorly related and that longitudinal and multivariate approaches are
required to capture meaningful aspects of *in vivo* oxidative physiology.

Figure 5: Pearson's correlation for pairs of oxidative stress markers. The model estimates are shown along with 95% confidence intervals. The dashed line indicates where Pearson's correlation is zero. The paired oxidative stress biomarkers are shown on the y axis and Pearson's correlation (back transformed from Fishers Z) is shown on the x axis. The red point indicates the model estimate for the intercept-only model along with the 95% confidence intervals. "k" represents the number of effect sizes.

Implications of low repeatability for statistical power: The case of oxidative stress and telomere length

Our meta-analysis revealed that oxidative stress markers exhibit low within-individual repeatability, raising critical concerns about their reliability as biomarkers of physiological condition and fitness. High intra-individual variation reduces the ability to detect biologically meaningful patterns and undermines statistical power. To quantify the implications of this low repeatability, we conducted an individual-based simulation to estimate the sample sizes required to detect a relationship between oxidative stress and telomere length in three biological scenarios: in the first scenario we assume that oxidative stress is fully causal in telomere attrition. The two additional scenarios depict moderate and weak causal effects of oxidative stress on telomere attrition. All simulations were parameterised using biological parameters within the natural range of a hypothetical great tit population (see supplementary materials).

Our simulations revealed that when oxidative stress repeatability is low (ICC = 0.2), a value
consistent with our empirical estimates), sample sizes n > 100 are required to achieve >80%
statistical power (Fig. 5A). A previous meta-analysis examining oxidative stress and telomere
length correlations (n = 37 studies) found that only 16 studies included more than 100 individuals
(17). This power deficit is even more pronounced under scenarios involving moderate or weak
true effects of oxidative stress on telomere attrition (Fig. 5A).

Depending on the study system, it may be efficient to increase the statistical power by increasing
the number of repeated measurements within individuals as opposed to increasing the sample
size. Our simulations show that increasing the number of repeated oxidative stress
measurements per individual can reduce the total sample size needed to achieve acceptable
power (Fig. 5B). Under the scenario in which the true effect of oxidative stress on telomere
attrition is moderate, taking three to four repeated measurements per individual reduces the
required total sample size by 25-30%, beyond which there are diminishing returns from further
increasing the per animal sampling effort. Hence increasing the number of repeated samples can
be an effective way of increasing statistical power when the study population is amenable to
recaptures. Moreover, such longitudinal studies would provide much needed opportunity for
investigating the causes of intraindividual variation. This could ultimately promote developing
methods that yield higher conditional repeatability which may be more effective in increasing
statistical power.

Figure 6: Simulated power of telomere length and oxidative stress correlations against sample size. A) Power is shown for different scenarios of true biological effects of oxidative stress on telomere attrition and for different individual repeatability (intraclass correlations) of oxidative stress. The coloured lines represent the mean value of each repeatability (ICC) group that was simulated. Shaded areas show the 95% confidence intervals. The red dashed line marks the 80% statistical power threshold. Where the coloured line crosses the 80% threshold this would indicate the minimum sample size required to have this level of power to detect the relationship between telomere length and oxidative stress. **B)** Relationship between sample size and repeated measures needed to achieve statistical power above 80% for detecting a correlation between telomere length and oxidative stress, using a scenario where the oxidative stress individual repeatability is low (ICC = 0.2) and causality of the oxidative stress effect on telomere length is moderate. The different colours represent the number of repeated measures per individual. The red dashed line marks the 80% power threshold. Shaded contours reflect the 95% confidence intervals. **C)** The points show the mean sample size required at each number of oxidative stress samples per individual to achieve >80% power to detect the correlation between telomere length and oxidative stress as shown in B. The curve merely illustrates the trend.

273
274
275 **Discussion**
276
The heuristic value of a trait as a health biomarker hinges on its ability to capture the links
between environmental conditions and fitness. Our meta-analysis reveals that markers of
oxidative stress exhibit high intraindividual variation, relative to interindividual variation, implying
these markers have low statistical power for capturing individual differences in health and fitness.
Furthermore, our moderator analysis indicates that this variability is broadly consistent across
different oxidative stress biomarkers, taxa, sexes, and population sources. We have previously
argued that strong biomarkers of fitness may be rare, because important fitness traits are likely
phenotypically robust and therefore largely invariable with respect to environmental conditions
(5). Here we show that the opposite can also be true in the sense that high intraindividual
variation induces a 'detection' problem when the source of this variation is unknown, complicating
the search for reliable bioindicators. Hence the ideal biomarker appears to comprise of traits that
are moderately repeatable, sensitive to environmental conditions, and moderately predictive of
fitness. Markers of oxidative stress might meet these criteria, if we can enhance our
understanding of the underlying causes of intraindividual variation.

Despite the low overall repeatability, several studies included in our meta-analysis reported
relatively high repeatability of oxidative stress measurements (R > 0.6). What do these studies
have in common that may explain their individual repeatability? While it is not possible to make
any definite conclusions at this point, we made some tentative observations. First, MDA,
particularly when quantified via high-performance liquid chromatography (HPLC), exhibited
relatively high repeatability in both wild (31,45) and controlled settings (46), suggesting its
potential as a reliable biomarker of oxidative status. High and moderate repeatability was also
observed for other biomarkers such as vitamin E and SOD under experimental conditions (47,48),
as well as OXY, SOD, and peroxidation index in wild populations (28,49,50). In controlled or
manipulated conditions, such as dietary interventions or supplementation, high repeatability may
reflect reduced environmental noise or greater physiological stability. Conversely, in wild
populations, high repeatability tended to occur in individuals experiencing prolonged and
predictable physiological demands, such as migration, lactation, and overwinter survival
(31,46,50,51).These findings suggest that repeatable oxidative phenotypes may arise when
individuals experience sustained metabolic challenges. Notably, high repeatability was detected
in both short-term and long-term sampling designs, indicating that repeatability can be high even
at extended sampling interval duration (46–48,50,51). Overall, the lack of consistent repeatability
patterns, even within individual studies, highlights the context dependent nature of oxidative
stress. This underscores the need for standardised methodologies, and experimental designs that
account for ecological and life-history context to better understand individual variation in oxidative
stress.

The timescale of within-individual variation in oxidative stress is important. Even if a true
biological relationship exists between oxidative stress and fitness, high temporal variability in
oxidative stress measurements over short intervals can obscure meaningful links to body
condition and fitness. At present, we are not aware of dedicated studies on the fine scaled
temporal patterns of within-individual variability of oxidative stress through the life course.
However, the stability of oxidative stress has been studied in detail in cell cultures. These in vitro
studies show dynamic instability, leading to unpredictable shifts over time, even under
standardised conditions (52–55). It remains unclear how well these temporal patterns of oxidative
stress in cell cultures that are kept in highly standardized conditions translate to the whole
organism. Among the 10 studies (32,46,48,50,56–61) in our meta-analysis that assessed
repeated measurements over short timescales (e.g. 1 month or less), most found high variability

and low repeatability of oxidative stress markers. Only three of these studies, two of which were
under controlled conditions with multiple repeated measurements, reported high repeatability
(46,48,50). These findings suggest that oxidative stress markers are highly plastic over brief
intervals in the majority of cases. To better characterise and understand the causes of this
variation, we need longitudinal studies that collect many repeated measurements throughout the
life course to obtain fine scale oxidative stress profiles within individuals. This would enable
identifying the drivers of intraindividual variation in oxidative stress potentially improving the utility
of oxidative stress measurements as a biomarker of body condition and fitness.

Our analysis revealed that correlations among different oxidative stress markers were generally
very weak, reinforcing previous findings (39,45).These poor correlations pose a conceptual
challenge because it is generally assumed that different markers of oxidative stress reflect
physiological processes that are dynamically linked (6,19,62). Our findings suggest we need to
step away from this univariate approach as the weak correlations we observed in our meta-
analysis indicate that markers of oxidative stress are largely decoupled. The cause of this
decoupling could be due to the high intraindividual variability itself: When two traits are poorly
repeatable their correlation can be expected to be weak due to the multiplication of poor
repeatability scores when correlating the two traits. On the other hand, when two traits
dynamically fluctuate in time in a synchronous manner then they could be more strongly
correlated than the product of their respective intraclass correlations and we found some
evidence for this regarding the correlation between reactive oxygen metabolites and total
antioxidant capacity (Fig. 5). These observations call for further research into the question of to
what extent different markers of oxidative stress are dynamically linked or whether they may
reflect distinct biochemical processes as previously suggested (38,52,53,63,64). Hence, we need
longitudinal multi-marker approaches which remain comparatively underutilised in ecological and
physiological studies (38,39).

Finally, the phenomenon of decoupling as described above could be relevant outside the context
of oxidative stress. For example, many behavioural traits exhibit high intraindividual variation
(65,66), suggesting that consistent behavioural correlations (i.e., syndromes) may be rare unless
behavioural traits are changing across time and contexts synchronously (65). The loss of
correlations among suites of traits with age could be indicative of a loss of function or
homeostasis, which could be linked to ageing or disease. It will be important to study these
multivariate phenotypic changes in relation to fitness to unravel their functional consequences,
and oxidative stress may lend itself very well for this purpose.

356
357
358 **Materials and Methods**
359
Literature search

We developed a search string using keywords derived from a review of the oxidative stress
literature as follows:

("Oxidative stress" OR "Antioxidant capacity" OR "Antioxidant*" OR "Oxidative
damage") AND ("Repeated*" OR "Repeated measure" OR "Repeatability*" OR
"Longitudinal*" OR "Longitudinal study" OR "Within Individual" OR "Within Individual
study") AND ("Wildlife*" OR "Wild animal" OR "Wild population" OR "Wild*" OR "Bird*"
OR "Avian*" OR "Invertebrate*" OR "Fish*" OR "Herpetofauna*" OR "Vertebrate*" OR
"Mammal*") NOT ("Human*" OR "Woman*" OR "Women*" OR "Man*" OR "Children*" OR
"Patients*" OR "People*" OR "Men*" OR "Child*")

This search string was used to collect literature from both the Web of Science core collection
database (last search 7[th] of December 2023) and Scopus database (last search 4[th] of June

2024). After removing duplicates, we recorded the number of results for each iteration following
the Preferred Reporting Items for Systematic Reviews and Meta-analysis (PRISMA) framework.
The PRISMA diagram summarising the selection process is shown in Figure S2. We were able to
include effect sizes from 22 studies (28,31,32,46–51,56–61,67–73).

Inclusion criteria

To be included in the meta-analysis, studies had to meet the following criteria:

1)   Measured at least one biomarker of oxidative stress (oxidative damage or antioxidant
defence).
2)   Took at least two repeated oxidative stress measurements from the same individual,
with the time interval between measurements reported.
3)   Focused on non-human organisms.
4)   Provided access to raw individual-level data, either publicly available or obtained via
author correspondence.
We included both experimental and observational studies across various environments to ensure
broad ecological and taxonomic representation.

Screening process

388   We used a two-step screening process to identify eligible studies. First, titles and abstracts were
389   screened using the "MetaGear" (74) package in R (75) to identify studies involving oxidative
390   stress in non-human organisms. Because repeat sampling was not always explicitly stated in
391   abstracts, we next screened the full texts of selected studies, verifying in the methods sections
392   that repeated measurements on the same individuals were conducted.

Data extraction

From each eligible study, we extracted biological and methodological information, including
species, taxonomic class, sample size, number of samples per individual (mean), time between
repeated samples (mean), oxidative stress biomarkers measured, environment, and study type
(manipulated vs. non-manipulated). For experimental studies, we separated treatment and control
groups, and calculated repeatability estimates independently for each group. To ensure
comparability across study types, control groups from experimental studies were pooled with non-
experimental studies and were referred to as non-manipulated, whereas treatment groups within
experimental studies were referred to as manipulated.

Environment was categorised as wild, captive, wild captive (wild individuals brought into captivity
for the purposes of the study), or livestock (Animals used for agricultural purposes, including
horses, cattle and sheep). Biomarkers were classified based on whether they measured oxidative
damage (lipid peroxidation, protein damage, DNA damage, or nonspecific damage) or antioxidant
defences (enzymatic or non-enzymatic) (see Table S5).

Repeatability estimation

To calculate within-individual repeatability of oxidative stress measurements, we obtained the raw
data from each study. For studies where raw data was not directly available, we contacted the
authors (Table S4). We estimated repeatability by determining the intraclass correlation
coefficient (ICC) using ANOVAs based on equations 4 and 5 from Nakagawa & Schielzeth (2010)
(76), along with the standard error and 95% confidence intervals. This ANOVA-based method is
recommended when repeatability is very low (<0.05) which was often the case in our dataset, as
the linear mixed model (LMM) approach tends to overestimate low repeatability values (26,76).
While we used the ANOVA-based approach for our main analyses, we confirmed the consistency
of the results with LMM-derived estimates (Fig. S3). Importantly, rather than calculating

conditional repeatability by including covariates within the repeatability model itself, we stratified
the data by relevant variables, specifically by sex, oxidative stress biomarker, and study type
(manipulation vs. non-manipulation) to calculate separate ICC values for each subset. This
approach allowed us to subsequently use these variables as moderators in the meta-analysis to
explore potential sources of variation in repeatability. This yielded 123 effect sizes from 22
species across two classes from 22 studies that were published between 2013 and 2023 (species
listed in Table S6).

Oxidative biomarker correlations.

425 For studies measuring multiple oxidative stress biomarkers within the same individuals, we
426 calculated Pearson's correlation coefficients between biomarker pairs using the cor function in R.
427 These values were included in an additional meta-analysis to assess the consistency of
428 biomarker co-variation across studies.

Meta-analysis approach.

All analysis were formed in R version 4.3.3 (75).

All meta-analyses were conducted using a multi-level meta-analysis structure with the metafor
package (77). ICC and Pearson's r values were standardised using Fishers Z transformation (76)
and then were back transformed for interpretation in figures and tables. Corresponding sampling
variances were calculated using the escalc function.

We first estimated the overall ICC using an intercept-only random effects model. We then tested
whether variation in repeatability could be explained by moderators including study type,
environment, taxonomic class, oxidative stress biomarker, mean time between measurements
(log-transformed), and the average number of samples per individual (log-transformed).
Subgroup analysis further compared between oxidative damage and antioxidant defence
categories using the respective biomarker type as a moderator in each analysis (Table S5).

The meta-analysis model exploring biomarker correlations used the same model structure, with
biomarker pair identity as a categorical moderator.

All models included study ID and observation ID as random effects to account for non-
independence and hierarchical structure (76,78). Total relative heterogeneity and the contribution
of each random effect was quantified using the i2_ml function from the OrchaRd package (79).

Lastly, we evaluated potential publication bias by fitting two multilevel meta-regression models,
with the square root of inverse sampling variance and mean-centred year of publication as
moderators to test for small study effects and time lag bias respectively (Fig S4) (78).

The methods for the simulation can be found in the supplementary materials, Appendix two.

Conditional repeatability comparison

To evaluate whether controlling for biological covariates alters repeatability estimates of oxidative
stress biomarkers, we performed an additional meta-analysis using a subset of studies (n = 8) for
which the raw data included information about age, and in some cases, sex and treatment group.
For each oxidative stress biomarker in these studies, we estimated repeatability under four model
conditions: (1) unadjusted (raw, no covariates), (2) age adjusted, (3) age and sex adjusted, and
(4) age and treatment group adjusted.

Repeatability was calculated using the rptR (76,80) package in R, employing an LMM-based
estimation approach. This method allows for covariates to be included as fixed effects within the
model, providing estimates of conditional repeatability. We then performed a meta-analysis using

repeatability model type as a moderator to assess whether controlling for these covariates
influenced repeatability estimates.

462
463
Availability of data and materials

465 The code and data used in this study will be made available online following peer review at the
466 time when the manuscript is accepted for publication

Funding

468 This project was funded through a grant from Wild Animal Initiative to Davide M Dominoni. Jelle
469 Boonekamp was supported through a grant from the Leverhulme Trust (RPG-2024-207).

Acknowledgments

471 Thanks to all of the authors who responded to our data enquiries, including Quentin Schull, Sarah
472 Guindre-Parker, Helga Sauerwein, Melissah Rowe, Brenna Gormally, Aurora Ramierez-Perez,
473 Mireia Blanco, Shelia Stivanin, Valeria Marasco, and Martin Tesicky.

Author contributions

475 Rachel Reid, Davide M Dominoni and Jelle Boonekamp conceived of and designed the study.
476 Rachel Reid and Jelle Boonekamp developed the simulation. Rachel Reid ran the meta-analyses
477 and drafted the manuscript with help from all co-authors. All authors read and approved the final
478 version.

479
References
481

1. Pole A, Dimri M, P. Dimri G, Department of Biochemistry and Molecular Medicine, School of
Medicine and Health Sciences, The George Washington University, Washington DC, USA.
Oxidative stress, cellular senescence and ageing. AIMS Molecular Science. 2016;3(3):300–
24.

2. Leyane TS, Jere SW, Houreld NN. Oxidative Stress in Ageing and Chronic Degenerative
Pathologies: Molecular Mechanisms Involved in Counteracting Oxidative Stress and Chronic
Inflammation. IJMS. 2022 June 30;23(13):7273.

3. Gil Del Valle L, Gravier Hernández R, Delgado Roche L, León Fernández OS. Oxidative
Stress in the Aging Process: Fundamental Aspects and New Insights. In: Hepel M,
Andreescu S, editors. ACS Symposium Series [Internet]. Washington, DC: American
Chemical Society; 2015 [cited 2026 Jan 12]. p. 177–219. Available from:
https://pubs.acs.org/doi/abs/10.1021/bk-2015-1200.ch006

4. Kregel KC, Zhang HJ. An integrated view of oxidative stress in aging: basic mechanisms,
functional effects, and pathological considerations. American Journal of Physiology-
Regulatory, Integrative and Comparative Physiology. 2007 Jan;292(1):R18–36.

5. Monaghan P, Metcalfe NB, Torres R. Oxidative stress as a mediator of life history trade-offs:
mechanisms, measurements and interpretation. Ecology Letters. 2009 Jan;12(1):75–92.

6. Costantini D, Rowe M, Butler MW, McGraw KJ. From molecules to living systems: historical
and contemporary issues in oxidative stress and antioxidant ecology. Functional Ecology.
2010 Oct;24(5):950–9.

7. Costantini D, Verhulst S. Does high antioxidant capacity indicate low oxidative stress?
Functional Ecology. 2009 June;23(3):506–9.

8. Von Zglinicki T. Oxidative stress shortens telomeres. Trends in Biochemical Sciences. 2002
July;27(7):339–44.

9. Itri R, Junqueira HC, Mertins O, Baptista MS. Membrane changes under oxidative stress: the
impact of oxidized lipids. Biophys Rev. 2014 Mar;6(1):47–61.

10. Sharma A, Gupta P, Prabhakar PK. Endogenous Repair System of Oxidative Damage of
DNA. CCB. 2019 July 12;13(2):110–9.

11. Campisi J. Cellular senescence and apoptosis: how cellular responses might influence aging
phenotypes. Experimental Gerontology. 2003 Jan;38(1–2):5–11.

12. Chen JH, Hales CN, Ozanne SE. DNA damage, cellular senescence and organismal ageing:
causal or correlative? Nucleic Acids Research. 2007 Nov 26;35(22):7417–28.

13. Luo J, Mills K, Le Cessie S, Noordam R, Van Heemst D. Ageing, age-related diseases and
oxidative stress: What to do next? Ageing Research Reviews. 2020 Jan;57:100982.

14. Liguori I, Russo G, Curcio F, Bulli G, Aran L, Della-Morte D, et al. Oxidative stress, aging,
and diseases. CIA. 2018 Apr;Volume 13:757–72.

15. Reichert S, Stier A. Does oxidative stress shorten telomeres *in vivo*? A review. Biol Lett.
2017 Dec;13(12):20170463.

16. Speakman JR, Blount JD, Bronikowski AM, Buffenstein R, Isaksson C, Kirkwood TBL, et al.
Oxidative stress and life histories: unresolved issues and current needs. Ecology and
Evolution. 2015 Dec;5(24):5745–57.

17. Wilbourn RV, Moatt JP, Froy H, Walling CA, Nussey DH, Boonekamp JJ. The relationship
between telomere length and mortality risk in non-model vertebrate systems: a meta-
analysis. Phil Trans R Soc B. 2018 Mar 5;373(1741):20160447.

18. Armstrong E, Boonekamp J. Does oxidative stress shorten telomeres in vivo? A meta-
analysis. Ageing Research Reviews. 2023 Mar;85:101854.

19. Smith SM, Nager RG, Costantini D. Meta-analysis indicates that oxidative stress is both a
constraint on and a cost of growth. Ecology and Evolution. 2016 May;6(9):2833–42.

20. Boonekamp JJ, Dijkstra R, Dijkstra C, Verhulst S. Canalization of development reduces the
utility of traits as fitness biomarkers: feather fault bars in nestling birds. Blanckenhorn W,
editor. Functional Ecology. 2017 Mar;31(3):719–27.

21. Wilson AJ. How should we interpret estimates of individual repeatability? Evolution Letters.
2018 Feb 1;2(1):4–8.

22. Nespolo RF, Franco M. Whole-animal metabolic rate is a repeatable trait: a meta-analysis.
Journal of Experimental Biology. 2007 June 1;210(11):2000–5.

23. Taff CC, Schoenle LA, Vitousek MN. The repeatability of glucocorticoids: A review and meta-
analysis. General and Comparative Endocrinology. 2018 May;260:136–45.

24. Schoenemann KL, Bonier F. Repeatability of glucocorticoid hormones in vertebrates: a meta-
analysis. PeerJ. 2018 Feb 21;6:e4398.

25. Cauchoix M, Chow PKY, Van Horik JO, Atance CM, Barbeau EJ, Barragan-Jason G, et al.
The repeatability of cognitive performance: a meta-analysis. Phil Trans R Soc B. 2018 Sept
26;373(1756):20170281.

26. Kärkkäinen T, Briga M, Laaksonen T, Stier A. Within-individual repeatability in telomere
length: A meta-analysis in nonmammalian vertebrates. Molecular Ecology. 2022
Dec;31(23):6339–59.

27. Cláudia Norte A, Sheldon B, Paulo Sousa J, Albino Ramos J. Repeatability and Method-
Dependent Variation of Blood Parameters in Wild-Caught Great Tits *Parus major*. Acta
Ornithologica. 2008 June;43(1):65–75.

28. Récapet C, Arrivé M, Doligez B, Bize P. Antioxidant capacity is repeatable across years but
does not consistently correlate with a marker of peroxidation in a free-living passerine bird. J
Comp Physiol B [Internet]. 2019 Mar 8 [cited 2026 Jan 9]; Available from:
http://link.springer.com/10.1007/s00360-019-01211-1

29. A. Cohen A, Hau M, Wikelski M. Stress, Metabolism, and Antioxidants in Two Wild Passerine
Bird Species. Physiological and Biochemical Zoology. 2008 July;81(4):463–72.

30. Alonso-Álvarez C, Pérez-Rodríguez L, García JT, Viñuela J, Mateo R. Age and Breeding
Effort as Sources of Individual Variability in Oxidative Stress Markers in a Bird Species.
Physiological and Biochemical Zoology. 2010 Jan;83(1):110–8.

31. Bodey TW, Cleasby IR, Blount JD, Vigfusdottir F, Mackie K, Bearhop S. Measures of
oxidative state are primarily driven by extrinsic factors in a long-distance migrant. Biol Lett.
2019 Jan;15(1):20180750.

32. Cram DL, Blount JD, Young AJ. The oxidative costs of reproduction are group-size
dependent in a wild cooperative breeder. Proc R Soc B. 2015 Nov 22;282(1819):20152031.

33. Jenni-Eiermann S, Jenni L, Smith S, Costantini D. Oxidative Stress in Endurance Flight: An
Unconsidered Factor in Bird Migration. McGraw K, editor. PLoS ONE. 2014 May
15;9(5):e97650.

34. Slos S, Stoks R. Predation risk induces stress proteins and reduces antioxidant defense.
Functional Ecology. 2008 Aug;22(4):637–42.

35. Van De Crommenacker J, Komdeur J, Burke T, Richardson DS. Spatio-temporal variation in
territory quality and oxidative status: a natural experiment in the Seychelles warbler
(Acrocephalus sechellensis): Territory quality-related oxidative costs in a wild passerine.
Journal of Animal Ecology. 2011 May;80(3):668–80.

36. Chainy GBN, Paital B, Dandapat J. An Overview of Seasonal Changes in Oxidative Stress
and Antioxidant Defence Parameters in Some Invertebrate and Vertebrate Species.
Scientifica. 2016;2016:1–8.

37. Rathwa SD, Vasava AA, Pathan MM, Madhira SP, Patel YG, Pande AM. Effect of season on
physiological, biochemical, hormonal, and oxidative stress parameters of indigenous sheep.
Vet World. 2017 June;10(6):650–4.

38. Goldfarb AH, Garten RS, Waller J, Labban JD. Day to Day Variability and Reliability of Blood
Oxidative Stress Markers within a Four-Week Period in Healthy Young Men. Journal of
Biomarkers. 2014 Oct 15;2014:1–7.

39. Mallard AR, Hollekim-Strand SM, Ingul CB, Coombes JS. High day-to-day and diurnal
variability of oxidative stress and inflammation biomarkers in people with type 2 diabetes
mellitus and healthy individuals. Redox Report. 2020 Jan 1;25(1):64–9.

40. García-Closas R, Berenguer A, Tormo MJ, Sánchez MJ, Quirós JR, Navarro C, et al. Dietary
sources of vitamin C, vitamin E and specific carotenoids in Spain. Br J Nutr. 2004
June;91(6):1005–11.

41. Landete JM. Dietary Intake of Natural Antioxidants: Vitamins and Polyphenols. Critical
Reviews in Food Science and Nutrition. 2013 Jan;53(7):706–21.

42. Beaulieu M, Costantini D. Biomarkers of oxidative status: missing tools in conservation
physiology. Conservation Physiology. 2014 May 12;2(1):cou014–cou014.

43. Hõrak P, Cohen A. How to measure oxidative stress in an ecological context: methodological
and statistical issues. Functional Ecology. 2010 Oct;24(5):960–70.

44. Costantini D. Understanding diversity in oxidative status and oxidative stress: the
opportunities and challenges ahead. Journal of Experimental Biology. 2019 July
1;222(13):jeb194688.

45. Christensen LL, Selman C, Blount JD, Pilkington JG, Watt KA, Pemberton JM, et al. Plasma
markers of oxidative stress are uncorrelated in a wild mammal. Ecology and Evolution. 2015
Nov;5(21):5096–108.

46. Orquera-Arguero KG, Casasús I, Ferrer J, Blanco M. Beef cows' performance and metabolic
response to short nutritional challenges in different months of lactation. Research in
Veterinary Science. 2023 June;159:26–34.

47. Marasco V, Stier A, Boner W, Griffiths K, Heidinger B, Monaghan P. Environmental
conditions can modulate the links among oxidative stress, age, and longevity. Mechanisms of
Ageing and Development. 2017 June;164:100–7.

48. Velázquez-Cantón E, Cruz-Rodríguez NDL, Zarco L, Rodríguez A, Ángeles-Hernández JC,
Ramírez-Orejel JC, et al. Effect of Selenium and Vitamin E Supplementation on Lactate,
Cortisol, and Malondialdehyde in Horses Undergoing Moderate Exercise in a Polluted
Environment. Journal of Equine Veterinary Science. 2018 Oct;69:136–44.

49. Těšický M, Krajzingrová T, Świderská Z, Syslová K, Bílková B, Eliáš J, et al. Longitudinal
evidence for immunosenescence and inflammaging in free-living great tits. Experimental
Gerontology. 2021 Oct;154:111527.

50. Eikenaar C, Winslott E, Hessler S, Isaksson C. Oxidative damage to lipids is rapidly reduced
during migratory stopovers. Galván I, editor. Functional Ecology. 2020 June;34(6):1215–22.

51. Christensen LL, Selman C, Blount JD, Pilkington JG, Watt KA, Pemberton JM, et al. Marker-
dependent associations among oxidative stress, growth and survival during early life in a wild
mammal. Proc R Soc B. 2016 Oct 12;283(1840):20161407.

52. Musakhanian J, Rodier JD, Dave M. Oxidative Stability in Lipid Formulations: a Review of the
Mechanisms, Drivers, and Inhibitors of Oxidation. AAPS PharmSciTech. 2022 May
20;23(5):151.

53. Ayuso M, Van Cruchten S, Van Ginneken C. A Medium-Throughput System for In Vitro
Oxidative Stress Assessment in IPEC-J2 Cells. IJMS. 2020 Oct 1;21(19):7263.

54. Shiva Shankar Reddy CS, Subramanyam MVV, Vani R, Asha Devi S. In vitro models of
oxidative stress in rat erythrocytes: Effect of antioxidant supplements. Toxicology in Vitro.
2007 Dec;21(8):1355–64.

55. Halliwell B. Oxidative stress in cell culture: an under-appreciated problem? FEBS Letters.
2003 Apr 10;540(1–3):3–6.

56. Noguera JC, Kim SY, Velando A. Family-transmitted stress in a wild bird. Proc Natl Acad Sci
USA. 2017 June 27;114(26):6794–9.

57. Viblanc VA, Schull Q, Roth JD, Rabdeau J, Saraux C, Uhlrich P, et al. Maternal oxidative
stress and reproduction: Testing the constraint, cost and shielding hypotheses in a wild
mammal. Gremillet D, editor. Functional Ecology. 2018 Mar;32(3):722–35.

58. Schull Q, Viblanc VA, Stier A, Saadaoui H, Lefol E, Criscuolo F, et al. The oxidative debt of
fasting: evidence for short to medium-term costs of advanced fasting in adult king penguins.
Journal of Experimental Biology. 2016 Jan 1;jeb.145250.

59. Sauerwein H, Blees T, Zamarian V, Catozzi C, Müller U, Sadri H, et al. Acute phase proteins
and markers of oxidative status in water buffalos during the transition from late pregnancy to
early lactation. Veterinary Immunology and Immunopathology. 2020 Oct;228:110113.

60. Gormally BMG, Estrada R, Yin H, Romero LM. Recovery from repeated stressors:
Physiology and behavior are affected on different timescales in house sparrows. General and
Comparative Endocrinology. 2019 Oct;282:113225.

61. Kolling GJ, Stivanin SCB, Gabbi AM, Machado FS, Ferreira AL, Campos MM, et al. Milk
production and hematological and antioxidant profiles of dairy cows supplemented with
oregano and green tea extracts as feed additives. Revista Brasileira de Zootecnia. 2022 Nov
22;51:e20210150.

62. Skrip MM, McWilliams SR. Oxidative balance in birds: an atoms-to-organisms-to-ecology
primer for ornithologists. J Field Ornithol. 2016 Mar;87(1):1–20.

63. Ji LL, Yeo D. Oxidative stress: an evolving definition. Fac Rev [Internet]. 2021 Feb 9 [cited
2026 Jan 9];10. Available from: https://facultyopinions.com/prime/reports/b/10/13/

64. Lushchak VI. Free radicals, reactive oxygen species, oxidative stress and its classification.
Chemico-Biological Interactions. 2014 Dec;224:164–75.

65. Bell AM, Hankison SJ, Laskowski KL. The repeatability of behaviour: a meta-analysis. Animal
Behaviour. 2009 Apr;77(4):771–83.

66. Polverino G, Buchholz KM, Goulet CT, Michelangeli M, Chapple DG. Temporal repeatability
of behaviour in a lizard: implications for behavioural syndrome studies. Evol Ecol. 2023
June;37(3):401–18.

67. Hau M, Haussmann MF, Greives TJ, Matlack C, Costantini D, Quetting M, et al. Repeated
stressors in adulthood increase the rate of biological ageing. Front Zool. 2015;12(1):4.

68. Herborn KA, Daunt F, Heidinger BJ, Granroth-Wilding HMV, Burthe SJ, Newell MA, et al.
Age, oxidative stress exposure and fitness in a long-lived seabird. Williams T, editor.
Functional Ecology. 2016 June;30(6):913–21.

69. Guindre-Parker S, Rubenstein DR. No short-term physiological costs of offspring care in a
cooperatively breeding bird. Journal of Experimental Biology. 2018 Jan 1;jeb.186569.

70. Guindre-Parker S, Baldo S, Gilchrist HG, Macdonald CA, Harris CM, Love OP. The oxidative
costs of territory quality and offspring provisioning. J of Evolutionary Biology. 2013
Dec;26(12):2558–65.

71. Rowe M, Pierson KL, McGraw KJ. Exploratory behavior is associated with plasma carotenoid
accumulation in two congeneric species of waterfowl. Behavioural Processes. 2015
June;115:181–90.

72. Skrip MM, Seeram NP, Yuan T, Ma H, McWilliams SR. Dietary antioxidants and flight
exercise in female birds affect allocation of nutrients to eggs: how carry-over effects work.
Journal of Experimental Biology. 2016 Sept 1;219(17):2716–25.

73. Vitikainen EIK, Cant MA, Sanderson JL, Mitchell C, Nichols HJ, Marshall HH, et al. Evidence
of Oxidative Shielding of Offspring in a Wild Mammal. Front Ecol Evol. 2016 May 18;4:58.

74. Lajeunesse MJ. Facilitating systematic reviews, data extraction and meta-analysis with the
METAGEAR package for R. Fitzjohn R, editor. Methods Ecol Evol. 2016 Mar;7(3):323–30.

75. R Core Team. R: A language and environment for statistical computing. 2024;

76. Nakagawa S, Schielzeth H. Repeatability for Gaussian and non-Gaussian data: a practical
guide for biologists. Biological Reviews. 2010 Nov;85(4):935–56.

77. Viechtbauer W. Conducting Meta-Analyses in *R* with the **metafor** Package. J Stat Soft
[Internet]. 2010 [cited 2026 Jan 9];36(3). Available from: http://www.jstatsoft.org/v36/i03/

78. Nakagawa S, Lagisz M, Jennions MD, Koricheva J, Noble DWA, Parker TH, et al. Methods
for testing publication bias in ecological and evolutionary meta-analyses. Methods Ecol Evol.
2022 Jan;13(1):4–21.

79. Nakagawa S, Lagisz M, O'Dea RE, Pottier P, Rutkowska J, Senior AM, et al. orchaRd 2.0:
An R package for visualising meta-analyses with orchard plots. Methods Ecol Evol. 2023
Aug;14(8):2003–10.

80. Stoffel MA, Nakagawa S, Schielzeth H. rptR: repeatability estimation and variance
decomposition by generalized linear mixed-effects models. Goslee S, editor. Methods Ecol
Evol. 2017 Nov;8(11):1639–44.