Generation of miR-141/200c conditional knockout mice from knockout-first, reporter-tagged parent and functional validation of the floxed allele

Sanjeev Kumar Yadav[1#], Rashmi Srivastava[1#], Mary-Katherine Cormier[1], Katie Lowther[2], Siu-Pok Yee[2,*], and Rajkumar Verma[1,*]

[1]Department of Neurosciences, UConn Health Farmington CT 06030

[2]Center for Mouse Genome Modification, UConn Health, Farmington CT 06030

[#]equal contribution

*Corresponding author

Rajkumar Verma

Assistant professor

Email: raverma@uchc.edu

ORCID ID: https://orcid.org/0000-0002-1123-6508

Sui-Pok Yee

Associate Professor

Email: syee@uchc.edu

ORCID: https://orcid.org/0000-0002-2427-401X

ABSTRACT

MicroRNAs (miRNAs) of the miR-200 family specifically miR-141 and miR-200c regulate neurogenesis, differentiation, and epithelial mesenchymal transitions in development. Dysregulation of these miRNAs is associated with several diseases including cancer and stroke. The Mirc13tm1Mtm/Mmjax mouse line, which targets the miR-141/200c cluster, was originally generated and described by Park et al. 2012 as a knockout-first, reporter-tagged insertion with conditional potential (conditional-ready) mouse line. To harness the full potential of this mouse line, it would require a two-step breeding process: breed with FLP mice to excise lacZ/neo cassettes followed by breeding with Cre to delete the floxed miRNA cluster(Park, et al. 2012). However, many studies either bypassed removal of the lacZ/Neo cassettes and treated the mouse line as Mirc13 knockouts or bred directly with Cre mouse lines, which could lead to unpredictable recombination and genotypes. In this study, we show that the presence of the Neo cassette influences expression of neighboring genes. We demonstrate that it is essential to follow the two-step breeding plan.

KEYWORDS: miR-141/200c cluster, lacZ/neo cassettes, Mirc13, microRNA, FLPo deleter

INTRODUCTION

microRNAs (miRNAs) are small, noncoding RNAs (~22 nucleotides) that regulate gene expression post-transcriptionally and play pivotal roles in development and disease. Originally discovered in C. elegans(Lee, et al. 1993; Wightman, et al. 1993) and later shown to be evolutionarily conserved across species(Pasquinelli, et al. 2000), miRNAs are now recognized as critical regulators in cardiovascular(Small, et al. 2010; van Rooij and Kauppinen 2014), neurological(Wu, et al. 2016; Verma, et al. 2018), and oncogenic pathways(Humphries and Yang 2015; Jin, et al. 2017), holding significant potential for therapeutic applications(Janssen, et al. 2013; Verma, et al. 2018) While cell-based studies have elucidated many of these mechanisms, in vivo models remain essential to fully understand the functions of miRNAs under physiological and pathological contexts.

Park et al. established a key genetic resource for in vivo miRNA investigation by generating conditional knockout-first mouse lines for conserved miRNAs, including the miR-141/200c cluster (Mirc13tm1Mtm/Mmjax)(Park, et al. 2012). This cluster, located on mouse chromosome 6, encodes two members of the miR-200 family implicated in neurogenesis(Choi, et al. 2008; Beclin, et al. 2016), differentiation(Zhang, et al. 2015), epithelial–mesenchymal transition(Tseng, et al. 2017), cancer(Chen, et al. 2013; Jin, et al. 2017), stroke(Small, et al. 2010; Yadav, et al. 2024), and Alzheimer's disease(Wu, et al. 2016). To make miR-141/200c cluster KO mice, Park et al. used a targeting vector to incorporate a lacZ reporter and β-actin-neo selection cassette flanked by FRT sites and loxP sites surrounding the miR-141/200c cluster. This enabled a two-step recombination strategy: first, FLP-mediated removal of the lacZ/neo cassette generates a clean conditional (floxed) allele, and then Cre-mediated excision removes the miR-141/200c cluster(Park, et al. 2012). However, many subsequent studies bypassed the essential FLP recombination step(Guan, et al. 2018; Hoefert, et al. 2018; Ji, et al. 2018; Liu, et al.

2018; Wu, et al. 2019; Xu, et al. 2025). In such cases, the lacZ/neo selection cassette

remains in the targeted allele, which can transcriptionally silence neighboring

genes(Testa, et al. 2004; Raja, et al. 2024) or alter local expression patterns due to the

strong β-actin promoter driving the cassette(Pham, et al. 1996), thereby confounding

phenotypic interpretation. Other studies attempted cassette removal using FLPe deleter

strains; however, several commonly used FLPe lines exhibit variable recombination

efficiency, which may lead to incomplete or mosaic excision depending on the deleter

strain and genomic target locus(Tran, et al. 2017; Mostofa, et al. 2022; Tran, et al. 2024).

Consequently, a small proportion of animals may retain the selection cassette despite

FLPe breeding, resulting in heterogeneous genotypes within experimental cohorts.

Several studies have reported that residual selection cassettes can unintentionally

suppress the expression of nearby genes(Olson, et al. 1996; Pham, et al. 1996; Skarnes,

et al. 2011). In the case of the miR-141/200c cluster, genes located in close proximity—

including *Ptpn6, Phb2, Atn1, Eno2,* and *Eng1* (www.Ensembl.org)—may be affected.

Because these genes are known to play important roles in neuronal function and

mitochondrial integrity(Whitton, et al. 1993; Bernabeu, et al. 2009; Xu, et al. 2014; Isgrò,

et al. 2015; He, et al. 2023), unintended suppression could lead to misleading

interpretations or the apparent loss of phenotype in knockout mice generated without prior

removal of the neo cassette.

To overcome these limitations, we optimized the original "knockout-first" design

suggested by Park et al. 2012(Park, et al. 2012) by employing high-efficiency FLPo

deleter mice instead of FLPe or β-actin-FLP strains(Raymond and Soriano 2007; Wu, et

al. 2009; Kranz, et al. 2010). FLPo recombinase exhibits greater thermostability and

recombination efficiency in mammalian cells(Raymond and Soriano 2007; Wu, et al.

2009; Kranz, et al. 2010), ensuring complete excision of the lacZ/neo cassette and

restoration of an unperturbed floxed allele before Cre-mediated deletion. This refinement

minimizes off-target silencing, preserves adjacent gene expression, and produces

genetically clean global or tissue-specific miR-141/200c knockouts. Here, we present a

validated, optimized, and detailed breeding strategy using FLPo deleter mice that ensures

complete removal of the lacZ/neo cassette and precise deletion of the miR-141/200c

cluster. This work establishes a robust framework for generating unambiguous cell-

specific or conditional knockouts of the miR-141/200c cluster, facilitating accurate studies

of miRNA function in neural and systemic disease models.

MATERIAL AND METHODS

Mouse Strains

All mouse strains used in this study were obtained from the Jackson Laboratory (Bar

Harbor, ME): Mirc13^{tm1Mtm}/Mmjax line (MMRRC stock #34657(Park, et

al. 2012)) (hereon referred as *Mirc13^{tm1}* line); B6.129S4-

Gt(ROSA)26Sor^{tFLP})Sor/J (ROSA-FLPo; Jax# 012930)(Raymond and

Soriano 2007); 129S1/Sv-Hprt1$^{tm1(CAG-cre)Mnn}$/J) (Hprt-Cre; Jax#

004302)(Tang, et al. 2002) and B6.129P2-Lyz2$^{tm1(cre)Ifo}$/J) (LysMCre;

Jax # 004781)(Clausen, et al. 1999). All mice were maintained under standard

conditions at ambient temperature and humidity, with ad libitum access to pellet food

and water. All animal studies were approved by the Institutional Animal Care and Use

Committee at the University of Connecticut Health Center.

Genotyping of Mouse Strains

Tail biopsies (~4 mm) were collected in DNase-free microtubes and incubated in 50 ml of

alkali buffer (25mM NaOH and 0.2mM EDTA) at 100°C for one hour and an equal volume

of neutralization buffer (40mM Tris buffer, pH 7.5) was added. One microliter of the DNA

was used for PCR genotyping with 2× JumpStart™ Taq ReadyMix (Sigma-Aldrich

P2893). Primer pairs are listed in Table 1. PCR was performed according to the

manufacturer's instructions. PCR products were fractionated on 2% agarose gels

prepared in 1× Tris-Acetate-EDTA (TAE) buffer (Bio-Rad 1610743) with SYBR™ Safe

DNA stain (Invitrogen S33102) and visualized under UV illumination.

β-Galactosidase (lacZ) Staining

A β-galactosidase staining kit (Sigma-Aldrich GALS-1KT) was used to assess lacZ

reporter expression. Mice were deeply anesthetized with Avertin (250 mg/kg, i.p.),

perfused with ice-cold PBS, and brains were collected for cryosections (10 µm, Superfrost

Plus slides). Sections were fixed with fixation solution (provided with kit) for 10 min, and

LacZ expression was determined using a β-galactosidase staining kit according to

manufacturer's instructions. Slides were mounted in 70% glycerol and imaged under

bright-field microscopy.

RNA Isolation, cDNA Synthesis, and Real-Time PCR

Total RNA was extracted from the olfactory bulb, frontal and parietal cortex,

hippocampus, cerebellum, heart, lung, and liver using TRIzol or miRVana kits (Ambion

Life Technologies, 15596018, AM1560). cDNA synthesis for miRNAs and mRNAs was

performed using the TaqMan MicroRNA Reverse Transcription Kit (Thermo Fisher

4366596) and iScript cDNA Synthesis Kit (Bio-Rad 1708891), respectively. TaqMan

assays included miR-141-3p (000463), miR-200c-3p (002300), U6snRNA (001973),

Ptpn6 (Mm00469153_m1), *Phb2* (Mm00476104_m1), *Atn1* (Mm00492256_m1), *Eno2*

(Mm00469062_m1), *Emg1* (Mm00496596_m1), and *Gapdh* (Mm99999915_g1) (Thermo

Fisher Scientific). PCR was performed using TaqMan Universal PCR Master Mix II (No

UNG; Thermo Fisher 4440040). Relative expression was determined by the 2^−ΔΔCt

method using U6 and Gapdh as normalizers for miRNA and mRNA targets, respectively.

RNA Fluorescence In Situ Hybridization (RNA-FISH)

RNA-FISH was performed using Molecular Instruments protocols. Fixed tissue sections were pre-hybridized at 37 °C and incubated overnight with 16 nM of the miR-141-3p probe. Following stringent SSCT washes, signal amplification was performed using snap-cooled hairpins (h1/h2; 6 pmol each) incubated overnight at room temperature. Sections were mounted with antifade DAPI UltraCruz Mounting Solution (Santa Cruz SC-24941) and imaged by fluorescence microscopy.

Statistical Analysis

Data are presented as mean ± SD. Relative expression values derived from the $2^{-\Delta\Delta Ct}$ method were used for statistical analysis. Statistical analyses were performed using GraphPad Prism (San Diego, CA). Comparisons between two groups were conducted using a two-tailed unpaired Student's *t*-test, while multiple group comparisons were analyzed using one-way ANOVA followed by Bonferroni post hoc correction. Data distribution was assessed for normality prior to applying parametric tests. A *p* value < 0.05 was considered statistically significant. Blinding was implemented during data acquisition and analysis where feasible. Each dot In graphs represent an individual mouse.

Target	Primer (F/R)	Primer Sequence (5'-3')	Amplicon size (bp)
Mirc13	Jax 11311 (F)	CAACAGCCTCTGACCTTTAACC	340 (WT)
	Jax 11312 (R)	CCTTCTGGGCAGACAAGAATAC	432 (floxed)
Mirc13	mir5Fn3 (F)	CCGGCAGGTGAAGGAGGAAG	1315 (WT)
	Jax 11312 (R)	CCTTCTGGGCAGACAAGAATAC	352 (KO)
lacZ	oIMR3054 (lacZ F)	ATCCTCTGCATGGTCAGGTC	315
	oIMR0040 (LacZ R)	CGTGGCCTGATTCATTCC	
neo	oIMR6916 (Neo F)	CTTGGGTGGAGAGGCTATTC	280
	oIMR6917 (Neo R)	AGGTGAGATGACAGGAGATC	
Flpo	oIMR8545 (F)	AAAGTCGCTCTGAGTTGTTAT	300
	oIMR8052 (R)	GCGAAGAGTTTGTCCTCAACC	
Rosa26	oIMR8545 (F)	AAAGTCGCTCTGAGTTGTTAT	603
	oIMR8546 (R)	GGAGCGGGAGAAATGGATATG	
Cre	oIMR1084 (F)	GCGGTCTGGCAGTAAAAACTA TC	100
	oIMR1085 (R)	GTCAAACAGCATTGCTGTCAC TT	

180   **Table1**. List of different sets of primer sequences (with amplicon size) used in PCR

181   amplification

RESULTS

Genomic Organization of the Mirc13 Mouse Lines and Genotyping of the Mouse Lines

The Mirc13[tm1Mtm]/Mmjax line (MMRRC stock #34657) mouse line, hereon referred as Mirc13[tm1] line, available from the Jax MMRCC is a knockout-first, reporter-tagged insertion with conditional potential (conditional-ready) mouse line. The miR-141/200c (Mirc13) cluster is located within intron 1 of the lncRNA, *GM15884*. As shown in Fig 1A, the targeted allele contains an *engrailed 2* (En2) splice acceptor followed by an internal ribosomal entry site (ires) sequence coupled to lacZ reporter sequence and polyadenylation signal sequence. This mouse line was originally generated by gene targeting in embryonic stem (ES) cells via homologous recombination. A β-actin-Neo cassette was placed 3' of the lacZ polyA sequence in the targeting vector to confer G418 selection to help screen for targeted ES cell clones. The Neo cassette and the lacZ reporter was separated by a single loxP site. The lacZ/Neo sequences were flanked by a pair of Frt sites and inserted into intron 1 of *GM15884* 5' of the floxed *Mirc13* cluster. With the polyadenylation sequence placed 3' of the lacZ sequence, it will terminate any *GM15884* 3'-downstream transcription, including the Mirc13 cluster, therefore, this mouse line is a double knockout of both *GM15884* and *Mirc13* cluster (knockout first). We first performed PCR genotyping of Mirc13[tm1/+] pups using primer pairs as shown in Table 1. As shown in Fig 2A, we detected amplicon of 340 bp and 432 bp specific to the Mirc13 WT and floxed allele, respectively. We were also able to amplify fragments of 315 bp and 280 bp specific to lacZ and Neo, respectively, confirming the genotype of the Mirc13[tm1/+] mice we obtained from MMRCC/Jax. We then bred the Mirc13[tm1/+] pups together to obtain Mirc13[tm1/tm1] pups for expression analyses as described below. PCR genotyping only detected the 432 bp fragment specific to the Mirc13 floxed allele as well as the lacZ and

neo specific amplicons. The absence of the Mirc13 WT specific amplicon of 340 bp

confirmed WT that these were homozygous Mirc13tm1 mice (Fig 2A).

Fig. 1 Genomic organization of Mirc13tm1 allele and after Flpo and Cre mediated

recombination. **A** The Mirc13tm1 mouse line was originally designed as a a knockout

first, reporter-tagged insertion with conditional potential (conditional-ready) mouse line

for Mirc13 (miR-141/200c cluster). Insertion of the en-lacZ gene trap cassette with

prevent any subsequent 3'-downstream transcription leading to the lack of GM15884

and Mirc13 expression. **B** When these mice were bred with Rosa26-Flpo, Flpo will

facilitate excision of the LacZ/Neo cassette to generate the Mirc13fl allele with wildtype

level of expression of both GM15884 and mir141/200c. **C** When Mirc13fl mice were

bred with Cre, Cre will mediate excision of the Mirc13 cluster sequence to generate

Mirc13- allele with wildtype GM15884 expression and the lack of miR141/200c

expression.

As described above, the lacZ/Neo sequences are flanked by a pair of Frt sites,

which can be excised by Flp recombinase resulting in the generation of functional *Mirc13*

conditional knockout (conditional later) mice expressing both *GM15884* and miR-141/200c as shown in Fig 1B. To accomplish this, we bred Mirc13^{tm1/+} mice with Rosa26^{Flpo/+} mice (Jax #012930). As shown in Fig 2B, genotyping results showed the absence of lacZ specific amplicon of 315 bp (lane 1 and 2) thereby confirming successful excision of the lacZ/Neo cassette and generation of Mirc13^{fl/+} pups. To generate Mirc13 knockout (KO) mice, we bred Mirc13^{fl/+} pups with Hprt^{CAG-Cre/+} mice (Fig1C). To detect the knockout allele, a forward primer (mir5Fn3) was designed to target the region upstream of the frt/loxP site and was paired with the Jax 11312 reverse primer downstream of the 3' loxP site. This primer pair will generate two amplicons, one specific to the WT allele (1.3 kb) and one specific to the knockout allele that is smaller in size due to Cre-mediated recombination (353 bp). As shown in Fig 2C, PCR genotyping showed that we were able to amplify the predicted fragments of 1.3 kb (lane 1) or 352 bp (lanes 2 and 3) specific to Mirc13 WT and KO alleles, respectively. We anticipated that we should obtain either Mirc13^{+/+} or Mirc13^{+/-} pups from this breeding pair. However, we did not detect both amplicons in heterozygous Mirc13^{+/-} mice. This is most likely because our PCR conditions favors amplification of the smaller amplicon specific to the knockout allele, and the pups shown in Fig 2C lanes 2 and 3, were Mirc13^{+/-} mice. We further bred Mirc13^{+/-} mice together and mice were genotyped using the same primers described above for the knockout allele and also with the original Jax primer pairs specific to the 3' loxP site in order to detect the presence or absence of the WT allele. We were able to obtain Mirc13^{-/-} mice as shown by PCR genotyping that was positive only for the KO-specific amplicon and no WT-specific amplicon was observed (Fig 2D, lane 2). The other littermates were

either Mirc13$^{+/-}$, which were positive for both amplicons (Fig 2D lanes 1 and 4) or Mirc13$^{+/+}$

as revealed by the presence of the WT-specific amplicon only (Fig 2D lane 3).

Fig. 2 PCR Genotyping of mice with different *Mirc13* alleles. **A** Generation of *Mirc13$^{tm1/tm1}$*

mice by breeding *Mirc13$^{tm1/+}$* mice, which contain lacZ/Neo cassette and floxed *Mirc13*

sequence, together; **B** Removal of lacZ and Neo cassette by breeding *Mirc13$^{tm1/+}$* mice

with *Rosa26$^{FLPo/+}$* mice. Positive *Mirc13$^{fl/+}$* mice, lacking the lacZ/Neo cassette (315 bp

amplicon), were marked by a red box; **C** Generation of heterozygous *Mirc13$^{+/-}$* (KO) mice

by breeding *Mirc13$^{fl/fl}$* with *Hprt$^{CAG-Cre/+}$* mice. Positive *Mirc13$^{+/-}$* mice, lacking a 432 bp

amplicon specific to the *Mirc1fl^{1/+}* allele, were marked by a red box; **D** Heterozygous

Mirc13^{+/-} mice were bred together to generate homozygous *Mirc13^{-/-}* global homozygous

KO mice. These mice were identified by the presence of a KO-specific amplicon (352 bp)

and the loss of wt specific amplicon (340 bp) and were marked by the red box. PCR were

performed using primer pairs as shown in Table 1 and amplicons specific for lacZ is 315

bp, *Mirc^+* is 340 bp, *Mirc13^{fl}* is 432 bp, *Mirc13^-* is 352 bp, Rosa26 is 603 bp, Flpo is 300

bp and Cre is 100 bp.

Expression Analysis of Homozygous Mirc13^{tm1} Mice

The En2 splice acceptor is designed to facilitate splicing between exon 1 of

GM15884 with the ires-lacZ reporter sequence. Thus, the targeted allele will generate a

fusion transcript consisting of exon 1 of the disrupted endogenous *GM15884*, a lncRNA,

ires and lacZ coding sequence. PCR genotyping confirmed presence of the lacZ reporter

(Fig 2A). Since the ires-lacZ gene trap cassette is inserted into intron 1 of the endogenous

GM15884 and placed under the *GM15884* transcriptional regulatory sequence, lacZ

expression will faithfully recapitulate spatiotemporal expression of the endogenous

GM15884 LncRNA, which should also reveal the spatiotemporal expression of the miR-

141/200c cluster (Mirc13) located within intron 1 of *GM15884*. However, the

polyadenylation sequence placed 3' with respect to the lacZ sequence will prevent further

3'-downstream transcription, including the Mirc13 cluster, to accomplish the knockout first

task. This mouse line is practically a double knockout of both *GM15884* and Mirc13

cluster.

We first examined lacZ expression, which should reveal endogenous expression of

GM15884 and miR-141-3p and miR-200c-3p, by x-gal staining of b-galactosidase, the

gene product of lacZ. Strong b-galactosidase activity was detected in the olfactory bulb,

with weaker staining near the lung–tracheal junction, and minimal expression in the heart

and liver (Fig. 3A). To further characterize expression of Mirc13, we isolated total RNA from the brain olfactory bulb and lungs of Mirc13$^{tm1/tm1}$ and Mirc13$^{+/+}$ mice and performed RT-qPCR to examine the level of miR-141-3p and miR-200c-3p expression. Our results showed that miR-141-3p and miR-200c-3p were highly expressed in the olfactory bulb and lung ($\Delta CT < 5$), whereas heart and liver exhibited lower expression levels ($\Delta CT > 10$) in Mirc13$^{+/+}$ mice, consistent with the miRNA expression atlas(Rishik, et al. 2025) (Fig. 3B and C). A significant and near-complete reduction of miR-141-3p (Fig 3D &E and miR-200c-3p (Fig. 3F & G) expression was observed in the olfactory bulb and lungs in Mirc13$^{tm1/tm1}$ mice. Data are presented as mean ± SD (*$p < 0.05$ vs. WT control).

Fig. 3 Expression analyses of *Mirc13$^{tm1/tm1}$* mice. **A**. LacZ expression as revealed by x–gal staining of olfactory bulb (OB) and lung (LG); **B**, **C** Basal expression of miR-141-3p and miR-200c-3p was analyzed by RT-qPCR in olfactory bulb, lung, heart and liver of wildtype mice. Relative expressions of miR-141-3p (**D** **&E**) and miR-200c-3p (**F & G**) were assessed by RT-qPCR in the brain olfactory bulb and lungs of Mirc13$^{tm1/tm1}$ KO mice compared to (Mirc13$^{+/+}$) littermate control mice. Data are presented as mean ± SD (*$p <$

0.05 vs. Mirc13$^{+/+}$ littermate control). Blue column denotes Mirc13$^{+/+}$ and yellow column

denotes Mirc13$^{tm1/tm1}$.

Presence of β-actin-Neo cassette leads to Transcriptional Interference of

Neighboring Genes

The Mirc13^{tm1} mouse line was originally generated by gene targeting in

embryonic stem cells via homologous recombination. A lacZ polyA sequence was

included in the targeting vector to confer G418 positive selection to screen for targeted

ES cell clones. However, the Neo cassette containing a strong ubiquitous β-actin

promoter was not removed prior to generation of the Mirc13^{tm1} mouse line. It has been

reported that insertion of a strong promoter within a gene would not only disrupt

endogenous expression, but it could also potentially lead to dysregulation of

neighboring genes and resulted in unanticipated phenotypic alterations(Olson, et al.

1996; Pham, et al. 1996; Skarnes, et al. 2011).

To further investigate the influence of the b-actin-Neo cassette on neighboring

gene expression, we identified 5 genes, Ptpn6, Phb2, Atn1, Eno2 and Emg1 that are

important for brain function, located within 50 kb from Mirc13. Total RNA was isolated

from olfactory bulb of Mirc13$^{+/+}$ and Mirc13$^{tm1/tm1}$ mice and the levels of neighboring gene

expression were characterized by RT-qPCR. As shown in Fig 4A-C , expression of Ptpn6,

Phb2, and Atn1 was significantly decreased (±>1.5 folds threshold, p<0.05) in

Mirc13$^{tm1/tm1}$ mice as compared to Mirc13$^{+/+}$ mice, while Eno2 and Emg1 showed a similar

decreasing trend (Fig 4D and E). In Mirc13$^{-/-}$ mice, the expression Ptpn6, Phb2, Atn1,

Eno2, and Emg1 did not show significantly change (±>1.5 folds threshold, p<0.05) as

compared to Mirc13$^{fl/fl}$ mice (Fig 4F to J). Our results showed that the presence of b-actin-

Neo cassette within GM15884 affects neighboring gene expression and it is therefore

essential to excise the cassette prior to obtaining transparent and interpretable subsequent data.

Fig. 4 Expression analysis of Mirc13 neighboring genes. Total RNA was isolated from olfactory bulb (OB) of Mirc13$^{tm1/tm1}$ and Mirc13$^{-/-}$ mice and corresponding negative littermate (Mirc13$^{+/+}$ and Mirc13$^{fl/fl}$). Expression of Ptpn6 (**A** and **F**), Phb2 (**B** and **G**), Atn1 (**C** and **H**), Eno2 (**D** and **I**), and Emg1 (**E** and **J**) were assessed by RT-qPCR. Data are presented as mean ± SD (*p < 0.05 vs. Mirc13$^{+/+}$ control. Light blue column denotes RNA obtained from Mirc13$^{+/+}$; Yellow column denotes RNA obtained from Mirc13$^{tm1/tm1}$; dark blue column denotes RNA from Mirc13$^{fl/fl}$ and red column denotes RNA isolated from Mirc13$^{-/-}$ mice.

Expression Analysis of miR-141-3p and miR-200c-3p in Mirc13$^{-/-}$ mice

We further characterized expression of miR-141-3p and miR-200c-3p in Mirc13$^{-/-}$ and their corresponding Mirc13$^{fl/fl}$ mice in the brain. We isolated total RNA from different regions of the brain, including olfactory bulb (OB), frontal cortex (FC), parietal cortex (PC), hippocampus (HP), and cerebellum (CB), from mice of both genotypes. Our results

showed both miR-141-3p and miR-200c-3p were highly expressed in the OB (ΔCT < 5; it is to be noted that lower ΔCT indicate higher expression), while their expression in other regions was very low (ΔCT > 10) (Fig 5A and B). Notably, the OB of Mirc13$^{-/-}$ mice showed no detectable expression of miR-141-3p by RNA-FISH (Fig 5C). This finding was further confirmed by RT-qPCR, which demonstrated a marked reduction of both miRNAs (miR-141-3p & miR-200c-3p) in the OB almost negligible (Fig 5D and I), along with significantly decreased expression in the FC (Fig 5E and J), PC (Fig 5F and K), HP (Fig 5G and L), and CB (Fig 5H and M) of Mirc13$^{-/-}$ mice as compared to Mirc13$^{fl/fl}$ mice.

Fig. 5 Expression analysis of miR-141-3p and miR-200c-3p in Mirc13$^{fl/fl}$ and Mirc13$^{-/-}$ mice. Total RNA was isolated from olfactory bulb (OB), frontal cortex (FC), parietal cortex (PC), hippocampus (HP), and cerebellum (CB) of Mirc13$^{fl/fl}$ and Mirc13$^{-/-}$ mice. Basal expression levels of miR-141-3p (**A**) and miR-200c-3p (**B**) were assessed by RT-qPCR

using RNA isolated from Mirc13$^{fl/fl}$ mice. (**C**) RNA-FISH was performed to detect

expression of miR-141-3p in OB of Mirc13$^{fl/fl}$ and Mirc13$^{-/-}$ mice. Total RNAs isolated from

OB (**D** and **I**); FC (**E** and **J**); PC (**F** and **K**); HP (**G** and **L**) and CB (**H** and **M**) were examined

for the relative level of miR-141-3p and miR-200c-3p, respectively, by RT-qPCR from

Mirc13$^{fl/fl}$ (dark blue column) and Mirc13$^{-/-}$ (red column) mice. Data are presented as mean

± SD (*$p < 0.05$ vs. Mirc13$^{fl/fl}$ control).

DISCUSSION

Generation of the Mirc13^{tm1} line by Park et al. 2012provided the first genetically

engineered model for the miR-141/200c cluster(Park, et al. 2012). In their design, a lacZ

reporter in association with the En2 splice acceptor and ires sequence, which is similar

to the gene trap design(Skarnes, et al. 1992), was aimed to reveal spatial expression of

the disrupted endogenous GM15884. Expression of the lacZ reporter is under the

transcriptional control of the endogenous GM15884 gene, and with the En2 splice

acceptor, lacZ sequence will fuse with exon 1 coding sequence of GM15884. Thus, x-gal

staining of the lacZ gene product, β-galactosidase, will faithfully recapitulate the

spatiotemporal expression of GM15884(Skarnes, et al. 1992) as well as the miR-

141/200c cluster. GM15884 is a lncRNA gene predicted by bioinformatics as described

in the Mouse Genome Database(Eppig, et al. 2005)

(https://www.informatics.jax.org/marker/key/326668). However, when we performed a

blast search of the mouse Expressed Taq Sequence database (dbEST)(Boguski, et al.

1993), we were able to identify a mouse cDNA clone, BB612857, of 646 bp. 5' sequence

of BB612857 is identical to predicted exon 1 and 2 of GM15884 the rest of the 3'-

downstream sequence identical to predicted GM15884 intron 2 sequence. It is well

established that EST are derived from transcribed genes (the transcriptome) and dbEST

was used to identify transcribed sequences without requiring a fully sequenced

genome(Boguski, et al. 1993). Identification of BB612857 confirms the identity of

GM15884. This result, together with the tissue-specific lacZ expression, suggests that

expression of GM15884/BB612857, as well as Mirc13, is under the control of a tissue-

specific Pol II promoter, and our lacZ expression observed in the olfactory bulb and lung

in the mice corroborated with the finding of Park et al.

Since the Mirc13^{tm1} mouse line was original generated by homologous

recombination using ES cells, A β-actin-neo selectable marker was included in the original

targeting vector and was retained in the Mirc13^{tm1} mouse line. It has been reported that

the Neo cassette with a strong ubiquitous promoter could influence neighboring gene

expression(Pham, et al. 1996) and therefore alter phenotype of the mice(Olson, et al.

1996; Pan, et al. 2016). Park et al explicitly recommended removal of Neo cassette prior

to subsequent use of the mouse line. The Neo cassette together with the lacZ sequence

were flanked by FRT sites and could be excised easily by breeding the Mirc13^{tm1} mice

with FLP mice. Additional methods by which one could excise the Frt flanked lacZ/Neo

cassette is either by electroporation of Flpo mRNA into one-cell embryos(Hashimoto and

Takemoto 2015) or infection of one-cell embryos with recombinant adeno-associated

viruses expressing Flpo(Nickl, et al. 2024), followed by transferring manipulated embryos

into pseudopregnant females for subsequent development. Overall, it is important to

generate a functional floxed Mirc13 allele prior to using Cre recombinase for either tissue-

specific or global deletion of the miR-141/200c cluster in order to avoid interference from

the lacZ/neo elements, including unanticipated knockout of endogenous GM15884 and

altering the expression of neighboring genes(McBurney, et al. 2002; Branda and Dymecki

2004). Our neighboring gene expression analyses further confirm the influence of

presence of β-actin-neo cassette on expression of many neighboring genes such as

ptpn6, phb2 and *atn1* in the Mirc13$^{tm1/tm1}$ mice. The genes *ptpn6, phb2,* and *atn1* each

play critical roles in brain function. *Ptpn6* regulates microglial signaling and helps restrain

neuroinflammation; its loss can drive excessive immune activation and neuronal

injury(He, et al. 2023). *Phb2* maintains mitochondrial integrity and neuronal survival, so

deletion can impair energy metabolism and cognitive functions(Xu, et al. 2014). *Atn1* acts

as a transcriptional co-regulator in neuronal development and disruption can lead to

abnormal differentiation, connectivity, or neurodegeneration(Whitton, et al. 1993). Thus,

downregulation of these genes as shown in Mirc13$^{tm1/tm1}$ mice could have significant

consequences for neuronal health and brain function. There is no change in expression

of these genes in Mirc13$^{fl/fl}$ mice, suggesting that it is essential to remove the lacZ/β-actin-

Neo cassettes to obtain fully functional Mirc13 cKO mice for meaningful and precise

analyses of their functional roles.

Despite the clear breeding plan recommended by Park et al(Park, et al. 2012),

several subsequent studies did not adopt the necessary breeding plan. They treated the

Mirc13^{tm1} mice, which contain lacZ/β-actin-neo cassettes, as Mirc13 cKO animals and

bred them directly with Cre deleters (Guan, et al. 2018; Hoefert, et al. 2018; Ji, et al. 2018;

Liu, et al. 2018; Wu, et al. 2019; Xu, et al. 2025). However, Mirc13^{tm1} mice contain three

colinear loxP sequences in the same allele (Fig 1A). It is well established that in the

presence of Cre, this would lead to different combinations of paired loxP sequences and

result in different excisions of loxP flanked sequences. As shown in Fig 6, this could

generate three different genotypes dependent on the different pairings of loxP sequences:

A). excision of the β-actin-Neo cassette alone; B) excision of Mirc13 alone; and C)

excision of β-actin-Neo cassette and Mirc13 together. Nevertheless, in all of these cases,

both the endogenous GM15884 and Mirc13 will not express and it is still not possible to

decipher the precise functional role of miR-141/200c. Other labs have bred Mirc13^{tm1}

pups with β-actin-Flpe mice prior to breeding them with Cre deleters. However, FLPe is

not codon-optimized for expression in mammals and the level of FLP protein expression

is compromised and excision of the b-actin-neo cassette may not take place in one-cell

embryo leading to mosaic animals(Tran, et al. 2017; Mostofa, et al. 2022; Tran, et al.

2024). We used Rosa26-FLPo mice, which would overcome the issue of low levels of the

FLP gene product(Raymond and Soriano 2007; Wu, et al. 2009; Kranz, et al. 2010). Our

breeding plan, as shown in Fig 1, is to remove the lacZ/neo cassettes using ROSA26-

FLPo mice, generating a functional Mirc13 floxed allele prior to using Cre mice, such as

the germline Hprt$^{CAG-Cre}$ mice, to achieve global deletion of the Mirc13 allele. These steps

ensure that any observed phenotype will be attributed directly to loss of the miR-141/200c

cluster, without any confounding contributions from transcriptional silencing or residual

neomycin cassette effects.

This breeding plan is not merely technical but critical for precise evaluation of the

miR-141/200c functional roles. Studies of the miR-200 family—including their roles in late

postnatal neurogenesis(Beclin, et al. 2016), stem cell differentiation(Zhang, et al. 2015),

and cancer biology(Chen, et al. 2013; Jin, et al. 2017; Tseng, et al. 2017) rely strongly on

precise gene dosage and context. Using animals with imprecise genotypes will only

hamper validities of the data leading to spurious conclusion and discrepancy among

different reports. By adhering to the breeding plan according to the design of the original

targeting vector, our work provides a framework to generate unambiguous global and

functional conditional knockouts of the miR-141/200c cluster.

Fig. 6 Breeding Mirc13^{tm1} mice containing three colinear loxP sites with Cre mice could lead to three different genotypes. **A** Excision of the β-actin-Neo cassette; **B** excision of β-actin-Neo cassette together with Mirc13 sequence; and **C** deletion of Mirc13 sequence. In all three different genotypes, both GM15884 and Mirc13 are not expressed due to the presence of the lacZ polyA sequence.

ACKOWLEDGEMENT

Author contributions

RV conceptualized the project. SKY, RS, RV and SPY wrote the manuscript. MKC, RS, SKY and KL performed the experiments. RV and SPY supervised the work. RV secured the funding for the project. All the authors approved the work.

Ethics declarations

Ethical Approval

All experiments were approved and conducted in compliance with the protocols approved (Number: AP-201043-0926) by the Institutional Animal Care and Use Committee (IACUC) at UConn Health. Animal data was reported following the ARRIVE Guidelines.

Declaration of Conflicting Interests

The authors declared no conflicts of interest.

Funding

This work was supported by NIH, United States (1R21NS114981 and R01 NS140176) and REP Convergence (OVPR UConn, United States) grant to R.V.

Data Availability

The data supporting the findings of this study will be available on request from the corresponding author.

REFERENCES

Beclin C, Follert P, Stappers E, Barral S, Coré N, Chevigny A, Magnone V, Lebrigand K, Bissels U, Huylebroeck D, et al. 2016. Corrigendum: miR-200 family controls late steps of postnatal forebrain neurogenesis via Zeb2 inhibition. Sci Rep 6:39368.

Bernabeu C, Lopez-Novoa JM, Quintanilla M. 2009. The emerging role of TGF-beta superfamily coreceptors in cancer. Biochim Biophys Acta 1792:954-973.

Boguski MS, Lowe TM, Tolstoshev CM. 1993. dbEST--database for "expressed sequence tags". Nat Genet 4:332-333.

Branda CS, Dymecki SM. 2004. Talking about a revolution: The impact of site-specific recombinases on genetic analyses in mice. Dev Cell 6:7-28.

Chen D, Zhang Y, Wang J, Chen J, Yang C, Cai K, Wang X, Shi F, Dou J. 2013. MicroRNA-200c overexpression inhibits tumorigenicity and metastasis of CD117+CD44+ ovarian cancer stem cells by regulating epithelial-mesenchymal transition. J Ovarian Res 6:50.

Choi PS, Zakhary L, Choi WY, Caron S, Alvarez-Saavedra E, Miska EA, McManus M, Harfe B, Giraldez AJ, Horvitz HR, et al. 2008. Members of the miRNA-200 family regulate olfactory neurogenesis. Neuron 57:41-55.

Clausen BE, Burkhardt C, Reith W, Renkawitz R, Förster I. 1999. Conditional gene targeting in macrophages and granulocytes using LysMcre mice. Transgenic Res 8:265-277.

Eppig JT, Bult CJ, Kadin JA, Richardson JE, Blake JA, Anagnostopoulos A, Baldarelli RM, Baya M, Beal JS, Bello SM, et al. 2005. The Mouse Genome Database (MGD): from genes to mice--a community resource for mouse biology. Nucleic Acids Res 33:D471-475.

Guan T, Dominguez CX, Amezquita RA, Laidlaw BJ, Cheng J, Henao-Mejia J, Williams A, Flavell RA, Lu J, Kaech SM. 2018. ZEB1, ZEB2, and the miR-200 family form a counterregulatory network to regulate CD8(+) T cell fates. J Exp Med 215:1153-1168.

Hashimoto M, Takemoto T. 2015. Electroporation enables the efficient mRNA delivery into the mouse zygotes and facilitates CRISPR/Cas9-based genome editing. Sci Rep 5:11315.

He Y, Nan D, Wang H. 2023. Role of Non-Receptor-Type Tyrosine Phosphatases in Brain-Related Diseases. Mol Neurobiol 60:6530-6541.

Hoefert JE, Bjerke GA, Wang D, Yi R. 2018. The microRNA-200 family coordinately regulates cell adhesion and proliferation in hair morphogenesis. J Cell Biol 217:2185-2204.

Humphries B, Yang C. 2015. The microRNA-200 family: small molecules with novel roles in cancer development, progression and therapy. Oncotarget 6:6472-6498.

Isgrò MA, Bottoni P, Scatena R. 2015. Neuron-Specific Enolase as a Biomarker: Biochemical and Clinical Aspects. Adv Exp Med Biol 867:125-143.

Janssen HL, Reesink HW, Lawitz EJ, Zeuzem S, Rodriguez-Torres M, Patel K, van der Meer AJ, Patick AK, Chen A, Zhou Y, et al. 2013. Treatment of HCV infection by targeting microRNA. N Engl J Med 368:1685-1694.

Ji ML, Jiang H, Zhang XJ, Shi PL, Li C, Wu H, Wu XT, Wang YT, Wang C, Lu J. 2018. Preclinical development of a microRNA-based therapy for intervertebral disc degeneration. Nat Commun 9:5051.

Jin T, Suk Kim H, Ki Choi S, Hye Hwang E, Woo J, Suk Ryu H, Kim K, Moon A, Kyung Moon W. 2017. microRNA-200c/141 upregulates SerpinB2 to promote breast cancer cell metastasis and reduce patient survival. Oncotarget 8:32769-32782.

Kranz A, Fu J, Duerschke K, Weidlich S, Naumann R, Stewart AF, Anastassiadis K. 2010. An improved Flp deleter mouse in C57Bl/6 based on Flpo recombinase. Genesis 48:512-520.

Lee RC, Feinbaum RL, Ambros V. 1993. The C. elegans heterochronic gene lin-4 encodes small RNAs with antisense complementarity to lin-14. Cell 75:843-854.

Liu B, Du R, Zhou L, Xu J, Chen S, Chen J, Yang X, Liu DX, Shao ZM, Zhang L, et al. 2018. miR-200c/141 Regulates Breast Cancer Stem Cell Heterogeneity via Targeting HIPK1/β-Catenin Axis. Theranostics 8:5801-5813.

McBurney MW, Mai T, Yang X, Jardine K. 2002. Evidence for repeat-induced gene silencing in cultured Mammalian cells: inactivation of tandem repeats of transfected genes. Exp Cell Res 274:1-8.

Mostofa MG, Tran M, Gilling S, Lee G, Fraher O, Jin L, Kang H, Park YK, Lee JY, Wang L, et al. 2022. MicroRNA-200c coordinates HNF1 homeobox B and apolipoprotein O functions to modulate lipid homeostasis in alcoholic fatty liver disease. J Biol Chem 298:101966.

Nickl P, Jenickova I, Elias J, Kasparek P, Barinka C, Kopkanova J, Sedlacek R. 2024. Multistep allelic conversion in mouse pre-implantation embryos by AAV vectors. Sci Rep 14:20160.

Olson EN, Arnold HH, Rigby PW, Wold BJ. 1996. Know your neighbors: three phenotypes in null mutants of the myogenic bHLH gene MRF4. Cell 85:1-4.

Pan Y, Zhang L, Liu Q, Li Y, Guo H, Peng Y, Peng H, Tang B, Hu Z, Zhao J, et al. 2016. Insertion of a knockout-first cassette in Ampd1 gene leads to neonatal death by disruption of neighboring genes expression. Sci Rep 6:35970.

Park CY, Jeker LT, Carver-Moore K, Oh A, Liu HJ, Cameron R, Richards H, Li Z, Adler D, Yoshinaga Y, et al. 2012. A resource for the conditional ablation of microRNAs in the mouse. Cell Rep 1:385-391.

Pasquinelli AE, Reinhart BJ, Slack F, Martindale MQ, Kuroda MI, Maller B, Hayward DC, Ball EE, Degnan B, Müller P, et al. 2000. Conservation of the sequence and temporal expression of let-7 heterochronic regulatory RNA. Nature 408:86-89.

Pham CT, MacIvor DM, Hug BA, Heusel JW, Ley TJ. 1996. Long-range disruption of gene expression by a selectable marker cassette. Proc Natl Acad Sci U S A 93:13090-13095.

Raja R, Dumontier E, Phen A, Cloutier JF. 2024. Insertion of a neomycin selection cassette in the Amigo1 locus alters gene expression in the olfactory epithelium leading to region-specific defects in olfactory receptor neuron development. Genesis 62:e23594.

Raymond CS, Soriano P. 2007. High-efficiency FLP and PhiC31 site-specific recombination in mammalian cells. PLoS One 2:e162.

Rishik S, Hirsch P, Grandke F, Fehlmann T, Keller A. 2025. miRNATissueAtlas 2025: an update to the uniformly processed and annotated human and mouse non-coding RNA tissue atlas. Nucleic Acids Res 53:D129-d137.

Skarnes WC, Auerbach BA, Joyner AL. 1992. A gene trap approach in mouse embryonic stem cells: the lacZ reported is activated by splicing, reflects endogenous gene expression, and is mutagenic in mice. Genes Dev 6:903-918.

Skarnes WC, Rosen B, West AP, Koutsourakis M, Bushell W, Iyer V, Mujica AO, Thomas M, Harrow J, Cox T, et al. 2011. A conditional knockout resource for the genome-wide study of mouse gene function. Nature 474:337-342.

Small EM, Frost RJ, Olson EN. 2010. MicroRNAs add a new dimension to cardiovascular disease. Circulation 121:1022-1032.

Tang SH, Silva FJ, Tsark WM, Mann JR. 2002. A Cre/loxP-deleter transgenic line in mouse strain 129S1/SvlmJ. Genesis 32:199-202.

Testa G, Schaft J, van der Hoeven F, Glaser S, Anastassiadis K, Zhang Y, Hermann T, Stremmel W, Stewart AF. 2004. A reliable lacZ expression reporter cassette for multipurpose, knockout-first alleles. Genesis 38:151-158.

Tran M, Gilling S, Wu J, Wang L, Shin DJ. 2024. miR-141/200c contributes to ethanol-mediated hepatic glycogen metabolism. Mol Metab 84:101942.

Tran M, Lee SM, Shin DJ, Wang L. 2017. Loss of miR-141/200c ameliorates hepatic steatosis and inflammation by reprogramming multiple signaling pathways in NASH. JCI Insight 2.

Tseng JH, Bisogna M, Hoang LN, Olvera N, Rodriguez-Aguayo C, Lopez-Berestein G, Sood AK, Levine DA, Jelinic P. 2017. miR-200c-driven Mesenchymal-To-Epithelial Transition is a Therapeutic Target in Uterine Carcinosarcomas. Sci Rep 7:3614.

van Rooij E, Kauppinen S. 2014. Development of microRNA therapeutics is coming of
age. EMBO Mol Med 6:851-864.

Verma R, Ritzel RM, Harris NM, Lee J, Kim T, Pandi G, Vemuganti R, McCullough LD.
2018. Inhibition of miR-141-3p Ameliorates the Negative Effects of Poststroke Social
Isolation in Aged Mice. Stroke 49:1701-1707.

Whitton C, Palmer E, Alkuraya F. 1993. ATN1-Related Neurodevelopmental Disorder. In:
Adam MP, Bick S, Mirzaa GM, Pagon RA, Wallace SE, Amemiya A, editors.
GeneReviews(®). Seattle (WA): University of Washington, Seattle

Copyright © 1993-2026, University of Washington, Seattle. GeneReviews is a registered
trademark of the University of Washington, Seattle. All rights reserved.

Wightman B, Ha I, Ruvkun G. 1993. Posttranscriptional regulation of the heterochronic
gene lin-14 by lin-4 mediates temporal pattern formation in C. elegans. Cell 75:855-862.

Wu MJ, Chen YS, Kim MR, Chang CC, Gampala S, Zhang Y, Wang Y, Chang CY, Yang
JY, Chang CJ. 2019. Epithelial-Mesenchymal Transition Directs Stem Cell Polarity via
Regulation of Mitofusin. Cell Metab 29:993-1002.e1006.

Wu Q, Ye X, Xiong Y, Zhu H, Miao J, Zhang W, Wan J. 2016. The Protective Role of
microRNA-200c in Alzheimer's Disease Pathologies Is Induced by Beta Amyloid-
Triggered Endoplasmic Reticulum Stress. Front Mol Neurosci 9:140.

Wu Y, Wang C, Sun H, LeRoith D, Yakar S. 2009. High-efficient FLPo deleter mice in
C57BL/6J background. PLoS One 4:e8054.

Xu L, Griffiths B, Sun X, Greer M, Chang H, McMillin E, Manis E, Russo I, Dean CAJ,
Stary CM. 2025. Conditional neuronal deletion of microRNA-141/200c cluster, but not
microRNA-181a/b-1 cluster, is protective against experimental stroke in male mice.
Physiol Rep 13:e70505.

Xu T, Fan X, Tan Y, Yue Y, Chen W, Gu X. 2014. Expression of PHB2 in rat brain cortex
following traumatic brain injury. Int J Mol Sci 15:3299-3318.

Yadav SK, Dhuri K, Gamiotea-Turro D, Cormier MK, Patel V, Yadawa AK, Pathuri M,
Bahal R, Verma R. 2024. Exploring the therapeutic potential of sγPNA-141:
Pharmacodynamics and mechanistic insights during ischemic stroke recovery. Mol Ther
Nucleic Acids 35:102355.

Zhang J, Gao Y, Yu M, Wu H, Ai Z, Wu Y, Liu H, Du J, Guo Z, Zhang Y. 2015. Retinoic
Acid Induces Embryonic Stem Cell Differentiation by Altering Both Encoding RNA and
microRNA Expression. PLoS One 10:e0132566.