1   **Using Tools as Cues for Motor Adaptation in Virtual Reality**

6   Andrew Michael King[1-2], Jacob J. Boulrice[1-2], Shanaathanan Modchalingam[1,3], Laura

7   Mikula[1], Bernard Marius 't Hart[1], Denise Y P Henriques[1-3]

9   1-Centre for Vision Research, York University

10   2-Psychology, York University

11   3-Kinesiology and Health Science, York University

12   Toronto, ON, M6J1P3

14   Corresponding author: Denise Y P Henriques, deniseh@yorku.ca, Kinesiology and Health

15   Science, York University, 4700 Keele St, Toronto, ON, Canada, M3J1P3. 416-736-2100

Abstract:

Humans are adept at using multiple tools, often switching between them even when each
involves distinct and potentially conflicting motor demands. This study investigated how
different features of a tool influence the formation of distinct motor memories during dual
adaptation to opposing visuomotor perturbations. Using an immersive virtual reality (VR) setup,
participants performed an aiming task in which they launched a ball toward a target using virtual
tools, each consistently associated with a different visuomotor rotation. We manipulated tool
features across three groups: one in which tools differed only in colour (Colour Control), one in
which they differed in shape but involved similar movement patterns (Motor Congruent), and
one in which they differed both in shape and in how they were operated—for example, a paddle-
like forward swing to propel the ball versus a draw-and-release motion similar to a slingshot
(Motor Incongruent). A fourth control group that adapted to a single perturbation with a single
tool at a time was also included. Only the Motor Incongruent group demonstrated robust dual
adaptation and clear aftereffects, comparable to those observed during single-tool learning.
These results suggest that distinct modes of tool operation play a critical role in supporting the
formation and retention of separate internal models during sensorimotor adaptation.

New & Noteworthy

Humans often switch between tools with conflicting motor demands. Using immersive
virtual reality, we tested whether visual features or operational differences support dual
adaptation to opposing visuomotor perturbations. Only when tools differed in both shape
and mode of operation did participants show robust adaptation and aftereffects,
comparable to single-tool learning. These findings highlight the critical role of
movement-relevant cues in forming distinct motor memories for flexible tool use.

Introduction

Humans excel at using a variety of tools, often switching fluidly between them—even
when they require opposing motor actions. This flexibility suggests that the motor
system can form and retrieve distinct internal models to govern tool-specific actions
without interference. However, the conditions under which such separate motor
memories can be acquired and maintained remain unclear.

Sensorimotor adaptation studies have shown that learning a novel mapping between
motor commands and their expected sensory outcomes—such as adapting to a rotated
visual cursor—relies on sensory prediction errors that update internal models of
movement [1–4]. However, when individuals are required to adapt to two opposing
perturbations within the same session, performance often suffers due to interference. In
this context, interference refers to the tendency for learning one perturbation (e.g., a
clockwise rotation) to partially overwrite, disrupt, or destabilize the motor memory
formed for the opposing perturbation (e.g., a counterclockwise rotation). As a result, the
motor system fails to maintain two distinct internal models, leading to reduced retention,
slower learning, or a blending of the two learned states into a compromised
intermediate solution. This limitation suggests that, without additional contextual cues to
signal when each perturbation should be applied, the nervous system treats the
conflicting experiences as arising from a single task environment and updates a single
control policy. By contrast, when clear contextual cues are provided—such as distinct
visual tools, postures, or task rules—the motor system can assign each perturbation to
a separate internal model, thereby reducing interference and enabling robust dual
adaptation [5–14].

Prior work has demonstrated that intrinsic contextual cues—those tied to the
motor execution itself, such as movement direction, posture, or lead-in movements—
support dual adaptation by recruiting distinct control policies [5,6,11–21] In contrast, extrinsic
cues—such as color or tool shape—often fail to enable interference-free learning unless
made task-relevant or tightly associated with specific sensorimotor demands [15,17,22,23].

Tools present a particularly interesting case, as they combine both extrinsic
visual features and intrinsic motor properties. Some studies have shown that tools can
support dual adaptation when they differ in their control dynamics or when paired with
distinct contact goals [9,10,12,24]. However, the relative importance of visual versus motoric
features in supporting the formation of separate internal models remains unresolved.

In this study, we investigated whether visual-identity and operational features of
tools can serve as effective contextual cues for dual adaptation in a visuomotor rotation
task. Using immersive virtual reality (VR), participants adapted to two opposing rotations
that altered the direction of ball motion, with each rotation consistently paired with a
distinct tool. Importantly, each tool launched the ball in a unique manner, requiring
participants to engage with it using different movement dynamics—for example, one
tool involved a smooth paddle-like swing to propel the ball, whereas another required a
draw-and-release action similar to a slingshot. Across experimental groups, we
systematically manipulated whether these tool pairs differed only in their visual shape
(providing purely extrinsic cues), in both shape and the specific movement required to
operate them (combining extrinsic and intrinsic cues), or solely in color (a minimal
extrinsic cue without functional differences). By doing so, our goal was to determine
whether movement-specific operational properties are necessary to support robust dual
adaptation, or whether visual features alone can serve as sufficient contextual signals in
the absence of distinct motor requirements.

Methods

Participants

A total of 144 right-handed undergraduate students (98 female, mean age = 21.98 ±
4.62 years) participated in the study. Participants belonged to one of four groups: Motor
Incongruent (n = 40), Motor Congruent (n = 40), Colour Control (n = 44), and Single
Adaptation (n = 20). All had normal or corrected-to-normal vision and provided informed
consent. Participants received course credit for participation. The study protocol was
approved by York University's Human Participants Review Sub-committee.

Apparatus and Virtual Environment

Participants were seated at a table and wore an Oculus Rift CV1 virtual reality headset
(resolution: 1080 × 1200 pixels per eye; refresh rate: 90 Hz). Head and hand
movements were tracked using three external Oculus sensors, which sampled at 60 Hz
with sub-millimetre precision (<1 mm). Participants used a hand-held Oculus Touch
controller to interact with objects in a custom-built virtual environment developed in
Unity. The virtual scene was designed to match the physical dimensions of the table
(3.5 m wide × 2 m deep) to provide consistent spatial alignment between real and virtual
space (as with Modchalingam et al., 2024, 2025). Trial presentation and data collection
were managed using the Unity Experimenter Framework[26], which supported
synchronized stimulus control and response logging throughout the experiment.

Task and Stimuli

Participants performed a tool-mediated ball-launching task. On each trial, they used one of three virtual tools—a paddle, slingshot, or curling rod (Fig 1C)—to launch a virtual ball (5 cm diameter) toward a target (5 cm) positioned 85 cm away in one of four directions (75°, 85°, 95°, 105° polar angle, Fig 1A). Tools differed in shape (Fig 1C-D) and pre-launch movement (Fig 1D, 4B): the paddle and curling rod involved a forward strike, while the slingshot required a backward pull and release. Each Group only trained with two of the three tools, were used to manipulate motor congruency between tools (as per Fig 1D).

Ball trajectories after launch was governed by Unity's physics engine, PhysX, with a maximum launch-velocity cap of 22 cm/s. Flanking walls on the virtual table ensured that the ball was released from the same restricted area for all tools, and the tool positions were constrained so they could not extend beyond this launch zone At the end of each trial, participants received visual feedback showing the ball's trajectory and a performance score (0-10 points), based on the minimum distance between the ball and the target (e.g., Fig 1B). A green timer bar (Fig 1A-B) prompted participants to launch the ball within 1 s (3 s during practice). Late responses or shots deviating more than 20 cm from the target received a score of 0.

Experimental Design

The experiment employed a between-subjects design with four participant groups, see Fig 1D. Three groups completed a dual adaptation (DA) condition, in which each participant used two distinct tools consistently associated with opposing visuomotor perturbations (clockwise [CW] or counterclockwise [CCW]). The Motor Congruent group trained with the paddle and curling rod, which required similar forward launching movements. The Motor Incongruent group trained with the paddle and slingshot, which required contrasting movement patterns (forward vs. backward), introducing greater sensorimotor dissimilarity between tools. The Colour Control group used two visually distinct paddles (red and blue) that differed only in colour and not in movement dynamics, allowing us to isolate the role of visual cues alone. A fourth group, the Single Adaptation (SA) group, served as a control group and completed two separate adaptation blocks with a single tool and perturbation pairing per block, with no interleaving of perturbation.

All participants completed four experimental phases: a Practice phase, a Baseline phase, a Training phase, and a Washout phase, as illustrated in Fig 2A. The Practice phase familiarized participants with the mechanics of the ball-launching task using unperturbed feedback (24 trials per tool). The Baseline phase consisted of 80 trials with veridical feedback (40 per tool) to assess initial performance (illustrated in Fig 2B, top

row and Fig 2C, left panel). In the Training phase, a 30° rotation (CW or CCW) was
imposed on the ball trajectory (illustrated in Fig 2B, bottom row and Fig 2C, right panel).
In the dual adaptation groups (Fig 2A, top row), each rotation was consistently paired
with one of the two tools, and tools alternated every 8 trials in mini-blocks throughout
the phase (160 trials total; 80 per tool). Each block of these 8 trials is represented by a
single rectangle in Fig 2A, and with different shades of same colour associated with
different tools. In contrast, participants in the Single Adaptation group (Fig 2A, bottom
row) completed 80 consecutive trials with one tool and rotation before switching to the
other tool and opposing perturbation for an additional 80 trials. To compensate for the
visuomotor rotation introduced during the Training phase, participants had to adjust their
hand movements in the direction opposite to the imposed rotation in order to accurately
aim the ball at the target. The order of tool and perturbation was counterbalanced
across participants. During the final Washout phase, participants completed 48 aligned
trials (24 per tool) —with veridical feedback identical to Baseline— to assess
aftereffects and de-adaptation. Background textures differed between aligned and
rotated trials (Fig 2B), but participants were not informed of these changes, nor were
they explicitly told about the visuomotor rotations. Short rest breaks were given between
phases, and participants kept their headsets on throughout the session.

Procedure

The participant's task was to use the provided tool to shoot a virtual ball toward a visible
target. To initiate each trial, participants moved the VR hand-held controller to the tool's
location. As the controller and hand had no visible representation aside from the tool
itself, the controller vibrated when it came within 1 cm of the tool to provide haptic
confirmation of proximity. Upon reaching the tool, participants pressed and held the
trigger button to "grab" it, which initiated a brief countdown signaled by a green
horizontal bar displayed on the screen (Fig 1A–B). This countdown lasted one second
during experimental trials and three seconds during practice trials, after which
participants were prompted to quickly and accurately shoot the ball toward the target.
Shooting the ball involved intuitive, tool-specific movements; for example, the paddle
and rod required a forward push, while the slingshot required a backward pulling motion.

Each trial ended when the ball either hit the target, slowed to a speed below 1 cm/s, or
reached a radial distance of 120 cm from the launch point. At that point, a white line
appeared to indicate the ball's path, providing participants with visual feedback on their
accuracy (Fig 1B). To motivate participants, a score between 0 and 10 was awarded on
each trial based on their minimum error—that is, the smallest distance between the ball
and the target during the trajectory. An additional 5 bonus points were awarded if the
ball hit the center of the target. The cumulative score and number of remaining trials
were displayed throughout the session on an in-VR scoreboard (Fig 1A–B). Trials in

which the ball missed the target by more than 20 cm or was launched after the
countdown expired received a score of 0.

Data Analysis

This study investigated how tool identity (extrinsic cues) and tool operation (intrinsic
cues) contribute to the concurrent learning of tool-pair associations, each linked to
different visuomotor perturbations. By manipulating the congruency between the
movements required to operate each tool and its visual identity, we evaluated whether
these characteristics support the formation of separate motor memories under
conditions of dual adaptation.

The primary dependent variable was angular error, defined as the difference between
the direction of the ball's trajectory and the target direction. To allow comparisons
across the two opposing perturbation directions, angular errors were standardized by
reversing the sign for one of the rotations. In this convention, a positive error of 30
degrees indicates no compensation for the rotation, and values closer to zero reflect
greater compensation. Complete compensation was defined as errors within ±3.3°,
corresponding to the diameter of the ball. Ball trajectory distance was also recorded,
and trials were excluded if the ball failed to travel at least 17 cm (20% of the required
distance) or if angular errors exceeded ±60°. These criteria resulted in the removal of 96
trials (0.07% of all trials) due to distance and 586 trials (0.41%) due to excessive
angular error.

To control for baseline tool or target biases, we computed the median angular error for
each tool–target pair during the Baseline phase and subtracted these values from
Training phase trials. For all statistical analyses, we focused on performance with the
red paddle, which was the only tool common across all four experimental groups. This
ensured comparability across the different cue conditions. While the second tool in each
pairing was not included in the main analyses, performance for both tools is presented
in Fig 3 for descriptive comparison. As expected, there were no substantial performance
differences between tools during the Baseline phase or in the first block of Training trials
prior to introducing the second tool and perturbation. Across groups, paddle
performance (shown in purple in Fig 3) fell within the 95% confidence intervals of the
other tool's errors, suggesting no tool-specific performance biases at the start of
adaptation.

We observed a small but significant effect of tool order during training. Specifically,
participants who encountered the paddle as the second tool showed larger mean errors
(34.23°) on the first paddle trial than those who used the paddle first (27.98°),
suggesting a 7° interference effect from prior exposure to the opposing perturbation.
This tool order effect was significant ($F_{(1,134)} = 7.478$, $p < 0.01$), but it did not interact
with group ($F_{(3,134)} = 0.67$, $p = 0.57$). Because tool order and associated rotation

direction were counterbalanced across participants, we collapsed across tool order in
our primary analyses to simplify comparisons. While this approach may reduce
sensitivity to detect between-group differences in learning rates due to increased
variance, it provides a more balanced assessment of overall group performance across
both tools, and our relatively large sample sizes help mitigate this reduced sensitivity.

To compare adaptation across the four groups, we conducted analyses at both the trial
level and the block level using angular errors from the paddle. At the trial level, we
compared the first and final trials of the Training phase to capture the full extent of
individual trial-by-trial adaptation. At the block level, we averaged angular errors across
all eight trials within each block and compared the first and final blocks of training. This
dual-level approach allowed us to assess not only the immediate change in
performance within a single trial but also broader patterns of learning across repeated
exposures to the same tool–rotation pairing.

Statistically, we used a 4 (group: Congruent, Incongruent, Colour, Single) × 2 (time:
early vs. late) linear mixed-effects model (LMER) separately for (1) the first and last
trials of training and (2) the mean errors from the first and last blocks. This enabled us
to examine overall group differences in the magnitude of adaptation, as well as any
interactions indicating group-specific learning profiles. Where significant group × time
interactions were found, we performed one-way ANOVAs on the first trial and first block
to confirm that all groups started with comparable performance levels. As expected,
there were no significant group differences in the first trial ($F(3,138) = 0.43$, $p = 0.72$) or
first block ($F(3,140) = 1.53$, $p = 0.21$), supporting the assumption that all groups began
training with similar baseline-corrected performance. These results were consistent
even when the analysis was restricted to the three dual adaptation groups (all $p > 0.05$).
We then compared performance in the final training block across groups using a one-
way ANOVA to test whether any differences emerged over the course of training.

When significant effects were detected, Tukey post hoc comparisons were used to
identify specific group differences. This stepwise approach ensured that any observed
group differences in final performance could be interpreted as the result of learning
under different cue conditions, rather than pre-existing performance differences or
random variation in early trials.

To examine within- and between-block adaptation, we analyzed performance on the first
trial (T1) and final trial (T8) of each 8-trial block. This allowed us to quantify both short-
term learning and retention after exposure to the opposing perturbation. Although full T1
and T8 data are plotted across all blocks in Fig 4A–C, we focused our statistical
comparisons on the first and final training blocks (Fig 3A). We used two 3 (group) × 2
(block) LMERs—one for T1 and one for T8—to assess learning and retention across the
Training phase. Any significant group × block interactions were followed by one-way

ANOVAs across groups on the first and final blocks to isolate the source of the
interaction. Post hoc comparisons were conducted as needed.

To assess aftereffects and de-adaptation, we compared angular errors for the paddle
between the final block of Baseline and the first block of Washout across all four groups
using a 4 × 2 LMER. Significant interactions were followed by paired t-tests within each
group to assess whether errors increased in the expected direction following removal of
the perturbation. To control for multiple comparisons, a Bonferroni correction was
applied to the four planned t-tests, resulting in an adjusted significance threshold of p =
0.0125.

Data and scripts can be found https://osf.io/khms8

Github: https://github.com/maceagle12/ToolUseData

Results

To evaluate whether tool-related cues are sufficient for dual adaptation, we first
describe the overall adaptation patterns across all tool pairings before focusing on
performance with the common paddle tool. Each group used a distinct combination of
two tools, with each tool consistently paired with a unique visuomotor perturbation. Our
central aim was to determine whether adaptation to one tool would interfere with
learning the opposing perturbation associated with the second tool—or whether certain
contextual cues would allow participants to maintain separate motor memories and
facilitate dual adaptation. To enable direct comparison across conditions, we focus our
main statistical analyses on the red paddle, which was used in all four groups. By
analyzing how performance with this shared tool changed in the presence of different
paired tools, we could isolate the influence of contextual cue type—tool identity, control
dynamics, or color—on adaptation.

General Performance Patterns Across Tool Pairings

Figure 3 shows angular launch errors across the final three Baseline blocks, all ten
Training blocks, and the first three Washout blocks for each group, plotted separately
for each tool. To compare across opposing perturbation directions, errors from one tool
in each pair were sign-reversed so that zero always reflects full compensation. Although
tools were alternated in blocks of 8 trials, we plot their trial sequences as overlapping to
visualize adaptation across the experiment.

As expected, all groups performed comparably at Baseline (all $p > 0.18$). Upon
introduction of the visuomotor rotation, each group initially exhibited large deviations
consistent with the imposed perturbation (first trial $F_{(3,138)} = 0.43$, $p = 0.72$; first block
$F_{(3,140)} = 1.53$, $p = 0.21$) (First trial means: Motor Incongruent (m= 29.7, SE=
1.92),Motor Congruent (m= 31.2, SE= 1.92), Colour control (m= 32.7, SE = 1.85),
Single adaptation (m = 30.8 , SE = 2.72); First Block means: Motor Incongruent (m=22.7,
SE = 1.11), Motor Congruent (m = 26.0, SE = 1.11), Colour Control (m = 24.9, SE =
1.05), Single Adaptation (m = 23.5, SE =1.56). All groups reduced their errors over
successive trials, indicating adaptation. However, a distinctive zig-zag pattern emerged
across Training for DA groups, reflecting interference: within-block improvements did
not fully persist into the next block with the same tool, at least for the Motor Congruent
and Control Control group, suggesting the intervening tool disrupted retention.

The Motor Incongruent group also showed a zig-zag pattern, but with clearer evidence
of retention across successive blocks using the same tool, suggesting reduced
interference compared to the other dual-adaptation groups. First-trial errors decreased
across successive paddle blocks, suggesting greater retention across tool switches.

This pattern was less apparent in the Colour and Motor Congruent groups, where
within-block learning was not maintained between blocks. These differences are
analyzed in more detail below.

Dual Adaptation: Impact of Tool Pairing

To quantify group differences in adaptation, we examined paddle performance during
Training. There was a significant main effect of trial, with angular error decreasing from
the first to the final trial ($F_{(1,138)} = 79.31$, $p < 0.001$; Fig. 4A), and a significant main
effect of block, with angular error decreasing from the first to the final block ($F_{(1,147)} =$
61.02, $p < 0.001$; Fig. 4B). However, the magnitude of this change differed across
groups, as indicated by significant trial × group and block × group interactions (trial ×
group: $F_{(3,139)} = 2.84$, $p < 0.05$; block × group: $F_{(3,148)} = 13.33$, $p < 0.001$). Post hoc
tests revealed that only the Motor Incongruent group (red) did not differ significantly
from the Single Adaptation group (green) by the end of Training ($p = 0.928$) as
illustrated in green in Fig 4. In contrast, both the Colour (blue) and Congruent (orange)
groups showed significantly less improvement (both $p < 0.001$), with no difference
between them ($p = 0.999$). These results indicate that only Motor Incongruent tool
pairings enabled sufficient contextual separation to support dual adaptation.

Retention Across Alternating Blocks

To investigate how adaptation was retained across tool switches, we examined
performance across the ten alternating training blocks. Specifically, we plotted launch
errors for the first trial (T1) of each block (Fig 5A) and the last trial (T8) of each block
(Fig5C). This allowed us to assess, respectively, how much adaptation was retained at
the beginning of each block and how much cumulative learning was achieved within
each block over time. Visual inspection of Fig 5A shows a progressive reduction in initial
error across blocks for the Motor Incongruent group (red), indicating that participants
retained some adaptation from previous encounters with the same tool—even after
using the other, differently perturbed tool in the intervening block. In contrast, the Motor
Congruent (orange) and Colour Control (blue) groups show relatively flat trajectories
across blocks, with little reduction in initial error, suggesting weaker retention across
tool switches.

In Fig 5C, which depicts the final trial of each block, a similar—but less pronounced—
trend is observed for the Motor Incongruent group: errors decrease modestly over the
first few blocks. In contrast, the other two groups show no change across all 10 blocks.
However, there is greater variability in the end-of-block data compared to the first trials,
and the downward trend is less pronounced.

To quantify these retention effects, we focused on comparing only the first and last
blocks of training (Blocks 1 and 10) for both T1 and T8 trials in the Dual Adaptation

conditions—where the greatest change in performance would be expected due to
repeated exposure (Figs 5B and 5D). Comparing launch error for T1 in the initial and
final blocks (Fig 5B) reveals a significant reduction in error from Block 1 to Block 10
$(F(1,120) = 7.12, p < 0.05)$. This confirms that some adaptation achieved within each
block is retained in subsequent blocks, for the same tool. However, this retention was
group-dependent, as revealed by a significant group × block interaction for T1 $(F(1,120)$
$= 3.45, p < 0.05)$. Because the groups did not differ in their T1 performance on the first
block (as previously reported), this interaction reflects differences in performance during
the final block. A follow-up ANOVA on the final block confirmed a main effect of group
on T1 $(F(2,119) = 6.87, p < 0.01)$. Tukey comparisons showed that the Motor
Incongruent group retained more learning across blocks than both the Colour Control (p
< 0.01) and Motor Congruent groups $(p < 0.01)$. In fact, only the Motor Incongruent
group shows a visible reduction in error from Block 1 to Block 10: the red data point in
Fig 5B falls below the confidence interval of their initial value.

When comparing T8 trials across blocks (Fig 5D), no significant change in error was
detected $(F(1,120) = 1.96, p = 0.16)$, and no group × block interaction was found
$(F(1,120) = 1.84, p = 0.16)$. This suggests that the cumulative adaptation achieved by
the end of the first block (before any interference from the second tool) was not
substantially improved after additional practice. This lack of statistical significance may
reflect increased inter-subject variability, rather than the absence of a true effect.
particularly in the Motor Incongruent group where the mean trend did suggest a
reduction in error.

Overall, these findings suggest that tool-pairings involving Motor Incongruent motor
actions, which provided both intrinsic and extrinsic cues, supported greater retention of
context-specific learning across alternating blocks. The Motor Incongruent group's
performance—particularly in the first trial of each block—shows evidence of dual
adaptation that accumulates across training despite frequent switching between tools
with opposing perturbations.

Aftereffects Following Adaptation

To evaluate persistent adaptation, we compared angular errors in the first Washout
block to the final Baseline block, as illustrated in Fig 6. A significant group × phase
interaction $(F(3,140) = 20.60, p < 0.001)$ indicated that aftereffects differed by condition.
Only the Motor Incongruent (red) and Single Adaptation (green) groups showed
significant deviations during Washout (both $p < 0.001$, Bonferroni corrected), consistent
with implicit motor memory formation. The Congruent and Colour groups showed no
such aftereffects (both $p > 0.0125$), reinforcing the conclusion that visual or identity cues
alone were insufficient to support robust durable dual adaptation.

Discussion

The present study demonstrates that distinct tool operations—not visual features
alone—enable successful dual adaptation to opposing visuomotor perturbations in
immersive VR. When tools differed in both appearance and how they were operated
(Motor Incongruent group), participants adapted as effectively as in the Single
Adaptation group, despite alternating between tools with opposing perturbations every
eight trials. In contrast, when tools differed only in color (Colour group) or had distinct
visual appearances but shared similar tool wielding patterns (Motor Congruent group),
participants exhibited limited retention across blocks and failed to show aftereffects—
suggesting substantial interference and a lack of distinct motor memory formation.
These findings build on prior work emphasizing the importance of intrinsic contextual
cues—those related to how a tool is operated—in enabling dual adaptation (e.g., Ayala
et al., 2015b; Heald et al., 2018, 2021; Howard et al., 2015; Proud et al., 2019). Our
results further support the conclusion that extrinsic cues are insufficient unless tightly
coupled with distinct control policies [5–8,11–14,20,21,29,30]. Previous studies testing extrinsic
cues typically relied on minimal differences—such as changes in color or shape—that
were not directly relevant to the task. In contrast, our study is one of the first to employ
rich, task-relevant extrinsic cues: the paddle and shuffle tools were visually distinct and
differed slightly in their method of operation. Despite this, our findings show that even
these salient visual distinctions, in the absence of clearly differentiated tool-use
dynamics, do not support robust dual adaptation.

Notably, adaptation in the Motor Incongruent group was characterized by progressive
improvement in the first trial of each block of the same tool, suggesting increasing
retention despite blocks of interleaved perturbations linked with the second tool. This
pattern was absent in the Colour and Congruent groups, where adaptation had to be re-
learned in each block. These results align with the notion that interference in dual
adaptation arises when the same control policy is reused across contexts [30], and that
differences in tool operation help the motor system retrieve and retain the appropriate
internal model.

Our aftereffects findings further underscore the importance of intrinsic cues related to
tool operation. In both the Motor Incongruent and Single Adaptation conditions,
participants exhibited significant reach deviations when the visuomotor perturbation was
removed for the common paddle—indicating the formation of implicit motor memories.
In contrast, no aftereffects were observed in the Colour Control or Motor Congruent
groups, where limited separation between internal models likely prevented the
consolidation of implicit adaptation.

Interestingly, the slingshot tool—despite supporting effective adaptation during
training—did not produce observable aftereffects in either the Dual Adaptation or Single

Adaptation conditions (see Fig 6). This discrepancy may reflect that the slingshot was

somewhat easier to learn, or alternatively, that participants relied on explicit strategies

due to its constrained and repetitive draw-and-release action. The uniformity of this

movement pattern may have encouraged the use of simple aiming heuristics or rule-

based strategies, thereby reducing reliance on an internal forward model [31,32]. This may

also account for the slightly greater compensation observed during training with the

slingshot, although the performance advantage over the paddle was modest—

approximately 5–10%. In contrast, the paddle, which permitted a wider range of

movement directions and magnitudes, likely required more flexible motor recalibration

and thus engaged implicit learning mechanisms that persisted even after the

perturbation was removed.

This study extends dual adaptation research by explicitly manipulating tool dynamics

within an immersive VR environment, where spatial cues and the surrounding visual

context were held constant. The use of VR provided a unique advantage by allowing us

to control for environmental variables while systematically testing both tool-specific

dynamics and richer, task-relevant extrinsic cues, such as distinct visual tool identities

(e.g., Modchalingam et al., 2025). Our findings more conclusively show that even these

salient visual features (like tool identity) —despite being relevant to the task—are not

sufficient to support dual adaptation in the absence of distinct operational demands.

Unlike prior studies that relied on workspace separation [14,15,17,22,33,34]), explicit lead-in

movements ([6–8,11,21], or movement planning manipulations ([30]), we demonstrate that

tool-specific operational demands alone can serve as effective contextual signals. Our

design represents an important step toward mimicking naturalistic tool use and supports

the idea that tools are encoded primarily by their functional demands rather than their

visual appearance [35–39].

The COIN model [28,40] offers a useful framework for interpreting our results by proposing

that the motor system forms and retrieves internal models based on probabilistic

inference about the current context. Rather than switching between fixed motor

memories, the brain continuously evaluates sensory cues and prior experience to

estimate which motor memory—or combination of memories—is appropriate at a given

moment. In our study, only the Motor Incongruent group demonstrated clear evidence of

dual adaptation, consistent with the idea that distinct tool operations for launching the

ball provided strong, unambiguous cues to signal different contexts. In contrast, tool

identity alone (as in the Color Control and Motor Congruent groups) may have lacked

the contextual specificity required to support reliable contextual inference, resulting in

interference. It remains possible that with extended training, these extrinsic cue

conditions might eventually support the formation of separate motor memories; however,

blocks of eight training trials per visual tool appeared insufficient. Overall, our findings

align with the COIN model's central claim that the reliability and task-relevance of

contextual cues are critical determinants of whether distinct internal models are formed
and retrieved.

Our findings also suggest that the effectiveness of contextual cues in supporting dual
adaptation may not stem solely from prompting the formation of entirely distinct internal
models, but rather from reducing interference from the competing memory—thereby
enhancing retention across blocks with the same cue. As shown in Figures 3 and 4, the
Motor Incongruent group exhibited increasingly stable performance with each return to a
given tool, as interference between the opposing mappings diminished with additional
training. Notably, this did not lead to greater learning *within* each block—at least not in a
clearly observable way. This suggests that the cues or contexts did not facilitate faster
or more complete formation of separate motor memories during each exposure, but
instead supported gradual strengthening and maintenance of those memories through
repeated retrieval with the same cue. From this perspective, dual adaptation may
sometimes reflect improved protection of a motor memory in the face of an opposing
perturbation, rather than the creation of new internal models, underscoring the role of
cue distinctiveness in limiting interference between competing mappings.

This also suggests that paradigms in which cues for different perturbations are
presented in a randomly interleaved fashion—without a minimum number of
consecutive trials per context—may fail to elicit dual adaptation not because the cues
are inherently ineffective, but because they do not allow sufficient time to establish a
stable memory trace for each condition. When exposure to a given cue-perturbation
pairing is too brief or interrupted too frequently, the motor system may be unable to
reinforce the association strongly enough to support retention. Mini-blocked designs,
like the one used here and by others who demonstrate dual adaptation[14,16,28,40–42], may
be especially important for scaffolding the initial stabilization of multiple memories, even
if eventual flexible switching is the end goal. Concurrent adaptation to opposing
perturbations that alternate randomly typically results in weaker dual adaptation, unless
the contextual cues are highly salient or strongly associated with each perturbation
[11,12,15,22,24]. This highlights the importance of considering not just the nature of
contextual cues, but also the structure and temporal continuity of exposure when
designing dual adaptation paradigms.

In conclusion, our results highlight the importance of movement-relevant and
operational cues for organizing multiple motor memories. The findings have practical
implications for tool-based training and rehabilitation, suggesting that designing tools
with distinct control demands may help reduce interference and enhance learning in
complex motor tasks.

References:

1.   Huang, V. S., Haith, A., Mazzoni, P. & Krakauer, J. W. Rethinking Motor Learning
and Savings in Adaptation Paradigms: Model-Free Memory for Successful
Actions Combines with Internal Models. *Neuron* **70**, 787–801 (2011).

2.   Wolpert, D. M., Miall, R. C. & Kawato, M. Internal models in the cerebellum.
Trends Cogn Sci **2**, 338–347 (1998).

3.   Lalazar, H. & Vaadia, E. Neural basis of sensorimotor learning: modifying internal
models. *Curr Opin Neurobiol* **18**, 573–581 (2008).

4.   Krakauer, J. W., Hadjiosif, A. M., Xu, J., Wong, A. L. & Haith, A. M. Motor learning.
Compr Physiol **9**, 613–663 (2019).

5.   Howard, I. S., Ingram, J. N., Franklin, D. W. & Wolpert, D. M. Gone in 0.6
seconds: the encoding of motor memories depends on recent sensorimotor states.
Journal of Neuroscience. **32**, 12756–12768 (2012).

6.   Howard, I. S., Ford, C., Cangelosi, A. & Franklin, D. W. Active lead-in variability
affects motor memory formation and slows motor learning. *Sci Rep* **7**, 7806
(2017).

7.   Howard, I. S., Wolpert, D. M. & Franklin, D. W. The effect of contextual cues on
the encoding of motor memories. *J Neurophysiol* **109**, 2632–2644 (2013).

8.   Howard, I. S., Wolpert, D. M. & Franklin, D. W. The value of the follow-through
derives from motor learning depending on future actions. *Current Biology* **25**,
397–401 (2015).

9.   McGarity-Shipley, M. R. *et al.* Motor memories in manipulation tasks are linked to
contact goals between objects. *J Neurophysiol* **124**, 994–1004 (2020).

10.   Heald, J. B., Ingram, J. N., Flanagan, J. R. & Wolpert, D. M. Multiple motor
memories are learned to control different points on a tool. *Nat Hum Behav* **2**, 300–
311 (2018).

11.   Ayala, M. N. & Henriques, D. Y. P. Differential contributions of implicit and explicit
learning mechanisms to various contextual cues in dual adaptation. *PLoS One* **16**,
e0253948 (2021).

12.   Ayala, M. N. & Henriques, D. Y. P. Context-dependent concurrent adaptation to
static and moving targets. *PLoS One* **13**, e0192476 (2018).

13.   Ayala, M. N., 't Hart, B. M. & Henriques, D. Y. P. Concurrent adaptation to
540   opposing visuomotor rotations by varying hand and body postures. *Exp Brain Res*
541   **233**, 3433–3445 (2015).

542   14.   Vinh, M. A. & Cressman, E. K. Implicit contributions to dual visuomotor adaptation.
543   *Exp Brain Res* **243**, (2025).

544   15.   Baldeo, R. & Henriques, D. Dual adaptation to opposing visuomotor rotations with
545   similar hand movement trajectories. *Exp Brain Res* **227**, 231–241 (2013).

546   16.   Forano, M., Schween, R., Taylor, J. A., Hegele, M. & Franklin, D. W. Direct and
547   indirect cues can enable dual adaptation, but through different learning processes.
548   *J Neurophysiol* **126**, 1490–1506 (2021).

549   17.   Woolley, D. G., Tresilian, J. R., Carson, R. G. & Riek, S. Dual adaptation to two
550   opposing visuomotor rotations when each is associated with different regions of
551   workspace. *Exp Brain Res* **179**, 155–165 (2007).

552   18.   Hegele, M. & Heuer, H. Implicit and explicit components of dual adaptation to
553   visuomotor rotations. *Conscious Cogn* **19**, 906–917 (2010).

554   19.   Wang, L. & Müsseler, J. Concurrent adaptation to opposite visual distortions:
555   Impairment and cue. *Psychol Res* **78**, 453–464 (2014).

556   20.   Sarwary, A. M. E., Stegeman, D. F., Selen, L. P. J. & Medendorp, W. P.
557   Generalization and transfer of contextual cues in motor learning. *J Neurophysiol*
558   **114**, 1565–1576 (2015).

559   21.   Howard, I. S., Franklin, S. & Franklin, D. W. Asymmetry in kinematic
560   generalization between visual and passive lead-in movements are consistent with
561   a forward model in the sensorimotor system. *PLoS One* **15**, e0228083 (2020).

562   22.   Hinder, M. R., Woolley, D. G., Tresilian, J. R., Riek, S. & Carson, R. G. The
563   efficacy of colour cues in facilitating adaptation to opposing visuomotor rotations.
564   *Exp Brain Res* **191**, 143–155 (2008).

565   23.   Modchalingam, S., Ayala, M. N. & Henriques, D. Y. P. Movement-goal relevant
566   object shape properties act as poor but viable cues for the attribution of motor
567   errors to external objects. *PLoS One* **19**, e0300020 (2024).

568   24.   Proud, K. *et al.* Separate motor memories are formed when controlling different
569   implicitly specified locations on a tool. *J Neurophysiol* **121**, 1342–1351 (2019).

570   25.   Modchalingam, S., King, A., 't Hart, B. M. & Henriques, D. Y. P. Visual properties
571   of action consequences and not environments affect context-specific motor
572   learning. *J Neurophysiol* **133**, 1176–1190 (2025).

26. Brookes, J., Warburton, M., Alghadier, M., Mon-Williams, M. & Mushtaq, F.
Studying human behavior with virtual reality: The Unity Experiment Framework.
Behav Res Methods **52**, 455–463 (2020).

27. Ayala, M. N., 't Hart, B. M. & Henriques, D. Y. P. Concurrent adaptation to
opposing visuomotor rotations by varying hand and body postures. *Exp Brain Res*
233, 3433–3445 (2015).

28. Heald, J. B., Lengyel, M. & Wolpert, D. M. Contextual inference underlies the
learning of sensorimotor repertoires. *Nature* **600**, 489–493 (2021).

29. Howard, I. S. & Franklin, D. W. Adaptive tuning functions arise from visual
observation of past movement. *Sci Rep* **6**, 28416 (2016).

30. Sheahan, H. R., Franklin, D. W. & Wolpert, D. M. Motor Planning, Not Execution,
Separates Motor Memories. *Neuron* **92**, 773–779 (2016).

31. McDougle, S. D., Ivry, R. B. & Taylor, J. A. Taking Aim at the Cognitive Side of
Learning in Sensorimotor Adaptation Tasks. *Trends Cogn Sci* **20**, 535–544 (2016).

32. Tsay, J. S. *et al.* Fundamental processes in sensorimotor learning: Reasoning,
refinement, and retrieval. *Elife* **13**, (2024).

33. Schween, R., Langsdorf, L., Taylor, J. A. & Hegele, M. How different effectors and
action effects modulate the formation of separate motor memories. *Sci Rep* **9**,
17040 (2019).

34. Schween, R., Taylor, J. A. & Hegele, M. Plan-based generalization shapes local
implicit adaptation to opposing visuomotor transformations. *J Neurophysiol* **120**,
2775–2787 (2018).

35. Gallivan, J. P., Logan, L., Wolpert, D. M. & Flanagan, J. R. Parallel specification
of competing sensorimotor control policies for alternative action options. *Nat*
Neurosci **19**, 320–326 (2016).

36. Gallivan, J. P., Chapman, C. S., Wolpert, D. M. & Flanagan, J. R. Decision-
making in sensorimotor control. *Nat Rev Neurosci* **19**, 519–534 (2018).

37. Gallivan, J. P., Adam McLean, D., Valyear, K. F. & Culham, J. C. Decoding the
neural mechanisms of human tool use. *Elife* **2013**, (2013).

38. Pazen, M. *et al.* Predictive perception of self-generated movements:
Commonalities and differences in the neural processing of tool and hand actions.
Neuroimage **206**, (2020).

39. Vingerhoets, G. Knowing about tools: Neural correlates of tool familiarity and
experience. *Neuroimage* **40**, 1380–1391 (2008).

40.   Heald, J. B., Lengyel, M. & Wolpert, D. M. Contextual inference in learning and
608   memory. *Trends Cogn Sci* **27**, 43–64 (2023).

609   41.   Cunningham, H. A. & Welch, R. B. Multiple concurrent visual-motor mappings:
610   implications for models of adaptation. *J Exp Psychol Hum Percept Perform* **20**,
611   987–999 (1994).

612   42.   Schween, R., Taylor, J. A. & Hegele, M. Plan-based generalization shapes local
613   implicit adaptation to opposing visuomotor transformations. *J Neurophysiol* **120**,
614   2775–2787 (2018).

Figure legends

Figure 1. Experimental apparatus. A) Virtual environment in the VR experiment at the beginning
of each trial. Once participants grabbed the tool, the timer would begin to deplete, prompting
them to complete their shot toward the target. All four potential target locations — 75^o, 85^o, 95^o,
and 105^o — are shown. B) The virtual environment in the VR experiment at the end of each trial.
Participants score and trials remaining would be displayed on the scoreboard. In this example,
the 75^o target is shown. C) Rendering of the tools in the virtual environment. D) Illustration of
the tool pairs used in each group. Arrows indicate whether a forward or backward movement was
required to shoot the ball. Each Dual Adaptation group (Motor Incongruent, Motor Congruent,
Colour Control) trained with two opposing rotations, each paired with a distinct tool. For the
Single Adaptation groups (shown on the right), only one tool and one rotation direction were
used/presented for a long training block.

Figure 2. Layout of the experimental design. A) Participants in all groups completed 4 different
experimental phases. Every vertical bar represents one block of 8 tool-specific trials. In Baseline
and Washout phases, no perturbation was imposed on the ball (aligned trials). The Training
phase involved a 30^o angular perturbation on the ball's movement direction (rotated trials). In the
Dual Adaptation conditions, participants alternated between tools every block. In the Single
Adaptation condition, participants only alternated tools in the Baseline and Washout phases.
During Training, these participants used one tool for 80 consecutive trials at a time. Tool order
and rotation direction were counterbalanced across all participants. B) Visualization of aligned
and rotated trials for the slingshot, paddle, and curling rod trials. C) Illustration of different trial
types. In aligned trials, the ball rolls forward in the intended aiming direction, while in rotated
trials the ball travels either 30° CW or CCW depending on the tool used.

Figure 3. Launch angle error across tool trial numbers, averaged across participants for each tool
pair, shown for the four groups: A) Motor Incongruent (purple for paddle, red for slingshot), B)
Motor Congruent (purple for paddle, orange for curling rod), C) Colour-Control (purple for
paddle, blue for blue paddle), and D) Single Adaptation (purple for paddle, green for slingshot).
Although tools alternated in blocks of 8 trials, errors for both tools are plotted as overlapping
across the corresponding trial number for each tool. These ten blocks are numbered on the top of
panel A. The horizontal gray bar (±3° centered on the x-axis) indicates the range within which
the 3.3° ball will hit the 3.3° target, relative to the launch point, for a successful hit. Shaded
vertical areas on the left and right of each plot mark aligned trials, with the Baseline phase on the
left and the Washout phase on the right. Only the final 24 trials of Baseline for each tool are
displayed. The unshaded central area represents the Training phase, where ball motion was
deviated 30-degree, with block numbers for this phase shown above panel A. Although the
rotations applied to the ball were in opposite directions for the two tools, the sign of one rotation
was reversed to facilitate comparison. 95% confidence intervals are shown in matching colour
for the two tools of each group.

Figure 4. Angular launch error when adapting to the 30° rotation of ball direction using the
paddle, shown for each participant (hollow circles) and for each of the four groups (full circle),
with 95% confidence intervals across participants. (A) Launch errors averaged over the first and
last trials of the Training phase. (B) Launch errors averaged over the first and last blocks of 8
trials during Training. Dark circles indicate the group mean error, with 95% confidence intervals
across participants. The gray horizontal bar shows the range required for a successful hit, as
illustrated in Figure 3.

Figure 5. Launch errors for first trial (T1) (top row) and final trial (T8) (bottom row) of every
block (A, C) and the first and last block (B, D) for each participant (hollow circles), and
averaged across participants (solid circles, with bars representing 95% confidence interval) for
the three Dual adaptation groups; Motor Incongruent (red), Motor Congruent (orange), Colour
Control (blue). Grey horizontal bar as in Figure 3.

Figure 6. Mean launch error (filled circles) for paddle across all four groups across the Last
block of Baseline and the First Block of Washout. Individual participant error (hollow circles)
and 95% CIs (coloured rectangles) are shown for all trials.

Funding: NSERC grant to DYPH. CFREF trainee funding to AK, SM, and LM.

A

Scoreboard

Surface

Timer

Possible Target Locations

Ball

Tool

Flanking Wall

B

Score

Ball Path

C

40 Cm

Flanking Wall

Launch Point

Slingshot (Elastic)

7 cm

6 cm

Curling Rod

Slingshot (Pouch)

Paddle

D

Tools Pairings

Motor Incongruent   Motor Congruent   Colour Control   Single Adaptation

↑ Forward Movement   ↓ Backward Movement

A

Dual Adaptation

Phase   **Practice**   **Baseline**   **Training**   **Washout**

Blocks

1 block = 8 trials

- ■ Aligned Tool 1
- ■ Aligned Tool 2
- ■ Rotated Tool 1
- ■ Rotated Tool 2

Single Adaptation

Phase   **Practice**   **Baseline**   **Training**   **Washout**

Blocks

1   49   129   211   291 340

Trial #

B

| Slingshot | Paddle | Curling Rod |

Aligned

Rotated

C

Aligned   Rotated

30° CCW   30° CW

Tool 1   Tool 2   Tool 1   Tool 2

Aftereffects per group