Pulmonary Hypertension Promotes Neuroinflammation and Neurodegeneration

Priscilla M. De La Cruz[1,2], Angelia Lockett[1], Marta T. Gomes[1], Somanshu Banerjee[3], Asif Razee[3], Amanda

Fisher[1], Todd Cook[1], Christopher D. Lloyd[4], Shino Magaki[5], Soban Umar[3], Adrian L. Oblak[4], Roberto F.

Machado[1*]

[1] Division of Pulmonary, Critical Care, Sleep and Occupational Medicine, Indiana University, Indianapolis

[2] Division of Pulmonary, Critical Care, and Sleep Medicine, University of New Mexico, Albuquerque

[3] Department of Anesthesiology, University of California Los Angeles Health, Los Angeles

[4] Stark Neuroscience Research Institute, Indianapolis

[5] Department of Pathology and Laboratory Medicine, University of California, Los Angeles

The authors declare no competing interests

*Correspondence to: Roberto Machado, 980 W Walnut Street, Room C400, Walther Hall, R3, Indianapolis, IN

46202, robmacha@iu.edu, 317-278-0047(Phone), 317-278-7030 (Fax)

Author Contributions:

PD- Designed and performed experiments, data analysis, manuscript preparation, AL-Performed and

interpreted cell culture experiments, MG-Performed and interpreted cell culture experiments, SB- Performed

antibody validations, immunostaining and microscopy, AR- Performed antibody validations, immunostaining

and microscopy, AF- Managed animal colonies, TC- Performed and interpreted right heart catheterization, CL-

Assisted with design of immunohistochemistry experiments, SM- Experimental design, technical expertise, SU-

Experimental design, manuscript preparation, AO- Assisted with design of immunohistochemistry experiments,

RM- Experimental design, manuscript preparation, and correspondence.

Funding: NIH T32-HL091816(PD), AHA-CDA 24CDA1274310(SB), NIH NHLBI R01HL160138(SU),

NIH/NHLBI R01HL127342, R01HL111656, and R01HL158108 (RFM)

Abstract

Introduction: Pulmonary arterial hypertension (PAH) is associated with neurocognitive deficits and abnormal brain MRI. Little is known about the mechanisms underlying these clinical observations. TDP-43 is a proteinopathy associated with frontotemporal lobar degeneration (FTLD), Alzheimer's Disease, and Amyotrophic Lateral Sclerosis (ALS). In this study, we hypothesize PAH will result in gliosis, reduced neuronal density, and increased TDP-43 mislocalization. **Methods:** Sprague Dawley rats were randomly assigned to receive Vehicle (DMSO) Monocrotaline, or Sugen/Hypoxia to induce PH. Right heart catheterization was used to confirm PAH. Brain tissue was fixed and probed for microglia (Iba1), astrocytes (GFAP), neurons (NeuN), and TDP-43. Human PH vs control brain tissue was also probed for NeuN and TDP-43. **Results and Conclusions:** We identify an increase in microglia and astrocyte density in the frontal cortex along with reduced neuronal density and neuronal TDP-43 mislocalization in rat models of PH. In addition, human PH frontal cortex demonstrated neuronal TDP-43 mislocalization. This is the first evidence of TDP-43 proteinopathy in PH.

Keywords
Pulmonary Hypertension; Neurodegeneration; Neuroinflammation; TDP-43;

Abbreviations

Pulmonary Hypertension (PH)

TAR DNA-binding protein 43 (TDP-43)

Amyotrophic Lateral Sclerosis (ALS)

Frontotemporal Lobar Degeneration (FTLD)

Pulmonary artery endothelial cell (PAEC)

Central nervous system (CNS)

Dimethyl sulfoxide (DMSO)

Monocrotaline (MCT)

Right Ventricular Systolic Pressure (RVP)

Phosphate Buffered Saline-Triton X (PBST)

Ionized calcium-binding adaptor molecule 1 (Iba1)

Glial Fibrillary Acidic Protein (GFAP)

Neuron-Specific Nuclear Protein (NeuN)

Polymerase Chain Reaction (PCR)

Sugen/Hypoxia (SuHx)

Differentially expressed genes (DEGs)

False discovery rate (FDR)

Introduction

Pulmonary hypertension (PH) is a severe condition, defined by an elevated mean pulmonary arterial pressure exceeding 20 mmHg, and affects approximately 1% of the global population[1]. Clinical studies indicate that patients with PH exhibit cognitive impairments, including challenges in executive function, memory, calculation, and visuospatial abilities[2–4]. A small imaging study reveals a reduction in grey matter volume on brain magnetic resonance imaging when compared to healthy age-matched control subjects, indicating potential neuronal loss [5]. Research on the pathophysiological effects of PH on the central nervous system remains limited. Animal studies have indicated increased concentrations of microglia and astrocytes in the hypothalamus and thoracic spinal cord in subjects with PH compared to control groups, which may correlate with the severity of PH[6,7]. The integration of clinical and animal data suggests the presence of neuroinflammatory and neurodegenerative processes.

TAR DNA-binding protein 43 (TDP-43), a critical regulator of RNA metabolism, has been extensively implicated in neurodegenerative disorders such as amyotrophic lateral sclerosis (ALS) and frontotemporal lobar degeneration (FTLD)[8–10]. Pathological mislocalization and aggregation of TDP-43 disrupt RNA splicing, translation, and intracellular transport—processes essential to both neuronal and endothelial cell function[11,12]. Recent data suggest a potential link between TDP-43 and inflammation, implicating it as a convergent node in pulmonary hypertension (PH)-associated multi-organ dysfunction. Notably, pathologic extranuclear TDP-43 aggregation occurs under pro-inflammatory conditions such as stroke and traumatic brain injury[11,12]. The role of TDP-43 in PH-associated neurodegeneration has not yet been explored.

In this study, we investigated the role of pulmonary hypertension (PH) in cortical neuroinflammation and neurodegeneration through complementary methodologies. We characterized brain tissue from animal models of PH to evaluate astrocyte, microglia, and neuronal densities, as well as TDP-43 proteinopathy. Additionally, we examined neuronal density and TDP-43 proteinopathy in brain tissue samples from PH patients. Given the emerging evidence linking TDP-43 to endothelial dysfunction, we conducted a secondary analysis of pulmonary artery endothelial cell (PAEC) transcriptomic data to identify potential TDP-43-regulated genes. While our studies on animal and human tissues concentrated on central nervous system (CNS) pathology,

these transcriptomic insights suggest a vascular component to the neuroinflammatory processes observed in

PH. We hypothesize that PH induces cortical neuroinflammation, neurodegeneration, and TDP-43

proteinopathy.

Methods

Animal Models of Pulmonary Hypertension and Hemodynamic Measurements

All experimental procedures were sanctioned by the Institutional Animal Care and Use Committee at

Indiana University. Following standard protocol as previously described[13], male and female Sprague Dawley

rats (190-200 g) were procured from Charles River and randomly allocated to receive either Vehicle (Dimethyl

sulfoxide (DMSO), intraperitoneal (ip) vehicle injection as a control), Monocrotaline (MCT; ip, 60 mg/kg of body

weight), or Sugen (subcutaneously, 20 mg/kg body weight). Rats administered Sugen were housed in a 10%

oxygen chamber for three weeks, followed by two weeks under normoxia conditions. At five weeks post-

injection, the rats underwent open field testing, as previously described[14]. Briefly, the rats were placed in an

open field maze, and measurements of distance traveled, time spent in the center, time at rest, and average

ambulatory velocity were recorded using Any-maze software. Right Ventricular Systolic Pressure (RVP) was

assessed via right heart catheterization employing a Millar pressure transducer catheter at the time of

euthanasia. The Fulton index was calculated by measuring the ratio of the right ventricle weight to the sum of

the weights of the left ventricle and septum (RV/(LV+S)). Brain tissue and plasma were collected at the time of

euthanasia.

Rat Brain Tissue Immunofluorescence Staining

The brains were meticulously dissected along the intracerebral fissure, with one half snap-frozen in

liquid nitrogen and stored at -80°C for subsequent RNA analyses. The remaining half was allocated for

immunofluorescence studies and was fixed with 4% paraformaldehyde overnight, followed by a 24-hour

treatment with 30% sucrose. The brains were then stored in optimal tissue cutting medium at -80°C until

sectioning. Coronal sections were prepared at a thickness of 30 micrometers and preserved in cryoprotectant

solution (FD NeuroTechnologies, Inch, Cat#PC102). The sections underwent heat-mediated antigen retrieval

for 30 minutes at 80°C (Universal HEIR antigen retrieval reagent, Abcam#ab208572). Subsequently, the

sections were washed with 0.3% Phosphate Buffered Saline-Triton X-100 (PBST) and subjected to a 10-

minute treatment with 0.3% Hydrogen Peroxide block. Following additional washing, the sections were

incubated with 10% Donkey Serum (Abcam, #NC1697010) for 1 hour at room temperature. The sections were

then incubated overnight with primary antibodies (Supplemental Table 1). After the overnight incubation, the

sections were washed with PBST, blocked with 10% Donkey Serum, and incubated with secondary antibodies

(1:1000) for 2 hours. The sections were then washed with PBST and mounted on slides using ProLong Gold

Antifade Mountant with DNA Stain DAPI (Invitrogen, #P36931). Imaging was conducted using the Leica

Thunder System (20x magnification unless specified), and the images were analyzed using Imaris (Oxford

Instruments) to quantify cells positively expressing both DAPI and cell-specific marker.

Rat Brain Quantitative Polymerase Chain Reaction (PCR)

RNA was extracted from hemibrains utilizing the RNeasy Plus Mini Kit (Qiagen, # 74134) in accordance

with the manufacturer's protocol. The quality of the RNA was evaluated using the NanoDrop One.

Subsequently, RNA was transcribed into cDNA employing the High-Capacity cDNA Reverse Transcription Kit

(Applied Biosystems, # 4368814) with the Bio-rad T100 Thermal Cycler. Relative quantitative RT-PCR was

conducted using the Roche Lightcycler 480 and the PowerUP SYBR Green Master Mix chemistry (Applied

Biosystems, # A25742) to quantify cytokine gene expression. The primers used were as follows: *IL-6* (Forward

'TCCTACCCCAACTTCCAATGCTC', Reverse 'TTGGATGGTCTTGGTCCTTAGCC'), *IL-1β* (Forward

'CACCTCTCAAGCAGAGCACAG', Reverse 'GGGTTCCATGGTGAAGTCAAC'), *TNF-α* (Forward

'AAATGGGCTCCCTCTCATCAGTTC', Reverse 'TCTGCTTGGTGGTTTGCTACGAC'), and *Gapdh* (Forward

'CATCACTGCCACCCAGAAGACTG', Reverse 'ATGCCAGTGAGCTTCCCGTTCAG'). The fold change was

calculated using the $2^{-\Delta\Delta CT}$ method, with results normalized to *Gapdh*.

Human Cortical Tissue Immunohistochemistry

In accordance with 45 CFR 46, our study does not involve human subjects but used deidentified human

tissues provided by UCLA Pathology departmental honest broker, the UCLA Translational Pathology Core Lab

(IRB# 11-002504). Paraffin-embedded 5 micrometer thick human brain cortex sections (Control n=3, PH n=4)

were rehydrated, and antigen retrieval was performed by heat-induced epitope retrieval at pH 6 (C9999,

Sigma). Tissue sections were blocked for an hour at room temperature in PBS with 5% goat serum, 1% BSA,

0.1% Triton, 0.05% Tween, and 0.3 M glycine. Tissue sections were incubated overnight with primary

antibodies diluted 1:200 in PBS with 1% BSA and 0.5% Triton at 4°C. Secondary antibodies were incubated for

an hour at room temperature and diluted 1:1000 in PBS with 0.05% Tween. Sections were incubated with

ProLong™ Gold antifade mountant with DNA Stain DAPI and cover slips were applied. All images were

acquired using a confocal microscope (40x magnification) (Nikon). A total of 4-5 images were acquired from

each slide. Images were analyzed using Imaris (Oxford Instruments).

Secondary Analysis of TDP-43-Regulated Genes in Pulmonary Artery Endothelial Cells

Raw gene expression count data were procured from the NCBI Gene Expression Omnibus (GEO)

under accession number GSE243193, encompassing RNA-sequencing data from human pulmonary artery

endothelial cells (PAECs) isolated from patients with group 1 pulmonary hypertension (PH) (n=20) and controls

(n=3) [15]. Group 1 PH patients were selected as they more closely align with the animal models represented in

this study. The dataset was imported into R (version 4.2.0) and underwent preprocessing to eliminate genes

with missing or NA values across samples[16]. Following quality control, expression counts were converted into a

matrix. Samples were categorized as "PH" or "Control" based on their column names, and group labels were

encoded as a factor variable for subsequent differential expression analysis. Gene expression normalization

and differential analysis were conducted using the edgeR and limma packages[17,18]. Expression counts were

filtered using filterByExpr() to retain genes with adequate counts across samples. Library sizes were

normalized using the trimmed mean of M-values (TMM), and log2-transformed counts per million (CPM) were

calculated via voom to model the mean-variance relationship[19]. Linear modeling was executed using lmFit(),

followed by empirical Bayes moderation using eBayes() to enhance variance estimation. Differentially

expressed genes (DEGs) were identified based on false discovery rate (FDR)-adjusted p-values < 0.05. Gene

annotation was performed using the org.Hs.eg.db Bioconductor package to map ENTREZ identifiers to gene

symbols and gene names[20]. To evaluate the enrichment of TDP-43 regulatory activity in PH-associated gene

expression changes, a curated list of TDP-43 RNA targets was obtained from POSTAR3 CLIP-seq datasets[21].

POSTAR-3 utilizes data from the ENCODE project which identified TDP-43 targets under baseline conditions.

Although these targets were identified in non-diseased immortalized cells, they provide a reference map of

physiologic TDP-43 binding sites, which may be dysregulated or lost in disease-specific contexts such as

nuclear exclusion[22].The overlap between differentially expressed genes (FDR < 0.05) and TDP-43 targets was

calculated. A Fisher's exact test was conducted to assess whether TDP-43 targets were statistically

overrepresented among DEGs compared to the background set of all expressed genes. A 2×2 contingency

table was constructed with counts of genes categorized as TDP-43 target vs. non-target and DEG vs. non-

DEG. The resulting odds ratio and p-value were utilized to determine enrichment significance. A volcano plot

was generated using ggplot2, with upregulated and downregulated genes highlighted. DEGs that overlapped

with TDP-43 targets[23].

Statistical Analysis

Statistical analyses were performed utilizing GraphPad Prism (version 8.4.3; GraphPad Software) and R

(version 4.2.0), as delineated in the preceding section. To compare experimental groups (e.g., SuHx vs. Vehicle

and MCT vs. Vehicle), an unpaired two-tailed Student's t-test was employed. Linear regression analysis was

conducted to assess correlations between variables. Pearson's correlation coefficient (r) and the coefficient of

determination (R^2) were calculated and reported. Data are presented as mean ± standard error of the mean

(SEM) unless otherwise specified. A p-value < 0.05 was considered indicative of statistical significance.

Results

SuHx-Induced Pulmonary Hypertension is Associated with Elevated RVP and Hypertrophy

Sugen/hypoxia (SuHx) and monocrotaline (MCT) are established models for the study of pulmonary

hypertension (PH) and were employed to induce PH in rats[24]. The presence of pulmonary hypertension was

confirmed through measurements of right ventricular pressure (RVP) and right ventricular hypertrophy (Fulton

Index). The SuHx model (n=6) elicited a pronounced PH phenotype, evidenced by a significant increase in RVP

(mean difference of 52.33 ± 11.84 mmHg compared to vehicle controls, n=6; p=0.0013) and the Fulton index

(mean difference of 0.2428 ± 0.03486, p<0.0001) (**Figure 1**). Conversely, the MCT model (n=6) demonstrated

a trend towards increased RVP (mean difference of 17.33 ± 6.98 mmHg versus controls, p=0.0522) and a modest

yet statistically significant increase in the Fulton Index (mean difference of 0.09815 ± 0.03066 versus controls,

p=0.0198) (**Supplemental Figure 1**). Although this study was not designed or powered to assess sex differences

(n=3 males and n=3 females per group), exploratory analysis indicated a sex-dependent effect in both models.

In SuHx-treated rats, males exhibited higher RVP (mean: 102.67 mmHg) and Fulton Index (mean: 0.53)

compared to females (mean RVP: 52 mmHg; mean Fulton index: 0.41). A similar pattern was observed in MCT-

treated animals, with males showing higher RVP (mean: 56 mmHg) and Fulton Index (mean: 0.36) compared to

females (mean RVP of 27 mmHg; mean Fulton index of 0.28). These findings align with existing literature on sex

differences in experimental PH and highlight the necessity of considering sex as a biological variable in preclinical

studies[25]. No statistically significant sex differences were observed in any of the subsequent measurements.

This suggests a need for further exploration of the impact of sex as a biological variable in neuroinflammation in

PH. Due to the relatively mild phenotype induced by MCT in our study (likely due to the modest effect on female

mice), all MCT-associated findings, including cardiovascular and central nervous system assessments, are

included in the supplemental materials.

No Behavioral Abnormalities are Observed in Open Field Testing

Open field testing (OFT) is widely used to assess sickness behavior, locomotor activity, and anxiety-like

behavior in rodent models of systemic disease, including those of sepsis-associated encephalopathy, heart

failure, and frontotemporal lobar degeneration[26–28]. In this study, OFT was employed to evaluate whether PH

was associated with changes in exploratory behavior or activity levels. SuHx-treated rats showed no significant

differences in total distance traveled, time spent in the center of the arena, time spent at rest, or average

ambulatory velocity compared to vehicle-treated controls (**Figure 2**). Similarly, no statistically significant

differences were observed between MCT-treated rats and controls (**Supplemental Figure 2**). OFT data did not

correlate significantly with RVP or Fulton in either model (data not shown).

Experimental PH Induces Neuroinflammation

Neuroinflammation is recognized as an early and critical risk factor for neurodegeneration[29,30]. In this study, we measured cortical density of microglia and astrocytes in two rodent models of PH. In SuHx-treated rats, microglial density was significantly increased ($3.875 \times 10^{-5} \pm 1.504 \times 10^{-5}$ cells/μm^2 compared to vehicle-treated controls (n = 6; p = 0.0276). Astrocyte density was also elevated in SuHx animals ($4.487 \times 10^{-5} \pm 1.806 \times 10^{-5}$ cells/μm^2, p = 0.0323) (**Figure 3**). Notably, neither microglia nor astrocytic density correlated with RVP or Fulton Index in the SuHx group (data not shown). In the MCT-treated rats, a significant increase in microglial density (0.2320 ± 0.009541 cells/μm^2) compared to vehicle-treated controls (n = 6; p <0.0001) was observed (**Supplemental Figure 3**). Similarly, astrocyte density was elevated in MCT-treated animals relative to control ($4.822 \times 10^{-5} \pm 1.309 \times 10^{-5}$ cells/μm^2, p= 0.0043) (**Supplemental Figure 3**). Unlike the SuHx-treated animals, cortical microglia density positively correlated with the RVP (r= 0.7670, 95% CI: 0.3449 to 0.9311, p=0.0036) and Fulton Index (r= 0.7990, 95% CI: 0.4156 to 0.9413, p=0.0018). Despite increased glial density, there was no increase in cortical expression of pro-inflammatory cytokines—including IL-1β, TNF-α, or IL-6 mRNA— between SuHx- or MCT-treated animals and their respective controls, with SuHx data presented in **Figure 4** and MCT data presented in **Supplemental Figure 4**

Experimental PH Induces Neuronal Loss and TDP-43 Proteinopathy

Neurodegeneration is characterized by the progressive loss of neurons and associated structural and functional decline[31]. A hallmark of several neurodegenerative diseases is TDP-43 proteinopathy, defined by the pathological mislocalization of TAR DNA-binding protein 43 (TDP-43) from the nucleus to the cytoplasm. While traditionally linked to conditions such as amyotrophic lateral sclerosis (ALS) and frontotemporal lobar degeneration (FTLD), TDP-43 pathology has also been identified in proinflammatory states, including traumatic brain injury and stroke[8,9,11,12]. In the SuHx model, we observed a significant reduction in cortical neuronal density compared to vehicle controls (n = 6), with a mean difference of $-0.0001113 \pm 2.937 \times 10^{-5}$ cells/μm^2 (p = 0.0035). SuHx-treated animals also exhibited a significant increase in neuronal extranuclear TDP-43, with a mean difference of 0.2592 ± 0.0432 (p < 0.0001). Linear regression analysis revealed that right ventricular pressure (RVP) significantly predicted neuronal density (R^2 = 0.5825, p = 0.0039) and the proportion of neurons with extranuclear TDP-43 (R^2 = 0.4029, p = 0.0266) (**Figure 5**). Similarly, the Fulton index strongly

predicted both neuronal density (R^2 = 0.6898, p = 0.0008) and extranuclear TDP-43 burden (R^2 = 0.6001, p =

0.0031) (data not shown).

In contrast, MCT-treated animals showed no significant change in neuronal density relative to controls

(n = 6). However, MCT treatment was associated with a significant increase in neuronal extranuclear TDP-43

(mean difference: 0.2553 ± 0.0762, p = 0.0194) (**Supplemental Figure 5**). Notably, in the MCT group, neither

RVP nor Fulton index correlated with neuronal density or extranuclear TDP-43.

Clinical PH Induces Cortical TDP-43 Proteinopathy

To investigate the potential impact of pulmonary hypertension (PH) on neurodegeneration and TDP-43

proteinopathy in humans, we analyzed postmortem cortical brain tissue from individuals with PH (n = 4) and

non-PH controls (n = 3). Quantitative analysis revealed no significant difference in cortical neuronal

density between the two groups. However, neuronal extranuclear TDP-43 was significantly increased in PH

subjects compared to controls, with a mean difference of 46.27 ± 12.96 (p = 0.0230) (**Figure 6**). Demographics

available in supplemental table 2.

PH Disproportionately Affects Expression of TDP-43 Regulated Genes in Pulmonary Artery Endothelial

Cells (PAECs)

Emerging evidence suggests that TDP-43 is essential for endothelial cell function and RNA

homeostasis[10,32]. This may represent a critical mechanistic link between TDP-43 proteinopathy and pulmonary

hypertension (PH), implicating endothelial dysregulation as a contributor to the neurodegenerative processes

observed in PH. To explore this hypothesis, we performed a secondary analysis of single-cell RNA sequencing

data (GSE243193) from pulmonary artery endothelial cells[15]. We identified TDP-43 target transcripts using

POSTAR3 CLIP-seq data and applied Fisher's exact test to determine whether PH disproportionately

dysregulates TDP-43–regulated genes[21]. Top downregulated genes were manually annotated using

GeneCards[33]. This analysis revealed a significant overrepresentation of TDP-43–associated transcripts among

the downregulated genes in PH (Fisher's exact test: $p < 2.2 \times 10^{-16}$, OR: 0.193, 95% CI: 0.139-0.269),

suggesting selective vulnerability of this RNA-regulatory pathway. Notably, many of the top downregulated

TDP-43 targets are involved in RNA processing (EIF4A1, HNRNPH2), vesicular trafficking (SNX15, CHMP3,

RAB4B), and inflammation (MIF) (**Figure 7**).

Discussion

Pulmonary hypertension (PH) is increasingly acknowledged as a systemic condition that affects not

only the pulmonary vasculature and the right ventricle but also encompasses cognitive, affective, and

neuroinflammatory consequences[2–4,7]. While prior clinical studies have documented neurocognitive symptoms

in PH patients, the underlying pathophysiology linking pulmonary vascular disease to central nervous system

(CNS) injury has remained poorly defined. Although previous clinical studies have documented neurocognitive

symptoms in patients with PH, the pathophysiological mechanisms linking pulmonary vascular disease to

central nervous system (CNS) injury remain inadequately defined. In this study, we present preclinical and

translational evidence suggesting that gliosis and TDP-43 proteinopathy are contributing factors to the

development of neurodegeneration in PH.

Our findings elucidate significant distinctions between the MCT and SuHx models of pulmonary

hypertension (PH). Both models indicate a sex-related disparity in PH and vascular remodeling, aligning with

existing literature[34]. As previously reported, the MCT induced a less severe PH phenotype in females, which

resulted in right ventricular hypertrophy without elevated right ventricular pressure[34]. Notably, there are

differences in central nervous system (CNS) findings between these models. Both models exhibit gliosis and

TDP-43 proteinopathy; however, the more severe SuHx model also shows reduced neuronal density. The

severity of PH correlates with TDP-43 proteinopathy and neuronal loss in the SuHx model, whereas in the

MCT model, it predicts microglial activation. This suggests that CNS findings may occur on a spectrum, with

milder forms of PH inducing neuroinflammation, while more severe PH leads to neuronal loss. The SuHx and

MCT models of pulmonary hypertension exhibit distinct pathological mechanisms, which may have implications

for extrapulmonary involvement, including the brain. The SuHx model combines VEGF receptor inhibition with

chronic hypoxia to induce severe, angio-obliterative pulmonary vascular remodeling. This model simulates a

progressive, endothelial-driven vasculopathy, which may potentially exacerbate blood-brain barrier (BBB)

dysfunction and cerebral microvascular injury.[35]. Monocrotaline, in contrast, induces pulmonary hypertension

through a singular, systemic endothelial toxin that preferentially damages the pulmonary vasculature via

hepatic metabolism. Although effective in promoting vascular remodeling and right ventricular hypertrophy,

monocrotaline is associated with systemic toxicity, including hepatic and renal injury, and may exert more

variable effects on the cerebral vasculature. [35]. Given the critical role of hypoxia in influencing neurovascular

integrity and mitochondrial metabolism, the Sugen hypoxia model may more accurately represent brain-

relevant systemic stressors, such as chronic cerebral hypoxemia. In contrast, the MCT model may complicate

interpretation due to its extensive systemic toxicities. Our limited clinical data revealed an increase in TDP-43

proteinopathy among patients with PH compared to control subjects.

TDP-43, a nuclear RNA-binding protein with established roles in neurodegenerative diseases such as

ALS and frontotemporal dementia, has recently been implicated in vascular and endothelial cell biology[10,32] In

these studies, the depletion of endothelial TDP-43 resulted in diminished expression of tight junction proteins,

such as claudin-5, heightened permeability, and the activation of microglia and astrocytes.[36,37]. Although our

studies on animal and human tissues concentrated on neuronal pathology, we hypothesized that endothelial

dysfunction might contribute to PH-associated neuroinflammation through TDP-43–mediated post-

transcriptional dysregulation. To investigate this hypothesis, we conducted a secondary analysis of single-cell

RNA sequencing data from pulmonary artery endothelial cells (GSE243193), focusing on transcripts known to

be regulated by TDP-43 using POSTAR3 CLIP-seq data[15,21]. This analysis demonstrated that PH

disproportionately downregulates TDP-43 target genes in endothelial cells, with key affected transcripts

involved in RNA metabolism (EIF4A1, HNRNPH2), vesicular trafficking (CHMP3, SNX15, RAB4B),

mitochondrial homeostasis (MRPS24, PLSCR3), and inflammatory signaling (MIF)[33]. These findings support a

model in which TDP-43–associated transcriptomic dysregulation may impair endothelial function in PH,

potentially contributing to downstream neuroinflammation and blood-brain barrier vulnerability. While we did

not directly assess blood-brain barrier (BBB) integrity in this study, the observed gene expression changes

implicate pathways known to affect endothelial stability. Importantly, these data do not suggest that TDP-43

dysfunction is merely a consequence of chronic hypoxia or right heart failure. Rather, they propose that TDP-43 dysregulation is a shared molecular feature of both pulmonary vascular and neurologic injury in PH. This hypothesis aligns with recent literature demonstrating the role of TDP-43 in oxidative stress responses and RNA transport in both neurons and endothelial cells[9,10,32,38].

Our study has several limitations. Firstly, although the neuroinflammatory and degenerative alterations were pronounced in both animal and human PH brain tissue, we did not directly demonstrate functional impairments, such as behavioral or cognitive deficits. Further work is needed to phenotype cognitive impairment in PH animal models. While an open field test is beneficial for identifying locomotion, anxiety, and sickness behavior in animal models, it may not effectively simulate the loss of executive function observed in frontotemporal lobar degeneration (FTLD), which is also challenging to differentiate from motor dysfunction in a clinical context [39]. Next, quantitative PCR analysis did not reveal a significant difference in proinflammatory cytokine mRNA levels, indicating that further investigation is required to determine the timing of neuroinflammation in relation to the onset of pulmonary hypertension. An additional limitation is the use of a limited data tissue biobank utilized to investigate TDP-43 in PH patients. This dataset does not include information regarding PH duration or severity which could influence the results. In addition, while our transcriptomic analysis indicates endothelial contributions to the TDP-43 signature, further research is necessary to validate these targets at the protein level and to assess causality using endothelial-specific TDP-43 models. Lastly, our study was not powered to assess sex as a biological variable which may be an important direction for future studies. In sum, the temporal sequence of neuroinflammation, TDP-43 mislocalization, endothelial dysfunction, and CNS injury remains to be elucidated.

Despite these limitations, our findings establish a foundational basis for a novel conceptual model of PH as a multisystem disorder characterized by mechanistic links to RNA dysregulation and neurodegeneration. Targeting TDP-43 or its downstream pathways may present therapeutic opportunities to mitigate both vascular and neurological complications in this population. Future research focused on elucidating the cell-specific roles of TDP-43 within the lung-brain axis may yield further insights into the systemic nature of pulmonary vascular disease.

Conclusion:

This study identifies TDP-43 dysregulation as a potential mechanistic link between pulmonary hypertension and central nervous system injury. Through the integration of animal models, human brain tissue, and endothelial transcriptomic analysis, we demonstrate that PH is associated with neuroinflammatory and neurodegenerative changes, alongside selective downregulation of TDP-43–regulated genes in pulmonary artery endothelial cells. These findings support a novel model in which TDP-43 contributes to multisystem dysfunction in PH, bridging vascular and neurologic pathology. Elucidating the role of TDP-43 across organ systems may open new avenues for targeted therapies aimed at mitigating both cardiopulmonary and cognitive complications of PH.

Acknowledgements: This work was supported by the Behavioral Phenotyping Core (BPC) at the Indiana University School of Medicine.

References

1. Hoeper MM, Humbert M, Souza R, et al. A global view of pulmonary hypertension. *Lancet Respir Med*. 2016;4(4):306-322. doi:10.1016/S2213-2600(15)00543-3

2. Heller S, See C, Singh I, Fredericks CA. Cognitive impairment in pulmonary arterial hypertension. *Ann Clin Transl Neurol*. 2023;10(10):1899-1903. doi:10.1002/acn3.51867

3. White J, Hopkins RO, Glissmeyer EW, Kitterman N, Elliott CG. Cognitive, emotional, and quality of life outcomes in patients with pulmonary arterial hypertension. *Respir Res*. 2006;7(1):55. doi:10.1186/1465-9921-7-55

4. Yuan P, Li J, Liu J, et al. Cognitive Dysfunction in Patients with Pulmonary Hypertension. *Am J Respir Crit Care Med*. 206(10):1289-1293. doi:10.1164/rccm.202204-0726LE

5. Roy B, Vacas S, Ehlert L, McCloy K, Saggar R, Kumar R. Brain Structural Changes in Patients with Pulmonary Arterial Hypertension. *J Neuroimaging Off J Am Soc Neuroimaging*. 2021;31(3):524-531. doi:10.1111/jon.12840

6. Sharma RK, Oliveira AC, Kim S, et al. Involvement of Neuroinflammation in the Pathogenesis of Monocrotaline-Induced Pulmonary Hypertension. *Hypertens Dallas Tex 1979*. 2018;71(6):1156-1163. doi:10.1161/HYPERTENSIONAHA.118.10934

7. Razee A, Banerjee S, Hong J, et al. Thoracic Spinal Cord Neuroinflammation as a Novel Therapeutic Target in Pulmonary Hypertension. *Hypertension*. 2023;80(6):1297-1310. doi:10.1161/HYPERTENSIONAHA.122.20782

8. Jo M, Lee S, Jeon YM, Kim S, Kwon Y, Kim HJ. The role of TDP-43 propagation in neurodegenerative diseases: integrating insights from clinical and experimental studies. *Exp Mol Med*. 2020;52(10):1652-1662. doi:10.1038/s12276-020-00513-7

9. Zuo X, Zhou J, Li Y, et al. TDP-43 aggregation induced by oxidative stress causes global mitochondrial imbalance in ALS. *Nat Struct Mol Biol*. 2021;28(2):132-142. doi:10.1038/s41594-020-00537-7

10. Cheemala A, Kimble AL, Burrage EN, et al. Amyotrophic lateral sclerosis and frontotemporal dementia mutation reduces endothelial TDP-43 and causes blood-brain barrier defects. *Sci Adv*. 2025;11(16):eads0505. doi:10.1126/sciadv.ads0505

11. Thammisetty SS, Pedragosa J, Weng YC, Calon F, Planas A, Kriz J. Age-related deregulation of TDP-43 after stroke enhances NF-κB-mediated inflammation and neuronal damage. *J Neuroinflammation*. 2018;15(1):312. doi:10.1186/s12974-018-1350-y

12. Bjorklund GR, Wong J, Brafman D, Bowser R, Stabenfeldt SE. Traumatic brain injury induces TDP-43 mislocalization and neurodegenerative effects in tissue distal to the primary injury site in a non-transgenic mouse. *Acta Neuropathol Commun*. 2023;11(1):137. doi:10.1186/s40478-023-01625-7

13. Chen J, Lockett A, Zhao S, et al. Sphingosine Kinase 1 Deficiency in Smooth Muscle Cells Protects against Hypoxia-Mediated Pulmonary Hypertension via YAP1 Signaling. *Int J Mol Sci*. 2022;23(23):14516. doi:10.3390/ijms232314516

14. Seibenhener ML, Wooten MC. Use of the Open Field Maze to Measure Locomotor and Anxiety-like Behavior in Mice. *J Vis Exp JoVE*. 2015;(96):52434. doi:10.3791/52434

15. Singh N, Eickhoff C, Garcia-Agundez A, et al. Transcriptional profiles of pulmonary artery endothelial cells in pulmonary hypertension. *Sci Rep*. 2023;13(1):22534. doi:10.1038/s41598-023-48077-6

16. R: The R Project for Statistical Computing. Accessed June 30, 2025. https://www.r-project.org/

17. Robinson MD, McCarthy DJ, Smyth GK. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics*. 2010;26(1):139-140. doi:10.1093/bioinformatics/btp616

18. Ritchie ME, Phipson B, Wu D, et al. limma powers differential expression analyses for RNA-sequencing and microarray studies. *Nucleic Acids Res*. 2015;43(7):e47. doi:10.1093/nar/gkv007

19. Law CW, Chen Y, Shi W, Smyth GK. voom: precision weights unlock linear model analysis tools for RNA-seq read counts. *Genome Biol*. 2014;15(2):R29. doi:10.1186/gb-2014-15-2-r29

20. org.Hs.eg.db. Bioconductor. Accessed June 30, 2025. http://bioconductor.org/packages/org.Hs.eg.db/

21. Zhao W, Zhang S, Zhu Y, et al. POSTAR3: an updated platform for exploring post-transcriptional regulation coordinated by RNA-binding proteins. *Nucleic Acids Res*. 2022;50(D1):D287-D294. doi:10.1093/nar/gkab702

22. Van Nostrand EL, Pratt GA, Yee BA, et al. Principles of RNA processing from analysis of enhanced CLIP maps for 150 RNA binding proteins. *Genome Biol*. 2020;21(1):90. doi:10.1186/s13059-020-01982-9

23. Wickham H. *Ggplot2: Elegant Graphics for Data Analysis*. Springer; 2009. doi:10.1007/978-0-387-98141-3

24. Boucherat O, Agrawal V, Lawrie A, Bonnet S. The Latest in Animal Models of Pulmonary Hypertension and Right Ventricular Failure. *Circ Res*. 2022;130(9):1466-1486. doi:10.1161/CIRCRESAHA.121.319971

25. Bal E, Ilgin S, Atli O, Ergun B, Sirmagul B. The effects of gender difference on monocrotaline-induced pulmonary hypertension in rats. *Hum Exp Toxicol*. 2013;32(7):766-774. doi:10.1177/0960327113477874

26. Swiergiel AH, Dunn AJ. Effects of Interleukin-1β and Lipopolysaccharide on Behavior of Mice in the Elevated Plus-Maze and Open Field Tests. *Pharmacol Biochem Behav*. 2007;86(4):651-659. doi:10.1016/j.pbb.2007.02.010

27. Dridi H, Liu Y, Reiken S, et al. Heart failure-induced cognitive dysfunction is mediated by intracellular Ca2+ leak through ryanodine receptor type 2. *Nat Neurosci*. 2023;26(8):1365-1378. doi:10.1038/s41593-023-01377-6

28. Frew J, Nygaard HB. Neuropathological and behavioral characterization of aged Grn R493X progranulin-deficient frontotemporal dementia knockin mice. *Acta Neuropathol Commun*. 2021;9:57. doi:10.1186/s40478-021-01158-x

29. Zhang W, Xiao D, Mao Q, Xia H. Role of neuroinflammation in neurodegeneration development. *Signal Transduct Target Ther*. 2023;8(1):1-32. doi:10.1038/s41392-023-01486-5

30. Kempuraj D, Thangavel R, Natteru P, et al. Neuroinflammation Induces Neurodegeneration. *J Neurol Neurosurg Spine*. 2016;1(1):1003.

31. Przedborski S, Vila M, Jackson-Lewis V. Series Introduction: Neurodegeneration: What is it and where are we? *J Clin Invest*. 2003;111(1):3-10. doi:10.1172/JCI17522

32. Arribas V, Onetti Y, Ramiro-Pareta M, et al. Endothelial TDP-43 controls sprouting angiogenesis and vascular barrier integrity, and its deletion triggers neuroinflammation. *JCI Insight*. 2024;9(5). doi:10.1172/jci.insight.177819

33. GeneCards - Human Genes | Gene Database | Gene Search. Accessed June 30, 2025. https://www.genecards.org/

34. Bal E, Ilgin S, Atli O, Ergun B, Sirmagul B. The effects of gender difference on monocrotaline-induced pulmonary hypertension in rats. *Hum Exp Toxicol*. 2013;32(7):766-774. doi:10.1177/0960327113477874

35. Wu XH, Ma JL, Ding D, Ma YJ, Wei YP, Jing ZC. Experimental animal models of pulmonary hypertension: Development and challenges. *Anim Models Exp Med*. 2022;5(3):207-216. doi:10.1002/ame2.12220

36. Cheemala A, Kimble AL, Tyburski JD, et al. Loss of Endothelial TDP-43 Leads to Blood Brain Barrier Defects in Mouse Models of Amyotrophic Lateral Sclerosis and Frontotemporal Dementia. *bioRxiv*. Published online December 14, 2023:2023.12.13.571184. doi:10.1101/2023.12.13.571184

37. Arribas V, Onetti Y, Ramiro-Pareta M, et al. Endothelial TDP-43 controls sprouting angiogenesis and vascular barrier integrity, and its deletion triggers neuroinflammation. *JCI Insight*. 2024;9(5). doi:10.1172/jci.insight.177819

38. Colombrita C, Zennaro E, Fallini C, et al. TDP-43 is recruited to stress granules in conditions of oxidative insult. *J Neurochem*. 2009;111(4):1051-1061. doi:10.1111/j.1471-4159.2009.06383.x

39. Huey ED, Goveia EN, Paviol S, et al. Executive dysfunction in frontotemporal dementia and corticobasal syndrome. *Neurology*. 2009;72(5):453-459. doi:10.1212/01.wnl.0000341781.39164.26

1A

1B

Figure 1. Sugen-hypoxia induces pulmonary hypertension and vascular remodeling.
Rats treated with Sugen-hypoxia (SuHx; n = 6) exhibit increased right ventricular pressure
(RVP), with a mean difference of 52.33 ± 11.84 mmHg compared to vehicle controls (n = 6;
p = 0.0013) (**1A**). SuHx-treated rats also demonstrate right ventricular hypertrophy, with a mean
Fulton Index difference of 0.2428 ± 0.03486 compared to controls (p < 0.0001) (**1B**). Data
analyzed using a two-tailed unpaired t-test and are presented as mean ± SEM. Female animals
are depicted in blue.

Figure 2. Sugen hypoxia does not induce sickness behavior in rats undergoing open field test.
Rats exposed to Sugen hypoxia (SuHx, n = 9) did not exhibit significant differences in behavioral
parameters compared to vehicle-treated controls (n = 10). Specifically, there were no significant
changes in total distance traveled (**2A**), time spent in the center of the open field test (**2B**), time spent at
rest (**2C**), or average ambulatory velocity (**2D**). Data were analyzed using an unpaired two-tailed t-test
and are presented as mean ± SEM.

Figure 3. Sugen hypoxia increases cortical gliosis. Representative immunofluorescence images of frontal cortex microglia (IBA1/DAPI) (**3A**) and astrocytes (GFAP/DAPI) (**3B**) in rats exposed to sugen/hypoxia (SuHx) or vehicle. SuHx-treated rats (n = 6) exhibited a significantly increased microglial density ($3.875 \times 10^{-5} \pm 1.504 \times 10^{-5}$) compared to vehicle-treated controls (n = 6; p = 0.0276) (**3C**). Similarly, astrocyte density was elevated in SuHx-treated animals ($4.487 \times 10^{-5} \pm 1.806 \times 10^{-5}$) relative to controls (p = 0.0323) (**3D**). Data are presented as mean ± SEM and were analyzed using a two-tailed unpaired t-test.

Figure 4. Sugen/hypoxia does not increase cortical cytokine expression. Quantitative PCR analysis of cortical cytokine mRNA expression in rats treated with sugen/hypoxia (SuHx; n = 6) or vehicle (n = 6) revealed no significant differences in IL-1β (**4A**), TNF-α (**4B**), or IL-6 (**4C**) transcript levels. Data are presented as mean ± SEM and were analyzed using a two-tailed unpaired t-test.

Figure 5. Sugen/hypoxia treatment reduces cortical neuronal density and induces neuronal TDP-43 mislocalization. Representative immunofluorescence images of cortical neurons stained for NeuN (neuronal marker) and DAPI (nuclear marker), TDP-43 and DAPI, and merged channels in rats exposed to vehicle at 40x magnification (**5A**) compared to sugen/hypoxia-treated rats (SuHx). The yellow arrow highlights dysmorphic neurons in SuHx-treated rats (**5B**). Quantification reveals a significant reduction in neuronal density in SuHx-treated rats (n=6) relative to vehicle controls (n=6), with a mean difference of $-0.0001113 \pm 2.937 \times 10^{-5}$ (p = 0.0035) (**5C**). Linear regression analysis indicates that right ventricular pressure significantly predicts neuronal density, accounting for 58% of the variance (R^2 = 0.5825, p = 0.0039). (**5D**). SuHx-treated animals also exhibit significantly increased neuronal TDP-43 nuclear exclusion compared to controls, with a mean difference of 0.2592 ± 0.04323 (p < 0.0001) (**5E**). Data are presented as mean ± SEM and analyzed using unpaired two-tailed t-tests unless otherwise specified.

Figure 6. Clinical pulmonary hypertension is associated with increased neuronal TDP-43 mislocalization. Representative immunohistochemistry of frontal cortex tissue from patients with pulmonary hypertension (PH) and age-matched control subjects was used to assess neuronal density and TDP-43 localization (**6A**). Quantitative analysis revealed no significant difference in cortical neuronal density between PH and control groups (**6B**). However, PH subjects (n = 4) demonstrated significantly increased neurons with extranuclear TDP-43compared to controls (n = 3), with a mean difference of 46.27 ± 12.96 (p = 0.0230; **6C**). Scale bar = 50 μm. Data are presented as mean ± SEM and analyzed using a two-tailed unpaired t-test.

Antibody	Dilution	Manufacturer	Product ID
Ionized calcium-binding adaptor molecule 1 (Iba1)	1:500	Synaptic Systems	#234009
Glial Fibrillary Acidic Protein (GFAP)	1:1000	Invitrogen	#13-0300
TDP-43	1:500	Proteintech	#10782-2
Neuron-Specific Nuclear Protein (NeuN)	1:1000	Antibodies.com	#A270544

Supplemental Table 1. Antibodies, dilution, manufacturer, and product ID utilized in immunohistochemistry experiments.

Age	Sex	PH Etiology	PH Severity (RVP)	Comorbidities
69	F	Control	UA	UA
27	F	Control	UA	UA
64	F	Control	UA	UA
83	M	CTEPH	UA	UA
69	M	PoPH	43 mmHg	ESRD, NASH cirrhosis
47	M	PoPH	52 mmHg	AUD, cirrhosis, DM2, hepatic encephalopathy
43	F	SLE-PH	93 mmHg	stroke, ITP, DVT

Supplemental Table 2. Pulmonary hypertension subject demographics. Abbreviations include; unavailable (UA), Chronic Thromboembolic Pulmonary Hypertension (CTEPH), Portal pulmonary hypertension (PoPH), end stage renal disease (ESRD), non-alcoholic steatohepatitis (NASH), alcohol use disorder (AUD), diabetes mellitis 2 (DM2), autoimmune thrombocytopenic purpura (ITP), deep venous thrombosis (DVT).

7A

7B

Gene	Name	Function
RFLNA	Refilin A	Actin filament organization
EIF4A1	Eukaryotic Translation Initiation Factor 4A1	RNA polymerase II regulation
MIF	Macrophage Migration Inhibitory Factor	Inflammation
PIGY	Phosphatidylinositol Glycan Anchor Bio-synthesis Class Y	Biosynthesis of glycosylphosphatidylinositol (GPI)
RAB4B	Ras-Related GTP-Binding Protein 4b	Involved in glucose import
HNRNPH2	Heterogeneous Nuclear Ribonucleoprotein H2	Pre-mRNA processing
MOB4	MOB Family Member 4	Oocyte maturation
SULT1A4	Sulfotransferase Family 1A Member 4	Sulfate conjugation
SNX15	Sorting Nexin 15	Intracellular traficking
NBL1	DAN Family BMP Antagonist	BMP antagonist
MRPS24	Mitochondrial Ribosomal Protein S24	Protein synthesis
PLSCR3	Phospholipid Scramblase 3	Apoptosis
CHMP3	Charged Multivesicular Body Protein 3	Sort proteins into multivesicular body pathway

Figure 7. Differential expression of TDP-43 target genes in pulmonary artery endothelial cells from PH dataset. Volcano plot showing differential gene expression from single-cell RNA sequencing of pulmonary artery endothelial cells (PAECs) in PH (n=20) vs control (n=3) (GSE243193). Fisher exact test confirmed that TDP-43–regulated genes were disproportionately downregulated in PH (p < 2.2 × 10^{-16}, OR: 0.193, 95% CI: 0.139-0.269) (**7A**). Manual annotation of top dyresgulated genes provided (**7B**).

Supplemental Figure 1. Monocrotaline induces vascular remodeling.
Rats treated with monocrotaline (MCT; n = 6) trend to an increased right ventricular pressure (RVP), with a mean difference of 17.33 ± 6.98 mmHg compared to vehicle controls (n = 6; p = 0.0522) (**S1A**). MCT-treated rats demonstrate right ventricular hypertrophy, with a mean Fulton Index difference of 0.09815 ± 0.03066 compared to controls (p = 0.0198) (**S1B**). Data analyzed using a two-tailed unpaired t-test and are presented as mean ± SEM. Female animals depicted in blue.

Supplemental Figure 2. Monocrotaline does not induce sickness behavior in rats undergoing open field test. Rats exposed to Monocrotaline (MCT, n = 10) did not exhibit significant differences in behavioral parameters compared to vehicle-treated controls (n = 10). Specifically, there were no significant changes in total distance traveled (**S2A**), time spent in the center of the open field test (**S2B**), time spent at rest (**S2C**), or average ambulatory velocity (**S2D**). Data were analyzed using an unpaired two-tailed t-test and are presented as mean ± SEM.

Supplemental Figure 3. Monocrotaline increases cortical gliosis. Representative immunofluorescence images of frontal cortex microglia (IBA1/DAPI) (**S3A**) and astrocytes (GFAP/DAPI) (**S3B**) in rats exposed to monocrotaline (MCT) or vehicle. MCT-treated rats (n = 6) exhibited a significantly increased microglial density (0.2320 ± 0.009541) compared to vehicle-treated controls (n = 6; p <0.0001). (**S3C**). Similarly, astrocyte density was elevated in MCT-treated animals (4.822 x 10^{-5} ± 1.309 x 10^{-5}) relative to controls (p = 0.0043) (**S3D**). Data are presented as mean ± SEM and were analyzed using a two-tailed unpaired t-test. Error bars are shown for all groups; however, the control group's error bars are not visible at this scale due to extremely low variance (mean = 0.0001573, SEM = 0.0000048) and large difference from the MCT group. A text label is included to indicate the control mean ± SEM.

Supplemental Figure 4. Monocrotaline does not increase cortical cytokine expression. Quantitative PCR analysis of cortical cytokine mRNA expression in rats treated with monocrotaline (MCT; n = 6) or vehicle (n = 6) revealed no significant differences in IL-1β (**S4A**), TNF-α (**S4B**), or IL-6 (**S4C**) transcript levels. Data are presented as mean ± SEM and were analyzed using a two-tailed unpaired t-test.

Supplemental Figure 5. Monocrotaline treatment reduces cortical neuronal density and induces neuronal TDP-43 mislocalization. Representative immunofluorescence images of cortical neurons stained for NeuN (neuronal marker) and DAPI (nuclear marker), TDP-43 and DAPI, and merged channels in rats exposed to vehicle at 40x magnification (**S5A**) compared to monocrotaline-treated rats (MCT) (**S5B**). Quantification reveals no change in cortical neuronal density in MCT-treated rats (n=6) compared to vehicle controls (n=6) (**S5C**).   MCT-treated animals exhibit significantly increased neuronal nuclear TDP-43 exclusion compared to controls, with a mean difference of 0.2553 ± 0.07623 (p = 0.0194) (**S5D**). Data are presented as mean ± SEM and analyzed using unpaired two-tailed t-test.

482