Title Page

A short informative title

Rank-biased access to females in periovulatory periods but comparable copulation rates across

ranks in male bonobos at Wamba

A short running title

Rank-biased access to females in periovulatory periods but comparable copulation rates across

ranks in male bonobos at Wamba

Author name

Kazuya Toda[1], Furuichi Takeshi[2]

Affiliation

[1]Research Center for Integrative Evolutionary Science, The Graduate University for Advanced

Studies, Sokendai, Japan

[2]Wildlife Research Center, Kyoto University, Inuyama, Aichi, Japan

Email address

[1]toda.kazuya.h17@kyoto-u.jp

[2]furuichit@gmail.com

Ethical statements

Our field studies of free-ranging bonobos at Wamba in the Luo Scientific Reserve were approved

by the Ministry of Scientific Research and the Center for Research on Ecology and Forestry in

the Democratic Republic of Congo. This study adhered to the Animal Research Guidelines of The

Graduate University for Advanced Studies (SOKENDAI), Japan, revised on March 23, 2021.

Keywords

Pan paniscus, prolonged sexual receptivity, non-conceptive copulation, intermale mating

competition, operational sex ratio

Abstract

Objectives

Bonobos (*Pan paniscus*) are unique among non-human primates in that females exhibit prolonged

sexual receptivity to males during the postpartum infertile stage. Although high synchrony in

female sexual receptivity has been hypothesized to constrain dominant males' monopolization of

reproductive success, recent genetic studies indicate pronounced male reproductive skew in this

species. We investigated how prolonged sexual receptivity in female bonobos affects male

behavioral patterns.

Materials and Methods

Behavioral observations were conducted on a free-ranging bonobo group at Wamba in the Luo

Scientific Reserve, Democratic Republic of the Congo. Periovulatory periods (POPs) were

estimated from the timing of detumescence marking the end of the maximal swelling phase,

following Ryu et al. (2022). For each male, we calculated the proportion of copulations with POP

females. Using all-day focal follows targeting mature males, we tested for rank-related bias in

their copulation rates and evaluated the effects of the party-level operational sex ratio (OSR; the

ratio of mature males to sexually receptive females) on the focal male's probability of copulation

and intermale aggression.

Results

Higher-ranking males were more likely to copulate with POP females, yet males of all ranks

engaged in copulation on most of focal observation days (twice per day, on average). Lower party-

level OSR values increased the focal male's probability of copulations while reducing intermale

aggression.

Discussion

Prolonged sexual receptivity in female bonobos—yielding a pool of sexually receptive but non-

POP females—may broaden male copulation opportunities beyond dominance rank and reduces

intermale mating competition, even when POPs are detectable.

1. Introduction

Sexual receptivity refers to the physiological, morphological, and behavioral readiness of a female

to accept a male and engage in copulation (Dixson 1998). In most mammals, estrus is the recurring

phase of sexual receptivity and typically coincides with the periovulatory period (POP) (Conaway

1971; Weir and Rowlands 1973). Estrous signals—visual, olfactory, or acoustic—advertise

fertility, attract males, and facilitate conception (Johns 1980; Semple and McComb 2000; Moreira

et al. 2015). However, in some species—particularly among simian primates—female sexual

receptivity extends beyond the fertile window (Beach 1976; Small 1988; Rooker and Gavrilets

2020). In species with promiscuous mating systems, copulating with multiple males can obscure

paternity and thereby reduce the risk of male-inflicted infanticide (e.g., bottlenose dolphins

[*Tursiops* spp.]: Connor et al. 1996; Hanuman langurs [*Semnopithecus entellus*]: Heistermann et

al. 2001; capuchin monkeys [*Cebus capucinus*]: Carnegie et al. 2005). Humans exhibit the most prolongation of female sexual receptivity among mammals (Symons 1979; Hrdy 1981), and ovulatory cues are sufficiently subtle that the POP is difficult for males to detect (Haselton and Gildersleeve 2011). A prominent hypothesis concerning concealed ovulation in humans posits that an obscured fertile window encourages males to maintain pair bonds rather than pursue multiple partners, thereby increasing paternity certainty and eliciting paternal investment in females and their offspring (Alexander and Noonan 1979; Symons 1979; Rodriguez-Girones and Enquist 2001). Thus, the evolution of prolonged female sexual receptivity may have helped shape a social environment advantageous to females.

Female chimpanzees (*Pan troglodytes*) and bonobos (*Pan paniscus*) are nonseasonal breeders with interbirth intervals of approximately 4–5 years (Thompson and Sabbi 2024). Across ovulatory cycles, females exhibit conspicuous changes in the size, shape, and firmness of the sexual skin around the anogenital region, reflecting fluctuations in estrogen and progesterone levels (Graham et al. 1972; Heistermann et al. 1996; Hashimoto 2023). Sexual skin tumescence ("sexual swelling") functions as an estrous signal indicating sexual receptivity (Furuichi 1987; Nishida et al. 1990). Females of both species continue to show several cycles of sexual swelling after conception and continue to copulate during pregnancy (Tutin and McGinnis 1981; Furuichi 1987; Takahata, Ihobe and Idani 1996; Wallis and Goodall 1993; Douglas et al. 2016). Moreover, in each ovulatory cycle, sexual swellings often persist for more than 10 days (Hasegawa and Hiraiwa-Hasegawa 1983; Wallis 1985; Furuichi 1987; Deschner et al. 2003; Paoli et al. 2009; Douglas et al. 2016), exceeding the fertile window, given that sperm remain viable for up to three days in the female reproductive tract whereas eggs are viable for only ~24 hours postovulation (Wilcox, Weinberg, and Baird 1995; Dunson, Baird, and Wilcox 1999). Prolonged sexual receptivity outside the fertile window in the genus of *Pan* is therefore thought to promote paternity

confusion and reduce the risk of male-inflicted infanticide (Zinner and Deschner 2000; Furuichi

and Hashimoto 2002; Wrangham 2002).

The two *Pan* species differ markedly in the timing of postpartum resumption of female

sexual receptivity. In bonobos, females resume sexual swellings at a mean of 8.1 months

postpartum while still nursing their infants (Hashimoto et al. 2022). By contrast, female eastern

chimpanzees typically do not resume sexual swellings until after weaning, at approximately four

years of offspring age (Gombe: 46.8 months [Wallis 1997]; Mahale: 55.2 months [Nishida et al.

2003]; Kanyawara: 40.8 months [Thompson and Ellison 2017]), and conception often follows

within a year of sexual receptivity resumption (Nishida et al. 1990; Wallis 1997; Furuichi and

Hashimoto 2002; Thompson 2013). Female western chimpanzees resume earlier (Tai: 21.6 and

28.8 months postpartum [Deschner and Boesch 2007]), yet female bonobos still resume

13 months earlier on average. In bonobos, the period during which sexual swellings occur prior

to conception spans roughly 3.5 years, calculated by subtracting postpartum amenorrhea (8.1

months) and gestation (7.9 months) from the mean interbirth interval of 4.8 years for surviving

offspring (Furuichi et al. 1998). The physiological basis of these interspecific differences remains

poorly understood; one possibility is that the sexual skin of female bonobos is more sensitive to

estrogen than that of female chimpanzees, facilitating earlier resumption of sexual receptivity

(Hashimoto et al. 2022).

High synchrony in female bonobos' sexual receptivity has been hypothesized to hinder

dominant males' monopolization of reproductive opportunities and thereby reduce intermale

mating competition (Furuichi 2006). The operational sex ratio (OSR)—the ratio of mature males

to sexually receptive females present at a given time—is widely used as an indicator of the

intensity of intermale contest over mates (Kvarnemo and Ahnesjo 1996; Weir, Grant, and

Hutchings 2011). In species characterized by multi-male/multi-female groups, long interbirth

intervals, and nonseasonal breeders, such as hominid apes (Thompson and Sabbi 2024), the OSR

is typically high. For example, the OSRs in chimpanzee groups are as high as 4.2 at Mahale and

12.3 at Gombe (reviewed by Furuichi and Hashimoto 2002). By contrast, the OSR in a bonobo

group at Wamba is 2.8, substantially lower than in chimpanzees, reflecting a relatively higher

proportion of sexually receptive females (Furuichi and Hashimoto 2002). Moreover, the mean

number of females per party—a flexible subgroup whose membership varies in size and

composition over time (Nishida 1968; Kuroda 1979)—is higher in bonobos at Wamba than in

eastern chimpanzees at Kalinzu (Mulavwa et al. 2008), whereas it is similar between bonobos at

LuiKotale and western chimpanzees at Taï (Surbeck et al. 2021). High rates of interfemale

associations in bonobos may therefore further reduce intermale mating competition (Furuichi

2023).

Several studies indicate that intermale mating competition appears both less frequent and

less intense in bonobos than in chimpanzees. In eastern chimpanzees, a substantial share of male

aggression occurs in mating contexts: 17/100 events (17.0%) at Gombe and 36/100 events

(36.0%) at Kanyawara (Goodall 1986; Muller 2002). By contrast, in bonobos at Wamba, the

proportion of male aggression during copulations or approaches to sexually receptive females is

lower: 2/36 (5.6%), 7/73 (9.6%), and 5/83 events (6.0%) (Furuichi 1989; Shibata and Furuichi

2024; this study). Higher proportions reported for bonobos at Lomako (37/150 events [24.7%],

Hohmann and Fruth 2003) and LuiKotale (95 of 431 events [22.0%], Surbeck et al. 2011) likely

reflect definitional differences from those studies at Wamba: Hohmann and Fruth (2003) analyzed

only contact aggression, which may be disproportionately expressed during intermale mating

competition, and Surbeck et al. (2011) broadly classified all antagonistic interactions occurring

within 15 minutes before or after copulation as aggression in mating contexts. Additionally, severe

intermale aggression—particularly lethal aggression both within and between groups—

characterizes chimpanzees and has not been documented in bonobos (Furuichi 2011; Wilson et al.

2014; Pusey 2022; Shibata and Furuichi 2024). Skeletal evidence from museum specimens

likewise indicates higher rates of trauma in chimpanzees than in bonobos (Jurmain 1997).

However, genetic paternity analyses from multiple bonobo groups at LuiKotale, Wamba,

and Kokolopori consistently show pronounced male reproductive skew (Surbeck et al. 2017;

Ishizuka et al. 2018; Mouginot et al. 2023). Across groups, the most successful males sired a

larger share of offspring than in chimpanzees (Mouginot et al. 2023: 69 ± 11% in bonobos; 44 ±

25% in chimpanzees). Male reproductive success also covaries with dominance rank; alpha males

in two study groups at Wamba achieved the highest reproductive success (Ishizuka et al. 2018),

and alphas in two of three study groups at Kokolopori sired more offspring than expected

(Mouginot et al. 2023). These genetic outcomes suggest that male bonobos can discriminate POP

from non-POP females with some accuracy, even when both are scored by observers as maximally

swollen. Consistent with this possibility, Ryu et al. (2022) have shown that males' intensive

following and mate guarding are concentrated on POP females, identified via urinary hormone

profiles. Male chimpanzees likewise appear able to detect the POPs (Goodall 1986; Wallis 1992;

Deschner et al. 2003; Thompson 2005), although the specific cues used remain untested in either

Pan species. In addition to the power of dominant males to exclude male competitors (Kano 1996;

Hohmann and Fruth 2003; Surbeck et al. 2012; Mouginot et al. 2024), the ability to detect POP

females despite high synchrony in female sexual receptivity may help explain the marked male

reproductive skew.

Despite pronounced inequality in male reproductive success in bonobos, why is

intermale mating competition not expressed in more severe forms of aggression? One

possibility is that the presence of multiple sexually receptive females dilutes competition by

distributing male copulation attempts. Additionally, if a broad range of males can obtain

copulation opportunities—even though most are nonconceptive—the proximate motivation

to compete over POP females may be attenuated. To evaluate the effects of prolonged sexual

receptivity in female bonobos on male behaviors, we conducted all-day focal observations

of males in a free-ranging bonobo group at Wamba. Specifically, this study addressed the

following three questions:

(1) Do dominant males have priority access to POP females? Given the pronounced male

reproductive skew, higher-ranking males are expected to copulate more frequently with

POP females. To test this prediction, we estimated POPs from daily assessments of each

female's sexual swelling status and, for each male, calculated the proportions of his

copulations that involved POP females.

(2) Do males of different ranks have comparable opportunities to copulate? If sexually

receptive but non-POP females attract males and provide many males with copulation

opportunities, copulation rates should differ little across dominance ranks. To test this

prediction, we examined a correlation of dominance ranks with copulation rates derived

from all-day focal follows, which reduce sampling bias relative to short focal or ad

libitum sampling.

(3) Do within-party changes in OSR influence intermale mating competition? If male

sexual interest becomes more dispersed at the number of sexually receptive females

increases, then lower OSR values should allow males to copulate more easily without

contest. To test this prediction, we used continuous records of party composition during

all-day focal follows and analyzed how the numbers of mature males and sexually

receptive females, and the presence of POP females affected the focal male's probability

of copulation and his involvement in intermale aggression.

2. Materials and Methods

2.1 Study site and subjects

Behavioral and ecological studies of free-ranging bonobos at Wamba, in the Luo Scientific

Reserve, Democratic Republic of the Congo, began in 1974 (Kano 1992). By 1976, all individuals

in the main study group, "E," were identified (Kuroda 1979). Subsequent work showed that the

E group actually comprised two independent groups; from 1984 onward tjere were designated

"E1" and "E2" (Furuichi 1987; Furuichi and Ihobe 1994). Demographic data—births, deaths,

immigration, and emigration—have since been collected primarily for E1. Fieldwork at Wamba

was interrupted in 1991–1994 owing to political instability and again in 1996–2002 during the

Congo Wars (Furuichi et al. 2012). Artificial provisioning was initially used to facilitate detailed

behavioral observations but was discontinued in 1996.

For this study, we collected data on the E1 group from December 1, 2022 to May 2, 2023

(153 days). During this period, E1 ranged from 45 to 48 individuals: 7 adult males (≥15 years), 5

adolescent males (8–14 years old), 16 adult females (≥15 years or older or parous), and 4

adolescent females (8–14 years and nulliparous). Following the resumption of fieldwork in 2004,

all E1 individuals were reidentified by 2007; at that time, the group comprised of 24 individuals

(10 adult males, 6 adult females, and 2 adolescent females; Hashimoto et al. 2008). E1's size

increased gradually after 2009 (Fig. S1), largely due to the immigration of nulliparous females

who subsequently reproduced (Toda et al. 2023).

2.2 Behavioral observations

Trained local assistants followed one party of the E1 group continuously from the time the

bonobos descended from their night beds (approximately 0600 h) until they constructed new night

beds (approximately 1700 h). The first author (KT) and a local assistant, Lokemba Batsindelia

(LB), conducted focal animal sampling (Altmann 1974) throughout the day to collect continuous

behavioral data from 6 adult males, 5 adolescent males, 15 adult females, and 4 adolescent

females in E1. One adult male (TN; estimated to be 55 years) was excluded from focal sampling

because he had been missing from E1 for five months and was absent at the start of this study.

Another adult male (TW; estimated to be 51 years) was included in focal follows, but his data

were excluded from analyses because he has not been observed copulating or exhibiting erections

since at least 2015 (Toda, personal observation).

Each day, we selected a focal subject from party members using a predetermined random

order. A focal follow continued until one of the following conditions was met: the focal subject

constructed a new night bed; darkness near sunset (approximately 1700 h) precluded

identification ; or the subject was out of sight for ≥2 consecutive hours. Focal follows with ≥4

hours of observation were included in our dataset for analyses (i.e., treated as all-day focal data).If

a focal subject was lost for ≥2 consecutive hours before reaching 4 hours, a second focal follow

was initiated on the same day only if it could start by 1200 h. If the predetermined focal subject

could not be located within 30 minutes of the start of the party follow, we selected the next

available individual in the predetermined order from among party members present at that time.

From the midpoint of the study onward, we prioritized undersampled subjects to balance

observation effort. To ensure independence among all-day focal follows, at least 20 days elapsed

before resampling the same individual.

During each focal follow, we recorded behaviors, including resting (e.g., sitting, lying, self-

grooming), feeding (e.g., taking, chewing, swallowing food), moving (e.g., walking, running,

climbing, jumping, brachiating), and social interactions (e.g., playing, social grooming, sexual

interactions, agonistic interactions), along with the duration of each activity and the identities of

participants. We also noted party composition—defined as the set of adult and adolescent

individuals present during focal follows—at 1-hour intervals from 0600 h to 1700 h using the 1-

hour party method (Hashimoto, Furuichi, and Tashiro 2001). In addition, all observed copulatory

and antagonistic interactions, including events not involving the focal subject, were recorded ad

libitum during focal and party follows.

In total, 349.6 hours of observation were collected from ten males (excluding TN and TW)

over 41 days of all-day focal follows conducted by KT (193.2 hours) and LB (156.4 hours). Mean

focal time per day was 8.5 hours (SD = 1.8; range = 4.2–11.0 hours), with a median start time of

0630 h (range = 0600–1142 h) and a median end time of 1700 h (range = 1315–1732 h). The

median focal time per subject was 35.4 hours (range = 25.1–44.7 hours; Table 1). Party

composition was recorded for 422 sessions (not included if the focal male was out of sight for the

entire hour. Of these sessions, 413 in which the focal male was visible for more than 15 minutes

were retained for analyses.

2.3 Definition of copulation and aggression

Copulation refers to heterosexual interactions in which males mounted females, involving the

insertion of the penis into the vagina, accompanied by thrusting movements. Because ejaculation

was often not visually discernible, such intercourse was recorded as copulation regardless of

whether ejaculation could be confirmed (Hashimoto 1997; Ryu et al. 2022). Aggression refers to

directed antagonistic interactions, including physical contacts (e.g., punching, kicking, pushing,

tackling, biting) and threat displays (e.g., chasing, charging, charging while branch-dragging,

branch-shaking) (Nishida et al. 1999). Undirected displays lacking a clear target were not

classified as aggression. We also recorded the target's responses to aggression, including

submissive behaviors (e.g., fleeing, grinning, screaming) and retaliations toward the aggressor.

To avoid inflating event counts, successive copulations or aggressions involving the same dyad

within 10 minutes were treated as a single event.

2.4 Evaluation of male dominance ranks

To determine the dominance hierarchy of the E1 group, we analyzed 297 aggression events: 124

recorded during all-day focal follows of males, 51 during all-day focal follows of females, and

122 collected ad libitum. In total, 201/297 events (67.7%) elicited submissive responses from the

target, allowing us to identify winners and losers. We then computed David's scores (David 1987)

for all adults and adolescents from the observed interaction matrix, using dyadic dominance

indices that incorporate win-lose proportions and correct for unequal opportunities (de Vries,

Stevens, and Vervaecke 2006). Although our primary aim was to evaluate male ranks, both sexes

were included because male bonobos are co-dominant with females (Furuichi 1997; Surbeck and

Hohmann 2013). Male dominance rank derived from David's score (Table 1) was not significantly

associated with age (Spearman's rank correlation test: $\rho = -0.26$, $S = 360$, $p = 0.417$).

2.5 Estimation of female fertile windows

Daily presence of each individual in E1 has been recorded since September 2009 (Sakamaki et al.

2015). In addition, adult and adolescent females' sexual swellings were scored daily on a scale of

1 to 3 based on tumescence (Furuichi 1987; Ryu et al. 2015; Ryu et al. 2022): 1 = no swelling, 2

= intermediate swelling, 3 = maximal swelling.

Douglas et al. (2016) reported that the maximal swelling phase (MSP) in female bonobos

at Luikotale is highly variable within and among individuals (mean ± SD = 10.6 ± 6.8 days; range

= 1–31 days) and that ovulation timing inferred from urinary hormones varies substantially with

respect to both MSP onset and end. By contrast, at Wamba, Ryu et al. (2022) found that ovulation

timing peaks four days before MSP end—13 of 14 ovulations occurred toward MSP end (see also

Hashimoto et al. 2022)—although it remains unpredictable from MSP alone. In primates, the

fertile window is often defined as the day of ovulation and the three preceding days (Deschner et

al. 2003; Douglas et al. 2016), consistent with sperm survival up to ~3 days in the female tract

and eggs remaining viable for ~24 hours post-ovulation (Wilcox, Weinberg, and Baird 1995;

Dunson, Baird, and Wilcox 1999), Applying this criterion, Ryu et al. (2022) estimated that the

probability of sperm coinciding with eggs (i.e. fertility) exceeds 0.3 from 7 to 2 days before

detumescence (i.e., MSP end) and falls to ~0 on the day before prior to detumescence.

Although ovulation does not occur during postpartum amenorrhea, female bonobos

nevertheless exhibit sexual swellings. To distinguish POP from non-POP females, we

operationally defined the POP as the 6-day window from 7 to 2 days before MSP end, following

Ryu et al. (2022). Gaps in swelling score records arose when females were absent from party

follows or observations were missed. If a female's swelling score was identical immediately

before and after a gap of ≤3 days, we interpolated the same score across the gap. When MSP end

was obscured by a gap, POP status was treated as indeterminate. If swelling scores dropped from

3 to 2 for a single day and then returned to score 3, the transient score-2 day was included in the

MSP. If an MSP lasted <6 days, no POP was assigned for that MSP.

2.6 Data analyses

Statistical analyses were conducted in R version 4.4.2 (R Core Team 2024) using RStudio version

12.0.467 (Posit Team 2024). We fit generalized linear models (GLMs) for non-Gaussian responses

with the 'glm' function in the 'stats' package (R Core Team 2024) and generalized linear mixed

models (GLMMs) to account for repeated measures within individuals and across observational

days (Bolker et al. 2009) with the 'glmer' function in the 'lme4' package (Bates et al. 2015).

Because covariate-response relationships can be nonlinear, we included quadratic terms for all

covariates. To reduce collinearity between linear and quadratic terms of the same predictor, all

covariates were z-transformed (mean = 0, SD = 1) prior to squaring (Schielzeth 2010). The

variables included in each model are described below.

To test the first prediction—that higher-ranking males copulate more frequently with POP

females—we evaluated the effect of male dominance rank on the proportion of copulations with

POP females versus non-POP females. For each male, we aggregated counts of copulations with

POP females and with non-POP females across all events recorded via focal and ad libitum

sampling during the study period. These two-column responses (POP, non-POP), constructed with

the 'cbind' function, were fit with a binomial GLM using a logit link (Zuur et al. 2009). Male rank

(1st–10th) and age (9–48 years) were included as covariates.

To test the second prediction—that male copulation rates differ little across dominance

ranks—we examined the rank effect on copulation rates. Because male chimpanzees, in particular

dominant males, show greater sexual interest in adult than adolescent females (Muller, Thompson,

and Wrangham 2006), we tallied, for each focal day, the focal male's copulations with adult and

adolescent females separately. These daily counts were fit with two Poisson GLMMs using a log

link, each including an offset for observation time to account for variation among focal follows

(Zuur et al. 2009). Male rank and age were included as covariates in both models. In addition, the

number of non-mother females with maximal swellings (hereafter termed as MS females) on each

focal day (adults: 0–6; adolescents: 0–4) was included as a covariate to capture day-to-day

variation in potential copulation opportunities. A random intercept for focal male identity was

included in both models to account for multiple observations from the same individual.

To test the third prediction—that lower OSR should allow males to copulate more easily

without conflict—we assessed how temporal variation in OSR and the presence of POP females

affected the focal male's probability of copulation and intermale aggression. For each 1-hour focal

session from 0600 to 1700 h (11 sessions per day), we coded whether ≥ 1 copulation and ≥ 1

intermale aggression involving the focal male occurred (1) or not (0). When an interaction

spanned consecutive sessions, it was assigned to the earlier session based on its start time. These

binary responses were fit with two binomial GLMMs using a logit link. Fixed effects included

the number of mature males except for TW (1–10 individuals), the number of MS females (0–8

individuals), and the presence of non-mother POP females (yes/no) in the party during each

session. We also included the duration the focal male spent resting during the session (0–60

minutes) to control for variation in inactivity and the duration he was out of sight (0–45 minutes)

to control for variation in effective observational time. Random intercepts for focal male identity,

observational date, and hour of day (i.e., session) were included in both models.

We fit GLMs and GLMMs by maximum likelihood (Zuur et al. 2009) and used the Akaike's

Information Criterion (AIC) for model selection (Burnham, Anderson, and Huyvaert 2011). The

model with the lowest AIC among all subsets of fixed effects was taken as the best model, and

we also examined models with ΔAIC < 2 relative to the best model (Burnham and Anderson 2004).

For each best model, we evaluated residual normality, dispersion, and outliers using the

'simulateResiduals' function in the 'DHARMa' package (Hartig 2024); none showed violations

(Fig. S2–S6). We assessed residual independence with the 'check_autocorrelation' function in the

'performance' package (Lüdecke et al. 2021) and found no significant autocorrelation (minimum

p-value = 0.176). Multicollinearity was evaluated with the 'check_collinearity' function in the

'performance' package (Lüdecke et al. 2021); variance inflation factors (VIFs) were low in all

best models (maximum VIF = 3.07).

3. Results

3.1 Frequency and types of male copulation and intermale aggression

Overall, we observed 67 copulations involving the focal male during 41 all-day focal follows, averaging 0.19 events per observation hour. Individual mean copulation rates ranged from 0.06 to 0.30 events per hour. On 38/41 focal days (92.7%), at least one MS female was in the same party as the focal male; on 33/38 of those days (86.8%), focal males copulated at least once. Including 206 copulations recorded ad libitum, a total of 273 copulations were documented over the study period: 64 (23.4%) with POP females, 186 (68.1%) with non-POP females, and 23 (8.4%) with females of undetermined fertility status. Visible sperm expulsion from the vagina immediately after copulation was confirmed in 17 events (6.2%); eight with POP females, six with non-POP females, and three with females of undetermined status. Ejaculation was more frequently observed in copulations with POP females (8/64 = 12.5%) than overall (17/273 = 6.2%; exact binomial test: $p < 0.001$).

We also recorded 83 intermale aggressions involving the focal male during all-day focal follows, corresponding to a mean of 0.24 events per observation hour. Individual mean aggression rates ranged from 0.08 to 0.42 events per observation hour. The dyad most frequently involved was the maternal brothers, KT and KY (21.7% of all intermale aggressions). Intermale aggression most often occurred when targets were resting on the ground (24.1%), followed by when targets were feeding near aggressors (18.1%). Only five aggressions (6.0%) occurred in mating contexts—when targets approached or solicited MS females, or during/immediately after copulation—and three of these (60.0%) involved contact aggression. Given that the overall proportion of contact aggression in 15/83 (18.1%), contact aggression was more likely to occur in mating contexts than overall (exact binomial test: $p = 0.044$).

3.2 Variation in the frequency of female maximal swellings and the timing of periovulatory periods

(POPs) after parturition

Based on daily presence records, the mean numbers of MS females per observation day were 2.8

adults (range = 0–6, N = 16) and 1.7 adolescents (range = 0–4, N = 4). Across individuals, the

percentage of observational days on which each female exhibited maximal swellings ranged from

0 to 73.2%. Excluding three pregnant females (Ik, Sc, and Ec) who gave birth during the study

period, this percentage was significantly higher in adolescents (56.1 ± 13.5 SD %, N = 4) than in

adults (28.7 ± 18.6 SD %, N = 13) (Wilcoxon rank-sum test: W = 46, p = 0.023).

We documented 30 MSPs that lasted >6 days. Of these, 21 POPs were identified—based

on MPS end—for 10 females (8 adults, 2 adolescents; Table S1). Copulation with mature males

occurred at least once in 18/21 POPs (85.7%). Among adults, the 15 POPs accompanied by

copulation occurred at 22–91 months postpartum; no female lost her most recent offspring during

the study period.

3.3 Effect of male dominance rank on access to females in periovulatory periods (POPs)

Figure 1 shows, for each male, the proportion of his copulations that involved POP females. The

alpha male, KT, copulated with POP females in 58.1% of his copulations (18/31 events). His

mother, Ki, was the alpha female (Table S1). His maternal brothers, KY and NB—ranked second

and fourth, respectively—also showed higher proportions (42.9% and 50.0%) than the across-

male mean of 23.2% (range = 4.3–58.1%, N = 10). We observed Ki intervene in her sons' agonistic

interactions four times; on one occasion, she intercepted another male's copulation attempt.

Among males whose mothers were not top-ranking, the third-ranking male, JO, copulated with

POP females at a higher proportion (22.2%) than the four lower-ranking males, HC, SE, YD, and

NI (range = 4.3–15.8%). Two males without resident mothers, DI and GC, each had only one

copulation with POP females (5.6% and 6.7%, respectively).

The best-fitting binomial GLM for the binary response (POP vs. non-POP) retained male

rank (linear term only) as the sole predictor (Table 2A). Male rank had a significant positive effect

on the odds that a copulation involved a POP female (estimate = 0.97, SE = 0.19, z = 5.06, p <

0.001), indicating that higher-ranking males copulated more frequently with POP females than

lower-ranking males.

3.4 Relationship of male dominance ranks with copulation opportunities

Figure 2 shows, for each male, the rate of copulations with adult females per observation hour.

The best-fitting Poisson GLMM for counts of copulations with adult females included the number

of non-mother adult MS females (linear and quadratic terms; Table 2B). The linear term was

positive (estimate = 0.94, SE = 0.28, z = 3.34, p < 0.001), whereas the quadratic term was negative

(estimate = –0.47, SE = 0.23, z = 2.03, p = 0.043), indicating that copulation rates increased with

the number of non-mother adult MS females but then peaked out (inverted U-shaped pattern). A

competing model that included male rank (linear term only) had a ΔAIC = 2.0, and the rank effect

was not significant (estimate = –0.03, SE = 0.16, z = –0.17, p = 0.867; Table S2A).

Rates of copulations with adolescent females per observation hour are also showed for each

male in Figure 2. The best-fitting Poisson GLMM for counts of copulations with adolescent

females was the null model with random intercept only (Table 2C), indicating that none of the

candidate predictors improved model fit. A model including male rank (linear term only) had

ΔAIC = 1.8 and showed a non-significant rank effect (estimate = 0.10, SE = 0.23, Z = 0.45, p =

0.657; Table S2B).

3.5 Effects of within-party operational sex ratio dynamics and the presence of females in

 periovulatory periods (POPs) on intermale mating competition

Across 1-hour sessions in which the focal male was visible for >15 minutes, the mean (± SD)

numbers of party members present were 3.0 ± 1.7 adult males, 2.2 ± 1.3 adolescent females, 6.4

± 3.6 adult females, and 1.5 ± 1.2 adolescent females. Sessions included, on average, 2.9 non-

mother MS females (range = 0–9) and 0.6 non-mother POP females (range = 0–4). At least one

non-mother POP female was present in 156/413 sessions (37.8%). Copulations and intermale

aggressions were recorded in 60 and 68 sessions, respectively (14.5% and 16.4%). The focal male

spent, on average, 23.1 minutes resting and 8/0 minutes out of sight per session.

Figure 3 shows the proportion of sessions in whether the focal male copulated, plotted

against (A) the numbers of mature males and (B) the number of MS females. The best-fitting

binomial GLMM for copulation occurrence included the following predictors (Table 2D): number

of mature males (linear term only), number of non-mother MS females (linear and quadratic

terms), presence of non-mother POP females (yes/no), and resting time (linear and quadratic

terms). The number of mature males had a negative effect on the focal male's copulation

probability (estimate = –0.83, SE = 0.23, z = –3.68, p < 0.001). The number of non-mother MS

females had a positive linear effect (estimate = 1.36, SE = 0.30, z = 4.60, p < 0.001) and a negative

quadratic effect (estimate = –0.44, SE = 0.15, z = –2.92, p = 0.004), consistent with an inverted-

U relationship: copulation probability increased with more non-mother adult MS females, but

peaked before the maximum observed counts. The presence of non-mother POP females had no

significant effect (estimate = –0.53, SE = 0.35, z = –1.54, p = 0.124).

Figure 3 also shows the proportion of sessions with intermale aggression involving the the

focal male, plotted against (C) the numbers of mature males and (D) the number of MS females.

The best-fitting binomial GLMM for aggression occurrence included the following predictors

(Table 2E): number of mature males (linear term only), number of MS females (linear term only),

and resting time (linear and quadratic terms). The number of mature males had a positive effect

on the focal male's aggression probability (estimate = 1.06, SE = 0.22, z = 4.74, p < 0.001),

whereas the number of MS females had a negative effect (estimate = –0.56, SE = 0.22, z = –2.58,

p = 0.010). A competing model that included presence of POP females (yes/no) had a ΔAIC = 2.0,

and the effect was not significant (estimate = 0.04, SE = 0.38, z = 0.10, p = 0.919; Table S2C).

Together, these results indicate that the focal male was more likely to engage in copulation

and less likely to be involved in intermale aggression in parties with lower OSRs—i.e., fewer

other males relative sexually receptive females—irrespective of the presence of POP females.

4. Discussion

4.1 Validity of the empirical method for inferring female fertility status from the end of the

maximal swelling phase (MSP)

Because we conducted all-day focal follows, routine urine collection sufficient to determine

ovulation timing was not feasible. As an alternative, we applied an empirical method based on

Ryu et al. (2022) to infer periovulatory periods (POPs) of female bonobos from the end of their

MSPs. MSPs occurring before 22 months postpartum did not exceed 6 days, and the postpartum

POP timing we estimated for adult females (offspring age 22–91 months) largely overlapped the

postpartum ovulation timing identified from urinary hormone profiles (24–63 months; Hashimoto

et al. 2022). These congruences suggest that clearly non-ovulatory MSPs were unlikely to have

been misclassified as POPs. Moreover, the pattern detected with this method—higher-ranking

males copulating more often with POP females—accords with genetic evidence of pronounced

male reproductive skew in bonobos, supporting its usefulness for rough among-male comparisons

of reproductive opportunity. Nonetheless, the method does not pinpoint the exact fertile window

and should be interpreted as an approximate proxy.

4.2 Biased reproductive opportunities toward dominant males in bonobos

Consistent with our first prediction, we found that higher-ranking males were more likely to copulate with POP females. This pattern indicates that dominant males enjoy priority access to POP females, aligning with the genetic evidence for pronounced male reproductive skew in bonobos (Surbeck et al. 2017; Ishizuka et al. 2018; Mouginot et al. 2023). It also accords with Ryu et al. (2022), who showed that male bonobos can detect females' POPs to some accuracy and dominants invest greater effort in guarding POP females. In our observations, overt interference with copulations by dominant males was infrequent; nevertheless, dominants may have restricted subordinates' access to POP females, thereby limiting their reproductive opportunities.

For male bonobos, maternal presence is a key determinant of dominance rank and access to sexually receptive females (Kano 1996; Furuichi 1997; Surbeck et al. 2011; Shibata and Furuichi 2023), thereby strongly affecting male reproductive success (Surbeck et al. 2019). In this study, the three sons of the alpha female (Ki)—ranked first (KT), second (KY), and fourth (NB) among E1 males—exhibited higher proportions of copulation with POP females than males whose mothers were ranked lower. In bonobo groups characterized by strong female-female cohesion (Furuichi 1989; Parish 1996; Moscovice et al. 2017), maternal presence likely increases sons' proximity to clusters of females (including their mothers), thereby facilitating access to copulation opportunities (Surbeck et al. 2011; Yokoyama and Furuichi 2022). Moreover, because female bonobos occupy the top position in the dominance hierarchy (Furuichi 1997; Surbeck and Hohmann 2013; Tokuyama et al. 2016), dominant mothers can intervene directly in intermale mating competition on behalf of their sons (Furuichi 1997; Surbeck et al. 2011; Shibata and Furuichi 2023).

Bonobos rarely form intermale coalitions (Furuichi and Ihobe 1994; Shibata and Furuichi

2024; Mouginot et al. 2024), and cooperative relationships between maternal brothers have not

been documented. At Wamba, Ihobe (1992) noted frequent spatial proximity among maternal

brothers, but this likely reflected simultaneous association with their mothers rather than genuine

affiliative bonds between brothers. In this study, we observed a single instance of coalitionary

aggression by Ki's sons against a common target; otherwise, they did not form enduring coalitions.

Instead, KT persistently directed aggression toward his younger brother, KY, throughout the study

period and occasionally toward his older brother, NB. In this case, the maternal brothers appear

to have been reproductive competitors, much like other males. By contrast, although

documentation on brotherhood in chimpanzees is limited, Bygott (1979) described four cases at

Gombe in which males with supportive elder brothers attained alpha status, suggesting that the

presence of maternal brothers can confer a reproductive advantage in chimpanzees.

4.3 Relationship between copulation opportunities and dominance ranks in male bonobos

Despite pronounced male reproductive skew, we found that overall copulation rates did not differ

appreciably across dominance ranks. This result supports our second prediction that rank-related

bias in copulation opportunity is weak in male bonobos and is consistent with the idea that the

availability of sexually receptive females during the postpartum amenorrhea provides many males

with copulation opportunities. Although dominant males have priority access to females (Kano

1996; Ryu et al. 2022), subordinates may obtain copulation opportunities in contexts where

dominants cannot interfere or with females outside of dominants' sexual interest. Assuming that

POP detection is not absolute, subordinate males could benefit by maximizing copulation

frequency to increase their chances of participating in sperm competition, even if such efforts

rarely translate into paternity.

Our finding of comparable copulation rates across dominance ranks in male bonobos

contrasts with prior reports of higher copulation rates for dominant males (Kano 1996; Hohmann

and Fruth 2003; Surbeck et al. 2011; Ryu et al. 2022; Yokoyama and Furuichi 2022). A plausible

explanation for this discrepancy is methodological differences. Earlier studies were conducted

under artificial provisioning (Kano 1996) or relied on short-duration focal sampling (10−60

minutes) (Surbeck et al. 2011; Ryu et al. 2022; Yokoyama and Furuichi 2022). Such approaches

can bias observations toward centrally positioned individuals and larger parties, thereby missing

copulations at the periphery or in smaller parties (Altmann 1974; Drickamer 1974). Because

subordinate males tend to seek copulation opportunities while avoiding dominants (Furuichi and

Ihobe 1994; Kano 1996; Shibata and Furuichi 2023), unbiased estimates require sustained focal

follows of these males, including periods when they range alone or with only a few companions.

4.4 Impact of prolonged sexual receptivity of female bonobos on intermale mating competition

This study provides, to our knowledge, the first dataset that integrates hourly party compositions

with behavioral records from all-day focal follows to test how fluctuations in the operational sex

ratio (OSR) predict the occurrence of copulation and intermale aggression involving the focal

male. We found that a focal male's probability of copulation increased with more females

exhibiting maximal swellings (MS females) and with fewer mature males in the party, whereas

his probability of intermale aggression decreased under the same conditions. These results support

our third prediction that male bonobos obtain copulation opportunities more readily in parties

with lower OSRs—that is, fewer male competitors relative to sexually receptive females. The

presence of MS females during postpartum infertility appears to heighten male sexual motivation,

suggesting that males may experience proximate physiological and/or neurological rewards from

copulation even when conception is unlikely. In humans and rats, the post-ejaculatory state is

associated with reduced general arousal, likely mediated by prolactin release (Krüger et al. 2002;

Ågmo and Laan 2022); accordingly, frequent day-to-day access (approximately twice per day, on

average) to copulation could lessen males' incentive to engage in costly contests over POP

females. Non-invasive monitoring of hormonal activities in male bonobos would help clarify the

physiological effects of non-conceptive copulations on behavioral patterns.

Nevertheless, male sexual motivation for non-conceptive copulations may differ from that

for conceptive copulations. In several *Macaca* species—Tonkean macaques (*M. tonkeana*: Aujard

et al. 1998), Barbary macaques (*M. sylvanus*: Heistermann et al. 2008), and Japanese macaques

(*M. fuscata*: Garcia. Shimizu, and Huffman 2009)—the frequency of ejaculatory copulations

increases toward ovulation. In Tibetan macaques (*M. thibetana*), copulations observed outside the

mating season occur less frequently and feature a shorter mount duration, an absence of

vocalization pauses, and less frequent ejaculation (Li, Yin, and Zhou 2007). Similarly, in our

dataset—although the number of copulations with confirmed ejaculation was small—ejaculation

was observed more often with POP females than non-POP females. Given that ejaculates are

costly to produce (Thomsen et al. 2006), male bonobos might have refrained from ejaculating in

copulations with non-POP females. Further detailed observations, ideally coupled with hormonal

measures, are needed to assess qualitative differences in male sexual motivation between

conceptive and non-conceptive copulations.

Genetic evidence of pronounced male reproductive skew in bonobos predicts that males

will compete for access to POP females. However, in our dataset, the presence of POP females

within parties did not increase the frequency of intermale aggression involving the focal male.

This result contrasts with previous findings from LuiKotale (Surbeck et al. 2012) and Wamba

(Ryu et al. 2022). At LuiKotale (2007–2009), the presence of "potentially fertile" females—

defined as maximal swelling phases occurring within six months prior to conception, inferred by

counting back eight months from the subsequent birth—was associated with increased male

aggression (Surbeck et al. 2012). Similarly, at Wamba (2013–2015), male interventions in

copulation attempts, including aggression toward male competitors, were more frequent in parties

containing POP females identified from urinary hormone profiles (Ryu et al. 2022). Taken

together, those studies suggest that subordinate males can engage in direct contests with

dominants over POP females, whereas such contests were rare in our dataset.

One factor that could explain the discrepancy between our results and prior works is the

stability of within-group dominance hierarchies. Instability may weaken dominants' ability to

exclude male competitors, thereby allowing subordinates to increase their reproductive success,

as reported in Sanje mangabeys (*Cercocebus sanjei*: Fernandez 2017) and rhesus macaques

(*Macaca mulatta*: Milich et al. 2020). During our study period, the mother-son pair of Ki-KT

occupied the top ranks in the E1 group; accordingly, there seemed little scope for subordinates to

compete successfully with them over POP females. By contrast, during the period studied by Ryu

et al. (2022), alpha status in E1 was unstable: the young male JR (son of Jk) began challenging

the then-alpha male NB (son of Ki), and antagonistic interactions between them were frequent,

sometimes occurring between their mothers (see Shibata and Furuichi 2023). Likewise, during

the period studied by Surbeck et al. (2012), the Bompusa group exhibited an unbalanced hierarchy

in which mother and son did not jointly hold alpha status (Surbeck et al. 2013). Longitudinal

behavioral observations combined with paternity data will help clarify how hierarchy stability

shapes alpha male reproductive success.

5. Conclusions

Using behavioral data from free-ranging bonobos in the E1 group at Wamba (December 2022–

May 2023), this study presents three key findings: (1) higher-ranking males copulated more

frequently with POP females; (2) male copulation rates were broadly similar across dominance

ranks; and (3) lower party-level OSR values increased the focal male's probability of copulation

while reducing intermale aggression. Taken together, these results suggest that prolonged female

sexual receptivity—which creates frequent copulation opportunities for many males—may

attenuate males' proximate motivation to engage in direct contests with dominants over POP

females. Nevertheless, because these findings are derived from six months of observations in a

single group, additional multi-group, longitudinal datasets are necessary to assess generality. This

study emphasizes the importance of all-day focal follows and hormonal measures in elucidating

the reproductive strategies of male bonobos and clarifying the adaptive benefits of prolonged

female sexual receptivity.

6. Acknowledgements

We thank the staff of the Centre de Recherche en Écologie et Foresterie (CREF), Democratic

Republic of the Congo, for supporting our fieldwork at Wamba in the Luo Scientific Reserve—

especially Messrs. Bahanande Iyokango, Yembo Bafaluka, Batsina Bambambe, and Bakka

Batuafe. We are also grateful to the local assistants who helped with our fieldwork and to the

residents of Wamba for their hospitality. Fieldwork by K. Toda was supported by Grant-in-Aid

for Scientific Research (21K13670) from the Japanese Society for the Promotion of Science and

by the Canon Foundation in Europe–Kyoto University Japan–Africa Exchange Fellowship.

7. References

Ågmo, A., E. Laan. 2022. "Sexual incentive motivation, sexual behavior, and general arousal:

Do rats and humans tell the same story?" *Neuroscience & Biobehavioral*

Reviews 135:104595. https://doi.org/10.1016/j.neubiorev.2022.104595

Alexander, R. D. 1979. "Concealment of Ovulation, Parental Care, and Human Social

Evolution." In. *Evolutionary Biology and Human Social Behaviour: An Anthropological*

Perspective, edited by N. A. Chagnon, and W. G. Irons, 436–453, North Scituate, Duxbury

Press.

Altmann, J. 1974. "Observational study of behavior: sampling methods." *Behaviour* 49, no. 3–4

(1974): 227–266. https://doi.org/10.1163/156853974X00534

Aujard, F., M. Heistermann, B. Thierry, and J. K. Hodges. 1998. "Functional significance of

behavioral, morphological, and endocrine correlates across the ovarian cycle in semifree

ranging female Tonkean macaques". *American Journal of Primatology* 46, no. 4:285–309.

https://doi.org/10.1002/(SICI)1098-2345(1998)46:4<285::AID-AJP2>3.0.CO;2-8

Bates, D., M. Martin, B. Bolker, and S. Walker. 2015. "Fitting linear mixed-effects models using

lme4." *Journal of statistical software* 67: 1–48. https://doi.org/10.18637/jss.v067.i01

Beach, F. A. 1976. "Sexual attractivity, proceptivity, and receptivity in female mammals."

Hormones and behavior 7, no. 1:105–138. https://doi.org/10.1016/0018-506X(76)90008-8

Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.

S. S. White. 2009. "Generalized linear mixed models: a practical guide for ecology and

evolution". *Trends in Ecology & Evolution 24, no.* 3:127–135.

https://doi.org/10.1016/j.tree.2008.10.008

Burnham, K. P., D. R. Anderson. 2004. "Multimodel inference: understanding AIC and BIC in

model selection". *Sociological Methods & Research* 33, no. 2:261–304.

https://doi.org/10.1177/0049124104268644

Burnham, K. P., D. R. Anderson, K. P. Huyvaert. 2011. "AIC model selection and multimodel

inference in behavioral ecology: some background, observations, and comparisons".

Behavioral Ecology and Sociobiology 65:23–35. https://doi.org/10.1007/s00265-010-1029-6

Bygott, J. D. 1979. "Agonistic Behavior, Dominance, and Social Structure in Wild Chimpanzees

of the Gombe National Park." In *The Great Apes*, edited by D. A. Hamburg, and E.

R. McCown, 405–427, Menlo Park, Benjamin/Cummings.

Carnegie, S. D., L. M. Fedigan, T. E. Ziegler. 2005. "Behavioral indicators of ovarian phase in

white-faced capuchins (*Cebus capucinus*)". *American Journal of Primatology* 67, no. 1:51–

68. https://doi.org/10.1002/ajp.20169

Conaway, C. H. 1971. "Ecological adaptation and mammalian reproduction". *Biology of*

Reproduction 4, no. 3:239–247.

Connor, R. C., A. F. Richards, R. A. Smolker, and J. Mann. 1996. "Patterns of female

attractiveness in Indian Ocean bottlenose dolphins". *Behaviour* 133, no. 1–2:37–69.

David, H. A. 1987. "Ranking from unbalanced paired-comparison data". *Biometrika* 74, no.

2:432–436. https://doi.org/10.1093/biomet/74.2.432

de Vries, H., J. M. Stevens, and H Vervaecke. 2006. "Measuring and testing the steepness of

dominance hierarchies". *Animal Behaviour* 71, no. 3:585–592.

https://doi.org/10.1016/j.anbehav.2005.05.015

Deschner, T., C. Boesch. 2007. "Can the patterns of sexual swelling cycles in female Taï

chimpanzees be explained by the cost-of-sexual-attraction hypothesis?" *International Journal*

of Primatology 28:389–406. https://doi.org/10.1007/s10764-007-9120-1

Deschner, T., M. Heistermann, K. Hodges, and C. Boesch. 2003. "Timing and probability of

ovulation in relation to sex skin swelling in wild West African chimpanzees, *Pan troglodytes*

verus". *Animal Behaviour* 66, no. 3:551–560. https://doi.org/10.1006/anbe.2003.2210

Dixson, A. 1998. *Primate Sexuality*. New York, Oxford University Press.

Douglas, P. H., G. Hohmann, R. Murtagh, R. Thiessen-Bock, and T. Deschner. 2016. "Mixed

messages: wild female bonobos show high variability in the timing of ovulation in relation to

sexual swelling patterns". *BMC Evolutionary Biology* 16:1–17.

671 https://doi.org/10.1186/s12862-016-0691-3

672 Drickamer, L. C. 1974. "Social rank, observability, and sexual behaviour of rhesus monkeys

673 (*Macaca mulatta*)." *Reproduction* 37, no. 1:117–120. https://doi.org/10.1530/jrf.0.0370117

674 Dunson, D. B., D. D. Baird, A. J. Wilcox, and C. R. Weinberg. 1999. "Day-specific probabilities

675 of clinical pregnancy based on two studies with imperfect measures of ovulation". *Human*

676 *Reproduction* 14, no. 7:1835–1839. https://doi.org/10.1093/humrep/14.7.1835

677 Fernández, D. 2017. "Consequences of a male takeover on mating skew in wild Sanje

678 mangabeys". *American journal of primatology* 79, no, 7:e22532.

679 https://doi.org/10.1002/ajp.22532

680 Furuichi, T. 1987. "Sexual swelling, receptivity, and grouping of wild pygmy chimpanzee

681 females at Wamba, Zaire". *Primates* 28:309–318. https://doi.org/10.1007/BF02381014

682 Furuichi, T. 1989. "Social interactions and the life history of female *Pan paniscus* in Wamba,

683 Zaire". *International journal of Primatology* 10, no. 3:173–197.

684 https://doi.org/10.1007/BF02735199

685 Furuichi, T. 1997. "Agonistic interactions and matrifocal dominance rank of wild bonobos (*Pan*

686 *paniscus*) at Wamba". *International Journal of Primatology* 18:855–875.

687 https://doi.org/10.1023/A:1026327627943

688 Furuichi, T. 2006. "Evolution of the Social Structure of Hominoids: Reconsideration of Food

689 Distribution and the Estrus Sex Ratio". In *Human Origins and Environmental Backgrounds*,

690 edited by H. Ishida, R. Tuttle, M. Pickford, N. Ogihara, and M. Nakatsukasa, 235–248,

691 New York, Springer.

692 Furuichi, T. 2011. "Female contributions to the peaceful nature of bonobo

693 society". *Evolutionary Anthropology: Issues, News, and Reviews* 20, no. 4:131–142.

694 https://doi.org/10.1002/evan.20308

Furuichi, T. 2023. "Prolonged Sexual Receptivity in Females and Its Impact on the Evolution of

Bonobos". In *Bonobos and People at Wamba: 50 years of Research,* edited by T. Furuichi, G.

Idani, D. Kimura, H. Ihobe, and C. Hashimoto, 545–564. Singapore, Springer Nature.

https://doi.org/10.1007/978-981-99-4788-1_39

Furuichi, T., and C. Hashimoto. 2002. "Why Female Bonobos Have a Lower Copulation Rate

during Estrus than Chimpanzees". In *Behavioural Diversity in Chimpanzees and Bonobos*,

edited by C Boesch, and G. Hohmann, 156–167, Cambridge, Cambridge University Press.

https://doi.org/10.1017/CBO9780511606397

Furuichi, T. and H. Ihobe. 1994. "Variation in male relationships in bonobos and chimpanzees".

Behaviour 130, no. 3–4:211–228. https://doi.org/10.1163/156853994X00532

Furuichi, T., G. Idani, I. Ihobe, et al. 1998. "Population dynamics of wild bonobos (*Pan*

paniscus) at Wamba". *International Journal of Primatology* 19:1029–1043.

https://doi.org/10.1023/A:1020326304074

Furuichi, T., G. Idani, and I. Ihobe, et al. 2012. "Long-Term Studies on Wild Bonobos at

Wamba, Luo Scientific Reserve, DR Congo: Towards the Understanding of Female Life

History in a Male-Philopatric Species". In *Long-Term Field Studies of Primates*, edited by P.

Kappeler, D. Watts, 413–433, Berlin, Heidelberg, Springer. https://doi.org/10.1007/978-3-

642-22514-7_18

Garcia, C., K. Shimizu, and M. Huffman. 2009. "Relationship between sexual interactions and

the timing of the fertile phase in captive female Japanese macaques (*Macaca*

fuscata)". *American Journal of Primatology* 71, no. 10:868–879.

https://doi.org/10.1002/ajp.20717

Goodall, J. 1986. *The Chimpanzees of Gombe: Patterns of Behavior.* Cambridge, Harvard

University Press.

Graham, C. E., D. C. Collins, H. Robinson, and J. R. K. Preedy. 1972. "Urinary levels of

estrogens and pregnanediol and plasma levels of progesterone during the menstrual cycle of

the chimpanzee: relationship to the sexual swelling". *Endocrinology* 91, no 1:13–24.

https://doi.org/10.1210/endo-91-1-13

Hartig, F. 2024. "DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)

Regression Models. R package version 0.4.7". https://CRAN.R-

project.org/package=DHARMa

Hasegawa, T., and M. Hiraiwa-Hasegawa. 1983. "Opportunistic and restrictive matings among

wild chimpanzees in the Mahale Mountains, Tanzania. *Journal of Ethology* 1, no. 1:75–85.

https://doi.org/10.1007/BF02347833

Haselton, M. G., and K. Gildersleeve. 2011. "Can men detect ovulation?" *Current Directions in*

Psychological Science 20, no. 2:87–92. https://doi.org/10.1177/0963721411402668

Hashimoto, C. 1997. "Context and development of sexual behavior of wild bonobos (*Pan*

paniscus) at Wamba, Zaire". *International Journal of Primatology* 18:1–21.

https://doi.org/10.1023/A:1026384922066

Hashimoto, C. 2023. "Sexual Behaviors and Hormonal Background of Female Bonobos".

In *Bonobos and People at Wamba: 50 years of Research,* edited by T. Furuichi, G. Idani, D.

Kimura, H. Ihobe, and C. Hashimoto, 61–76. Singapore, Springer.

https://doi.org/10.1007/978-981-99-4788-1_39

Hashimoto, C., H. Ryu, K. Mouri, K. Shimizu, T. Sakamaki, and T. Furuichi. 2022. "Physical,

behavioral, and hormonal changes in the resumption of sexual receptivity during postpartum

infertility in female bonobos at Wamba". *Primates* 63, no. 2:109–121.

https://doi.org/10.1007/s10329-021-00968-w

Hashimoto, C., T. Furuichi, Y. Tashiro 2001. "What factors affect the size of chimpanzee parties

in the Kalinzu Forest, Uganda? Examination of fruit abundance and number of estrous

females." *International Journal of Primatology* 22, no. 6:947–959.

https://doi.org/10.1023/A:1012061504420

Hashimoto, C., Y. Tashiro, E. Hibino, et al. 2008. "Longitudinal Structure of a Unit-Group of

Bonobos: Male Philopatry and Possible Fusion of Unit-Groups". In *The Bonobos: Behavior,*

Ecology, and Conservation, edited by T. Furuichi, and J. Thompson, 107–119. New York,

Springer. https://doi.org/10.1007/978-0-387-74787-3_6

Heistermann, M., U. Möhle, H. Vervaecke, L. van Elsacker, and J. K. Hodges. 1996.

"Application of urinary and fecal steroid measurements for monitoring ovarian function and

pregnancy in the bonobo (*Pan paniscus*) and evaluation of perineal swelling patterns in

relation to endocrine events". *Biology of Reproduction* 55, no. 4:844–853.

Heistermann, M., T. Ziegler, C. P. van Schaik, K. Launhardt, P. Winkler, and J. K. Hodges.

2001. "Loss of oestrus, concealed ovulation and paternity confusion in free-ranging Hanuman

langurs". *Proceedings of the Royal Society of London. Series B: Biological Sciences* 268, no.

1484:2445–2451. https://doi.org/10.1098/rspb.2001.1833

Heistermann, M., K. Brauch, U. Möhle, D. Pfefferle, J. Dittami, and K. Hodges. 2008. "Female

ovarian cycle phase affects the timing of male sexual activity in free-ranging Barbary

macaques (*Macaca sylvanus*) of Gibraltar". *American Journal of Primatology* 70. no. 1. 44–

53. https://doi.org/10.1002/ajp.20455

Hohmann, G., and B. Fruth. 2003. "Intra- and inter-sexual aggression by bonobos in the context

of mating". *Behaviour* 140, no. 11/12. 1389–1413. https://www.jstor.org/stable/4536099

Hrdy, S. B. 1981. *The Woman That Never Evolved*. Cambridge, Harvard University Press.

Ihobe, H. 1992. "Male-male relationships among wild bonobos (*Pan paniscus*) at Wamba,

Republic of Zaire". *Primates* 33, no. 2: 163–179. https://doi.org/10.1007/BF02382747

Ishizuka, S., Y. Kawamoto, T. Sakamaki, et al. 2018. "Paternity and kin structure among

neighbouring groups in wild bonobos at Wamba". *Royal Society Open Science* 5, no.1:

171006. https://doi.org/10.1098/rsos.171006

Johns, M. A. 1980. "The Role of the Vomeronasal System in Mammalian Reproductive

Physiology". In *Chemical Signals*, edited by D. Müller-Schwarze, and R.M. Silverstein, 341–

364, Boston MA, Springer. https://doi.org/10.1007/978-1-4684-1027-3_21

Jurmain, R. 1997. "Skeletal evidence of trauma in African apes, with special reference to the

Gombe chimpanzees". *Primates* 38, no. 1: 1–14. https://doi.org/10.1007/BF02385918

Kano, T. 1992. *The Last Ape. Pygmy Chimpanzee Behavior and Ecology*. Stanford, Stanford

University Press.

Kano, T. 1996. "Male Rank Order and Copulation Rate in a Unit-Group of Bonobos at Wamba,

Zaire". In *Great Ape Societies*, edited by W. C. McGrew, L. F. Marchant, and T. Nishida,

135–145. Cambridge, Cambridge University Press.

https://doi.org/10.1017/CBO9780511752414.012

Krüger, T. H., P. Haake, U. Hartmann, M. Schedlowski, and M. S. Exton. 2002. "Orgasm-

induced prolactin secretion: feedback control of sexual drive?" *Neuroscience &*

Biobehavioral Reviews 26, no. 1:31–44. https://doi.org/10.1016/S0149-7634(01)00036-7

Kuroda, S. 1979. "Grouping of the pygmy chimpanzees". *Primates* 20:161–183.

https://doi.org/10.1007/BF02373371

Kvarnemo, C., I . Ahnesjo. 1996. "The dynamics of operational sex ratios and competition for

mates". *Trends in Ecology & Evolution* 11, no. 10:404–408. https://doi.org/10.1016/0169-

5347(96)10056-2

Li, J., H. Yin, and L. Zhou. 2007. "Non-reproductive copulation behavior among Tibetan

macaques (*Macaca thibetana*) at Huangshan, China". *Primates* 48:64–72.

https://doi.org/10.1007/s10329-006-0002-5

Lüdecke, D., M. S. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. "performance:

An R package for assessment, comparison and testing of statistical models". *Journal of Open*

Source Software 6, no.60:3139. https://doi.org/10.21105/joss.03139

Milich, K. M., A. Ruiz-Lambides, E. Maldonado, and D. Maestripieri. 2020. "Age negatively

impacts reproduction in high-ranking male rhesus macaques on Cayo Santiago, Puerto

Rico". *Scientific Reports* 10:13044. https://doi.org/10.1038/s41598-020-69922-y

Moreira, L. A. A., D. G. R. de Oliveira, M. B. C. de Sousa, D. M. A. Pessoa. 2015. "Parturition

signaling by visual cues in female marmosets (*Callithrix jacchus*)". *PLoS One* 10,

no.6:e0129319. https://doi.org/10.1371/journal.pone.0129319

Moscovice, L.R., P. H. Douglas, L. Martinez-Iñigo, M. Surbeck, L, Vigilant, G. Hohmann.

2017. "Stable and fluctuating social preferences and implications for cooperation among

female bonobos at LuiKotale, Salonga National Park, DRC". *American Journal of Physical*

Anthropology 163, no. 1:158–172. https://doi.org/10.1002/ajpa.23197

Mouginot, M., L. Cheng, M. L. Wilson. 2023. "Reproductive inequality among males in the

genus *Pan*". *Philosophical Transactions of the Royal Society B* 378, no. 1883:20220301.

https://doi.org/10.1098/rstb.2022.0301

Mouginot, M., M. L. Wilson, N. Desai, and M. Surbeck. 2024. "Differences in expression of

male aggression between wild bonobos and chimpanzees". *Current Biology* 34, No.8:1780–

1785. https://doi.org/10.1016/j.cub.2024.02.071

Mulavwa, M., T. Furuichi, K. Yangozene, et al. 2008. "Seasonal changes in fruit production and

party size of bonobos at Wamba". In *The Bonobos: Behavior, Ecology, and Conservation*,

edited by T. Furuichi, and J. Thompson, 121–134. New York, Springer.

https://doi.org/10.1007/978-0-387-74787-3_7

Muller, M. N. 2002. "Agonistic relations among Kanyawara chimpanzees". In *Behavioural*

Diversity in Chimpanzees and Bonobos, edited by C. Boesch, G. Hohmann, and L. Marchant

112–124, Cambridge, Cambridge University Press.

Muller, M. N., M. E. Thompson, and R. W. Wrangham. 2006. "Male chimpanzees prefer mating

with old females". *Current Biology* 16, no.22:2234–2238.

https://doi.org/10.1016/j.cub.2006.09.042

Nishida, T. 1968. "The social group of wild chimpanzees in the Mahali Mountains". *Primates* 9:

167–224. https://doi.org/10.1007/BF01730971

Nishida, T., H. Takasaki, and Y. Takahata. 1990. "Demography and Reproductive Profiles". In

The Chimpanzees of the Mahale Mountains: Sexual and Life History Strategies, edited by T.

Nishida, 63–97, Tokyo, University of Tokyo Press. Nishida, T., T. Kano, J. Goodall, W. C.

McGrew, and M. Nakamura. 1999. "Ethogram and ethnography of Mahale

chimpanzees". *Anthropological Science* 107, no.2:141–188.

https://doi.org/10.1537/ase.107.141

Nishida, T., N. Corp, M. Hamai, et al. 2003. "Demography, female life history, and reproductive

profiles among the chimpanzees of Mahale". *American Journal of Primatology* 59, no.3:99–

121. https://doi.org/10.1537/ase.107.141

Paoli, T. 2009. "The Absence of Sexual Coercion in Bonobos". In *Sexual Coercion in Primates*

and Humans: An Evolutionary Perspective on Male Aggression against Females, edited by

M. N. Muller, and R. W. Wrangham, 410–423, Cambridge, Harvard University Press

doi.org/10.4159/9780674054349-016

Parish, A. R. 1996. "Female relationships in bonobos (*Pan paniscus*) Evidence for bonding,

cooperation, and female dominance in a male-philopatric species". *Human Nature* 7, no. 1:

61–96.

Posit team. 2024. *RStudio: Integrated Development Environment for R*. Posit Software, PBC.

http://www.posit.co/

Pusey, A. E. 2022. "Warlike chimpanzees and peacemaking bonobos". *Proceedings of the*

National Academy of Sciences 119, no.31:e2208865119.

https://doi.org/10.1073/pnas.2208865119

R Core Team. 2024. R: *A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Rodríguez-Gironés, M. A., Enquist, M. 2001. "The evolution of female sexuality." *Animal*

Behaviour 61, no.4: 695–704. https://doi.org/10.1006/anbe.2000.1630

Rooker, K., and S. Gavrilets. 2020. "On the evolution of sexual receptivity in female

primates". *Scientific Reports* 10, no. 1:11945. https://doi.org/10.1038/s41598-020-68338-y

Ryu, H., C. Hashimoto, D. A. Hill, K. Mouri, K. Shimizu, and T. Furuichi. 2022. "Do males

know? Evidence-driven rainmaking of male bonobos to meet the fertile window of

females". *bioRxiv*, 2022-03.

Ryu, H., D.A. Hill, T. and Furuichi. 2015. "Prolonged maximal sexual swelling in wild bonobos

facilitates affiliative interactions between females". *Behaviour* 152:285–311. https://doi.

org/10.1163/1568539X-00003212

Sakamaki, T., I. Behncke, M. Laporte, M. Mulavwa, H. Ryu, H. Takemoto, N. Tokuyama, S.

Yamamoto, and T. Furuichi. 2015. "Intergroup Transfer of Females and Social Relationships

between Immigrants and Residents in Bonobo (*Pan paniscus*) Societies". In *Dispersing*

Primate Females: Life History and Social Strategies in Male-Philopatric Species, edited by

T. Furuichi, J, Yamagiwa, and F. Aureli, pp. 127–164, Tokyo, Springer.

Schielzeth, H. 2010. "Simple means to improve the interpretability of regression

coefficients". *Methods in Ecology and Evolution* 1, no.2:103–113.

https://doi.org/10.1111/j.2041-210X.2010.00012.x

Semple, S., and K. McComb. 2000. "Perception of female reproductive state from vocal cues in

a mammal species". *Proceedings of the Royal Society of London. Series B: Biological*

Sciences 267, no. 1444:707–712. https://doi.org/10.1098/rspb.2000.1060

Shibata, S., and T. Furuichi. 2023. "Intermale Relationships in Wild Bonobos at Wamba".

In *Bonobos and People at Wamba: 50 years of Research,* edited by T. Furuichi, G. Idani, D.

Kimura, H. Ihobe, and C. Hashimoto, 285–310. Singapore, Springer.

https://doi.org/10.1007/978-981-99-4788-1_23

Shibata, S., T. Furuichi. 2024. "Comparative analysis of intragroup intermale relationships: a

study of wild bonobos (*Pan paniscus*) in Wamba, Democratic Republic of Congo and

chimpanzees (*Pan troglodytes*) in Kalinzu Forest Reserve, Uganda". *Primates* 65, no. 4:243–

255.

Small, M. F. 1988. "Female primate sexual behavior and conception: Are there really sperm to

spare?" *Current Anthropology* 29, no.1:81–100. https://doi.org/10.1086/203615

Surbeck, M., C. Boesch, C. Crockford, et al. 2019. "Males with a mother living in their group

have higher paternity success in bonobos but not chimpanzees". *Current Biology* 29, no.10:

R354–R355.

Surbeck, M., C. Girard-Buttoz, L. Samuni, et al. 2021. "Attractiveness of female sexual

signaling predicts differences in female grouping patterns between bonobos and

chimpanzees". *Communications Biology* 4, no.1:1119. https://doi.org/10.1038/s42003-021-

02641-w

Surbeck, M., and G. Hohmann. 2013. "Intersexual dominance relationships and the influence of

leverage on the outcome of conflicts in wild bonobos (*Pan paniscus*)". *Behavioral Ecology*

and Sociobiology 67:1767–1780. https://doi.org/10.1007/s00265-013-1584-8

Surbeck, M., K. E. Langergraber, B. Fruth, L. Vigilant, and G. Hohmann. 2017. "Male

reproductive skew is higher in bonobos than chimpanzees". *Current Biology* 27, no.13:R640–

R641. https://doi.org/10.1016/j.cub.2017.05.039

Surbeck, M., R. Mundry, and G. Hohmann. 2011. "Mothers matter! Maternal support,

dominance status and mating success in male bonobos (*Pan paniscus*)". *Proceedings of the*

Royal Society B: Biological Sciences 278, no. 1705:590–598.

https://doi.org/10.1098/rspb.2010.1572

Surbeck, M., T. Deschner, G. Schubert, A. Weltring, and Hohmann, G. 2012. "Mate

competition, testosterone and intersexual relationships in bonobos, *Pan paniscus*". *Animal*

Behaviour 83, no. 3:659–669.

Symons, D. 1979. *The Evolution of Human Sexuality*. New York, Oxford University.

Takahata, Y., H. Ihobe, and G. I. Idani. 1996. "Comparing Copulations of Chimpanzees and

Bonobos: Do Females Exhibit Proceptivity or Receptivity?". In *Great Ape Societies,* edited

by W. C. McGrew, L. F. Marchant, and T. Nishida, 146–156, Cambridge, Cambridge

University Press.

Thompson, M. E. 2005. "Reproductive endocrinology of wild female chimpanzees (*Pan*

troglodytes schweinfurthii): methodological considerations and the role of hormones in sex

and conception". *American Journal of Primatology* 67, no. 1:137–158.

https://doi.org/10.1002/ajp.20174

Thompson, M. E. 2013. "Reproductive ecology of female chimpanzees". *American Journal of*

Primatology 75, no.3:222–237. https://doi.org/10.1002/ajp.22084

Thompson, M. E., and K. Sabbi. 2024. "Evolutionary Demography of the Great Apes". In

Human Evolutionary Demography, edited by O, Burger, R. Lee, and R. Sear, 423–474,

Cambridge UK, Open Book Publishers. https://doi.org/10.11647/obp.0251.19

Thompson, E. M., and P. T. Ellison. 2017. "Fertility and Fecundity". In *Chimpanzees and*

Human Evolution, edited by M. N. Muller, R. W. Wrangham, and D. Pilbeam, 217–258,

Cambridge, MA, Harvard University Press. doi.org/10.4159/9780674982642-006

Thomsen, R., J. Soltis, M. Matsubara, K. Matsubayashi, M. Onuma, and O. Takenaka. 2006.

"How costly are ejaculates for Japanese macaques?" *Primates* 47. no. 3:272–274.

https://doi.org/10.1007/s10329-005-0171-7

Toda, K., N. Tokuyama, and T. Sakamaki. 2023. "Multiple Phases of Natal Transfer Process in

Female Bonobos and Factors Underlying Each Phase: Findings from Long-Term

Observations in Wild Populations". In *Bonobos and People at Wamba: 50 years of Research,*

edited by T. Furuichi, G. Idani, D. Kimura, H. Ihobe, and C. Hashimoto, 181–209. Singapore,

Springer. https://doi.org/10.1007/978-981-99-4788-1_23

Tokuyama, N., and T. Furuichi. 2016. "Do friends help each other? Patterns of female coalition

formation in wild bonobos at Wamba". *Animal Behaviour* 119:27–35.

https://doi.org/10.1016/j.anbehav.2016.06.021

Tutin, C. E., and P. R. McGinnis. 1981. "Chimpanzee Reproduction in the Wild". In

Reproductive Biology of the Great Apes: Comparative and Biomedical Perspectives, edited

by C. E. Graham, 239–264. New York, Academic Press. https://doi.org/10.1016/B978-0-122-

95020-9.X5001-8

Wallis, J. 1985. "Synchrony of estrous swelling in captive group-living chimpanzees (*Pan*

troglodytes)". *International Journal of Primatology* 6, no. 3:335–350.

https://doi.org/10.1007/BF02745505

Wallis, J. 1992. "Chimpanzee genital swelling and its role in the pattern of sociosexual

behavior". *American Journal of Primatology* 28, no. 2:101–113.

https://doi.org/10.1002/ajp.1350280203

Wallis, J. 1997. "A survey of reproductive parameters in the free-ranging chimpanzees of

Gombe National Park". *Reproduction* 109, no. 2:297–307.

https://doi.org/10.1530/jrf.0.1090297

Wallis, J., and J. Goodall, 1993. "Anogenital swelling in pregnant chimpanzees of Gombe

National Park". *American Journal of Primatology* 31, no.2:89–98.

Weir, B. J., and I. W. Rowlands. 1973. "Reproductive strategies of mammals". *Annual Review of*

Ecology and Systematics 4:139–163.

Weir, L. K., J. W. Grant, and J. A. Hutchings. 2011. "The influence of operational sex ratio on

the intensity of competition for mates". *The American Naturalist* 177, no.2:167–176.

Wilcox, A. J., C. R. Weinberg, and D. D. Baird. 1995. "Timing of sexual intercourse in relation

to ovulation—effects on the probability of conception, survival of the pregnancy, and sex of

the baby". *New England Journal of Medicine* 333, no.23:1517–1521.

Wilson, M. L., C. Boesch, B. Fruth, et al. 2014. "Lethal aggression in *Pan* is better explained by

adaptive strategies than human impacts". *Nature* 513, no.7518:414–417.

https://doi.org/10.1038/nature13727

Wrangham, R. W. 2002. "The Cost of Sexual Attraction: Is There a Trade-Off in Female *Pan*

between Sex Appeal and Received Coercion". *Behavioural Diversity in Chimpanzees and*

Bonobos, edited by C. Boesch, and G. Hohmann, 204–216, Cambridge, Cambridge

University Press.

Yokoyama, T., and T. Furuichi. 2022. "Why bonobos show a high reproductive skew towards

high-ranking males: Analyses for association and mating patterns concerning female sexual

states". *Primates* 63, no. 5:483–494. https://doi.org/10.1007/s10329-022-01004-1

Zinner, D., and T. Deschner. 2000. "Sexual swellings in female hamadryas baboons after male

take-overs: "deceptive" swellings as a possible female counter-strategy against

infanticide". *American Journal of Primatology* 52, no.4:157–168.

https://doi.org/10.1002/1098-2345(200012)52:4<157::AID-AJP1>3.0.CO;2-L

Zuur, A. F., E. N. Ieno, N. Walker, A. A. Saveliev, G. M. Smith. 2009. *Mixed Effects Models and*

Extensions in Ecology with R. New York, Springer. https://doi.org/10.1007/978-0-387-87458-

6

8. Figure Legends

Fig.1 Proportion of copulations with POP females for each male. Males are ordered from left to

right by increasing dominance rank. The curve and shaded band represent the predicted proportion

and its 95% confidence interval from the best-fitting binomial generalized linear model (GLM).

Point sizes are proportional to the number of copulations for which the female's fertility status

(POP vs. non-POP) was determined.

Fig.2 Copulation rates (events per hour of focal follows) for each focal male. Males are ordered

from left to right by increasing dominance rank. Gray and white bars denote copulations with

adult females and adolescent females, respectively.

Fig.3 Effects of the number of mature males and the number of females with maximal swellings

(MS females) present in the party on the probability that the focal male engaged in copulation (A,

B) and was involved in intermale aggression (C, D) during 1-hour interval sessions. Point sizes

are proportional to the number of sessions observed at each value of (A, C) the number of mature

males and (B, D) the number of MS females. Curves and shaded bands represent predicted

probabilities with 95% confidence intervals from the best-fitting binomial generalized linear

mixed models (GLMMs). Colors indicate predictions evaluated at the 25th (red), 50th (blue), and

75th (yellow) percentiles of the other predictor (i.e., in panel A and C, predictors are computed at

these percentiles of the number of MS females; in panel B and D, at these percentiles of the

number of mature males).

  TABLE 1 Overview of male subjects in the E1 group at Wamba during the study period

  (December 2022–May 2023).

ID	Age[c] (yrs)	Male rank [David Score]	Observation hours during all-day focal sampling [day]	Number of copulation acts	
				Focal data [number of days copulation occurrence]	Focal + ad-libitum data [POP females, non-POPs]
TN[a]	52*	Not used [-2.9]	Not collected	Not collected	(5 [0, 5])[d]
TW[b]	49*	Not used [-12.9]	(39.2 [4])[d]	(0)[d]	(0)[d]
DI	48*	5th [4.6]	39.7 [4]	7 [3]	19 [1, 17]
NB	35	4th [7.2]	44.7 [5]	9 [5]	31 [14, 14]
GC	33*	8th [-11.2]	37.5 [4]	7 [4]	17 [1, 14]
KT	19	1st [53.4]	36.4 [4]	3 [3]	33 [18, 13]
JO	17	3rd [7.5]	36.7 [4]	11 [4]	38 [8, 28]
KY	14	2nd [10.0]	30.6 [4]	6 [3]	35 [15, 20]
HC	14	6th [-2.7]	32.0 [4]	6 [3]	25 [3, 19]
SE	12	7th [-6.1]	32.6 [4]	9 [3]	20 [3, 16]
NI	10	10th [-28.9]	25.1 [4]	7 [3]	23 [3, 20]
YD	9	9th [-27.0]	34.4 [4]	2 [2]	27 [1, 22]

[a]TN was not included in the focal subject, because he had been lost for a five month in E1 group

and was absent at the beginning of this study.

[b]TW had seemed erectile dysfunction before the study period, and thus data on him were not used

for the analysis.

[c]Asterisks in the column means estimated ages.

[d]Data in parentheses were not used for the analysis.

[g]Not included when female fertility status could not be determined

21　**TABLE 2** Parameter estimates from the GLM and GLMMs selected as the best models by AIC

Responses	Predictors	Estimate	SE	Z	*p*-value
(A) Binomial GLM					
Binary data of the proportion whether to copulate with POP or non-POP females	(Intercept)	−1.39	0.19	−7.33	
	Male rank (linear)	0.97	0.19	5.06	< 0.001***
(B) Poisson GLMM 1					
Count data of copulations with adult females for each focal follow	(Intercept)	−2.06	0.23	−8.9	
	Number of MS females (linear)	0.94	0.28	3.34	< 0.001***
	Number of MS females (quadratic)	-0.47	0.23	−2.03	0.043*
(C) Poisson GLMM 2					
Count data of copulations with adolescent females for each focal follow	(Intercept)	−2.06	0.23	−8.9	
(D) Binomial GLMM 1					
Binary data of whether copulation occurred or not per session during focal follow	(Intercept)	−1.16	0.30	−3.87	
	Number of mature males (linear)	−0.83	0.23	−3.68	< 0.001***
	Number of MS females (linear)	1.36	0.30	4.60	< 0.001***
	Number of MS females (quadratic)	−0.44	0.15	−2.92	0.004**
	Presence of POP females (yes/no)	−0.53	0.35	−1.54	0.124
	Resting time (linear)	−0.50	0.21	−2.40	0.017*
	Resting time (quadratic)	−0.45	0.20	−2.28	0.022*
(E) Binomial GLMM 2					
Binary data of whether intermale aggressions occurred or not per session during focal follow	(Intercept)	−1.54	0.29	−5.38	
	Number of mature males (linear)	1.06	0.22	4.74	< 0.001***
	Number of MS females (linear)	−0.56	0.22	−2.58	0.010**
	Resting time (linear)	0.18	0.20	0.90	0.369
	Resing time (quadratic)	−0.61	0.19	−3.24	0.001**

22　Predictors that significantly influence the responses are indicated in bold.