Title Page

A short informative title

Rank-biased access to females in the likely fertile period despite comparable copulation rates in

male bonobos at Wamba

A short running title

Male bonobo rank-dependent mating strategies

Author name

Kazuya Toda[1], Furuichi Takeshi[2]

Affiliation

[1]Research Center for Integrative Evolutionary Science, The Graduate University for Advanced

Studies, Sokendai, Japan

[2]Wildlife Research Center, Kyoto University, Inuyama, Aichi, Japan

Email address

[1]toda.kazuya.h17@kyoto-u.jp

[2]furuichit@gmail.com

Ethics statement

Our field studies of free-ranging bonobos at Wamba in the Luo Scientific Reserve were

approved by the Ministry of Scientific Research and the Center for Research on Ecology and

Forestry in the Democratic Republic of Congo. This study adhered to the Animal Research

Guidelines of The Graduate University for Advanced Studies (SOKENDAI), Japan, revised on

March 23, 2021.

Data availability statement

The data that support the findings of this study are available from the corresponding author

upon reasonable request.

Keywords

Pan paniscus, prolonged sexual receptivity, non-conceptive copulation, intermale mating

competition, operational sex ratio

Abstract

Objectives

Female bonobos (*Pan paniscus*) show prolonged sexual swelling and copulate across an

unusually extended timeframe, including the postpartum infertile stage. We tested the updated

prolonged sexual receptivity hypothesis that the simultaneous presence of multiple receptive

females disperses male mating effort and weakens male-male contest competition by examining

rank effects on copulation rates and mating partners.

Materials and Methods

In a free-ranging bonobo group at Wamba, we collected copulation and party-composition data

using all-day focal follows. Based on daily swelling scores, we operationally defined a likely

fertile period (LFP) from detumescence at the end of the maximal swelling phase. We analyzed

rank-dependent patterns in copulation rates and mating partners were analyzed using

generalized linear mixed models (GLMMs).

Results

Copulation rates did not differ detectably across male dominance ranks, with approximately

twice per day on average. In contrast, higher-ranking males were more likely to copulate with

LFP females. At the party level, lower-ranking males increasingly copulated with non-LFP

females as the number of LFP females present increased.

Discussion

These findings indicate that pronounced male reproductive skew in bonobos appears to arise

through rank-biased access to females in the fertile window despite broadly similar copulation

rates across males. Subordinate males' tendency to shift mating effort toward non-LFP females

when LFP females are available may reduce injury risks of direct contest competition while

facilitating dominant males to guard fertile females effectively. Therefore, prolonged sexual

receptivity in female bonobos could be a key factor yielding the condition where lower male

aggression and higher reproductive skew coexist.

1. Introduction

Sexual receptivity can be defined as the physiological, morphological, and behavioral readiness

of a mammalian female to accept a male and engage in copulation (Dixson 1998). In most

non-primate mammals, receptivity is largely confined to the periovulatory phase of the estrus

cycle and is tightly regulated by fluctuations in ovarian hormones such as estrogen and

progesterone (Conaway 1971; Weir and Rowlands 1973). Prosimian primates show a broadly

similar pattern, with female receptivity concentrated within a relatively narrow fertile window

(Small 1988). In contrast, across anthropoid primates, sexual behavior is often less tightly

governed by ovarian hormones, and females may remain receptive outside the fertile phase

(Wallen and Zehr 2004). Prolonged sexual receptivity can impose fitness costs, including

opportunity and energetic costs associated with sexual activity (Kunz et al. 2021), greater

exposure to male harassment and coercion that can result in injury (Muller et al. 2011), and

increased risk of sexually transmitted pathogens (Phillips-Conroy et al. 1994). Nonetheless,

these costs may be outweighed when prolonged sexual receptivity benefits females, for example

by confusing paternity to reduce infanticide risk and/or by eliciting paternal investment

(Fernandez-Duque et al. 2020; Rooker and Gavrilets 2020).

Across anthropoids, females vary widely in whether and how they advertise fertility to

males. In some polygynandrous taxa, females express graded visual cues of ovulation, such as

sexual skin tumescence or skin coloration, which can bias male mating effort toward fertile

periods while maintaining uncertainty across a prolonged receptive period. This combination

may allow females to increase conception probability while also promoting assurance with

paternity confusion (e.g., Engelhardt et al. 2004; Setchell and Wickings 2004; Brauch et al.

2007; Gesquiere et al. 2007; Dubuc et al. 2009 Higham et al. 2009). In contrast, monogamous

taxa typically show slight or absent visual ovulatory cues (i.e., concealed ovulation), and

females are receptive over a long range of their estrus cycles (e.g., Kendrick and Dixson 1983;

Barelli et al. 2008; Fernandez-Duque et al. 2011). A comparative phylogenetic analysis suggests

an evolutionary scenario that concealed ovulation among anthropoids tended to occur under

polygamous or polygynandrous mating systems as a counterstrategy against male infanticide

and thereafter can have facilitated the transition to monogamous (Sillén-Tullberg and Moller

1993). In humans, concealed ovulation with continuous sexual receptivity has been argued to

promote pair bonds and paternal investment (Alexander and Noonan 1979; Strassmann 1981;

Rodriguez-Girones and Enquist 2001). However, there are other hypotheses that concealed

ovulation in humans could have evolved as a side effect of the transition to bipedal locomotion

(e.g., reduced visibility of sexual swelling) or the adaptation to African savannah environment

(e.g., increased physiological cost of water retention), and so on (Pawłowski 1999).

Two *Pan* species, chimpanzees (*P. troglodytes*) and bonobos (*P. paniscus*), the closest

living relatives to humans, are non-seasonal breeders with interbirth intervals of approximately

4–6 years (Emery Thompson and Sabbi 2024). They live in multi-male/multi-female groups

with promiscuous mating systems, where females display sexual skin tumescence as visual

ovulatory cues, reflecting fluctuations in estrogen and progesterone levels (Graham et al. 1972;

Heistermann et al. 1996). This advertised ovulation is phylogenetically considered to have been

acquired by the common ancestor of the genus *Pan* after the divergence from the hominin

ancestor (Sillén-Tullberg and Moller 1993). Although chimpanzees and bonobos diverged at an

estimated 2.6–0.8 million years ago (Takemoto, Kawamoto, and Furuichi 2015; Takemoto et al.

2017), females of both *Pan* species are known to exhibit prolonged sexual receptivity from the

facts that their maximal swellings persist longer than the actual fertile window in ovulation

cycles and can appear after conception and during postpartum infertile phase (Deschner et al.

2003; Douglas et al. 2016).

However, female bonobos spend a remarkably longer period of displaying sexual swelling

and engaging in copulation during the infertile period than female chimpanzees (Furuichi 2011).

One factor reason is that the duration of maximal swelling phases in each cycle is about two

days longer on average in female bonobos than chimpanzees (13.5 ± 1.8 SD days for bonobos

and 11.3 ± 1.1 SD days for chimpanzees: reviewed by Ryu et al. 2025). Another factor is that the

span from parturition to the first observed sexual swelling postpartum is much shorter in female

bonobos than chimpanzees (0.7 ± 0.4 SD years for bonobos: Hashimoto et al. 2022; 1.8 ± 0.4

SD and 2.4 ± 1.5 SD years for western chimpanzees: Deschner and Boesch 2007; 3.4–4.6 years

on average for eastern chimpanzees: Wallis 1997; Nishida et al. 2003; Emery Thompson and

Ellison 2017). The physiological mechanisms underlying these interspecific differences remains

poorly understood; one possibility is that the threshold of estrogenic stimulation causing sexual

skin tumescence is lower in female bonobos than in female chimpanzees (Hashimoto et al.

2022).

The exceptionally early resumption of sexual swelling in female bonobos during

postpartum infertile stage has been hypothesized to contribute to the increased number of

receptive females in groups and thereby hinder few dominant males from monopolizing access

to mating partners (termed as "prolonged sexual receptivity" hypothesis: Furuichi 2023). The

operational sex ratio, calculated by the number of mature males per that of sexually receptive

females present at a given time, is widely used as an indicator of the intensity of male-male

contest competition over mates (Kvarnemo and Ahnesjo 1996; Weir, Grant, and Hutchings

2011). The operational sex ratio in a bonobo group at Wamba was reported to be lower than

those in chimpanzee groups at Mahale and Gombe (2.8 versus 4.2 and 12.3: Furuichi and

Hashimoto 2002). Moreover, although both *Pan* species exhibit a high degree of fission-fusion

dynamics in their grouping patterns (Nishida 1968; Kuroda 1979), the mean number of females

present within one party (i.e., a flexible subgroup whose membership varies in size and

composition over time) was higher in bonobos at Wamba (3.2 ± 1.1 SD adult females) than in

eastern chimpanzees at Kalinzu (1.2 ± 0.8 SD adult females) (Mulavwa et al. 2008), whereas it

was only slightly higher in bonobos at LuiKotale than in western chimpanzees at Taï (4.5 versus

3.7 and 3.3 on average: Surbeck et al. 2021). High party-level association among females in

bonobos may therefore further weaken male-male mating competition (Furuichi 2023).

Although male bonobos use aggression at a similar or higher frequency than male

chimpanzees (Shibata and Furuichi 2024; Mouginot, Cheng, and Wilson 2023), it is widely

accepted that the intensity of male aggression is substantially weaker in bonobos than in

chimpanzees (Tokuyama 2023). Male chimpanzees exhibit aggressive attitudes toward males in

neighboring groups and cooperate in lethal raids that can lead to territorial expansion (Watts et

al. 2006; Boesch et al. 2008; Mitani, Watts, and Amsler 2010). In some populations, male

chimpanzees also occasionally kill infants (Arcadi and Wrangham 1999; Wilson et al. 2014;

Lowe et al. 2020) and other male within their own groups (Fawcett and Muhumuza 2000; Watts

2004; Kaburu, Inoue, and Newton-Fisher 2013). By contrast, lethal aggression has been not

reported to date in male bonobos (Furuichi 2011; Wilson et al. 2014; Tokuyama 2023) but in

females (Pashchevskaya et al. 2025). Skeletal evidence from museum specimens likewise

indicates higher rates of trauma in chimpanzees than in bonobos (Jurmain 1997). Additionally,

in 693 within-group aggressive interactions observed from a bonobo group at Wamba, there

were no cases of males injuring conspecifics but two cases of female-inflicted injuries, even

though males exhibit aggressive behavior more often than females (Tokuyama, Sakamaki, and

Furuichi 2019). Furthermore, male bonobos have a smaller size of the canine tooth (a biological

weaponry) than male chimpanzees (Kelley 1995; Balolia et al. 2025). Also, canine dimorphism,

a generalized indicator of the strength of male-male contest competition over mates in primates

(Plavcan and van Schaik 1992; Plavcan, van Schaik, and Kappeler 1995), is less in bonobos

than chimpanzees (Kelley 1995; Balolia et al. 2025).

Previously, weak male-male mating competition in bonobos predicts that reproductive

success was widely distributed among males within group. However, recent genetic paternity

analyses from multiple bonobo groups at LuiKotale, Wamba, and Kokolopori consistently show

pronounced male reproductive skew (Surbeck et al. 2017; Ishizuka et al. 2018; Mouginot,

Cheng, and Wilson 2023). Across groups, the most successful males sired a larger share of

offspring than in chimpanzees (Mouginot, Cheng, and Wilson 2023: 69 ± 11% in bonobos; 44 ±

25% in chimpanzees). Male reproductive success also covaries with dominance rank; alpha

males in two study groups at Wamba achieved the highest reproductive success (Ishizuka et al.

2018), and alpha males in two of three study groups at Kokolopori sired more offspring than

expected (Mouginot, Cheng, and Wilson 2023). These genetic outcomes serve as evidence that

dominant males are able to monopolize access to fertile females despite noisy ovulatory signals

of sexual swelling during the infertile period. Following genetic research, one recent study

based on behavioral and hormonal data demonstrates that male bonobos extract usable ovulation

timing with some accuracy and concentrate mating effort on females in the periovulatory phase

(Ryu et al. 2025). In addition to high dominance rank empowering male bonobos to guard mates

by excluding subordinate competitors (Kano 1996; Hohmann and Fruth 2003; Surbeck et al.

2012; Mouginot et al. 2024), their ability to identify fertile females by effectively managing

graded ovulatory signals may help explain the pronounced male reproductive skew.

Pronounced inequality in male reproductive rates in bonobos raises the question of why

male-male mating competition does not escalate into severe aggression (Shibata and Furuichi

2023). One conceivable explanation is that less aggressive males are preferred as mating

partners by females or easy to receive permission of sharing the same space with with females

(Surbeck et al. 2012). However, it was reported from Kokolopori bonobos that the most

successful two males exhibited the highest rates of contact aggression and aggression against

females more often occurred by dominant males than subordinates (Mouginot, Cheng, and

Wilson 2023; Mouginot et al. 2024). This pattern suggests that female mate choice alone is

unlikely to account for why severe aggression was not favored by natural selection in

bonobos. Herein, we hypothesized that the simultaneous presence of multiple receptive

females disperses male copulation attempts irrespective of female fertility status and thereby

reduce the intensity of contest competition over mates, updating the prolonged sexual receptivity hypothesis (Furuichi 2023). Specifically. when multiple receptive females are available concurrently, subordinate males would adopt flexible mating strategies, conditionally shifting their mating effort toward females that entail lower risks of interference or aggression from dominant males. Such strategic allocation of mating effort could indirectly facilitate dominant males to monopolize access to fertile females.

It is a critical challenge to resolve the contradiction between lower male aggression and higher reproductive skew in bonobos for a better understanding of diverse social structures among primates. To examine the updated prolonged sexual receptivity hypothesis, we collected all-day focal data on copulation and party composition in a free-ranging bonobo group at Wamba. Specifically, this study addressed the following three predictions:

(1) Copulation rates are predicted to differ little among males of different ranks. If the simultaneous presence of multiple receptive females facilitates a wide range of males to engage in copulation, males of different dominance ranks would exhibit comparable copulation rates. To test this prediction, we investigated the relationship between males' ranks and their copulation rates using all-day focal data, which reduce sampling bias relative to short focal or ad libitum sampling.

(2) Dominant males are predicted to have priority access to fertile females. Given the pronounced male reproductive skew, dominant males should copulate more frequently with fertile females than subordinates. To test this prediction, we estimated a likely fertile period (LFP) of female bonobos from the end of maximal swelling phase (MSP) based on daily assessments of sexual swelling status and compared the proportion of copulations involving higher- and lower-ranking males between LFP and non-LFP, using all-day focal data on females.

217   (3) Subordinate males are predicted to have a tendency to target receptive females who

218   are less closely guarded by dominants. If males strategically allocate mating effort to

219   reduce the risks of interference or aggression, lower-ranking males would be more

220   likely than higher-ranking males to copulate with non-LFP females when LFP females

221   are present in the party. To test this prediction, we analyzed male-based all-day focal

222   data on copulation and party compositions to examine the effect of within-party male

223   rank on the probability of copulation with non-LFP females, while distinguishing

224   parties where LFP females were present from those where they were absent.

2. Materials and Methods

2.1 Study group and subjects

This study was conducted on a single habituated group of free-ranging bonobos (E1 group) at

Wamba in the Luo Scientific Reserve, Democratic Republic of the Congo, a site studied since

1974 (Kano 1992; Furuichi et al. 2012). Data collection spanned from December 1, 2022 and

May 2, 2023 (153 days). During this period, the E1 group comprised 45–48 individuals,

including 10 adult males, 2 adolescent males, 17 adult females, and 3 adolescent females. The

size of E1 group increased gradually after 2009, largely due to the immigration of nulliparous

females (Toda, Tokuyama, and Sakamaki 2023), and thus the number of adult and adolescent

females was highest ever (Figure S1). All adult and adolescent females had immigrated from

other groups. Five adult and two adolescent males resided with their mothers within group.

Age classes were defined as follows. Adult males: $\geq$12 years, given growth in male body

mass in zoo-housed bonobos ceases around 12 years (Berghaenel et al. 2023). Adolescent

males: 8–<12 years, in light of the rapid rise in urinary testosterone beginning around 7 years

(Behringer et al. 2014; Berghaenel et al. 2023) and the report that captive male bonobos can

reproduce from about 7 years of age (Thompson-Handler 1990). Adult females: ≥12 years or

parous, consistent with a mean age at first parturition of approximately 12 years at Wamba

(Toda, Tokuyama, and Sakamaki 2023). Adolescent females: nulliparous females aged 8–<12

years, as females at Wamba typically disperse from the natal group at about 7 years and often

engage in copulation after immigration (Toda et al. 2022).

All-day focal follows were conducted on adults and adolescents in the E1 group (Table 1:

males; Table 2: females). One adult male (TN) was not followed because he had been missing

from E1 for five months and was absent at the start of this study. One adult female (Aq) was

also excluded from the focal subject because she had not been habituated well and thus was

impossible to conduct focal follows without disturbing her natural behavior. Another adult

female (No) was last seen on December 31, 2022; consequently, only one all-day focal follow

was obtained for her.

2.2 All-day focal animal sampling

Trained research assistants followed a single party of the E1 group on a daily basis from the

time the bonobos descended from their night beds (approximately 0600 h) until they constructed

new night beds (about 1700 h). When the party being followed split into several parties,

observers, except for those who were engaging in focal animal sampling, typically followed the

largest party. The first author (KT) and a research assistant, Lokemba Batsindelia (LB),

conducted focal animal sampling (Altmann 1974) throughout the day to collect continuous

behavioral data and party composition on adults and adolescents.

On each observation day, we selected a focal subject from party members using a

predetermined random order. A focal follow continued until one of the following conditions was

met: the focal subject constructed a new night bed; darkness after sunset precluded

identification; or the subject was out of sight for ≥2 consecutive hours. Focal follows with ≥4

hours of observation time were treated as all-day focal data and included in analyses. If a focal

subject was lost for ≥2 consecutive hours before reaching 4 hours, a second focal follow was

initiated that day only if it could start by 1200 h. If the predetermined focal subject could not be

located within 30 minutes of the start of the party follow, we selected the next available

individual in the predetermined order from among party members present at that time. From the

midpoint of the study onward, we prioritized undersampled subjects to balance observation

effort. To ensure independence among all-day focal follows, ≥20 days elapsed before

resampling the same subject.

During each focal follow, we recorded a wide range of behaviors—resting, feeding,

moving, and social interactions—along with the duration of each activity and the identities of

participants. We also noted party composition—defined as the set of adult and adolescent

individuals present during focal follows—for each time window separated at fixed hourly

intervals (i.e., 0600–0700 h, 0700–0800 h, 0800–0900 h, …, 1600–1700 h), following the

one-hour party method (Hashimoto, Furuichi, and Tashiro 2001). Although focal follows

sometimes began before 0600 h or ended after 1700 h, 1-hour party data outside 0600–1700 h

were excluded from dataset because of concern that darkness reduces the accuracy of

identification.

In total, 1033.8 hours of all-day focal data were collected by KT (544.8 hours) and LB

(489.0 hours) over 116 days from 11 males (384.1 hours over 45 days; Table 1) and 19 females

(649.7 hours over 71 days; Table 2). Mean focal time per day was 8.9 hours (SD = 1.4; range =

4.2–11.0 hours), with a median start time of 0631 h (range = 0551–1142 h) and a median end

time of 1700 h (range = 1247–1743 h). Moreover, focal party composition was recorded for a

total of 1211 one-hour time windows (hereafter termed as OTWs) (males: 462 OTWs; females:

748 OTWs). Mean observation time per hourly time window was 51.8 minutes (SD = 13.3, range = 0.2–60.0 minutes).

2.3 Definition of copulation

Copulation was defined as a heterosexual interaction in which a male mounted a female with penile-vaginal intromission accompanied by thrusting. Because ejaculation was often not visually discernible, intercourse was recorded as copulation regardless of whether ejaculation could be confirmed (Hashimoto 1997; Ryu et al. 2025). We recorded all observable copulatory interactions between adolescent or adult individuals with ad libitum sampling. To avoid inflating counts, successive copulations involving the same dyad within 10 minutes were treated as a single copulation event.

During all-day focal follows, we observed 67 copulation events involving the focal male and 67 involving the focal female. In addition, we recorded 134 copulations ad libitum that did not involve the focal subject, yielding 268 copulations in total (19–38 events per male). Visible sperm expulsion from the vagina immediately after copulation was confirmed in 17/268 events (6.3%). With the exception of one older male (TW; estimated 51 years old), copulation was observed from all male subjects. TW has not been observed copulating or exhibiting erections since at least 2015 (Toda personal observation), and thus his focal data (38.5 hours and 42 OTWs) were excluded from analyses.

2.4 Definition of aggression

Aggression was defined as a directed aggressive interaction, including contact aggression (e.g., punching, kicking, pushing, tackling, biting) and threat display (e.g., chasing, charging, charging while branch-dragging, branch-shaking) (Nishida et al. 1999). Undirected display

lacking a clear target was not classified as aggression. We recorded all aggressive interactions

between adolescent or adult individuals and noted the target's response, including submissive

behavior (e.g., fleeing, grinning, screaming) and retaliation toward the aggressor. Successive

aggressive interactions involving the same dyad within 10 minutes were treated as a single

aggression event.

In total, we recorded 297 aggression events: 124 involving the focal male, 51 involving the

focal female, and 122 not involving the the focal subject. Of these, 138 were male-male (46.5%),

81 male-to-female (27.3%), 62 female-to-male (20.9%), and 16 female-female (5.4%). Contact

aggression occurred in 95/297 events (32.0%): initiated by males in 43/95 (45.2%) and by

females in 52/95 (54.7%). Submissive responses were observed in 201/297 events (67.7%),

allowing us to identify winners and losers.

2.5 Evaluation of male dominance ranks

To infer the dominance hierarchy of the E1 group, we calculated David's scores (David 1987)

from an interaction matrix comprising 201 aggressive interactions that elicited submissive

responses, using dyadic dominance indices that incorporate win-loss proportions and correct for

unequal interaction opportunities (de Vries 1998). Although our primary aim was to evaluate

male dominance ranks, both sexes were included in the same matrix because male bonobos are

codominant with females (Furuichi 1997; Surbeck and Hohmann 2013). We computed the

David's scores for each male ($N = 12$; Table 1) and each female ($N = 20$; Table 2) with the 'DS'

function in the 'EloRating' package (Neumann and Kulik 2024) in R version 4.4.2 (R Core

Team 2024). We also quantified hierarchy steepness based on David's scores (de Vries, Stevens,

and Vervaecke 2006) and the modified Landau's linearity index (de Vries 1995), and compared

the observed values with those from randomized matrices using the 'steepness' and 'h.index'

functions in the 'EloRating' package. Both steepness and linearity were greater than expected

(one-tailed $p = 0.001$ and $p = 0.020$, respectively).

The median rates of aggression from the focal male during all-day focal follows were

0.067 events/h towards males (range $= 0$–0.332, $N = 32$) and 0.052 events/h towards and

females (range $= 0$–0.221, $N = 31$). The former figure is quite similar as the frequency reported

from Mouginot et al. (2024) at Kokolopori (median $= 0.08$ events/h, range $= 0$–0.300) but the

latter figure appears about 10 times higher (median $= 0.005$ events/h, range $= 0$–0.027), yet

aggression elicited submissive responses from females was in the minority (11/31 events

[35.5%]). The correlation of male dominance ranks with the rate of aggression towards males

was not significant (Spearman's rank correlation test: $\rho = -0.55$, $p = 0.100$), but that towards

females was significant ($\rho = -0.74$, $p = 0.015$).

2.6 Assessment of sexual swelling states

We recorded the daily variation in sexual swelling states of female bonobos in the E1 group, by

scoring the size and firmness of the sexual skin. Following established protocols (Furuichi

1987; Ryu, Hill, and Furuichi 2015), each female was assigned a score of 1 to 3: 1 = no swelling,

2 = intermediate swelling, 3 = maximal swelling (Ryu et al. 2025). Score 1 was determined with

the minimum range of sexual skin size. However, the size alone did not always clearly separate

scores 2 and 3, we additionally relied on the firmness and shininess seen of the sexual skin to

distinguish these categories. Detumescence day was defined as the first day on which the

maximal swelling began to subside, accompanied by loss of firmness and shininess, and the

score shifted from 3 to 2.

To reduce observation error and scoring bias, researchers and trained assistants discussed

and reached consensus on each female's swelling score at a daily evening meeting. During the

study period, the mean number of females with maximal swelling in the followed party per

observation day was 4.5 (SD = 2.4, range = 0–10), comprising 3.1 adult females (SD = 1.8,

range = 0–7, N = 16) and 1.4 adolescent females (SD = 1.0, range = 0–3, N = 3).

The maximal swelling phase (MSP) was defined as the interval from the first day a

female was scored 3 to the last day scored 3 immediately before detumescence. Sample gaps in

daily scores arose when a female was absent from the followed party or when observations were

missed. If the score was identical immediately before and after a sample gap of ≤ 3 days, we

interpolated the same score across the gap. If the score dropped from 3 to 2 and returned to 3

within 4 days, the intervening period at score 2 was considered part of a continuous MSP,

following previous studies (Douglas et al. 2016; Ryu et al. 2025). Over the study period, we

identified 48 MSPs from 13 adult females and 3 adolescent females (Table 2).

2.7 Estimation of female fertile windows

The relationship between ovulation timing and the maximal swelling phase (MSP) in female

bonobos has been examined using ovarian steroid profiles measured from urine or fecal samples.

Some studies suggest that ovulation can occur outside the MSP and that its timing is difficult to

predict from MSP onset or end (Reichert et al. 2002; Douglas et al. 2016). In contrast, other

work indicates that MSP end tends to occur a few days after ovulation and that this temporal

relationship is less variable (Heistermann et al. 1996; Ryu et al. 2025). In field studies of

non-human primates, the fertile phase is often operationally defined as a 4-day window

comprising the day of ovulation and the three preceding days (Deschner et al. 2003; Douglas et

al. 2016). This definition is biologically motivated by the short post-ovulatory viability of the

ovum (generally < 24 h) and evidence from human fertility studies indicating that most

conceptions result from intercourse occurring within a few days prior to ovulation, with

comparatively fer pregnancies attributable to sperm older than three days (Wilcox, Weinberg,

and Baird 1995; Dunson et al. 1999). Applying this criterion, Ryu et al. (2025) suggested that

the probability of sperm-ovum overlap (i.e. fertility) remains high from 7 to 2 days before

detumescence (peaking 4–5 days before detumescence) and then drops to zero by the day before

detumescence. Ryu et al. (2025) also demonstrated that male bonobos concentrated mating

effort during the late MSPs and largely ceased such effort after detumescence.

It was impractical to implement the intensive sampling necessary to pinpoint ovulation for

each female in parallel with full-day focal follows. Instead, we operationally defined the likely

fertile period (LFP) as the 6-day window from 7 to 2 days before MSP end in ovulatory cycles,

following Ryu et al. (2025), which was conducted at the same field site and using the same

visual assessment method of sexual swelling states. Nonetheless, MSPs do not necessarily

coincide with ovulation (Douglas et al. 2016; Ryu et al. 2025), and distinguishing from

ovulatory from anovulatory cycles based on visual records of swelling states alone is difficult.

To increase the validity of the LFP assignment, we applied the following criteria.

[1] Early adolescent females were excluded from LFP candidates. In chimpanzees,

adolescent females experience a prolong period of subfecund swelling cycles after

menarche, likely due to anovulatory cycles and early abortions associated with an

immature hypothalamic-pituitary-ovarian axis (Young and Yerkes 1943; Wallis 1997;

Nishida et al. 2003). Captive bonobos also reportedly reach menarche at 8.1 ± 0.6 years

($N = 3$) and experience 4.4 ± 2.6 years of adolescent subfecundity ($N = 5$)

(Thompson-Handler 1990). Accordingly, we classified 11 MSPs from 3 adolescent

females (Tp, Bc, Ih) younger than 11 years into anovulatory.

[2] Adult females in early lactation were excluded from LFP candidates. In the E1 group,

the mean interbirth interval is 57.7 ± 9.6 months (range = 43–72, $N = 23$) (Hashimoto

et al. 2022). However, recently, the shorter interbirth interval at 31 and 32 months has

been recorded at Wamba (unpublished data). Subtracting gestation length (7.9 months;

Furuichi et al. 1998) suggests her conception at approximately 23–24 months

postpartum. In addition, the earliest ovulation estimated from urinary hormone profiles

was confirmed to be 24 months after parturition from a female in E1 (Hashimoto et al.

2022). Based on these early resumption cases, we classified 6 MSPs from 5 adult

females (Yk, Nv, Ot, Jj, Aq) with infants younger 24 months into anovulatory. On the

other hand, ovulation may resume rapidly after infant loss: one female that lost her

infant gave birth 9 months later (Cheng et al. 2022), implying potential resumption

within about a month of infant death. This study included one female (Mz) losing her

infant

[3] MSPs of short duration were considered as anovulatory. At Wamba, MSP duration in

ovulatory cycles (mean = 20.7 $\pm$ 6.4 SD, range = 14–38, N = 14) appears longer than

in all cycles combined (mean = 14.0 ± 11.2 SD, range = 1–45, N = 36) (Ryu 2017). We

therefore classified 5 MSPs lasting <8 days (mean ovulatory MSP duration minus 2 SD,

based on Ryu 2017) into anovulatory.

[4] MSPs followed by an unusually short interval to the next MSP were considered as

anovulatory. In female bonobos, luteal-phase length, which reflected by progesterone

from the corpus luteum and associated detumescence, lasts approximately 9–13 days

(Heistermann et al. 1996: 13.0 ± 0.7 SD, N = 6; Douglas et al. 2016: 9.5 ± 1.2 SD, N =

6; Ryu 2017: 10.6 ± 0.6 SD, N = 3). Accordingly, we classified 2 MSPs with an interval

of <9 days between MSP and and the onset of the next MSP as of interval to next MSP

onset (mean luteal phase length minus 2 SD, based on Ryu 2017) into anovulatory.

Overall, we classified 20/48 MSPs as ovulatory cycles in 9 adult females (No, Ki, Hs, Jk,

Sl, Fk, Pf, Db, Mz) (Table 2). However, in 5 MSPs, the exact day of detumescence could not be

determined because it fell within a sample gap. When detumescence was obscured by a gap,

females were treated as having indeterminate LFP status during the 6 days of the MSP

preceding the gap, whereas earlier days in that MSP were classified as non-LFP. Copulation

with adult or adolescent males was observed at least once during 14/15 LFPs.

2.8 Data analyses

Statistical analyses were conducted in R version 4.4.2 (R Core Team 2024) using RStudio

version 12.0.467 (Posit Team 2024). We fitted generalized linear mixed models (GLMMs) to

account for repeated measures within individuals (Bolker et al. 2009) with the 'glmer' function

in the 'lme4' package (Bates et al. 2015). All continuous predictors were z-transformed (mean =

0, SD = 1) to improve the model convergence, performance, and interpretability (Schielzeth

2010). Variables included in each model are described below.

To test the first prediction—that male copulation rates differ little across dominance

ranks—we examined the effect of male dominance ranks on copulation rates. This analysis used

all-day focal data on males (345.6 hours across 41 days), excluding one old male (TW). Because

adult male chimpanzees, particularly high-ranking males, show lower sexual interest in

adolescent than adult females (Muller, Thompson, and Wrangham 2006; Reddy et al. 2021), we

tallied, for each focal day, the focal male's copulations with adult and adolescent females

separately. These daily counts were analyzed using two Poisson GLMMs using a log link, each

including an offset for observation time to account for variation in focal follow duration (Zuur

et al. 2009). Male rank (1–10) was included as a predictor of interest. In addition, the number of

non-mother females with maximal swellings present on each focal day (adults: 0–6;

adolescents: 0–4) was included as a control predictor to capture day-to-day variation in

copulation opportunities. A random intercept for focal male identity was included in both

models.

To test the second prediction—that higher-ranking males copulate more frequently with

LFP females than lower-ranking males—we examined the effect of female LFP status on the

probability of copulation with higher- versus lower-ranking males. All-day focal data on females

included 22 days (194.8 hours) during which at least one copulation event was observed. After

excluding 2 days (17.6 hours) in which the focal female's LFP status was indeterminate, we

aggregated for each remaining day (N = 20 days), counts of copulations with high-ranking

males (ranks 1–5) and low-ranking males (ranks 6–10). These paired counts were fitted as a

two-column response using a binomial GLMM with a logit link (Zuur et al. 2009). Female LFP

status (LFP: N = 4 days; non-LFP: N = 16) was included as a predictor of interest, with a

random intercept for focal female identity. Also, the ratio of numbers of high- versus

low-ranking males observed during female focal follows was also included as a control

predictor.

To reinforce the result of the analysis above at which males were roughly divided into two

rank classes (higher and lower), we additionally examined the effect of male dominance ranks

on the proportion of copulations with LFP versus non-LFP females using all copulation events

(N = 268) recorded through both focal and ad libitum sampling. Of them, we excluded 11

copulations involving females of indeterminate LFP status, we aggregated, for each male,

counts of copulations with LFP females (N = 59 events) and non-LFP females (N = 198 events).

These two-column responses (LFP, non-LFP) were analyzed using a binomial GLM with a logit

link (Zuur et al. 2009), with male rank (1–10) included as a predictor of interest.

To test the third prediction—that subordinate males selectively target females who are

less closely guarded by dominant males as mating partners—we assessed the effect of

within-party male dominance ranks on the probability of copulation within each one-hour time

window (OTW) during male focal follows. Specifically, we fitted three separate binomial

GLMMs to examine (i) whether higher-ranking males within the focal party were more likely to

copulate, (ii) whether higher-ranking males within the focal party were more likely to copulate

with LFP females when LFP females were present, and (iii) whether lower-ranking males within

the focal party were more likely to copulate with non-LFP females when LFP females were

present. Herein, we excluded OTWs in which the focal male was out of sight for >30 minutes

(N = 41), because the number of bonobos was reported to increase with the amount of

observation time until 30 minutes and then become saturated (Mulavwa et al. 2008). Also,

OTWs in which the focal male was resting for >95% of observation time (N = 19) were

excluded to eliminate datapoints where the focal male was almost inactive by lying on the bed.

For each OTW, copulation occurrence was coded as 1 ($\geq$1 copulation involving the focal male)

or 0 (on copulation) and analyzed using binomial GLMMs. Copulation spanning consecutive

OTWs were assigned to the earlier OTW based on the start time of the interaction. A random

intercept for focal male identity was included in all models. Datasets and predictors for each

model were as follows.

(i). The first model used 360 OTWs collected over 41 days, including 58 OTWs with

copulation and 302 without. The number of males higher-ranking than the focal male

within the party (0–7), representing the focal male's within-party rank position, was

included as the predictor of interest. The numbers of non-mother non-LFP females

with maximal swelling (0–7) and non-mother LFP females (0–2) in the party were

included as control predictors.

(ii). The second model used 117 OTWs collected over 17 days in which LFP females

were present, including 10 OTWs with copulation with LFP females and 107 without,

excluding 31 OTWs in which females of indeterminate LFP status were present.

Predictors included the numbers of males higher-ranking than the focal male (0–7),

non-mother non-LFP females with maximal swelling (0–7), and non-mother LFP

females (1–2) within the party.

(iii). The third model used 329 OTWs collected over 38 days, where the LFP status of all

females were determined, including 42 OTWs with copulation with non-LFP females

and 287 without. In addition to the numbers of males higher-ranking than the focal

male (0–7), non-mother non-LFP females with maximal swelling (0–7), and

non-mother LFP females (0–2), we included an interaction between the number of

higher-ranking males and the number of non-mother LFP females.

For model selection, we compared the goodness of fit of a single candidate model

containing predictors of interest with that of the corresponding null model retaining the control

predictor and random-effects structure (Bolker et al. 2009), following a null-hypothesis

significance testing framework (Nickerson 2000). Model comparisons were performed using

simulated likelihood ratio tests based on parametric bootstrap (1000 iterations) implemented

with the 'simulateLRT' function in the 'DHARMa' package (Hartig 2024). Fixed-effects

estimates were interpreted only when the candidate model significantly outperformed the null

model. Confidence intervals for fixed effects were obtained via parametric bootstrap inference

(2000 iterations) using the 'bootMer' function in the 'lme4' package. For each fitted model,

residual diagnostics, including normality, dispersion, and the presence of outliers, were

evaluated (Harrison et al. 2018) using the 'simulateResiduals' function in the 'DHARMa'

package (Hartig 2024); no violations were detected (Figures S2–S8). Residual independence

was assessed using the 'check_autocorrelation' function in the 'performance' package (Lüdecke

et al. 2021); no significant autocorrelation was detected (minimum $p = 0.218$). Multicollinearity

was evaluated with the 'check_collinearity' function in the same package (Lüdecke et al. 2021);

variance inflation factors (VIFs) were low in all models (maximum VIF = 2.11).

2.9 Ethical statement

The field studies of wild bonobos were approved by the Ministry of Scientific Research and the

Center for Research on Ecology and Forestry in the Democratic Republic of Congo. This study

conformed to the Animal Research Guidelines in The Graduate University for Advanced Studies,

Sokendai, Japan, revised on March 23, 2021.

3. Results

3.1 Relationship between male copulation rate and dominance rank

On the majority of male focal days (38 of 41 days, 92.7%), at least one non-mother female in

the party exhibited maximal swelling. In addition, at least one non-mother female in the likely

fertile period (LFP) was present in 17 male focal days (41%). The mean number of non-mother

females with maximal swellings per day was 4.1 ± 2.4 SD (2.4 ± 1.9 SD for adult females and

1.6 ± 1.2 SD for adolescent females, respectively). Copulations were observed on 33 of 38 days

(86.8%) at which maxima swelling females are present. Across males ($N = 10$), the mean

copulation rate was 0.20 ± 0.08 events per focal observation hour (range = 0.06–0.29, $N = 10$

males; Table 1), comprising 0.12 ± 0.06 events/h with adult females (range = 0–0.21) and $0.07 \pm$

0.05 events/h with adolescent females (range = 0–0.19). Of 83 male-male aggressive

interactions involving the focal male, only 5 cases (6.0%) occurred in mating contexts when

targets approached or solicited maximally tumescent females or immediately after copulation.

Poisson GLMMs testing the effect of male rank on copulation counts did not differ from

their respective null models for either adult females (log-likelihood ratio = 0.19, $p = 0.503$) or

adolescent females (log-likelihood ratio = 0.13, p = 0.623). These results indicate that male

copulation rates were not dependent on dominance rank, consistent with the first prediction that

male copulation rates are broadly comparable across ranks.

3.2 Male dominance rank and access to females in the likely fertile period (LFP)

Mating partners differed markedly according to the LFP status during female focal follows.

Among 15 copulations involving two LFP females, 13 events (86.7%) involved high-ranking

males (ranks 1–5), and 5 events (33.3%) involved the alpha male. In contrast, among 48

copulations involving nine non-LFP females, 28 events (41.7%) were with high-ranking males,

and only 2 events (4.2%) involved the alpha male.

A binomial GLMM examining the effect of female LFP status on the proportion of

copulations with high- versus low-ranking males significantly outperformed the null model

(log-likelihood ratio = 3.3, p = 0.014). Females in the LFP were therefore more likely to

copulate with high-ranking males than were non-LFP females (Table 3A; Figure 1), supporting

the second prediction that high-ranking males have priority access to LFP females.

An additional analysis of all copulation events for which female LFP status could be

estimated (N = 257) further supported this copulation pattern. A binomial GLM examining the

effect of male rank on the proportion of copulations with LFP versus non-LFP females

significantly outperformed the null model (log-likelihood ratio = 25.9, p < 0.001).

Higher-ranking males were more likely than lower-ranking males to copulate with LFP females

(Table 3B; Figure 2).

As for all copulation events, the median proportion of copulations with LFP females

across males was 9.3% (range = 0–54.8%, N = 10 males). The alpha male (KT) copulated with

LFP females in 54.8% of his copulations. His maternal brothers (KY and NB), sons of the alpha

female   (Ki) (Table 2), also showed higher proportions (51.4% and 35.7%, respectively) than

the across-male median. We observed Ki supported her sons in aggressive interactions with

other individuals than maternal brothers on four occasions; including one instance in which she

intercepted another male's copulation attempt. The dyad most frequently involved in aggressive

interactions was the maternal brothers, KT and KY (27 of 297 events [9.1%]), and all of them

were directed from the older (KT) to younger (KY) brothers. We also observed Ki intervened in

their aggressive interactions on six occasions and seemingly tried to stop the aggression of KT.

Among males whose mothers were not top-ranking, the third-ranking male (JO) copulated

with LFP females at a higher proportion (19.4%) than did four lower-ranking males with

resident mothers (HC, SE, YD, and NI; range = 3.7–10.0%). However, the mother of JO (Jk) did

not support her son in aggressive interactions with other males throughout the study period. Two

males without resident mothers (DI and GC) copulated with LFP females once or not at all.

Among all copulation events, sperm expulsion was confirmed in 6 of 59 copulations with

LFP females (10.2%) and in 8 of 162 with non-LFP females (4.9%). Ejaculation was therefore

observed approximately twice as frequently during copulations with LFP females as during

copulations with non-LFP females, although this difference was not statistically significant

(Fisher's Exact Test: $p = 0.209$).

3.3 Effects of within-party male dominance ranks on mating partners

One-hour time windows (OTW) during male focal follows ($N = 391$) consisted of 4.4 ± 2.4

males (range = 1–9) and 9.1 ± 4.1 females (0–19). The mean number of non-mother females

with maximal swellings was 2.9 ± 1.9 females (range = 0–9), consisting of 2.5 ± 1.7 non-LFP

and 0.4 ± 0.6 LFP females. Of these OTWs, at least one non-LFP female with maximal swelling

was present in 354 OTWs (90.5%), and at least one LFP female was present in 129 OTWs

(33.0%).

The first model examining the effect of the number of males higher ranking than the focal

male (i.e., within-party male rank) on the occurrence of copulations within one-hour time

windows (OTWs) did not outperform the null model (log-likelihood ratio = 1.6, $p = 0.075$). This

result indicates that within-party male rank alone did not affect the probability of copulation,

following the first prediction.

In contrast, the second model examining the effect of within-party male rank, expressed

as the number of males ranking higher than the focal male within the party, on the occurrence of

copulations with LFP females within OTWs significantly outperformed the null model

(log-likelihood ratio = 5.7, $p = 0.001$). Within-party higher-ranking males were more likely than

lower-ranking males to copulate with LFP females when such females were present in the party

(Table 3C; Figure 3), consistent with the second prediction.

Finally, the third model examining the interaction between within-party male rank and the

number of LFP females present on the occurrence of copulations with non-LFP females within

OTWs significantly outperformed the null model (log-likelihood ratio = 7.3, $p = 0.002$). The

significant interaction indicates that, as the number of LFP females increased, within-party

lower-ranking males were more likely than higher-ranking males to copulate with non-LFP

females (Table 3D; Figure 4). This pattern supports the third prediction that subordinate males

selectively target non-LFP females as mating partners when LFP females are available.

4. Discussion

4.1 Rank-independent male copulation frequency with an increased number of receptive females

All-day focal follows of male bonobos revealed no detectable relationship between male

dominance rank and copulation frequency with either adult or adolescent females. This finding

contrasts with prior reports of higher copulation rates for dominant males (Kano 1996;

Hohmann and Fruth 2003; Surbeck, Mundry, and Hohmann 2011; Yokoyama and Furuichi

2022; Ryu et al. 2025). One plausible explanation for this discrepancy is methodological

heterogeneity among studies. These previous studies were conducted under provisioning or

relied on short-duration focal follows (10–60 minutes). Such approaches can over-represent

centrally positioned individuals and larger parties, thereby missing copulations occurring at the

periphery or during small-party ranging (Altmann 1974; Drickamer 1974). This concern is

particularly relevant if subordinate males pursue mating opportunities while avoiding

interference by dominants (Furuichi and Ihobe 1994; Kano 1996; this study). Under these

condition, all-day focal follows are more likely to yield less biased estimates of mating activity

across ranks. Nevertheless, because our inference is based on six months of observations in a

single group, additional multi-group and long-term datasets using comparable all-day protocols

are needed to assess the generality of rank-independent copulation frequency.

Across group-living mammals, male dominance rank may predict mating success when

ovulation timing is temporally asynchronous among females and the operational sex ratio is

male-biased (i.e., when few females are receptive relative to the number of mature males)

(Emlen and Oring 1977; Clutton-Brock 1988). Such conditions can generate strong reproductive

skew if dominant males exclude subordinates from mating opportunities. In our study, however,

multiple receptive females were often simultaneously present (mean = 2.9 females per one-hour

time window), a context that likely dispersed male mating effort and reduced the scope for

rank-based monopolization. This interpretation is consistent with the expectation that female

reproductive synchrony weakens priority-of-access dynamics by increasing the number of

simultaneously receptive females (Bulger 1993; Ostner, Nunn, and Schulk 2008; Soulsbury

2011), and with evidence from seasonally breeding mammals in which high synchrony can

reduce mating skew (e.g., Japanese macaques, *Macaca fuscata*: Soltis, Thomsen, and Takenaka

2001, rhesus macaques, *Macaca mulatta*: Dubuc et al. 2011; Barbary macaques, *Macaca*

sylvanus: Bissonnette, Bischofberger, and van Schaik 2011; Tibetan macaques, *Macaca*

thibetana: Zhang et al. 2024, raccoons, *Procyon lotor*: Gehrt and Fritzell 1999; Soay sheep,

Ovis aries: Preston et al. 2005).

4.2 Rank-biased access to females in the likely fertile period and predictability of ovulation from

the visual signals

Despite the absence of rank effects on males' copulation frequencies, high-ranking males

accounted for a disproportionate share of copulations with females in the likely fertile period

(LFP), based on both female-based focal follows and ad libitum records. This pattern is

consistent with genetic evidence for pronounced male reproductive skew in bonobos (Surbeck

et al. 2017; Ishizuka et al. 2018; Mouginot, Cheng, and Wilson 2023) and with behavioral and

hormonal evidence that males can extract usable information about ovulation timing from

graded swelling cues and concentrate mating effort in periovulatory periods (Ryu et al. 2025).

Although overt, direct disruption of copulations by dominant males was infrequent in our

observations, dominants may nonetheless have constrained subordinates' access to LFP females

through close following and positional control. Comparable patterns have been reported in other

non-human primates in which females remain receptive across extended period that includes

infertile days, yet male mating effort is biased toward the periovulatory phase (e.g.,

chimpanzees: Deschner et al. 2004; mandrills, *Mandrillus sphinx*: Setchell, Charpentier, and

Wickings 2005; savannah baboons, *Papio cynocephalus*: Alberts, Buchan, and Altmann 2006;

sifakas, *Propithecus verreauxi*: Mass, Heistermann, and Kappeler 2009; rhesus macaques:

Dubuc et al. 2012; crested macaques, *Macaca nigra*: Higham et al. 2021). Together, these

findings underscore that variation in male reproductive success can be shaped more by access to

females within the fertile window than by total copulation counts.

In addition to male rank, maternal effects may have reinforced the rank bias in access to

LFP females in this group. The alpha male (KT) showed the highest proportion of copulations

with LFP females, and his maternal brothers (KY and NB), all sons of the alpha female (Ki),

also showed much higher proportions than the across-male median. We also observed Ki

supporting her sons during conflicts with other males, including one instance in which she

intercepted another male's copulation attempt. This pattern is consistent with evidence that the

presence of mothers, and particularly their dominance status, enhances sons' mating and

reproductive success in bonobos by increasing access to females and enabling direct maternal

intervention during male-male competition (Kano 1996; Furuichi 1997; 2011; Surbeck, Mundry,

and Hohmann 2011; Surbeck et al. 2019; Yokoyama and Furuichi 2022; Shibata and Furuichi

2024). In contrast, the elevated LFP access of these maternal brothers is unlikely to reflect

cooperative alliances among them, because aggressive interactions were most frequent within

this set of brothers and coalitionary support was rare in our observations. Taken together, these

results suggest that rank-biased access to females in the fertile window may be reinforced by

maternal support rather than by coalition formation among maternal brothers.

Because ejaculates are energetically costly to produce (Dewsbury 1982; Thomsen et al.

2006; Parker 2016), males are expected to allocate mating effort strategically by prioritizing

copulations with females in the periovulatory period while reducing ejaculatory investment in

non-conceptive copulations (e.g., Aujard et al. 1998; Li, Yin, and Zhou 2007; Heistermann et al.

2008; Garcia, Shimizu, and Huffman 2009). In contrast to a strictly deterministic allocation rule,

our observations indicate that even the alpha male bonobo copulated with non-LFP females in

nearly half of his observed copulation events, and ejaculation was occasionally confirmed

during copulations with non-LFP females. These results suggest that male bonobos' assessment

of female fertility is probabilistic rather than categorical. At the same time, our findings are

broadly consistent with conditional allocation: within parties, higher-ranking males were less

likely to copulate with non-LFP females when LFP females were present, and ejaculation was

observed approximately twice as often during copulations with LFP females as during

copulations with non-LFP females, although this difference was not statistically significant.

Taken together, these patterns suggest that male bonobos bias mating effort toward females

estimated to have higher conception probability while maintaining occasional ejaculatory

investment in females with lower conception probability, plausibly reflecting uncertainty in

fertility assessment rather than indiscriminate mating.

In this study, female fertility status was inferred from daily swelling scores by

operationally defining the LFP based on the empirical relationship between detumescence

timing and hormone-estimated ovulation reported by Ryu et al. (2025). This operational

definition yielded a rank-biased pattern of copulations with LFP females that is congruent with

paternity pattens reported by genetic studies of bonobos. Thus, the swelling-based proxy can be

useful for estimating the fertile window when intensive hormonal sampling is infeasible.

However, because swelling patterns can vary substantially in their relationship to ovulation in

bonobos, the LFP should be interpreted as an approximate indicator rather than a precise fertile

window.

4.3 Within-party rank dynamics and alternative mating strategies by subordinate males

Party-level analyses provide additional insight into how male bonobos allocate mating effort on

short temporal scales. Although within-party male dominance rank did not affect the probability

of copulation per se, it strongly influenced the fertility status of mating partners. Higher-ranking

males preferentially copulated with LFP females, whereas lower-ranking males increasingly copulated with non-LFP females as the number of LFP females in the party increased. This pattern suggests that subordinate males can secure mating opportunities by targeting females that are less likely to be guarded by dominant males. Such an alternative tactic may increase subordinates' participation in sperm competition, even if it rarely translates into paternity. Furthermore, over longer time horizons, avoiding direct contests with dominant males may also reduce injury risk and extend reproductive lifespan. This could be especially relevant in bonobos, where female dominance rank tends to increase with age and maternal rank can shape sons' future competitive prospects (Furuichi 1997; Tokuyama and Furuichi 2016; Toda and Furuichi 2020; Shibata and Furuichi 2023).

Comparable patterns have been reported in chimpanzees, where subordinate males maintain mating activity by shifting effort toward females that are less closely guarded by dominant males (Nishida 1997; Matsumoto-Oda 1999; Deschner et al. 2004). Unlike bonobos, however, subordinate chimpanzee males sire a non-trivial proportion of offspring (Constable et al. 2001; Boesch et al. 2006; Wroblewski et al. 2009; Newton-Fisher et al. 2010). One tactic subordinate males adopt is to form a temporary exclusive association with a given female by leading her away from other males and to mate with her repeatedly over several days during the fertile phase (i.e., consortship; Tutin 1979; Matsumoto-Oda 1999; Constable et al. 2001; Wroblewski et al. 2009; Newton-Fisher et al. 2010; Bray, Pusey, and Gilby 2016). The fission-fusion grouping pattern may afford subordinate males more opportunities to practice the consortship tactic, particularly in chimpanzee groups where females spend more time alone (e.g., Gombe), than in more spatially cohesive groups such as baboons and macaques (Wroblewski et al. 2009). However, consortship does not appear to play a major role in bonobo reproduction (Kano 1992; Takahata, Ihobe, and Idani 1999; Surbeck et al. 2017). In bonobos, the presence of

maximally tumescent females serves as a key driver increasing female-female gregariousness

(Surbeck et al. 2021). This social context may make it easier for males to locate and monitor

receptive females, potentially increasing the extent to which dominant males can monopolize

reproductive opportunities (Mouginot, Cheng, and Wilson 2023).

Even if the ultimate reproductive payoff of alternative tactics differs between species,

subordinate males in bonobos may have more frequent opportunities to obtain mating partners

than subordinate males in chimpanzees. because bonobo parties often include more receptive

females relative to the number of mature males (Furuichi and Hashimoto 2002; Surbeck et al.

2021). In our study, 92.6% of all-day focal follows of male bonobos (38 of 41 days) in the E1

group included at least one maximally tumescent female, which was more than twice the

percentage reported for all-day focal follows of male chimpanzees in the Kanyawara group

(38.8%; 33 of 85 days; Georgiev et al. 2014). In addition, the mean number of maximally

tumescent females present on a given day was substantially higher in E1 bonobos (4.1 ± 2.4 SD)

than Kanyawara chimpanzees (0.5 ± 0.1 SE). Although the E1 group during our study period

contained the largest number of females ever (Figure S1), these comparisons suggest that the

earlier resumption of sexual receptivity postpartum in female bonobos than chimpanzees may

provide mature males of all ranks with frequent mating opportunities.

4.4 Implications for female mate choice and potential benefits from prolonged sexual signaling

Together, our results indicate that pronounced reproductive skew among male bonobos is

shaped primarily through rank-biased access to fertile females, potentially reinforced by

maternal support and high female gregariousness. Female mate choice may also contribute to

this skew in bonobos, where females can ignore or refuse male courtship (Furuichi and

Hashimoto 2004; Hashimoto and Furuichi 2006) and sexual coercion appears to be substantially

limited (Paoli 2009). If male dominance rank covaries with aspects of phenotypic quality, such

as disease resistance, weapon traits, or behavioral dispositions, female preferences for dominant

males could yield indirect fitness benefits consistent with good genes models (Kirkpatrick 1996;

Møller and Alatalo 1999). Comparative evidence from other primates suggests that male rank

can correlate with traits plausibly linked to health condition or competitive ability, including

immune function and oxidative status (rhesus macaques: Georgiev et al. 2015), canine

dimensions (anubis baboons, *Papio anubis*: Galbany et al. 2015), and personality profiles

(chimpanzees: Weiss et al. 2023). Identifying whether dominance rank in male bonobos tracks

measurable indicators phenotypic quality will be important for evaluating the role of female

mate choice in generating male reproductive inequality.

Our findings also support the updated prolonged sexual receptivity hypothesis that a

higher number of receptive females relative to the number of mature males disperses male

mating effort across multiple females and thereby mitigates male-male contest competition over

mates. Females may benefit from weakening male-male contest competition, potentially

reducing risks linked to infanticide and/or sexual coercion. However, prolonged sexual

receptivity in female bonobos likely has functions beyond shaping male mating strategies.

Maximally tumescent females attract other females as well as males, increasing female-female

gregariousness (Surbeck et al. 2021) and eliciting affiliative interactions among females (Ryu,

Hill, and Furuichi 2015; Anzà, Demuru, and Palagi 2021). Moreover, female bonobos frequently

form coalitions during aggressive interactions and maintain high dominance relative to males

(Tokuyama and Furuichi 2016; Samuni and Surbeck 2023; Surbeck et al. 2025). Given that

strong social bonds among unrelated females is a central feature of bonobo society (Moscovice

et al. 2017; Furuichi 2023; Samuni and Surbeck 2023), prolonged sexual receptivity may have

been favored primarily because it promotes female-female affiliation and cooperation, with any

reduction in male-male contest competition emerging as a secondary consequence.

This study adds to evidence that female bonobos remain receptive for prolonged periods,

yet ovulation cues are sufficiently informative for males to bias mating effort toward the fertile

phase (Ryu et al. 2025) and for paternity to be strongly skewed toward dominant males

(Mouginot et al. 2024). This combination is not necessarily incompatible with a role of

prolonged sexual receptivity in paternity confusion. Even when paternity is concentrated among

dominant males, frequent non-conceptive copulations with multiple subordinates may still

create uncertainty about paternity and reduce incentives for aggression toward infants (Palombit

2015). In parallel, prolonged sexual receptivity can yield additional benefits by facilitating

socio-sexual interactions that strengthen affiliation and cooperation among unrelated females

(Ryu, Hill, and Furuichi 2015; Moscovice et al. 2019; Surbeck et al. 2021; Furuichi 2023). An

unresolved question is whether females obtain additional benefits from males through

non-conceptive copulations, such as increased tolerance around food resources. Although

anecdotal, bonobos have been reported to exchange sex for food (Kuroda 1984; Van

Krunkelsven et al. 1996; but Yamamoto 2015). Further evidence on the benefits females receive

through non-conceptive sexual interactions may help explain interspecific variation in the

duration and dynamics of sexual receptivity among anthropoid primates.

5. Acknowledgements

We thank the staff of the Centre de Recherche en Écologie et Foresterie (CREF), Democratic

Republic of the Congo, for supporting our fieldwork at Wamba in the Luo Scientific

Reserve—especially Messrs. Bahanande Iyokango, Yembo Bafaluka, Batsina Bambambe, and

Bakka Batuafe. We are also grateful to the local assistants who helped with our fieldwork and to

the residents of Wamba for their hospitality. Fieldwork by K. Toda was supported by

Grant-in-Aid for Scientific Research (21K13670) from the Japan Society for the Promotion of

Science and by the Canon Foundation in Europe–Kyoto University Japan–Africa Exchange

Fellowship.

6. References

Alberts, S. C., J. C. Buchan, and J. Altmann. 2006. "Sexual Selection in Wild Baboons: From

Mating Opportunities to Paternity Success." *Animal Behaviour* 72: 1177–1196.

https://doi.org/10.1016/j.anbehav.2006.05.001.

Alexander, R. D., K. M. Noonan. 1979. "Concealment of Ovulation, Parental Care, and Human

Social Evolution." In. *Evolutionary Biology and Human Social Behaviour: An*

Anthropological Perspective, edited by N. A. Chagnon, and W. G. Irons, 436–453, North

Scituate, Duxbury Press.

Altmann, J. 1974. "Observational Study of Behavior: Sampling Methods." *Behaviour* 49, no.

3–4 (1974): 227–266. https://doi.org/10.1163/156853974X00534

Anzà, S., E. Demuru, and E. Palagi. 2021. "Sex and Grooming as Exchange Commodities in

Female Bonobos' Daily Biological Market." *Scientific Reports*.

https://doi.org/10.1038/s41598-021-98894-w.

Arcadi, A. C., and R. W. Wrangham. 1999. "Infanticide in Chimpanzees: Review of Cases and a

New Within-Group Observation from the Kanyawara Study Group in Kibale National Park."

Primates 40, no. 2: 337–351.

Aujard, F., M. Heistermann, B. Thierry, and J. K. Hodges. 1998. "Functional Significance of

Behavioral, Morphological, and Endocrine Correlates Across the Ovarian Cycle in Semifree

Ranging Female Tonkean Macaques". *American Journal of Primatology* 46, no. 4:285–309.

https://doi.org/10.1002/(SICI)1098-2345(1998)46:4<285::AID-AJP2>3.0.CO;2-8

Balolia, K. L., K. Baughan, and J. S. Massey. 2025. "Relative Facial Width, and Its Association

With Canine Size and Body Mass Among Chimpanzees and Bonobos: Implications for

Understanding Facial Width-to-Height Ratio Expression Among Human Populations."

American Journal of Biological Anthropology 186, no. 1: e25040.

https://doi.org/10.1002/ajpa.25040.

Barelli, C., M. Heistermann, C. Boesch, and U. H. Reichard. 2008. "Mating Patterns and Sexual

Swellings in Pair-Living and Multimale Groups of Wild White-Handed Gibbons, *Hylobates*

lar." *Animal Behaviour* 75, no. 3: 991–1001. https://doi.org/10.1016/j.anbehav.2007.08.012.

Bates, D., M. Martin, B. Bolker, and S. Walker. 2015. "Fitting Linear Mixed-Effects Models

Using lme4." *Journal of statistical software* 67: 1–48. https://doi.org/10.18637/jss.v067.i01

Behringer, V., T. Deschner, C. Deimel, J. M. G. Stevens, and G. Hohmann. 2014. "Age-Related

Changes in Urinary Testosterone Levels Suggest Differences in Puberty Onset and Divergent

Life History Strategies in Bonobos and Chimpanzees." *Hormones and Behavior* 66: 525–533.

https://doi.org/10.1016/j.yhbeh.2014.07.011.

Berghaenel, A., J. M. G. Stevens, G. Hohmann, T. Deschner, and V. Behringer. 2023. "Evidence

for Adolescent Length Growth Spurts in Bonobos and Other Primates Highlights the

Importance of Scaling Laws." *eLife* 12: e86635. https://doi.org/10.7554/eLife.86635.

Bissonnette, A., N. Bischofberger, and C. P. van Schaik. 2011. "Mating Skew in Barbary

Macaque Males: The Role of Female Mating Synchrony, Female Behavior, and Male–Male

Coalitions." *Behavioral Ecology and Sociobiology* 65, no. 1: 167–182.

https://doi.org/10.1007/s00265-010-1023-z.

Boesch, C., G. Kohou, H. Néné, and L. Vigilant. 2006. "Male Competition and Paternity in Wild

Chimpanzees of the Taï Forest." *American Journal of Physical Anthropology* 130, no. 1:

103–115. https://doi.org/10.1002/ajpa.20341.

Boesch, C., C. Crockford, I. Herbinger, R. Wittig, Y. Moebius, and E. Normand. 2008.

"Intergroup Conflicts Among Chimpanzees in Taï National Park: Lethal Violence and the

Female Perspective." *American Journal of Primatology* 70, no. 6: 519–532.

https://doi.org/10.1002/ajp.20524.

Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.

S. S. White. 2009. "Generalized Linear Mixed Models: A Practical Guide for Ecology and

Evolution". *Trends in Ecology & Evolution 24, no.* 3:127–135.

https://doi.org/10.1016/j.tree.2008.10.008

Brauch, K., D. Pfefferle, K. Hodges, U. Möhle, J. Fischer, and M. Heistermann. 2007. "Female

Sexual Behavior and Sexual Swelling Size as Potential Cues for Males to Discern the Female

Fertile Phase in Free-Ranging Barbary Macaques (*Macaca sylvanus*) of Gibraltar." *Hormones

and Behavior* 52, no. 3: 375–383. https://doi.org/10.1016/j.yhbeh.2007.06.001.

Bray, J., A. E. Pusey, and I. C. Gilby. 2016. "Incomplete Control and Concessions Explain

Mating Skew in Male Chimpanzees." *Proceedings of the Royal Society B: Biological

Sciences* 283, no. 1842: 20162071. https://doi.org/10.1098/rspb.2016.2071.

Bulger, J. B. 1993. "Dominance Rank and Access to Estrous Females in Male Savanna

Baboons." *Behaviour* 127, no. 1/2: 67–103.

Cheng, L., A. Shaw, A. Menante, L. Samuni, and M. Surbeck. 2022. "Social Relationships

Between Female Bonobos (*Pan paniscus*) in the Kokolopori Bonobo Reserve, Democratic

Republic of Congo." *Primates*. https://doi.org/10.1007/s10329-022-00986-2.

Clutton-Brock, T. H. 1988. *Reproductive Success: Studies of Individual Variation in Contrasting

Breeding Systems*. Chicago, University of Chicago Press.

Conaway, C. H. 1971. "Ecological Adaptation and Mammalian Reproduction". *Biology of

Reproduction* 4, no. 3:239–247.

Constable, J. L., M. V. Ashley, J. Goodall, and A. E. Pusey. 2001. "Noninvasive Paternity

Assignment in Gombe Chimpanzees." *Molecular Ecology* 10: 1279–1300.

https://doi.org/10.1046/j.1365-294X.2001.01262.x.

David, H. A. 1987. "Ranking From Unbalanced Paired-Comparison Data". *Biometrika* 74, no.

2:432–436. https://doi.org/10.1093/biomet/74.2.432

de Vries, H. 1995. "An Improved Test of Linearity in Dominance Hierarchies Containing

Unknown or Tied Relationships." *Animal Behaviour* 50: 1375–1389.

de Vries, H. 1998. "Finding a Dominance Order Most Consistent with a Linear Hierarchy: A

New Procedure and Review." *Animal Behaviour* 55: 827–843.

de Vries, H., J. M. Stevens, and H Vervaecke. 2006. "Measuring and Testing the Steepness of

Dominance Hierarchies". *Animal Behaviour* 71, no. 3:585–592.

https://doi.org/10.1016/j.anbehav.2005.05.015

Deschner, T., C. Boesch. 2007. "Can the Patterns of Sexual Swelling Cycles in Female Taï

Chimpanzees be Explained by the Cost-Of-Sexual-Attraction Hypothesis?" *International*

Journal of Primatology 28:389–406. https://doi.org/10.1007/s10764-007-9120-1

Deschner, T., M. Heistermann, K. Hodges, and C. Boesch. 2004. "Female Sexual Swelling Size,

Timing of Ovulation, and Male Behavior in Wild West African Chimpanzees." *Hormones and*

Behavior 46: 204–215. https://doi.org/10.1016/j.yhbeh.2004.03.013.

Deschner, T., M. Heistermann, K. Hodges, and C. Boesch. 2003. "Timing and Probability of

Ovulation in Relation to Sex Skin Swelling in Wild West African Chimpanzees, *Pan*

troglodytes verus". *Animal Behaviour* 66, no. 3:551–560.

https://doi.org/10.1006/anbe.2003.2210

Dewsbury, D. A. 1982. "Ejaculate Cost and Male Choice." *The American Naturalist* 119, no. 5:

601–610.

Dixson, A. 1998. *Primate Sexuality: Comparative Studies of the Prosimians, Monkeys, Apes,*

and Humans. New York, Oxford University Press.

Douglas, P. H., G. Hohmann, R. Murtagh, R. Thiessen-Bock, and T. Deschner. 2016. "Mixed

Messages: Wild Female Bonobos Show High Variability in the Timing of Ovulation in

Relation to Sexual Swelling Patterns". *BMC Evolutionary Biology* 16:1–17.

https://doi.org/10.1186/s12862-016-0691-3

Drickamer, L. C. 1974. "Social Rank, Observability, and Sexual Behaviour of Rhesus Monkeys

(*Macaca mulatta*)." *Reproduction* 37, no. 1:117–120. https://doi.org/10.1530/jrf.0.0370117

Dubuc, C., L. Muniz, M. Heistermann, A. Engelhardt, and A. Widdig. 2011. "Testing the

Priority-of-Access Model in a Seasonally Breeding Primate Species." *Behavioral Ecology*

and Sociobiology 65: 1615–1627.

Dubuc, C., L. Muniz, M. Heistermann, A. Widdig, and A. Engelhardt. 2012. "Do Males Time

Their Mate-Guarding Effort With the Fertile Phase in Order to Secure Fertilisation in Cayo

Santiago Rhesus Macaques?" *Hormones and Behavior* 61: 696–705.

https://doi.org/10.1016/j.yhbeh.2012.03.003.

Dubuc, C., L. J. N. Brent, A. K. Accamando, M. S. Gerald, A. MacLarnon, S. Semple, M.

Heistermann, and A. Engelhardt. 2009. "Sexual Skin Color Contains Information About the

Timing of the Fertile Phase in Free-ranging *Macaca mulatta*." *International Journal of*

Primatology 30: 777–789. https://doi.org/10.1007/s10764-009-9369-7.

Dunson, D. B., D. D. Baird, A. J. Wilcox, and C. R. Weinberg. 1999. "Day-Specific

Probabilities of Clinical Pregnancy Based on Two Studies With Imperfect Measures of

Ovulation". *Human Reproduction* 14, no. 7:1835–1839.

https://doi.org/10.1093/humrep/14.7.1835

Emery Thompson, M., and K. Sabbi. 2024. "Evolutionary Demography of the Great Apes". In

Human Evolutionary Demography, edited by O, Burger, R. Lee, and R. Sear, 423–474,

Cambridge UK, Open Book Publishers. https://doi.org/10.11647/obp.0251.19

Emery Thompson, M., and P. T. Ellison. 2017. "Fertility and Fecundity". In *Chimpanzees and*

Human Evolution, edited by M. N. Muller, R. W. Wrangham, and D. Pilbeam, 217–258,

Cambridge, MA, Harvard University Press. doi.org/10.4159/9780674982642-006

Emlen, S. T., and L. W. Oring. 1977. "Ecology, Sexual Selection, and the Evolution of Mating

Systems." *Science* 197: 215–223.

Engelhardt, A., J.-B. Pfeifer, M. Heistermann, C. Niemitz, J. A. R. A. M. van Hooff, and J. K.

Hodges. 2004. "Assessment of Female Reproductive Status by Male Longtailed Macaques,

Macaca fascicularis, Under Natural Conditions." *Animal Behaviour* 67: 915–924.

Fawcett, K. A., and G. Muhumuza. 2000. "Death of a Wild Chimpanzee Community Member."

American Journal of Primatology 51, no. 4: 243–247.

https://doi.org/10.1002/1098-2345(200008)51:4<243::AID-AJP3>3.0.CO;2-P.

Fernandez-Duque, E., K. Burke, K. Schoenrock, C. K. Wolovich, and C. R. Valeggia. 2011.

"Hormonal Monitoring of Reproductive Status in Monogamous Wild Female Owl Monkeys

(*Aotus azarai*) of the Argentinean Chaco." *Folia Primatologica* 82: 143–153.

https://doi.org/10.1159/000333081.

Fernandez-Duque, E., M. Huck, S. Van Belle, and A. Di Fiore. 2020. "The Evolution of

Pair-Living, Sexual Monogamy, and Cooperative Infant Care: Insights From Research on

Wild Owl Monkeys, Titis, Sakis, and Tamarins." *Yearbook of Physical Anthropology* 171:

1–56. https://doi.org/10.1002/ajpa.24017.

Furuichi, T. 1987. "Sexual Swelling, Receptivity, and Grouping of Wild Pygmy Chimpanzee

Females at Wamba, Zaire". *Primates* 28:309–318. https://doi.org/10.1007/BF02381014

Furuichi, T. 1997. "Agonistic Interactions and Matrifocal Dominance Rank of Wild Bonobos

(*Pan paniscus*) at Wamba". *International Journal of Primatology* 18:855–875.

https://doi.org/10.1023/A:1026327627943

Furuichi, T. 2011. "Female Contributions to the Peaceful Nature of Bonobo

Society". *Evolutionary Anthropology: Issues, News, and Reviews* 20, no. 4:131–142.

https://doi.org/10.1002/evan.20308

Furuichi, T. 2023. "Prolonged Sexual Receptivity in Females and Its Impact on the Evolution of

Bonobos". In *Bonobos and People at Wamba: 50 years of Research,* edited by T. Furuichi, G.

Idani, D. Kimura, H. Ihobe, and C. Hashimoto, 545–564. Singapore, Springer Nature.

https://doi.org/10.1007/978-981-99-4788-1_39

Furuichi, T., and C. Hashimoto. 2002. "Why Female Bonobos Have a Lower Copulation Rate

during Estrus than Chimpanzees". In *Behavioural Diversity in Chimpanzees and Bonobos*,

edited by C Boesch, and G. Hohmann, 156–167, Cambridge, Cambridge University Press.

https://doi.org/10.1017/CBO9780511606397

Furuichi, T., and C. Hashimoto. 2004. "Sex Differences in Copulation Attempts in Wild

Bonobos at Wamba." *Primates* 45: 59–62. https://doi.org/10.1007/s10329-003-0055-7.

Furuichi, T. and H. Ihobe. 1994. "Variation in Male Relationships in Bonobos and

Chimpanzees". *Behaviour* 130, no. 3–4:211–228. https://doi.org/10.1163/156853994X00532

Furuichi, T., G. Idani, I. Ihobe, et al. 1998. "Population Dynamics of Wild Bonobos (*Pan

paniscus*) at Wamba". *International Journal of Primatology* 19:1029–1043.

https://doi.org/10.1023/A:1020326304074

Furuichi, T., G. Idani, and I. Ihobe, et al. 2012. "Long-Term Studies on Wild Bonobos at Wamba,

Luo Scientific Reserve, DR Congo: Towards the Understanding of Female Life History in a

Male-Philopatric Species". In *Long-Term Field Studies of Primates*, edited by P. Kappeler, D.

Watts, 413–433, Berlin, Heidelberg, Springer. https://doi.org/10.1007/978-3-642-22514-7_18

Galbany, J., J. Tung, J. Altmann, and S. C. Alberts. 2015. "Canine Length in Wild Male

Baboons: Maturation, Aging and Social Dominance Rank." *PLoS ONE* 10, no. 5: e0126415.

https://doi.org/10.1371/journal.pone.0126415.

Garcia, C., K. Shimizu, and M. Huffman. 2009. "Relationship Between Sexual Interactions and

the Timing of the Fertile Phase in Captive Female Japanese Macaques (*Macaca*

fuscata)". *American Journal of Primatology* 71, no. 10:868–879.

https://doi.org/10.1002/ajp.20717

Gehrt, S. D., and E. K. Fritzell. 1999. "Raccoon Mating System." Animal Behaviour 57, no. 3:

601.

Georgiev, A. V., A. F. Russell, M. Emery Thompson, E. Otali, M. N. Muller, and R. W.

Wrangham. 2014. "The Foraging Costs of Mating Effort in Male Chimpanzees (*Pan*

troglodytes schweinfurthii)." *International Journal of Primatology* 35: 725–745.

https://doi.org/10.1007/s10764-014-9788-y.

Georgiev, A. V., M. P. Muehlenbein, S. P. Prall, M. Emery Thompson, and D. Maestripieri. 2015.

"Male Quality, Dominance Rank, and Mating Success in Free-ranging Rhesus Macaques."

Behavioral Ecology 26, no. 3: 763-772. https://doi.org/10.1093/beheco/arv008.

Gesquiere, L. R., E. O. Wango, S. C. Alberts, and J. Altmann. 2007. "Mechanisms of Sexual

Selection: Sexual Swellings and Estrogen Concentrations as Fertility Indicators and Cues for

Male Consort Decisions in Wild Baboons." *Hormones and Behavior* 51: 114–125.

https://doi.org/10.1016/j.yhbeh.2006.08.010.

Graham, C. E., D. C. Collins, H. Robinson, and J. R. K. Preedy. 1972. "Urinary Levels of

Estrogens and Pregnanediol and Plasma Levels of Progesterone During the Menstrual Cycle

of the Chimpanzee: Relationship to the Sexual Swelling". *Endocrinology* 91, no 1:13–24.

https://doi.org/10.1210/endo-91-1-13

Harrison, X. A., L. Donaldson, M. E. Correa-Cano, J. Evans, D. N. Fisher, C. E. D. Goodwin, B.

S. Robinson, D. J. Hodgson, and R. Inger. 2018. "A Brief Introduction to Mixed Effects

Modelling and Multi-Model Inference in Ecology." *PeerJ* 6: e4794.

https://doi.org/10.7717/peerj.4794.

Hartig, F. 2024. "DHARMa: Residual Diagnostics for Hierarchical (Multi-Level/Mixed)

Regression Models. R package version 0.4.7".

https://CRAN.R-project.org/package=DHARMa

Hashimoto, C. 1997. "Context and Development of Sexual Behavior of Wild Bonobos (*Pan

paniscus*) at Wamba, Zaire". *International Journal of Primatology* 18:1–21.

https://doi.org/10.1023/A:1026384922066

Hashimoto, C., H. Ryu, K. Mouri, K. Shimizu, T. Sakamaki, and T. Furuichi. 2022. "Physical,

Behavioral, and Hormonal Changes in the Resumption of Sexual Receptivity during

Postpartum Infertility in Female Bonobos at Wamba". *Primates* 63, no. 2:109–121.

https://doi.org/10.1007/s10329-021-00968-w

Hashimoto, C., and T. Furuichi. 2006. "Comparison of Behavioral Sequence of Copulation

Between Chimpanzees and Bonobos." *Primates* 47: 342–345.

https://doi.org/10.1007/s10329-005-0144-x.

Hashimoto, C., T. Furuichi, Y. Tashiro 2001. "What Factors Affect the Size of Chimpanzee

Parties in the Kalinzu Forest, Uganda? Examination of Fruit Abundance and Number of

Estrous Females." *International Journal of Primatology* 22, no. 6:947–959.

https://doi.org/10.1023/A:1012061504420

Heistermann, M., U. Möhle, H. Vervaecke, L. van Elsacker, and J. K. Hodges. 1996.

"Application of Urinary and Fecal Steroid Measurements for Monitoring Ovarian Function

and Pregnancy in the Bonobo (*Pan paniscus*) and Evaluation of Perineal Swelling Patterns in

Relation to Endocrine Events". *Biology of Reproduction* 55, no. 4:844–853.

Heistermann, M., K. Brauch, U. Möhle, D. Pfefferle, J. Dittami, and K. Hodges. 2008. "Female
Ovarian Cycle Phase Affects the Timing of Male Sexual Activity in Free‐Ranging Barbary
macaques (*Macaca sylvanus*) of Gibraltar". *American Journal of Primatology* 70. no. 1.
44–53. https://doi.org/10.1002/ajp.20455

Higham, J. P., S. Semple, A. MacLarnon, M. Heistermann, and C. Ross. 2009. "Female
Reproductive Signaling, and Male Mating Behavior, in the Olive Baboon." *Hormones and
Behavior* 55: 60–67. https://doi.org/10.1016/j.yhbeh.2008.08.007.

Higham, J. P., M. Heistermann, M. Agil, D. Perwitasari-Farajallah, A. Widdig, and A.
Engelhardt. 2021. "Female Fertile Phase Synchrony, and Male Mating and Reproductive
Skew, in the Crested Macaque." *Scientific Reports*.

Hohmann, G., and B. Fruth. 2003. "Intra- and Inter-Sexual Aggression by Bonobos in the
Context of Mating". *Behaviour* 140, no. 11/12. 1389–1413.
https://www.jstor.org/stable/4536099

Ishizuka, S., Y. Kawamoto, T. Sakamaki, et al. 2018. "Paternity and Kin Structure Among
Neighbouring Groups in Wild Bonobos at Wamba". *Royal Society Open Science* 5, no.1:
171006. https://doi.org/10.1098/rsos.171006

Jurmain, R. 1997. "Skeletal Evidence of Trauma in African Apes, With Special Reference to the
Gombe Chimpanzees". *Primates* 38, no. 1: 1–14. https://doi.org/10.1007/BF02385918

Kaburu, S. S. K., S. Inoue, and N. E. Newton-Fisher. 2013. "Death of the Alpha:
Within-Community Lethal Violence Among Chimpanzees of the Mahale Mountains National
Park." *American Journal of Primatology* 9999: 1–9. https://doi.org/10.1002/ajp.22135.

Kano, T. 1992. *The Last Ape. Pygmy Chimpanzee Behavior and Ecology*. Stanford, Stanford
University Press.

Kano, T. 1996. "Male Rank Order and Copulation Rate in a Unit-Group of Bonobos at Wamba,

Zaire". In *Great Ape Societies*, edited by W. C. McGrew, L. F. Marchant, and T. Nishida,

135–145. Cambridge, Cambridge University Press.

https://doi.org/10.1017/CBO9780511752414.012

Kelley, J. 1995. "Sexual Dimorphism in Canine Shape Among Extant Great Apes." *American*

Journal of Physical Anthropology 96: 365–389.

Kendrick, K. M., and A. F. Dixson. 1983. "The Effect of the Ovarian Cycle on the Sexual

Behaviour of the Common Marmoset (*Callithrix jacchus*)." *Physiology & Behavior* 30, no. 5:

735–742.

Kirkpatrick, M. 1996. "Good Genes and Direct Selection in the Evolution of Mating

Preferences." *Evolution* 50, no. 6: 2125–2140.

Kunz, J. A., G. J. Duvot, M. A. van Noordwijk, E. P. Willems, M. Townsend, N. Mardianah, S.

S. U. Atmoko, E. R. Vogel, T. P. Nugraha, M. Heistermann, M. Agil, T. Weingrill, and C. P.

van Schaik. 2021. "The Cost of Associating With Males for Bornean and Sumatran Female

Orangutans: A Hidden Form of Sexual Conflict?" *Behavioral Ecology and Sociobiology* 75: 6.

https://doi.org/10.1007/s00265-020-02948-4.

Kuroda, S. 1979. "Grouping of the Pygmy Chimpanzees". *Primates* 20:161–183.

https://doi.org/10.1007/BF02373371

Kuroda, S. 1984. "Interaction Over Food Among Pygmy Chimpanzees." In *The Pygmy*

Chimpanzee, edited by R. L. Susman, 301–324. New York: Plenum Press.

Kvarnemo, C., I . Ahnesjo. 1996. "The Dynamics of Operational Sex Ratios and Competition

for Mates". *Trends in Ecology & Evolution* 11, no. 10:404–408.

https://doi.org/10.1016/0169-5347(96)10056-2

Li, J., H. Yin, and L. Zhou. 2007. "Non-Reproductive Copulation Behavior Among Tibetan

Macaques (*Macaca thibetana*) at Huangshan, China". *Primates* 48:64–72.

https://doi.org/10.1007/s10329-006-0002-5

Lowe, A. E., C. Hobaiter, C. Asiimwe, K. Zuberbühler, and N. E. Newton-Fisher. 2020.

"Intra-Community Infanticide in Wild, Eastern Chimpanzees: A 24-Year Review." *Primates*

61: 69–82. https://doi.org/10.1007/s10329-019-00730-3.

Lüdecke, D., M. S. Ben-Shachar, I. Patil, P. Waggoner, and D. Makowski. 2021. "performance:

An R Package for Assessment, Comparison and Testing of Statistical Models". *Journal of*

Open Source Software 6, no.60:3139. https://doi.org/10.21105/joss.03139

Mass, V., M. Heistermann, and P. M. Kappeler. 2009. "Mate-Guarding as a Male Reproductive

Tactic in *Propithecus verreauxi*." *International Journal of Primatology* 30: 389–409.

https://doi.org/10.1007/s10764-009-9345-2.

Matsumoto-Oda, A. 1999. "Female Choice in the Opportunistic Mating of Wild Chimpanzees

(*Pan troglodytes schweinfurthii*) at Mahale." *Behavioral Ecology and Sociobiology* 46:

258–266.

Mitani, J. C., D. P. Watts, and S. J. Amsler. 2010. "Lethal Intergroup Aggression Leads to

Territorial Expansion in Wild Chimpanzees." *Current Biology* 20, no. 12: R507–R508.Møller,

A. P., and R. V. Alatalo. 1999. "Good-Genes Effects in Sexual Selection." *Proceedings of the*

Royal Society of London. Series B: Biological Sciences 266: 85–91.

Moscovice, L. R., M. Surbeck, B. Fruth, G. Hohmann, A. V. Jaeggi, and T. Deschner. 2019.

"The Cooperative Sex: Sexual Interactions Among Female Bonobos Are Linked to Increases

in Oxytocin, Proximity and Coalitions." *Hormones and Behavior* 116: 104581.

https://doi.org/10.1016/j.yhbeh.2019.104581.

Moscovice, L. R., P. H. Douglas, L. Martinez-Iñigo, M. Surbeck, L, Vigilant, G. Hohmann.

2017. "Stable and Fluctuating Social Preferences and Implications for Cooperation Among

Female Bonobos at LuiKotale, Salonga National Park, DRC". *American Journal of Physical Anthropology* 163, no. 1:158–172. https://doi.org/10.1002/ajpa.23197

Mouginot, M., L. Cheng, M. L. Wilson. 2023. "Reproductive Inequality Among Males in the Genus *Pan*". *Philosophical Transactions of the Royal Society B* 378, no. 1883:20220301. https://doi.org/10.1098/rstb.2022.0301

Mouginot, M., M. L. Wilson, N. Desai, and M. Surbeck. 2024. "Differences in Expression of Male Aggression Between Wild Bonobos and Chimpanzees". *Current Biology* 34, No.8:1780–1785. https://doi.org/10.1016/j.cub.2024.02.071

Mulavwa, M., T. Furuichi, K. Yangozene, et al. 2008. "Seasonal Changes in Fruit Production and Party Size of Bonobos at Wamba". In *The Bonobos: Behavior, Ecology, and Conservation*, edited by T. Furuichi, and J. Thompson, 121–134. New York, Springer. https://doi.org/10.1007/978-0-387-74787-3_7

Muller, M. N., M. E. Thompson, S. M. Kahlenberg, and R. W. Wrangham. 2011. "Sexual Coercion by Male Chimpanzees Shows That Female Choice May Be More Apparent Than Real." *Behavioral Ecology and Sociobiology* 65: 921–933. https://doi.org/10.1007/s00265-010-1093-y.

Muller, M. N., M. E. Thompson, and R. W. Wrangham. 2006. "Male Chimpanzees Prefer Mating With Old Females". *Current Biology* 16, no.22:2234–2238. https://doi.org/10.1016/j.cub.2006.09.042

Neumann, C., and L. Kulik. 2024. "Package 'EloRating': Animal Dominance Hierarchies by Elo Rating." R package version 0.46.18. https://github.com/gobbios/EloRating.

Newton-Fisher, N. E., M. Emery Thompson, V. Reynolds, C. Boesch, and L. Vigilant. 2010. "Paternity and Social Rank in Wild Chimpanzees (*Pan troglodytes*) From the Budongo Forest, Uganda." *American Journal of Physical Anthropology* 142: 417–428.

https://doi.org/10.1002/ajpa.21241.

Nickerson, R. S. 2000. "Null Hypothesis Significance Testing: A Review of an Old and

Continuing Controversy." *Psychological Methods* 5, no. 2: 241–301.

https://doi.org/10.1037//1082-989X.5.2.241.

Nishida, T. 1968. "The Social Group of Wild Chimpanzees in the Mahali

Mountains". *Primates* 9: 167–224. https://doi.org/10.1007/BF01730971

Nishida, T. 1997. "Sexual Behavior of Adult Male Chimpanzees of the Mahale Mountains

National Park, Tanzania." *Primates* 38: 379–398

Nishida, T., N. Corp, M. Hamai, et al. 2003. "Demography, Female Life History, and

Reproductive Profiles Among the Chimpanzees of Mahale". *American Journal of*

Primatology 59, no.3:99–121. https://doi.org/10.1537/ase.107.141

Nishida, T., T. Kano, J. Goodall, W. C. McGrew, and M. Nakamura. 1999. "Ethogram and

Ethnography of Mahale Chimpanzees." *Anthropological Science* 107, no. 2: 141–188.

Ostner, J., C. L. Nunn, and O. Schülke. 2008. "Female Reproductive Synchrony Predicts

Skewed Paternity Across Primates." *Behavioral Ecology* 19, no. 6: 1150–1158.

https://doi.org/10.1093/beheco/arn093.

1145

Palombit, R. A. 2015. "Infanticide as Sexual Conflict: Coevolution of Male Strategies and

Female Counterstrategies." *Cold Spring Harbor Perspectives in Biology* 7: a017640.

https://doi.org/10.1101/cshperspect.a017640.

Paoli, T. 2009. "The Absence of Sexual Coercion in Bonobos". In *Sexual Coercion in Primates*

and Humans: An Evolutionary Perspective on Male Aggression against Females, edited by M.

N. Muller, and R. W. Wrangham, 410–423, Cambridge, Harvard University Press

doi.org/10.4159/9780674054349-016

Parker, G. A. 2016. "The Evolution of Expenditure on Testes." *Journal of Zoology* 298: 3–19.

https://doi.org/10.1111/jzo.12297.

Pashchevskaya, S., B. Fruth, S. L. Kjærland, L. Berton, L. Vigilant, and G. Hohmann. 2025.

"Coalitionary Intra-Group Aggression by Wild Female Bonobos." *Current Biology*.

https://doi.org/10.1016/j.cub.2025.08.010.

Pawłowski, B. 1999. "Loss of Oestrus and Concealed Ovulation in Human Evolution: The Case

Against the Sexual-Selection Hypothesis." *Current Anthropology* 40, no. 3: 257–275.

https://doi.org/10.1086/200017.

Phillips-Conroy, J. E., C. J. Jolly, B. Petros, J. S. Allan, and R. C. Desrosiers. 1994. "Sexual

Transmission of SIVagm in Wild Grivet Monkeys." *Journal of Medical Primatology* 23: 1–7.

Plavcan, J. M., and C. P. van Schaik. 1992. "Intrasexual Competition and Canine Dimorphism in

Anthropoid Primates." *American Journal of Physical Anthropology* 87: 461–477.

Plavcan, J. M., C. P. van Schaik, and P. M. Kappeler. 1995. "Competition, Coalitions and

Canine Size in Primates." *Journal of Human Evolution* 28: 245–276.

Posit Team. 2024. *RStudio: Integrated Development Environment for R*. Posit Software, PBC.

http://www.posit.co/

Preston, B. T., I. R. Stevenson, J. M. Pemberton, D. W. Coltman, and K. Wilson. 2005. "Male

Mate Choice Influences Female Promiscuity in Soay Sheep." *Proceedings of the Royal*

Society B: Biological Sciences 272. https://doi.org/10.1098/rspb.2004.2977.

R Core Team. 2024. R: *A Language and Environment for Statistical Computing*. R Foundation

for Statistical Computing. https://www.R-project.org/

Reddy, R. B., K. E. Langergraber, A. A. Sandel, L. Vigilant, and J. C. Mitani. 2021. "The

Development of Affiliative and Coercive Reproductive Tactics in Male Chimpanzees."

Proceedings of the Royal Society B: Biological Sciences 288: 20202679.

https://doi.org/10.1098/rspb.2020.2679.

1178

Reichert, K. E., M. Heistermann, J. K. Hodges, and G. Hohmann. 2002. "What Females Tell

Males About Their Reproductive Status: Are Morphological and Behavioural Cues Reliable

Signals of Ovulation in Bonobos (*Pan paniscus*)?" *Ethology* 108: 583–600.

Rodríguez-Gironés, M. A., Enquist, M. 2001. "The Evolution of Female Sexuality." *Animal*

Behaviour 61, no.4: 695–704. https://doi.org/10.1006/anbe.2000.1630

Rooker, K., and S. Gavrilets. 2020. "On the Evolution of Sexual Receptivity in Female

Primates". *Scientific Reports* 10, no. 1:11945. https://doi.org/10.1038/s41598-020-68338-y

Ryu, H. 2017. "Mechanisms and Socio-Sexual Functions of Female Sexual Swelling, and Male

Mating Strategies in Wild Bonobos." PhD diss., Kyoto University.

Ryu, H., D.A. Hill, T. and Furuichi. 2015. "Prolonged Maximal Sexual Swelling in Wild

Bonobos Facilitates Affiliative Interactions Between Females". *Behaviour* 152:285–311.

https://doi. org/10.1163/1568539X-00003212

Ryu, H., C. Hashimoto, D. A. Hill, K. Mouri, K. Shimizu, and T. Furuichi. 2025. "Male Bonobo

Mating Strategies Target Female Fertile Windows Despite Noisy Ovulatory Signals During

Sexual Swelling." *PLoS Biology* 23, no. 12: e3003503.

https://doi.org/10.1371/journal.pbio.3003503.

Samuni, L., and M. Surbeck. 2023. "Cooperation Across Social Borders in Bonobos." *Science*

382, no. 6672. https://doi.org/10.1126/science.adg0844.

Schielzeth, H. 2010. "Simple Means to Improve the Interpretability of Regression

Coefficients". *Methods in Ecology and Evolution* 1, no.2:103–113.

https://doi.org/10.1111/j.2041-210X.2010.00012.x

Setchell, J. M., and E. J. Wickings. 2004. "Social and Seasonal Influences on the Reproductive

Cycle in Female Mandrills (*Mandrillus sphinx*)." *American Journal of Physical Anthropology*

125: 73–84. https://doi.org/10.1002/ajpa.10375.

Setchell, J. M., M. Charpentier, and E. J. Wickings. 2005. "Mate Guarding and Paternity in

Mandrills: Factors Influencing Alpha Male Monopoly." *Animal Behaviour* 70: 1105–1120.

https://doi.org/10.1016/j.anbehav.2005.02.021.

Shibata, S., and T. Furuichi. 2023. "Intermale Relationships in Wild Bonobos at Wamba".

In *Bonobos and People at Wamba: 50 years of Research,* edited by T. Furuichi, G. Idani, D.

Kimura, H. Ihobe, and C. Hashimoto, 285–310. Singapore, Springer.

https://doi.org/10.1007/978-981-99-4788-1_23

Shibata, S., T. Furuichi. 2024. "Comparative analysis of intragroup intermale relationships: a

study of wild bonobos (*Pan paniscus*) in Wamba, Democratic Republic of Congo and

chimpanzees (*Pan troglodytes*) in Kalinzu Forest Reserve, Uganda". *Primates* 65, no.

4:243–255.

Sillén-Tullberg, B., and A. P. Møller. 1993. "The Relationship Between Concealed Ovulation

and Mating Systems in Anthropoid Primates: A Phylogenetic Analysis." *The American*

Naturalist 141, no. 1: 1–25.

Small, M. F. 1988. "Female Primate Sexual Behavior and Conception: Are There Really Sperm

to Spare?" *Current Anthropology* 29, no.1:81–100. https://doi.org/10.1086/203615

Soltis, J., R. Thomsen, and O. Takenaka. 2001. "The Interaction of Male and Female

Reproductive Strategies and Paternity in Wild Japanese Macaques, *Macaca fuscata*." *Animal*

Behaviour 62: 485–494. https://doi.org/10.1006/anbe.2001.1774.

Soulsbury, C. D. 2010. "Ovulation Mode Modifies Paternity Monopolization in Mammals."

Biology Letters 6: 39–41. https://doi.org/10.1098/rsbl.2009.0703.

Strassmann, B. I. 1981. "Sexual Selection, Paternal Care, and Concealed Ovulation in Humans."

Ethology and Sociobiology 2: 31–40.

Surbeck, M., C. Boesch, C. Crockford, et al. 2019. "Males With a Mother Living in Their Group Have Higher Paternity Success in Bonobos but Not Chimpanzees". *Current Biology* 29, no.10: R354–R355.

Surbeck, M., C. Girard-Buttoz, L. Samuni, et al. 2021. "Attractiveness of Female Sexual Signaling Predicts Differences in Female Grouping Patterns Between Bonobos and Chimpanzees". *Communications Biology* 4, no.1:1119. https://doi.org/10.1038/s42003-021-02641-w

Surbeck, M., and G. Hohmann. 2013. "Intersexual Dominance Relationships and the Influence of Leverage on the Outcome of Conflicts in Wild Bonobos (*Pan paniscus*)". *Behavioral Ecology and Sociobiology* 67:1767–1780. https://doi.org/10.1007/s00265-013-1584-8

Surbeck, M., K. E. Langergraber, B. Fruth, L. Vigilant, and G. Hohmann. 2017. "Male Reproductive Skew is Higher in Bonobos than Chimpanzees". *Current Biology* 27, no.13:R640–R641. https://doi.org/10.1016/j.cub.2017.05.039

Surbeck, M., L. Cheng, M. Kreyer, G. Gort, R. Mundry, G. Hohmann, and B. Fruth. 2025. "Drivers of Female Power in Bonobos." *Communications Biology*. https://doi.org/10.1038/s42003-025-07900-8.

Surbeck, M., R. Mundry, and G. Hohmann. 2011. "Mothers Matter! Maternal Support, Dominance Status and Mating Success in Male Bonobos (*Pan paniscus*)". *Proceedings of the Royal Society B: Biological Sciences* 278, no. 1705:590–598. https://doi.org/10.1098/rspb.2010.1572

Surbeck, M., T. Deschner, G. Schubert, A. Weltring, and Hohmann, G. 2012. "Mate Competition, Testosterone and Intersexual Relationships in Bonobos, *Pan paniscus*". *Animal Behaviour* 83, no. 3:659–669.

Takahata, Y., H. Ihobe, and G. Idani. 1999. "Do Bonobos Copulate More Frequently and

Promiscuously Than Chimpanzees?" *Human Evolution* 14, no. 3: 159–167.

Takemoto, H., Y. Kawamoto, and T. Furuichi. 2015. "How Did Bonobos Come to Range South

of the Congo River? Reconsideration of the Divergence of *Pan paniscus* From Other *Pan*

Populations." *Evolutionary Anthropology: Issues, News, and Reviews* 24, no. 5: 170–184.

https://doi.org/10.1002/evan.21456.

Takemoto, H., Y. Kawamoto, S. Higuchi, E. Makinose, J. A. Hart, T. B. Hart, T. Sakamaki, N.

Tokuyama, G. E. Reinartz, P. Guislain, J. Dupain, A. K. Cobden, M. N. Mulavwa, K.

Yangozene, S. Darroze, C. Devos, and T. Furuichi. 2017. "The Mitochondrial Ancestor of

Bonobos and the Origin of Their Major Haplogroups." *PLoS ONE* 12, no. 5: e0174851.

https://doi.org/10.1371/journal.pone.0174851.

Thompson-Handler, N. E. 1990. "The Pygmy Chimpanzee: Sociosexual Behavior, Reproductive

Biology and Life History Patterns." PhD diss., Yale University.

Thomsen, R., J. Soltis, M. Matsubara, K. Matsubayashi, M. Onuma, and O. Takenaka. 2006.

"How Costly are Ejaculates for Japanese Macaques?" *Primates* 47. no. 3:272–274.

https://doi.org/10.1007/s10329-005-0171-7

Toda, K., and T. Furuichi. 2020. "Low Resistance of Senior Resident Females Toward Female

Immigration in Bonobos (*Pan paniscus*) at Wamba, Democratic Republic of the Congo."

International Journal of Primatology 41, no. 2: 415–427.

https://doi.org/10.1007/s10764-019-00126-6.

Toda, K., K. Mouri, H. Ryu, T. Sakamaki, N. Tokuyama, T. Yokoyama, S. Shibata, M.-L. Poiret,

K. Shimizu, C. Hashimoto, and T. Furuichi. 2022. "Do Female Bonobos (*Pan paniscus*)

Disperse at the Onset of Puberty? Hormonal and Behavioral Changes Related to Their

Dispersal Timing." *Hormones and Behavior* 142: 105159.

https://doi.org/10.1016/j.yhbeh.2022.105159.

Toda, K., N. Tokuyama, and T. Sakamaki. 2023. "Multiple Phases of Natal Transfer Process in

Female Bonobos and Factors Underlying Each Phase: Findings from Long-Term

Observations in Wild Populations". In *Bonobos and People at Wamba: 50 years of Research,*

edited by T. Furuichi, G. Idani, D. Kimura, H. Ihobe, and C. Hashimoto, 181–209. Singapore,

Springer. https://doi.org/10.1007/978-981-99-4788-1_23

Tokuyama, N. 2023. "Aggressive Behaviors and Social Dominance in Bonobos". In *Bonobos*

and People at Wamba: 50 years of Research, edited by T. Furuichi, G. Idani, D. Kimura, H.

Ihobe, and C. Hashimoto, 247–265. Singapore, Springer.

https://doi.org/10.1007/978-981-99-4788-1_21

Tokuyama, N., and T. Furuichi. 2016. "Do Friends Help Each Other? Patterns of Female

Coalition Formation in Wild Bonobos at Wamba". *Animal Behaviour* 119:27–35.

https://doi.org/10.1016/j.anbehav.2016.06.021

Tokuyama, N., T. Sakamaki, and T. Furuichi. 2019. "Inter-Group Aggressive Interaction Patterns

Indicate Male Mate Defense and Female Cooperation Across Bonobo Groups at Wamba,

Democratic Republic of the Congo." *American Journal of Physical Anthropology* 170:

535–550. https://doi.org/10.1002/ajpa.23929.

Tutin, C. E. G. 1979. "Mating Patterns and Reproductive Strategies in a Community of Wild

Chimpanzees (*Pan troglodytes schweinfurthii*)." *Behavioral Ecology and Sociobiology* 6:

29–38.

Van Krunkelsven, E., J. Dupain, L. Van Elsacker, and R. F. Verheyen. 1996. "Food Calling by

Captive Bonobos (*Pan paniscus*): An Experiment." *International Journal of Primatology* 17,

no. 2: 207–217.

Wallen, K., and J. L. Zehr. 2004. "Hormones and History: The Evolution and Development of

Primate Female Sexuality." *The Journal of Sex Research* 41, no. 1: 101–112.

Wallis, J. 1997. "A Survey of Reproductive Parameters in the Free-Ranging Chimpanzees of Gombe National Park". *Reproduction* 109, no. 2:297–307. https://doi.org/10.1530/jrf.0.1090297

Watts, D. P. 2004. "Intracommunity Coalitionary Killing of an Adult Male Chimpanzee at Ngogo, Kibale National Park, Uganda." *International Journal of Primatology* 25, no. 3: 507–521.

Watts, D. P. 2015. "Mating Behavior of Adolescent Male Chimpanzees (*Pan troglodytes*) at Ngogo, Kibale National Park, Uganda." *Primates* 56: 163–172. https://doi.org/10.1007/s10329-014-0453-z.

Weir, B. J., and I. W. Rowlands. 1973. "Reproductive Strategies of Mammals". *Annual Review of Ecology and Systematics* 4:139–163.

Weir, L. K., J. W. Grant, and J. A. Hutchings. 2011. "The Influence of Operational Sex Ratio On the Intensity of Competition for Mates". *The American Naturalist* 177, no.2:167–176.

Weiss, A., J. T. Feldblum, D. M. Altschul, D. A. Collins, S. Kamenya, D. Mjungu, S. Foerster, I. C. Gilby, M. L. Wilson, and A. E. Pusey. 2023. "Personality Traits, Rank Attainment, and Siring Success Throughout the Lives of Male Chimpanzees of Gombe National Park." *PeerJ* 11: e15083. https://doi.org/10.7717/peerj.15083.

Wilcox, A. J., C. R. Weinberg, and D. D. Baird. 1995. "Timing of Sexual Intercourse in Relation to Ovulation—Effects on the Probability of Conception, Survival of the Pregnancy, and Sex of the Baby". *New England Journal of Medicine* 333, no.23:1517–1521.

Wroblewski, E. E., C. M. Murray, B. F. Keele, J. C. Schumacher-Stankey, B. H. Hahn, and A. E. Pusey. 2009. "Male Dominance Rank and Reproductive Success in Chimpanzees, *Pan troglodytes schweinfurthii*." *Animal Behaviour* 77: 873–885.

https://doi.org/10.1016/j.anbehav.2008.12.014.

Yamamoto, S. 2015. "Non-Reciprocal but Peaceful Fruit Sharing in Wild Bonobos in Wamba."

Behaviour 152: 335–357. https://doi.org/10.1163/1568539X-00003257.

Wilson, M. L., C. Boesch, B. Fruth, et al. 2014. "Lethal Aggression in *Pan* is Better Explained

by Adaptive Strategies than Human Impacts". *Nature* 513, no.7518:414–417.

https://doi.org/10.1038/nature13727

Yokoyama, T., and T. Furuichi. 2022. "Why Bonobos Show a High Reproductive Skew Towards

High-Ranking Males: Analyses for Association and Mating Patterns Concerning Female

Sexual States". *Primates* 63, no. 5:483–494. https://doi.org/10.1007/s10329-022-01004-1

Young, W. C., and R. M. Yerkes. 1943. "Factors Influencing the Reproductive Cycle in the

Chimpanzee; The Period of Adolescent Sterility and Related Problems." *Endocrinology* 33,

no. 3: 121–154.

Zhang, Q.-X., Y. Zhu, D.-P. Xia, X. Wang, and J.-H. Li. 2024. "Male Consort Behavior Is

Driven by Female Mate Choice and Receptive Synchrony in Tibetan Macaques (*Macaca*

thibetana)." *Global Ecology and Conservation* 54: e03158.

https://doi.org/10.1016/j.gecco.2024.e03158.

Zuur, A. F., E. N. Ieno, N. Walker, A. A. Saveliev, G. M. Smith. 2009. *Mixed Effects Models and*

Extensions in Ecology with R. New York, Springer.

https://doi.org/10.1007/978-0-387-87458-6

1340

7. Figure Legends

Figure 1. Comparisons in the proportion of copulations involving high-ranking males between

females in the likely fertile period (LFP) and non-LFP females. Open circles represent observed

proportions for each female focal follow, with circle size indicating the total number of

copulations. Filled circles denote model-estimated means, and vertical error bars indicate

confidence intervals.

1347

Figure 2. Relationship between male dominance rank and proportion of copulations involving

females in the likely fertile period (LFP). Male dominance rank is ordered such that lower

values indicate higher-ranking males and higher values indicate lower-ranking males. Open

circles represent observed proportions for each male, with circle size indicating the total number

of copulations. The solid line shows the model-predicted relationship, and the shaded area

represents the 95% confidence interval.

1354

Figure 3. Relationship between within-party male dominance rank and the probability of

copulation with females in the likely fertile period (LFP). Predicted copulation probabilities are

shown separately for one-hour time windows (OTWs) containing one (left panel) or two (right

panel) females in the LFP. The size of open circles indicates observed total counts of whether

copulation occurred for each OTW. Solid lines show model-predicted probabilities, and shaded

areas represent 95% confidence intervals on the probability scale.

1361

Figure 4. Relationship between within-party male dominance rank and the probability of

copulation with females outside the likely fertile period (LFP), changing with the number of

LFP females. Predicted copulation probabilities are shown separately for one-hour time

windows (OTWs) containing zero (left panel), one (center panel), or two (right panel) females

in the LFP. The size of open circles indicates observed total counts of whether copulation

occurred for each OTW. Solid lines show model-predicted probabilities, and shaded areas

represent 95% confidence intervals on the probability scale.

TABLE 1. Basic information about male subjects in the E1 group (December 2022–May 2023).

ID	Age[c] (yrs)	Resident mother[d]	Within-sex dominance rank [David Score]	Focal observation hours and days	Counts of copulation events during focal follows	
					with adults	with adolescents
TN[a]	52*	–	Not used [-2.9]	Not collected	–	–
TW[b]	49*	–	Not used [-12.9]	39.2 h (4 d)[e]	0	0
DI	48*	–	5th [4.6]	39.7 h (4 d)	4	3
NB	35	Ki	4th [7.2]	44.7 h (5 d)	5	4
GC	33*	–	8th [-11.2]	37.5 h (4 d)	3	4
KT	19	Ki	1st [53.4)	36.4 h (4 d)	3	0
JO	17	Jk	3rd [7.5]	36.7 h (4 d)	4	7
KY	14	Ki	2nd [10.0]	30.6 h (4 d)	5	1
HC	14	Hs	6th [-2.7]	32.0 h (4 d)	5	1
SE	12	Sl	7th [-6.1]	32.6 h (4 d)	7	2
NI	10	Nv	10th [-28.9]	25.1 h (4 d)	5	2
YD	9	Yk	9th [-27.0]	34.4 h (4 d)	0	2

[a]TN was not included in the focal subject, because he had been lost for a five month in E1 group

and was absent at the beginning of this study.

[b]TW had seemed erectile dysfunction before the study period, and thus data on him were not

used for the analysis.

[c]Asterisks in the column means estimated ages.

[c]See Table 2.

[e]Data in parentheses were not used for the analysis.

[g]Not included when female fertility status could not be determined

26 **TABLE 2.** Basic information about female subjects in the E1 group (December 2022–May
27 2023).

ID	Age (yrs)[d]	Within-sex dominance rank [David Score]	Length after last parturition (months)[e]	Number of maximal swelling phases	Number of likely fertile periods	Focal observation hours (and days)
No[a]	52*	12th [-2.3]	92	1	0	9.0 h (1 d)
Ki	49*	1st [41.5]	43	3	2	38.0 h (4 d)
Hs	42*	2nd [10.2]	30	4	2	40.7 h (4 d)
Yk	40*	4th [5.3]	16	1	0	35.7 h (4 d)
Jk	35*	9th [-0.29]	28	4	3	38.1 h (4 d)
Sl	31*	6th [4.7]	22	3	2	35.9 h (4 d)
Nv	29*	8th [2.8]	5	0	0	38.3 h (4 d)
Ot	26*	10th [-0.96]	10	3	0	34,5 h (4 d)
Fk	25*	5th [4.8]	33	3	2	36.2 h (4 d)
Pf	19*	3rd [5.3]	25	3	3	35.9 h (4 d)
Ik[b]	16*	15th [-6.0]	26	0	0	34.5 h (4 d)
Sc[b]	15*	11th [-1.2]	44	0	0	29.5 h (4 d)
Db	14*	7th [4.1]	35	3	0	27.1 h (3 d)
Jj	12*	17th [-10.1]	6	2	0	38.3 h (4 d)
Aq	12*	13th [-2.4]	4	4	0	Not collected
Ec[b]	11*	14th [-4.0]	–	0	0	38.6 h (4 d)
Mz[c]	10	18th [-10.8]	6	3	1	36.5 h (4 d)
Tp	10*	20th [-11.7]	–	5	0	35.5 h (4 d)
Bc	10*	19th [-11.3]	–	2	0	33.0 h (4 d)
Ih	9	16th [-8.9]	–	4	0	34.2 h (4 d)

[a]No disappeared from E1 after December 31, 2023 and is estimated to be dead due to her old
age.

[b]Ik, Sc, and Ec were pregnant and gave birth to their infants during the study period.

[c]Mz had gave birth to her first infant at 9.6 years of age but lost it at 4 months after parturition.

[d]Asterisks in the column means estimated ages.

[e]At the beginning of this study period (December 6th, 2022).

46 **TABLE 3.** Results of model analyses for copulation frequencies and patterns

Models	Responses	Predictors	Estimate	SE	Z	P
A: Binomial GLMM (Figure 1)	Proportion of copulations with high- versus low-ranking males	(Intercept)	–0.30	0.30	–1.01	0.313
		Female LFP status (yes, no)	2.02	0.83	2.43	0.015*
		Ratio of numbers of high- versus low-ranking males observed	0.28	0.27	1.01	0.311
B: Binomial GLM (Figure 2)	Proportion of copulations with LFP versus non-LFP females	(Intercept)	–1.91	0.25	–7.75	< 0.001***
		Male rank (1–10)	–1.41	0.24	–5.81	< 0.001***
C: Binomial GLMM (Figure 3)	Probability of copulations with LFP females within OTWs	(Intercept)	–3.30	0.64	–5.15	< 0.001***
		Number of males higher-ranking than the focal male (0–7)	–1.69	0.63	–2.70	0.007**
		Number of LFP females (1–2)	0.93	0.43	2.19	0.028*
		Number of non-LFP females (0–7)	–0.47	0.44	–1.07	0.287
D: Binomial GLMM (Figure 4)	Probability of copulations with non-LFP females within OTWs	(Intercept)	–2.37	0.25	–9.67	< 0.001***
		Number of males higher-ranking than the focal male (0–7)	–0.02	0.20	–0.12	0.906
		Number of LFP females (0–2)	–0.85	0.29	–2.95	0.003**
		Number of non-LFP females (0–7)	0.80	0.21	3.73	< 0.001***
		Interaction between number of males higher-ranking than the focal male and number of LFP females	0.66	0.24	2.78	0.005**

47 All continuous predictors are z-transformed (mean of 0 and a standard deviation of 1).
48 Asterisks indicate significance levels, *: $P < 0.05$, **: $P < 0.01$, ***: $P < 0.001$.
49 GLM = generalized linear model; GLMM = generalized linear mixed model; LFP = likely fertile period; OTW = one-hour time window.