Gene expression profiling of bovine raw milk as a new tool to monitor the inflammatory status of the udder : a pilot study

C. Gitton[a], Y. Le Vern[a], M. Gaborit[b], R. Prado Martins[a,1], P. Germon[a]*

[a] ISP UMR 1282, INRAE, Université François Rabelais de Tours, Nouzilly, France
[b] UEP UE 0326, INRAE, Borculo, Le Pin au Haras, France.

[1] Present address: PFIE UE 1277, INRAE, Nouzilly, France

* Corresponding author
Email: pierre.germon@inrae.fr

Abstract

Inflammation of the mammary gland, or mastitis, is a common disease of dairy cows with important consequences on the economy of dairy farms, well-being of cows and antibiotic usage. Monitoring of the inflammatory status of mammary gland currently relies on milk somatic cell counting. In order to improve this monitoring, methods aiming at differentiating cell types among milk cells are being developed. The objective of the present study was to explore the use of gene expression profiling from raw milk samples to define the inflammatory status of udder quarters. A pilot study allowed us to select 36 genes among 95 whose expression were associated with one of three milk cell types, i.e. neutrophils, macrophages or T lymphocytes, purified by cell sorting. Based on expression data measured by RT-qPCR for these genes, we were able to identify four different types of milk samples, each with specific properties. One cluster was characterized by increased expression of genes typifying neutrophils, a second one was identified by increased expression of genes typifying macrophages while the two others were distinguished based on the expression of genes associated with T lymphocytes. Clustering of samples based on gene expression was shown to be comparable to that obtained by flow cytometry analysis. Our results highlight the potential of profiling of milk samples based on gene expression for the analysis of the inflammatory status of udders.

Keywords

Mastitis, inflammation, milk, gene expression profiling

Introduction

Mastitis remains a major issue in dairy farming. Upon entry and recognition of bacterial pathogens by the host immune cells, in particular mammary epithelial cells and macrophages, an inflammatory response is triggered which leads to recruitment of immune cells in the mammary gland.

As a consequence, somatic cell counts (SCC) in milk is routinely used as an indirect indicator
of the sanitary status of mammary glands.
SCC values below 100,000 cells per ml are generally considered as an indication of a healthy
mammary quarter (Hillerton; Harmon, 1994). On the opposite, SCC values above 200,000
cells/ml are generally taken as an indication of an on-going inflammatory response or a quarter
recovering from infection (Hillerton; Smith and Hillerton, 2001). In between these two SCC values
one cannot state with confidence whether a quarter is healthy or not. In addition, setting SCC
threshold values to distinguish healthy from infected quarters should also take into account
different criteria such as the lactation stage and parity (Mondini et al., 2025). In any case, these
two values are not definite indicators of a health status: mastitic quarters below SCC values of
100,000 could be still be infected by mastitis pathogens (Schwarz et al., 2010).
As a consequence, several improvements have been tested, in particular the differential cell
counts that determines the type of cells present in milk upon inflammation (Rivas et al., 2001;
Pilla et al., 2013). Cells present in milk belong to different cell types including neutrophils,
macrophages, lymphocytes and epithelial cells. The proportion and quantities of each of these
cell population is highly variable. For instance, the strong inflammation observed during clinical
mastitis is characterized by a majority of neutrophils which can amount to almost 90% of cells
present in milk (Sordillo et al., 1997).
Reports on the cell composition of mammary gland secretions from healthy animals show
contrasting results. Several studies, using electronic microscope examinations or flow cytometry
analyses of milk samples from healthy quarters, revealed mostly macrophages (50- 80%) with
10-20 % lymphocytes (Lee et al., 1980; Lun et al., 2007; Souza et al., 2020; Farschtschi et al.,
2021; Silva et al., 2021). Yet, contrasting results have been presented in which T lymphocytes
are the most prevalent cell type in the absence of infections with almost 50% T lymphocytes and
around 20-30 % macrophages (Park et al., 1992; Dosogne et al., 2003; Schwarz et al., 2011).
These discrepancies might be due to different methodologies, to the potential confusion
between macrophages and epithelial cells after staining and light microscopy examination
(Leitner et al., 2000). The cell composition is also highly dependent on the inflammatory status of
the mammary gland, with increased prevalence of neutrophils and decreasing percentages for
lymphocytes, and on the lactation status with increased values in late lactation for macrophages
(Leitner et al., 2000; Schwarz et al., 2011).
Flow cytometry analyses identified different subpopulations of T lymphocytes, in particular
CD4+, CD8+ as well as $\gamma\delta$ T cells, along with different macrophage classical (CD14+ CD16-) and
non-classical (CD14- CD16+) subpopulations (Farschtschi et al., 2022). While CD4 and CD8 T
lymphocytes constitute approximately 20-30 % of all T-cells, $\gamma\delta$ T cells are also present in milk in
significant numbers (10-30% of all lymphocytes in milk) (Taylor et al., 1994; Leitner et al., 2000;
Farschtschi et al., 2022). The proportion of these subpopulations among T lymphocytes depends
on the mammary gland health status: indeed, the CD4/CD8 ratio increases upon infection or
inflammation (Park et al., 1992; Asai et al., 1998; Rivas et al., 2000; Mehrzad et al., 2008; Tassi
et al., 2013) as well as in late lactation (Taylor et al., 1994). While B-cells are only present in
limited numbers in healthy mammary glands, their proportion among lymphocytes was shown to
increase upon inflammation (Schwarz et al., 2013).
Based on such evidence that the proportion of the different cell types is related to the health
status of a mammary gland, differential cell count strategies have been explored in order to not
only rely on SCC values to characterize diseased quarters (Rivas et al., 2001; Schwarz et al.,
2011). One such Differential Somatic Cell Count (DSCC) method discriminates between
neutrophils and macrophages while lymphocytes are counted along with neutrophils (Damm et
al., 2017; Halasa and Kirkeby, 2020). Four groups based on SCC and DSCC values were
determined. Similar to previous studies, this method highlights variations depending on parity and
lactation status (Schwarz et al., 2020b; Zecconi et al., 2023). Although both indicators are linked
to the recruitment of leukocytes in mammary gland secretions, their dynamics was found to be
slightly different, with DSCC values decreasing faster than SCC in the resolutive phase of
inflammation (Kirkeby et al., 2020). DSCC can thus be considered as an additional indicator for
the control of mastitis. Yet, the use of DSCC will require the definition of suitable thresholds in
order to improve its predictive value (Schwarz et al., 2020b). In fact, a recent survey suggest very
little added value of DSCC over SCC in terms of positive- and ngetive- predictive values for the
detection of mastitis cases due to major pathogens (Fonseca et al., 2025a). Furthermore, the

definition of suitable thresholds for SCC/DSCC combinations will have to take into account a
potential herd effect since authors have also observed significantly different values between
different herds, showing that the herd effect could be more important that the pathogen (Kirkeby
et al., 2021).
103
High resolution differential cell count based on flow cytometry has also been proposed
recently to better characterize the inflammation/infectious status of mammary glands (Farschtschi
et al., 2022). A more global approach has also been tested using single-cell RNAseq highlighting
the presence of different cell populations and subpopulations in a healthy milk sample (Becker et
al., 2021; Wiarda et al., 2025). Altogether these different methods allow the identification of cell
populations present in milk.
We hypothesized that another strategy to characterize the inflammatory status of mammary
glands could be based on a functional assessment of the cells present in mammary gland
secretion. In order to do so, we first identified a subset of genes specific for macrophages,
neutrophils or T lymphocytes, three of the main cell populations present in milk. We then
undertook a gene-expression profiling analysis of these selected genes based on RNA samples
collected from unprocessed raw milk. Gene expression profiles were determined allowing
clustering of milk samples and the clustering obtained was then compared to results obtained by
flow cytometry.

Materials and methods

Collection of milk samples

Quarter foremilk samples (40 ml) from all four quarters were collected under aseptic
conditions as recommended by the National Mastitis Council guidelines, from 28 Holstein (HF)
and 10 Normande (NO) cows at the experimental farm of Le Pin-au-Haras (Unité Expérimentale
du Pin-au-Haras). No criteria was applied for the selection of cows from which milk samples were
collected. More specifically, after teat cleaning, the first streams of milk were discarded,
premilking disinfectant solution was applied and then drier, teat apex was disinfected by scrubing
with a 70% ethanol moistened gauze pad and milk was collected in 50ml tubes held at 45° angle
to avoid contamination by dust materials. Milk was then transported to the laboratory at room
temperature within 3h.
A 500 µL portion of each milk sample was subjected to a Fossomatic model 90 (Foss Electric,
Hillerod, Denmark) to determine the milk SCC. The somatic cell score (SCS) was calculated with
the formula : SCS = Log2(SCC / 100,000) + 3. One milk sample from each cow was selected,
except for four cows for which two samples were collected from two different quarters: the only
criteria for selection of milk samples was to have a panel of milk samples representing a large
range of SCC values. These quarter milk samples originated from cows that were at different
stages of lactation (days-in-milk ranging from 35 to 391 days) and with different SCC values
(ranging from 17,000 to 3,603,000 cells/ml). None of the samples were from clinical mastitis
cases. Details are presented in **Supplementary table S1**.
138
Bacteriological examination of the selected quarter milk samples was performed by plating
50 µL of milk on sheep blood agar plates (ThermoFicher Columbia agar with Sheep Blood PLUS
#PB5039A) and recording bacterial colonies after 24 and 48 hours of incubation at 37°C. Basic
tests (colony morphology and colour, hemolysis, Gram-staining, catalase) were performed in
order to qualify the bacteriological status of milk samples. Samples were considered
contaminated when more that 3 different colony types were observed.

Isolation and labelling of milk cells

Milk cells were collected from the selected milk samples: An equal volume of 1× DPBS
(reference, Sigma D8537) was added to milk samples. The diluted milk sample was left at room
temperature for 15 minutes and then centrifuged at room temperature for 10 minutes at 1400g.

The cream layer and the supernatant were removed and the cell pellet was then washed once in
10ml of room temperature (DPBS, 2mM EDTA, 1% horse serum) and then centrifuged 5 minutes
at 500g. The cells were then resuspended in FACS buffer (DPBS, 2mM EDTA, 10% horse
serum). Cells were then labelled prior to flow cytometry analysis or cell sorting.
Cells were quantified using a LUNA-FL cell counter (Logos Biosystems) according to the
manufacturer's instructions. One million cells prepared as indicated above were transferred to a
1.5ml tube, centrifuged and resuspended in FACS buffer. Cells were labeled for 30 min with
primary antibodies (30 min, 4°C). Primary antibodies used in the present study were CD14-PE-
AF750 (Bio-Rad, reference MCA1568P750, clone Tük4), CD45-PE (Bio-Rad, reference
MCA2220PE, clone 1.11.32), G1 (Kingfisher Biotech, reference WSC0608B-100, clone CH138A)
and CD3 (Kingfisher Biotech, reference WS0561B-100, clone MM1A). After washing in FACS
buffer, cells were incubated (30 min, 4°C) with the secondary antibodies BV421 labelled rat anti-
mouse IgM antibody (BD Biosciences, reference 562595, clone R6-60.2) and Alexa Fluor™ 488
labelled Goat anti-Mouse IgG1 (Thermo Fisher Scientific, reference A21121) in FACS buffer,
matching respectively G1 and CD3 primary antibodies. After washing in FACS buffer, cells were
resuspended in DPBS containing the Zombie Aqua viability dye (BioLegend, reference 423101)
for 30 minutes at 4°C. After a final wash in FACS buffer, cells were resuspended in FACS buffer.
166

Cells sorting of purified specific leucocytes populations

Milk cells from three quarter milk samples were collected and labelled as described above.
These three samples were selected based on their SCC, representative of low, intermediate and
high SCC quarters. Labelled milk cells were then sorted on a MoFlo Astrios EQ (Beckman
Coulter) high speed cell sorter equipped with 4 lasers to obtain, among live cells, purified
macrophages (CD45+ G1- CD14+), T lymphocytes (CD45+ CD3+) and neutrophils(CD45+ G1+).
The average sorting speed was 10,000 cells/s.

Flow cytometry analysis of milk cells

To determine the proportion of leucocyte subpopulations in milk samples, milk cells were
isolated and labelled as described above. Milk cells were then analyzed on a LSR Fortessa™ X-
20 Flow cytometer (Becton Dickinson) and results were analyzed with the Kaluza software v2.1
(Beckman Coulter). Among live cells, macrophages were identified as CD45+ G1- CD14+ cells,
neutrophiles as CD45+ G1+ cells and T lymphocytes as CD45+ CD3+ cells. Our gating strategy
is exemplified in **Supplementary Figure S1**.

RNA extraction and high-throughput RT-qPCR

For sorted cells, immediately after collection, cells were centrifuged 10 minutes at 1000g,
resuspended in 1 ml of Trizol and stored at -80°C until RNA was extracted as described below.
For milk samples, 200 microliters of milk were mixed with 800 µl of Trizol (Thermofisher) in a
lysing matrix D tube (MP Bio) and homogenized for 45 seconds – speed 6 in a FastPrep-24
apparatus. Samples were then stored at -80°C until further processed.
Upon thawing of milk and purified cells samples mixed with Trizol, 200 µl chloroform was
added, the tube was vortexed for 10 seconds and then centrifuged at 12000g for 15min at 4°C.
The aqueous phase was collected, mixed with an equal volume of 70% ethanol, loaded on a
NucleoSpin RNA columns (Macherey Nagel) and RNA was then purified and treated with DNAse
I as indicated by the manufacturer. Total RNA (50 ng) was reverse transcribed to cDNA using 5x
iScript reverse transcription supermix (Biorad) according to manufacturer's instructions. Ten-fold
diluted cDNA were preamplified according to Fluidigm's protocol (quick reference PN 100-5875
B1) using PreAmp master mix and a mix of all forward and reverse primers at 500nM each.
Preamplification program was 2 minutes 95°C, followed by 16 cycles of (95°C for 15s, 60°C for 4
min). Preamplified cDNA were then diluted 5-fold with (10 mM Tris, pH 8.0, 1 mM EDTA) buffer.
Gene expression level were measured from pre-amplified cDNA on 96.96 GE Dynamic Array IFC

using the Fluidigm BioMark^{TM} HD System. Primers used in this study are listed in
Supplementary Table S1.

Data analysis

A reference sample was built from the arithmetic mean of Ct for each gene of analyzed
samples. Fold changes compared to this reference sample were calculated by the ∆∆Ct method
using *ACTB* and *GAPDH* as reference genes (Livak and Schmittgen, 2001). Heatmaps were
generated using the pheatmap package (v1.0.10) in RStudio (1.1). Likelyhood Ratio Test was
performed using the GTest function of the R package DescTools (v0.99). PCA analyses were
performed using the PCA function of the FactoMineR package (v2.9). Mann-Whitney and
Kruskall-Wallis tests were performed using the stats package (v4.5.1). Relatedness of
dendrogram based on gene expression and flow cytometry proportions were analyzed using the
dendextend package (v1.19.1). Scripts used in the analysis and raw qPCR data are available at
https://github.com/pgermon37/milk_RNA_profile.
211

212   **Results**

Identification of a gene panel able to discriminate macrophages, neutrophils and T
lymphocytes

The aim of our study was to investigate if profiling of milk samples based on gene expression
could provide relevant information regarding the inflammatory status of mammary gland quarters.
For such an analysis to take into account the contribution of different leucocyte populations to the
gene expression profile of raw milk samples, we first needed to select a panel of genes that
showed different levels of expression depending on the cell type.
Therefore, we sorted macrophages neutrophils (live CD45+ G1+ cells), (live CD45+ G1-
CD14+ cells), and T lymphocytes (live CD45+ CD3+ cells) by flow cytometry from three quarter
milk samples having somatic cell counts ranging from 45,000 to 389,000 cells/ml
(**Supplementary Table S2**).
The expression of a panel of 95 genes by these purified sorted cells was investigated by high-
throughput RT-qPCR. Genes were selected based on their expected role in the function of cells
types studied and based on the laboratory's expertise in bovine immunity. Only genes yielding a
specific amplification with no primer dimers, as evaluated by melting curve analysis, in all
samples were selected which restricted the number of genes to 78. Expression results for these
78 genes in the three selected cell types were first analyzed by hierarchical clustering (**Figure**
1A) and Principal Component Analysis (**Figure 1B**). Both analyses showed clear separation of
samples based on cell types. The hierarchical clustering analysis allowed the identification of 6
clusters of genes with specificities for one or two of the cell types. For example, the cluster
identified as "Lc" contains genes that are more expressed in T lymphocytes than in the two other
cell types. A second cluster specifically more expressed in neutrophils, labeled "PMN2", can also
be clearly observed, as well as a cluster labeled "Macro" associated with higher expression in
macrophages. The list of genes from these six clusters are presented in **Supplementary Table**
S3.

Figure 1: Heatmap and PCA of gene expression of sorted milk macrophages, neutrophils and T lymphocytes. Cells from three milk samples from three different cows (one smaple per cow) were collected by centrifugation, labelled with antibodies against CD3 (T lymphocytes), CD14 (macrophages) and G1 (neutrophils) and sorted on a MoFlo Astrios EQ cell sorter. RNA was extracted from purified cell populations and analyzed by high-throughput RT-qPCR. Gene expression values were calculated against a theoretical control sample and analyzed by principal component analysis (A) and hierarchical clustering (B).

Based on these results, because they showed clearly distinct patterns of expression between the three leucocyte populations investigated, we selected the 41 genes belonging to clusters "Lc" (11 genes), "PMN2" (15 genes) and "Macro" (15 genes) for the investigation of expression profiles in RNA from raw milk samples.

Comparison of clustering based on gene expression and flow cytometry profiles

We then analyzed quarter milk samples from 28 Holstein (HF), which included the three quarter milk samples used for sorting the purified cell types, and from 10 Normande (NO) cows. One sample per cow was selected so that the panel of samples covered a range of SCC value from 17,000 to 3,603,000 cells/ml, with only 6 samples above the 400,000 cells/ml threshold. Cows were at different stages of lactation (days-in-milk ranging from 35 to 391 days). Detailed description of the different milk samples are given in **Supplementary Table S2**.

SCC distribution of selected milk samples are depicted in **Supplementary Figure S2**. We could not detect any statistically significant difference in terms of SCC between samples from Holstein and Normande cows (Mann-Whitney p value = 0.29). The breed criteria was thus not explored further in downstream analyses.

For each sample, the gene expression profile after RNA extraction was determined by high-throughput RT-qPCR. Among the 41 genes selected above, expression values were obtained for all samples for only 36 genes (excluded genes were *CD4*, *CXCL10*, *CCL20*, *IFNB1* and *TNFa*). Analysis by hierarchical clustering of expression values for these 36 genes, performed without any a priori, clearly identified four clusters of samples (**Figure 2A**). The PCA analysis showed a clear separation of these four clusters (**Figure 2B**). Clusters 1 and 2 encompass samples with SCC values generally below 100,000 cells/ml whereas cluster 3 and 4 samples are characterized by significantly higher SCC (**Figure 3**).

274
275

Figure 2: Heatmap and PCA of gene expression of raw milk samples. Milk samples were aseptically collected from 38 individual quarters. RNA was extracted directly from raw milk samples and analyzed by high-throughput RT-qPCR. Gene expression values were calculated against a theoretical control sample and analyzed by hierarchical clustering (A) and principal component analysis (B). Color code for clusters is indicated on the figure.

Figure 3: log(SCC) values of milk samples from the four different clusters. The log(SCC) of each of the milk samples analyzed is represented and samples are grouped and colored depending on the cluster to which they were allocated in the hierarchical clustering analysis. Color code is identical to that of Figure 2. *

 indicates statistical significance (p < 0,05). p-values were calculated using Mann and Whitney after global
comparison using a Kruskal and Wallis test.

As can be seen in **Figure 2**, our method was able to differentiate different categories of
samples in the SCC range of 50,000 to 200,000 with samples belonging to clusters 1, 2 and 3. In
this SCC range we thus had samples with significantly different immune status: even though the
SCC of cluster 3 samples is low, these samples are characterized by a higher expression of
"Macro" genes and are thus quarters where macrophages could be present in higher proportions
or be more transcriptionally active. As expected, high SCC milk samples belonged to cluster 4
characterized by higher expression of "PMN2" genes associated with the neutrophil cell
population. Samples belonging to clusters 1 and 2 are characterized by a higher level of
expression of "Lc" genes but can be distinguished from one another by different levels of
expression of "Macro" and "PMN2" genes maybe suggesting different immune status.
We could not detect any statistically significant difference in lactation stage between the
different clusters (**Supplementary Figure S3** – Kruskal-Wallis p value = 0.35).

299
300   For each of these milk samples, the proportion of macrophages, neutrophils and T
301 lymphocytes were obtained by flow cytometry (see **Supplementary Table S2**). Analysis of these
302 results by Principal Component Analysis showed separation of samples from the 4 clusters
303 identified previously by gene-expression profiling with some outliers (**Figure 4**).
304
305

Figure 4: Principal component analysis of proportions of milk samples analyzed by flow cytometry.
Proportions of macrophages, neutrophils and T-lymphocytes among cells from collected milk samples was
determined by flow cytometry. Principal Component Analysis of the cell composition was performed on
the percent values for each of the three cell types analyzed. Dots are colored depending on the cluster to

which they were allocated based on the hierarchical clustering analysis of gene expression data. Color
code is identical to that of Figure 2 and is indicated on the figure.

To quantitatively estimate the relatedness of analyses based on gene expression and flow
cytometry, we analyzed the relatedness of dendrograms obtained with both methods calculating
Baker's index (data not shown). This analysis showed that the two trees share significant
similarities: the null hypothesis that the two trees are not similar can be rejected with a risk below
0.01.

Clustering based on gene expression profile differentiates samples with similar SCC.

Classical bacteriological analysis was performed on all selected milk samples. Absence of
bacteria was observed in 13 samples while, when bacteria were present, only Gram-positive
pathogens were detected (16 samples). Bacteriological status could not be determined for 5
samples and 4 samples could not be examined due to technical reasons (**Supplementary Table**
S2). When the clustering was analyzed relative to the presence of bacteria in the milk, the
number of "No growth" sample decreases while the number of samples showing growth of
"Gram-positive" bacteria increases (**Figure 5**). As expected, cluster 4 samples have a tendency
to contain bacteria (Likelihood Ratio Test p-value = 0.06).

Figure 5: Barplot representing the number of samples in bacteriology positive or negative samples.
Bacteriological status of each milk sample was analyzed by plating 50 microliter of each milk sample on
sheep-blood agar plates. Samples with more that 3 types of colonies were considered as contaminated.
Gram staining was performed on colonies growing on plates. NA indicates absence of bacteriological
analysis.

Discussion

The objective of the present study was to investigate a new method for the functional
characterization of the inflammatory status of mammary gland quarters. As an alternative to the
characterization of cells present in milk, we investigated the gene expression profile of milk
samples as a surrogate for the functionality of cells present in milk.
In order to determine if this strategy would allow us to characterize the inflammatory status of
low cell counts samples, for which current strategies provide low positive predictive values of
being infected, we focused our analyses on samples with rather low SCC values: the median
value for SCC is 95 500 (Q1-Q3 : 38 250 – 218 000). In order to improve the diagnostic of
inflammatory quarters, analyses were done on quarter milk samples. Indeed, analyses of
composite samples show reduced predictive power compared to analyses based on quarter milk
samples (Dufour and Dohoo, 2012; Fonseca et al., 2025c). Importantly, it should be noted that
none of the milk samples analyzed were from clinical cases, even the two samples with the high
SCC values.
347
Based on the expression level of 36 genes from which expression could be measured in all
samples, we identified four different clusters of milk samples, each with specific gene expression
profiles. Cluster 4 samples are characterized by increased expression from neutrophil and
macrophage related genes. Cluster 3 samples are typified by increased expression of
macrophage related genes while clusters 1 and 2 samples are characterized by higher
expression of genes typifying T lymphocytes. This clustering based on gene expression actually
overlaps with that obtained from flow cytometry: consistent with the increased expression of
neutrophil gene set, cluster 3 and 4 samples are the ones with the highest number of neutrophils.
Compared to clusters 1 and 2, clusters 3 and 4 can therefore be considered as inflamed quarters.
Nevertheless, cluster 3 and 4 samples are different: cluster 3 samples show higher values for
macrophage related genes but less expression of neutrophil related genes compared to cluster 4
samples. Hence, cluster 3 samples could originate from mammary gland quarters in the process
of resolution with increased proportions of dead neutrophils, could be representatives of
chronically infected cows or be specific to a particular pathogen. With the available data, we
cannot decide which hypothesis is the most likely. The number of samples we analyzed is clearly
not sufficient to firmly establish a link between a particular profile and a specific pathogen.
Our results also clearly distinguished cluster 1 from cluster 2 samples, two types of milk
samples with low somatic cell counts. These low SCC milk samples are indeed not identical in
terms of inflammatory status. Although they show similar expression of genes typifying T
lymphocytes, they showed different expression of genes typifying macrophages or neutrophils. In
particular cluster 2 samples express higher levels of the neutrophil-associated genes while
cluster 1 samples showed higher expression of macrophage-associated genes. They
consequently represent two different inflammatory statuses. Because of higher expression of pro-
inflammatory markers (CXCL8 for example), whose expression is clearly induced in inflamed
mammary glands (Bannerman et al., 2004), we hypothesize that cluster 2 samples are samples
where a recruitment of neutrophils has been initiated. Overall, cluster 2 samples indeed have
higher percentage of neutrophils (71 % +/- 25% vs. 39% +/- 25%, $p < 0.01$). On the contrary, we
hypothesize that cluster 1 samples correspond to healthy samples with limited inflammation.
Compared to cluster 2 samples, cluster 1 samples have higher expression levels of macrophage-
associated genes: considering these samples as healthy is consistent with the observation of
higher proportions of macrophages in healthy mammary glands (Lee et al., 1980; Damm et al.,
2017; Halasa and Kirkeby, 2020).
380
The method described in this pilot study has several advantages over current methods to
qualify the inflammatory status of mammary gland quarters. Compared to clustering strategies
proposed in other studies, such as grouping samples based on SCC values and bacteriological
status or on SCC and DSCC values (Schwarz et al., 2020a; Souza et al., 2020), the clusters we

have identified were obtained without any a priori, are only based on intrinsic properties of the
milk samples linked to macrophages, neutrophils and T lymphocytes and are not conditioned by
user-defined thresholds. It should be noted that even in low SCC milk samples a clear gene
expression pattern could be observed. Therefore, contrary to DSCC which requires SCC values
above 50,000 cells/ml, a low SCC is not a limit to the use of our method (Damm et al., 2017).
In addition, the protocol presented here does not rely on preliminary centrifugation of milk
samples which, as can be the case for flow cytometry methods, might fail to recover all cells; in
particular macrophages that can remain in the cream layer after centrifugation (Lee et al., 1980).
We therefore believe that it could bring an added value to the definition of inflammatory status
of mammary gland quarters. More specifically, this method could be used to investigate in more
details the response of mammary gland quarters to infection. For instance, neutrophil subtypes in
different inflammatory conditions could be investigated which would extend our observations of
increased proportions of a neutrophil subset expressing MHCII surface molecules during
subclinical mastitis (Rambault et al., 2023). Moreover, in vaccine based studies were a local
immune response is expected, the gene panel could be adapted to monitor more precisely the
response of T-lymphocytes such as the Th17 response that has been shown to be beneficial to
the clearance of bacteria (Herry et al., 2017; Cebron et al., 2020).
For these more detailed analyses to be possible, new developments will be required to
increase the panel of genes analyzed in order to better distinguish additional cell types or
subtypes. Single-cell RNAseq analyses have recently been performed on milk cells (Wiarda et
al., 2025). These data confirmed some of the associations described herein: for example, CXCL8
and TLR4 were also identified as specific for granulocytes/neutrophils. CD14 was identified as
associated to macrophages, as expected for a macrophage specific gene (Grandoni et al., 2023).
More detailed analyses of these data and of the cell clusters observed could provide usefull
information regarding additionnal genes specific for different cell types or even subtypes.
410
The method described herein nevertheless has certain limitations. The main limit is that it is
not easily accessible to on-farm diagnostic of mammary gland inflammation and requires specific
equipments. Its interest is limited for the moment to laboratory studies trying to better qualify the
inflammatory status of the udder. We also failed to clearly quantify epithelial cells by flow
cytometry. Although these cells are likely to represent only a low proportion of milk cells, their
proportions could potentially modify the clustering observed by flow cytometry. Nevertheless, it is
likely that the gene expression profiling of milk samples is actually susceptible to the presence of
epithelial cells in milk (Boutinaud et al., 2015). It is also possible, since fat globules have been
shown to contain RNA molecules originating from the milk producing epithelial cell, that the gene
expression profile also reflects the functionality of mammary epithelial cells (Canovas et al.,
2014). As a consequence, RNA quantification from raw milk might also reflect both the presence
and/or functionality of epithelial cells.
423
The present report should be considered as a pilot study that was limited to only a small
number of samples. In order to validate the usefulness of our method, more broader studies
would be required. A more detailed investigation of the relationship between gene expression
clusters and the bacteriological status of mammary gland quarters would definitely support the
predictive value of these analyses. So far, only an tendency for an increased proportion of
samples with Gram+ bacteria in cluster 4 samples was observed, consistent with the fact that
these samples are the ones with the highest SCC. It also remains to be determined whether the
gene expression profile varies depending on parity, days-in-milk as these parameters are clearly
associated with variations in SCC and DSCC (Fonseca et al., 2025b). Complementary analyses
should also take into account the variability of milk cellular composition at different stages during
milking (Ostensson et al., 1988). The current profiles identified were based on foremilk samples
which are likely to be enriched in SCC in samples with more than 100,000 cells/ml (Sarikaya and
Bruckmaier, 2006).
437

To conclude, we set up a new way of monitoring the inflammatory status of mammary gland
quarters during lactation based on the profiling of gene expression from raw milk with minimal
processing before RNA extraction. We were able to distinguish four types of milk samples that
corresponded to different inflammatory status. Potential refinements include the addition of new
genes to better distinguish quarters with low SCC values, improved qualification of the
inflammatory status of mammary glands along with more in-depth characterization of bacteria
present in milk samples.
Although not applicable as such in commercial farms, this method paves the way to the
development of new medium- or high- throughput phenotyping strategies. Such methods will
provide useful information for the implementation of selective dry cow therapy or for the routine
analysis of health status of mammary glands.

Acknowledgements

450   The authors would like to thank the animal staff of the "Unité Expérimentale du Pin-au-Haras"
451 (INRAE, Gouffern-en-Auge, France) and of the "Unité Expérimentale de Physiologie animale de
452 l'Orfrasière" (INRAE, Nouzilly, France) for their help in collect ing milk samples. We are grateful to
453 Aude Remot for critical reading of the manuscript.

Ethics approval

455 Not applicable. Milk samples were collected from cows as part of a routine milking.

Funding

457 This work received funding from APIS-GENE (Masticells project).

Conflict of interest disclosure

459   The authors declare that they comply with the PCI rule of having no financial conflicts of
460 interest in relation to the content of the article.

Data, scripts, code, and supplementary information availability

462 Data are available online: https://doi.org/10.5281/zenodo.18633548.
463 Supplementary information is available online: https://doi.org/10.5281/zenodo.18631973

Author ORCIDs

465 Christophe Gitton   0000-0001-7986-0414
466 Yves Le Vern   0009-0009-3930-6107
467 Marie Gaborit   0009-0008-5461-1196
468 Rodrigo Prado Martins   0000-0002-8213-4015
469 Pierre Germon   0000-0003-3078-876X

References

471
472

473 Asai, K., Kai, K., Rikiishi, H., Sugawara, S., Maruyama, Y., Yamaguchi, T., Ohta, M., Kumagai, K., 1998.
474   Variation in CD4+ T and CD8+ T lymphocyte subpopulations in bovine mammary gland secretions

during lactating and non-lactating periods. Veterinary Immunology and Immunopathology 65, 51-61. doi:10.1016/s0165-2427(98)00176-7.

Bannerman, D.D., Paape, M.J., Lee, J.W., Zhao, X., Hope, J.C., Rainard, P., 2004. Escherichia coli and *Staphylococcus aureus* elicit differential innate immune responses following intramammary infection. Clinical and Diagnostic Laboratory Immunology 11, 463-472. doi:10.1128/CDLI.11.3.463-472.2004.

Becker, D., Weikard, R., Hadlich, F., Kuhn, C., 2021. Single-cell RNA sequencing of freshly isolated bovine milk cells and cultured primary mammary epithelial cells. Sci Data 8, 177. doi:10.1038/s41597-021-00972-1.

Boutinaud, M., Herve, L., Lollivier, V., 2015. Mammary epithelial cells isolated from milk are a valuable, non-invasive source of mammary transcripts. Front Genet 6, 323. doi:10.3389/fgene.2015.00323.

Canovas, A., Rincon, G., Bevilacqua, C., Islas-Trejo, A., Brenaut, P., Hovey, R.C., Boutinaud, M., Morgenthaler, C., VanKlompenberg, M.K., Martin, P., Medrano, J.F., 2014. Comparison of five different RNA sources to examine the lactating bovine mammary gland transcriptome using RNA-Sequencing. Sci Rep 4, 5297. doi:10.1038/srep05297.

Cebron, N., Maman, S., Walachowski, S., Gausseres, B., Cunha, P., Rainard, P., Foucras, G., 2020. Th17-related mammary immunity, but not a high systemic Th1 immune response is associated with protection against *E. coli* mastitis. npj Vaccines 5, 108. doi:10.1038/s41541-020-00258-4.

Damm, M., Holm, C., Blaabjerg, M., Bro, M.N., Schwarz, D., 2017. Differential somatic cell count-A novel method for routine mastitis screening in the frame of Dairy Herd Improvement testing programs. Journal of Dairy Science 100, 4926-4940. doi:10.3168/jds.2016-12409.

Dosogne, H., Vangroenweghe, F., Mehrzad, J., Massart-Leen, A.M., Burvenich, C., 2003. Differential leukocyte count method for bovine low somatic cell count milk. Journal of Dairy Science 86, 828-834. doi:10.3168/jds.S0022-0302(03)73665-0.

Dufour, S., Dohoo, I.R., 2012. Monitoring dry period intramammary infection incidence and elimination rates using somatic cell count measurements. Journal of Dairy Science 95, 7173-7185. doi:10.3168/jds.2012-5839.

Farschtschi, S., Hildebrandt, A., Mattes, M., Kirchner, B., Pfaffl, M.W., 2022. Using High-Resolution Differential Cell Counts (HRDCCs) in Bovine Milk and Blood to Monitor the Immune Status over the Entire Lactation Period. Animals (Basel) 12 doi:10.3390/ani12111339.

Farschtschi, S., Mattes, M., Hildebrandt, A., Chiang, D., Kirchner, B., Kliem, H., Pfaffl, M.W., 2021. Development of an advanced flow cytometry based high-resolution immunophenotyping method to benchmark early immune response in dairy cows. Sci Rep 11, 22896. doi:10.1038/s41598-021-02405-w.

Fonseca, M., Kurban, D., Roy, J.-P., Santschi, D.E., Molgat, E., Dufour, S., 2025a. Usefulness of differential somatic cell count for udder health monitoring: Diagnostic performance of somatic cell count and differential somatic cell count for diagnosing intramammary infections in dairy herds with automated milking systems. Journal of Dairy Science 108, 3929-3941. doi:10.3168/jds.2024-25404.

Fonseca, M., Kurban, D., Roy, J.P., Santschi, D.E., Molgat, E., Dufour, S., 2025b. Usefulness of differential somatic cell count for udder health monitoring: Effect of intramammary infections, days in milk, quarter location, and parity on quarter-level differential somatic cell count and somatic cell score in apparently healthy dairy cows. Journal of Dairy Science 108, 3878-3899. doi:10.3168/jds.2024-25401.

Fonseca, M., Kurban, D., Roy, J.P., Santschi, D.E., Molgat, E., Yang, D.A., Dufour, S., 2025c. Usefulness of differential somatic cell count for udder health monitoring: Identifying referential values for differential somatic cell count in healthy quarters and quarters with subclinical mastitis. Journal of Dairy Science 108, 3917-3928. doi:10.3168/jds.2024-25403.

Grandoni, F., Fraboni, D., Canonico, B., Papa, S., Buccisano, F., Schuberth, H.-J., Hussen, J., 2023. Flow Cytometric Identification and Enumeration of Monocyte Subsets in Bovine and Water Buffalo Peripheral Blood. Current Protocols 3, e676. doi:https://doi.org/10.1002/cpz1.676.

Halasa, T., Kirkeby, C., 2020. Differential Somatic Cell Count: Value for Udder Health Management.
Front Vet Sci 7, 609055. doi:10.3389/fvets.2020.609055.
Harmon, R.J., 1994. Physiology of mastitis and factors affecting somatic cell counts. Journal of Dairy
Science 77, 2103-2112. doi:10.3168/jds.S0022-0302(94)77153-8.
Herry, V., Gitton, C., Tabouret, G., Reperant, M., Forge, L., Tasca, C., Gilbert, F.B., Guitton, E., Barc, C.,
Staub, C., Smith, D.G.E., Germon, P., Foucras, G., Rainard, P., 2017. Local immunization impacts
the response of dairy cows to *Escherichia coli* mastitis. Sci Rep 7, 3441. doi:10.1038/s41598-017-
03724-7.
Hillerton, J., Redefining mastitis based on somatic cell count. Book of abstracts of the p. 4–6.
Kirkeby, C., Schwarz, D., Denwood, M., Farre, M., Nielsen, S.S., Gussmann, M., Toft, N., Halasa, T.,
2021. Dynamics of somatic cell count (SCC) and differential SCC during and following
intramammary infections. Journal of Dairy Science 104, 3427-3438. doi:10.3168/jds.2020-19378.
Kirkeby, C., Toft, N., Schwarz, D., Farre, M., Nielsen, S.S., Zervens, L., Hechinger, S., Halasa, T., 2020.
Differential somatic cell count as an additional indicator for intramammary infections in dairy
cows. Journal of Dairy Science 103, 1759-1775. doi:10.3168/jds.2019-16523.
Lee, C.S., Wooding, F.B., Kemp, P., 1980. Identification, properties, and differential counts of cell
populations using electron microscopy of dry cows secretions, colostrum and milk from normal
cows. Journal of Dairy Research 47, 39-50. doi:10.1017/s0022029900020860.
Leitner, G., Shoshani, E., Krifucks, O., Chaffer, M., Saran, A., 2000. Milk leucocyte population patterns
in bovine udder infection of different aetiology. J Vet Med B Infect Dis Vet Public Health 47, 581-
589. doi:10.1046/j.1439-0450.2000.00388.x.
Livak, K.J., Schmittgen, T.D., 2001. Analysis of relative gene expression data using real-time
quantitative PCR and the 2(-Delta Delta C(T)) Method. Methods 25, 402-408.
doi:10.1006/meth.2001.1262.
Lun, S., Astrom, G., Magnusson, U., Ostensson, K., 2007. Total and differential leucocyte counts and
lymphocyte subpopulations in lymph, afferent and efferent to the supramammary lymph node,
during endotoxin-induced bovine mastitis. Reprod Domest Anim 42, 126-134. doi:10.1111/j.1439-
0531.2006.00741.x.
Mehrzad, J., Janssen, D., Duchateau, L., Burvenich, C., 2008. Increase in Escherichia coli inoculum
dose accelerates CD8+ T-cell trafficking in the primiparous bovine mammary gland. Journal of
Dairy Science 91, 193-201. doi:10.3168/jds.2007-0096.
Mondini, S., Gislon, G., Zucali, M., Sandrucci, A., Tamburini, A., Bava, L., 2025. Factors influencing
somatic cell count and leukocyte composition in cow milk: A field study. Journal of Dairy Science
108, 2721-2733. doi:10.3168/jds.2024-25357.
Ostensson, K., Hageltorn, M., Astrom, G., 1988. Differential cell counting in fraction-collected milk
from dairy cows. Acta Veterinaria Scandinavica 29, 493-500. doi:10.1186/BF03548647.
Park, Y.H., Fox, L.K., Hamilton, M.J., Davis, W.C., 1992. Bovine mononuclear leukocyte subpopulations
in peripheral blood and mammary gland secretions during lactation. Journal of Dairy Science 75,
998-1006. doi:10.3168/jds.S0022-0302(92)77842-4.
Pilla, R., Malvisi, M., Snel, G.G., Schwarz, D., Konig, S., Czerny, C.P., Piccinini, R., 2013. Differential cell
count as an alternative method to diagnose dairy cow mastitis. Journal of Dairy Science 96, 1653-
1660. doi:10.3168/jds.2012-6298.
Rambault, M., Gilbert, B.F., Roussel, P., Tessier, A., David, V., Germon, P., Winter, N., Remot, A.,
2023. MHC-II expressing neutrophils circulate in blood and milk during mastitis and show high
microbicidal activity. Journal of Dairy Science
Rivas, A.L., Quimby, F.W., Blue, J., Coksaygan, O., 2001. Longitudinal evaluation of bovine mammary
gland health status by somatic cell counting, flow cytometry, and cytology. Journal of Veterinary
Diagnostic Investigation 13, 399-407. doi:10.1177/104063870101300506.
Rivas, A.L., Quimby, F.W., Coksaygan, O., Olmstead, L., Lein, D.H., 2000. Longitudinal evaluation of
CD4+ and CD8+ peripheral blood and mammary gland lymphocytes in cows experimentally
inoculated with Staphylococcus aureus. Canadian Journal of Veterinary Research 64, 232-237.

Sarikaya, H., Bruckmaier, R.M., 2006. Importance of the sampled milk fraction for the prediction of
total quarter somatic cell count. Journal of Dairy Science 89, 4246-4250. doi:10.3168/jds.S0022-
0302(06)72470-5.
Schwarz, D., Diesterbeck, U.S., Failing, K., Konig, S., Brugemann, K., Zschock, M., Wolter, W., Czerny,
C.P., 2010. Somatic cell counts and bacteriological status in quarter foremilk samples of cows in
Hesse, Germany--a longitudinal study. Journal of Dairy Science 93, 5716-5728.
doi:10.3168/jds.2010-3223.
Schwarz, D., Diesterbeck, U.S., Konig, S., Brugemann, K., Schlez, K., Zschock, M., Wolter, W., Czerny,
C.P., 2011. Flow cytometric differential cell counts in milk for the evaluation of inflammatory
reactions in clinically healthy and subclinically infected bovine mammary glands. Journal of Dairy
Science 94, 5033-5044. doi:10.3168/jds.2011-4348.
Schwarz, D., Kleinhans, S., Reimann, G., Stuckler, P., Reith, F., Ilves, K., Pedastsaar, K., Yan, L., Zhang,
Z., Valdivieso, M., Barreal, M.L., Fouz, R., 2020a. Investigation of dairy cow performance in
different udder health groups defined based on a combination of somatic cell count and
differential somatic cell count. Prev Vet Med 183, 105123. doi:10.1016/j.prevetmed.2020.105123.
Schwarz, D., Rivas, A.L., Konig, S., Diesterbeck, U.S., Schlez, K., Zschock, M., Wolter, W., Czerny, C.P.,
2013. CD2/CD21 index: a new marker to evaluate udder health in dairy cows. Journal of Dairy
Science 96, 5106-5119. doi:10.3168/jds.2013-6804.
Schwarz, D., Santschi, D.E., Durocher, J., Lefebvre, D.M., 2020b. Evaluation of the new differential
somatic cell count parameter as a rapid and inexpensive supplementary tool for udder health
management through regular milk recording. Prev Vet Med 181, 105079.
doi:10.1016/j.prevetmed.2020.105079.
Silva, V.M., Souza, M.T., Blagitz, M.G., Souza, F.N., Batista, C.F., Alves, A.J., Fernandes, A.C.C.,
Sanchez, E.M.R., Ordinola-Ramirez, C.M., da Costa, L., Della Libera, A., 2021. Milk lymphocyte
profile and macrophage functions: new insights into the immunity of the mammary gland in
quarters infected with Corynebacterium bovis. BMC Vet Res 17, 282. doi:10.1186/s12917-021-
02989-5.
Smith, K.L., Hillerton, J.E., 2001. Guidelines on Normal and Abnormal Raw Milk Based on Somatic Cell
Counts and Signs of Clinical Mastitis. National Mastitis Council Research Committee
Sordillo, L.M., Shafer-Weaver, K., DeRosa, D., 1997. Immunobiology of the mammary gland. Journal
of Dairy Science 80, 1851-1865. doi:10.3168/jds.S0022-0302(97)76121-6.
Souza, F.N., Blagitz, M.G., Batista, C.F., Takano, P.V., Gargano, R.G., Diniz, S.A., Silva, M.X., Ferronatto,
J.A., Santos, K.R., Heinemann, M.B., De Vliegher, S., Della Libera, A., 2020. Immune response in
nonspecific mastitis: What can it tell us? Journal of Dairy Science 103, 5376-5386.
doi:10.3168/jds.2019-17022.
Tassi, R., McNeilly, T.N., Fitzpatrick, J.L., Fontaine, M.C., Reddick, D., Ramage, C., Lutton, M.,
Schukken, Y.H., Zadoks, R.N., 2013. Strain-specific pathogenicity of putative host-adapted and
nonadapted strains of Streptococcus uberis in dairy cattle. Journal of Dairy Science 96, 5129-5145.
doi:10.3168/jds.2013-6741.
Taylor, B.C., Dellinger, J.D., Cullor, J.S., Stott, J.L., 1994. Bovine milk lymphocytes display the
phenotype of memory T cells and are predominantly CD8+. Cellular Immunology 156, 245-253.
doi:10.1006/cimm.1994.1169.
Wiarda, J.E., Davila, K.M.S., Trachsel, J.M., Loving, C.L., Boggiatto, P., Lippolis, J.D., Putz, E.J., 2025.
Single-cell RNA sequencing characterization of Holstein cattle blood and milk immune cells during
a chronic Staphylococcus aureus mastitis infection. Sci Rep 15, 12689. doi:10.1038/s41598-025-
96657-5.
Zecconi, A., Zaghen, F., Meroni, G., Sora, V., Martino, P.A., Laterza, G., Zanini, L., 2023. Early Milk
Total and Differential Cell Counts as a Diagnostic Tool to Improve Antimicrobial Therapy
Protocols. Animals (Basel) 13 doi:10.3390/ani13071143.

A-

B-

A-

B-