Development of a vehicle for non-invasive oral administration to facilitate behavioral experiments in fish

Sayaka Matsuo[1] · Takehiko Yokoyama[1,2] · Shun Satoh[3,4] · Yuki Yoshio[3] · Kazuya Fukuda[1*]

[1] School of Marine Biosciences, Kitasato University, Sagamihara, Kanagawa 252-0373, Japan

[2] Current address: Graduate School of Agricultural Science, Tohoku University, Sendai, Miyagi, 980-8572, Japan

[3] Graduate School of Science, Kyoto University, Yoshida-honmachi, Sakyo-ku, Kyoto, Japan

[4] The Hakubi Center for Advanced Research, Kyoto University, Yoshida-honmachi, Sakyo-ku, Kyoto, Japan

*Correspondence: Kazuya Fukuda (E-mail: fukuda.kazuya@kitasato-u.ac.jp, Tel: +81-42-778-8009). 1-15-1 Kitazato, Minami-ku, Sagamihara, Kanagawa 252-0373, Japan

ORCiD ID

Takehiko Yokoyama: 0000-0002-1991-9100

Shun Satoh: 0000-0003-1284-5249

Yuki Yoshio: 0009-0007-1325-2310

Kazuya Fukuda: 0000-0002-4355-1545

Total text pages: 10

Numbers of figures: 1

Abstract

Pharmacological approaches are increasingly used in fish ethological studies to investigate the proximate mechanisms underlying behavior. However, an injection-based administration can induce stress or injury, potentially interfering with natural behavioral patterns. Here, we present a non-invasive vehicle for oral administration. To demonstrate its utility, the oxytocin receptor antagonist atosiban was administered orally using this vehicle and subsequently detected in blood plasma via high-performance liquid chromatography. Our method offers an easy-to-use and minimally disruptive tool for fish behavioral experiments.

Keywords

Fish, Oral administration, Atosiban, Non-invasive administration, High-performance liquid chromatography

Introduction

In recent years, ethological studies have increasingly adopted integrative approaches that consider not only the ultimate factors of behavior but also its proximate mechanisms (Davies et al. 2012; Rubenstein and Alcock 2019). This broader perspective has enabled researchers to gain a more comprehensive understanding of the evolution of behavior and to identify conserved patterns across taxa. It has also led to novel insights that are difficult to obtain through observational studies alone. For example, Matsumoto et al. (2018) investigated filial cannibalism in male blenniid fish, *Rhabdoblennius nitidus*, and successfully resolved a theoretical paradox in behavioral ecology by incorporating an endocrinological approach. Similarly, Nowicki et al. (2020) compared the expression patterns of nonapeptides, dopamine, and opioid receptor genes in the forebrains of pair-bonded and solitary butterflyfish species and found that these systems modulate sociality as is also observed in other vertebrates. These studies illustrate that approaches bridging proximate and ultimate factors hold great potential to open new directions for research.

An accessible approach to bridge these levels is to use pharmacological manipulation within traditional behavioral experiments, in which candidate regulatory systems (e.g., neural, endocrine, or physiological bases) are enhanced or inhibited to

assess effects on behavior. This approach has been widely used; for example, in studies

of territorial behavior (Santangelo and Bass 2006), paternal care (O'Connell et al. 2012),

shoaling (Ataei Mehr et al. 2020), and reproductive behavior (Almeida et al. 2023). In

most cases, pharmacological manipulations involve injecting compounds into subjects

via intraperitoneal, intracerebroventricular, or intramuscular routes. However, because

such injection-based methods are inherently invasive, they are sometimes unsuitable

for behavioral experiments depending on the experimental design. For example, when

experiments must be conducted over extended periods, treatments are required to be

administered repeatedly, which can induce substantial stress or physical injury. These

consequences may alter the physiological state and confound behavior (e.g., Sneddon

et al. 2014). To avoid these problems, non-invasive compound delivery methods are

needed.

One non-invasive method is oral administration, which has been used in long-

term treatments (e.g., Madrid et al. 2021; Lu and Patton 2022; Ochocki and Kenney

2023). However, oral administration of a considerable number of compounds remains

challenging compared with injection-based methods due to various physiological and

biochemical barriers, including low permeability and degradation in gastrointestinal

fluids (Song et al. 2004; Lou et al. 2023). In fish, oral administration faces similar

limitations, and efforts have been made to improve its efficacy (e.g., Amezawa et al.

2018). A practical and accessible strategy is to optimize the composition of the vehicle

mixed with the compounds to enhance delivery efficiency, as the vehicle's composition

is known to influence oral delivery outcomes (e.g., Tønsberga 2010; Serdoz et al. 2010).

Establishing an effective and easy-to-use vehicle system could provide a valuable tool

for behavioral experiments in ethological studies of fish.

In this study, we examined whether an oligopeptide compound could enter

systemic circulation following oral administration in combination with a vehicle modified

with reference to pharmaceutical research. As a model fish, we used the convict cichlid

(*Amatitlania nigrofasciata*), a species employed in various behavioral studies, including

those on reproductive behavior (e.g., Santangelo 2005), proactive prosociality (Satoh et

al. 2021), and affective state (Laubu et al. 2019). As a test case, we selected atosiban, a

receptor antagonist of the neuropeptide oxytocin, which has been widely studied in the

context of social behavior. In some fish, atosiban introduced into systemic circulation via

intramuscular or intraperitoneal injection has been shown to affect behavior, with its

effects expected to result from function within the central nervous system (e.g., Cardoso

et al. 2015; Ataei Mehr et al. 2020). However, to our knowledge, it has not yet been used

in behavioral experiments via oral administration in teleost fish. Therefore, we used high-

performance liquid chromatography (HPLC) to determine whether orally administered

atosiban, given with the modified vehicle, transfers into the systemic circulation. Based

on these results, we propose a novel vehicle and oral administration method suitable for

use in ethological studies of fish.

Materials and methods

Six male convict cichlids (58–99 mm SL) were collected from Okinawa-jima Island,

Japan. Fish were housed in a 50 L tank at Kitasato University, where the water

temperature was maintained at 26 ℃ under a 14 h light/10 h dark photoperiod, and fed a

commercial fish food (Hikari Carnivore, Kyorin CO., LTD.), hereafter "food", twice daily.

To prepare the medicated food for oral administration, atosiban (T5148,

TargetMol) was first dissolved in dimethyl sulfoxide (DMSO; 045-24511, Fujifilm Wako

Pure Chemical Co.) and then mixed with either a modified vehicle or a control solution.

Following Kim et al. (2017), the vehicle composition was modified. The vehicle

composition was modified following Kim et al. (2017) and consisted of atosiban, 50

µg/µL; DMSO, 5% (v/v); polyethylene glycol 400 (PEG400; N0443, Tokyo Chemical

Industry Co., Ltd.), 40% (v/v); Tween 80 (35703-62, Nacalai Tesque, Inc.), 5% (v/v); and

reverse osmosis water. The control solution contained 50 µg/µL atosiban and 5% (v/v)

DMSO in reverse osmosis water to control for the absence of other excipients. Food

pellets were cut into pieces approximately 3 mm in diameter and 4 mm in length, each

infused with 10 µL of either the atosiban-containing vehicle or control solution, and air-

dried at room temperature for one hour. All medicated food was prepared immediately

before oral administration.

The procedures for oral administration and sample collection were as follows.

After a three-day fasting period, fish were divided into two groups. Three individuals

were each fed three pieces of vehicle-treated food, and the remaining three were each fed

three pieces of control-treated food. After 55 min, each fish was deeply anesthetized with

benzocaine, and blood was collected from the caudal vessel. Blood samples were

immediately centrifuged at 10,000 × g for 15 min at 4 °C to obtain serum. The serums

from three fish that had been fed the same treated food were pooled. Following the

addition of 50 µl of 1 M perchloric acid (162-00695, Fujifilm Wako Pure Chemical Co.)

to the pooled sample, it was centrifuged again at 15,000 × g for 20 minutes at 4°C, and

the supernatant was collected. The supernatant was then passed through a 0.22 µm

hydrophilic nylon filter (SLNY1322N, Hawach Scientific), and concentrated to

approximately 50 µl using a centrifugal concentrator. The concentrated samples were then

injected into the HPLC system. Standard solutions of atosiban (0.5 and 5 µg/ml dissolved

in ultra-pure water) were used to detect atosiban in samples.

The chromatographic analysis was performed with reference to Raju et al. (2014)

with a JASCO (Tokyo, Japan) HPLC system consisting of a PU-4180 quaternary gradient

pump with a degasser, a CO-4060 column oven, an AS-4050 autosampler with a cooling

system, an UV-4075 UV/VIS detector, and a ChromNAV data processor. The analytical

column was a reversed-phase ODS-120H (4.6 × 250 mm) (Tosoh, 0023415). Elution was

performed with a mixture of solvent A (30 mM potassium phosphate buffer at pH 3.2)

and solvent B (acetonitrile:milli-Q water = 80:20) at 40 °C; the flow rate was set at 1.0

mL・min^{-1} and detention wavelength was set at 220 nm. The elution gradient was set as

follows: 0–5 min, 20–20% solvent B in solvent A; 5–12 min, 20-70% solvent B in solvent

A.

Results

HPLC analysis detected a peak corresponding to orally administered atosiban with the

same retention time as the atosiban standard only in the group that ingested atosiban with

the modified vehicle (Fig. 1). These results demonstrate that oral administration using the

proposed vehicle delivered atosiban into systemic circulation.

Discussion

Ethological studies by adaptationist thinking have long contributed to understanding the

evolutionary background (ultimate factor) of social behaviors across vertebrates, and

some theoretical and empirical frameworks of ethological studies are broadly applicable

to adjacent research fields (Davies et al. 2012; Rubenstein and Alcock 2019). One of the

key emerging questions is whether behaviors that evolved under similar ultimate factors

have entirely different phylogenetic origins and mechanisms, or whether they are, at least in part, underpinned by evolutionarily homologous mechanisms. Indeed, commonalities across vertebrates have been identified not only in the phenotype but also in some aspects of the underlying regulatory mechanisms for some social behaviors, such as territoriality (Oldfield et al. 2015), maternal and paternal care behaviors (Dulac et al. 2014), and monogamous mating system (Young et al. 2019). These examples illustrate how an integrative approach can reveal that some of the core mechanisms regulating these social behaviors are conserved across vertebrates. In recent years, behavioral studies focusing on psychological/ cognitive aspects in fish, such as prosociality (Satoh et al. 2021), self-recognition (Kohda et al. 2019, 2022; Kobayashi et al. 2024), and affective states (Laubu et al. 2019), have gained increasing attention. Notably, similarities have been found between fish and mammals in mechanisms underlying some cognitive abilities (Calvo and Schluessel 2021) and fear contagion (Akinrinade et al. 2023). As such findings accumulate, the importance of integrative research is likely to become even more pronounced. The method proposed in this study offers a practical tool for incorporating non-invasive pharmacological manipulation into ethological experiments without the need for complex procedures or costly equipment. PEG400 and Tween 80 are excipients used for oral administration, which are generally regarded as nontoxic and nonirritant in mammals (e.g., Rowe et al. 2009; Gullapalli and Mazzitelli 2015; Ma et al. 2017). They have also been shown to enhance the permeability of the administered compound (Tønsberga 2010), and their application has also been extended to fish (e.g., Kim et al. 2017). We expect the vehicle proposed here will be particularly useful for long-term behavioral studies and for experiments involving sensitive species or delicate behaviors, where minimally disruptive conditions are essential. We expect the vehicle proposed here will be particularly useful for long-term behavioral studies and for experiments involving sensitive species or delicate behaviors, where minimally disruptive conditions.

On the other hand, we acknowledge that several considerations should be noted in applying our method. First, although we used atosiban as a model compound, the extent to which this vehicle improves oral delivery efficiency for other substances should be validated individually. For example, while atosiban is an oligopeptide and hydrophilic, enhancements in permeability with PEG400 and Tween 80 are often reported for hydrophobic compounds (e.g., Tønsberga 2010). Second, if a greater transfer into

systemic circulation via the oral route is required, alternative strategies may be necessary.
In pharmacology, several sophisticated approaches for improving oral bioavailability
have been developed (Brayden et al. 2020; Cui et al. 2022; Lou et al. 2023). As our
primary aim was to minimize procedural complexity, it would be valuable to consider
incorporating such methods.

Therefore, the present findings should be regarded as a preliminary proof-of-
concept. Elucidating the underlying mechanisms was beyond the scope of this study and
will require future research. Nevertheless, we hope that our results will encourage the
future development of non-invasive pharmacological methods for application in
ethological experiments.
199

200 **Acknowledgments**
201 We are grateful to C. Ogawa for their support during our study.
202

203 **Author contributions**
204 Conceptualization, Methodology: SM, TY, SS, YY, KF; Investigation: SM, TY, KF;
205 Writing - original draft preparation: SM, KF; Writing - review and editing: TY, SS, YY,
206 KF; Funding acquisition: KF; Resources: SM, SS, KF; Supervision: TY, SS, KF
207

208 **Funding**
209 This work was financially supported by JSPS KAKENHI Grant Numbers 24932659,
210 20346357, 23721203 to KF, and the Hakubi Project of Kyoto University.
211

212 **Data availability**
213 The datasets are available from the corresponding author on reasonable request.
214

215 **Declarations**
216 **Conflict of interest**
217 The authors declare that they have no conflicts of interest.
218

219 **Ethical approval**
220 At Kitasato University, there are currently no institutional ethical regulations for fish.

221 Therefore, all procedures adhered to the ASAB/ABS guidelines for the treatment of
222 animals in behavioral research.
223

References

Akinrinade I, Kareklas K, Teles MC, Reis TK, Gliksberg M, Petri G, Levkowitz G,
Oliveira RF (2023) Evolutionarily conserved role of oxytocin in social fear
contagion in zebrafish. Science 379:1232–1237

Almeida O, Félix AS, Oliveira RF (2023) Interaction between vasotocin and gonadal
hormones in the regulation of reproductive behavior in a cichlid fish. Acta Ethol
26:31–43

Amezawa K, Yazawa R, Takeuchi Y, Yoshizaki G (2018) Spawning induction of blue
mackerel *Scomber australasicus* and eastern little tuna *Euthynnus affinis* by oral
administration of a crude gonadotropin-releasing hormone analogue. Fish Sci
84:495–504

Ataei Mehr B, Garner SR, Neff BD (2020) Effect of isotocin on shoaling behaviour of
the guppy (*Poecilia reticulata*). Anim Cogn 23:1093–1100

Brayden DJ, Hill TA, Fairlie DP, Maher S, Mrsny RJ (2020) Systemic delivery of peptides
by the oral route: Formulation and medicinal chemistry approaches. Adv Drug Deliv
Rev 157:2–36

Calvo R, Schluessel V (2021) Neural substrates involved in the cognitive information
processing of fishes. Anim Cogn 24:923–946

Cardoso SC, Bshary R, Mazzei R, Paitio JR, Oliveira RF, Soares MC (2015) Arginine
vasotocin modulates associative learning in a mutualistic cleaner fish. Behav Ecol
Sociobiol 69:1173–1181

Cui J, Wen Z, Zhang W, Wu W (2022) Recent advances in oral peptide or protein-based
drug liposomes. Pharmaceuticals 15:1072

Davies NB, Krebs JR, West SA (2012) An introduction to behavioural ecology, 4th edn.
Wiley-Blackwell, Oxford

Dulac C, O'Connell LA, Wu Z (2014) Neural control of maternal and paternal behaviors.
Science 345:765–770

Gullapalli RP, Mazzitelli CL (2015) Polyethylene glycols in oral and parenteral
formulations–A critical review. Int J Pharm 496:219–239

Kim SS, Im SH, Yang JY, Lee YR, Kim GR, Chae JS, Shin DS, Song JS, Ahn S, Lee BH,
Woo JC, Ahn JH, Yun CS, Kim P, Kim HR, Lee KR, Bae MA (2017) Zebrafish as a
screening model for testing the permeability of blood–brain barrier to small
molecules. Zebrafish 14:322–330

Kobayashi T, Kohda M, Awata S, Bshary R, Sogawa S (2024) Cleaner fish with mirror
self-recognition capacity precisely realize their body size based on their mental
image. Sci Rep 14:20202

Kohda M, Hotta T, Takeyama T, Awata S, Tanaka H, Asai J, Jordan AL (2019) If a fish
can pass the mark test, what are the implications for consciousness and self-
awareness testing in animals? PLoS Biol 17:e3000021

Kohda M, Sogawa S, Jordan AL, Kubo N, Awata S, Satoh S, Kobayashi T, Fujita A,
Bshary R (2022) Further evidence for the capacity of mirror self-recognition in
cleaner fish and the significance of ecologically relevant marks. PLoS Biol
20:e3001529

Laubu C, Louâpre P, Dechaume-Moncharmont F-X (2019) Pair-bonding influences
affective state in a monogamous fish species. Proc R Soc B 286:20190760

Lou J, Duan H, Qin Q, Teng Z, Gan F, Zhou X, Zhou X (2023) Advances in oral drug
delivery systems: challenges and opportunities. Pharmaceutics 15:484

Lu Y, Patton EE (2022) Long-term non-invasive drug treatments in adult zebrafish that
lead to melanoma drug resistance. Dis Model Mech 15: dmm049401

Ma BL, Yang Y, Dai Y, Li Q, Lin G, Ma YM (2017) Polyethylene glycol 400 (PEG400)
affects the systemic exposure of oral drugs based on multiple mechanisms: taking
berberine as an example. RSC Adv 7:2435–2442

Madrid RRM, Mertins O, Tavares-Dias M, Flores-Gonzales AP, Patta ACMF, Ramirez
CAB, Rigoni VLS, Mathews PD (2021) High compliance and effective treatment of
fish endoparasitic infections with oral drug delivery nanobioparticles: Safety of
intestinal tissue and blood parameters. J Fish Dis 44:1819–1829

Matsumoto Y, Tateishi T, Terada R, Soyano K, Takegaki T (2018) Filial cannibalism by
male fish as an infanticide to restart courtship by self-regulating androgen levels.
Curr Biol 28:2831–2836.e3

Nowicki JP, Pratchett MS, Walker SPW, Coker DJ, O'Connell LA (2020) Gene
expression correlates of social evolution in coral reef butterflyfishes. Proc R Soc B

287:20200239

O'Connell LA, Matthews BJ, Hofmann HA (2012) Isotocin regulates paternal care in a monogamous cichlid. Horm Behav 61:725–733

Ochocki AJ, Kenney JW (2023) A gelatin-based feed for precise and non-invasive drug delivery to adult zebrafish. J Exp Biol 226:jeb245186

Oldfield RG, Harris RM, Hofmann HA (2015) Integrating resource defence theory with a neural nonapeptide pathway to explain territory-based mating systems. Front Zool 12:S16

Raju VVSSA, Mohamud AB, Pathi PJ, Raju NA (2014) Estimation of atosiban acetate in parenterals by RP-HPLC. J Pharm Sci Innov 3:39–42

Rowe RC, Sheskey PJ, Quinn ME (2009) Handbook of Pharmaceutical Excipients, 6th edn. Pharmaceutical Press, London / American Pharmacists Association, Washington, DC

Rubenstein DR, Alcock J (2019) Animal behavior: an evolutionary approach, 11th edn. Manhattan, New York

Santangelo N (2005) Courtship in the monogamous convict cichlid: what are individuals saying to rejected and selected mates? Anim Behav 69:143–149

Santangelo N, Bass AH (2006) New insights into neuropeptide modulation of aggression: field studies of arginine vasotocin in a territorial tropical damselfish. Proc R Soc B 273:3085–3092

Satoh S, Bshary R, Shibasaki M, Inaba S, Sogawa S, Hotta T, Awata S, Kohda M (2021) Prosocial and antisocial choices in a monogamous cichlid with biparental care. Nat Commun 12:1775

Serdoz F, Voinovich D, Perissutti B, Grabnar I, Hasa D, Ballestrazzi R (2010) An innovative oxytetracycline self-emulsifying formulation for fish diets: preparation, characterisation and oral bioavailability in rainbow trout (*Oncorhyncus mykiss*) and in European sea bass (*Dicentrarchus labrax*). J Drug Del Sci Tech 20:431–437

Sneddon LU, Elwood RW, Adamo SA, Leach MC (2014) Defining and assessing animal pain. Anim Behav 97:201–212

Song NN, Zhang SY, Liu CX (2004) Overview of factors affecting oral drug absorption. Asian J Drug Metab Pharmacokinet 4:167–176

Tønsberg H, Holm R, Boll JB, Jacobsen J, Müllertz A (2010) Effects of polysorbate 80

on the in-vitro precipitation and oral bioavailability of halofantrine from

polyethylene glycol 400 formulations in rats. J Pharm Pharmacol 62:63–70

Young RL, Ferkin MH, Ockendon-Powell NF, Orr VN, Phelps SM, Pogány Á, Richards-

Zawacki CL, Summers K, Székely T, Trainor BC, Urrutia AO, Zachar G, O'Connell

LA, Hofmann HA (2019) Conserved transcriptomic profiles underpin monogamy

across vertebrates. Proc Natl Acad Sci USA 116:1331–1336

Fig1. Chromatograms of (**a**) a blood extract from fish that ingested atosiban with the
modified vehicle solution, and of the 0.5 µg/mL atosiban standard, and (**b**) a blood extract
from fish that ingested atosiban with the control solution, and of the 5 µg/mL atosiban
standard. Black arrowheads indicate the peak of atosiban. Cyan and magenta lines
represent the chromatogram of the blood extract and standard, respectivery.