Electrophysiological Characteristics of Epidural Spinal Signals in Preclinical Models of Spinal Cord Stimulation

Running Title: Spinal Signals in Preclinical SCS Models

Kimberley Ladner, MRes[1], Eline M. Versantvoort, MSc[1], Marjolein E. G. Thijssen, MSc/Meng[2], Dave Mugan, BSc, MBA[1,2], Stuart N. Baker, PhD[3], Alexander Kraskov, PhD[3], Quoc C. Vuong, PhD[3,4], Mahima Sharma, PhD[5], Marom Bikson, PhD[5], Birte E. Dietz, PhD[1,2], Stefano Palmisani, MD[6], Ilona Obara, PhD[1]*

[1]School of Pharmacy and Translational and Clinical Research Institute, Newcastle University, Newcastle-upon-Tyne, UK; [2]Saluda Medical Europe Ltd, Harrogate, UK; [3]Biosciences Institute, Newcastle University, Newcastle-upon-Tyne, UK; [4]School of Psychology, Newcastle University, Newcastle-upon-Tyne, UK; [5]Neural Engineering Laboratory, Department of Biomedical Engineering, The City College of the City University of New York, City College Center for Discovery and Innovation, USA; [6]Guy's and St. Thomas' NHS Foundation Trust, London, UK

Source(s) of Financial Support: This study was supported by Saluda Medical and statutory funding from Newcastle University. Equipment was provided by Saluda Medical Pty Ltd, Sydney, Australia.

Authorship Statement: KL conducted the study, including data collection, analysis, and interpretation, and wrote the manuscript. EMV participated in data collection, analysis, interpretation, and reviewed the manuscript. MEGT contributed to data collection and assisted in manuscript review. DM designed the study and reviewed and edited the manuscript. SNB and AK were involved in data collection and interpretation and also

reviewed the manuscript. QCV performed data analysis and interpretation and contributed to manuscript review. MS and MB participated in data interpretation and manuscript revision. BED and SP contributed to manuscript review. IO designed and conducted the study, including data collection, analysis, and interpretation, and wrote the manuscript. All authors read and approved the final manuscript.

Conflict of Interest: MEGT, DM, and BED are employees of Saluda Medical and hold stock in the company. The City University of New York holds patents related to brain stimulation, with MB listed as an inventor. MB also holds equity in Soterix Medical Inc., and has provided consultancy, expert witness support, received grants, assigned inventions, and/or served on the Scientific Advisory Board of the following organizations: SafeToddles, Boston Scientific, GlaxoSmithKline, Biovisics, Axonics, Mecta, SigmaStim, Lumenis, Wave Neuroscience, Neurolief, Allergan (AbbVie), and Ceragem Clinical. SP has received consultancy fees and conducts research funded by Saluda Medical and Nevro. IO has received travel and research grants from Saluda Medical. None of the authors have a commercial interest in the material presented in this study. No other relationships exist that could be perceived as potential conflicts of interest.

***Corresponding Author:** Ilona Obara, School of Pharmacy, Faculty of Medical Sciences, Newcastle University, King George VI Building, Newcastle-Upon-Tyne, NE1 7RU, UK.
Phone: +44 (0) 7464616175, Email: ilona.obara@ncl.ac.uk

ABSTRACT

Objectives: Epidural stimulation of the spinal cord evokes distinct electrophysiological responses that can be recorded epidurally. Here, we characterized evoked compound action potentials (ECAPs), doublets (secondary or tertiary ECAPs, likely of different physiological origin than primary ECAPs), evoked synaptic activity potentials (ESAPs), and electromyographic (EMG) signals in preclinical models. Our objective was to clarify the features and distinct physiological origins of these signals, in order to advance mechanistic studies and support clinical applications of spinal cord stimulation (SCS) therapy.

Materials and Methods: Adult male Sprague-Dawley rats (300-440 g) were implanted with two epidural leads (caudal and rostral; each with eight electrodes) and received monopolar, biphasic stimulation (200 µs pulse width) at 2 and 50 Hz, with current increased stepwise to motor threshold. Rhesus macaques (11.5 and 10.2 kg) were implanted with a single 12-electrode epidural lead and stimulated using either tripolar, triphasic pulses at 10 Hz (100 µs) or tripolar, biphasic pulses at 3 Hz (80 µs) up to 3×ECAP threshold. Recordings were taken from non-stimulating electrodes.

Results: ECAPs and EMG signals were recorded across multiple spinal segments in both rats (L1-T7) and macaques (L2-T11). Doublets presented as complex waveforms with multiple negative peaks, two in rats and three in macaques, likely representing distinct ECAPs at T11-T6 in a rat and L1-T11 in macaques. ESAPs, detectable in rats, showed anatomical specificity, over the L1/T13 vertebrae with peak responses at L1. Signal analysis included activation thresholds, amplitudes, latencies, and conduction velocities.

Conclusions: This study outlines electrophysiological signals evoked by SCS in terms of their waveform, recruitment thresholds, and putative physiological origins. We propose that, to the extent these signals reflect different aspects of spinal processing and may serve as

biomarkers of dysregulated nociceptive pathways, as well as indicators of SCS efficacy or potential side effects.

Keywords: evoked compound action potentials, doublets, evoked synaptic activity potentials, spinal cord stimulation, rat, macaque

INTRODUCTION

Over the past decade, the technology to record electrophysiological signals from the epidural space in both humans and animal models during dorsal column (DC) stimulation has advanced spinal cord stimulation (SCS) therapy and research [1-5]. In particular, the recording of evoked compound action potentials (ECAPs), which reflect the synchronous activation of DC fibers, have enabled the development of a closed-loop SCS system [6-8]. ECAPs, when integrated into the closed-loop system, provide a real-time and pragmatic biomarker of spinal cord activation. This enables continuous monitoring of neural activation and dynamic adjustment of stimulation parameters to maintain DC activation, which has been linked to improved outcomes in pain patients [9-12]. Beyond their clinical utility, ECAP recordings also offer insights into the neurophysiological mechanisms underlying SCS, providing a direct and dynamic view of neural modulation during stimulation [13, 14].

Recent studies have identified additional electrophysiological signals recorded via epidural SCS electrodes that reflect neurophysiological responses to stimulation distinct from ECAPs [15, 16]. Sharma et al. (2023) identified a novel electrophysiological signal during epidural SCS in rats, termed evoked synaptic activity potentials (ESAPs), which reflect synaptic activity within dorsal horn neurons following DC activation [15]. ESAPs consist of several slow waveforms, most notably the S1 component, which appears after the triphasic ECAP signal (P1-N1-P2). The synaptic origin of ESAPs was validated by administering a glutamatergic synapse blocker (6-Cyano-7-nitroquinoxaline-2,3-dione; a selective competitive AMPA receptor antagonist) which significantly reduced the S1 wave without affecting the ECAP [15]. A separate fast evoked potential, referred here to as a secondary ECAP, or doublet, occurring ~1-2 ms after the primary ECAP has been reported by Gmel et al. (2023) in response to SCS in humans. They proposed the involvement of post-synaptic dorsal column (PSDC) fibers, which may be activated either synaptically by primary afferents or directly by stimulation pulses [16]. These secondary responses exhibited distinct conduction velocities, suggesting they arise from a separate neural population than those

contributing to the primary ECAP [16]. Herein, we discuss the possibility that these likely second-order signals propagate through one of several spinal tracts [17]. Finally, electromyographic (EMG) signals, recorded from epidurally placed electrodes across multiple species, represent muscular responses to epidural stimulation, arising either from a Hoffmann reflex or direct activation of motor neurons [18].

The novel evoked spinal potentials, like ESAPs and doublets, are compelling because they reflect neural processing distinct from ECAPs, offering complementary biomarkers for therapeutic engagement and mechanistic insight. However, epidural recordings during SCS are also susceptible to myogenic signals and various stimulation artifacts, and therefore rigorous signal processing is essential to accurately distinguish neurophysiological responses from myogenic or electrical noise [19, 20].

To advance the understanding of electrophysiological signals evoked by SCS, we analyzed experimental data from both rats and macaques and reviewed the existing literature on these signals. Our objectives were to systematically characterize signal features across species, discuss their potential as biomarkers and mechanistic indicators of SCS efficacy, and identify critical factors that influence their interpretation and applicability.

MATERIALS AND METHODS

Animals

Rat. Naïve adult male Sprague-Dawley rats (n = 25; 8-10 weeks; 300-440 g; Charles River Laboratories, Kent, UK) were acclimatized to the colony room for at least seven days after arrival and prior to the start of any experimental procedures. Rats were housed in groups of 3-4 animals per polyethene cage (Comparative Biology Centre, Newcastle University, Newcastle upon Tyne, UK). They were maintained on a 12-h day/night cycle (lights on at 6:00 am; lights off at 6:00 pm), and under controlled temperature (21 °C) and humidity (55%). Standard rodent food and water was accessible *ad libitum*.

Macaque. Adult male rhesus macaques (n = 2; ~7 years; 11.5 and 10.2 kg; Centre for Macaques, Salisbury, UK) were pair-housed in spacious pens exceeding UK and EU welfare standards, with access to natural daylight (Comparative Biology Centre, Newcastle University, Newcastle upon Tyne, UK). These animals were part of a separate, unrelated study involving unilateral corticospinal tract lesions, induced either by intracortical injections of endothelin-1 or by a thermocoagulation of the internal capsule. Water was available *ad libitum,* and food access was regulated to maintain motivation during the behavioral components of the primary study.

All experiments were conducted in accordance with the UK Animals (Scientific Procedures) Act 1986 under Home Office Project Licences P6694C943 and PP8237166, with local ethical approval from the Animal Welfare and Ethical Review Body (AWERB). This study adheres to the Animal Research: Reporting of In Vivo Experiments (ARRIVE) guidelines. Animal welfare was carefully monitored throughout the study, and all efforts were made to minimize suffering and reduce the number of animals used.

Lead Implantation and SCS Delivery

Rat. General anesthesia was induced via a nose cone using 5% isoflurane in oxygen (2 L/min flow rate). After induction, urethane (1.5 g/kg, i.p.) was administered alongside glycopyrrolate (0.015 mg/kg, s.c.) and anesthesia was maintained with 0-0.5% isoflurane in oxygen (2 L/min flow rate). All rats were implanted with two custom-made eight-electrode SCS leads (each electrode: 0.3×1.0 mm; Saluda Medical, Sydney, Australia) placed in the dorsal epidural space. Electrode placement was verified by X-ray imaging (Orange 1040HF, EcoRay, Seoul, South Korea) (Figure 1a). Each lead featured electrodes spaced 4 mm apart (center-to-center). Implantation procedures followed protocols established in our previous studies [1, 10]. One lead was positioned caudally, targeting spinal segments L2-T12, while the second lead was placed rostrally, spanning T12/T11 to T6 (Figure 2a). Both leads were connected to a custom-designed Multi-Channel System MKII (Saluda Medical, Sydney, Australia) as described in our earlier work [1, 2, 10]. Monopolar, biphasic stimulation was delivered in a stepwise manner with increasing current up to visible motor threshold (observed as a contraction in back or leg muscles), using a pulse width of 200 µs, at frequencies of 2 (n = 14) or 50 Hz (n = 11) (Figure 2a). Stimulation was delivered through each electrode consecutively, while recordings were simultaneously acquired from the remaining non-stimulating electrodes. Both the return and reference electrodes were placed subcutaneously. Therefore, none of the recorded signals were attributed to the reference electrode. None of the animals were allowed to recover from anesthesia at the end of the experiments and were humanely euthanized.

Recordings were averaged across approximately two traces at 2 Hz or fifty traces at 50 Hz per current level. Prior to analysis, the data were upsampled fivefold to improve temporal resolution; no offline filtering was applied. Signal processing and characterization were performed using a custom-built analysis toolbox (MATLAB 2013 release; MathWorks, Inc.) along with custom scripts developed in MATLAB 2024.

Macaque. Anesthesia was initially induced with ketamine (10 mg/kg, i.m.), supplemented by medetomidine (3 µg/kg, i.m.) and midazolam (0.3 mg/kg, i.m.) or propofol (1.3-4.4 mg/kg, i.m.). General anesthesia was maintained using sevoflurane (2-3% in 100% oxygen) and alfentanil (0.4-0.57 µg/kg/min, i.v.). To enhance analgesia, meloxicam (0.3 mg/kg, i.m.) and paracetamol (25 mg/kg, i.m.) were administered. Antibiotic prophylaxis was provided with either cefotaxime (20 mg/kg, i.v.) or co-amoxiclav (12.5-20 mg/kg, i.v.), and methylprednisolone (5.4 mg/kg/h, i.v.) was given to minimize cerebral edema. Fluid balance was maintained with Hartmann's solution (approximately 5 mL/kg/h, i.v.), with adjustments to maintain a total infusion rate of ~10 mL/kg/h, accounting for concurrent drug administration. A tracheotomy was performed to enable positive pressure artificial ventilation. A central arterial line was inserted via the carotid artery for continuous blood pressure monitoring, and the bladder was catheterized. Body temperature was maintained using a thermostatically controlled heat pad and a warm air blanket. A laminectomy was performed to expose the cervical and lumbar spinal cord. Following this, sevoflurane was discontinued (reduced to 0%), and anesthesia was transitioned to continuous infusions of alfentanil (0.4-1.2 µg/kg/min, i.v.), ketamine (6-10 mg/kg/h, i.v.), and midazolam (0.3 mg/kg/h, i.v.), a regimen previously shown to support stable anesthesia while preserving central nervous system activity. Throughout the procedure, physiological parameters were closely monitored, including pulse oximetry, heart rate, arterial blood pressure, core and peripheral temperatures, and end-tidal CO_2. Rapid increases in heart rate or blood pressure in response to noxious stimuli, or gradual upward trends in these measures, were interpreted as signs of inadequate anesthesia, prompting supplemental doses of injectable agents as needed. Macaques were implanted with a twelve-electrode SCS lead (1.3 mm diameter; Saluda Medical, Sydney, Australia) positioned in the dorsal epidural space. Lead placement was verified by X-ray imaging (Orange 1040HF, EcoRay, Seoul, South Korea) (Figure 1b). Electrodes were spaced 7 mm apart (center-to-center), spanning spinal vertebral levels T11 to L3 (Figure 3a). SCS was delivered using a clinical-grade neuromodulation system (Evoke System, Saluda Medical, Sydney, Australia). In line with clinically used settings and previous

studies [16], and to avoid the second cathode effect [21], tripolar, triphasic stimulation (with the active electrode located between two return electrodes) at 10 Hz with 100 μs pulse width or tripolar, biphasic stimulation at 3 Hz with 80 μs pulse width (Figure 3a) was delivered in a stepwise manner with increasing current up to 2 or 3×ECAP threshold. Stimulation was delivered through each electrode consecutively, while recordings were simultaneously acquired from the remaining non-stimulating electrodes. The position of the return electrodes for each recording are indicated in Figure 3. Neuromuscular blockade (atracurium; 0.75 mg/kg/h, i.v.) was administered during the acquisition of all electrophysiological signals except EMG, to minimize movement artifacts. None of the animals were allowed to recover from anesthesia at the end of the experiments and were humanely euthanized.

Recordings were averaged across approximately three traces at 3 Hz or ten traces at 10 Hz per current level. Data were analyzed without filtering or upsampling using a custom application (*Plot Builder*) in combination with custom scripts developed in MATLAB 2024 (MathWorks, Inc.).

In both species, the threshold for each electrophysiological signal was defined as the lowest stimulation current at which the signal could be reliably detected [1,10]. Signal amplitudes (mV) were calculated as the voltage difference between characteristic peaks: P2-N1 for ECAPs, $P2_d$-$N1_d$ for doublets (d refers to secondary ECAP), the maximum positive peak minus the preceding negative peak for EMG signals. For ESAPs, amplitude was defined as the voltage measured at the maximum point of the S1 peak. Conduction velocity (m/s) was estimated by measuring the latency of characteristic peaks, N1 for ECAPs and $N1_d$ for doublets, across spatially separated recording electrodes.

Figure 1. Representative lead positioning in the spinal cords of a rat and macaque. **a.** A representative dorsal X-ray image from a rat showing the placement of two epidural leads and confirming coverage from vertebral levels L2-T12 (lead 1) and T11-T6 (lead 2). Each electrode on the lead, along with the corresponding vertebral segments, is labelled (lead 1: 1-6, lead 2: 9-14). **b.** A representative dorsal X-ray image from a macaque showing the epidural lead covering the L3 to T11 vertebral levels. Each electrode and corresponding vertebral level are labelled (1-12). Scale bar: 10 mm.

RESULTS

ECAP

ECAPs were reliably evoked and recorded across multiple segmental levels of the spinal cord during SCS in both rats (L1-T8) and macaques (L2-T11). As the earliest electrophysiological responses detected in epidural spinal recordings, ECAPs were distinguished by their short latencies relative to other spinal signals described in this study (Figures 2b and 3b). Morphologically, ECAPs exhibited a characteristic triphasic waveform, consisting of an initial positive peak (P1), followed by a negative peak (N1), and a second positive peak (P2), with a total duration of approximately 1 ms from P1 to P2 (Figures 2b and 3b). The ECAP threshold was 32.00 ± 3.10 µA in rats (n = 18) and 375.00 ± 25.00 µA in macaques (n = 2).

As illustrated in Figure 4a for rats, ECAP amplitude increased linearly with stimulation intensity following suprathreshold activation, up to the motor threshold. Although the waveform remained consistent across recording sites (L1-T8), electrodes located farther from the stimulation source exhibited longer latencies and reduced amplitudes (Figure 2b and 2c). Comparable results were observed in macaques (Figure 3b, 3c and 5a). Average conduction velocities were 30.70 ± 1.81 m/s in rats (n = 18) (Figure 4b) and 82.75 ± 2.46 m/s in macaques (n = 2) (Figure 5b).

Doublet

Doublets appeared as complex signals characterized by multiple negative peaks, two in the rat and three in the macaque, representing two (or three) separate ECAPs. In Figure 2c distinct primary (N1) and secondary ($N1_d$) peaks are observed at levels T11 to T8, whereas beyond T8 only the continuation/propagation of $N1_d$ peak was observable and the primary N1 peak could no longer be observed. In Figure 3c, N1 and $N1_d$ peaks are clearly distinguishable at levels L2 to T12, but only $N1_d$ peaks are observable at T12 to T11 (distances of 63 and 70 mm away from stimulation site). The secondary ECAP, elicited 1.65

ms in the rat and 1.36 ms in the macaque after the primary ECAP was reliably detected in both species (Figures 2c and 3c; asterisk). In macaques, a tertiary ECAP was also observed, emerging around 2 ms after the primary ECAP and 1 ms after the secondary ECAP (Figure 3c; downward triangle). In one macaque, secondary and tertiary ECAPs were observed at a distance between 28 and 70 mm away from stimulation site (Figure 3c). In the second macaque, secondary ECAPs were also observed between 28 and 70 mm away, however tertiary ECAPs were only observed beyond approximately 77 mm away from stimulation (data not shown).

Doublets were only evoked when stimulation was applied to the lower thoracic to upper lumbar spinal regions (shown in Figure 2c as L2 and Figure 3c as L3). They were recorded exclusively in the orthodromic direction (rostral), predominantly within the lower to upper thoracic spinal cord (T11-T6 in Figure 2c), though they were also detected in the upper lumbar region in the macaques (L1-T11 in Figure 3c). Both the secondary and tertiary ECAPs retained the triphasic morphology characteristic of the primary ECAP. The threshold for secondary ECAP was 143 µA (n = 1) in the rat, corresponding to a primary ECAP threshold of 105 µA (n = 1), recorded on the same electrode. In macaques, secondary ECAP thresholds were 550 µA (n = 1) and 950 µA (n = 1), while primary ECAP thresholds on the same electrode were 550 µA (n = 1) and 400 µA (n = 1), respectively. In the macaque exhibiting tertiary ECAPs at 35 mm, the threshold was approximately 700 µA (n = 1).

Their amplitudes (measured as $P2_d$-$N1_d$ in mV for the secondary, and $P2_d$-$N1_t$ in mV for the tertiary; t refers to tertiary ECAP) increased in response to rising stimulation current (Figure 4c and 5c). Unlike the primary ECAP, the amplitudes of these later components increased with recording distance (Figures 2c and 3c). Similar to the primary ECAP, both secondary and tertiary components demonstrated clear propagating behavior, with latencies increasing at electrodes positioned farther from the stimulation site (Figure 4d and 5d). Conduction velocity of the secondary ECAP was 33.62 m/s (n = 1) in the rat, while primary ECAP

conduction velocity was 15.28 m/s (n = 1) (Figure 4d). In macaques, secondary ECAP conduction velocities measured 81.54 m/s and 108.07 m/s, with corresponding primary ECAP velocities of 54.81 m/s and 46.74 m/s (n = 1) (Figure 5d). In one of the macaques, tertiary ECAPs exhibited a conduction velocity of 79.49 m/s (n = 1) (calculated from $N1_t$ peaks).

ESAP

ESAPs were detectable in rats (Figure 2d; star) but were not observed in macaques. Morphologically, ESAPs were defined by a positive S1 wave that appeared at a longer latency (~3-4 ms) following a secondary negative peak (N2) in the epidural recordings. The S1 wave exhibited a prolonged duration of approximately 6 ms, in contrast to the ~1 ms duration of ECAPs. The average ESAP threshold was 91.00 ± 12.00 µA (n = 4).

Similar to ECAPs, the S1 wave amplitude increased in response to rising stimulation current (Figure 4e). However, unlike ECAPs, ESAP amplitudes did not consistently decrease with increasing distance from the stimulation site. Instead, ESAPs exhibited a spatially localized maximum, with the largest signals consistently recorded at a specific electrode (e.g., electrode 3 located 8 mm from the stimulation site at L2; Figure 2d). ESAPs were only detectable when the recording contact was positioned over the L1/T13 vertebrae, with peak responses consistently observed at the L1 level, suggesting that anatomical specificity may be a defining characteristic of ESAPs. Furthermore, ESAP latencies did not increase with distance from the stimulation site, consistent with their non-propagating nature (Figure 4f).

EMG

EMG signals were detected in epidural recordings and were distinct from ECAPs, doublets and ESAPs, in morphology and stimulus-response characteristics. EMG signals appeared as either biphasic or triphasic waveforms 3-5 ms after the stimulus, which is approximately 2-4

ms after the primary ECAP (Figure 2e and 3d). These signals had a slightly longer duration than ECAPs (~2 ms).

Similar to ECAPs, EMGs were evoked and recorded across multiple spinal levels in the rat (L1-T7; Figure 2e) and the macaque (L2-L3; Figure 3d). The EMG threshold was 245 µA in the rat (n = 1) and 900 µA in the macaque (n = 1).

A key distinguishing feature of EMG signals was their step-like recruitment pattern, characterized by the abrupt appearance of EMG responses once the stimulation threshold was reached, rather than a gradual increase with stimulation intensity. In both species, EMG amplitudes were consistent across recording sites, showing no change in amplitude (in channels where the signal was detected) as the distance from the stimulation site increased (Figure 2e and 3d). Unlike ECAPs and doublets, EMGs were non-propagating, meaning their latency remained constant regardless of the distance of the recording electrode from the stimulation site.

Figure 2. Epidural spinal electrophysiological signals evoked by spinal cord stimulation in rats. **a.** Schematic illustration of the experimental electrode setup, showing the distances (in mm) between the stimulating and recording electrodes, along with their corresponding vertebral level placements. Two leads were implanted epidurally: one caudally (T12-L2) and one rostrally (T11-T6). Stimulation was delivered through each electrode consecutively, while recordings were simultaneously acquired from the remaining electrodes. Return and reference electrodes were placed subcutaneously. Data presented in the figure refer to stimulation delivered through the most caudal electrode and recordings obtained from all remaining electrodes on both leads. Monopolar, biphasic stimulation was delivered in a stepwise manner (1 µA) with increasing current up to visible motor threshold, using a pulse width of 200 µs, at frequencies of 2 or 50 Hz. *Created with BioRender.com.* Evoked compound action potential (ECAP; **b**), doublet (**c**), evoked synaptic activity potential (ESAP; **d**), and electromyography (EMG) signals (**e**) recorded at distances of 4-24 mm (ECAP, ESAP, and EMG) or 24-44 mm (doublet) from the stimulating electrode. The N1 peaks are marked with dots (**b-e**), the secondary N1 peaks (N1$_d$) with asterisks (**c**), the S1 waves with stars (**d**), and the EMG maxima with upward triangles (**e**). Each trace represents the average of two recordings at 0.23 mA (**b-d**) and fifty recordings at 0.30 mA (**e**), each evoked by separate stimulation pulses delivered at motor threshold within a 1-s period in a single rat. Insets highlight the characteristic morphology of each signal type. The solid line represents the lead implanted caudally, while the dotted line represents the lead implanted rostrally.

Figure 3. Epidural spinal electrophysiological signals evoked by spinal cord stimulation in macaques. **a.** Schematic illustration of the experimental electrode setup, showing the vertebral level placements. The lead was implanted epidurally (L3-T11). For the signals shown, stimulation at 1.5×Evoked Compound Action Potential (ECAP) threshold was delivered through different electrodes in a consecutive manner, while recordings were obtained from the remaining electrodes. The stimulation electrode is indicated for each signal in the schematic illustrations of the leads. Return electrodes are represented by black rectangles (as shown in **b-d**). Reference electrode was positioned on a second rostrally implanted lead at the furthest away contact. Stimulation was applied as tripolar, triphasic pulses at 10 Hz (100 μs) (**b, c**) or tripolar, biphasic pulses at 3 Hz (80 μs) (**d**). *Created with BioRender.com.* Evoked compound action potential (ECAP; **b**), doublet (**c**),

and electromyography (EMG) signals (**d**) were recorded at distances of 28-49 mm (ECAP), 28-70 mm (doublet), and 42-63 mm (EMG) from the stimulating electrode. The N1 peaks are indicated by dots (**b-d**), the secondary N1 peaks (N1$_d$) by asterisks (**c**), the tertiary N1 peaks by downward triangles (N1$_t$) (**c**), and the EMG maxima by upward triangles (**d**). Each trace represents the average from a single macaque, recorded at a current of 0.75 mA (1.5×ECAPT) for panels **b-c**, and 1.7 mA for panel **d**. Insets emphasize the characteristic morphology of each signal type.

Figure 4. Spinal cord stimulation (SCS)-evoked neural responses in rats. **a.** Mean evoked compound action potential (ECAP) amplitude as a function of stimulation current during 2 Hz and 50 Hz SCS recorded orthodromically 4 mm from the stimulation site (n = 18). **b.** Mean ECAP distance vs. latency during 2 Hz and 50 Hz SCS across rats (n = 18; mean. 30.70 ± 1.81 m/s) calculated from N1 peaks. The inset shows ECAP distance vs. latency from a rat exhibiting doublets in panel c. Filled circles represent N1 latencies calculated from the caudally implanted lead, whereas open circles represent N1 latencies calculated from the rostrally implanted lead. The slope of the linear fit indicates the conduction velocity. **c.** Doublet amplitude vs. stimulation current during 2 Hz SCS (n = 1), recorded orthodromically ~28 mm from stimulation site. Gray shows primary ECAP, and yellow shows secondary ECAP. **d.** Doublet distance vs. latency during 2 Hz SCS (n = 1). The slope of the linear fits indicates the conduction velocities. Primary and secondary ECAP conduction velocities were 15.28 and 33.62 m/s, respectively. Primary ECAP conduction velocity was calculated from N1 peaks, while secondary ECAP conduction velocity was calculated

from $N1_d$ peaks. In the inset, filled circles represent N1 latencies calculated from the caudally implanted lead, open circles represent N1 latencies calculated from the rostrally implanted lead, whereas yellow filled circles represent $N1_d$ latencies recorded from the rostrally implanted lead. **e**. Mean evoked synaptic activity potential (ESAP) amplitude vs. stimulation current during 2 Hz and 50 Hz SCS across rats (n = 4), recorded orthodromically 8 mm from stimulation site. **f.** Mean ESAP distance vs. latency during 2 Hz and 50 Hz SCS across rats (n = 4), recorded orthodromically. All data were collected with 200 µs pulse width. Stimulation current was normalized between relevant threshold (0) and motor threshold (1) (**a, c, e**). Distance vs. latency were plotted and conduction velocity was calculated at motor threshold. Standard error of the mean (± SEM) is indicated by the shaded region (**a**, **e**) and error bars (**b**, **f**).

Figure 5. Spinal cord stimulation (SCS)-evoked neural responses in macaques. **a.** Mean evoked compound action potential (ECAP) amplitude as a function of stimulation current during 10 Hz SCS (n = 2), recorded orthodromically from the electrode 28 mm away from stimulation site. Stimulation current was normalized between relevant threshold (0) and 2×ECAP threshold (1). **b.** Mean ECAP distance vs. latency during 10 Hz SCS across macaques (n = 2; mean. 82.75 ± 2.46 m/s), calculated from N1 peaks. Gray dots and dashed line show group mean. The slope of the linear fit indicates the conduction velocity. **c.** Doublet amplitude vs. stimulation current during 10 Hz SCS (n = 1), recorded orthodromically 28 mm from stimulation site. Gray shows primary ECAP data, and yellow shows secondary ECAP data. Stimulation current was normalized between relevant threshold (0) and 3×ECAP threshold (1). **d.** Doublet distance vs. latency during 10 Hz SCS (n = 1). The slope of the linear fits indicates the conduction velocities. Primary and secondary ECAP conduction velocities were 46.74 and 108.07 m/s, respectively. Primary ECAP conduction velocity was calculated from N1 peaks, while secondary ECAP conduction velocity was calculated from $N1_d$ peaks. All data were collected with 100 μs (**a-d**) pulse width. Standard error of the mean (± SEM) is indicated by the shaded region (**a**).

DISCUSSION

We present here a summary of the distinct electrophysiological signals evoked and recorded epidurally during SCS, each distinguishable by unique waveform features and stimulus-response characteristics, including latency and amplitude. This study highlights their defining properties and provides representative examples to facilitate interpretation (Table 1). By incorporating data from both rats and macaques, we enable cross-species comparisons that may offer translational insights. Below, we further discuss the relevance of each signal to spinal processing. Unlike ECAPs, other evoked signals may be affected by aberrant spinal circuitry, particularly in maladaptive states such as neuropathic pain, and thus may offer a deeper understanding not only of the mechanisms underlying SCS but also of the pathophysiology of chronic pain.

ECAP

ECAPs recorded in this study are consistent with those reported previously across multiple species, including rodents, large animals (e.g., sheep and pig), non-human primates and humans. The characteristic triphasic morphology (P1-N1-P2) observed here aligns with typical extracellular recordings of axonal CAPs, where the P1 peak reflects the initial depolarization, the N1 valley corresponds to axonal depolarization driven by Na^+ influx, and the P2 peak reflects the subsequent hyperpolarization driven by K^+ efflux [2]. This morphology is conserved across species, supporting the cross-species reliability of ECAP recordings [2].

Consistent with previous studies, ECAPs recorded during SCS primarily reflect the activation of fast-conducting DC fibers, specifically large-diameter myelinated Aβ afferents, as confirmed by conduction velocity measurements [1-5, 14]. A novel observation in this study is the apparent slowing of ECAP conduction velocity with increasing distances, observed in both rats (Figure 4b and 4d) and macaques (Figure 5b and 5d). Similar findings have been reported in anaesthetized cats, where the conduction velocity of group I and II afferent fibers

progressively decreased as they ascended the DC [22]. One possible explanation for this deceleration is that larger-diameter axons terminate at lower spinal levels; less than 25% of lumbar DC axons reach the brainstem [23]. Supporting this, a study by Yamamoto et al. (1996) in rats demonstrated that only 50.1% of axons present in the DC at the L3 level ascend to the T6 level, with most large-diameter fibers terminating or thinning out between these regions [24]. We also observed a step change in conduction velocity in both models, consistent with a previous report in sheep, where such discontinuities were hypothesized to result from the segmental termination of fibers at discrete spinal levels [13]. Further investigation is needed to confirm these findings and to characterize this phenomenon in human subjects.

In addition to conduction velocity, changes in ECAP morphology, such as amplitude and width, may provide insight into DC fiber dynamics and their modulation by SCS parameters. For example, high-frequency SCS at 200, 500, and 1000 Hz reduced ECAP amplitude and increased peak latencies and ECAP width, effects likely reflecting asynchronous firing of DC axons [14]. These findings highlight ECAPs as an indicator of DC activation and support their utility in advancing both the mechanistic understanding and therapeutic optimization of SCS.

Doublet

Here, we observed secondary ECAPs in both rodents and macaques that share core characteristics with previously described PSDC signals in humans [16]. Additionally, in macaques, we identified a tertiary ECAP which exhibited characteristics closely resembling those of secondary ECAPs, albeit occurring at a later latency. Maruyama et al. (1982) have previously described a similar signal recorded with epidural electrodes in response to cauda equina stimulation in humans [25].

Doublets reported in this study were exclusively evoked when stimulating at lower vertebral levels and recorded at upper levels, in the orthodromic (rostral) direction. These responses exhibited higher conduction velocities compared to the N1 component of primary ECAPs, consistent with previous findings in humans [16]. The presence of these additional negative peaks suggests activation of spinal projection neurons, which are likely recruited following electrical stimulation of DC axons and may represent output from endogenous spinal processing. The administration of a neuromuscular blocker in the macaques used when obtaining all recordings except from the EMG signal in this study, effectively rules out any potential contamination of the signals by myogenic activity. Moreover, our interpretation aligns with previous studies reporting PSDC neurons to be predominantly found in the lumbar and cervical enlargements in animals [26, 27]. While doublets described here are consistent with the hypothesized activation of the PSDC pathway, contributions from alternative spinal pathways involving postsynaptic activation, such as the spinothalamic tract [17], cannot be ruled out, particularly given that doublets exhibit higher activation thresholds compared to primary ECAPs.

Gmel et al. (2023) found no evidence that secondary ECAP signals in humans resulted from direct activation of nerve rootlets, based on propagation times for the secondary ECAP [16]. A comparable calculation, extrapolating the two trendlines to their y-intercepts, shows that in the rat the intercepts differ by 61.6 mm. This would require stimulation pulses to activate a fiber bundle located 6.16 cm away, without affecting any intervening fibers. In the monkey, the intercepts are 134 mm apart, implying a required activation distance of 13.4 cm under the same unlikely condition. Furthermore, care must be taken with stimulating contact configurations (cathodes and anodes) and charge delivery, since second-cathode effects can manifest as multimodal signals [21, 28].

ESAP

First identified and characterized by Sharma et al. (2023) using epidural electrodes in rat, the ESAPs we report here, also recorded in rats, are consistent with those previously described [15]. Features of the S1 component are reported to be consistent with intracellularly recorded monosynaptic responses of dorsal horn neurons in *in vitro* models following DC stimulation [29]. Although not demonstrated in the present study, the pharmacological sensitivity of ESAPs to AMPA receptor blockade provides strong evidence that they originate from synaptic, rather than axonal, activity [15]. Sharma et al. further verified that the S1-wave was not a stimulation artifact and/or a reflection of hindlimb/trunk EMG [15]. Importantly, the spatial specificity of ESAP detection, unlike the widespread propagating nature of ECAPs, suggest that ESAPs reflect local connectivity (circuits) (Table 1) [15]. Additionally, Sharma et al. demonstrated that ESAPs, but not ECAPs, exhibited gradual adaptation in response to 50 Hz stimulation [15]. This habituation is consistent with, though not exclusive to, synaptic processing mechanisms and mirrors findings from superficial dorsal horn interneuron responses to 50 Hz SCS [15, 30-32]. Together, these observations support the hypothesis that ESAPs may serve as an epidurally measurable marker of dorsal horn activity in rodents.

However, the absence of ESAPs in our macaque model may reflect species-dependent differences in spinal processing, limitations in recording sensitivity (e.g., CSF thickness separating the dorsal horn from recording electrodes), or differences in experimental design, particularly the stimulation waveform used. This highlights the need for further investigation into the conditions required to detect these signals in higher-order species. Notably, classical slow-wave signals were consistently observed in human spinal cord neurophysiology studies [30, 33, 34] and indeed formed the foundation for the development of gate-control theory and, ultimately, SCS [35].

EMG

Building on the identification of neural signals such as ECAPs, doublets and ESAPS, it is also essential to distinguish EMG signals which can be detected via the epidural lead. EMGs are often detected within the same latency as the other neural responses and can introduce a source of signal contamination, especially when interpreting overlapping components [28]. However, the non-propagating nature of EMGs, along with their step-like appearance, higher activation thresholds and consistent amplitude across recording sites, help differentiate them from neural signals such as ECAPs, doublets and ESAPs. EMG signals can be observed across multiple species, including rodents, large animals (e.g., sheep and pig), non-human primates and humans, confirming their relevance in both preclinical and clinical contexts.

It remains to be confirmed whether EMG signals induced by SCS result from direct activation of motor neuron axons, as they exit the anterior spinal cord via the ventral roots, or from monosynaptic activation of motor neurons following stimulation of proprioceptive afferents in the dorsal roots. In peripheral nerve studies, the H-wave typically has a threshold lower than the M-wave, reflecting the greater excitability of afferent fibers compared to efferent fibers [36]. Given the dorsal roots' closer proximity to the stimulating electrodes relative to motor axons in the ventral roots, SCS-induced EMG signals in humans and large animal models are most likely reflex-mediated. In rodents, however, the deep DCs contain descending corticospinal fibers that may be directly activated at higher SCS intensities, subsequently activating motoneurons [37]. Thus, similar to ESAPs and potentially doublets, EMG signals likely represent an output of spinal processing.

CONCLUSIONS

Epidural spinal recordings, including ECAPs, doublets, ESAPs, and EMGs provide insights into the neural processes engaged by SCS. This study summarizes key characteristics of these signals, as outlined in Table 1, to support their differentiation and classification in future studies based on waveforms, recruitment thresholds, and physiological origins. As ESAPs, EMG signals, and potentially doublets, represent outputs of spinal processing

detectable via epidural recordings, they may offer insights into the pathophysiology of conditions involving disrupted spinal function. One example is diabetic neuropathic pain, which may arise from spinal cord disinhibition [38]. Continued investigation into the origins, functional relevance, and clinical significance of these signals may provide critical insights into SCS mechanisms, inform therapy optimization, and guide the development of more effective SCS interventions. Species-specific patterns, potentially reflecting neuroanatomical or technical differences that influence signal detectability in preparations of varying size, emphasize the importance of cross-species analyses to improve the translational relevance of neuromodulation research.

Data statement

The datasets used and/or analyzed during the current study are available from the corresponding author on reasonable request.

Declaration of generative AI and AI-assisted technologies in the writing process

During the preparation of this work, the authors used ChatGPT in order to improve the readability and language of the manuscript. After using this tool, the authors reviewed and edited the content as needed and take full responsibility for the content of the publication.

Glossary

ARRIVE	Animal Research: Reporting in *In Vivo* Experiments
AWERB	Animal Welfare Ethical Review Body
CAP	Compound Action Potential
DC	Dorsal Column
ECAP	Evoked Compound Action Potential
ECAPT	Evoked Compound Action Potential Threshold
EMG	Electromyography
ESAP	Evoked Synaptic Activity Potential
N1	First Negative Peak
P2	Second Positive Peak

SCS   Spinal Cord Stimulation

SEM   Standard   Error   of   the   Mean

Table 1. A summary of the defining characteristics of electrophysiological signals detected in epidural spinal recordings, including evoked compound action potentials (ECAPs), doublets, evoked synaptic activity potentials (ESAPs), and electromyographic (EMG) signals, as identified in current and previous

Characteristics	ECAP	Doublet	ESAP*	EMG
Morphology	Triphasic signal (P1, N1 and P2 peaks)	A secondary ECAP following a primary ECAP signal (N1$_d$ and P2$_d$ peaks)	S1 wave occurs after the N2 peak, following the ECAP signal	Biphasic or triphasic signal
Latency	Occurs first**	~1-2 ms after N1	~3-4 ms after N1	~3-4 ms after N1
Duration	~1 ms	Like primary ECAP (~1 ms)	Prolonged duration (~6 ms)	~2 ms
Threshold	Lower than ESAP threshold and significantly lower than EMG threshold	Similar to ECAP threshold or higher	Higher than ECAP threshold but significantly lower than EMG threshold	Significantly higher than ECAP and ESAP thresholds
Region to Evoke	All vertebral levels	Lower thoracic and upper lumbar vertebral levels	Lower thoracic and upper lumbar vertebral levels	All vertebral levels
Region to Record	All vertebral levels	Upper thoracic vertebral levels	Upper lumbar vertebral levels (~T13-L2)	All vertebral levels
Direction of Observation	Orthodromic and antidromic	Orthodromic	Orthodromic and antidromic	Orthodromic and antidromic
Propagation	Yes	Yes	No	No
Latency × Distance	Peak latencies increase with greater electrode distance	Peak latencies increase with greater electrode distance. Secondary ECAP peak latencies 'catch up' with primary latencies as distance increases orthodromically	Peak latencies remain constant with greater electrode distance	Peak latencies remain constant with greater electrode distance
Latency × Current	Peak latencies remain constant with greater current	Peak latencies remain constant with greater current	S1 latency decreases with greater current	Peak latencies remain constant with greater current
Amplitude × Distance	Amplitude (P2-N1) decreases with greater electrode distance	Amplitude (P2$_d$-N1$_d$) increases with greater electrode distance	Region specific	Amplitude remains constant across varying electrode distance
Amplitude × Current	Amplitude (P2-N1) increases with greater current	Amplitude (P2$_d$-N1$_d$) increases with greater current	Amplitude (S1 peak) increases with greater current	Step response instead of linear growth with current

studies, with species indicated by the corresponding icons.

*Tested but not identified in rhesus macaques. **ECAP latency typically ranges ~0.3-1.5 ms, depending on factors like electrode placement, neural conduction velocity, stimulation parameters and recording set-up.

REFERENCES

1. Dietz BE, Mugan D, Vuong QC, Obara I. Electrically Evoked Compound Action Potentials in Spinal Cord Stimulation: Implications for Preclinical Research Models. Neuromodulation. 2022;25(1):64-74.

2. Parker JL, Karantonis DM, Single PS, Obradovic M, Laird J, Gorman RB, et al. Electrically evoked compound action potentials recorded from the sheep spinal cord. Neuromodulation. 2013;16(4):295-303.

3. Sagalajev B, Zhang T, Abdollahi N, Yousefpour N, Medlock L, Al-Basha D, et al. Absence of paresthesia during high-rate spinal cord stimulation reveals importance of synchrony for sensations evoked by electrical stimulation. Neuron. 2023;112:1-17.

4. Gmel GE, Santos Escapa R, Parker JL, Mugan D, Al-Kaisy A, Palmisani S. The Effect of Spinal Cord Stimulation Frequency on the Neural Response and Perceived Sensation in Patients With Chronic Pain. Front Neurosci. 2021;15:625835.

5. Parker JL, Karatonis DM, Single PS, Obradovic M, Cousins MJ. Compound action potential recorded in the human spinal cord during neurostimulation for pain relief. Pain 2012;153(3):593-601.

6. Heijmann L, Joosten EA. Mechanisms and mode of action of spinal cord stimulation in chronic neuropathic pain. Postgrad Med. 2020;132:17-21.

7. Vallejo R, Chakravarthy K, Will A, Trutnau K, Dinsmoor D. A New Direction for Closed-Loop Spinal Cord Stimulation: Combining Contemporary Therapy Paradigms with Evoked Compound Action Potential Sensing. J Pain Res. 2021;14:3909-18.

8. Nijhuis H, Hofsté W, Huygen F, al e. First report on real-world outcomes with evoked compound action potential (ECAP)-controlled closed-loop spinal cord stimulation for treatment of chronic pain. Pain Ther. 2023;12(5):1221-33.

9. Parker J, Karantonis D, Single P. Hypothesis for the mechanism of action of ECAP-controlled closed-loop systems for spinal cord stimulation. Healthc Technol Lett. 2020;7(3).

10. Versantvoort EM, Dietz BE, Mugan D, Vuong QC, Luli S, Obara I. Evoked compound action potential (ECAP)-controlled closed-loop spinal cord stimulation in an experimental model of neuropathic pain in rats. Bioelectron Med. 2024;10(1):2.

11. Mekhail N, Levy R, Deer TR, Kapural L, Li S, Amirdelfan K, et al. ECAP-controlled closed-loop versus open-loop SCS for the treatment of chronic pain: 36-month results of the EVOKE blinded randomized clinical trial. Reg Anesth Pain Med. 2024;49(5).

12. Muller L, Pope J, Verrills P, Petersen E, Kallewaard JW, Gould I, et al. First evidence of a biomarker-based dose-response relationship in chronic pain using physiological closed-loop spinal cord stimulation. Reg Anesth Pain Med. 2024;0:1-7.

13. Parker JL, Obradovic M, Hesam Shariati N, Gorman RB, Karantonis DM, Single PS, et al. Evoked Compound Action Potentials Reveal Spinal Cord Dorsal Column Neuroanatomy. Neuromodulation. 2020;23(1):82-95.

14. Ladner K, Versantvoort EM, Mugan D, Vuong QC, Dietz BE, Hu A, et al. Preclinical Insights into the Effects of Frequency and Pulse Width on Evoked Compound Action Potential Morphology during Spinal Cord Stimulation. 2025.

15. Sharma M, Bhaskar V, Yang L, FallahRad M, Gebodh N, Zang T, et al. Novel Evoked Synaptic Activity Potentials (ESAPs) Elicited by Spinal Cord Stimulation. eNeuro. 2023;10(5).

16. Gmel GE, Santos Escapa R, Benkohen TE, Mugan D, Parker JL, Palmisani S. Postsynaptic dorsal column pathway activation during spinal cord stimulation in patients with chronic pain. Frontiers in Neuroscience. 2023;Volume 17 - 2023.

17. Wang L-H, Ding W-Q, Sun Y-G. Spinal ascending pathways for somatosensory information processing. Trends in Neuroscience. 2022;45(8):594-607.

18. Lindley D. Short-Term Outcomes of a High-Volume,Low-Concentration Bolus Starting DoseTechnique With Ziconotide: A Case Series. Neuromodulation. 2021;24(7):1209-14.

19. Deshmukh A, Settell M, Cheng K, Knudsen B, Trevathan J, LaLuzerne M, et al. Epidural spinal cord recordings (ESRs): sources of neural-appearing artifact in stimulation evoked compound action potentials. Journal of Neural Engineering. 2025;22(1).

20. Single P, Scott J. Cause of Pulse Artefacts Inherent to the Electrodes of Neuromodulation Implants. IEEE Transactions on Neural Systems and Rehabilitation Engineering. 2018;26(10):2078-83.

21. Brucker-Hahn MK, Zander HJ, Will AJ, Vallabh JC, Wolff JS, Dinsmoor DA, et al. Evoked compound action potentials during spinal cord stimulation: effects of posture and pulse width on signal features and neural activation within the spinal cord. Journal of Neural Engineering. 2023;20.

22. Fern R, Harrison PJ, Riddell JS. The dorsal column projection of muscle afferent fibres from the cat hindlimb. Journal of Physiology. 1988;401(1):97-113.

23. Horch KW, Burgess PR, Whitehorn D. Ascending collaterals of cutaneous neurons in the fasciculus gracilis of the cat. Brain Research. 1976;117(1):1-17.

24. Yamamoto K, Ohnishi A. Ascending myelinated axons of primary sensory neurons of the sciatic nerve in the posterior column of the rat spinal cord. Acta Neuropathologica. 1996;92:27-34.

25. Maruyama Y, Shimoji K, Shimizu H, Kuribayashi H, Fujioka H. Human spinal cord potentials evoked by different sources of stimulation and conduction velocities along the cord. Journal of Neuroscience. 1982;48(5).

26. Giesler GJ, Nahin RL, Madsen AM. Postsynaptic dorsal column pathway of the rat. I. Anatomical studies. J Neurophysiol. 1984;51(2):260-75.

27. Enevoldson TP, Gordon G. Postsynaptic dorsal column neurons in the cat: a study with retrograde transport of horseradish peroxidase. Exp Brain Res. 1989;75(3):611-20.

28. Deshmukh A, Settell M, K. C, B. K, Trevathan J, LaLuzerne M, et al. Epidural spinal cord recordings (ESRs): sources of neural-appearing artifact in stimulation evoked compound action potentials. Journal of Neural Engineering. 2025;22(1).

29. Baba H, Yoshimura M, Nishi S, Shimoji K. Synaptic responses of substantia gelatinosa neurones to dorsal column stimulation in rat spinal cord in vitro. Journal of Physiology. 1994.

30. Tomita M, Shimoji K, Denda S, Tobita T, Uchiyama S, Baba H. Spinal tracts producing slow componenets of spinal cord potentials evoked by descending volleys in man. Electroencephalography and clinical Neuroscience. 1996;100(1):68-73.

31. Sdrulla A, Xu Q, He S, Tiwari V, Yang F, Zhang C, et al. Electrical stimulation of low-threshold afferent fibers induces a prolonged synpatic depression in lamina II dorsal horn neurons to high-threshold afferent inputs in mice. Pain. 2015;156:1008-17.

32. Fan W, Sullivan SJ, Sdrulla A. Dorsal column and root stimulation at Aβ-fiber intensity activate superficial dorsal horn glutamatergic and GABAergic populations. Molecular Pain. 2022;18:1-13.

33. Yates BJ, Thompson FJ, Mickle JP. Origin and properties of spinal cord field potentials. Neurosurgery. 1982;11(3):439-50.

34. Wall PD. The origin of a spinal-cord slow potential. Journal of Physiology. 1962;164(3):508-26.

35. Bikson M, Sharma M. A Brief History of Slow Spinal Potentials, Gate Theory of Pain, and Spinal Cord Stimulation. Neuromodulation. 2024;27(5):946-8.

36. Palmieri RM, Ingersoll CD, Hoffman MA. The Hoffmann Reflex: Methodologic Considerations and Applications for Use in Sports Medicine and Athletic Training Research. Journal of Athletic Training. 2004;39(3):268-77.

37. Krisa L, Runyen M, Detloff MR. Translational Challenges of Rat Models of Upper Extremity Dysfunction After Spinal Cord Injury. Topics in Spinal Cord Injury Rehabilitation. 2018;24(3):195-205.

38. Marshall A, Kalteniece A, Ferdousi M, Azmi S, Jude EB, Adamson C, et al. Spinal disinhibition: evidence for a hyperpathia phenotype in painful diabetic neuropathy. Brain Communications. 2023;5(2).