Title: Specific and versatile monoclonal antibodies for hantavirus research

Running title: Novel anti-hantavirus antibodies

Authors: Autumn LaPointe[a], Kimberly Martinez[a], Christina Shou[a], Inessa Manuelyan[b], Jason

Botten[b,c], Alison M. Kell[a#]

Affiliations:

[a] Department of Molecular Genetics and Microbiology, University of New Mexico School of

Medicine, Albuquerque, New Mexico, USA

[b] Department of Medicine, Division of Pulmonary and Critical Care Medicine, Robert Larner,

M.D. College of Medicine, University of Vermont, Burlington, VT, USA.

[c] Department of Microbiology and Molecular Genetics, Robert Larner, M.D. College of Medicine,

University of Vermont, Burlington, VT, USA.

[#] amkell@salud.unm.edu

Abstract word count: 249

Text word count: 2,673

Abstract

Rodent-borne hantaviruses pose a continual public health threat to humans through zoonotic

transmission, with case fatality rates of up to 50% in some cases. Human infections can lead to

hemorrhagic fever with renal syndrome (HFRS) or hantavirus cardio-pulmonary syndrome

(HCPS), depending on the viral species. Despite the morbidity and mortality associated with this

family of viruses, no anti-viral therapeutics or vaccines are available to treat and prevent

hantavirus disease. The relative shortage of commercially available reagents to study

hantavirus infections *in vitro* and *in vivo* likely contributes to the challenges in developing viral

countermeasures. This report describes the generation of a panel of mouse monoclonal

antibodies that collectively recognize the four viral proteins of Seoul virus (*Orthohantavirus*

seoulense), an Old World hantavirus with worldwide distribution and the causative agent of

HFRS. We have validated the specificity and versatility of these antibodies against a subset of

Old World and New World hantaviruses in assays relying on antigen recognition in denatured or

native conformations. We present several antibodies that specifically recognize the Seoul virus

nucleoprotein and polymerase protein in Western blotting and immunostaining assays. We also

identified three novel antibodies directed against the glycoprotein complex that are capable of

binding to the N-terminal glycoprotein of all hantaviruses tested. These antibodies are freely

available to all hantavirus researchers to add to the small, but growing, collection of reliable and

available reagents to be used to study hantavirus biology, identify novel antiviral compounds,

and measure viral prevalence in the laboratory and the field.

Importance

Pathogenic hantaviruses cause severe hemorrhagic disease and pose a significant public

health threat world-wide. Insufficient research into the biology of these viruses has slowed the

development of effective direct-acting antivirals and vaccines. Here, we describe the generation

and validation of novel, specific monoclonal antibodies for the detection of Seoul virus proteins

in vitro. These reagents can be used to fill in critical gaps in knowledge regarding hantavirus

entry, protein expression, and particle generation.

<u>Introduction</u>

Hantaviruses (genus *Orthohantavirus*, order *Elliovirales*, class *Bunyaviricetes*) are zoonotic

segmented, negative-sense RNA viruses that can cause severe hemorrhagic disease in

humans. Old World hantaviruses (OW) are distributed throughout Southeast Asia, Russia, and

Europe and cause a form of hantavirus disease termed Hemorrhagic Fever with Renal

Syndrome (HFRS). New World hantaviruses (NW) are found in rodent reservoir populations in

the Americas and are responsible for Hantavirus Cardio-Pulmonary Syndrome (HCPS).

Infections with OW hantaviruses are more common with high morbidity but lower case-fatality

rates (1-15%) compared to NW hantavirus infections (30-45%). Despite the public health

concern, there are no FDA approved vaccines or targeted therapeutics for hantavirus infections.

The type species, *Orthohantavirus hantanense* (Hantaan virus, HTNV), was first isolated in

1978, and NW hantaviruses were discovered in 1993 (1-4). Yet much of the basic biology of this

family of viruses remains underexplored. This can be explained, in part, due to a scarcity of

tractable tools and models to study hantavirus infections *in vitro* and *in vivo*, especially

compared to other segmented viruses, such as influenza. We report here the generation of

monoclonal antibodies targeting the OW *Orthohantavirus seoulense* (Seoul virus, SEOV)

nucleoprotein, glycoprotein complex, and polymerase and demonstrate their use for Western

blotting and immunostaining assays. We tested these antibodies for specificity against SEOV

and other NW and OW hantaviruses, finding cross-reactivity only in a subset of anti-glycoprotein

antibodies. Several of the antibodies generated can be used to detect denatured proteins in cell

lysates as well as native proteins in immunoassays. These tools will be made available for the

research community through public repositories and have widespread applicability to test

antiviral efficacy, study pathogenesis in culture models, and to explore virus-host interactions for

this understudied virus family.

Results

Antibody development

Through Genscript, we purchased custom-generated hybridomas from mice immunized against

the four SEOV proteins: nucleoprotein (N), N-terminal glycoprotein (G_n), C-terminal glycoprotein

(G_c), and polymerase (L). Mice were immunized with linear peptides from Gn

(WRKKANQESANQNSC), Gc (CGDPGDVMGPKDKPF), or L (LEKKVIPDHPSGKTC). For N,

the entire recombinant protein was synthesized as the target antigen for immunization. From the

immunization scheme, we received five hybridomas per protein target, selected for antigen

reactivity by peptide ELISA. Supernatants from the hybridomas were purified on protein A/G

columns, concentrated, and tested for reactivity and specificity by Western blot and

immunostaining assays, as described below.

Nucleoprotein

The hantavirus nucleoprotein coats the viral RNA, plays a critical role in viral transcription and

translation, and is abundantly expressed in infected cells. We tested antibodies from each of the

five hybridomas generated against the SEOV nucleoprotein for reactivity by Western blot (WB),

immunofluorescence (IFA), and focus-forming unit assay (FFU). To test for specificity against

SEOV N, we transfected HEK293T cells with plasmids expressing the N protein from the OW

hantaviruses SEOV and HTNV, as well as the NW hantaviruses *Orthohantavirus*

sinnombreense (Sin Nombre, SNV) and *Orthohantavirus andesense* (Andes virus, ANDV).

Although expression for HTNV N was low, all ectopically-expressed proteins were detected in

the lysates by their HA tag but only SEOV N was detected by the monoclonal antibodies (Fig

1a). Next, we tested the specificity of these antibodies in immunofluorescence assays (IFA).

HEK293T cells were transfected with the N expression plasmids used above for SEOV, HTNV,

SNV, and ANDV. Two days post-transfection, the cells were fixed, permeabilized, and probed

with the monoclonal antibodies followed by an anti-mouse IgG fluorescent secondary. We

observed that all transfected N proteins were detectable with the anti-HA primary antibody, but

only SEOV N was detected with the anti-N monoclonal antibodies (Fig 1b, c). Fig 1c shows the

absence of N staining for HTNV, SNV, and ANDV N using a representative anti-N, clone 17E11.

Importantly, very low background staining was detected in mock transfected cells, supporting

the utility of these antibodies for IFA (Supp Fig 1a).

We then asked whether these antibodies could detect N during infection. We infected Vero E6

cells with SEOV, HTNV, SNV, or ANDV for 12 days and performed WB on collected cell lysates

(Fig 2a). Our standard hantavirus propagation protocols call for 12-day growth in Vero E6 cells

as the peak of viral production. Therefore, this time point was chosen to be reasonably assured

of abundant viral protein in cell lysates for all hantaviruses tested. Again, we see that the SEOV

N antibodies are highly specific for SEOV N and can be used to assess SEOV infection status in

susceptible cells. Similar results were observed when probing hantavirus-infected Vero E6 cells

for IFA (Fig 2b). Again, we noted very low background staining for these antibodies on mock-

infected cells (Supp Fig 1b). Finally, we tested the five anti-N antibodies for their efficacy in FFU

assays. Hantaviruses are typically non-cytopathic in healthy cells and therefore do not form

plaques. Instead, virus titrations can be executed by immunostaining. Vero E6 cells were

infected with serial dilutions of SEOV for 1 hour, after which a methylcellulose overlay was

added, and cells were incubated for 7 days post-infection. Cells stained with each of the anti-N

antibodies showed well-defined foci of infection with very low background staining of uninfected

monolayers (Fig 2c). These monoclonal antibodies, therefore, are specific for the SEOV

nucleoprotein and are effective in denaturing WB conditions and IF assays.

Glycoproteins

The hantavirus M segment encodes a precursor glycoprotein that is co-translationally cleaved at

a conserved WAASA site into two mature glycoproteins (G_N and G_C) (5). These glycoproteins

form trimeric complexes in the ER consisting of a G_N dimer and a G_C monomer (known as

glycoprotein complex, GPC), and are trafficked to the Golgi (6-11). Co-expression of both

proteins is necessary for trafficking from the ER to the Golgi, as individual expression results in

retention of the proteins in the ER (12-17). We immunized mice with peptides from either G_N or

G_C, but found that none of the antibodies raised against G_C-immunization were able to detect

glycoproteins from any of the viruses tested either in transfected or virus-infected cells (data not

shown). For these experiments, we transfected HEK293T cells with plasmids expressing the M

segment sequence for both G_N and G_C (GPC) for SEOV, HTNV, and ANDV. These plasmids

encode for the M segment coding region with an HA tag at the C-terminus. Thus, upon

expression and cleavage of the polyprotein, only the 52-58kDa G_C protein will be detectable with

an anti-HA antibody. SNV GPC was omitted because this plasmid failed to express detectable

HA-tagged G_C. Curiously, we found that several anti-G_N antibodies detected a protein migrating

at the same size as our viral G_N (~72-74kDa) in transfected HEK293T cells (Fig 3a). This cross-

reactive protein was only detected in HEK293T and not in other commonly used cells for

hantavirus research (A549, Vero E6, HUVEC-C, and primary rat lung microvascular endothelial

cells) (Fig 3b). Surprisingly, we found that three of our tested anti-G_N antibodies (1B12, 11A5,

and 12D10) detected the G_N from HTNV, SNV, and ANDV-infected Vero E6 cell lysates by WB

(Fig 3c). Unfortunately, none were able to recognize native GPC by IFA (data not shown).

Polymerase

The RNA-dependent RNA polymerase protein of hantaviruses (L) is expressed at low, almost

undetectable levels in cultured cells due to its presumed toxicity (18, 19). Even overexpression

of plasmid derived L protein has proven difficult without mutations that inhibit its transcriptional

activity (18). We generated HA-tagged plasmids for ectopic expression of SEOV L and SNV L in

HEK293T cells to test the anti-L antibodies in WB and IF assays (Fig 4a, b). Two of the four

antibodies tested were able to detect SEOV L by both assays using transfected HEK293T cells,

but none were cross-reactive against SNV L. We also fund that clone 13G8 was able to detect

SEOV L from infected Vero E6 cell lysates, but not from cells infected with HTNV, SNV, or

ANDV (Fig 4c). Despite considerable effort, we were unable to detect L expression in SEOV-

infected cells by IFA (data not shown).

Conclusions:

We have characterized novel monoclonal antibodies for their specificity and versatility for

detection of hantavirus proteins in exogenous and endogenously expressed conditions. These

tools for SEOV detection and quantification should replace less specific and less reliable

commercial sources for hantavirus antibodies. We are actively working to make these reagents

available to the hantavirus research community though trusted biorepositories, such as BEI

Resources.

Methods:

Cell culture

Vero E6 cells (ATCC, CRL-1586), A549 (ATCC, CCL-185), and HEK293T (ATCC, CRL-3216)

cells were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with 10%

heat-inactivated FBS, 1% pen/strep, 1% nonessential amino acids, 2.5% HEPES. Primary rat

microvascular endothelial cells (RLMVEC, VEC Technologies) were cultured in MCDB-131 base

medium (Corning) supplemented with EGMTM-2 Endothelial SingleQuots Bullet Kit (Lonza, CC-

4176) and 10% heat-inactivated FBS. Human umbilical endothelial cells (HUVEC-C; ATCC,

CRL-1730) were cultured in Lonza EGM-Plus (Lonza, CC-4542) supplemented with bullet kit

(Lonza, CC-5036) and 10% heat-inactivated FBS. All cells were cultured in tissue culture-

treated plastics and plate for endothelial cells were coated with rat-tail collagen (VWR, 47747-

218).

<u>Viruses and *in vitro* infections</u>

Seoul virus (strain SR11), Hantaan virus (strain 76-118) Sin Nombre virus (strain NMR11), and

Andes virus (CHI7913) were propagated on Vero E6 cells (ATCC, CRL-1586) for 12 days, with

a maximum of 3 passages. Infectious virus was isolated by harvesting supernatant and

centrifuging at 1000 rpm for 10 minutes to remove cellular debris. For virus infections, cells were

seeded in cell culture vessels 18-24 hours prior to infection at a target density of 70%. Virus

stock was diluted to the target MOI using serum-free Dulbecco's modified Eagle's medium

(DMEM) supplemented with 1x pen/strep, 1% nonessential amino acids, and 2.5% HEPES.

Virus was allowed to adsorb for 1 hour at 37ºC. Cells were washed twice with sterile PBS

solution (FisherScientific, SH30264FS), and appropriate culture medium was added for the

duration of the experiment.

<u>Antibody preparation</u>

Hybridomas were cultured in Dulbecco's modified Eagle's medium (DMEM) supplemented with

10% heat-inactivated FBS, 1% pen/strep, 1% nonessential amino acids, 1% L-Glutamine, 2.5%

HEPES, and 1x hybridoma fusion and cloning supplement (HFCS; MilliporeSigma

11363735001) for the first passage. Hybridomas were then maintained in in Dulbecco's

modified Eagle's medium (DMEM) supplemented with 10% heat-inactivated FBS, 1% pen/strep,

1% nonessential amino acids, 1% L-Glutamine, and 2.5% HEPES. Supernatants were collected

every 2-3 days when cells were at 90% confluency. They were then centrifuged at 1,000xg for

10min to remove cell debris and stored at 4C until purified. To purify antibody, a 2mL Pierce

protein A/G agarose column (Thermo Scientific, 20422, 29922) was prepared according to

manufacturer instructions. Supernatants from the same hybridoma culture were pooled together

and 1/3 volume of Pierce protein A/G IgG binding buffer (Thermo Scientific, 54200) was added

to the supernatants. Supernatant and binding buffer were run through the column according to

manufacturer instructions. Antibody was eluted from the column using 8mL of Pierce IgG elution

buffer (Thermo Scientific, 21004) and neutralized with 400uL 1M Tris pH 7.5. Amicon Ultra-15

centrifugal filters, 30kDa MW (MilliporeSigma, UFC903024), were used to concentrate the

purified antibody down to 200uL. Antibody concentration was determined via Pierce BCA

Protein Assay Kit (ThermoFisher Scientific, 23225) and stocks stored at 1ug/mL.

<u>Plasmids</u>

To validate the detection of hantavirus proteins by each monoclonal antibody, we subcloned

each of the described proteins from SEOV, HTNV, SNV, and ANDV into our previously

described pCAGGS expression vector (20, 21). This vector contains the coding sequence of

each viral protein followed by a C-terminal hemagglutinin (HA) epitope tag (YPYDVPDYA). Viral

protein coding sequences were PCR amplified from viral cDNA and subcloned into the vector

using Gateway Technology (Invitrogen) following manufacturer's instructions. The nucleotide

sequence of each plasmid was verified by DNA sequencing (Plasmidsaurus). Viral protein

sequences were subcloned from the following virus strains: SEOV strain SR-11, HTNV strain

76-118, SNV strain NMR11, and ANDV strain CHI-7913. For the SEOV L protein plasmid, the

previously reported K44A mutation was made to increase expression of L (18).

<u>Plasmid transfections</u>

HEK 293T cells were seeded in 10cm dishes at 2×10^5 cells/mL. Transfections were

accomplished 24hrs later using jetPRIME (VWR, 89129-924) according to manufacturer's

specifications. Briefly, 5ug (GPC and L) or 10ug (N) of the indicated expression plasmid were

diluted in 500uL jetPRIME buffer and 10uL of jetPRIME reagent was added. The mixture was

incubated at room temperature for 10min and was then added dropwise to the 10cm dishes of

cells. Lysates were harvested 48hrs post-transfection in RIPA buffer (50mM Tris HCl pH 7.4,

150mM NaCl, 1% (v/v) NP-40, 0.5% (w/v) Na-deoxycholate) and clarified through 25,000rcf

centrifugation for 15 minutes at 4°C. Protein was quantified via Pierce BCA Protein Assay Kit

(ThermoFisher Scientific, 23225). 30ug of protein was loaded per sample into an 8% (L) or 10%

(N and GPC) polyacrylamide gel and, after denaturing electrophoresis, transferred to a 0.45μm

nitrocellulose membrane (VWR, 10120-006). Membranes were blocked at room temperature in

10% FCS in PBS-T. Primary antibodies (1ug/mL) were incubated at 1:500 at 4° overnight. HRP-

conjugated secondary antibodies against primary antibody species were incubated 1:10,000 for

1 hour at room temperature. Blots were imaged on BioRad Chemidoc MP Imaging System

using chemiluminescence Pierce Substrate for Western Blotting (VWR, PI80196).

Generation of infected-cell lysates

Cell lysates to be interrogated for protein expression by immunoblot analysis were harvested in

protein lysis buffer 12 days post-infection and prepared as previously described (22). Briefly,

protein was harvested in RIPA buffer (50mM Tris HCl pH 7.4, 150mM NaCl, 1% (v/v) NP-40,

0.5% (w/v) Na-deoxycholate) and clarified through 25,000rcf centrifugation for 15 minutes at

4°C. Protein was quantified via Pierce BCA Protein Assay Kit (ThermoFisher Scientific, 23225).

30ug of protein was loaded per sample into an 8% (L) or 10% (N and GPC) polyacrylamide gel

and, after denaturing electrophoresis, transferred to a 0.45μm nitrocellulose membrane (VWR,

10120-006). Membranes were blocked at room temperature in 10% FCS in PBS-T. Primary

antibodies (1ug/mL) were incubated at 1:500 at 4° overnight. HRP-conjugated secondary

antibodies against primary antibody species were incubated 1:10,000 for 1 hour at room

temperature. Blots were imaged on BioRad Chemidoc MP Imaging System using

chemiluminescence Pierce Substrate for Western Blotting (VWR, PI80196).

Immunofluorescence assay

Vero E6 cells were seeded in 48 well plates at 2.5x10^3 cells/well. When cells had reached 70-

80% confluency, they were infected with either SEOV, HTNV, ANDV, or SNV at an MOI of 0.1.

Virus was allowed to adsorb for 1hr at 37° C, then cells were washed 2x with 1x PBS and

complete DMEM was added. At 4 days post-infection, supernatants were removed and cells

were washed 1x with 1x PBS. Cells were fixed with 95% ethanol:5% acetic acid for 10min at -

20° C. Cells were permeabilized with 0.01% Triton X-100 in PBS for 15min at room temperature

and then blocked with 3% FBS in PBS for at least 30min. Primary antibodies (1ug/mL) were

incubated at 1:500 at 4°C overnight. Goat α mouse 647 fluorescent secondary antibodies

(Invitrogen A21235) were incubated at 1:400 for at least 2hrs at room temperature. Nuclei were

stained with DAPI at 1:1000 for 5min at room temperature. Stained cells were visualized using

the 20x objective on the EVOS Cell Imaging System (ThermoScientific). Microscope settings for

fluorescence intensity were set on mock-transfected/infected cells to remove background and

remained constant for all imaging of transfected/infected cells.

Focus Forming Unit Assay

Infectious virus was quantified using immunostaining as previously described (23). Vero E6 cells

were infected with 100μL of culture supernatants and incubated for 2 hours at 37ºC. After

incubation, a 2% carboxymethylcellulose overlay containing supplemented DMEM (2%

pen/strep, 2% nonessential amino acids, 2% HEPES, 4% heat-inactivated FBS). Cells were

incubated for 7 days at 37ºC. After incubation, cells were fixed with 95% EtOH:5% Acetic Acid

for 10 minutes at -20ºC, blocked with 3% FBS in PBS, and probed for SEOV N (α SEOV

nucleocapsid, 17E11 or others presented here) or HTNV N (anti-HTNV nucleocapsid 76–118,

BEI resources NR-12152, cross-reacts with HTNV, SNV, and ANDV) at 4º for 24 hours. HRP

conjugated secondary (donkey α rabbit for α HTNV, Jackson Immunoresearch 711-035-152; or

goat α mouse for α SEOV, Jackson Immunoresearch 115-035-003) for 2 hours at room

temperature. Foci were then stained using Vector VIP Substrate Kit (Vector Laboratories, SK-

4600) and counted under a light microscope to calculate titer.

Acknowledgments:

Work performed by KM and CS was supported by the National Institutes of Health NIAID grant

1R21AI175721-01A1 granted to AMK and JB. ATL was supported by a post-doctoral fellowship

by the National Institutes of Health NIGMS T32AI007538. The funders had no role in study

design, data collection and analysis, decision to publish, or preparation of the manuscript.

The authors also thank Dr. Rebekah Honce for technical support in plasmid preparation and

expression.

References:

1.   Nichol ST, Spiropoulou CF, Morzunov S, Rollin PE, Ksiazek TG, Feldmann H, et al. Genetic
identification of a hantavirus associated with an outbreak of acute respiratory illness. Science.
1993;262(5135):914-7. doi: 10.1126/science.8235615. PubMed PMID: 8235615.
2.   Marshall E. Hantavirus outbreak yields to PCR. Science. 1993;262(5135):832, 4-6. doi:
10.1126/science.8235602. PubMed PMID: 8235602.
3.   Lee HW, Lee PW, Baek LJ, Song CK, Seong IW. Intraspecific transmission of Hantaan
virus, etiologic agent of Korean hemorrhagic fever, in the rodent Apodemus agrarius. Am J Trop
Med Hyg. 1981;30(5):1106-12. doi: 10.4269/ajtmh.1981.30.1106. PubMed PMID: 6116436.
4.   Lee HW, Lee PW, Johnson KM. Isolation of the etiologic agent of Korean Hemorrhagic
fever. J Infect Dis. 1978;137(3):298-308. doi: 10.1093/infdis/137.3.298. PubMed PMID: 24670.
5.   Lober C, Anheier B, Lindow S, Klenk HD, Feldmann H. The Hantaan virus glycoprotein
precursor is cleaved at the conserved pentapeptide WAASA. Virology. 2001;289(2):224-9. Epub
2001/11/02. doi: 10.1006/viro.2001.1171. PubMed PMID: 11689045.
6.   Petazzi RA, Koikkarah AA, Tischler ND, Chiantia S. Detection of Envelope Glycoprotein
Assembly from Old-World Hantaviruses in the Golgi Apparatus of Living Cells. J Virol.
2021;95(4). Epub 2020/11/27. doi: 10.1128/JVI.01238-20. PubMed PMID: 33239451; PubMed
Central PMCID: PMC7851546.
7.   Cifuentes-Munoz N, Salazar-Quiroz N, Tischler ND. Hantavirus Gn and Gc envelope
glycoproteins: key structural units for virus cell entry and virus assembly. Viruses.
2014;6(4):1801-22. Epub 2014/04/24. doi: 10.3390/v6041801. PubMed PMID: 24755564;
PubMed Central PMCID: PMC4014721.

8.   Schmaljohn CS, Schmaljohn AL, Dalrymple JM. Hantaan virus M RNA: coding strategy,
nucleotide sequence, and gene order. Virology. 1987;157(1):31-9. Epub 1987/03/01. doi:
10.1016/0042-6822(87)90310-2. PubMed PMID: 3103329.
9.   Antic D, Wright KE, Kang CY. Maturation of Hantaan virus glycoproteins G1 and G2.
Virology. 1992;189(1):324-8. Epub 1992/07/01. doi: 10.1016/0042-6822(92)90709-x. PubMed
PMID: 1604818.
10.   Shi X, Elliott RM. Analysis of N-linked glycosylation of hantaan virus glycoproteins and
the role of oligosaccharide side chains in protein folding and intracellular trafficking. J Virol.
2004;78(10):5414-22. Epub 2004/04/29. doi: 10.1128/jvi.78.10.5414-5422.2004. PubMed
PMID: 15113920; PubMed Central PMCID: PMC400336.
11.   Serris A, Stass R, Bignon EA, Muena NA, Manuguerra JC, Jangra RK, et al. The Hantavirus
Surface Glycoprotein Lattice and Its Fusion Control Mechanism. Cell. 2020;183(2):442-56 e16.
Epub 2020/09/17. doi: 10.1016/j.cell.2020.08.023. PubMed PMID: 32937107; PubMed Central
PMCID: PMC7572791.
12.   Ruusala A, Persson R, Schmaljohn CS, Pettersson RF. Coexpression of the membrane
glycoproteins G1 and G2 of Hantaan virus is required for targeting to the Golgi complex.
Virology. 1992;186(1):53-64. Epub 1992/01/01. doi: 10.1016/0042-6822(92)90060-3. PubMed
PMID: 1727611.
13.   Pensiero MN, Hay J. The Hantaan virus M-segment glycoproteins G1 and G2 can be
expressed independently. J Virol. 1992;66(4):1907-14. Epub 1992/04/01. doi:
10.1128/JVI.66.4.1907-1914.1992. PubMed PMID: 1548747; PubMed Central PMCID:
PMC288978.
14.   Shi X, Elliott RM. Golgi localization of Hantaan virus glycoproteins requires coexpression
of G1 and G2. Virology. 2002;300(1):31-8. Epub 2002/08/31. doi: 10.1006/viro.2002.1414.
PubMed PMID: 12202203.
15.   Spiropoulou CF, Goldsmith CS, Shoemaker TR, Peters CJ, Compans RW. Sin Nombre virus
glycoprotein trafficking. Virology. 2003;308(1):48-63. Epub 2003/04/23. doi: 10.1016/s0042-
6822(02)00092-2. PubMed PMID: 12706089.
16.   Deyde VM, Rizvanov AA, Chase J, Otteson EW, St Jeor SC. Interactions and trafficking of
Andes and Sin Nombre Hantavirus glycoproteins G1 and G2. Virology. 2005;331(2):307-15. Epub
2005/01/05. doi: 10.1016/j.virol.2004.11.003. PubMed PMID: 15629773.
17.   Sperber HS, Welke RW, Petazzi RA, Bergmann R, Schade M, Shai Y, et al. Self-association
and subcellular localization of Puumala hantavirus envelope proteins. Sci Rep. 2019;9(1):707.
Epub 2019/01/27. doi: 10.1038/s41598-018-36879-y. PubMed PMID: 30679542; PubMed
Central PMCID: PMC6345964.
18.   Heinemann P, Schmidt-Chanasit J, Günther S. The N terminus of Andes virus L protein
suppresses mRNA and protein expression in mammalian cells. J Virol. 2013;87(12):6975-85.
Epub 2013/04/10. doi: 10.1128/JVI.00043-13. PubMed PMID: 23576516; PubMed Central
PMCID: PMC3676134.
19.   Meier K, Thorkelsson SR, Durieux Trouilleton Q, Vogel D, Yu D, Kosinski J, et al. Structural
and functional characterization of the Sin Nombre virus L protein. PLoS Pathog.
2023;19(8):e1011533. Epub 20230807. doi: 10.1371/journal.ppat.1011533. PubMed PMID:
37549153; PubMed Central PMCID: PMC10406178.

20.   Cornillez-Ty CT, Liao L, Yates JR, Kuhn P, Buchmeier MJ. Severe acute respiratory
syndrome coronavirus nonstructural protein 2 interacts with a host protein complex involved in
mitochondrial biogenesis and intracellular signaling. J Virol. 2009;83(19):10314-8. Epub
20090729. doi: 10.1128/JVI.00842-09. PubMed PMID: 19640993; PubMed Central PMCID:
PMC2748024.
21.   King BR, Hershkowitz D, Eisenhauer PL, Weir ME, Ziegler CM, Russo J, et al. A Map of the
Arenavirus Nucleoprotein-Host Protein Interactome Reveals that Junín Virus Selectively Impairs
the Antiviral Activity of Double-Stranded RNA-Activated Protein Kinase (PKR). J Virol.
2017;91(15). Epub 2017/07/12. doi: 10.1128/JVI.00763-17. PubMed PMID: 28539447; PubMed
Central PMCID: PMC5512243.
22.   Klimaj S, LaPointe A, Martinez K, Acosta EH, Kell AM. Seoul orthohantavirus evades
innate immune activation by reservoir endothelial cells. bioRxiv. 2024:2024.06.18.599607. doi:
10.1101/2024.06.18.599607.
23.   Kraus AA, Priemer C, Heider H, Kruger DH, Ulrich R. Inactivation of Hantaan virus-
containing samples for subsequent investigations outside biosafety level 3 facilities.
Intervirology. 2005;48(4):255-61. Epub 2005/05/28. doi: 10.1159/000084603. PubMed PMID:
15920350.

Figure legends:

Figure 1. Antibodies specifically recognize SEOV N in transfected cells. HEK293T cells

expressing HA-tagged N protein from SEOV, HTNV, SNV, and ANDV were interrogated by

Western blotting and IFA. A) 30ug of cell lysates were loaded into 10% polyacrylamide gels in

denaturing conditions and probed for N expression using anti-HA and newly generated anti-

SEOV N antibodies. Actin served as a loading control. B) HEK293T cells transfected with SEOV

N plasmid, probed using DAPI (nuclei, blue) and anti-HA or antibodies against SEOV N (red). C)

HEK293T cells transfected with N plasmids from the indicated virus and probed with anti-HA

antibodies or SEOV N antibody 17E11 (red) and DAPI (nuclei, blue). Cells visualized on the

EVOS Cell Imaging System.

Figure 2. Antibodies specifically recognize SEOV N in hantavirus infected cells. Vero E6

cells were infected with SEOV, HTNV, ANDV or SNV and interrogated by Western blotting (A),

IFA (B), and FFU assay (C). A) 30ug cell lysates were loaded into 10% polyacrylamide gels in

denaturing conditions and probed for N expression using antibody clones against N. Actin

serves as loading control. B) SEOV-infected Vero E6 cells probed with anti-N antibodies (red)

and DAPI (nuclei, blue). Cells visualized on the EVOS Cell Imaging System. C) Focus forming

unit assay using Vero E6 cells infected with SEOV and incubated with a 2% methylcellulose

overlay for 7 days. Cells were probed with anti-N antibody clones, HRP-conjugated secondary

antibody, and visualized for foci of stained cells, representing viral infectious units. Mock-

infected cells represent non-specific background antibody interactions.

Figure 3. Several anti-G_N antibodies recognize G_N from OW and NW hantaviruses. A)

HEK293T cells were transfected with plasmids expressing the indicated hantavirus GPCs from

SEOV, HTNV, or ANDV. Each GPC encodes a C-terminal HA tag which, upon cleavage allows

for the detection of expressed G_C only (52-58 kDa). Cell lysates were interrogated by Western

blotting for recognition by anti-G_N antibodies or the anti-HA antibody. B) Un-transfected cell

lysates from several cell types commonly used in hantavirus research were interrogated by

Western blotting for cross-reactivity with anti-G_N antibodies. C) Vero E6 cells infected with

SEOV, HTNV, ANDV, SNV, or mock infected were analyzed by Western blot for detection of G_N

by anti-G_N monoclonal antibodies. Actin served as a loading control.

Figure 4. Monoclonal antibodies specifically recognize SEOV L protein. A) HEK293T cells

transfected with plasmids expressing HA-tagged SEOV L or SNV L were analyzed by Western

blotting using anti-L monoclonal antibodies. B) SEOV L plasmid transfected HEK293T cells

were probed with anti-HA antibody or clones 13G8 or 17F3 (red) and DAPI (nuclei, blue). Cells

visualized on the EVOS Cell Imaging System. C) Vero E6 cells infected with SEOV, HTNV,

ANDV, SNV, or mock-infected were analyzed by Western blot for detection by anti-L

monoclonal antibodies. Actin served as a loading control.

Supplementary Figure 1. Anti-N antibodies exhibit low background staining in mock

transfected or mock infected cells. Mock-transfected HEK293T cells (A) and mock-infected

Vero E6 cells (B) probed with anti-HA and anti-N monoclonal antibodies (red) and DAPI (nuclei,

blue). Cells visualized on the EVOS Cell Imaging System.

Supplementary Figure 2. Anti-L antibodies exhibit low background staining in mock

transfected cells. Mock-transfected HEK293T cells probed with anti-HA and anti-L monoclonal

antibodies (red) and DAPI (nuclei, blue). Cells visualized on the EVOS Cell Imaging System.