Title: Polysialic acid is a versatile marker for retinal Müller glia in common vertebrate model organisms and systems.

Running Title: PolySia labels vertebrate Müller glia

Authors: A. Nicol San Juan[1], Allison D. N. N. Tran[1], Debarati Hore[1], Linnea M. Kriese[1], Natalie H. N. Van Engelen[1], Maria Sharkova[2], Nicole C. L. Noel[3], Carmanah Hunter[4], Olga Spencer Carreira[5], Jennifer C. Hocking[2,6], Lisa M. Willis[4], Brittany J. Carr[1,2]

[1] Department of Ophthalmology and Visual Sciences, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, T6G 2E1, Canada

[2] Department of Cell Biology, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, T6G 2R8, Canada

[3] Institute of Ophthalmology, University College London, London, EC1V 9EL, United Kingdom

[4] Department of Biological Sciences, Faculty of Science, University of Alberta, Edmonton, T6G 2R3, Canada

[5] iNOVA4Health, NOVA Medical School, Universidade Nova de Lisboa, Lisbon, Portugal

[6] Division of Anatomy, Department of Surgery, Faculty of Medicine and Dentistry, University of Alberta, Edmonton, T6G 2B7, Canada

A. Nicol San Juan (aldeanic@ualberta.ca)

Allison D. N. N. Tran (adtran1@ualberta.ca)

Linnea M. Kriese (lkriese@ualberta.ca)

Natalie H. N. Van Engelen (nvanenge@ualberta.ca)

Debarati Hore (hore1@ualberta.ca)

Maria Sharkova (sharkova@ualberta.ca)

Nicole C. L. Noel (n.noel@ucl.ac.uk)

Carmanah Hunter (carmanah@ualberta.ca)

Olga Spencer Carreira (olga.carreira@nms.unl.pt)

Jennifer C. Hocking (jhocking@ualberta.ca)

Lisa M. Willis (lisa.willis@ualberta.ca)

Corresponding Author: *Brittany J. Carr (bcarr1@ualberta.ca)

Conflict of Interest Disclosure

The authors declare no conflict of interest.

Keywords

polysialic acid (polySia), GFP-Endoneuraminidase double mutant (GFP-EndoN$_{DM}$), Müller glia,

retinal development, vertebrate retina, mAb735, Neu5Ac

Highlights

- PolySia is a highly conserved Müller glia marker in vertebrate retina and retinal

organoids

- Labeling Müller glia with markers of polySia, facilitates the analysis of fine Müller glia

processes in the IPL sublaminae, whole Müller glia morphology, and degenerative

gliosis

- PolySia can be labeled with an engineered lectin conjugate (GFP-EndoN$_{DM}$) and a

commercially-available monoclonal antibody

Abstract (250/250 words)

Müller glia are retinal support cells that play crucial roles in tissue structure, waste management, and repair. A challenge for the field has been to find Müller glia markers that detect non-reactive cells or work well in non-mammalian models. We introduce two novel markers for identifying reactive and non-reactive Müller glia in vertebrate retinas: a modified enzyme lectin (GFP-EndoN$_{DM}$) and a monoclonal antibody (mAb735). These markers recognize polysialic acid (polySia), which is a highly conserved glycosylation modification in humans and vertebrates. In the retina, polySia is present on Müller glia predominantly in the form of polySia-NCAM. We used GFP-EndoN$_{DM}$ and mAb735 to investigate polySia distribution in Müller glia of fish, amphibians, reptiles, birds, rodents, retinal organoids, and humans. In adult retinas of most species, polySia was localized to the Müller glia and spanned outer to inner retina, with inner plexiform layer (IPL) sublaminae ramifications. Gliosis was also detectable in degenerating murine retinas. Notable species differences were that only outer retinal regions of Müller glia were labelled in adult zebrafish, whereas the outer Müller glia body up to the first IPL sublamina was labelled in adult turquoise killifish. There was no significant retinal polySia labeling in larval zebrafish, but it was present in the brain. Larval turquoise killifish have polySia throughout the retina, similar to other adult vertebrates. Labelling polySia expands the scientific toolbox for Müller glia markers, and offers a versatile way to visualize and monitor structural changes in non-reactive and reactive Müller glia across most vertebrate species.

1. Introduction

1.1 Müller glia

Müller glia are the principal macroglial cells in the vertebrate retina. They are homologous to the radial glia of the brain and spinal cord, and are integral supporters of retinal architecture, metabolism, and homeostasis. Müller glia have many functions including water and ion regulation[1], retinoid metabolism and visual pigment recycling[2], and energy production[3]. In addition, Müller glia also participate in phagocytosis and retinal waste removal, both during development[4] and in response to retinal degenerative disease[5]. Although they are one of the last cell types to differentiate during retinal development[6], Müller glia are essential in guiding retinal organization and maintaining retinal laminations by supporting retinal plasticity, neuronal pruning, and retinal information processing[1,4,7,8]. Selective destruction of the Müller glia results in retinal dysplasia, photoreceptor death, retinal vascular abnormalities, and retinal pigment epithelium (RPE) dysfunction[9–11]. From apical to basal retina, Müller glia occupy nearly the entire neural retinal space, and interact with or comprise significant structural components of the retina, including the outer segment layer (OS), the external limiting membrane (ELM), the outer nuclear layer (ONL), the outer plexiform layer (OPL), the inner nuclear layer (INL), the inner plexiform layer (IPL), the ganglion cell layer (GCL), the nerve fibre layer (NFL), and the internal limiting membrane (ILM). The Müller glia apical processes, which extend beyond the ELM into the OS space, may interact with RPE microvilli to aid in cone stability or chromophore metabolism and recycling[12]. Müller glia interact with outer and inner retinal neurons in both plexiform layers, and contribute to retinal homeostasis by supporting the neurons with trophic molecules, removing waste, ion regulation, neurotransmitter release, and regulation of ATP[13]. Müller glia contribute to retinal integrity not only by forming connections with retinal neurons[14], but also by formation of the ELM and the ILM[15]. The ELM and ILM are not true membranes, but instead comprise tight and adherent junctions between adjacent Müller glia and the

photoreceptors, which form a type of semi-permeable membrane that can regulate molecule

transport and add structural support to the retina[16,17]. The ELM is formed as a junction with the

photoreceptor inner segments, whereas the ILM is formed by the Müller glia endfeet at the

vitreoretinal interface. Finally, in some species like fish and amphibians, Müller glia support

retinal regeneration and repair by de-differentiating into retinal progenitors, acting as stem cells

to re-populate retinal neurons lost to injury[18–20].

In addition to important roles in retinal homeostasis, Müller glia become reactive in response to

retinal degenerative pathology by undergoing morphological, physiological, and biochemical

changes[21,22]. These gliotic alterations are intended to protect the retina, by releasing neurotropic

factors, scavenging free radicals, modulating glutamate uptake, and facilitating the formation of

glial scars, which isolate diseased tissue from healthy tissues[21–23]. Sometimes, Müller glia can

instead be harmful, particularly by contributing to the development of edema and preventing

retinal repair and remodeling as a result of glial scarring[24–27]. Two well known indicators of

Müller glial reactivity and gliosis are the upregulation of the intermediate filament protein, glial

fibrillary acidic protein (GFAP) and changes in glutamine synthetase (GS), a Müller cell-specific

enzyme normally involved in neurotransmitter recycling[28–30]. Consequently, these two Müller glia

markers are the most popular for use in biomedical research. GFAP and GS are less-than-ideal

tools for revealing the morphology of the entire Müller glia cell, however, especially the fine

processes in the plexiform layers[31,32]. They are also not very effective markers for Müller glia

under homeostatic conditions, because they sometimes do not have significant expression of

GFAP or GS[33]. Thus, there is significant room for improvement in the development or

characterization of a versatile Müller glia marker for use across various stages of retinal health,

and in different species.

1.2 PolySia-NCAM

Glycosylation is the post-translational process of covalently attaching carbohydrates to a target protein or macromolecule, with virtually all cell surface proteins undergoing modification with glycans to form a glycocalyx around the cell membrane[34]. Protein glycosylation increases proteomic diversity, provides biochemical stability, and, in some cases, can confer additional functions to proteins. The glycocalyx on the outside of the cell, which can be up to 500 nm thick, is involved with cell-cell and cell-extracellular matrix communication, cell migration and adhesion, and pathogen-host interactions[35,36]. Sialic acids or neuraminic acids comprise a family of 9-carbon carboxylated sugars, named 3-deoxy-D-*glycero*-D-*galacto*-non-2-ulosonic acids. *N*-acetylneuraminic acid (Neu5Ac) and *N*-glycolylneuraminic acid (Neu5Gc) are the most common sialic acids in vertebrates and differ from one another only in the presence of an additional hydroxyl group in Neu5Gc. In mammals, the Neu5Gc modification is created via hydroxylation of Neu5Ac by the cytidine monophosphate-N-acetylneuraminic acid hydroxylase (CMAH) enzyme[37]. Certain species, including humans, deer, chickens, and brown bats, have lost the ability to make Neu5Gc due to dysfunctional or non-functional CMAH[38]. Regardless of whether a species can make Neu5Gc, the dominant sialic acid in the brain and retina is almost exclusively Neu5Ac[39]. These sialic acids frequently occupy the terminal position on various glycoproteins and glycolipids in mono-, oligo-, or polymeric forms[34]. Polysialic acid (polySia) is a is negatively charged linear homopolymer of 8 or more sialic acids which, in the vertebrate nervous system, are connected though α2,8-linkages. PolySia is reported to modify properties of cell-cell associations, including signaling, adhesion, and migration[40]. It also plays a significant role in cancer metastasis, and has been investigated in the context of cancer vaccines[41,42].

In the central nervous system, polySia is found bound primarily to the membrane protein neural cell adhesion molecule (NCAM)[43,44]. NCAM is found in neurons, glial cells, and Schwann cells[45–

[47], and can be post-translationally modified with polySia to form polySia-NCAM[48,49]. It is thought

that polySia-NCAM reduces the strength of the adhesion forces between cells by steric

hindrance, making it harder for cells come into contact and decreasing cellular adhesion[48,50].

This decreased adhesion property of polySia-NCAM has led to its implication in the regulation of

cell migration, synapse formation, and neuronal plasticity[51]. Correspondingly, polySia

expression in the brain is high during embryonic development[44,52,53], but it also remains present

at lower expression levels in adults, particularly in areas important for neuro-plasticity or neuro-

progenitor populations[54–58]. Retinal expression of polySia-NCAM is less well studied. NCAM

alone has been reported to be expressed in adult rodent Müller glia and putative optic nerve

astrocytes, and the inner and outer plexiform layers in bovine and frog retina[52,59–61]. Retinal

polySia expression has been investigated previously in the rat, salamander and zebrafish[60,62,63].

Interestingly, only the rat paper identified the polySia to be localized to the Müller glia. In

salamander and zebrafish, polySia expression was reported to be localized primarily to the

ONL, near the cone photoreceptor cell bodies. In the salamander, polySia expression was also

reported within the INL, IPL, GCL, and NFL[62].

The high conservation of polySia structure and its localization to the cell membrane makes it an

attractive target for labeling Müller glia, enabling the study of glial morphology and behavior in

the vertebrate retina. We found that polySia is an effective and versatile marker for Müller glia

in a wide variety of vertebrate species, from amphibians to humans, as well as in retinal

organoids. We also found notable differences in polySia localization on the Müller glia of fishes

compared to all other vertebrates. Contrary to reports of significant differences in polySia

expression during development in the brain, there is a less apparent change in polySia

expression in retinal Müller glia expression of polySia between embryonic and adult retinas. The

exception is again fishes; polySia is absent in larval zebrafish larval and less confined in larval

killifish compared to adult killifish.

2. Materials and Methods

2.1 Human and Animal Ethics Statements

Human post-mortem eye tissues were obtained through collaborators under approved protocols.

Retinal tissues were donated from 3 men, aged late-sixties to early eighties, with no known prior

history of ocular disease (Lions Eye Bank, University of Iowa). The use of these tissues for this

study was approved by the University of Alberta (Pro00147177), and carried out in accordance

with the Health Canada and Public Health Agency of Canada (PHAC) Research Ethics Board

(REB), guided by the principles of the second edition of the Tri-Council Policy Statement: Ethical

Conduct for Research Involving Humans (TCPS 2). This study was performed in accordance

with the ethical standards as laid down in the 1964 Declaration of Helsinki and its later

amendments.

Animal experiments were approved by the Canadian Council on Animal Care (CCAC) and

Animal Care and Use Committees (ACUC). We examined adult retinas of 8 different vertebrate

non-human species: African turquoise killifish (*Nothobranchius furzeri*), zebrafish (*Danio rerio*),

African clawed frog (*Xenopus laevis*), hummingbird (*Calypte anna*), fire skink (*Lepidothyris*

fernandi), zebra finch (*Taeniopygia guttata*), mouse (*Mus musculus*), and rat (*Rattus*

norvegicus). Adult frog eyes were obtained under the University of Alberta approved animal

ethics protocol (AUP00004203) and adult zebrafish eyes were obtained under the University of

Alberta approved ethics protocol (AUP00001476). Killifish experimental procedures were

conducted in accordance with the UK Home Office Animals (Scientific Procedures) Act of 1986

and killifish eyes were obtained under the animal licenses 70/7391 and PP2133797.

All other animal tissues were donated to the Carr lab from various researchers after the animals

were euthanized by appropriate personnel under appropriate protocols (mouse, rat) or obtained

under appropriate wildlife permits and approved protocols (fire skink, hummingbird, zebra finch).

We also investigated two cephalopod species. Octopus (*Octopus bimaculoides*) and squid

(*Euprymna berryi*) were acquired from the Cephalopod Resource Center (Woods Hole, MA,

USA) and kept at the University of Oregon, until fixed tissues were sent to the Carr lab.

Cephalopod studies were conducted with approved protocols from the University of Oregon

Animal Care Services, in compliance with the Association for Assessment and Accreditation of

Laboratory Animal Care International guidelines. Animal husbandry and protocols were carried

out in accordance with published guidelines for the care and welfare of cephalopods in the

laboratory[64]. Figures are representative images from an n = 3-4 for all species, except for the

fire skink (n = 1), zebra finch (n = 2), octopus (n = 1), and squid (n = 1).

Most tissues were from wildtype animals. Exceptions were 2/3 of the adult Sprague Dawley rats,

which were Thy1-GFP transgenics, the postnatal rats were culls from a connexion (Cx) 40

heterozygous breeding (25% $Cx^{+/+}$, 25% $Cx40^{-/-}$, 50% $Cx40^{+/-}$), and the postnatal mice, which

were floxed $R26^{lsl}$-Cmah mice[65]. Thy1 is a surface glycosylphosphatidylinositol-anchored protein

with proposed roles in the control of cellular differentiation, cell adhesion, cell signalling, and

immunity[66,67]. In the adult retina, THY1 protein is reported to be expressed in a subset of retinal

amacrine and ganglion cells[68–70], which we confirmed in this study. The presence of Thy1-GFP

did not obviously affect retinal polySia expression patterns compared to the wildtype rat

(**Supplementary Fig. S1**). Cx40 is associated with the development of retinal vasculature[71],

and we did not observe any significant differences in retinal vasculature or retinal labeling

patterns between the postnatal rats investigated in this study (n = 6). Dams of postnatal mice

were not fed tamoxifen during pregnancy, so the floxed R26[lsl]-Cmah mice should represent a

wildtype phenotype. We observed no differences in the labelling patterns between littermates (n

= 7).

2.2 Organoid Sources and Differentiation

Retinal organoids were obtained from collaborators under approved protocols (NOVA Medical

School, Universidade Nova de Lisboa). Two human inducible pluripotent stem cell (hiPSC) lines

were used in this study: a Parkinson's patient commercialized cell line (ND50040,

NHCDR/RUCDR) and a healthy control cell line (IMR90-4, WiCell). Cell lines were maintained

and expanded to 80% confluency, after which a retinal organoid differentiation protocol was

implemented as described previously[72]. Briefly, hiPSCs were lifted (Versene, Thermo Fisher

Scientific) and split at 3000 cells/well in a Nunclon Sphera 96-well U-bottom plate (Thermo

Fisher Scientific) using mTeSR™ Plus medium supplemented with 10 µM of ROCK inhibitor

(Focus Biomolecules). The cells were grown as aggregates for 6 days as an adaptation period

to Neural Induction Medium (NIM) containing DMEM/F12, 1% N2 supplement, 1□×□non-

essential amino acids, 1□×□GlutaMax (Gibco, Thermo Fisher Scientific), and 2 mg/ml Heparin

(Sigma-Aldrich). On day 6, 1.5 nM of bone morphogenic protein-4 (BMP4, Peprotech) was

added to fresh NIM, and on day 7, EBs were transferred to growth factor–reduced matrigel-

coated 6-well plates (Corning). The medium was replaced by half-fresh NIM on days 9, 12, and

15. On day 16, the medium was replaced by retinal differentiation medium (RDM) containing

DMEM:F12 (3:1), 2% B27 minus vitamin A, 1□×□non-essential amino acids, 1□×□GlutaMax,

1□×□antibiotic/antimycotic (Gibco, Thermo Fisher Scientific). This medium was changed every

2–3 days. At day 30, optic vesicles were manually dissected using a surgical scalpel (SM65A,

Swann-Morton Ltd) under the microscope EVOS XL core (Thermo Fisher Scientific). Dissected

organoids were maintained in suspension flasks (Sarstedt) in 3D-retinal differentiation medium

(3D-RDM) containing DMEM:F12 (3:1), 2% B27 minus vitamin A, 1□×□non-essential amino

acids, 1□×□GlutaMax, 1□×□antibiotic/antimycotic, 5% FBS, 1:1000 chemically defined lipid

supplement (Gibco, Thermo Fisher Scientific), 100 µM taurine (Sigma-Aldrich), and 1 µM of all-

trans retinoic acid (RA, Sigma-Aldrich) until day 120. Fixed and cryoprotected organoid tissues

(day 120) were shipped to us, and then we sectioned and labelled them according to the

protocol described below.

2.3 Retinal Tissue Preparation and Cryosectioning

Whole eyes prepared in the lab were removed and fixed in 4% paraformaldehyde (PFA) + 3%

sucrose in 0.1 M sodium phosphate buffer (PB, pH 7.4) (0.774M Na_2HPO_4 + 0.226M NaH_2PO_4,

pH 7.4) overnight at 4°C. Eyes obtained under wildlife permits (fire skink, hummingbird, zebra

finch) were perfused with 4% PFA and then kept in fixative for various periods, from 4 days

(hummingbird, fire skink) to 2 years (zebra finch). Cephalopod eyes were obtained by deeply

anesthetizing them in artificial seawater (ASW; 460mM $NaCl_2$, 10mM KCl, 10mM glucose,

10mM HEPES, 55mM $MgCl_2$, 11mM $CaCl_2$, 2mM glutamine, pH 7.4) supplemented with 110mM

$MgCl_2$, followed by rapid decapitation and fixation in 4% PFA in ASW at 4°C. Frog eyes labeled

with GFAP primary antibody (**Fig. 4E**) were fixed in 80% methanol + 20% dimethylsulfoxide at -

20°C (Dent's fixative protocol) overnight since 4% PFA + 3% sucrose fixation eliminated

antibody binding. After fixation, all eyes were processed the same way. Post-fixed eyes were

cryoprotected in increasing gradients of sucrose (10%, 20%, and then 30%) diluted in 0.1 M PB.

Fully cryoprotected eyes were then embedded in Tissue Tek O.C.T. (*S*akura Finetek, Torrance,

CA, USA) and sagittal sections were cut at a thickness of 12-16 µm. Sections were picked up on

room temperature SuperFrost slides (Thermo Fisher Scientific, Waltham, Massachusetts, USA)

and then left on a warm (42°C) hot plate for 8-10 min to increase tissue adherence to the slides.

Prepared slides were stored at -20°C until required.

2.4 Immunofluorescence and Epi-Fluorescence Microscopy

Frozen slides were thawed at room temperature, tissues were outlined with a hydrophobic

marker (ImmEdge™, MJC Biolynx, Brockville, ON, Canada), and then slides were washed 3x8

mins in 1X phosphate buffered saline (PBS, pH 7.4) (137 mM NaCl + 2.7 mM KCl + 10 mM

Na_2HPO_4 + 1.8 mM KH_2PO_4) at room temperature (~20°C). After washing, slides were blocked

in blocking buffer (1% goat serum + 0.1% Triton X-100 in 1X PBS) for 30 minutes in a humid

chamber at room temperature.

Slides were washed again 3x8 min in 1X PBS and then sections were labeled with GFP-

$EndoN_{DM}$ (10 µg/mL; Dr. Lisa Willis[73], University of Alberta) or a primary antibody to polySia,

mAb735 (1 µg/mL; Absolute Antibody, AB00240-2.0) in dilution buffer (0.1% goat serum + 0.1%

TX-100 in 1X PBS) overnight in a humid chamber at room temperature. After the overnight

incubation, slides were washed three times for 8 min each in 1X PBS, and then counter labels

and the secondary antibody for mAb735 (AF488 goat anti-mouse, Jackson Immunoresearch,

West Grove, PA, USA) were applied in dilution buffer and incubated for 4–6 h at 20°C. All

tissues were counter labeled with Hoechst 33342 (Millipore Sigma, Burlington, Massachusetts,

USA) and wheat germ agglutinin (Thermo Fisher, Waltham, Massachusetts, USA) to visualize

nuclei and photoreceptor outer segments, respectively. Slides were washed a final time for 3x8

min in 1X PBS and coverslipped using Mowiol Cover Slip Mounting Medium without DABCO[74].

For double labeling experiments, retinal sections were incubated with GFP-$EndoN_{DM}$ and the

primary antibody overnight at the same time. Antibodies used in this study and their

concentrations are listed in **Supplementary Table 1**.

Slides were imaged using either a 1) Leica DM6000B epi-fluorescence microscope with a 20x

objective (air, N.A. 0.50), Leica K3M Camera (3072 px x 2048 px), and Leica Software; 2) a

Zeiss Airyscan LSM900 with a 40x objective (water, N.A. 1.2; killifish); or a 3) Zeiss LSM 700

302   with a 63x objective (Oil, N.A. 1.40; zebrafish). Post-imaging processing was performed using

303   Affinity Photo 2 (Serif, West Bridgford, UK) and FIJI[75].

2.5 GFP-EndoN$_{DM}$ and mAb735 Characterization

Endoneuraminidase F (EndoN) is a polySia hydrolase that was originally isolated from the

tailspike of an *E. coli* K1-specific bacteriophage[76,77]. A double mutant (EndoN$_{DM}$) was

subsequently engineered to be a catalytically dead lectin that can detect and bind to polySia

chains with high affinity ($K_D = 10^{-8}$ M) without hydrolysing it[78]. GFP-tagged EndoN$_{DM}$ (GFP-

EndoN$_{DM}$) is an simple and efficient marker that has been used for the detection of polySia in

numerous cell culture systems and applications[42,76,79,80]. mAb735 is a primary mouse

monoclonal antibody (Absolute antibody, AB00240) that specifically recognizes α2,8-linked

polySia chains of 8 or longer[81,82]. In adult mice, mAb735 labels polySia-NCAM, attributed to

Müller glia and astrocytes in the retina and optic nerve[52]. mAb735 has also previously been

reported to label Müller glia in adult rat retina[60].

To validate that retinal GFP-EndoN$_{DM}$ and mAb735 positive signal in our vertebrate retinas were

due to binding to polySia, slides with mouse retina were incubated for 30 min at 37°C with the

active polySia hydrolase EndoN[77]. PBS was used as a negative control to keep slides hydrated

in the absence of the hydrolase. After the 30 min incubation, slides were washed 3x8 min with

1X PBS and then labeled with GFP-EndoN$_{DM}$ or mAb735. GFP-EndoN$_{DM}$ and mAb735 labeling

were eliminated by pre-treatment with polySia hydrolase (**Supplementary Fig. S2B,D**). We

found that there was non-specific secondary antibody labeling in the intraretinal and vitreal

blood vessels, the RPE, the choroid, and the sclera in mice and rats, this is expected due to

cross-reactivity of mouse secondary antibodies to rodent endogenous blood plasma IgGs

(**Supplementary Fig. S2E**). This cross reactivity was not seen in non-rodent species.

3. Results

3.1 GFP-EndoN$_{DM}$ and mAb735 localization in developing non-human vertebrate retinas

To determine the extent of polySia expression in developing non-human vertebrate retinas, we investigated the pattern of GFP-EndoN$_{DM}$ and mAb735 labeling in the early free-swimming larval retinas of zebrafish (*Danio rerio*) (~5 days post fertilization (dpf)), turquoise killifish (*Nothobranchius furzeri*) (golden eye stage, ~21 dpf), and African clawed frog (*Xenopus laevis*) (N.F stage 40, ~5 dpf)[83–85]. We also looked at postnatal (P3) mouse (*Mus musculus*) and (P1) rat (*Rattus norvegicus*) retina, before retinal lamination is complete. GFP-EndoN$_{DM}$ and mAb735 labeling patterns were comparable (**Fig. 1**). PolySia was detected in the retina and brain of larval killifish (**Fig. 1A-B, I**) and larval frog (**Fig. 1E-F,K**). It was also detected in the retinas of postnatal P3 mouse (**Fig. 1G,L**) and postnatal P1 rat (**Fig. 1H,M**), but not in larval zebrafish retina (**Fig. 1D,J**). Although there was no GFP-EndoN$_{DM}$ or mAb735 signal in the larval zebrafish retina, there was significant positive GFP-EndoN$_{DM}$ signal in the larval zebrafish brain (**Fig. 1C**). In larval frog and killifish retinas, GFP-EndoN$_{DM}$ and mAb735 positive signal was present in the entire Müller glial cell body, from the ELM to the endfeet, but was strongest near the ONL and the OPL, and the NFL. In P1 rat and P3 mouse, the retina is not yet fully laminated, but there was a strong GFP-EndoN$_{DM}$ and mAb735 positive band near the lower one-third of the retina (IPL) and diffuse labeling through the neuroblast layer (NBL) up to the developing ELM (**Fig. 1E,F; J,K**).

3.2 GFP-EndoN$_{DM}$ and mAb735 localization in adult non-human vertebrate and cephalopod retinas

To investigate polySia expression in adult non-human vertebrate retinas, we investigated the localization of GFP-EndoN$_{DM}$ and mAb735 in 8 different species: turquoise killifish, zebrafish,

hummingbird (*Calypte anna*), zebra finch (*Taeniopygia guttata*), fire skink (*Lepidothyris*

fernandi), African clawed frog, mouse, and rat. GFP-EndoN$_{DM}$ (**Fig. 2**) and mAb735 (**Fig. 3**)

positive labeling patterns were comparable, and resembled the labeling patterns expected from

Müller glia. PolySia expression in adult fishes was significantly different than all other

vertebrates examined, and slightly different from each other. PolySia expression in killifish was

limited to the ONL, the OPL, the INL, and the S1 sublaminae of the IPL; there was also some

low intensity signal in the NFL (**Fig. 2A, Fig. 3A**). Expression of polySia in zebrafish was more

confined, and signal was observed only in the ONL, the OPL, and the very top of the INL (**Fig.**

2B, Fig. 3B). In adult frogs, polySia expression extended from the ONL to the NFL. PolySia in

the INL exhibited a stratified pattern with arborisations in the S1, S3, and S5 IPL sublamina

(**Fig. 2C, Fig. 3C**). In frogs, polySia signal intensity was brightest in the NFL for GFP-EndoN$_{DM}$

labeling (**Fig. 2C**), whereas the ONL labeling was brightest when using mAb735 (**Fig. 3C**),

though the overall expression patterns were the same. For the fire skink, polySia expression

was strongest in the OPL and the NFL, and IPL sublaminae appeared to be thicker than those

observed in frog, mice, and rats (**Fig. 2D, Fig. 3D**). There was greater background fluorescence

observed in fire skink retinas labeled with mAb735, compared to GFP-EndoN$_{DM}$. PolySia

expression in birds was similar to frogs and lizards. GFP-EndoN$_{DM}$ and mAb735 labeled the

ONL, OPL, IPL, and NFL, as well as S1, S3, and S5 IPL sublaminae (**Fig. 2E-F, Fig. 3E-F**).

Background fluorescence and NFL signal was more intense in retinas labeled with mAb735

compared to GFP-EndoN$_{DM.}$ Finally, rodents had similar polySia expression patterns to frogs,

lizards, and birds (**Fig. 2G-H, Fig 3. G-H**). Whether labeled with GFP-EndoN$_{DM}$ or mAb735,

polySia expression extended from the ONL, throughout the retina, to the NFL. A notable

difference between rodents and other animals was the presence of polySia labeling in the apical

processes of the Müller glia, which extend beyond the ELM into the OS space. IPL S1, S3, and

S5 sublamina were also especially apparent in rodents compared to lizards and birds.

The labeling patterns of GFP-EndoN$_{DM}$ and mAb735 in adult retinas fixed less than 24 hrs were

equivalent, but mAb735 was less effective at labeling tissues that had been fixed for significant

periods of time, particularly the fire skink and the zebra finch, which had high background

fluorescence and less definitive Müller glia-like labeling patterns (**Fig. 3D,F**). mAb735 also

appeared to have greater non-specific labeling of photoreceptor outer segments in killifish (**Fig.**

3A), frog (**Fig. 3C**), and zebra finch (**Fig. 3F**). Other slight variations in signal localization were

observed between GFP-EndoN$_{DM}$ and mAb735 due to cross-reactivity of the secondary

antibody with rodent sclera, RPE, and intraretinal blood vessels (**Fig. 2G-H, Fig. 3G-H,**

Supplementary Fig. S2E).

In addition to vertebrate retinas, we also had the opportunity to examine potential polySia

expression in the retina and optic lobe of two different cephalopods: *Octopus bimaculoides* and

squid (*E. berryi*). Unlike vertebrate retinas and brain, there was no significant labeling of

cephalopod retinas or optic lobe using GFP-EndoN$_{DM}$ (**Supplementary Fig. S3**).

3.3 GFP-EndoN$_{DM}$ co-labeling with NCAM and established Müller glia markers

NCAM is the most common carrier of polySia in the CNS and Müller glia are known to express

NCAM in zebrafish[63,86], so we performed co-labeling with NCAM and GFP-EndoN$_{DM}$ in adult frog

to determine whether polySia expression in the retina was confined to the Müller glia. Co-

labeling of NCAM and GFP-EndoN$_{DM}$ demonstrated strong co-localization within the processes

in the outer nuclear layer, cell membranes in the inner nuclear layer, and in the nerve fibre layer

(**Fig. 4A-C**). GFP-EndoN$_{DM}$ labeling was higher intensity in the outer nuclear layer compared to

the rest of the retina and compared to NCAM labeling, which was more consistent across the

whole retina. We also investigated the co-localization of GFP-EndoN$_{DM}$ with vimentin in adult

frog. Vimentin and GFAP label just the Müller glia cytoskeleton (**Fig. 4D-E**). Vimentin labeling

co-localized with GFP-EndoN$_{DM}$ in the inner plexiform layer, where the GFP-EndoN$_{DM}$ signal

appeared to associated with the vimentin cytoskeleton (**Fig. 4F**).

Finally, we performed double-labeling with GS and GFP-Endo-N$_{DM}$ in larval killifish, adult

zebrafish, adult frog, adult hummingbird, adult mouse, and adult rat. This was done to verify that

polySia co-localizes with a known Müller glia marker in multiple species. GS labeled the tips of

the Müller glia at the ELM, Müller glial extensions in the ONL and INL, and their endfeet with

high intensity in all species examined (**Fig. 4G-L**). GS labeling of the ONL and ELM was

particularly intense in the hummingbird (**Fig. 4J**). GFP-EndoN$_{DM}$ labeling co-localized with GS at

the ELM, ONL, INL, and NFL. The intensity of GS was higher in the ONL and the NFL than

GFP-EndoN$_{DM}$. There was also GFP-EndoN$_{DM}$ signal in the apical processes of mouse and rat

Müller glia (**Fig. 4K-L**). The most apparent co-localization of GS and GFP-EndoN$_{DM}$ occurred in

the ELM, ONL, INL, and NFL. There was less co-localization of GS and GFP-EndoN$_{DM}$ in the

OPL and the IPL, where GFP-EndoN$_{DM}$ signal was more intense and demonstrated IPL

sublaminae organization where GS did not exhibit significant IPL sublaminae signal. We did not

perform double-labeling with mAb735 because all primary antibodies were mouse monoclonal

antibodies and the labeling patterns for GFP-EndoN$_{DM}$ and mAb735 were effectively

indistinguishable.

3.4 GFP-EndoN$_{DM}$ labeling in a mouse model of severe retinal degeneration

To test whether polySia labeling of Müller glia could also visualize gliosis, we obtained eyes

from aryl hydrocarbon receptor-interacting protein-like 1 knockout (*Aipl1*$^{-/-}$) mice, aged P97.

These mice are a model for human Leber Congenital Amaurosis (LCA), and have been reported

previously to exhibit rapid photoreceptor degeneration, loss of electroretinogram (ERG), and

signs of Müller gliotic changes, as early as P14[87]. We were able to visualize gliotic changes in

these mice using GFP-EndoN$_{DM}$ and mAb735 (**Fig. 5**). Double-labeling with GS showed strong

correlation between high intensity labeling patterns near the outer plexiform layer and in the

endfeet. As seen with healthy tissues, GFP-EndoN$_{DM}$ had increased signal around cell bodies in

the inner nuclear layer and the IPL sublaminae labeling remained intact (**Fig. 5A-B,D**).

3.5 GFP-EndoN$_{DM}$ localization in aged human retina and retinal organoids

Finally, we tested GFP-EndoN$_{DM}$ localization in human retina and patient-derived retinal

organoids; mAb735 was not tested due to tissue scarcity. The labeling pattern of GFP-EndoN$_{DM}$

is very similar to what is seen in other vertebrate retinas, including the IPL sublaminae (**Fig. 6A-**

C). The bright band of labeling at the ELM was not as distinct as observed in other vertebrates,

but was most visible in tissues sampled from just outside of the macula (**Fig. 6B**). One sample

also exhibited choroidal Schwann cells that labeled intensely with GFP-EndoN$_{DM}$ (**Fig. 6A, D-E**).

There was significant red and green autofluorescence in the RPE (**Fig. 6A-E).**

Wildtype and Parkinson's patient-derived retinal organoids also exhibited strong expression of

GFP-EndoN$_{DM}$. Labeling was dispersed throughout the organoid in intensely-labelled areas that

resembled plaques, laminations, and the external limiting membrane (**Fig. 7**). ELM-like

structures were especially apparent along the outer organoid surface, where photoreceptor

outer segments should be, and near photoreceptor rosettes that were present inside of the

organoids. It also appeared as if more plaques were present in patient-derived organoids

compared to healthy ones, although this was not quantified.

4. Discussion

4.1 PolySia during development and homeostasis across vertebrate species

In this study, we showed that GFP-EndoN$_{DM}$ and mAb735, which label polySia, are effective

Müller glia markers in most common model organisms, retinal organoids, and human tissues.

We report that polySia is expressed throughout the entirely of the Müller glia cell body, from

ELM to endfeet, with fine banding in the S1, S3, and S5 IPL sublaminae in healthy retinas of

humans and non-fish vertebrates. Mice and rats also exhibit significant polySia expression in

the Müller glia apical processes that extend beyond the ELM. Interestingly, polySia labeling in

zebrafish and killifish is restricted to a very defined region of the glia in the outer retina. The

reason for this restriction of expression compared to other vertebrates remains unknown.

Our data agree with previous reports of polySia-NCAM expression in the retina of adult

zebrafish[63], salamander (*Pleurodeles waltl*)[62], frog (*Lithobates pipiens*)[61], chicken[88], and rats[60].

These data are, to our knowledge, the first reports of polySia expression in larval and adult

killifish retinas. We found that α2,8-Neu5Ac polySia is not significantly expressed in cephalopod

retinas or optic lobes. There is little literature on sialic acids in mollusks, as it is generally

accepted that they do not express them. However, one previous report indicated the presence

of Neu5Ac in *Octopus vulgaris* and *Todarodes pacificus* hepatopancreatic and cerebral ganglia

tissues, on the basis of negative ion electrospray ionization mass spectrometry (ESI-MS) and

fluorometric high-performance liquid chromatography (HPLC)[89].

We believe that retinal polySia detected by GFP-EndoN$_{DM}$ and mAB735 corresponds primarily

to Müller glia on the basis of co-labeling experiments with NCAM, vimentin, and GS, which

demonstrated significant overlap with GFP-EndoN$_{DM}$. It should be noted, however, that although

the predominant form of CNS-associated polySia is polySia-NCAM, other retinal proteins with

polySia modification include an alpha subunit of voltage-gated sodium channels, CD36, synaptic

cell adhesion molecule 1 (SynCAM 1), and neuropillin-2[90–93]. SynCAM 1 and voltage-gated

sodium channels are particularly abundant in the retina, and could contribute to polySia labeling

near the photoreceptor nuclei and the ganglion cell axons[94–97]. Retinal SynCAM 1 does not

appear to co-localize with Müller glia or GFAP, however[94,95].

While the presence of polySia-NCAM in the brain during development and adulthood has been

well-studied[44,52,53,58], there are comparatively fewer studies on changes in polySia-NCAM

expression in the retina over time or with disease. Most studies in the brain report that polySia

levels are high during embryonic development when cell migration and plasticity is most active,

and then levels decrease as animals age, with the exception of some areas where plasticity and

neurogenesis remain important[54–58,98]. These developmental changes are attributed to roles of

polySia in cell migration[99], axon growth, guidance and fasciculation[100–102], synaptic plasticity[103],

and dendritic sprouting[104,105]. Most of the animals assessed in our report were adults; however,

we were able to compare retinal polySia labeling between early developmental and adult life

stages for killifish, zebrafish, frogs, mice, and rats. PolySia was present in Müller glia during

early retinal development, before the retina is fully laminated, and in adulthood. In most

vertebrate species examined, the intensity of polySia labeling within the retina did not change

significantly between larval/post-natal and adult life stages. The exception to this is the bony

fishes. PolySia was absent in free-swimming larval zebrafish and restricted to the outer retina in

adults. On the other hand, polySia expression was more widespread in the larval killifish retina

including the ONL, OPL, INL, IPL, and NFL, compared to the adult killifish retina where it was

restricted to the outer retina and the S1 IPL sublamina. We observed polySia labelling in the

larval zebrafish, killifish, and frog brains in agreement with previous reports of high polySia

expression in the vertebrate brain during embryonic development[44,52,53,58]. Zebrafish are popular

aquatic model organisms for the study of retinal development and disease[106], and interest in

killifish as a model of retinal and brain disease is increasing due to their unique lifecycle and fast

aging[107–109], so the finding that polySia expression in the Müller glia of fishes differs from other

vertebrates, even amphibians, is a noteworthy consideration for future studies using these

animals.

The IPL is divided into ON- (S3, S4, and S5) and OFF- (S1 and S2) sublaminae, which

represent termination layers of rod and cone bipolar-specific circuitry. Here, we report polySia

labeling of fine arborisations in the S1, S3, S5 IPL sublaminae in frog, lizard, birds, rodents, and

humans. Our data align with previous reports of Müller glia fine processes in the IPL sublaminae

in a number of other species including cat, dog, echidna, guinea pig, rabbit, pigeon, chicken,

and monkey[88,110–112]. Recently, a transgenic mouse was used to confirm that the Müller glia-

associated IPL sublaminae are located above, between, and below the ON and OFF cholinergic

acetyltransferase (ChAT) positive amacrine cell layers[32], in the S1 OFF-sublamina and the S3

and S5 ON-sublaminae[113,114]. Although fish also have these IPL sublaminae, and S1 was

labeled with GFP-EndoN$_{DM}$ in killifish, S3 and S5 were not significantly labeled using polySia in

either fish species examined. This is likely a difference in protein modification with polySia

between fishes and other vertebrates, rather than significant differences in retinal circuitry

between bony fishes and other vertebrates.

In human tissues, the labeling of the ELM was not as apparent as other vertebrates examined.

This could be due to location within the eye, as tissue sections from just outside of the macula

had the best labeling of the ELM. It could also be due to age, since these tissue sections

represented aged humans (69-81 years), in comparison to the early to mid adulthood tissue

sections that were taken from all other vertebrates examined. There was also significant red-

green autofluorescence in the RPE of the human retinas, which is typical of aged retinas due to

the build-up of lipofuscins and other indigestible protein and lipid waste products[115,116]. One

human retina also demonstrated additional labeling in a subset of choroidal cells that was not

seen in other species. We suspect that these were Schwann cells because they did not co-

localize with choroidal melanocytes, and non-myelinating Schwann cells are present in the

choroid and have been reported previously to express polySia-NCAM[46,117]. Schwann cells are a

type of glial cell that is not usually found in healthy retinas, but instead are usually associated

with peripheral nervous system development and repair[118]. They can infiltrate into the retina

under conditions of photoreceptor degeneration and high oxidative stress, however, and

injection of Schwann cells has been shown to be protective against retinal ganglion cell axon

degeneration in Royal College of Surgeons rats[119,120].

PolySia in disease states

We report here, that polySia labeling is also useful for studying gliosis, which is a reactive

response observed in Müller glia and astrocytes to various forms of nervous system injury and

disease[22,121,122]. In cases of severe reactive gliosis, Müller cells lose functionality and form glial

scars, inhibiting neuronal regeneration and survival. These gliotic scars were detected using

GFP-EndoN$_{DM}$ in *Aipl1$^{-/-}$* mice, and were comparable to previously published results in the same

mouse model using GFAP and GS to detect gliosis[87].

Finally, we showed that polySia is an effective marker for Müller glia-like cells in retinal

organoids derived from human cell lines, and that the labeling patterns of polySia in retinal

organoids closely resembled those of vimentin[72]. A recent study investigated the impacts of the

loss of Neu5Ac on cortical organoid development and found that organoids with mutations in *N*-

acetylneuraminic acid synthase, an enzyme critical for the synthesis of polySia, had significantly

reduced proliferation, organization, and synaptic connectivity[123]. Although the role of polySia in

developing and adult Müller glia is still unknown, a number of studies have shown that

intravitreal injections of polySia are neuroprotective under conditions of AMD-like disease and

kainic acid-induced retinal ganglion cell excitotoxity[124–126]. It is possible that the Müller glia may

mediate this effect.

Antibodies vs. lectins

Because Müller glia play such a critical role in retinal physiology, it is important to have effective

tools to study them. Common labeling tools, such as primary antibodies against GFAP,

vimentin, and GS can sometimes have limitations in their applicability across various animal

models. First, there can be significant differences in protein structure between species, which

can affect the ability of antibodies to bind to their target. Most commercially available antibodies

are made to cross-react with human, mouse, and rat proteins, because they are the most

common animal and cell model organisms. While effective in these models, commercial

antibodies rarely adequately label protein targets from other species, such as frogs, fish, or

birds. The structure of polySia is highly conserved across vertebrates, eliminating these issues

with poor protein structure conservation across species.

Second, some antigens are poorly immunogenic, of limited availability, or are unstable, and

raising antibodies against them is therefore difficult. For example, *Xenopus laevis* may not

express GFAP, and instead expresses a protein that cross-reacts with both GFAP and vimentin

antibodies[127]. Furthermore, antibodies for unique model organisms are difficult, time consuming,

and expensive to try to generate, particularly if the sequence of the proteins of interest is

unknown or poorly annotated. For instance, the killifish genome currently lacks a curated

database like those available for most other model organisms such as Xenbase.org, ZFIN.org,

Flybase.org, and the mouse genome database (MGD), which makes designing custom

antibodies difficult.

Third, the molecular targets of GFAP, vimentin, and GS may not be ideal for visualization of

whole Müller glia cell bodies and fine processes. GFAP and vimentin are cytoskeletal markers,

and therefore do not label cell membranes or Müller glial arborisations. In fact, GFAP labels only

~15% of the volume of astrocytes, and has similar efficacy for Müller glia[31]. GS is a cytoplasmic

marker and is therefore not detected in Müller glia cell membranes or fine IPL arborisations,

resulting in missing morphological information. Labeling of GS can also be dependent on the

upregulation of the target protein glutamine synthetase, such as during disease, for best

visualization[29,128,129]. Nevertheless, we consider GS to be the better antibody for Müller glia

visualization. The GS polyclonal antibody used here cross-reacts with many different species,

and it better highlights Müller glial morphology than either vimentin or GFAP antibodies.

As for the polySia markers tested here, we consider GFP-EndoN$_{DM}$ to be a better marker than

mAb735. GFP-EndoN$_{DM}$ is a conjugate, therefore does not have cross-reactivity problems due

to non-specific binding of the mouse secondary antibody to endogenous IgGs. It can also be

used in double-labeling experiments with any species of primary antibody or other lectin, leading

to greater versatility in experimental design. Further advantages to GFP-EndoN$_{DM}$ are that it is

inexpensive to synthesize in large quantities, far more stable than antibodies, resistant to heat

treatment and detergents, and can be engineered to have different conjugates. For example,

the GFP can be switched out for a biotinylated version of EndoN$_{DM}$ that could be coupled to any

fluorescent streptavidin or streptavidin-coated agarose beads if one wanted to enrich Müller glia

from a dissociated retina sample[79]. Finally, binding of GFP-EndoN$_{DM}$ to polySia appears to be

resistant to formaldehyde over-fixation, as evidenced by its successful use in our study on

tissues that have been fixed for a significant amount of time (lizard and zebra finch) compared

to poorer mAb735 labeling of the same tissues. Direct conjugates can sometimes have low

signal due to 1:1 binding with the target, unlike the signal amplification from combined use of

primary and secondary antibodies. However, because GFP-EndoN$_{DM}$ binds to the α2,8 linkages

of polySia chains, it has the opportunity to bind tens to hundreds of times along these chains,

resulting in significant signal amplification similar to primary-secondary systems.

Conclusion

Here, we report the versatility of polySia as a marker of Müller glia in amphibian, avian,

mammalian, human, and retinal organoid tissues. Labeling polySia is convenient, simple, and

useful for the study of Müller glia across different species, developmental stages, and disease

states. We suggest GFP-EndoN$_{DM}$ and mAb735 as suitable alternatives or supplemental

markers to conventional Müller glia markers such as GFAP, vimentin, and GS in models or

experimental paradigms where using these reagents is inconvenient or not possible.

Acknowledgements

We thank the donors to the Iowa Lions Eye bank for their generosity and commitment to

science, for they have given the incredible gift of a part of themselves so that we may learn

more about the human body. Thank you to the following persons for providing human eyes,

animal eyes, and retinal organoids, without which this project would not have been possible:

Maryam Hejazi (University of Alberta; adult wildtype mice), Dr. Daniel Hass (University of

Washington, Hurley Lab; *Aipl1*$^{-/-;-/+}$ mice), Dr. Christine Webber (University of Alberta; adult rat),

Nichole Vestby (University of Alberta, Director of Operations NCAS), Dr. Matthew Macauley

(University of Alberta; postnatal mouse), Dr. Francis Plane (University of Alberta; postnatal rat),

Dr. Zach Hall (in memoriam; University of Alberta; larval zebrafish), Dr. Cristian Gutierrez

(University of Alberta, Wylie Lab; hummingbird, fire skink), Dr. Judit Pungor (University of

Oregon, Niell Lab; cephalopods), Dr. Sandra Tenreiro (NOVA Medical School, Universidade

Nova de Lisboa; retinal organoids), and Dr. Robert F. Mullins (University of Iowa; human

retinas). We also wish to acknowledge the Marine Resources Center (Woods Hole, MA, USA)

for the husbandry and care of cephalopods for the greater research community.

Funding Statement

We wish to acknowledge and thank the donors of Macular Degeneration Research, a program

of the BrightFocus Foundation, for their generous support of this research program. We also

acknowledge and wish to thank the donors of Fighting Blindness Canada, for the generosity and

commitment to funding vision science. This research was funded by a BrightFocus Foundation

Macular Degeneration Postdoctoral Award (BJC; M2021001F), a BrightFocus Foundation

Macular Degeneration Early Career Investigator Award (BJC; M2024011N), a Fighting

Blindness Canada Early Career Researcher Grant, and University of Alberta Startup Funds,

funded by the Royal Alexandra Hospital Foundation (BJC). ADNNT and ANSJ were funded by

an Undergraduate Research Initiative award (URI), and ADNNT was also funded by a Women

and Children Health Research Institute (WCHRI) summer research student incentive. LMK was

funded by an NSERC summer student research award. NCLN was supported by a BrightFocus

Macular Degeneration Postdoctoral Fellowship (M2022002F), a Banting Postdoctoral

Fellowship (CIHR), a Moorfields Eye Charity Springboard Award (SB-24B-106), a Fight for Sight

Seed Grant, a Macular Society Seedcorn, and an Odette Maymon ECR Research Fund Grant.

NCLN works under the supervision of Ryan B. MacDonald, whose lab is supported by a BBSRC

David Phillips Fellowship (BB/S010386/1) and a BBSRC Partnering award (BB/V018078/1).

Moorfields Eye Charity funds the Institute of Ophthalmology Killifish system (GR001613). The

authors declare no conflict of interest.

References

1. Reichenbach, A. & Bringmann, A. New functions of Müller cells. *Glia* **61**, 651–678 (2013).
2. Xue, Y. *et al.* The role of retinol dehydrogenase 10 in the cone visual cycle. *Sci. Rep.* **7**, 2390 (2017).
3. Poitry-Yamate, C., Poitry, S. & Tsacopoulos, M. Lactate released by Muller glial cells is metabolized by photoreceptors from mammalian retina. *J. Neurosci.* **15**, 5179–5191 (1995).
4. Tworig, J. M. & Feller, M. B. Müller Glia in Retinal Development: From Specification to Circuit Integration. *Front. Neural Circuits* **15**, 815923 (2022).
5. Sakami, S., Imanishi, Y. & Palczewski, K. Müller glia phagocytose dead photoreceptor cells in a mouse model of retinal degenerative disease. *FASEB J.* **33**, 3680–3692 (2019).
6. Bassett, E. A. & Wallace, V. A. Cell fate determination in the vertebrate retina. *Trends Neurosci.* **35**, 565–573 (2012).
7. Reichenbach, A. *et al.* What do retinal Müller (glial) cells do for their neuronal 'small siblings'? *J. Chem. Neuroanat.* **6**, 201–213 (1993).
8. Newman, E. & Reichenbach, A. The Müller cell: a functional element of the retina. *Trends Neurosci.* **19**, 307–312 (1996).
9. Dubois-Dauphin, M. *et al.* Early postnatal Müller cell death leads to retinal but not optic nerve degeneration in NSE-Hu-Bcl-2 transgenic mice. *Neuroscience* **95**, 9–21 (1999).
10. Byrne, L. C. *et al.* AAV-Mediated, Optogenetic Ablation of Müller Glia Leads to Structural and Functional Changes in the Mouse Retina. *PLoS ONE* **8**, e76075 (2013).
11. Shen, W. *et al.* Conditional Müller Cell Ablation Causes Independent Neuronal and Vascular Pathologies in a Novel Transgenic Model. *J. Neurosci.* **32**, 15715–15727 (2012).
12. NILSSON, S. E. RECEPTOR CELL OUTER SEGMENT DEVELOPMENT AND ULTRASTRUCTURE OF THE DISK MEMBRANES IN THE RETINA OF THE TADPOLE (RANA PIPIENS). *J Ultrastruct Res* **11**, 581–602 (1964).
13. Bringmann, A. & Wiedemann, P. Müller Glial Cells in Retinal Disease. *Ophthalmologica* **227**, 1–19 (2012).
14. Pfeiffer, R. L. *et al.* Pathoconnectome Analysis of Müller Cells in Early Retinal Remodeling. *Adv. Exp. Med. Biol.* **1185**, 365–370 (2019).
15. Reichenbach, A. *et al.* The structure of rabbit retinal Müller (glial) cells is adapted to the surrounding retinal layers. *Anat. Embryol. (Berl.)* **180**, 71–79 (1989).
16. Omri, S. *et al.* The outer limiting membrane (OLM) revisited: clinical implications. *Clin. Ophthalmol. Auckl. NZ* **4**, 183–195 (2010).
17. Zhang, K. Y. & Johnson, T. V. The internal limiting membrane: Roles in retinal development and implications for emerging ocular therapies. *Exp. Eye Res.* **206**, 108545 (2021).
18. Hamon, A. *et al.* Linking YAP to Müller Glia Quiescence Exit in the Degenerative Retina. *Cell Rep.* **27**, 1712-1725.e6 (2019).
19. Hamon, A., Roger, J. E., Yang, X. & Perron, M. Müller glial cell-dependent regeneration of the neural retina: An overview across vertebrate model systems. *Dev. Dyn.* **245**, 727–738 (2016).
20. Wan, J. & Goldman, D. Retina regeneration in zebrafish. *Curr. Opin. Genet. Dev.* **40**, 41–47 (2016).

21.   Bringmann, A. *et al.* Cellular signaling and factors involved in Müller cell gliosis: Neuroprotective and detrimental effects. *Prog. Retin. Eye Res.* **28**, 423–451 (2009).

22.   Bringmann, A. *et al.* Müller cells in the healthy and diseased retina. **25**, 397–424 (2006).

23.   Bringmann, A. *et al.* Role of retinal glial cells in neurotransmitter uptake and metabolism. *Neurochem. Int.* **54**, 143–160 (2009).

24.   Navneet, S., Wilson, K. & Rohrer, B. Müller Glial Cells in the Macula: Their Activation and Cell-Cell Interactions in Age-Related Macular Degeneration. *Investig. Opthalmology Vis. Sci.* **65**, 42 (2024).

25.   Edwards, M. M. *et al.* Subretinal Glial Membranes in Eyes With Geographic Atrophy. *Investig. Opthalmology Vis. Sci.* **58**, 1352 (2017).

26.   Edwards, M. M. *et al.* Idiopathic preretinal glia in aging and age-related macular degeneration. *Exp. Eye Res.* **150**, 44–61 (2016).

27.   Palko, S. I. *et al.* Compartmentalized citrullination in Muller glial endfeet during retinal degeneration. *Proc. Natl. Acad. Sci.* **119**, e2121875119 (2022).

28.   Bignami, A. & Dahl, D. The radial glia of Müller in the rat retina and their response to injury. An immunofluorescence study with antibodies to the glial fibrillary acidic (GFA) protein. *Exp. Eye Res.* **28**, 63–69 (1979).

29.   Eisenfeld, A. J., Bunt-Milam, A. H. & Sarthy, P. V. Müller cell expression of glial fibrillary acidic protein after genetic and experimental photoreceptor degeneration in the rat retina. *Invest. Ophthalmol. Vis. Sci.* **25**, 1321–1328 (1984).

30.   Chen, H. & Weber, A. J. Expression of glial fibrillary acidic protein and glutamine synthetase by Müller cells after optic nerve damage and intravitreal application of brain-derived neurotrophic factor. *Glia* **38**, 115–125 (2002).

31.   Bushong, E. A., Martone, M. E., Jones, Y. Z. & Ellisman, M. H. Protoplasmic Astrocytes in CA1 Stratum Radiatum Occupy Separate Anatomical Domains. *J. Neurosci.* **22**, 183–192 (2002).

32.   Wang, J. *et al.* Anatomy and spatial organization of Müller glia in mouse retina. *J. Comp. Neurol.* **525**, 1759–1777 (2017).

33.   Sarthy, P. V., Fu, M. & Huang, J. Developmental expression of the Glial fibrillary acidic protein (GFAP) gene in the mouse retina. *Cell. Mol. Neurobiol.* **11**, 623–637 (1991).

34.   Stanley, P., Moremen, K. W., Lewis, N. E., Taniguchi, N. & Aebi, M. N-Glycans. in *Essentials of Glycobiology* (eds Varki, A. et al.) (Cold Spring Harbor Laboratory Press, Cold Spring Harbor (NY), 2022).

35.   Sato, C. & Kitajima, K. Polysialylation and disease. *Mol. Aspects Med.* **79**, 100892 (2021).

36.   Marki, A., Esko, J. D., Pries, A. R. & Ley, K. Role of the endothelial surface layer in neutrophil recruitment. *J. Leukoc. Biol.* **98**, 503–515 (2015).

37.   Ito, T. *et al.* Recognition of N-glycolylneuraminic acid linked to galactose by the alpha2,3 linkage is associated with intestinal replication of influenza A virus in ducks. *J. Virol.* **74**, 9300–9305 (2000).

38.   Nemanichvili, N. *et al.* Wild and domestic animals variably display Neu5Ac and Neu5Gc sialic acids. *Glycobiology* (2022) doi:10.1093/glycob/cwac033.

39.   Davies, L. R. L. *et al.* Metabolism of vertebrate amino sugars with N-glycolyl groups: resistance of α2-8-linked N-glycolylneuraminic acid to enzymatic cleavage. *J. Biol. Chem.* **287**, 28917–28931 (2012).

40.   Angata, K. & Fukuda, M. Roles of Polysialic Acid in Migration and Differentiation of Neural Stem Cells. in *Methods in Enzymology* vol. 479 25–36 (Elsevier, 2010).

41.   Heimburg-Molinaro, J. *et al.* Cancer vaccines and carbohydrate epitopes. *Vaccine* **29**, 8802–8826 (2011).

42.   Hunter, C. *et al.* Attenuation of Polysialic Acid Biosynthesis in Cells by the Small Molecule Inhibitor 8-Keto-sialic acid. *ACS Chem. Biol.* **18**, 41–48 (2023).

43.   Finne, J. Occurrence of unique polysialosyl carbohydrate units in glycoproteins of developing brain. *J. Biol. Chem.* **257**, 11966–11970 (1982).

44.   Rothbard, J. B., Brackenbury, R., Cunningham, B. A. & Edelman, G. M. Differences in the carbohydrate structures of neural cell-adhesion molecules from adult and embryonic chicken brains. *J. Biol. Chem.* **257**, 11064–11069 (1982).

45.   Pestronk, A., Schmidt, R. E., Bucelli, R. & Sim, J. Schwann cells and myelin in human peripheral nerve: Major protein components vary with age, axon size and pathology. *Neuropathol. Appl. Neurobiol.* **49**, e12898 (2023).

46.   Roche, P.-H. *et al.* Expression of cell adhesion molecules in normal nerves, chronic axonal neuropathies and Schwann cell tumors. *J. Neurol. Sci.* **151**, 127–133 (1997).

47.   Edelman, G. M. & Crossin, K. L. CELL ADHESION MOLECULES: Implications for a Molecular Histology. *Annu. Rev. Biochem.* **60**, 155–190 (1991).

48.   Rutishauser, U. Polysialic acid in the plasticity of the developing and adult vertebrate nervous system. *Nat. Rev. Neurosci.* **9**, 26–35 (2008).

49.   Mindler, K., Ostertag, E. & Stehle, T. The polyfunctional polysialic acid: A structural view. *Carbohydr. Res.* **507**, 108376 (2021).

50.   Rutishauser, U. & Landmesser, L. Polysialic acid in the vertebrate nervous system: a promoter of plasticity in cell-cell interactions. *Trends Neurosci.* **19**, 422–427 (1996).

51.   Sato, C. & Kitajima, K. Disialic, oligosialic and polysialic acids: distribution, functions and related disease. *J. Biochem. (Tokyo)* **154**, 115–136 (2013).

52.   Bartsch, U., Kirchhoff, F. & Schachner, M. Highly sialylated N-CAM is expressed in adult mouse optic nerve and retina. *J. Neurocytol.* **19**, 550–565 (1990).

53.   Hildebrandt, H. *et al.* Polysialic acid on the neural cell adhesion molecule correlates with expression of polysialyltransferases and promotes neuroblastoma cell growth. *Cancer Res.* **58**, 779–784 (1998).

54.   Amoureux, M.-C. *et al.* Polysialic Acid Neural Cell Adhesion Molecule (PSA-NCAM) is an adverse prognosis factor in glioblastoma, and regulates olig2 expression in glioma cell lines. *BMC Cancer* **10**, 91 (2010).

55.   Petridis, A. K., El Maarouf, A. & Rutishauser, U. Polysialic acid regulates cell contact-dependent neuronal differentiation of progenitor cells from the subventricular zone. *Dev. Dyn.* **230**, 675–684 (2004).

56.   Rutishauser, U. & Landmesser, L. Polysialic acid in the vertebrate nervous system: a promoter of plasticity in cell-cell interactions. *Trends Neurosci.* **19**, 422–427 (1996).

57.   Durbec, P. & Cremer, H. Revisiting the function of PSA-NCAM in the nervous system. *Mol. Neurobiol.* **24**, 53–64 (2001).

58.   Yang, Y. *et al.* Comparative Studies of Polysialic Acids Derived from Five Different Vertebrate Brains. *Int. J. Mol. Sci.* **21**, 8593 (2020).

59.   Murphy, J. A., Hartwick, A. T. E., Rutishauser, U. & Clarke, D. B. Endogenous Polysialylated Neural Cell Adhesion Molecule Enhances the Survival of Retinal Ganglion Cells. *Invest. Ophthalmol. Vis. Sci.* **50**, 861–869 (2009).

60.   Sawaguchi, A. *et al.* Multistratified Expression of Polysialic Acid and Its Relationship to VAChT-containing Neurons in the Inner Plexiform Layer of Adult Rat Retina. *J. Histochem. Cytochem.* **47**, 919–927 (1999).

61.   Fliesler, S. J., Cole, G. J. & Adler, A. J. Neural cell adhesion molecule (NCAM) in adult vertebrate retinas: Tissue localization and evidence against its role in retina-pigment epithelium adhesion. *Exp. Eye Res.* **50**, 475–482 (1990).

62.   Becker, C. G., Becker, T., Schmidt, A. & Roth, G. Polysialic acid expression in the salamander retina is inducible by thyroxine. *Dev. Brain Res.* **79**, 140–146 (1994).

63.   Kustermann, S., Hildebrandt, H., Bolz, S., Dengler, K. & Kohler, K. Genesis of rods in the zebrafish retina occurs in a microenvironment provided by polysialic acid-expressing Müller glia. *J. Comp. Neurol.* **518**, 636–646 (2010).

64.   Fiorito, G. *et al.* Guidelines for the Care and Welfare of Cephalopods in Research -A consensus based on an initiative by CephRes, FELASA and the Boyd Group. *Lab. Anim.* **49**, 1–90 (2015).

65.   Enterina, J. R. *et al.* Coordinated changes in glycosylation regulate the germinal center through CD22. *Cell Rep.* **38**, 110512 (2022).

66.   Morris, R. J. Thy-1, a Pathfinder Protein for the Post-genomic Era. *Front. Cell Dev. Biol.* **6**, 173 (2018).

67.   Hagood, J. S. Thy-1 as an Integrator of Diverse Extracellular Signals. *Front. Cell Dev. Biol.* **7**, 26 (2019).

68.   Barnstable, C. J. & Dräger, U. C. Thy-1 antigen: A ganglion cell specific marker in rodent retina. *Neuroscience* **11**, 847–855 (1984).

69.   Raymond, I. D., Vila, A., Huynh, U.-C. N. & Brecha, N. C. Cyan fluorescent protein expression in ganglion and amacrine cells in a thy1-CFP transgenic mouse retina. *Mol. Vis.* **14**, 1559–1574 (2008).

70.   Partida, G. J., Stradleigh, T. W., Ogata, G., Godzdanker, I. & Ishida, A. T. Thy1 Associates with the Cation Channel Subunit HCN4 in Adult Rat Retina. *Investig. Opthalmology Vis. Sci.* **53**, 1696 (2012).

71.   Haefliger, J.-A. *et al.* Targeting Cx40 (Connexin40) Expression or Function Reduces Angiogenesis in the Developing Mouse Retina. *Arterioscler. Thromb. Vasc. Biol.* **37**, 2136–2146 (2017).

72.   de Lemos, L. *et al.* Modelling neurodegeneration and inflammation in early diabetic retinopathy using 3D human retinal organoids. *Vitro Models* **3**, 33–48 (2024).

73.   Tajik, A., Phillips, K. L., Nitz, M. & Willis, L. M. A new ELISA assay demonstrates sex differences in the concentration of serum polysialic acid. *Anal. Biochem.* **600**, 113743 (2020).

74.   Mowiol mounting medium. *Cold Spring Harb. Protoc.* **2006**, pdb.rec10255 (2006).

75.   Schindelin, J. *et al.* Fiji⬚: an open-source platform for biological-image analysis. **9**, (2019).

76.   Jokilammi, A. *et al.* Construction of antibody mimics from a noncatalytic enzyme–detection of polysialic acid. *J. Immunol. Methods* **295**, 149–160 (2004).

77.   Rutishauser, U., Watanabe, M., Silver, J., Troy, F. A. & Vimr, E. R. Specific alteration of NCAM-mediated cell adhesion by an endoneuraminidase. *J. Cell Biol.* **101**, 1842–1849 (1985).

78.   Pelkonen, S., Aalto, J. & Finne, J. Differential activities of bacteriophage depolymerase on bacterial polysaccharide: binding is essential but degradation is inhibitory in phage infection of K1-defective Escherichia coli. *J. Bacteriol.* **174**, 7757–7761 (1992).

79.   Hunter, C. *et al.* Site-specific immobilization of the endosialidase reveals QSOX2 is a novel polysialylated protein. *Glycobiology* **34**, cwae026 (2024).

80.   Yu, C.-C. *et al.* A glyco-gold nanoparticle based assay for α-2,8-polysialyltransferase from Neisseria meningitidis. *Chem. Commun.* **49**, 10166 (2013).

81.   Bitter-Suermann, D. & Roth, J. Monoclonal antibodies to polysialic acid reveal epitope sharing between invasive pathogenic bacteria, differentiating cells and tumor cells. *Immunol. Res.* **6**, 225–237 (1987).

82.   Frosch, M., Görgen, I., Boulnois, G. J., Timmis, K. N. & Bitter-Suermann, D. NZB mouse system for production of monoclonal antibodies to weak bacterial antigens: isolation of an IgG antibody to the polysaccharide capsules of Escherichia coli K1 and group B meningococci. *Proc. Natl. Acad. Sci.* **82**, 1194–1198 (1985).

83.   Kimmel, C. B., Ballard, W. W., Kimmel, S. R., Ullmann, B. & Schilling, T. F. Stages of embryonic development of the zebrafish. *Dev. Dyn.* **203**, 253–310 (1995).

84.   Api, M. *et al.* Effects of Parental Aging During Embryo Development and Adult Life: The Case of *Nothobranchius furzeri*. *Zebrafish* **15**, 112–123 (2018).

85.   Nieuwkoop, P. D. & Faber, J. Normal table of Xenopus laevis (Daudin): a systematical and chronological survey of the development from the fertilized egg till the end of metamorphosis. (1994).

86.   Lenkowski, J. R. & Raymond, P. A. Müller glia: Stem cells for generation and regeneration of retinal neurons in teleost fish. *Prog. Retin. Eye Res.* **40**, 94–123 (2014).

87.   Singh, R. K., Kolandaivelu, S. & Ramamurthy, V. Early Alteration of Retinal Neurons in *Aipl1*$^{-/-}$ Animals. *Investig. Opthalmology Vis. Sci.* **55**, 3081 (2014).

88.   Daniloff, J., Chuong, C., Levi, G. & Edelman, G. Differential distribution of cell adhesion molecules during histogenesis of the chick nervous system. *J. Neurosci.* **6**, 739–758 (1986).

89.   Saito, M., Kitamura, H. & Sugiyama, K. Occurrence of gangliosides in the common squid and pacific octopus among protostomia. *Biochim. Biophys. Acta BBA - Biomembr.* **1511**, 271–280 (2001).

90.   Zuber, C., Lackie, P. M., Catterall, W. A. & Roth, J. Polysialic acid is associated with sodium channels and the neural cell adhesion molecule N-CAM in adult rat brain. *J. Biol. Chem.* **267**, 9965–9971 (1992).

91.	Galuska, S. P. *et al.* Synaptic cell adhesion molecule SynCAM 1 is a target for polysialylation in postnatal mouse brain. *Proc. Natl. Acad. Sci. U. S. A.* **107**, 10250–10255 (2010).

92.	Curreli, S., Arany, Z., Gerardy-Schahn, R., Mann, D. & Stamatos, N. M. Polysialylated neuropilin-2 is expressed on the surface of human dendritic cells and modulates dendritic cell-T lymphocyte interactions. *J. Biol. Chem.* **282**, 30346–30356 (2007).

93.	Yabe, U., Sato, C., Matsuda, T. & Kitajima, K. Polysialic acid in human milk. CD36 is a new member of mammalian polysialic acid-containing glycoprotein. *J. Biol. Chem.* **278**, 13875–13880 (2003).

94.	Ribic, A., Liu, X., Crair, M. C. & Biederer, T. Structural organization and function of mouse photoreceptor ribbon synapses involve the immunoglobulin protein synaptic cell adhesion molecule 1. *J. Comp. Neurol.* **522**, 900–920 (2014).

95.	Fujita, E., Urase, K., Soyama, A., Kouroku, Y. & Momoi, T. Distribution of RA175/TSLC1/SynCAM, a member of the immunoglobulin superfamily, in the developing nervous system. *Dev. Brain Res.* **154**, 199–209 (2005).

96.	Pietri, T., Easley-Neal, C., Wilson, C. & Washbourne, P. Six *cadm/synCAM* genes are expressed in the nervous system of developing zebrafish. *Dev. Dyn.* **237**, 233–246 (2008).

97.	Wang, J., Ou, S.-W. & Wang, Y.-J. Distribution and function of voltage-gated sodium channels in the nervous system. *Channels Austin Tex* **11**, 534–554 (2017).

98.	Kiss, J. Z., Troncoso, E., Djebbara, Z., Vutskits, L. & Muller, D. The role of neural cell adhesion molecules in plasticity and repair. *Brain Res. Rev.* **36**, 175–184 (2001).

99.	Ono, K., Tomasiewicz, H., Magnuson, T. & Rutishauser, U. N-CAM mutation inhibits tangential neuronal migration and is phenocopied by enzymatic removal of polysialic acid. *Neuron* **13**, 595–609 (1994).

100.	Silver, J. & Rutishauser, U. Guidance of optic axons in vivo by a preformed adhesive pathway on neuroepithelial endfeet. *Dev. Biol.* **106**, 485–499 (1984).

101.	Tang, J., Landmesser, L. & Rutishauser, U. Polysialic acid influences specific pathfinding by avian motoneurons. *Neuron* **8**, 1031–1044 (1992).

102.	Yin, X., Watanabe, M. & Rutishauser, U. Effect of polysialic acid on the behavior of retinal ganglion cell axons during growth into the optic tract and tectum. *Development* **121**, 3439–3446 (1995).

103.	Muller, D. *et al.* PSA–NCAM Is Required for Activity-Induced Synaptic Plasticity. *Neuron* **17**, 413–422 (1996).

104.	Muller, D., Stoppini, L., Wang, C. & Kiss, J. Z. A role for polysialylated neural cell adhesion molcule in lesion-induced sprouting in hippocampal organotypic cultures. *Neuroscience* **61**, 441–445 (1994).

105.	Aubert, I., Ridet, J. L., Schachner, M., Rougon, G. & Gage, F. H. Expression of L1 and PSA during sprouting and regeneration in the adult hippocampal formation. *J. Comp. Neurol.* **399**, 1–19 (1998).

106.	Noel, N. C. L., MacDonald, I. M. & Allison, W. T. Zebrafish Models of Photoreceptor Dysfunction and Degeneration. *Biomolecules* **11**, 78 (2021).

107. Bergmans, S. *et al.* Age-related dysregulation of the retinal transcriptome in African turquoise killifish. *Aging Cell* **23**, e14192 (2024).

108. Vanhunsel, S. *et al.* The killifish visual system as an in vivo model to study brain aging and rejuvenation. *Npj Aging Mech. Dis.* **7**, 22 (2021).

109. Noel, N. C. *et al.* Characterising Photoreceptor Subtypes in the Turquoise Killifish (Nothobranchius furzeri). *Invest. Ophthalmol. Vis. Sci.* **64**, 3735–3735 (2023).

110. Distler, C. & Dreher, Z. Glia Cells of the Monkey Retina—II. Müller Cells. *Vision Res.* **36**, 2381–2394 (1996).

111. Dreher, Z., Robinson, S. R. & Distler, C. Müller cells in vascular and avascular retinae: A survey of seven mammals. *J. Comp. Neurol.* **323**, 59–80 (1992).

112. Dreher, Z., Distler, C. & Dreher, B. Vitread proliferation of filamentous processes in avian Müller cells and its putative functional correlates. *J. Comp. Neurol.* **350**, 96–108 (1994).

113. Masland, R. H. The fundamental plan of the retina. *Nat. Neurosci.* **4**, 877–886 (2001).

114. Haverkamp, S. & Wässle, H. Immunocytochemical analysis of the mouse retina. *J. Comp. Neurol.* **424**, 1–23 (2000).

115. Brunk, U. T. & Terman, A. Lipofuscin: mechanisms of age-related accumulation and influence on cell function. *Free Radic. Biol. Med.* **33**, 611–619 (2002).

116. Călin, E. F. *et al.* Lipofuscin: a key compound in ophthalmic practice. *Romanian J. Ophthalmol.* **65**, 109–113 (2021).

117. Forestier, N. L., Lescs, M. & Gherardi, R. K. Anti-NKH-1 antibody specifically stains unmyelinated fibres and non-myelinating Schwann cell columns in humans. *Neuropathol. Appl. Neurobiol.* **19**, 500–506 (1993).

118. Bosch-Queralt, M., Fledrich, R. & Stassart, R. M. Schwann cell functions in peripheral nerve development and repair. *Neurobiol. Dis.* **176**, 105952 (2023).

119. Lashay, A. *et al.* Role of Schwann Cells in Preservation of Retinal Tissue Through Reduction of Oxidative Stress. *Med. Hypothesis Discov. Innov. Ophthalmol. J.* **8**, 323–332 (2019).

120. Dezawa, M. & Adachi-Usami, E. Role of Schwann cells in retinal ganglion cell axon regeneration. *Prog. Retin. Eye Res.* **19**, 171–204 (2000).

121. Garcia-Garcia, D., Locker, M. & Perron, M. Update on Müller glia regenerative potential for retinal repair. *Curr. Opin. Genet. Dev.* **64**, 52–59 (2020).

122. Poonam, S. & Rajesh, R. Retina regeneration: lessons from vertebrates. *Oxf. Open Neurosci.* 1–21 (2022).

123. Bu, Q. *et al.* Neurodevelopmental defects in human cortical organoids with *N*-acetylneuraminic acid synthase mutation. *Sci. Adv.* **9**, (2023).

124. Krishnan, A. *et al.* PolySialic acid-nanoparticles inhibit macrophage mediated inflammation through Siglec agonism: a potential treatment for age related macular degeneration. *Front. Immunol.* **14**, 1237016 (2023).

125. Karlstetter, M. *et al.* Polysialic acid blocks mononuclear phagocyte reactivity, inhibits complement activation, and protects from vascular damage in the retina. *EMBO Mol. Med.* **9**, 154–166 (2017).

126. Lobanovskaya, N. & Zharkovsky, A. A role of PSA-NCAM in the survival of retinal ganglion cells (RGCs) after kainic acid damage. *NeuroToxicology* **72**, 101–106 (2019).

127. Martinez-De Luna, R. I. *et al.* Müller glia reactivity follows retinal injury despite the absence of the glial fibrillary acidic protein gene in Xenopus. *Dev. Biol.* **426**, 219–235 (2017).

128. Gorovits, R., Avidan, N., Avisar, N., Shaked, I. & Vardimon, L. Glutamine synthetase protects against neuronal degeneration in injured retinal tissue. *Proc. Natl. Acad. Sci.* **94**, 7024–7029 (1997).

129. Lewis, G. P. & Fisher, S. K. Up-Regulation of Glial Fibrillary Acidic Protein in Response to Retinal Injury: Its Potential Role in Glial Remodeling and a Comparison to Vimentin Expression. in *International Review of Cytology* vol. 230 263–290 (Elsevier, 2003).

Figure 1. GFP-EndoN$_{DM}$ and mAb735 labeling in developing killifish (A-B,I), zebrafish (C-D, J), frog (E-F,K), mouse (G,L), and rat (H,M). PolySia is present in the retina and brain of killifish and frog (A-B, E,F), in the brain of zebrafish (C), and in the retina of postnatal mice and rats (G,H,L,M). Minor differences in labeling, such as increased scleral reactivity observed in mAb735-labeled mice (L) are the result of cross-reactivity of the secondary antibody (Supplementary Fig. S2). *Labels*: Green, GFP-EndoN$_{DM}$ (A-H), mAb735 (I-M); Blue, Hoechst. *Abbreviations*: OS, outer segments; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer; NFL, nerve fiber layer; NBL, neuroblast layer. Scale bar = 50 μm. n = 3-7.

Figure 2. GFP-EndoN$_{DM}$ reactivity in adult killifish (A), zebrafish (B), frog (C), fire skink (D), hummingbird (E), zebra finch (F), mouse (G), and rat (H). GFP-EndoN$_{DM}$ expression in adult killifish or zebrafish is most concentrated to the outer retina (ONL, OPL; A,B). In the killifish, the GFP-EndoN$_{DM}$ labeling extends to OFF sublamina S1 (A). All other species examined showed GFP-EndoN$_{DM}$ expression throughout the entirely of the retina, likely Müller glia, from ELM to endfeet (C-H). GFP-EndoN$_{DM}$ also exhibited increased intensity in bands in the IPL that corresponded with the S1 (OFF), S3 (ON), and S5 (ON) IPL sublaminae (C-H). *Labels*: Green, GFP-EndoN$_{DM}$; Blue, Hoechst. *Abbreviations*: OS, outer segment layer; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer; NFL, nerve fibre layer. Scale bar = 50 μm. n = 3-4 except fire skink (n = 1) and zebra finch (n = 2).

Figure 3. mAb735 reactivity in adult killifish (A), zebrafish (B), frog (C), fire skink (D), hummingbird (E), zebra finch (F), mouse (G), and rat (H). The labeling pattern of mAb735 in each species was the same as observed using GFP-EndoN$_{DM}$, including restriction of signal to the outer retina in killifish and zebrafish, and labeling of the IPL S1, S3, and S5 sublaminae in all other species. mAb735 was less effective than GFP-EndoN$_{DM}$ in highly fixed tissues such as the fire skink (D) and zebra finch (F) because background fluorescence was much higher. mAb735 also appeared to have greater non-specific labeling of photoreceptor outer segments in killifish (A), frog (C), and zebra finch (F). There was also non-specific cross reactivity of the secondary antibody (AF488 anti-mouse) in mouse and rat sclera, RPE, and intraretinal blood vessels (G-H). *Labels*: Green, mAb735; Blue, Hoechst. *Abbreviations*: Chr, choroid; RPE, retinal pigment epithelium; OS, outer segment layer; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer; NFL, nerve fibre layer. Scale bar = 50 µm. n = 3-4 except fire skink (n = 1) and zebra finch (n = 2).

Figure 4. Co-labeling experiments with GFP-EndoN$_{DM}$ and various other Müller glia markers. GFP-EndoN$_{DM}$ co-localized with NCAM (C) and vimentin (F) in frogs, and GS in larval killifish (G), adult zebrafish (H), adult frog (I), adult hummingbird (J), adult mouse (K), and adult rat (L). There were some differences in signal intensity between GFP-EndoN$_{DM}$ and GS. Generally, the ONL, and NFL were more intensely labeled with GS, and the OPL, IPL, and apical processes in mice and rats were more intensely labeled with GFP-EndoN$_{DM}$. GS and GFP-EndoN$_{DM}$ signals localized to the same spaces within the retina, except the apical processes (GFP-EndoN$_{DM}$ in mice and rats only). *Labels*: Green, GFP-EndoN$_{DM}$ (C,F, G-L), vimentin (D), GFAP (E); Magenta, NCAM (C), vimentin (F), GS (G-L). *Abbreviations*: GS, glutamine synthetase; ELM, external limiting membrane; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer; NFL, nerve fibre layer. Scale bar = 50 µm. n = 3.

Figure 5. Detection of gliosis in a 97 day old *Aipl1*[-/-] mouse by GFP-EndoN$_{DM}$ (A), mAb735 (B), glutamine synthetase (GS; C), and double-labeling with GFP-EndoN$_{DM}$ + GS (D). *Aipl1*[-/-] mice had complete loss of their outer segment and outer nuclear layer, and gliotic scarring in the OPL was visible using all markers (white arrows). GS and GFP-EndoN$_{DM}$ labeling of scarring was co-localized (D). *Labels*: Green, GFP-EndoN$_{DM}$ (A,D), mAb735 (B); Magenta, GS; Blue, Hoechst. *Abbreviations*: GS, glutamine synthetase; ELM, external limiting membrane; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer; NFL, nerve fibre layer. Scale bar = 50 μm. n = 3.

Figure 6. GFP-EndoN$_{DM}$ labeling in post-mortem human retina sections that are outside of the macula (A), close to the macula (B), and close to the optic nerve (C). The labeling pattern is very similar to what is seen in other vertebrate retinas, including the IPL sublaminae, but the bright band of labeling at the ELM is not as distinct as observed in other vertebrates in sections taken close to the macula and the optic nerve. Choroidal Schwann cells labeled brightly with GFP-EndoN$_{DM}$ in one of the samples (A, D-E, white arrow), and did not co-localize with melanocytes (E, black cells, black arrowhead). There was significant red and green channel RPE autofluorescence observed in all samples (A-E, white arrowheads). Labels: Magenta, wheat germ agglutinin (WGA); Green, GFP-EndoN$_{DM}$; Blue, Hoechst. *Abbreviations*: Chr, choroid; RPE, retinal pigment epithelium; OS, outer segment layer; ONL, outer nuclear layer; OPL, outer plexiform layer; INL, inner nuclear layer; IPL, inner plexiform layer; GCL, ganglion cell layer; NFL, nerve fibre layer. Scale bar = 50 μm. n = 3.

Figure 7. GFP-EndoN$_{DM}$ labeling in cross sections of healthy (A) and patient-derived (B) retinal organoids, aged 120 days. Expression patterns resembled external limiting membrane-like labeling (white arrows), plaque-like areas of high intensity labeling (asterisks), and areas of high intensity surrounding rosettes (black arrowheads). *Labels*: Magenta, wheat germ agglutinin (WGA); Green, GFP-EndoN$_{DM}$; Blue, Hoechst. Scale bar = 50 μm. n = 3

turquoise killifish — A) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, IPL, GCL, NFL

zebrafish — B) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, IPL, GCL, NFL

frog — C) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, IPL, GCL, NFL

fire skink — D) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, GCL, NFL

hummingbird — E) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, IPL, GCL, NFL

zebra finch — F) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, IPL, GCL

mouse — G) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, IPL, GCL, NFL

rat — H) GFP-EndoN_DM; OS, ONL, OPL, INL, OFF S1, S2, S3, ON S4, S5, IPL, GCL, NFL

Healthy

A)

Patient-derived

B)