Integrative multi-omic and phenotypic analysis of open raceway pond production of *Monoraphidium*
minutum 26B-AM reveals distinct stress signatures for scale-up and infection.

Georgios Kepesidis [a], Jenna Schambach [b], Daniel Yang [a], Elise Wilbourn [c], Thomas Sheffield [a], Tyler Eckles
[c], Olivia Watt [c], Matthew P. Hirakawa [a], Todd W. Lane [c], Raga Krishnakumar [a+]

a Biosecurity and Bioassurance, Sandia National Laboratories, Livermore, 94550, U.S.A.

b Environmental Systems Biology, Sandia National Laboratories, Albuquerque, 87123, U.S.A.

c Bioresource & Environmental Security, Sandia National Laboratories, Livermore, 94550, U.S.A.

+ Corresponding author

Abstract

Background: Green microalgae, such as *Monoraphidium minutum* 26B-AM, have garnered significant commercial interest due to their high biomass production and lipid yield, providing promising candidates for various bioprocessing applications. However, the economic viability of large-scale algal cultivation in open raceway ponds is limited by biocontamination and environmental stressors, necessitating deeper understanding of the molecular mechanisms that underpin resilience and productivity in these systems. We hypothesized that the molecular signature associated with the cellular responses of *M. minutum* to environmental stressors will reveal critical information for the timely prediction of resilience and productivity in algal cultures within open pond systems.

Results: To test this hypothesis, we conducted a longitudinal multi-omic study, integrating transcriptomics, proteomics, metabolomics, and phenomics, to monitor the acclimation, growth dynamics, and pathogen responses of algal cultures in two 1000 L raceway ponds, before and after the introduction of a pathogen as a stressor. We identified a number of molecular patterns that correlate with changes in the algal environment, and we can track these changes within the ponds per time. Furthermore, we identify scale-up and infection-specific molecular pathways through integrated multi-omics, showing that most patterns are unique to each studied stressor/transition.

Conclusions: Ultimately, this study demonstrates the utility of multi-omics observations at scale, revealing unique signatures and laying the groundwork for developing molecular detection techniques and predictive models that can improve the sustainability and efficiency of large-scale algae biomass production.

Keywords

Multi-omics, algae, *Monoraphidium minutum*, ponds, transcriptomics, metabolomics, proteomics, biomarkers

1. Background

Microalgae are a polyphyletic group of unicellular photosynthetic aquatic organisms that have attracted commercial interest since the middle of the past century because of their potential to advance biotechnology and biomanufacturing processes. Compared to land plants, their biomass yield is greater [1], and their high content of useful metabolites, such as lipids, has increased their presence in multiple industrial bioprocesses, including the production of food, pharmaceuticals, cosmetics, bioplastics, and biofuels. The high production costs, however, curtail most of these applications, especially for low-cost and high-volume products such as biofuels, making the use of algae currently unfeasible and unsustainable and restricting their potential [2] [3] [4].

A low-energy and low-cost method for large-scale production of algal biomass, given the financial restrictions for algal cultivation mentioned earlier, is the paddlewheel-based open raceway pond cultivation system [5] [6]. These widely used systems allow the cultures to grow in direct relation to the

surrounding environment and the local weather conditions. Additionally, these ponds can be established
on land that is not suitable for terrestrial agriculture, thereby utilizing a resource that would otherwise
remain unused.

Despite the aforementioned advantages of these open systems, a number of barriers preclude the overall
economic viability of bioprocesses within these systems. One crucial barrier is the susceptibility of the
cultures to environmental "intruders", which threaten both the quality of the final product and the overall
survival of the pond cultures [7]. These events lead to pond crashes and a significant increase in
operational costs, strongly establishing the need for their timely prediction.

In combination with the growth of undesired microbial organisms, open pond systems are susceptible to
unpredictable changes in the weather [8], as are other site-specific environmental factors, such as sand
particles or dust, which change the composition of the culture media [9]. All these factors, which are
directly related to the immediate exposure of the cultures to their biotic and abiotic surroundings, can
lead to unpredictable outcomes in terms of biomass productivity [10]. In addition, the growth and
productivity are not always aligned with the observations made at smaller scales [11]. Understanding the
effects that all these variables have on culture outcomes is critical to optimally predict growth in industrial
settings. Consequently, optimal prediction can guide the design of resilient culture systems, ultimately
reducing the cost of each operation. Identification of exposure- and threat-agnostic biomarkers of stress
as expressed by the microalgae themselves would allow for detailed monitoring and remediation of open
cultures in real-time and understanding what molecular pathways might be involved in resilience.

In the current industrial landscape, enhancing the resilience (i.e., sustained productivity and adaptability
in the face of potential stressors) within expansive open pond systems and optimizing our predictive tools
are essential for maintaining the stability of algal production systems. By utilizing the algal cells as sensors
of their environment, we identified key genetic and molecular pathways that can be engineered to bolster
their resilience. Here, we performed an in-depth multi-omics analysis (transcriptomics, proteomics,
metabolomics and phenomics) in order to further understand the molecular dynamics behind the
acclimation, growth, and response to infection by a parasitoid amoeboaphelid species (PL3) that are
known to cause pond crashes in two 1000 L open raceway ponds under environmentally-simulated
conditions. For this study, we used the freshwater green microalga *Monoraphidium minutum* 26B-AM, a
species currently studied as a promising winter species for biofuel feedstock because of its elevated lipid
yield [12] [13] [14] [15] [16]. The genome of *M. minutum* has also been recently annotated [17]. Samples
were collected for all the different analyses during a 5-day pond run, followed by an infection with the
parasitoid amoeboaphelid species FDO1 [18] [19], which is known to cause pond crashes in real-life
scenarios. In addition to the samples used for the molecular observations, daily measurements of internal
conditions, as well as certain phenotypical outcomes, were also monitored to obtain a holistic view of the
dynamics between environmental signaling, genetic regulation and phenotypical outcomes in a
nonextensively studied species, such as *M. minutum*. By leveraging this rich dataset, we aimed to perform
in-depth molecular observations on the dynamic resilience of algal cultures over time, adding to the
ongoing efforts to increase stability and productivity in large-scale applications. This work reveals a
number of molecular signatures and regulatory elements of interest, and lays the groundwork for
proactive interventions that can sustain algal health and performance in the face of environmental
challenges. In this study, in addition to phenotypical observations, molecular markers evaluated via multi-
omics are indicative of distinct stages of open pond culture. We verified this method by demonstrating a
known link between light and redox signaling and photosynthesis and translation-related genes in *M.*

minutum, in accordance with similar observations in closely related green algal species, confirming the
applicability of the markers identified here across algal culture systems. Finally, we focused on the
molecular pathways associated with the scale-up acclimation and the introduction of the pathogen into
the system through integrated multi-omic analysis. While specific molecular networks of genes, proteins
and metabolites were identified for each transition, we discovered unique signatures in response to the
two stressors, revealing distinct signaling outcomes across stressors.

2. Methods

2.1 Strain and conditions

The *M. minutum* 26B-AM strain was initially collected and isolated as described previously [20], and it has
been utilized in our laboratory as part of the DISCOVR strain library. Seed cultures of *M. minutum* (100
ml) were maintained in sterile 250 ml Erlenmeyer flasks, containing DISCOVR medium [20] under a 16:8 h
light/dark cycle with continuous shaking at 110 rpm in a closed incubator with LED lighting. The light
intensity during the light cycle was 100 µmol photons m^{-2} s^{-1} and the temperature was set between a
maximum of 30 °C during the light period and a minimum of 23 °C during the dark period. The seed
cultures were diluted regularly to maintain exponential phase growth.

2.2 Scale-up

The culture scale-up is graphically summarized in Fig.1. The seed cultures of *M. minutum* described above
were used as inocula for 500 ml cultures grown in 1 L Erlenmeyer flasks. Progressively, the cultures were
transitioned to 2 L bottles, 8 L carboys with stirring and air bubbling and 20 L carboys with bubbling. Each
step lasted 10–15 days, until the inoculum was dense enough (approximately 2 million cells/ml) to be
diluted to the next scale. Up to the 2 L scale, the cultures were grown under the same conditions as the
initial seed cultures detailed in the previous section. For the 20 L carboys, the cultures were acclimated to
relatively high light intensities (1500 µmol photons m^{-2} s^{-1}). The 20 L carboys were thermally acclimated
between 22-28 °C under a 12/12 h dark/light cycle in a water bath, to match the conditions of the next
scale.

A

500ml

100ml

2L

8L

20L

Scale-up

1000L

B

Fig. 1: M.minutum scale-up. *A) Graphic representation of the scale-up. B) Picture of the 1000L open raceway pond on Day 0 of*
the culture run, after the inoculation of the media with the precultures.

2.3 Cultivation and sample collection at the 1000 L opened raceway ponds

The cultures from the 20 L carboys (3 from each pond) were used as inocula for the 1000 L open raceway
ponds (manuscript in preparation). The ponds were filled with tap water and the nutrients and salts were
added via the modified 10% BG-11 recipe, as described previously [20]. A total of 60 L of the inoculum
was added to each pond. The ponds were operated under a 12 h/12 h dark/light cycle, implemented with
LED light panels, placed over the ponds, reaching 1500 µmol photons m^{-2} s^{-1} during the light period, under
a sinusoidal light intensity program (measured in closest proximity to the source). The temperature was
maintained within a temperature range of 22-28 °C, with the highest temperature being reached in the
middle of the light period and the lowest in the middle of the dark period. Temperature control was
achieved through a heat-exchanger lining on the walls of the ponds. The cultures were constantly mixed
with a mechanical paddlewheel at 20 rpm. The pH was held constant at 7.2 via CO_2 sparging. The cells
were grown under these conditions for 5 days and samples for the different omic analyses were collected

on a daily basis one hour after the lights were turned on. The phenotypic and conditional measurements
were taken directly after that.

On the 5th day of the cultivation, 2/3 of each pond was drained and fresh water with new nutrients was
added. In parallel, an inoculum of the parasitoid amoeboaphelid species FDO1 [19] [18] was added at a
0.05% volume concentration, in order to study the dynamics of a pond crash caused by this strain. This
specific strain has been isolated by the Arizona Center for Algae Technology and Innovation as the cause
of *Monoraphdium minutum* pond crashes, and it has not yet been fully characterized. Under laboratory
conditions, it caused *M. minutum* cultures to crash in 4-5 days. The determination of culture crashes was
visual, when the ponds had completely transitioned from green to brown.

2.4 Phenotypic and conditional measurements

All phenotypic and conditional measurements were taken on a daily basis in technical triplicates unless
otherwise stated. The cell count and cell size were determined with the Countess™ 3 FL Automated Cell
Counter (Thermo Fisher Scientific, Waltham, MA, USA) on a daily basis, from a sample collected from the
ponds one hour before the lights were turned on.

To measure chlorophyll fluorescence, 200 µl pond samples (in technical triplicates) in 96-well transparent
plates were measured at an excitation wavelength of 450 nm and an emission wavelength of 685 nm with
a Spark® Multimode Microplate Reader (TECAN, Männedorf, Switzerland). The chlorophyll content of the
ponds is presented in relative units.

For the ash-free dry weight (AFDW) measurements, the steps described in [21] were followed.

The total carbohydrate concentration was determined via the Total Carbohydrate Quantification Assay
(Abcam, Cambridge, UK), following the manufacturer's instructions. Samples in technical triplicates were
collected from each pond on a daily basis, alongside the samples for the omic analyses.

Samples were collected in triplicate on a daily basis from each pond for the quantification of total H3
histones. This measurement was performed with the Histone H3 Total Quantification Kit (Abcam,
Cambridge, UK), following the manufacturer's protocol, and the colorimetric measurements were
conducted using the Spark® Multimode Microplate Reader (TECAN, Männedorf, Switzerland).

Samples for the determination of caspase activity were also collected in triplicate on a daily basis. For the
estimation of cell lysis and caspase activity, the EnzChek Caspase-3 Assay Kit #1 (Thermo Fisher Scientific,
Waltham, MA, USA) was used, with Z-DEVD-AMC as the substrate, according to the manufacturer's
protocol. Dissolved oxygen (%DO), pH, conductivity, temperature and light intensity were measured in
each pond daily, immediately after sample collection. For each variable, we took the average of 9 collected
data points. The %DO and temperature were measured with the Waterproof Dissolved Oxygen Meter Pen
(Sper Scientific Direct, Scottsdale, AZ, USA), the pH and conductivity were measured with an Ohaus Starter
AB33M1-F pH & Conductivity Bench Meter (Parsippany, NJ, USA), and the light intensity was measured
with an LI-1500 Light Sensor Logger (Lincoln, NE, USA).

2.5 Transcriptomics/ RNA extraction and cDNA library preparation

The samples collected for the transcriptomic analyses were centrifuged after collection at 5000 rpm for 2
minutes at 4 °C, and the resulting cell pellets were stored at -20 °C until RNA extraction was performed.

To initiate cell lysis, the frozen pellets were thawed on ice and resuspended in 750 μL of Qiagen RLT Buffer
supplemented with 1% β-mercaptoethanol from the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany).
The resuspended cells were then transferred to 2 mL tubes containing Lysing Matrix Y zirconium oxide
beads (MP Biomedicals, Santa Ana, CA, USA). Following a 10-second vortex, the samples were flash-frozen
in liquid nitrogen. After thawing on ice, the cells were lysed via a FastPrep-24 bead beater (MP
Biomedicals, Santa Ana, CA, USA) for 25 seconds at 40 Hz. The samples were subsequently centrifuged at
9000 rpm for 10 minutes at 4 °C, after which the clear supernatant was collected for RNA extraction. RNA
was extracted via the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) according to the manufacturer's
protocol. The quantity of RNA was assessed via a NanoDrop™ One (Thermo Scientific, Waltham, MA, USA),
whereas its integrity was evaluated via the 4200 TapeStation System (Agilent Technologies, Santa Clara,
CA, USA). RNA samples with a RIN value of 6.5 or higher were deemed to have acceptable integrity and
were subsequently processed for the cDNA library construction.

For the library preparation for RNA sequencing, the KAPA RNA HyperPrep Kit (Roche, Basel, Switzerland)
was utilized, following the manufacturer's instructions. cDNAs were ligated with Unique Dual-Indexed
Adapters from the KAPA Kit (Roche, Basel, Switzerland). The concentration of each library was determined
via a Qubit fluorometer (Thermo Fisher Scientific, Waltham, MA, USA), and quality was assessed via the
4200 TapeStation System (Agilent Technologies, Santa Clara, CA, USA). The libraries were pooled in equal
quantities, and the pooled samples were sent for sequencing at Azenta Genewiz, South Plainfield, NJ, USA.
The sequencing was performed on an Illumina NextSeq 500/550 (Illumina, Inc., San Diego, USA) using a
75-cycle run for paired-end reads.

Genes were filtered for at least 75% of the values to be above zero, run through principal component
analysis (PCA, with 10 components), and a standard scalar was applied to the PCs.

2.6 Proteomics

For the proteomic analysis (protein extraction, sample preparation, mass spectrometry and early data
processing), the pond-collected samples were sent to MS Bioworks, Ann Arbor, MI, USA.

For the sample preparation, the cells were washed twice with 1X PBS and lysed in a modified RIPA buffer
containing 50 mM Tris HCl (pH 8.0), 150 mM NaCl, 2.0% SDS, 0.1% TX100, and 1X Roche Complete Protease
Inhibitor (Roche, Basel, Switzerland). Lysis was performed using 1.4 mm stainless steel beads in a Next
Advance Bullet Blender (Next Advance, Troy, NY, USA) for two cycles of 3 minutes each. The samples were
then heated to 60 °C for 30 minutes and clarified by centrifugation at 13,200 rpm. Protein was precipitated
via trichloroacetic acid (TCA) overnight at -20 °C, after which the pellet was washed and resuspended in a
solution of 8 M urea, 50 mM Tris HCl (pH 8.0), and 1X Roche Complete Protease Inhibitor (Roche, Basel,
Switzerland). Protein quantification was performed via a Qubit fluorometer (Thermo Fisher Scientific,

Waltham, MA, USA), and a pooled sample was created from all the individual samples. For digestion, 2.5
µg of protein was treated overnight with trypsin. The samples were first reduced at room temperature
for 1 hour in 12 mM DTT, followed by alkylation for 1 hour at room temperature in 15 mM iodoacetamide.
Trypsin was added at an enzyme-to-substrate ratio of 1:20. Each sample was then acidified to 0.3% TFA
and subjected to solid-phase extraction (SPE) using a Waters µHLB (Waters, Pleasenton, CA, USA). The SPE
process involved activating the matrix with four additions of 500 µL of 70% acetonitrile, equilibrating with
four additions of 500 µL of 0.3% TFA, adding the sample, washing the wells with three additions of 500 µL
of 0.3% TFA, and eluting in 200 µL followed by an additional 400 µL of 60% acetonitrile in 0.3% TFA. The
samples were then frozen at -80 °C and lyophilized overnight.

For the mass spectrometry analysis, 500 ng of each sample was analyzed in triplicate via an LC-MS/MS
setup consisting of a Thermo Fisher Vanquish Neo UPLC system (Thermo Fisher Scientific, Waltham, MA,
USA) interfaced with a Thermo Fisher Orbitrap Astral (Thermo Fisher Scientific, Waltham, MA, USA).
Peptides were loaded onto a trapping column and eluted through a 75 µm analytical column at a flow rate
of 350 nL/min, with the column temperature maintained at 55 °C. A 30-minute gradient was used for
elution. The mass spectrometer was operated in data-independent acquisition (DIA) mode, capturing full
scan MS data at a resolution of 240,000 FWHM over the m/z range of 380-980, followed by 300 sequential
2 m/z precursor isolation windows, with product ions acquired at a resolution of 40,000 FWHM. The
maximum ion injection time (IIT) was set to 3.5 ms for DIA, and the normalized collision energy (NCE) was
set to 25. An injection of the pooled sample was included at both the beginning and end of the analysis
batch.

The initial data processing for the DIA results was performed via DIA-NN (v1.9.1) [22] , which facilitated
several key functions: conversion of RAW files to the QUANT format, alignment on the basis of retention
times, searching data via an in-silico spectral library generated from the FASTA file and iteratively from
the spectral library derived from RAW data, filtering of database search results at a 1% false discovery rate
(FDR) for both peptides and proteins, calculation of peak areas for detected peptides, and normalization
of the data. The data were subjected to percentile normalization for the PCA plots.

2.7 Metabolomics

For the metabolomics analysis, 1 ml from the samples taken from the ponds (containing both the media
and cells) was sent to Creative Proteomics, Shirley, NY, USA, for untargeted metabolomics analysis. For
the blank, samples with only media were taken from the ponds before the algal inoculation. For the
sample preparation, the samples were transferred into a tube for lyophilization. Following transfer, 500
µL of 80% methanol was added, and the mixture was vortexed for 30 seconds and then subjected to
sonication in an ice-water bath for 30 minutes. The samples were subsequently stored at -20 °C for 1 hour,
after which they were vortexed for an additional 30 seconds and centrifuged at 12,000 rpm for 10 minutes
at 4 °C, to lyse and remove the cells. From the resulting supernatant, 200 µL was transferred into a vial,
and 5 µL of a 0.14 mg/mL solution of DL-o-chlorophenylalanine was added. The final solution was filtered
through a 0.22 µm filter in preparation for LC-MS analysis. The separation of the compounds was achieved
via the Vanquish Flex UPLC system in conjunction with a Q Exactive Plus mass spectrometer (Thermo
Scientific, Waltham, MA, USA), which featured a heated electrospray ionization (ESI) source. The LC

separation utilized a Hypersil Gold C18 column (100 × 2.0 mm, 1.9 μm), with the mobile phase consisting
of solvent A (0.05% formic acid in water) and solvent B (acetonitrile) under a gradient elution profile: from
0 to 1 minute at 5% B, from 1 to 12.5 minutes increasing from 5% to 95% B, holding at 95% B from 12.5 to
13.5 minutes, returning to 95% B from 13.5 to 13.6 minutes, and finally returning to 5% B from 13.6 to 16
minutes. The flow rate was maintained at 0.3 mL/min, with the column temperature set at 40 °C and the
sample manager temperature at 4 °C. The mass spectrometry parameters for both ESI positive (ESI+) and
negative (ESI-) modes were configured as follows: for ESI+, the heater temperature was set to 300 °C, the
sheath gas flow rate was 45 arb, the auxiliary gas flow rate was 15 arb, the sweep gas flow rate was 1 arb,
the spray voltage was 3.0 kV, the capillary temperature was 350 °C, and the S-Lens RF level was 30%. For
ESI-, the heater temperature remained at 300 °C, with the same sheath and auxiliary gas flow rates as
those used for ESI+, but the spray voltage was adjusted to 3.2 kV, and the S-Lens RF level was increased
to 60%.

For the bioinformatic analysis (see section 2.8), we considered the metabolites whose abundance was
more than 10x greater than the abundance of the metabolites in the blank to be significant.

2.8 Bioinformatics

Principal component analysis (PCA) was performed in R with the factoextra package [23]. For 3D PCA
analysis, the first 3 components were plotted for each replicate per day, and the centroid point was
also plotted on the same axes. Day-specific standard deviation from the centroid (based on
replicates) is shown by a shaded sphere.

For RNA sequencing analysis, the following sequence processing steps and associated packages were
used: the fastq file was quality-checked with FastQC, and adapter trimming was performed with
Trimmomatic [24], t

The pseudoalignment of the transcripts to the transcriptome of *M. minutum* (as it was published and
publicly available by [17]) and the quantification of their abundance were performed with the Kallisto
package [25]. The integration of the Kallisto data and the differential gene expression analysis, including
the time course analysis, was performed with Sleuth [26]. The GEMMA package [27] was used for the
correlation of the gene expression with specified quantitative phenotypes.

Additional tools for the multi-omic analyses include ggplot2 [28] for graphic representation and pheatmap
[29] for the heatmap.

De novo motif enrichment analysis of the promoter regions (200 bps upstream of ATG) was performed
with the Homer Motif Discovery and Analysis software [30].

We used the Data Integration Analysis for Biomarker Discovery using Latent variable approaches
for Omics studies (DIABLO) framework from the R software package mixOmics (version 6.28.0) to combine
the transcriptomics, proteomics, and metabolomics datasets [31]. Specifically, we compared these data
across Day 0 and Day 1 and subsequently Day 4 and Day 5 to identify key features of culture scale up and
infection, respectively. The metabolomics and proteomics datasets were log transformed prior to this
analysis, and the transcriptomics data were normalized to transcripts per million (TPM) values. The

proteomic and transcriptomic datasets were further filtered via the nearZeroVar function from the caret
R package (version 7.0- 1), with a unique cutoff of 20 for the proteomics dataset and selecting transcripts
with variances of greater than or equal to 200 were selected. In the DIABLO design matrix, we prioritized
correlations between the three omics datasets by choosing a 0.6 weighted model. The features from
latent component 1, from the Day 0 and Day 1 comparison and from the Day 4 and Day 5 comparison
were then compiled for comparison across all four days via the R package pheatmap (version). The values
were scaled and centered prior to plotting, and both rows and columns were clustered via Euclidean
distance.

3. Results

3.1 Phenotypic and conditional measurements reveal distinct acclimation, growth and infection dynamics
in algae in open pond systems

We performed a longitudinal analysis of algal growth in two 1000 L open raceway ponds, and introduced
an amoeboaphelid pathogen on day 4 of the experiment, tracking a total of 12 variables. Phenotypic
measurements included cell count, AFDW, chlorophyll concentration, total carbohydrate content, total
H3 histone concentration, cell size, and caspase activity (Additional File 1). In addition, we measured five
environmental variables within the ponds, including %DO, pH, conductivity, temperature, and light
intensity.

Across the time-course, we observed 3 growth phases of interest (color-coded in Fig. 2A):

a) The acclimation or lag phase (cyan). This phase occurred immediately after the inoculation in
the open system with refreshed nutrients. This period lasted 2 days, and growth during this phase
was slower than that during the other phases.

b). The log phase (yellow). The cells were fully acclimated to the open-pond conditions and grown
exponentially. This phase lasted for 4 days before the dilution and the infection and 3 days after
dilution.

c) Crash phase (brown). This phase is caused by algal death, which leads to the complete overtake
of the ponds by the pathogen and other opportunistic species.

The ponds behave largely similarly, although we do observe a divergence of phenotypes at later stages of
infection, as documented in Fig. 2.

To understand whether stress conditions trigger programmed cell death, we measured caspase activity
throughout the experiment. Caspases are proteases that catalyze the breakdown of proteins during
programmed cell death [32]. We observed that caspase activity initially increased after pond inoculation
and then gradually decreased as the cells acclimated to their new environment (Additional File. 2).
Notably, we did not observe an increase in caspase activity after infection with amoeboaphelid, suggesting

that the reduction in the algal cell count during the postinfection phase is caused by caspase-independent
cell death [33] [34].

Fig. 2: Growth tracking of the M. minutum pond cultures. *A) The cell count between the three ponds. B) Chlorophyll concentration*
in relative units (RU) from florescence measurements. C) The Ash-free dry weight (AFDW) of algae cells in mg per ml of culture.

For an integrated view of the phenotypic and conditional profile of the longitudinal experiment, principal
component analysis (PCA) was performed with the 12 phenotypic and environmental variables.
Dimension 1 represented 49.4% of the explained variance (Fig. 3A) and separated the variables that are
indicative of growth, from those indicative of stress. This provides insight into the competing pressures
that the algae face during cultivation, and the contributions of these pressures to outcomes.

We observe that the three distinct phases (acclimation, log, and crash) are resolved away from each other
in the PCA biplot shown in Figure 3B, and we can discern a trajectory in PC space as the longitudinal
experiment progresses, which is shown by connecting temporally adjacent points.   While the largest
adjacent transition occurred during scale-up (Day 0 to Day 1), we also observed a significant shift upon
infection into a distinct PC space highlighted in brown in Figure 3B. This visualization further emphasizes

the distinction between the conditions and outcomes during these phases and suggests the presence of
distinct molecular profiles within the algae that likely precede outcomes. To this end, we performed
transcriptomic, proteomic and metabolomic analyses across the first 6 days of this experiment.

Fig. 3: PCA for 12 variables as measured at the two ponds. *A) PCA for the 12 variables used. The color of each line represents*
the contribution of each variable to the principal axes. Created with factoextra in R. B) Each day of the cultivation, plotted as
individuals. The arrows are indicating the trajectory of each day during the culture run.

3.2 Transcriptomic stability tracks with optimal growth and culture resiliency

With the goal of identifying molecular patterns within the algae that precede the observed phenotypic
outcomes, we performed transcriptomic analysis from Day 0 to Day 6 (Additional File 2). The samples thus

represented the acclimation period (day 0 and day 1), the preinfection log phase (day 2- day 4) and the
early infection days (days 5 and 6).

A considerable percentage of transcripts (12.3%) were significantly differentially expressed from day 0
(the inoculation day) to day 1, reflecting the transcriptional acclimation to the scale transition from the
20 L carboys to the open 1000 L ponds (Fig.4A). This difference in expression was multiple times greater
than any daily difference observed during the following days. This included days after infection where
only 0.7% of the transcripts were significantly differentially expressed.

We performed PCA on the transcriptomic data to determine whether the longitudinal trajectory was
comparable to that of our phenotypic and environmental variables (Fig. 4B and 4C). The transition from
Day 0 to Day 1 was the most striking, reflecting the rapid and large transcriptomic response to the 1000 L
pond conditions. The distributions of the days corresponding to the pre-infection log phase, on the other
hand, are closer in PC space, reflecting transcriptional stability during that growing stage. A clear
distinction between the distribution of the preinfection and postinfection days was also observed, as a
result of both the culture dilution and the introduction of the pathogen. In fact, unlike in Fig. 3B where
the measured environmental and phenotypic variables postinfection were in a distinct cluster away from
the scale-up, the early postinfection transcriptomic profile trends back toward the acclimation phase
profile but still remains distinct from them.

Fig. 4: Comparative analysis of the transcript abundances overlaid with cell growth results. *A) Growth curve of both ponds over the time course, with transcriptomic data collected on the same day as growth was measured (as shown by points on the curve). Percentages of differentially expressed genes between specific daily points within the cultivation are shown by the numbers. B) 3D PCA analysis of the transcriptomics, showing the distribution of all the technical replicates for each day from the two ponds. Centroids are indicated with a darker point, and standard deviation is indicated by the shaded circle. C) 2D representation of the PCA showing the average points for each day. The arrows indicate the trajectory of the time course.*

The annotation of *M. minutum* is incomplete, making traditional pathway analysis complicated. To further investigate the regulation behind those transcriptional changes we performed a motif analysis among the promoter regions of the genes that were significantly differentially expressed to identify regions that could be bound by conserved transcription factors. We focused on the transitions between day 0 and day 1 (Table 1), and between day 4 and day 5 (Table 2). The ten most enriched motifs that emerged from each group are shown in Table 1 and Table 2, alongside their best matches from other organisms. Most identified motifs are unique to each transition.

Among the acclimation phase gene promoters, we also observed two motifs (labeled with HOMER, such as ZNF549 and ZNF415) that are likely bound by zinc finger proteins. In addition, 40% of the motifs

correspond to a promoter motif from the diatom *Thalassiosira pseudonana*. A notably high percentage
(66.66%) of the promoter regions of the genes that were significantly differentially expressed after the
infection contained a motif that can be recognized by Auxin Response Factor 1 (ARF1), a transcription
factor identified in *Arabidopsis thaliana.*

Table 1: Enriched promoter motifs at the scale adaptation-related genes.

	Motif	% of Targets	P- value	Best match/ Details
1		43.38%	1e-12	PB0188.1_Tcf7l2_2/Jaspar (0.841)
2		40.40%	1e-9	Tp_0225(RRM)/Thalassiosira_pseudonana-RNCMPT00225-PBM/HughesRNA (0.730)
3		25.17%	1e-9	RTG3/Literature (Harbison)/Yeast(0.896)
4		32.78%	1e-8	ABF2/MA0266.2/Jaspar (0.781)
5		41.72%	1e-8	ZNF549/MA1728.2/Jaspar (0.753)
6		27.48%	1e-8	STB4/STB4_YPD/[] (Harbison)/Yeast(0.723)
7		34.44%	1e-8	Syncrip(RRM)/Xenopus_tropicalis-RNCMPT00281-PBM/HughesRNA(0.754)
8		21.85%	1e-7	ZBED4/MA2328.1/Jaspar (0.742)
9		18.54%	1e-7	RAP211(AP2EREBP)/colamp-RAP211-DAP-Seq (GSE60143)/Homer (0.722)
10		34.44%	1e-7	ZNF415(Zf)/HEK293-ZNF415.GFP-ChIP-Seq (GSE58341)/Homer (0.721)

Table 2: Enriched promoter motifs at the infection-related genes.

	Motif	% of Targets	P- value	Best match/ Details
1		66.67%	1e-9	ARF1(ABI3/VP1)/Arabidopsis thaliana/AthaMap (0.719)
2		64.41%	1e-9	gcm2/MA0917.1/Jaspar (0.770)
3		35.59%	1e-8	Abd-B/dmmpmm(Bergman)/fly(0.775)

4		32.33%	1e-8	RTG3/Literature (Harbison)/Yeast (0.882)
5		50.38%	1e-8	RGT1/Literature (Harbison)/Yeast (0.775)
6		28.57%	1e-8	Sp1(Zf)/Promoter/Homer (0.747)
7		40.85%	1e-7	bHLH78/MA0963.1/Jaspar (0.804)
8		26.07%	1e-7	HES2/MA0616.3/Jaspar (0.840)
9		23.06%	1e-7	ZmHOX2a (2) (HD-HOX)/Zea mays/AthaMap (0.795)
10		41.10%	1e-7	PBX1/MA0070.2/Jaspar (0.812)

3.3 Transcriptomic and phenotypic data reveal the light and dissolved oxygen (DO) regulation of protein translation and photosynthesis

One of the advantages of performing time-course transcriptomics on large open cultures in a poorly studied species such as *M. minutum* is that it allows for the extraction of reliable and useful information about specific environmentally triggered expressed genes. Light intensity and DO are two environmental factors known to influence outcomes in algae. To understand their role in the outcome of our cultures, we used a program called GEMMA that applies mixed linear models to connect phenotypes with large-scale molecular data [27]. We identified transcripts whose expression was positively or negatively correlated with light intensity and %DO across samples (Figs. 5 and 6).

The observation of the top up- and downregulated genes correlated with light intensity (Tables 3 and 4), revealed a strong correlation between high light intensity and the expression of genes related to structural components of ribosomes and other translation-related proteins. In addition, we noted a structural component of Photosystem I among the genes whose expression was downregulated in response to light, which is consistent with previous literature showing that photosynthetic organisms downregulate photosystem components in response to light stress [35].

To further understand the regulatory genomic elements that might contribute to light-dependent changes in gene expression, we performed a motif analysis on the promoter regions of all the genes whose expression was significantly up- or downregulated by light (Tables 5 and 6). Among the most significant motifs from the upregulated genes, two corresponded to transcription factors from two higher plant species, *Arabidopsis* and *Zea mays*.

Table 3: The annotations of the top 5 genes whose expression was upregulated with light.

JGI ID	beta	Gene product
427491	486.8095	ribosomal protein S9 component of cytosolic 80S ribosome and 40S small subunit
412255	479.0458	40S ribosomal protein S29-like
420537	442.9904	putative 40S ribosomal protein S3a
431691	419.257	Translation protein SH3-like domain-containing protein
420483	377.4611	component of cytosolic 80S ribosome and 60S large subunit

457 *Table 4: The annotations of the top 5 genes whose expression was downregulated with light.*

JGI ID	beta	Gene product
518027	-257.7188	heat shock cognate 70 kDa protein 4-like
474353	-195.5336	hypothetical protein
453947	-167.1721	eukaryotic translation initiation factor 5
437942	-161.7026	photosystem I protein P30
272372	-155.3597	hypothetical protein

459 *Table 5: Enriched promoter motifs at the light upregulated genes.*

	Motif	% of Targets	P- value	Best match/ Details
1		21.74%	1e-8	Unknown3/Arabidopsis-Promoters/Homer (0.710)
2		33.33%	1e-8	CAT8/MA0280.2/Jaspar (0.749)

	Motif	% of Targets	P-value	Best match/ Details
3		23.91%	1e-7	ZmHOX2a (2) (HD-HOX)/Zea mays/AthaMap (0.823)
4		23.91%	1e-7	AT5G04390/MA1736.2/Jaspar (0.752)
5		23.91%	1e-7	lsy-2/MA2181.1/Jaspar (0.874)
6		47.46%	1e-7	Sp1(Zf)/Promoter/Homer (0.783)
7		30.80%	1e-6	RGT1/Literature (Harbison)/Yeast (0.793)
8		30.43%	1e-6	insv/MA2315.1/Jaspar (0.821)
9		16.67%	1e-6	ZNF549/MA1728.2/Jaspar (0.779)
10		30.80%	1e-5	ZNF610/MA1713.2/Jaspar (0.721)

462 *Table 6: Enriched promoter motifs at the light downregulated genes.*

	Motif	% of Targets	P- value	Best match/ Details
1		36.93%	1e-10	br/MA0011.2/Jaspar (0.725)
2		41.83%	1e-9	PB0188.1_Tcf7l2_2/Jaspar (0.842)
3		32.35%	1e-8	ZNF331/MA1726.2/Jaspar (0.870)
4		23.53%	1e-8	MSN4(MacIsaac)/Yeast (0.777)
5		27.12%	1e-7	MET31(MacIsaac)/Yeast (0.683)

6		26.47%	1e-7	PHD1(MacIsaac)/Yeast (0.856)
7		17.97%	1e-6	PUF68(RRM)/Drosophila_melanogaster-RNCMPT00141-PBM/HughesRNA (0.805)
8		27.45%	1e-6	ZmHOX2a(2)(HD-HOX)/Zea mays/AthaMap (0.799)
9		17.32%	1e-6	WRKY1/MA0589.2/Jaspar (0.869)
10		41.50%	1e-6	TEAD2/MA1121.2/Jaspar (0.808)

A similar GEMMA analysis was repeated with the genes most correlated with %DO. Among the ten most upregulated and downregulated genes correlated with %DO (Fig. 6A, Tables 7 and 8), multiple genes related to photosynthesis and carbon fixation were identified. A phosphoribulokinase/uridine kinase, an enzyme responsible for a key step in the Calvin-Benson cycle, a light-harvesting protein of Photosystem I, and a protein involved in programmed cell death were among the most upregulated genes correlated with DO. We performed motif analysis on the promoter regions of all the genes correlated with DO (Tables 9 and 10). We observed enrichment of the binding site for the same transcription factor from the diatom *Thalassiosira pseudonana* at both the up- and downregulated genes from the DO.

% DO-Associated Transcripts

Fig. 6: GEMMA analysis, revealing genes correlated with %DO. *The volcano plot shows the significance and the extent of the*
correlation between the light intensity and the transcript abundance. The top 5 up- and downregulated genes by %DO are indicated.

Table 7: The annotations of the top 5 genes whose expression was upregulated with %DO.

JGI ID	beta	Gene product
413487	7.216919	Phosphoribulokinase / Uridine kinase family-domain containing protein
475645	6.064846	hypothetical protein
399114	5.610648	expressed protein
419767	4.78923	light-harvesting chlorophyll-a/b protein of photosystem I

374147	4.481988	Programmed cell death protein 2, C-terminal putative domain-domain containing protein

478   *Table 8: The annotations of the top 5 genes whose expression was downregulated with %DO.*

JGI ID	beta	Gene product
427491	-21.50261	ribosomal protein S9 component of cytosolic 80S ribosome and 40S small subunit
420537	-16.57716	putative 40S ribosomal protein S3a
425187	-11.31527	ribosomal protein L3 component of cytosolic 80S ribosome and 60S large subunit
420483	-9.651194	component of cytosolic 80S ribosome and 60S large subunit
439234	-9.335318	P-loop containing nucleoside triphosphate hydrolase protein

480   *Table 9: Enriched promoter motifs at the %DO upregulated genes.*

	Motif	% of Targets	P- value	Best match/ Details
1		23.02%	1e-8	GATA15(C2C2gata)/col-GATA15-DAP-Seq (GSE60143)/Homer (0.833)
2		28.68%	1e-8	ARR1/Literature (Harbison)/Yeast (0.743)
3		41.51%	1e-7	NKX2-3/MA0672.2/Jaspar (0.771)
4		21.89%	1e-6	ABF2/MA0266.2/Jaspar (0.803)
5		37.74%	1e-6	ZNF549/MA1728.2/Jaspar (0.751)
6		38.87%	1e-6	Tp_0225(RRM)/Thalassiosira_pseudonana-RNCMPT00225-PBM/HughesRNA (0.796)
7		57.36%	1e-6	ZNF415(Zf)/HEK293-ZNF415.GFP-ChIP-Seq (GSE58341)/Homer (0.760)

	Motif	% of Targets	P-value	Best match/ Details
8		17.74%	1e-6	Vdr/MA0693.4/Jaspar (0.846)
9		16.98%	1e-6	EFL-1(E2F)/cElegans-L1-EFL1-ChIP-Seq(modEncode)/Homer (0.719)
10		28.30%	1e-6	REB1/SacCer-Promoters/Homer (0.815)

482   *Table 10: Enriched promoter motifs at the %DO downregulated genes.*

	Motif	% of Targets	P- value	Best match/ Details
1		43.38%	1e-12	PB0188.1_Tcf7l2_2/Jaspar (0.841)
2		40.40%	1e-9	Tp_0225(RRM)/Thalassiosira_pseudonana-RNCMPT00225-PBM/HughesRNA (0.730)
3		25.17%	1e-9	RTG3/Literature (Harbison)/Yeast (0.896)
4		32.78%	1e-8	ABF2/MA0266.2/Jaspar (0.781)
5		41.72%	1e-8	ZNF549/MA1728.2/Jaspar (0.753)
6		27.48%	1e-8	STB4/STB4_YPD/ [] (Harbison)/Yeast (0.723)
7		34.44%	1e-8	Syncrip (RRM)/Xenopus_tropicalis-RNCMPT00281-PBM/HughesRNA (0.754)
8		21.85%	1e-7	ZBED4/MA2328.1/Jaspar (0.742)
9		18.54%	1e-7	RAP211 (AP2EREBP)/colamp-RAP211-DAP-Seq (GSE60143)/Homer (0.722)
10		34.44%	1e-7	ZNF415(Zf)/HEK293-ZNF415.GFP-ChIP-Seq (GSE58341)/Homer (0.721)

484   *3.4 Proteome composition shifts during algal growth and stress response.*

To gain additional molecular information from the algae as they sensed transitions in their environment, a comprehensive proteomic analysis was conducted on the two pond populations longitudinally. Similar to the transcriptomics data, PCA of the proteomics data (Additional File 3) revealed a highly dynamic phase during the first two days of the culture (Fig. 7). After entering the log phase, however, the proteome becomes progressively more stable, while at the same time maintaining a distinct profile between the pre- and postinfection periods. This was in contrast to the more dynamic changes in the transcriptome after infection, highlighted in the plasticity of the transcriptional regulation in relation to the environmental triggers in comparison with the stability of the protein expression. Further analysis of the correlations between specific transcripts and proteins will be discussed later in this article.

Fig. 7: PCA of the proteomics. *A) 3D PCA analysis of the proteomics, showing the distribution of all the technical replicates for each day from the two ponds. Centroids are indicated with a darker point, and standard deviation is indicated by the shaded circle. B) 2D representation of the PCA showing the average points for each day. The arrows indicate the trajectory of the time course.*

3.5 Metabolomic analysis reveals potential indicators of the stress response and pathogen infection.

PCA of the metabolites (Sup. Dataset 4) revealed distinct profiles during the three phases of cultivation similar to the transcriptomic and proteomic PCAs (Fig. 8A and 8B). Interestingly, day 3, which is in the middle of the log phase, stood out in the PCA. This contrasted with the observation made from transcriptomics. While log phase was linked with transcriptomic stability, the metabolic profile is very dynamic during the growth phase.

We performed clustering analysis of the longitudinal metabolomics data to further understand the metabolites that covaried (Fig. 8C). We used DPGP [36], a nonparametric clustering algorithm that can factor in longitudinal information, to separate metabolites into clusters on the basis of their patterns in abundance. We found that many metabolites either had their peak or trough on days 3 or 4, the midpoint of the log phase (clusters 2, 4, 6 and 8 and 9).   The most abundant cluster (Cluster 1) included metabolites whose abundance steadily increased over time. On the other hand, Cluster 5 showed the opposite trend, starting with high abundance and slowly decreasing in abundance over time.

Hierarchical clustering of the scaled and centered metabolite abundances in the heatmap, shown in Fig. 8D, highlights the emergence of three groupings of metabolites, and is in relative accordance with the DPGP clustering. The first group consists of the metabolites whose abundance peaks in the lag phase. The second group peaked in the log phase. The last group, which was the largest by number, contained the metabolites that reached their peak abundance after the infection. The last group included hormones and steroids, such as 5-Dihydrotestosterone, 7-Hydroxytestosterone, Estrone, Estriol and 19-Norandrostenedion.

Fig. 8: The metabolomic analysis from the pond cultures. *A) 3D PCA analysis of the metabolomics, showing the distribution of all the technical replicates for each day from the two ponds. Centroids are indicated with a darker point, and standard deviation is indicated by the shaded circle. B) 2D representation of the PCA showing the average points for each day. The arrows indicate the trajectory of the time course. C) Clustering analysis of the abundance of the metabolites throughout the studied period, via DPGP. D) Heatmap with the hierarchical clustering of the significantly produced metabolites throughout the run of the cultivation. It includes the metabolites which had more that 10x higher signal than the metabolites in the blank.*

3.6 Integrated multi-omic analysis revealed distinct molecular features related to scale transition and infection defense

To assess the global relationships among proteins, metabolites and transcripts across our longitudinal experiment, we performed an integrative analysis via the DIABLO framework in the mixOmics R package [31]. We performed this analysis using all omics data from days 0 and 1 to explore the molecular features of the scale-up stress, and separately used days 4 and 5 data to explore that of the infection stress. While we found that there was no overlap in the key proteins or transcripts output by the model between the

scale up and infection stress, here were two metabolites, guanine and 12-oxo-Phytodienoic acid, were
common across the two stressors, suggesting more convergence at the metabolite level than at the
molecular pathway level.

We took the molecular features from the first latent component of both analyses and combined them
into one heatmap to comprehensively visualize these results (Fig. 9). As expected, the columns clustered
by day, and we observed that the two transitions had almost entirely unique patterns. We observed a
cluster of transcripts (all significantly differentially expressed between days 0 and 1), proteins, and
myristic acid elevated on day 0 relative to the other days. Similarly, we observed another distinct cluster
of transcripts and proteins, as well as 1,5-hexadiene and 2,3-Diaminosalicylic acid, which were elevated
on Day 4 compared with the other days. However, a decrease in a different cluster of proteins, as well as
maltose and hypogeic acid, distinguished day 1 relative to the other days. These results suggest distinct
molecular and metabolic profiles in response to different stressors, identifying potential unique
biomarkers and target pathways for detection and intervention.

Fig. 9: Indicative signatures of scale adaptation and infection. *Heatmap containing the molecular features (genes, proteins and*
metabolites) which change significantly their abundance during the transition from day 0 to day 1 and/or from day 4 to day 5.
Their abundance is being indicated by a scaled blue-to-red scheme. Each of the 6 biological replicates for each condition are
indicated separately.

4. Discussion

4.1 Transcriptomics and metabolomics reveal the commonalities and differences between the molecular mechanisms governing scale-up stress and infection response

We performed a multi-omic analysis of *M. minutum* 26B-AM cultivated over 12 days in 1000 L open raceway ponds to explore the molecular signatures that manifest in response to stressors often encountered at a large scale. The three distinct growth phases we observed in the 1000 L ponds were in line with what is typically observed at a large scale, i.e., batch cultivations of algae [37] [38]. Culture scale-up is often evidenced by a lag phase, as we observed in our cultures, especially when moving from well-controlled laboratory cultures into open raceway ponds [37] [39] [40]. A change in scale represents a significant stressor for the cultures, introducing a combination of different environmental signals, including changes in light, radiation, heat, gas distribution, as well as microbial and environmental particle composition that requires an acclimation period, which is reflected in our growth curves over days 0 and 1.

Similarly, algal pests are common stressors experienced in open raceway ponds and active areas of study. Changes in phenotypic measurements in response to infection or even visual indications of infection stress, typically appear sometime after the initial infection. In our cultures, we introduced the amoebaphelid on day 4 but did not begin to observe a decline in growth, which was emblematic of a culture crash, until day 8, accompanied by a color change in the culture from light green to brown. Pond crashes, which are driven by algal pathogens and often damage crops overnight, account for almost 30% of algal crop losses annually [41], increasing the overall cost of algal applications. Despite the dramatic effects of those infections on ponds, the cellular mechanism of the infection which includes the insertion of the parasite in the host cell [19], allows a certain amount of time of incubation before a noticeable crash [42]. Cellular and molecular alterations are already in place and if they can be detected in a timely manner, the irreversible effects of a pond crash can potentially be avoided, or at least resource allocation decisions can be made much earlier.

Differential expression analysis revealed that culture scale up has a substantial effect on the transcriptome, as evidenced by the much larger portion of significantly differentially expressed transcripts between days 0 and 1 compared with the other days, including those impacted by infection. In addition, infection is not expected to trigger a wide immune response from the algal cells, reflected by transcriptional adaptation. Characterizing these pathways and adjusting conditions to prevent their activation could help acclimate the culture in a less stressful way and potentially make it more resilient to future stressors.

With respect to infection, current strategies for early indication of contamination include visual detection strategies such as flow cytometry [43], in situ microscopy [44], multispectral image analysis [45] and photosynthetic measurements [46] [47]. Those methodologies, although relatively quick, require a certain pathogen titer in the culture. Biomolecular approaches include 16S and 18S sequencing [48], and the detection of Volatile Organic Compounds (VOC), as indirect early indicators of an infection [49]. Here we

show that the algae can also be used as early indicators of infection, as demonstrated by the changes in
transcripts.

Because many of the transcripts in *M. minutum* are poorly annotated, we used motif analysis to further
explore the significantly differentially expressed genes. Most identified motifs were unique to each
transition, potentially indicating two distinct regulatory pathways between the scale adaptation and
pathogen stress. However, common elements between the two datasets exist, implying that there are
factors that could be used to coregulate responses to different stressors (either endogenously or via
synthetic biology). An example of a potential point of convergence are the motifs recognized by the
transcription factors mitochondrial retrograde 1 (RTG1) and 3 (RTG3), both of which were originally
identified in yeast and are involved in mitochondrial metabolic regulation.

Among the promoters of genes differentially expressed during scaling up, we also identified two motifs
labeled by HOMER, ZNF549 and ZNF415, which are likely bound by zinc finger proteins, a family of
transcription factors that have been shown to regulate stress responses in other algal species [50] [51]. In
addition, 40% of the motifs correspond to a promoter motif from the diatom *Thalassiosira pseudonana*
suggesting possible functional importance across diverse algal species. A notably high percentage
(66.66%) of the promoter regions of the genes that were significantly differentially expressed after the
infection contained a motif that can be recognized by Auxin Response Factor 1 (ARF1), a transcription
factor identified in *Arabidopsis thaliana.* ARF1 is responsive to the phytohormone auxin, which is
responsible for a wide range of developmental and stress responses in plants [52] [53]. Given the
evolutionary proximity between higher plants and green algae, it is likely that ARF1 may have similar
functions in algal responses post-infection and might suggest a possible defense mechanism. The
identification of these factors points to potential targets for synthetic biology in *M. minutum* in the future,
to either boost or tamper with specific molecular responses.

A survey of the metabolites released by the algae upon encountering various stressors also provides
valuable insights into predictive markers, pond health indicators, and infection indicators, while also
explaining the natural mechanisms of growth and defense in algae. We used PCA and DPGP to look at
global trends in metabolites over time. While log phase was linked with transcriptomic stability, the
metabolic profile is very dynamic during the growth phase.

These metabolites are highly important, both as early indicators of an upcoming crash, multiple days
before the effects of the infections can be visible, and as guides for the development of prophylactic
strategies for algal cultures. Among those metabolites, we identified 14-HDHA, a hydroxy
docosahexaenoic acid oxylipin which is derived from docosahexaenoic acid (DHA). The production of
oxylipins from algae is linked with oxidative stress [54], suggesting a similar regulatory pathway here for
pathogen defense. In this group, there are also multiple hormones and steroids (5-Dihydrotestosterone,
7-Hydroxytestosterone, Estrone, Estriol and 19-Norandrostenedion). While such compounds originate in
the aquatic environments from sources other than algae, many algal species are known to biotrasform
them or completely remove them from the system [55] [56] [57], with effects that can be either toxic or
growth promoting for the cells [58] [59]. The hypothesis that these compounds play a role in the
molecular signaling or progression of the infection is one that deserves further investigation for many
reasons, including the discovery of early biomarkers for an upcoming pond crash.

4.2 Transcriptomic and phenotypical observations reflect light and gas-regulated responses in M. minutum

Light and gas constitute the two cornerstone environmental factors for aquatic organisms, especially the
photosynthetic organisms such as green microalgae. Indeed, light and CO_2 availability are essential not
only for photosynthesis, the primary metabolic function of microalgae but also for photoprotection, lipid
composition, and carbon concentration [60] [61] [62] [63] [64]. The conserved and well-studied role of
light and CO_2 availability led us to explore which *M. minutum* genes are corelated with these genes, as a
mechanism by which to verify that our method identifies valid molecular signatures.

We observed that translation-related proteins were strongly correlated with high light intensity. Protein
translation is an essential regulatory step for phenotypic acclimation, and its control by light has been
observed in other algae, including the Chlorophyte *Ulva prolifera* [65], the Charophyte *Klebsormidium*
nitens [66], and *Chlorella sp.* [67]. In addition, we noted a structural component of Photosystem I among
the genes whose expression was downregulated in response to light. Photosystem I is a crucial component
of the electron transport chain and is responsible for the light-mediated transfer of electrons across the
thylakoid membrane. Excess light can cause oxidative stress and subsequent photoinhibition in
Photosystem I, and in many photosynthetic organisms, including in *Chlamydomonas reinhardtii*, light can
suppress the expression of structural proteins in Photosystem I to protect the machinery from this type
of damage [68]. Future work to test this hypothesis across algal species will be critical.

Motif analysis of the promoter regions of genes correlated with light highlighted two motifs related to
homeobox proteins in higher plants. The *Zea mays* homeobox 2a (ZmHOX2a) protein, whose expression
in maize is restricted to meristems, is involved in in plant development and growth [69]. In addition, this
motif might more broadly represent homeobox transcription factors, which are known to regulate
environmental responses in plants [70] [71], suggesting evolutionary conservation of light-induced gene
regulation through homeobox transcription factors across the plants and algae. A motif corresponding to
the same transcription factor is also found among the motifs from the promoters of the downregulated
genes by light, highlighting the involvement of these motifs in both the up- and downregulation responses
to light. Interestingly, 30.80% of the targets of the promoter regions from the up-regulated genes
contained a motif corresponding to the transcription factor RGT1 (Restores Glucose Transport) from
Saccharomyces cerevisiae. The protein regulates glucose transport though the expression of glucose
transporters [72], and the connection between light-induced stress and glucose transport has been
documented in other organisms [73] [74]. The motifs of the downregulated genes include one that is
recognized by the *Saccharomyces cerevisiae* stress-related multicopy suppressor of SNF1 mutation
protein 4 (Msn4), which controls the metabolic response under cellular stress conditions [75].

Among the genes correlated with DO, multiple genes are related to photosynthesis and carbon fixation.
A phosphoribulokinase/uridine kinase, an enzyme responsible for a key step in the Calvin-Benson cycle,
and a light-harvesting protein of Photosystem I are among the most upregulated genes. Genes related to
programmed cell death were positively correlated with DO, an observation that aligns with results from
other green algal species linking ROS signaling and programmed cell death [76] [77] [78] [79] [80]. These
observations highlight specific molecular changes experienced by large-scale cultures that are likely
directly related to environmental fluctuations. Our results also revealed that increased DO caused more
gene downregulation than gene upregulation overall, especially for the genes indicated in Table 8. Most
of these genes are involved with structural components of ribosomes, indicating a strong suppressive

relationship between O$_2$ levels and translational regulation, as was also observed in the case of genes
upregulated with light. In fact, the most downregulated gene from the DO was the most upregulated one
from light. Since high light is a stress factor, as is low oxygen, this reciprocal relationship between their
regulated genes aligns with expectations. The list of the top five downregulated genes includes a P-loop
containing nucleoside triphosphate hydrolase, which, in higher plants such as *Arabidopsis,* acts as a
signaling molecule involved in ribosome formation and attenuates ROS formation and oxidative stress
[81] [82].

Motif analysis of the promoter regions of all the genes correlated with DO revealed enrichment of the
binding site for the same transcription factor from the diatom *Thalassiosira pseudonana* that we observed
during scale acclimation. This putative transcription factor motif is found in 40.40% of the promoter
regions of the genes differentially expressed during scale acclimation (Table 1), which is a sign of a strong
contribution of the DO differences to the scale transition stress. Future efforts to identify this transcription
factor that is conserved across algae might be fruitful for understanding stress mechanisms and broad-
scale synthetic biology-based interventions. The second most enriched motif among the promoters of the
upregulated genes corresponds to the transcription factor Arrestin 1 (ARR1), which in higher plants is
responsive to the phytohormones Cytokinins and plays a fundamental role in numerous developmental
stages of the plant [83] .The motif bound by the yeast transcription factor Regulation of the Mitochondrial
Retrograde 3 (RTG3) is also found at the promoters of downregulated genes. This transcription factor
regulates in multiple metabolic pathways, including amino acid and nucleotide synthesis [84] indicating a
possible role of oxygen-regulated metabolic responses in *Monoraphidium*. Interestingly, motifs
corresponding to the same transcription factor were found among the photometers of the differentially
expressed genes during the acclimation period and after the infection with the amoeboapheids (Tables 1
and 2), which implies a role for transcription factors that bind this motif across stressors and
environmental changes.

We also noted that we identified overlap in the genes correlated with light and DO and those related to
stress responses (specifically responses to cold and salt), as determined by [17]. Specifically, of the 24
genes they outline as being annotated with a fasciclin-like domain, a domain with known roles in algal and
plant stress responses, we note 2 with expression coordinated with dissolved oxygen (annotation #
450375, 525465), 1 with expression coordinated with light (annotation # 480939) and 2 with expression
levels coordinated with both (annotation # 421681, 446308). These results support the validity of our
signature identification through multi-omic data integration in algae, and point towards the need for
follow-up on our stress signatures for identification of predictive markers for tracking large-scale pond
culture resiliency, and molecules for therapeutic intervention.

5. Conclusions

In this study, we used multi-omics to investigate the dynamic responses of the green alga *M. minutum* to
various environmental changes and stressors. A study of multi-omic analyses of M. *minutum* was recently
published, providing a molecular profile of this industrially significant species under high salt and cold
temperature stress [17]. We utilized a multi-omic approach in larger open and, thus, more heterogeneous
culture systems in order to provide a roadmap for the application of molecular observations at the cellular

level and data processing to address the significant challenges faced by the large-scale bioprocessing industry. Specifically, in the context of algae open pond systems, issues of unpredictability, inconsistency, and often deadly infections lead to crashes and severe crop losses [86] [42]. Our approach aims to address these issues while gaining information about the cellular responses of the species. We demonstrated the ability to integrate datasets of genes, proteins and metabolites that respond to specific stressors or correspond to certain culture stages and transitions, opening up possibilities for follow-up studies in the fields of strain engineering, biomarker development, bioprocessing, and basic cellular studies. We show that biomarker information can be extracted from this integrated data (transcripts, proteins, or metabolites) that indicate resilient growth and upcoming stressful events, such as prolonged acclimation periods during scale transitions and potential crashes due to the presence of pathogens. With respect to metabolites in particular, which are the most likely candidates for biomarker detection and immediate therapeutic intervention, specific hormones and steroids, such as 5-Dihydrotestosterone, 7-Hydroxytestosterone, Estrone, Estriol and 19-Norandrostenedion, are highly expressed during early infection. Hormones and steroids act as signaling molecules that regulating not only growth development but also stress adaptation [87] [88] [58]. Given that their abundance increased after the introduction of the pathogen, showcases that those compounds, produced by either the pathogen or the algae cells in response to their interaction with the pathogen, act as early regulators of a stress response and can be utilized as timely biomarkers for upcoming pond crashes. The detection of these markers provides preemptive information on impending phenotypic changes, which can be useful both for pond monitoring and eventual intervention and mitigation strategies. In the long term, if some of these molecules act in defense against algae, microbiomes can be engineered to produce these molecules as a prophylactic strategy during large-scale culture. The mechanism for this can be gleaned through the associated transcripts and proteins that are co-regulated.

In addition, when observing changes in transcription, proximal genomic elements can provide clues as to how algae can be engineered to optimize gene expression profiles and cell states. While there is not an abundance of information on transcription factor binding in *Monoraphidium* and related species, the presence of conserved sequences suggests not only that the binding of specific factors is likely to be pivotal in stress-related regulatory mechanisms but also that these mechanisms are likely conserved across algal species. These findings suggest that the factors identified here are likely to regulate stress response in other algal species and that mitigation and intervention strategies can be broadly applied. Future work will focus on testing across species to confirm these findings.

On a broader scale, the collection of molecular data from such expansive cultivation systems can pave the way for analyses aimed at developing predictive power with respect to culture resiliency, as well as potential targets for engineering this resiliency in the future. Indeed, we observed transcripts shared in both this study and the study conducted by Calhoun et al (2022), which indicates that more data generation in small- and large-scale systems, coupled with integrated analysis, could reveal novel health monitoring molecules in algal culture. The ultimate goal is for these models to be adaptable to multiple types of bioprocessing, beyond algae cultures, and this study provides a roadmap in that direction, using omic data to optimize large-scale applications.

Abbreviations

14-HDHA: 14-hydroxy-docosahexaenoic acid

AFDW: Ash-free dry weight

ARF1: ADP-ribosylation factor 1

ARR1: Arabidopsis response regulator 1

CCM: Carbon concentration mechanism

DIABLO: Data Integration Analysis for Biomarker discovery using Latent Components

DISCOVR: Development of integrated screening, cultivar optimization, and verification research

DPGP: Dirichlet Process Gaussian Process mixture model

Msn4: multicopy suppressor of SNF1 mutation protein 4

PCA: principal component analysis

ROS: reactive oxygen species

RuBisCO: Ribulose-1,5-bisphosphate carboxylase/oxygenase

RTG1: mitochondrial retrograde 1

RTG3: mitochondrial retrograde 3

SNF1: sucrose non-fermenting 1

Declarations

Ethics approval and consent to participate

N/A

Consent for publication

N/A

Availability of data and materials

The datasets supporting the conclusions of this article are available in the NCBI SRA repository, under
BioProject PRJNA1307767 and are included within the article and its additional files.

Competing interests

The authors declare that they have no competing interests.

Funding

This work was supported by the DOE Bioenergy Technology Office (BETO) and the Laboratory Directed
Research and Development program at Sandia National Laboratories, a multimission laboratory managed
and operated by National Technology and Engineering Solutions of Sandia LLC, a wholly owned subsidiary
of Honeywell International Inc. for the U.S. Department of Energy's National Nuclear Security
Administration under contract DE-NA0003525.

Authors' contributions

R.K. led the funding acquisition efforts and had the logistic and programmatic lead of the project. G.K and
R.K. conceptualized the project and designed the methodology and the experimental procedure. G.K.,
D.Y., E.W., T.E., O.W. and M.P.H. worked on the different aspects of the experimental procedure (scale-
up, pond running and maintenance, sampling and phenotyping, omics preparation and sequencing). G.K.,
J.S., D.Y., T.S. and R.K. worked on the computational and bioinformatic analysis. T.W.L. advised on the
experimental design and algal physiology. G.K, J.S. and R.K. wrote the original manuscript. All the authors
reviewed and edited the manuscript.

Acknowledgements

This paper describes objective technical results and analysis. Any subjective views or opinions that might
be expressed in the paper do not necessarily represent the views of the U.S. Department of Energy or the
United States Government. We thank Ryan W. Davis, Catherine Mageeney, Pamela Lane, and Carolyn
Fisher, Sharon Nademanee, and Damian Carrieri for their insightful comments on an earlier draft of this
manuscript.

Additional data

H3 histones

Additional Figure. 1: The caspase activity during the pond run. *The caspase activity for every day of the culture run in pond 1 (pink) and pond 2 (black).*

Additional Dataset 1: Phenomics

Additional Dataset 2: Transcriptomics

Additional Dataset 3: Proteomics

Additional Dataset 4: Metabolomics

References

1. Dismukes, G.C., et al., *Aquatic phototrophs: efficient alternatives to land-based crops for biofuels.* Current Opinion in Biotechnology, 2008. **19**(3): p. 235-240.
2. Cruce, J.R. and J.C. Quinn, *Economic viability of multiple algal biorefining pathways and the impact of public policies.* Applied Energy, 2019. **233-234**: p. 735-746.
3. Thanigaivel, S., et al., *The urge of algal biomass-based fuels for environmental sustainability against a steady tide of biofuel conflict analysis: Is third-generation algal biorefinery a boon?* Fuel, 2022. **317**: p. 123494.
4. Wiatrowski, M., et al., *Techno-economic assessment for the production of algal fuels and value-added products: opportunities for high-protein microalgae conversion.* Biotechnology for Biofuels and Bioproducts, 2022. **15**(1): p. 8.

5.   Costa, J.A.V., et al., *Chapter 9 - Open pond systems for microalgal culture*, in *Biofuels from Algae*
(Second Edition), A. Pandey, et al., Editors. 2019, Elsevier. p. 199-223.
6.   Jerney, J. and K. Spilling, *Large Scale Cultivation of Microalgae: Open and Closed Systems*, in
Biofuels from Algae: Methods and Protocols, K. Spilling, Editor. 2020, Springer New York: New
York, NY. p. 1-8.
7.   Lane, T.W., *Barriers to microalgal mass cultivation.* Current Opinion in Biotechnology, 2022. **73**:
p. 323-328.
8.   Quiroz, D., et al., *Geographical assessment of open pond algal productivity and evaporation losses*
across the United States. Algal Research, 2021. **60**: p. 102483.
9.   Echenique-Subiabre, I., et al., *Site-specific factors override local climatic conditions in determining*
microalgae productivity in open raceway ponds. Algal Research, 2023. **74**: p. 103235.
10.   Kleiman, R.M., et al., *Characterizing weather-related biophysical and financial risks in algal biofuel*
production. Applied Energy, 2021. **294**: p. 116960.
11.   Borowitzka, M. and A. Vonshak, *Scaling up microalgal cultures to commercial scale.* European
Journal of Phycology, 2017. **52**: p. 407-418.
12.   Patidar, S.K., et al., *Potential of Monoraphidium minutum for carbon sequestration and lipid*
production in response to varying growth mode. Bioresource Technology, 2014. **172**: p. 32-40.
13.   Dommange, X., P.A. Tanguy, and M. Jolicoeur, *Feasibility of lipid mechanical extraction from viable*
Monoraphidium minutum. Microalgae Biotechnol, 2015. **1**(1): p. 12-19.
14.   Bogen, C., et al., *Reconstruction of the lipid metabolism for the microalga Monoraphidium*
neglectum from its genome sequence reveals characteristics suitable for biofuel production. BMC
genomics, 2013. **14**: p. 1-18.
15.   Yu, X., et al., *Isolation of a novel strain of Monoraphidium sp. and characterization of its potential*
application as biodiesel feedstock. Bioresource Technology, 2012. **121**: p. 256-262.
16.   Pikoli, M.R., et al., *Characteristics of tropical freshwater microalgae Micractinium*
conductrix, Monoraphidium sp. and Choricystis parasitica, and
their potency as biodiesel feedstock. Heliyon, 2019. **5**(12).
17.   Calhoun, S., et al., *Multi-omics profiling of the cold tolerant Monoraphidium minutum 26B-AM in*
response to abiotic stress. Algal Research, 2022. **66**: p. 102794.
18.   Mikhailov, K.V., et al., *Genomic analysis reveals cryptic diversity in aphelids and sheds light on the*
emergence of Fungi. Current Biology, 2022. **32**(21): p. 4607-4619.e7.
19.   Letcher, P.M., et al., *Characterization of Amoeboaphelidium protococcarum, an Algal Parasite*
New to the Cryptomycota Isolated from an Outdoor Algal Pond Used for the Production of Biofuel.
PLOS ONE, 2013. **8**(2): p. e56232.
20.   Huesemann, M., et al., *DISCOVR strain pipeline screening – Part I: Maximum specific growth rate*
as a function of temperature and salinity for 38 candidate microalgae for biofuels production. Algal
Research, 2023. **71**: p. 102996.
21.   Van Wychen, S. and L.M.L. Laurens, *Determination of Total Solids and Ash in Algal Biomass:*
Laboratory Analytical Procedure (LAP). 2016, National Renewable Energy Lab. (NREL), Golden, CO
(United States): United States. p. Medium: ED; Size: 415 KB.
22.   Demichev, V., et al., *DIA-NN: neural networks and interference correction enable deep proteome*
coverage in high throughput. Nature Methods, 2020. **17**(1): p. 41-44.
23.   Kassambara, A. and F. Mundt. *Extract and Visualize the Results of Multivariate Data Analyses [R*
package factoextra version 1.0.7]. 2020.
24.   Bolger, A.M., M. Lohse, and B. Usadel, *Trimmomatic: a flexible trimmer for Illumina sequence data.*
Bioinformatics, 2014. **30**(15): p. 2114-20.
25.   Bray, N.L., et al., *Near-optimal probabilistic RNA-seq quantification.* Nature Biotechnology, 2016.
34(5): p. 525-527.

26.   Pimentel, H., et al., *Differential analysis of RNA-seq incorporating quantification uncertainty.*
Nature Methods, 2017. **14**(7): p. 687-690.

27.   Zhou, X. and M. Stephens, *Genome-wide efficient mixed-model analysis for association studies.*
Nat Genet, 2012. **44**(7): p. 821-4.

28.   Wickham, H., *ggplot2: Elegant Graphics for Data Analysis.* Springer-Verlag New York, 2016.

29.   Kolde, R., *pheatmap: Pretty Heatmaps.* 2025.

30.   Heinz, S., et al., *Simple combinations of lineage-determining transcription factors prime cis-*
regulatory elements required for macrophage and B cell identities. Mol Cell, 2010. **38**(4): p. 576-
89.

31.   Rohart, F., et al., *mixOmics: An R package for 'omics feature selection and multiple data*
integration. PLOS Computational Biology, 2017. **13**(11): p. e1005752.

32.   Dho, S.H., et al., *Caspases as master regulators of programmed cell death: apoptosis, pyroptosis*
and beyond. Experimental & Molecular Medicine, 2025. **57**(6): p. 1121-1132.

33.   Vavilala, S.L., et al., *KCl induces a caspase-independent programmed cell death in the unicellular*
green chlorophyte Chlamydomonas reinhardtii (Chlorophyceae). Phycologia, 2016. **55**(4): p. 378-
392.

34.   Zuppini, A., et al., *Chlorella saccharophila cytochrome f and its involvement in the heat shock*
response. Journal of Experimental Botany, 2009. **60**(14): p. 4189-4200.

35.   Teramoto, H., et al., *Light-intensity-dependent expression of Lhc gene family encoding light-*
harvesting chlorophyll-a/b proteins of photosystem II in Chlamydomonas reinhardtii. Plant Physiol,
2002. **130**(1): p. 325-33.

36.   McDowell, I.C., et al., *Clustering gene expression time series data using an infinite Gaussian*
process mixture model. PLOS Computational Biology, 2018. **14**(1): p. e1005896.

37.   Gao, S., et al., *DISCOVR strain screening pipeline – Part III: Strain evaluation in outdoor raceway*
ponds. Algal Research, 2023. **70**: p. 102990.

38.   Yang, H., Q. He, and C. Hu, *Feasibility of biodiesel production and CO2 emission reduction by*
Monoraphidium dybowskii LB50 under semi-continuous culture with open raceway ponds in the
desert area. Biotechnology for Biofuels, 2018. **11**(1): p. 82.

39.   Huesemann, M., et al., *A validated model to predict microalgae growth in outdoor pond cultures*
subjected to fluctuating light intensities and water temperatures. Algal Research, 2016. **13**: p. 195-
206.

40.   Matula, E.E. and D.J.A. Nabity, *Metabolic response of Chlorella vulgaris to a transient thermal*
environment for supporting simultaneous air revitalization and thermal control in a crewed
habitat. Acta Astronautica, 2021. **187**: p. 406-415.

41.   McBride, R.C., et al., *Contamination Management in Low Cost Open Algae Ponds for Biofuels*
Production. Industrial Biotechnology, 2014. **10**(3): p. 221-227.

42.   Harmon, V.L., et al., *Reliability metrics and their management implications for open pond algae*
cultivation. Algal Research, 2021. **55**: p. 102249.

43.   Ioc Coi Coi, M., et al., *Microscopic and molecular methods for quantitative phytoplankton analysis*

IOC. Manuals and guides.

44.   Havlik, I., et al., *Monitoring of microalgal cultivations with on-line, flow-through microscopy.* Algal
Research, 2013. **2**(3): p. 253-257.

45.   Murphy, T.E., K. Macon, and H. Berberoglu, *Rapid algal culture diagnostics for open ponds using*
multispectral image analysis. Biotechnology Progress, 2014. **30**(1): p. 233-240.

46.   Deore, P., et al., *Non-photochemical quenching, a non-invasive probe for monitoring microalgal*
grazing: an early indicator of predation by Oxyrrhis marina and Euplotes sp. Applied Phycology,
2020. **1**(1): p. 20-31.

47. Fundel, B., et al., *Can phaeopigments be used as markers for Daphnia grazing in Lake Constance?*
Journal of Plankton Research, 1998. **20**(8): p. 1449-1462.
48. Bartley, M.L., et al., *Effects of salinity on growth and lipid accumulation of biofuel microalga*
Nannochloropsis salina and invading organisms. Biomass and Bioenergy, 2013. **54**: p. 83-88.
49. Reese, K.L., et al., *Chemical Profiling of Volatile Organic Compounds in the Headspace of Algal*
Cultures as Early Biomarkers of Algal Pond Crashes. Scientific Reports, 2019. **9**(1): p. 13866.
50. Xie, J., et al., *Genome-Wide Identification and Functional Analysis of C2H2 Zinc Finger*
Transcription Factor Genes in the Intertidal Macroalga Pyropia haitanensis. Int J Mol Sci, 2025.
26(9).
51. Thiriet-Rupert, S., et al., *Transcription factors in microalgae: genome-wide prediction and*
comparative analysis. BMC Genomics, 2016. **17**(1): p. 282.
52. Liu, L., et al., *Enigmatic role of auxin response factors in plant growth and stress tolerance.* Front
Plant Sci, 2024. **15**: p. 1398818.
53. Ulmasov, T., G. Hagen, and T.J. Guilfoyle, *ARF1, a Transcription Factor That Binds to Auxin*
Response Elements. Science, 1997. **276**(5320): p. 1865-1868.
54. Linares-Maurizi, A., et al., *Stress-Induced Production of Bioactive Oxylipins in Marine Microalgae.*
Mar Drugs, 2024. **22**(9).
55. Lai, K.M., M.D. Scrimshaw, and J.N. Lester, *Biotransformation and bioconcentration of steroid*
estrogens by Chlorella vulgaris. Appl Environ Microbiol, 2002. **68**(2): p. 859-64.
56. Shi, W., et al., *Removal of estrone, 17alpha-ethinylestradiol, and 17beta-estradiol in algae and*
duckweed-based wastewater treatment systems. Environ Sci Pollut Res Int, 2010. **17**(4): p. 824-
33.
57. Liu, D., et al., *Biochemical responses of freshwater microalgae Chlorella sorokiniana to combined*
exposure of Zn(Ⅱ) and estrone with simultaneous pollutants removal. J Environ Manage, 2023.
348: p. 119392.
58. Yu, H., et al., *Exploring the molecular mechanism of Chlorella vulgaris in response to*
androstenedione exposure based on genes continuously up-regulated in transcription analysis.
Ecotoxicology and Environmental Safety, 2024. **271**: p. 115996.
59. Czarny, K., et al., *Toxicity of single steroid hormones and their mixtures toward the*
cyanobacterium Microcystis aeruginosa. Journal of Applied Phycology, 2019. **31**(6): p. 3537-3544.
60. Yamaoka, Y., et al., *Light, CO2, and carbon storage in microalgae.* Current Opinion in Plant Biology,
2025. **84**: p. 102696.
61. Tokutsu, R., et al., *Isolation of photoprotective signal transduction mutants by systematic*
bioluminescence screening in Chlamydomonas reinhardtii. Sci Rep, 2019. **9**(1): p. 2820.
62. Ruiz-Sola, M.Á., et al., *Light-independent regulation of algal photoprotection by CO2 availability.*
Nature Communications, 2023. **14**(1): p. 1977.
63. Maltsev, Y., et al., *Influence of Light Conditions on Microalgae Growth and Content of Lipids,*
Carotenoids, and Fatty Acid Composition. Biology (Basel), 2021. **10**(10).
64. Dao, O., et al., *The green algae CO2 concentrating mechanism and photorespiration jointly*
operate during acclimation to low CO2. Nature Communications, 2025. **16**(1): p. 5296.
65. Jahan, K., et al., *Transcriptomic Analysis Provides New Insights into the Tolerance Mechanisms of*
Green Macroalgae Ulva prolifera to High Temperature and Light Stress. Biology, 2024. **13**(9): p.
725.
66. Serrano-Pérez, E., et al., *Transcriptomic and Metabolomic Response to High Light in the*
Charophyte Alga Klebsormidium nitens. Frontiers in Plant Science, 2022. **13**.
67. Wang, B., et al., *Cell damage repair mechanism in a desert green algae Chlorella sp. against UV-B*
radiation. Ecotoxicology and Environmental Safety, 2022. **242**: p. 113916.

68.   Nama, S., et al., *High light induced changes in organization, protein profile and function of photosynthetic machinery in Chlamydomonas reinhardtii.* Journal of Photochemistry and Photobiology B: Biology, 2015. **152**: p. 367-376.

69.   Kirch, T., et al., *The two homeodomains of the ZmHox2a gene from maize originated as an internal gene duplication and have evolved different target site specificities.* Nucleic Acids Res, 1998. **26**(20): p. 4714-20.

70.   Bondino, H.G. and E.M. Valle, *A small intergenic region drives exclusive tissue-specific expression of the adjacent genes in Arabidopsis thaliana.* BMC Mol Biol, 2009. **10**: p. 95.

71.   Ejaz, M., et al., *ARABIDOPSIS THALIANA HOMEOBOX GENE 1 controls plant architecture by locally restricting environmental responses.* Proc Natl Acad Sci U S A, 2021. **118**(17).

72.   Polish, J.A., J.-H. Kim, and M. Johnston, *How the Rgt1 Transcription Factor of Saccharomyces cerevisiae Is Regulated by Glucose.* Genetics, 2005. **169**(2): p. 583-594.

73.   Valifard, M., et al., *The novel chloroplast glucose transporter pGlcT2 affects adaptation to extended light periods.* Journal of Biological Chemistry, 2023. **299**(6): p. 104741.

74.   Powner, M.B. and G. Jeffery, *Light stimulation of mitochondria reduces blood glucose levels.* Journal of Biophotonics, 2024. **17**(5): p. e202300521.

75.   Rajvanshi, P.K., M. Arya, and R. Rajasekharan, *The stress-regulatory transcription factors Msn2 and Msn4 regulate fatty acid oxidation in budding yeast.* J Biol Chem, 2017. **292**(45): p. 18628-18643.

76.   Al-Azab, A.J.S., et al., *Chronological transcriptome changes induced by exposure to cyanoacrylate resin nanoparticles in Chlamydomonas reinhardtii with a focus on ROS development and cell wall lysis-related genes.* Algal Research, 2022. **68**: p. 102884.

77.   Gao, Y. and D.L. Erdner, *Cell death responses to acute high light mediated by non-photochemical quenching in the dinoflagellate Karenia brevis.* Scientific Reports, 2022. **12**(1): p. 14081.

78.   Xu, C., et al., *Programmed cell death process in freshwater Microcystis aeruginosa and marine Phaeocystis globosa induced by a plant derived allelochemical.* Science of The Total Environment, 2022. **838**: p. 156055.

79.   Bai, F., et al., *Paraquat induces different programmed cell death patterns in Microcystis aeruginosa and Chlorella luteoviridis.* Ecotoxicology and Environmental Safety, 2023. **249**: p. 114429.

80.   Zhao, Y., et al., *ROS-mediated programmed cell death (PCD) of Thalassiosira pseudonana under the stress of BDE-47.* Environmental Pollution, 2020. **262**: p. 114342.

81.   Zhao, M.G., Q.Y. Tian, and W.H. Zhang, *Nitric oxide synthase-dependent nitric oxide production is associated with salt tolerance in Arabidopsis.* Plant Physiol, 2007. **144**(1): p. 206-17.

82.   Moreau, M., et al., *AtNOS/AtNOA1 is a functional Arabidopsis thaliana cGTPase and not a nitric-oxide synthase.* J Biol Chem, 2008. **283**(47): p. 32957-67.

83.   Sakai, H., et al., *ARR1, a Transcription Factor for Genes Immediately Responsive to Cytokinins.* Science, 2001. **294**(5546): p. 1519-1521.

84.   Hashim, Z., et al., *Metabolic profiling of retrograde pathway transcription factors rtg1 and rtg3 knockout yeast.* Metabolites, 2014. **4**(3): p. 580-98.

85.   Singh, A., et al., *DIABLO: an integrative approach for identifying key molecular drivers from multi-omics assays.* Bioinformatics, 2019. **35**(17): p. 3055-3062.

86.   Harun, R., et al., *Bioprocess engineering of microalgae to produce a variety of consumer products.* Renewable and Sustainable Energy Reviews, 2010. **14**(3): p. 1037-1047.

87.   Bajguz, A., *Brassinosteroids in Microalgae: Application for Growth Improvement and Protection Against Abiotic Stresses*, in *Brassinosteroids: Plant Growth and Development*, S. Hayat, et al., Editors. 2019, Springer Singapore: Singapore. p. 45-58.

88.   Stirk, W.A., et al., *Endogenous brassinosteroids in microalgae exposed to salt and low temperature*
1027   *stress.* European Journal of Phycology, 2018. **53**(3): p. 273-279.

1028