# The Drosophila connectome reveals Axo-Axonic Synapses on Descending Neurons that modulate the Giant Fiber System

Cesar Ceballos[1], Juan Lopez[1], Ty Roachford[1], Daniel Sanchez[1], Sabrina Jara[1], Kelli Robbins[1], Casey L Spencer[2], Rodney Murphey[1], Rodrigo FO Pena[1,*]

[1]Department of Biological Sciences, Florida Atlantic University, FL 33458, Jupiter, Florida, USA

[2]Harriet Wilkes Honors College, Florida Atlantic University, Jupiter, FL, 33458, USA

*penar@fau.edu

## Abstract

Axo-axonic synapses can veto, amplify, or synchronize spikes, yet their circuit-scale logic is unknown. Using the complete electron-microscopy connectome of the adult male Drosophila ventral nerve cord (MANC v1.2.1), we charted every axo-axonic input onto the 1,314 descending neurons that carry brain commands to the ventral nerve cord. Only 1.0 % of the 861,591 possible descending–descending neuron pairs form such contacts, but when present, synaptic strength grows linearly with partner number regardless of transmitter identity. By definition, any synapse connected to a descending neuron within the cord is axo-axonic. Thus, we uncovered the ascending-descending and interneurons-descending axo-axonic relationship. Neurons with many partners (high-degree nodes) integrate into the network without clustering into a densely interconnected 'rich-club' of hubs. The connectome identified an octet of ascending neurons (AN08B098) whose axo-axonic input to the Giant Fiber descending neurons (DNp01) predicted modulation of escape control. Immunostaining confirms their cholinergic identity, while optogenetic activation confirmed that this excitatory cohort increases Giant Fiber excitability, validating connectome-derived rules. Our work delivers the first comprehensive map of axo-axonic wiring in a complete ventral nerve cord connectome and constraints for models of fast motor control.

**Keywords:** Connectome, Drosophila, Axo-axonic, Giant Fiber escape circuit, Network topology, Electron-microscopy reconstruction, Motor control

**Introduction**

Neurons communicate through a variety of synaptic motifs. In flies, the canonical arrangement is axo-dendritic, but a second configuration, axo-axonic, can exert unusually powerful control over spike initiation and transmitter release[1,2]. Axo-axonic contacts (AACs) in invertebrates have been known for some time[2–6]. Similar studies have also been conducted in vertebrates[7–9]. In the mammalian cortex, exemplified by chandelier cells, recent studies show how inhibition modulates the pyramidal axons at their initial segments[10,11]. In fish, excitatory spiral fibers and M-cells control a similar escape circuit[12]. Although these examples suggest broader significance, the quantitative wiring logic of axo-axonic synapses remained poorly characterized, primarily due to the absence of a fully mapped system at the necessary synaptic resolution.

This limitation has been addressed by recent electron microscopy reconstructions of the complete adult male nerve cord and female brain (MANC[13–15] and FAFB[16]) in *Drosophila melanogaster*, now publicly available through the NeuPrint interface[17]. Early graph analyses of the FAFB data revealed rich-club hubs and other mesoscale motifs[16], but systematic exploration of the ventral nerve cord (VNC), the insect analogue of the vertebrate spinal cord, is only beginning. At the heart of the VNC, a pair of descending neurons, known as Giant Fibers (GFs), function as command neurons that receive visual input in the brain and drive jump and flight motor neurons[18,19]. These GFs receive a high number of AACs; thus, it is only natural to assume that an AAC-to-behavior modulation is in place. This system is comparable to the descending escape circuits in larger insects, which have been shown to be the target of axo-axonic connections, exemplified by the input to the descending contralateral movement detector (DCMD) neurons in the locust central nervous system[4].

Within the VNC, a single class of neurons is ideally positioned for studying axo-axonic wiring and their connection to the GFs: Descending neurons (DNs). Because DNs have their somata and dendrites in the brain yet project their axons into the VNC[20–23], any synapse made onto a DN within the cord is, by definition, axo-axonic. In fact, we have manually curated 1,000 ascending-descending neurons connections, where we observed clear presynaptic mitochondrial expression (93%), a hallmark of axonal terminals[24] (example shown in Fig. S1, and curation details in Table S1). The MANC connectome contains DN, ascending neurons (ANs), intrinsic interneurons (INs), and motor neurons (MNs), offering a comprehensive substrate for network-wide analysis of this motif. Although classic anatomical work suggested that each GF acts largely in isolation, our work using the MANC connectome reveals an unexpectedly rich set of axo-axonic inputs onto the GF axons, including direct contacts from other DNs, ANs, and INs. In the present study, we show that the GF receives 97 distinct axo-axonic partners, with some offering a crucial role for axo-axonic modulation, i.e., an octet of ascending neurons (AN08B098). Below, we will refer to these neurons of interest as Axc.

Past light-microscopic surveys catalogued individual DN types and noted scattered DN-to-DN
contacts[20], yet the circuit-level role of axo-axonic synapses has remained unexplored. The
outstanding issues are their prevalence, partner preference, and functional impact on fast behavior.
Specifically, it is unknown how frequently axo-axonic connections occur among the 861,591
possible DN pairs, and whether such a sparse motif can nevertheless mediate rapid, system-wide
control required for escape. We tackle these questions by mining the complete electron-
microscopy connectome of the adult male ventral nerve cord (MANC). Our analysis (i) enumerates
every axo-axonic input onto all 1,314 DNs, (ii) dissects the morphological relationship to DN
connectivity, (iii) shows that the DN network is small-world but not rich-club, partitioned into
thirteen functional communities, (iv) uses large-scale leaky integrate-and-fire simulations to
predict which highly connected AACs modulate the Giant Fiber escape pathway, and (v) confirms
those predictions with targeted optogenetic activation *in vivo*.
81
Our results provide the first system-level blueprint of axo-axonic wiring in the *Drosophila* adult
nervous system, suggesting general design principles for descending motor control that may extend
beyond insects. They also illustrate the functional power of the sparse axo-axonic network
described below and place the GFs at the center of a distributed control system rather than a simple
linear reflex arc.

**Results**
**Global Distribution of Axo-Axonic Inputs to Descending Neurons**.
To uncover the prevalence of axo-axonic connections between DNs in the VNC, we used the
MANC dataset available through the NeuPrint+ interface and API[25–27]. Figure 1*A-C* shows
morphological reconstructions of some exemplary DNs of interest from Neuroglancer: DNp01
(GFs, black dot), DNd03 (blue dot), and DNg02 (red dot, Fig. 1*A-C*). For each neuron,
Neuroglancer overlays the 3D-EM reconstruction on the VNC volume. The GFs, whose axons are
nearly unbranched (green or blue), receive one of the highest numbers of axo-axonic inputs (n =
97; see NeuPrint+) in the dataset. DNd03 is a highly arborized neuron with a similarly large input
count (n=160), while DNg02 has less arborization but similar axo-axonic inputs (n=110).
Interestingly, all three share similar synaptic strength despite differences in arborization (DNp01
2352, DNd03 2364, DNg02 1751; Fig. 1*H*). Neuroglancer presents transmission-EM micrographs
at 500 nm resolution taken from MANC; presynaptic active zones are marked by magenta dots,
postsynaptic densities by cyan dots, and partner axons are pseudo-colored to match the 3D
rendering (Fig. 1*D-F*).
102

**Figure 1. Descending neurons (DNs) and their axo-axonic inputs in the VNC.** *(A)*-*(C)* Exemplary Neuroglancer reconstructions of three DNs of interest. *(D)*-*(F)* EM images illustrate axo-axonic connections formed by the three reconstructed neurons: *(D)* shows DNp01 to DNp01 connectivity, *(E)* shows DNd03 connectivity with DNp42, and *(F)* shows DNg02 to DNg02 connectivity (same colors as in A-C). Red circles identify the locations of presynaptic T-bars; blue circles show postsynaptic densities. Circles were enlarged for visibility. *(G)* Histogram of the number of axo-axonic neurons innervating one DN. Presynaptic neurons can be DN, AN, or IN. *(H)* Histogram of the cumulative strength of the synaptic connections for each DN. Presynaptic neurons can be DN, AN, or IN. *(I)* Scatter plot and linear regression of the number of axo-axonic neurons vs their synaptic strength. $R^2 = 0.92$ and $p < 0.001$.

We quantified the distribution of axo-axonic connections made onto DNs reported by MANC. Figure 1*G* reveals a highly skewed distribution of partner count; the median DN is innervated by a single presynaptic neuron (inter-quartile range 1–3), and 11% of DNs receive input from only one cell with a synaptic strength of six or greater. A small set of outliers, however, attracts hundreds of partners, including DNp01 (GFs) with 97 distinct axo-axonic inputs. Strikingly, the GFs receive these connections despite possessing one of the simplest axonal arbors in the dataset (Fig. 1*A*). All other DNs exhibit more extensively branched and structurally complex axonal arbors, which are likely to account for their higher levels of innervation. Total input strength follows a similar heavy-tailed pattern (Fig. 1*H*). The modal DN receives fewer than 20 synapses, whereas the GF is again among the top of the list in terms of individual contacts. Across the population, partner count and cumulative strength are tightly coupled (Pearson r = 0.96, p < 0.001; Fig. 1*I*), indicating that neurons that recruit many partners also receive proportionally stronger innervation. Together, these findings identify the distribution of axo-axonic connections made with DNs in the VNC. We confirm that the synapses quantified in the graph analysis correspond to active zones and demonstrate how neurons with markedly different morphologies can converge on similar axo-axonic input profiles.

Using these findings, we next examine whether anatomical complexity serves as a strong, though not perfect, predictor of synaptic strength for axo-axonic inputs onto DNs (Fig. 2). Representative reconstructions illustrate the general pattern: neurons with short, unbranched axons receive few contacts, those with intermediate arborization receive proportionately more, and highly branched cells accumulate the greatest axo-axonic strength (Fig. 2*A-C*). A kernel density estimation (KDE) plot of all 1,314 DNs reinforces this observation, showing a diagonal ridge in which increasing cable length accompanies increasing total input strength (Fig. 2*F*). Linear, quadratic, and symbolic regressions all yield significant positive slopes, confirming a monotonic relationship (Fig. 2*G*). However, the shallow gradient and the presence of two striking outliers, a long axon with sparse innervation and a short axon with dense innervation, indicate that branch length alone does not determine synaptic investment (Fig. 2*D-E*). Thus, we find that a presynaptic neuron's morphology broadly predicts the degree of modulation received by a partner DN, although additional cell-type-specific factors likely refine the final allocation of axo-axonic synapses.

**Figure 2. Analysis of morphology and synaptic strength between DN-to-DN axo-axonic connections.** (*A-E*)
Examples of DNs include (legend colors indicate pre- and postsynaptic in the figure): (A) low connectivity with simple
morphology, (B) moderate connectivity and morphology, and (C) high connectivity with complex morphology.

Additionally, (D) low connectivity with complex morphology and (E) high connectivity with simple morphology
represent exceptions. These neurons generally align with the overall trend—a linear correlation between
morphological complexity and synaptic strength—while (D) and (E) are outliers. Dots indicate synapse locations. *F*)
Kernel density estimation (KDE) of the scatter data of total input strength vs cable length, *G*) Scatter plot of total input
strength vs cable length and the corresponding linear regression, polynomial regression of a quadratic function, and
symbolic regression.

Further exploration revealed a unique feature of connectivity. In the VNC, neurons will often form
synapses with multiple downstream partners (Fig. 3*A-B*). We found certain neurons, such as
members of IN21A026 and some members of AN08B098, had a disproportionately larger number
of connections made onto a single target type of DNs such as the GF than other neuron type. To
determine whether these neurons constitute a separable class, we quantified neuron output
specialization for every presynaptic neuron using three metrics: the fraction of total synapses made
onto the dominant DN, the entropy of the neuron type distribution, and the number of connections
of the second largest neuron type connection (off-target) relative to the largest DN connection (on-
target).

For each presynaptic neuron, we first quantified the percentage of strength of its outgoing synapses
that were made onto its most strongly targeted DN. Neurons were required to exceed a threshold
value and the percentage of strength of the dominant DN was permitted to exceed the threshold,
ensuring a uniquely dominant target. Initially, we assumed threshold values that separate two
populations: multi-target vs single-type biases. The final threshold was determined by optimizing
the minimization of an error parameter. A statistical comparison of the distributions using the
Kolmogorov-Smirnov test (with a significant p-value < 0.001) revealed two distinct distributions
of presynaptic neurons, which we defined as single-type biased and multi-target, with threshold
values of 0.27 in MANC v1.2.1 and 0.1 in male-cns v0.9. In addition, single-biased cannot send
more than 15 connections to other off-target DN to exclude external influence[28].

In both connectomes, we identified several single-type biased axo-axonic connections (Fig. 3*C*) to
DNs in both MANC v1.2.1 and male-cns v0.9. DNp01 receives the highest number of single-target
AACs in MANC (n = 16) and among the highest for male-cns (n = 11). Among these neurons, five
belong to the Axc type, and the remaining are labeled as "unknown".

183
**Figure 3. Neurons that provide exclusive axo-axonic connections to DNs.** (*A*) Sketch of single type biased vs (*B*)
multi-target neuron. Single-type biased neurons project a significant proportion of their outgoing connections to a
single DN, while unbiased neurons form several strong connections among partners. (*C*) Number of exclusive neurons
per DN for both MANC v1.2.1 (blue histogram) and Male-cns v0.9 (orange histogram). The Giant Fibers (DNp01;
marked by an asterisk) are among a small population of DNs that receive most single-type biased connections, i.e., as
seen in (*D*), there are approximately 16 neurons that innervate DNp01 in MANC and 11 in Male-cns. (*D*) For each
dataset, the fraction of outputs made from neurons onto their most strongly targeted DN type was computed per
neuron, and a dataset-specific cutoff was identified from the distribution as the point separating the population of
unbiased neurons and single-type biased neurons (MANC 0.27, male-cns 0.1). Single-type biased neurons across both
datasets had similar properties, including high target preference, reduced target entropy, and reduced off-target
connection strength (Kolmogorov-Smirnov test, $p < 0.001$). We further refined this by removing neurons whose off-
target connections had 15 or greater synapses, suggesting non-influence over their synaptic partners[28].
196
Compared with other highly connected DNs, the GFs attain an equivalent synaptic strength with a
simple morphology, further demonstrating how morphological complexity alone does not predict
axo-axonic input. These results identify the GFs as exceptional hubs for axo-axonic modulation
due to their significant connectivity despite their relatively simple morphology compared to other

highly connected DNs. However, these findings raise several new questions: Which specific AAC
populations account for the GF's unusually dense axo-axonic input? How do these contacts alter
GF excitability? Is the GF an outlier, or do similar axo-axonic motifs govern other rapid motor
pathways in the fly? We address these questions by investigating the strongest single-target AAC
population presynaptic to the GFs, the Axc type neurons.
206
**Sparse yet Globally Connected Axo-Axonic Network among Descending Neurons**.
To study how DNs communicate with other DNs through axons, we extracted every DN-to-DN
contact from the updated MANC dataset. The resulting adjacency matrix is strikingly sparse: only
about 1% of the possible ordered pairs carry even a single synapse (Fig. 4*A*). Moreover, partner
choice is not random: homotypic pairs (same DN type) and heterotypic pairs (different types) show
distinct transmitter profiles, as shown for acetylcholine, GABA, and glutamate, implying that
neurotransmitter identity contributes to the selectivity of these long-range axo-axonic contacts
(Fig. 4*D*). In homotypic DN connections, ~90% of the connections are cholinergic (excitatory),
while heterotypic DN connections are ~60% cholinergic [the remaining are GABA/glutamate
(inhibitory)]. This suggests that AACs between DNs, while rare, are predominantly excitatory.
connections.
218
We then asked whether the same wiring principle applies to non-descending inputs, namely
ascending neurons (ANs) and intrinsic interneurons (INs). Repeating the analysis with ANs and
INs as presynaptic partners to DNs again produced very sparse matrices: about 1.2% of all possible
AN-to-DN pairs and 0.7% of the IN-to-DN pairs contain at least one synapse (Fig. 4*B-C*). Figure
4*E* shows the neurotransmitter profile for AN-to-DN and IN-to-DN pairs. Both pair profiles were
similar, with nearly half of the neurons classified as cholinergic and the other half
GABA/glutamate. This suggests that AACs between AN-to-DN and IN-to-DN follow the same
rules for neurotransmitter identity as DN-to-DN AACs.
227

A — DN-DN connectivity matrix
B — AN-DN connectivity matrix
C — IN-DN connectivity matrix

D
Different DN — acetylcholine 1331 (58%), gaba 657 (29%), glutamate 304 (13%)
Same DN — acetylcholine 434 (91%), gaba 24 (5%), glutamate 19 (4%)

acetylcholine
gaba
glutamate

E
AN->DN — acetylcholine 1100 (51%), gaba 613 (29%), glutamate 435 (20%)
IN->DN — acetylcholine 3582 (60%), gaba 1347 (23%), glutamate 1038 (17%)

F1 — in-degree — Fraction vs # presynaptic DN
G1 — Fraction vs # presynaptic AN
H1 — Fraction vs # presynaptic IN
F2 — out-degree — Fraction vs # postsynaptic DN
G2 — Fraction vs # postsynaptic DN
H2 — Fraction vs # postsynaptic DN

J — Connectivity matrix
Presynaptic AN/DN/IN vs Postsynaptic DN
DNd03   DNg02   DNp01
IN / DN / AN

I — Rich club neurons
Rich club coefficient vs Degree threshold (k)

K — DN
L — DN
M — AN
N — AN
O — IN
P — IN

Q — % connections — DN->DN, AN->DN, IN->DN — ipsi / contra

R — DN->DN — acetylcholine 48%, gaba 38%, glutamate 13%
S — AN->DN — acetylcholine 48%, gaba 37%, glutamate 14%
T — IN->DN — acetylcholine 49%, gaba 35%, glutamate 16%

acetylcholine
gaba
glutamate

**Figure 4. Axo-axonic connectivity analysis.** (*A*) Connectivity matrix of pairs of DN-to-DN pairs. Only 1% of all possible connections were identified, suggesting that DN-to-DN axo-axonic connections are extremely rare. (*B*) Connectivity matrix of pairs of AN-to-DN pairs. Only 1.2% of all possible connections were found, which suggests that AN-to-DN axo-axonic connections are also exceptionally rare. (*C*) Connectivity matrix of pairs of IN-to-DN pairs. Only 0.7% of all possible connections were observed, indicating that IN-to-DN axo-axonic connections are significantly too uncommon. (*D*) Pie chart of predicted neurotransmitter of DN of the different type (left, e.g., DNp01 -> DNp02) and same type (right, e.g., DNp01 ->DNp01) that make pairs of axo-axonic connections. (*E*) Pie chart of predicted neurotransmitter of AN-to-DN and IN-to-DN axo-axonic connections. (*F*) Histograms of the number of (F1) presynaptic DN (in-degree) and (F2) postsynaptic DN (out-degree) for DN-to-DN pairs. (*G*) Histograms of the number of (G1) presynaptic AN and (G2) postsynaptic DN for AN-to-DN pairs. (*H*) Histograms of the number of (H1) presynaptic IN and (H2) postsynaptic DN for IN-to-DN pairs. (*I*) Rich-club analysis of the DN-to-DN axo-axonic network. (*J*) Connectivity matrix of pairs of AN/DN/IN-to-DN connections. Filled circles follow the same color scheme as in Fig. 1*A*-*C*. Cyan square identifies Axc-type neurons. Dashed green lines and gray tones for dots separate ANs, DNs, and INs. (*K*-*P*) Examples of each category from DN-to-DN, AN-to-DN, and IN-to-DN. (From left to right: (*K*) DNx01-DNx01, (*L*) DNb04-DNb04, (*M*) AN08B099-DNp01, (*N*) AN10B019-DNp01, (*O*) IN11A001-DNp01 and (*P*) IN06B059-DNp01). (*Q*) Percentage of contralateral and ipsilateral connections between DN-to-DN, AN-to-DN, and IN-to-DN. (*R*-*T*) Pie chart of predicted neurotransmitter of DN to contralateral DN, AN to contralateral DN, and IN to contralateral DN of axo-axonic connections.

Lastly, we show the distribution of the number of pre-synaptic DN, AN, or IN that project axo-axonic synapses to a post-synaptic DN (Fig. 4*F*-*H* top row) as well as the number of postsynaptic DN that are innervated axo-axonically by the same DN, AN, or IN (Fig. 4*F*-*H* bottom row). The distributions are similar across neuron types, showing a decay pattern resembling an exponential (Poisson) distribution, where a few neurons are highly connected, and a few neurons are sparsely connected. When we merge all three presynaptic classes into a single connectivity matrix (Fig. 4*J*,) the same pattern from Figure 4*A*-*C* emerges: extreme sparseness (see dots where our exemplary neurons are located; color scheme as in Fig. 1*A*.). Note that in Fig. 4*A* there is a clear diagonal line which indicates DNs synapsing onto their contralateral counterparts. We also highlight the degree of convergence across presynaptic classes and identify connectivity hubs based on their total input. Taken together, regardless of presynaptic identity, the findings demonstrate that the network's organization combines stringent partner selectivity with complete coverage of the descending system.

Global wiring principles were identified to explore more refined AAC spatial connectivity preferences and determine whether they are repeated across DNs, ANs, and INs. Figure 5 shows examples of the contralateral axo-axonic connections between DN-to-DN (Fig. 4*K*-*L*), AN-to-DN (Fig. 4*M*-*N*), and IN-to-DN (Fig. 4*O*-*P*). Figure 4*Q* shows the percentage of contralateral and ipsilateral connections. AACs between contralateral DNs are rare. However, ANs and INs projecting to contralateral DNs are frequent. Interestingly, the neurotransmitter profiles from our earlier findings are preserved regardless of the type of presynaptic neuron, where approximately half of the neurons are cholinergic (excitatory), and the other half is GABAergic/glutamatergic (inhibitory) (Fig. 4*R*-*T*).

Collectively, these findings suggest a coherent narrative. Whether the presynaptic axon descends, ascends, or remains intrinsic to the cord, axo-axonic wiring obeys the same two rules: partner selectivity is extremely high, and midline crossing is common only for neurons that share strong

275 reciprocal ties. The result is a VNC in which local subtypes can interact tightly with their mirror
276 images, yet the great majority of neurons remain insulated from direct axo-axonic influence.
277

**Graph-Theoretic Architecture of the Axo-Axonic Network**.
The connectivity matrix we derived was employed as a directed graph to examine the system-wide
arrangement of sparse axo-axonic interactions among DNs (Fig. 4*A*). The graph contains 884
nodes and 2,757 edges, giving a density of 0.007. Despite this low density, the average shortest-
path length is only 4.03, indicating that any two DNs can be reached through a small number of
intermediate axonal contacts. In fact, we can compare with the chance level: the formula $\langle L \rangle =$
$\ln(n)/\ln(\bar{k})$ is a well-known and widely used approximation for the average shortest-path length
in large random networks, such as the Erdős-Rényi model. Here, $n$ is the number of vertices in the
graph, $\bar{k}$ is the mean degree, which can be obtained by the ratio $m/n$ with $m$ being the edge count.
In our problem, $n = 884$ DNs and $m = 2,757$ axo-axonic contacts leave us with $\langle L \rangle \approx 5.9$.
Therefore, the average shortest path length is indeed shorter than you would expect by chance.
The clustering coefficient is likewise low at 0.013, indicating limited local redundancy.
Community detection identifies 13 separate modules, consistent with the network's functional
subdivision. Node degree ranges from 1 to 29 with a mean of 6.24, reflecting a modest hierarchical
spread.
293

The rich-club coefficient Φ(k) measures the density of connections among nodes whose total
degree (in-degree + out-degree) equals or exceeds a threshold k. Φ(k) = 1 indicates that the high-
degree subgraph is fully connected, whereas Φ(k) = 0 means that those nodes share no edges
beyond chance level. The rich-club coefficient in our data peaks at 0.076 for nodes with a degree
of at least 18, so high-degree hubs do not preferentially connect to each other; integration is
therefore distributed rather than centralized (Fig. 4*I*). The modest peak value and the absence of
a sustained plateau indicate that high-degree DNs do not preferentially interconnect to form a rich
club; instead, network integration is distributed rather than hub-dominated.
302

A slightly positive degree assortativity of 0.029, a measure that suggests a tendency of nodes to
connect with other nodes in a similar way, suggests a weak tendency for neurons of similar degree
to link. Finally, the maximum betweenness centrality of 0.049, a measure indicating how often a
node lies on the shortest path between all pairs of nodes in a network, highlights a small set of
bridge neurons that channel information between modules. Together, these metrics portray a
network that combines efficient global communication with modular specialization, an
architecture well-suited for coordinating diverse motor outputs.
310

**Axo-axonic modulators of the Giant Fiber escape circuit.**
The MANC connectome introduces several novel neuron types to the Giant Fiber escape circuit.
To understand and underscore the significance of AAC modulation of the circuit, it is important
to know which individual cells can exert a strong, targeted influence on a system such as the GF
system[18,19]. As shown in Figure 3, the GFs are among a small population of DNs that receive

single-target connections. A focused search within MANC revealed the Axc type neurons, a cluster
of eight neurons whose axo-axonic output is both nearly exclusive and among the strongest to the
GFs (Figs. 3-4). These Axc-type neurons are predicted to be cholinergic by the MANC
connectome. The BANC connectome demonstrates that these neurons do not connect to the GF in
the brain (see Fig. S3).
321
To test whether the connectome-predicted Axc type neurons truly form chemical AACs with the
GFs, we used split-Gal4 drivers to direct membrane-bound GFP to Axc axons, back-filled the GFs
with tetramethyl-rhodamine, and labelled presynaptic active zones with anti-Bruchpilot (nc82).
High-resolution confocal stacks revealed discrete GFP-positive boutons that colocalize with Brp
puncta on the descending GF tract in the first thoracic neuromere (Fig. 5*A*,*C*), confirming axo-
axonic contacts between Axc type neurons and GF. The split-Gal4 line being used labels
AN08B098. However, there are a few extra neurons labeled that can come from AN08B099 and
make weak connections to the GF. We have also provided (i) single-section confocal images with
optical enhancement showing the synaptic connectivity and (ii) similar EM reconstructions from
Neuprint to demonstrate that active synapse locations along the axon correspond to synapse
locations identified in the MANC connectome (Fig.5*G*-*O*). Immunostaining for Choline
Acetyltransferase (ChAT) revealed that Axc somata, as labelled through genetic expression of
mCD8::GFP, are ChAT-positive (Fig. 5*D*-*G*), establishing this cohort as likely cholinergic and
therefore excitatory.
336

**Figure 5. Axcs make cholinergic axo-axonic synapses on the GFs as predicted by the MANC connectome.**
(*A*) AN08B098-type neurons (green) form axo-axonic connections at distinct locations along Giant Fiber (red, GF) axons. The scale bar shown is 20 µm. (*B*) The AN08B098-GF connections reconstructed in Neuroglancer mirror the morphology seen with fluorescence in *A*. (*C*) We validated AN08B098-to-GF connectivity using anti-bruchpilot to stain for T-bars in active zones (white) along AN08B098-type neurons and colocalized the resulting fluorescence with the GFs (red), which were filled with tetramethylrhodamine. The scale bar shown is 20 µm. (*D*) Neuroglancer reveals presynaptic sites at similar locations seen in colocalized fluorescent image *C*. (*E-F*) XY, and XZ plane view of the preparation shown in *A* and *B*. The zoomed-in 6 µm inlays show AN08B098 forming a single synapse with the GF in E. The yellow arrows detail the precise location AN08B098 forms the Brp-positive chemical synapse (white) to the GFs in F. Scale bars shown are 5 µm. (*G-O*) EM images showing monosynaptic connections between single GF (green) and AN08B098 neurons identified by the following MANC id: (G-H) 21041, (I) 21589, (J) 23949, (K) 152261, (L) 16900, (M) 20444, (N) 22275, and (O) 24038. Pre-and postsynaptic sites are detected in EM slices using a 3D convolutional neural network to identify T-bars (cyan dots) and postsynaptic densities[29] (PSDs, magenta dots). (*P-S*) We expressed anti-choline acetyltransferase (anti-ChAT) and anti-GFP in AN08B098 neurons. Anti-ChAT

352 colocalizes to AN08B098 cells, with particularly bright staining in the cell bodies. This finding suggests acetylcholine
353 synthesis is present within these cells, validating connectome transmitter predictions for AN08B098.
354
Following this confirmation, we investigated whether these AACs could measurably influence GF
activity in a simulation of the complete VNC connectome. To test this, we stimulated all VNC
neurons with a Poisson distribution of excitatory synaptic currents at 30 Hz and 5 nS. Additionally,
one GF was stimulated more strongly at 10 Hz and 40 nS to establish a baseline level of excitability
(Fig. 6*A, where* spikes are indicated by large black dots at the bottom of the raster plot and arrows
underneath). When the eight cholinergic Axc neurons were activated, the GF crossed threshold
more often, indicating that the axo-axonic inputs are net excitatory (Fig. 6*B*). Additional
simulations with different scenarios where stimulation and deletion of non-Axc (nAxc)
populations demonstrate that the Axc cells are paramount for increased excitability (Figs. 6*B-E*).
GF subthreshold voltage histograms quantify the increased excitability (Fig. 6*F*), supported by
statistical significance (Fig. 6*H*). These simulations highlight a previously unrecognized cohort of
axo-axonic synaptic modulation, providing a rapid mechanism for amplifying the escape-circuit
output.

368
**Figure 6. Simulations using the whole VNC network.** (*A*) Stimulation of the GF and corresponding raster plot of
all the neurons in the VNC and voltage trace of GF response (spikes are shown as large black dots at the bottom of
the raster plot, also marked with an arrow). (*B*) Schematic of the GF stimulation by Axc type neurons and
corresponding raster plot of all the neurons in the VNC and voltage trace of GF response (GF spikes are represented
by large black dots at the bottom of the raster plot, while Axc spikes are represented by large red dots). (*C*) Similar to
previous plots, but now the stimulation is driving non-Axc (nAxc), and the Axc is removed from the simulation. (*D*)

Similar to previous plots, but now the stimulation is driving Axc cells, and nAxc are deleted. (*E*) Similar to *A*, but now
Axc cells are deleted. (*F*) Histogram comparison of subthreshold voltages during GF-only and GF+Axc stimulation.
(*G*) Similar to *F*, but now the histogram shows the difference between silencing nAxc and stimulation in Axc vs.
silencing Axc and stimulation in nAxc. (*H*) Violin plots showing statistical analysis of some of the comparisons. The
overall excitability of the GF increases when the Axc neurons are activated, leading to higher firing probability for
the GF.

As the control experiment, we also tested the possibility of silencing nAxc cells, which are defined here as presynaptic cells to the GF that contribute to polysynaptic pathways, and Axc cells selectively (Figs. 6*C-D*). The results from these figures demonstrate that when nAxc connections are silenced, the increased excitability persists, and when the Axc monosynaptic connections are silenced, but nAxc is retained, the excitability is decreased. Fig. 6*G-H* shows additional quantification of this Axc to GF excitability, with the only non-significant comparison being Axc stim vs. silencing nAxc + Axc stim, demonstrating that the possibility that Axc acts polysynaptically could be neglected.

**Testing connectome predictions *in vivo* via optogenetic stimulation of axo-axonic neurons.**

To assess the predictions of these connectome simulations, we tested the ideas *in vivo* using classic electrophysiology and optogenetics. We used the split-Gal4 line that labels the Axc neurons (BDSC #89945) to drive a UAS construct containing a mVenus-tagged red-shifted channelrhodopsin, CsChrimson (BDSC #55136). Expression of CsChrimson was verified with immunohistochemistry (Fig. S2A). We electrically stimulated the GFs at 24 Hz in flies expressing CsChrimson in Axc type neurons and recorded from the jump muscle (TTM) and flight muscle (DLM) as previously described[30], (Fig. 7*A-B*). We then set the electrical stimulation of the GF just below threshold. This near-threshold stimulation resulted in occasional (~20%) responses in both muscles (n = 5 specimens, Fig. 7*B* and 7*E*). However, when the same preparation was illuminated (590 nm, 0.37mW) during the stimulus train, the sub-threshold electrical stimulus elicited responses from both muscles nearly 100% of the time (n = 5 specimens, Fig. 7*D* and 7F). A one-way ANOVA found significant differences between responses before, during, and after light application ($p < 0.001$), but there was no significant difference for the control group ($p = 0.53$) (Fig. 7*G*). Post-hoc multiple comparisons revealed significant differences between responses at baseline (before light application) and during and after light application. We also found significant differences using a paired t-test for the same responses during and after light application ($p<0.001$), but not before ($p>0.26$).

**Figure 7. Optogenetic activation of the Giant Fiber system.** (*A*) Sketch of the electrophysiological recordings with
optogenetic activation of axo-axonic cells (Axc). (*B*) Example of raw traces for control (no light application). A
zoomed-in panel is presented, showing a difference in latencies of ~0.6 ms between TTM and DLM onset responses.
(*C*) Simulation of a computational model of the VNC connectome emulating the experimental results under
subthreshold conditions and light stimulation. Note that GF only spikes when the light is applied and is time-locked
with the electrical pulses. Action potentials are indicated with arrows. (*D*) Example of raw traces when light was
applied. A zoomed-in panel is presented, showing difference in latencies of ~0.6 ms between TTM and DLM onset
responses. (*E*) Spike probability per fly (n = 5) under no-light conditions. (*F*) Spike probability per fly (n = 5) under
light conditions (light was applied during the blue shaded region). (G) Average and SEM of spike probability (paired
t-test, * = $p$<0.001). **Spike initiation in the VNC**. (*H*) Sketch of the electrophysiological recordings with optogenetic
activation of the GF before and after decapitation. (*I*) Recordings from both TTM and DLM before decapitation. The
constant latency between TTM and DLM indicates the presence of GF action potentials. Traces indicate that both
muscles respond to the light. A zoomed-in panel shows traces with different speeds by checking the new constant
latency between TTM and DLM (0.77 ms, 0.81 ms, and 0.91 ms). (*J*) Response of the TTM to optical stimulation in
a specimen driving CsChrimson in the GF as above. The blue curve represents the response of the TTM in animals
with the head on, exhibiting a long tail of post-light response. (*K*) The black curve is after the head was removed to
remove the spike-initiating zone in the brain. The spiking behavior persists despite decapitation.
429
To verify that light stimulation does not elicit effects in the absence of CsChrimson expression or
function, we performed two control experiments. In the first experiment, flies expressed GFP
instead of CsChrimson (Fig. S2A); in the second, CsChrimson-expressing flies were raised without
dietary cis-trans-retinal, thereby preventing optogenetic activation (Fig. S2*B*). A one-way
ANOVA revealed no significant differences in stimulus-evoked responses across conditions (*p* =
1). Post-hoc comparisons revealed no significant differences between the baseline, light, and post-
light periods. Paired t-tests also found no significant differences between light and no-light
conditions during and after stimulation. These results confirm that the observed effects are
specifically attributable to CsChrimson activation.
439
The experiment in Fig. 7 demonstrates that the classic method of GF stimulation using extracellular
electrodes in the brain can be employed to assess the role of the Axcs in controlling GF excitability.
We do not directly monitor GF spikes but instead use ~~the~~ TTM responses as a reporter. The GF
extracellular stimulation activates both the TTM and the DLM with a short and constant latency
between them (~0.6 ms) as described in the early recordings of this type in the 1980s[32–34]. Action
potentials in the GF are indicated by synaptic potentials in both the TTM and the DLM as described
in a more recent protocol for the preparation[30]. It has long been assumed that the GF has the lowest
threshold for brain stimulation due to its large size and the presence of a spike initiation region in
the brain. Recently, a spike initiation zone (SIZ) has been confirmed to be located in the brain at
the dendrite-axon bifurcation[35].
450
How the Axc alters excitability depends in part on the site of spike initiation for the GF. In the
present experiments, the stimulating electrodes are in the brain where there are no monosynaptic
connections from Axc to GF[25]. The only monosynaptic Axc-to-GF connections are in the VNC
(Fig. 5, S3). How can the optical excitation of the Axcs in the VNC, where they make synaptic

contact, have a modulatory effect on the GF spike initiation zone (SIZ) in the brain, where they
do not make contact? We have now demonstrated that the GF also possesses a spike-initiating
region(s) in the VNC, where the Axc could influence spike initiation. We demonstrated this
alternative SIZ by driving CsChrimson in the GF, removing the head to remove the SIZ in the
brain, and testing headless animals with optical stimulation to reveal GF activity generated in the
VNC (Fig. 7J-*K*). GF activity driven by the ~~combined electrical and~~ light stimulation is revealed
by the response of both TTM and DLM with a constant latency between them indicative of GF
action potentials (Fig. 7*K*). Therefore, the Axcs can directly influence this SIZ in the VNC.
463
Unexpectedly, we also note a prolonged response of the GF after the light was turned off (Fig. 7D,
7F and 7G). Before light application, the TTMn response probability was low (~20%, black trace,
Fig. 7*G*), during light application the probability increased to almost 100% (blue trace, Fig. 7*G*)
and after light-off activity returned to baseline very slowly (>100 msec). This prolonged response
after light-off has been observed in several other situations for CsChrimson. For instance, at the
larval neuromuscular junction (NMJ), CsChrimson drove spiking in motor neurons (monitored by
muscle synaptic potentials) and exhibited a post-light response hundreds of milliseconds in
duration[31]. In a second example, descending neurons DNp07 and DNp10, when expressing
CsChrimson, exhibit a 100-400 msec response after light-off[36]. Finally, we tested this idea in the
GF by expressing CsChrimson in the GF and illuminating the preparation at 590 nM and observed
a long tail of TTM response of approximately 100 msec (Fig. 7*J,* blue trace). In brief, a variety of
examples show that CsChrimson when activated produces a strong response with a long slow
decay after the light is turned off and this accounts for the prolonged response seen in Fig 7*F-G*.
477
To confirm the parallel between the simulation and the *in vivo* experiment, we also developed a
simulation that more closely matched the *in vivo* experiments with optogenetics (Fig. 7*C*). A large-
scale network of the VNC, composed of leaky integrate-and-fire neuron models, was used to
simulate this experiment. The GF was electrically stimulated just below threshold at 24 Hz. In
these simulations, the GF only spikes in the presence of the light. Both the simulation and the
optogenetic results on the living animal suggest that the Axc-to-GF connection is excitatory,
consistent with the connectome and chemical predictions that the synapse is cholinergic. These
results agree with our simulation's predictions that the main effect of the optogenetic activation of
the cholinergic axo-axonic neurons (Axcs) is to increase the excitability of the GF (Fig. 7*C*).
487
488  Overall, these results support our prediction that the primary effect of optogenetic activation of the
489  cholinergic axo-axonic neurons (Axcs) is to increase the excitability of the GF. Our conclusion is
490  that the optically activated Axcs excite the GF via their synapses in the VNC and alter the
491  likelihood of GF action potentials.
492

**Discussion**

By analyzing all 1,314 brain-originating descending neurons, we move beyond scattered descriptions of axo-axonic contacts and derive circuit-level design rules for presynaptic modulation. The quantitative principles that emerged within this work are: (i) extreme sparseness showing that between 0.7 to 1.2% of possible DN-to-DN, AN-to-DN, and IN-to-DN pairs form axo-axonic synapses; (ii) a tight linear relation between synaptic strength and partner multiplicity; and (iii) a small-world architecture that distributes integration rather than concentrating it in a rich-club core. Together, these features constitute a wiring grammar for axo-axonic control in the adult *Drosophila* motor system.

Axo-axonic synapses have been identified in the *Drosophila brain*[37–39]. Excitatory cholinergic synapses, like those investigated here, have been observed in the fly olfactory system, where projecting neurons (PN) form lateral PN-to-PN connections, similar to the DN-to-DN connections we identified[37]. Axo-axonic connections have also been reported between olfactory sensory neurons (OSNs)[39]. Bristle mechanosensory neurons (BMNs) are also shown to be axo-axonically connected[40]. These similarities suggest that axo-axonic connectivity principles might be conserved across the VNC and brains in flies, although the functional role of AACs remains unexplored.

Notably, axo-axonic wiring has revealed an extraordinarily selective, yet globally percolating schema, in contrast to, for example, the serotonin axo-axonic configuration in the hippocampus, which comprises 30%[10,41] or in the cortex, which comprises 19%[10,42]. In the fly VNC, only ~1 % of the possible DN-to-DN pairs form axo-axonic synapses, an order of magnitude sparser than axo-dendritic contacts in either the MANC or FAFB brain data sets[14,43]. Despite their scarcity, these connections are sufficient to incorporate every DN into a single, strongly connected network, yielding short average path lengths (~4 hops) and thus preserving the ability for any axon to gain rapid, monosynaptic influence over remote peers. For comparison, the entire network has an average path length of 2.8 (~3 hops). This combination of stringent partner choice with network-wide reach resembles presynaptic inhibition motifs in mammalian spinal cord and chandelier-cell control of cortical pyramidal neurons, suggesting convergent architectural solutions across phyla[44,45]. What accounts for the sparsity of axo-axonic connections? One possibility is wiring economy: boutons on sister axons consume precious membrane and cytoplasmic resources[46]. Another is developmental lineage. Many high-degree bridge neurons derive from hemilineages that extend across neuromeres[47], hinting that growth-cone timing and Slit-Robo–mediated midline crossing[48] pre-pattern the opportunities for axo-axonic contact. Comparative EM work in other insects and in zebrafish larval spinal cord will clarify whether these archetypes represent a universal motor blueprint.

As previously found, rich clubs with dense interlinking of high-degree hubs are present in the fruit *Drosophila* connectome[16]. They are also widely present in vertebrates, such as in the cerebral cortex[49,50]. In contrast, the axo-axonic fly DN network revealed only a weak, transient rich-club

coefficient, suggesting that presynaptic modulation in the nerve cord is decentralized. Control must therefore emerge from many mid-degree "broker" neurons (with fewer connections) and with high betweenness rather than from a minority of super-hubs. Such a layout guards against single-point failure and allows different motor modules to be gated independently, consistent with behavioral data showing that flies can blend local reflexes with whole-body maneuvers[51].

In terms of functional implications, axo-axonic contacts positioned directly on descending axons are placed ideally to veto, amplify, or synchronize commands emanating from the brain before they fan out onto premotor interneurons and motoneurons. Our simulations demonstrate this principle: activating a cohort of eight cholinergic Axc type neurons, which exclusively target the Giant Fibers (GFs), accelerates their firing. Choline acetyltransferase (ChAT) immunostaining confirms that Axc type neurons are cholinergic, and optogenetic activation of these cells reliably enhances the GF's excitability, confirming the large-scale network of leaky-integrate-and-fire neuron simulations.

Our experimental evidence from optogenetic stimulation of the GF suggests that action potentials generated in the GF within the VNC propagate through the TTM via electrical synapses, causing them to fire simultaneously. These action potentials then propagate to the PSI through electrical synapses and via chemical synapses, inducing an action potential through the DLM. Finally, we observe the response of the muscles in both DLM and TTM, noting phase-locked responses with a delay consistent with the synaptic pathway. We repeated these experiments in headless animals to demonstrate that the source of the action potential could count for the VNC directly, suggesting the existence of a second spike initiation zone (SIZ) besides the already well-known one in the brain. Admittedly, in view of the AN08B098 synapses on or near the GF synaptic terminal bend, another possibility is that they elicit a subthreshold depolarization in it that is enough to pass current via its electrical output synapses with TTMn, thus stimulating the latter to threshold without eliciting retrograde spikes in the GF. In this case, a TTM action potential would not be an infallible readout of a GF spike, as is conventionally assumed, merely a readout of depolarizing synaptic current in the GF terminal. This latter hypothesis has not been tested in our experiments and is worth future experimentation.

Crucially, however, the GF is not unique; 17 other DNs receive >50 presynaptic partners that occupy bridge positions between functional modules. These neurons may be promising targets for future studies investigating the role of axo-axonic connections in precise behavior modulation. Future opto-physiological experiments probing these DNs will test whether distributed control is the norm rather than the exception.

Excitatory axo-axonic connections, exemplified by the Axc to GF projection described here, are not unique to this system and have been reported in other organisms, including the well-characterized Mauthner cells in zebrafish[12] and dopamine cells in mice[53]. In these examples, the

573   primary function of the excitatory axo-axonic synapse is to increase the release probability of the
574   axon terminal they innervate. This mechanism can even bypass upstream action potentials,
575   representing a hijacking mechanism that may have physiological roles. The findings presented in
576   this study may contribute to a broader understanding of axo-axonic connectivity principles across
577   species.
578
One limitation of our study is that the connectome used in the analysis was derived from a single
male specimen. New datasets are now being released that are sex-specific or can address
developmental differences, such as the Female Adult Nerve Cord Connectome (FANC) and the
Brain and Nerve Cord Connectome (BANC). All functional whole-network tests are *in silico*, and
although experimental verification is provided through optogenetics, the large-scale network based
on leaky integrate-and-fire neuron model respects synapse counts but ignores conductance
dynamics and neuromodulators influences. Furthermore, EM of the MANC connectome (v1.2.1)
did not detect gap-junctional coupling, suggesting that the present map under-reports electrical
axo-axonic contacts that could further reshape information flow[54–58].
588
Lastly, dendro-axonic synapses account for only 2% of all synapses[24,59]. Thus, we cannot rule out
the possibility that some of the synapses investigated in this study are dendro-axonic, although the
vast majority are axo-axonic. Nevertheless, we curated a large number of EM images (Fig. S1) to
demonstrate the presence of mitochondria in axo-axonic synapses of descending neurons,
suggesting that this structure can be used as a marker of axo-axonic synapses. Based on this, we
found that synapses of AN08B098 onto Giant Fiber contain presynaptic mitochondria, suggesting
that these synapses are axo-axonic.
596
In conclusion, our study presents the first comprehensive blueprint of axo-axonic connectivity
within a fully reconstructed adult ventral nerve cord system. The network revealed here offers
concrete parameters for next-generation models of rapid motor decision making in both
invertebrates and vertebrates. By formalizing direct axon-to-axon communications, the work
bridges a critical gap between connectome structure and motor function and opens new avenues
for dissecting the fastest circuits in the animal kingdom.
603
604
**Methods**
**Data acquisition and analysis**
Data for VNC neurons used in the study came from the publicly available *Drosophila* male adult
nerve cord connectome[14]. NeuPrint+ API[25,27] access is granted to registered users, and Python
accessibility is maintained through the neuPrint-python package[26]. The latest *Drosophila* MANC
data can be accessed here: https://www.janelia.org/project-team/flyem/manc-connectome.
611
**Connectivity analysis**

We retrieved all simple synaptic connections using neuprint.fetch_simple_connections[25–27]
separately for: outputs (presynaptic AN, IN, and DN as presynaptic) and Inputs (postsynaptic DN).
Connections were summed by partner neuron and filtered to those with total strength ≥ 6. For each
DN, we then counted the number of distinct AN, IN, and DN partners in both the output and input
directions.
618
**Network analysis**
Network analysis was performed using Python with the NetworkX[60] library to extract quantitative
measures of connectivity. From the structural connectivity matrix, we calculated a comprehensive
set of graph-theoretical metrics to characterize the network topology. These included basic
properties (number of nodes and edges, network density), integration measures (average path
length, diameter), segregation indices (clustering coefficient, modularity with community
detection using the Louvain algorithm), centrality metrics (degree, betweenness, closeness, and
eigenvector centrality), and organizational principles (degree assortativity, rich club coefficient).
627
The rich club analysis was specifically conducted across multiple degree thresholds to identify
potential hierarchical organization among high-degree hub regions. All weighted graph measures
incorporated the synaptic strengths from the original connectivity matrix. This multimodal graph-
theoretical approach allowed for a detailed characterization of both local and global properties of
the network's structural connectome, revealing important aspects of neural communication
efficiency, functional segregation, and network resilience.
634
**Statistical and visual analysis**
SWC analysis was conducted in Python using the Navis client[61]. We stored morphology and
input/output counts using the Pandas library. Relationships between morphology metrics (cable
length, branch points, node count) and partner counts were assessed with Pearson correlation and
linear regression (scipy.stats.pearsonr, linregress). Results were visualized with Matplotlib[62].
Network-style connection diagrams were rendered interactively with ipycytoscape.
641
**Software and libraries**
NeuPrint+ analyses were performed in Python 3.12.4 using the neuprint-python API. We used the
Janelia NeuPrint client (neuprint), and Navis (v1.10.0) for data access and network building; data
manipulation and grouping were done with pandas v2.2.2 and NumPy v1.26.4; graph-based
operations with NetworkX v3.2.1; interactive visualization with ipycytoscape v1.3.3 and
ipywidgets v8.1.5; plotting and PDF export with Matplotlib v3.10.1; and statistical analyses
(Pearson's r, linear regression) with SciPy[63] v1.13.1. Auxiliary parsing and utility functions relied
on Python's built-in json, re, itertools, and warnings modules.
650
**Computational model**

To study the impact of axo-axonic projections onto DNp01 (GF), we constructed an *in-silico* model
of the ventral nerve cord (VNC) by utilizing the entire MANC v1.2.1 connectome. First, we
scrubbed the connectivity data by omitting connections with a total strength less than 6, then we
excluded all connectivity data where the neurotransmitter type, neuron type, or neuron ID was
unknown or invalid. This resulted in a total of 23,437 valid neurons and 1,152,548 connections
between them, with 619,684 excitatory (cholinergic) and 532,864 inhibitory (glutamatergic or
GABAergic) connections. In the fly, cholinergic receptors are excitatory, whereas glutamatergic
and GABAergic receptors are inhibitory[64].

**Neuron model**
To achieve biological plausibility while avoiding computational intractability, leaky integrate-and-
fire (LIF) neuron models were used:

664
$$C_m \frac{dv}{dt} = -g_L(v - V_{rest}) - g_{ach}(v - E_{ach}) - g_{gaba}(v - E_{gaba}) - g_{glut}(v - E_{glut}), \quad (1)$$

Unlike other models[65], we employ distinct parameters for the GF compared to the rest of the
neurons in the connectome, i.e., the membrane resistance. In Eq. (1), $v$ is the neuron membrane
potential, $g_L$, $g_{ach}$, $g_{GABA}$, and $g_{glut}$ are the conductances for leak channels and acetylcholine
(nicotinic), GABA (GABA$_A$), and glutamate ionotropic receptors (nAChR, GABA$_A$R, and GluCl,
respectively). Similarly, $V_{rest}$, $E_{ach}$, $E_{GABA}$, and $E_{glut}$ are their respective reversal potentials.
Since GABA$_A$R and GluR are chloride-permeable channels, their reversal potentials were obtained
from chloride reversal potential[66]. $C_m$ is the membrane capacitance. These are parameterized with
values obtained from the literature, which investigates the biophysical properties of fruit fly
synapses. Furthermore, we collected physiologically relevant values for the membrane
capacitance, resistance, and leak reversal potential. Simulations of these neurons were made using
the BRIAN2 simulator [67].

**Synapse model**
For each of the synaptic currents ($x = ach, gaba, glut$), the conductance obeys

681
$$\frac{dg_x}{\tau_x} = -g_x, \quad\quad (2)$$

682    which increases by $J_x$ upon a presynaptic spike. The list of parameters is included in Table 1.

| Parameter | Symbol | Value | Ref. |
|---|---|---|---|
| Acetylcholine receptor (nAChR) conductance | $J_{ach}$ | 0.27 nS | 35 |

| | | | |
|---|---|---|---|
| nAChR decay time | $\tau_{ach}$ | 1.1 ms | [35] |
| nAChR reversal potential | $E_{ach}$ | 0 mV | [68] |
| GABA receptor (GABA$_A$R) conductance | $J_{GABA}$ | 0.8 nS | Estimated |
| GABA$_A$R decay time | $\tau_{GABA}$ | 5.4 ms | [69] |
| GABA$_A$R reversal potential | $E_{GABA}$ | -60 mV | [64,66] |
| Glutamate receptor (GluCl) conductance | $J_{glut}$ | 0.8 nS | Estimated |
| GluCl decay time | $\tau_{glut}$ | 5 ms | [70] |
| GluCl reversal potential | $E_{glut}$ | -60 mV | [66] |
| Membrane Resistance (GF) | $R_m$ | 25 MOhm | [35] |
| Membrane Resistance (other neurons) | $R_m$ | 300 MOhm | [71] |
| Resting membrane potential | $V_{rest}$ | -55 mV | [65,71] |
| Action potential threshold | $V_{th}$ | -40 mV | [71] |
| Reset potential | $V_{reset}$ | -55 mV | Estimated |
| Refractory period | $\tau_{ref}$ | 2 ms | Estimated |
| Membrane capacitance | $C_m$ | 72 pF | [71] |

**Table 1.** List of parameters used in the connectome simulation of the VNC in this paper. For each of our simulations, we ran the entire VNC network with a step-size of 0.1 ms during 2000 ms. Furthermore, for each simulation, each neuron in the VNC network was provided with an excitatory Poisson input of 30 Hz at a synaptic conductance of 5 nS to simulate typical endogenous activity driven by neurons in the fly brain. During stimulation experiments, neurons DNp01, DNg02, and/or DNd03 were stimulated with additional Poisson input at 10 Hz at a synaptic conductance of 40 nS. In Figure 8, the same network was used, but the GF was electrically stimulated just below threshold with a Poisson-distributed stimulation at 24 Hz.

**Fly Genetics**

Flies were raised at 25°C with a natural (12-hr) light cycle, and data were collected from both male and female adults 2-4 days old. The Gal4 drivers used in our study were selected through NeuronBridge[72]. Neurons of interest were identified through NeuPrint+, then identified using NeuronBridge software v3.2.6, and finally chosen based on similarity score and visual accuracy. A split-Gal4 driver specific to AN08B098 (Axc) was identified using NeuronBridge software v3.2.6 and chosen based on similarity score. SS25451 contains w[1118]; VT063643-p65.AD; VT059224-GAL4.DBD (BDSC #89495) (see Fig S4). Constructs used were 20XUAS-IVS-CsChrimson.mVenus (#55136), and 20XUAS-IVS-mCD8::GFP (BDSC #32194).

**Electrophysiology**

Flies were mounted dorsal side up in dental wax after $CO_2$ anesthesia. Muscle recordings were
obtained using glass microelectrodes placed directly into both left and right tergotrochanteral jump
muscles[18,30,55,58,73] (TTM). Microelectrodes were filled with O'Dowd's Drosophila saline[74].
Stimulation of the Giant Fibers was performed using 3-4 V applied extracellularly using Grass
stimulators with tungsten wire electrodes placed in the brain through the eyes, while a tungsten
wire electrode was placed in the fly's abdomen to ground the circuit[30]. The Axon Digidata 1440A
Data Acquisition System was used to digitize recordings which are then visualized using Clampex
software (version 10 & 11, Molecular Devices). Muscle responses were obtained by stimulating
at 24 Hz.

**Optogenetics**
We drove channelrhodopsin expression in neurons of interest using the UAS-Gal4 system. Flies
expressing csChrimson (BDSC #55136) were crossed with our driver line to produce: w1118;
VT063643-p65/+; 20XUAS-IVS-csChrimson.mVenus/VT059224-GAL4.DBD. Adults were
placed in the dark at 50% relative humidity on standard cornmeal-molasses food, supplemented
with all-trans-retinal (0.4 mM post-eclosion) for two days before testing. Light stimulation was
performed for 100 ms while flies are mounted in dental wax as described above. We used a 590
nm high-power fiber-coupled LED (1.5 mW maximum light intensity, Thorlabs M590F3) with an
optic fiber of diameter 200 μm and N.A. 0.22 to activate the mVenus csChrimson (M118L02). We
tested various placements for the optic fiber and found the strongest activation of csChrimson to
occur when the fiber was less than 1 cm from the sample and aimed at the neck. Light intensity
was controlled via T-Cube LED Driver (Thorlabs LEDD1B) and kept at maximum power during
testing. Each sample received electrical-only, optogenetic-only or paired opto-electrical
stimulation. If Axc is stimulated only, no response is recorded, meaning that csChrimson alone is
unable to elicit spikes in the GF.

We also conducted two control experiments for the optogenetic protocol (Fig. S2). The first control
used flies of the same genotype as the experimental group but were not raised on food
supplemented with all-trans-retinal. The second control involved flies that were raised on retinal-
supplemented food but lacked the UAS-csChrimson.mVenus transgene (w1118; VT063643/+;
VT059224-GAL4/VT059224-GAL4). The results are shown in Figure S2 B-C.

**Dye-injection**
The VNC and connective of young adult male flies were dissected and kept in cold O'Dowd's
saline within a Sylgard 184-filled dissection dish. Samples were immediately transferred to a
Nikon Eclipse E600FN fluorescent microscope. A tetramethylrhodamine-labeled dextran dye
(10,000MW, Invitrogen, D1817) solution was backfilled into a pulled glass electrode tip (~30 MΩ
resistance tip) and filled with O'Dowd's saline. Electrodes were placed into a 3D hydraulic
micromanipulator. The electrode was lowered into place, penetrating the GF axons near the neck
connective. A Gettings Model 5A amplifier was used to inject positive current pulses (between 1

nA and 5 nA), with a reference electrode placed into the saline-filled dish. GFs were filled with
dye for approximately 5 minutes each.

**Immunostaining**

A mouse primary antibody against ChAT (1:20, DSHB, 4B1) was used to confirm cholinergic
cells and coupled to secondary Goat anti-mouse AlexaFluor 647 (1:500, 115-605-003). A mouse
primary antibody against bruchpilot (1:50, DSHB, NC82) was used to label presynaptic chemical
active zones and coupled to secondary Goat anti-mouse AlexaFluor 647 (1:500, 115-605-003).
Primary rabbit anti-GFP (1:500, Invitrogen A11122) and secondary goat anti-rabbit (1:500,
Jackson Immuno, 111-225-144) were used to amplify the fluorescence of mCD8::GFP. mVenus-
tagged csChrimson. Fluorescence was amplified using a conjugated antibody (1:500, Millipore,
MABE1906). Primary antibody solutions contained 3% BSA and 0.3% Triton X-100 in PBS. The
secondary antibody solutions contained PBS.

Samples were dissected and processed according to the dye-injection methods above. Samples
were then immediately fixed in PBS + 4% PFA for 25 minutes at room temperature (22°C).
Samples were washed with PBS and placed on a rotator (60RPM) for 30 minutes. Samples were
permeabilized with PBS + 0.5% Triton X-100 for 45 minutes at 22°C. Primary antibodies were
added, and samples were stored at 4°C for 48 hours. Samples were then washed on a rotator with
PBS for 4 hours at 22°C, changing the solution every 30 minutes. Secondary antibodies were
added, and samples were stored at 4°C for 24 hours. Samples were then washed on a rotator with
PBS for 4 hours at 22°C, changing the solution every 30 minutes. Samples were dehydrated with
a series of ethanol steps, 50%, 70%, 90%, and 100% for 8 minutes each step. Samples were
trimmed and embedded in 100% methylsalycylate, covered with cover slips, sealed, and imaged
within 24 hours.

**Confocal Microscopy**

Samples were imaged on a Nikon AXR confocal using an NSPARC detector and a 60x oil
immersion objective. Z-dimension step size was 194 μm for all results displayed.  Laser channels
used were 488 nm, 561 nm, and 640 nm. Channels were imaged in series using filters to avoid
cross-excitation of close wavelengths, and 2x integration was used.

**Image Analysis**

Z-stack files were imported into Imaris software v10.2.0, and the fluorescence intensity of each
channel was adjusted for best fit. GF and AN08B098 (Axc) channels were surfaced using a
thresholding method that matched fluorescence in each Z dimension. Masks of each channel were
created for each surface to detect colocalization of Brp or ChAT. Images were acquired in volume
view and exported.

**Acknowledgments**

R.P. was funded by an IBRO Collaborative Research Grants. The authors thank the FAU Jupiter
Life Sciences Initiative (JLSI) for funding.
785

**Author contributions**
C.C.C. and R.P. were responsible for the conceptualization of the work. C.C.C., J.L., T.R., D.S.,
S.J., and C.S. analyzed the data and made the figures. R.P. worked on the graph theoretical
measures. D.S. performed and analyzed the electrophysiology recordings. S.J. and C.S. performed
immunofluorescence. K.R. maintained fly stocks and set up experiment crosses. T.R. built and ran
the network simulations. R.P., C.S., and R.M. supervised the research and secured funding. All
authors participated in the writing, reviewed, and approved the manuscript.

**Declaration of interests**
The authors declare no competing interests.

## References

1. Brette, R. Theory of axo-axonic inhibition. *PLoS Comput Biol* 21, e1013047 (2025).

2. Schneider-Mizell, C. M. *et al.* Structure and function of axo-axonic inhibition. *Elife* 10, (2021).

3. Stretton, A. O. W. *et al.* Structure and physiological activity of the motoneurons of the nematode Ascaris. *Proc Natl Acad Sci U S A* 75, (1978).

4. Pearson, K. G. & Goodman, C. S. Presynaptic inhibition of transmission from identified interneurons in locust central nervous system. *J Neurophysiol* 45, (1981).

5. Atwood, H. L. & Kwan, I. Reciprocal axo-axonal synapses between excitatory and inhibitory neurons in crustaceans. *Brain Res* 174, (1979).

6. Altrup, U., Speckmann, E. J. & Caspers, H. Differentiation between anti- and orthodromic responses to nerve stimulation in neurons with axo-axonal synapses (Helix pomatia). *Neurosci Lett* 11, (1979).

7. Andersen, P., Eccles, J. C., Schmidt, R. F. & Yokota, T. Depolarization Of Presynaptic Fibers In The Cuneate Nucleus. *J Neurophysiol* 27, (1964).

8. Wendell-Smith, C. P., Blunt, M. J., Baldwin, F. & Paisley, P. B. Neurone - Satellite cell relationship. *Nature* 205, (1965).

9. Gobel, S. & Dubner, R. Fine structural studies of the main sensory trigeminal nucleus in the cat and rat. *Journal of Comparative Neurology* 137, (1969).

10. Cover, K. K. & Mathur, B. N. Axo-axonic synapses: Diversity in neural circuit function. *Journal of Comparative Neurology* 529, 2391–2401 (2021).

11. Dudok, B. *et al.* Recruitment and inhibitory action of hippocampal axo-axonic cells during behavior. *Neuron* 109, (2021).

12. Lacoste, A. M. B. *et al.* A Convergent and Essential Interneuron Pathway for Mauthner-Cell-Mediated Escapes. *Current Biology* 25, 1526–1534 (2015).

13. Takemura, S.-Y. *et al.* A Connectome of the Male \textit{Drosophila} Ventral Nerve Cord. *bioRxiv* (2023).

14. Cheong, H. S. J. *et al.* Transforming descending input into behavior: The organization of premotor circuits in the Drosophila Male Adult Nerve Cord connectome. *bioRxiv* (2023).

15. Marin, E. C. *et al.* Systematic annotation of a complete adult male Drosophila nerve cord connectome reveals principles of functional organisation. *Elife* 13, (2024).

16. Lin, A. *et al.* Network statistics of the whole-brain connectome of Drosophila. *Nature 2024 634:8032* 634, 153–165 (2024).

17. Clements, J. *et al.* neuPrint: Analysis Tools for EM Connectomics. *bioRxiv* 2020.01.16.909465 (2020) doi:10.1101/2020.01.16.909465.

18. Allen, M. J., Godenschwege, T. A., Tanouye, M. A. & Phelan, P. Making an escape: Development and function of the Drosophila giant fibre system. *Seminars in Cell and Developmental Biology* vol. 17 Preprint at https://doi.org/10.1016/j.semcdb.2005.11.011 (2006).

19. Asinof, S. K. & Card, G. M. Neural Control of Naturalistic Behavior Choices. *Annu Rev Neurosci* 47, 369–388 (2024).

20. Cheong, H. S. J. *et al.* Transforming descending input into behavior: The organization of premotor circuits in the Drosophila Male Adult Nerve Cord connectome. *bioRxiv* (2023).

21.  Stürner, T. *et al.* Comparative connectomics of Drosophila descending and
ascending neurons. *Nature 2025* 1–15 (2025) doi:10.1038/s41586-025-08925-z.
22.  Takemura, S. *et al.* A Connectome of the Male Drosophila Ventral Nerve Cord.
*Elife* 13, (2024).
23.  Marin, E. C. *et al.* Systematic annotation of a complete adult male Drosophila nerve
cord connectome reveals principles of functional organisation. *Elife* 13, (2024).
24.  Schneider-Mizell, C. M. *et al.* Quantitative neuroanatomy for connectomics in
Drosophila. *Elife* 5, (2016).
25.  Plaza, S. M. *et al.* neuPrint: An open access tool for EM connectomics. *Front*
*Neuroinform* 16, (2022).
26.  Berg, S. & Schlegel, P. neuprint-python. Preprint at
https://doi.org/10.5281/zenodo.7592899 (2017).
27.  Clements, J. *et al.* neuPrint: Analysis Tools for EM Connectomics. *bioRxiv*
2020.01.16.909465-2020.01.16.909465 (2020) doi:10.1101/2020.01.16.909465.
28.  Bates, A. S. *et al.* Distributed control circuits across a brain-and-cord connectome.
Preprint at https://doi.org/10.1101/2025.07.31.667571 (2025).
29.  Huang, G. B., Scheffer, L. K. & Plaza, S. M. Fully-automatic synapse prediction
and validation on a large data set. *Front Neural Circuits* 12, (2018).
30.  Allen, M. J. & Godenschwege, T. A. Electrophysiological recordings from the
Drosophila giant fiber system (GFS). *Cold Spring Harb Protoc* 5, (2010).
31.  Klapoetke, N. C. *et al.* Independent optical excitation of distinct neural populations.
*Nat Methods* 11, (2014).
32.  Tanouye, M. A. & Wyman, R. J. Motor outputs of giant nerve fiber in Drosophila. *J*
*Neurophysiol* 44, (1980).
33.  Tomas, J. B. & Wyman, R. J. Mutations altering synaptic connectivity between
identified neurons in Drosophila. *Journal of Neuroscience* 4, (1984).
34.  Thomas, J. B. & Wyman, R. J. A mutation in Drosophila alters normal connectivity
between two identified neurones. *Nature* 298, (1982).
35.  Moreno-Sanchez, A. *et al.* Morphology and synapse topography optimize linear
encoding of synapse numbers in Drosophila looming responsive descending
neurons. Preprint at https://doi.org/10.7554/eLife.99277.1 (2024).
36.  Ache, J. M. *et al.* Neural Basis for Looming Size and Velocity Encoding in the
Drosophila Giant Fiber Escape Pathway. *Current Biology* 29, (2019).
37.  Scheffer, L. K. *et al.* A connectome and analysis of the adult drosophila central
brain. *Elife* 9, (2020).
38.  Manoim, J. E., Davidson, A. M., Weiss, S., Hige, T. & Parnas, M. Lateral axonal
modulation is required for stimulus-specific olfactory conditioning in Drosophila.
*Current Biology* 32, 4438-4450.e5 (2022).
39.  Gruber, L. *et al.* The unique synaptic circuitry of specialized olfactory glomeruli in
Drosophila melanogaster. Preprint at https://doi.org/10.7554/eLife.88824.1 (2023).
40.  Calle-Schuler, S. A., Santana-Cruz, A. E., Kmecová, L., Hampel, S. & Seeds, A.
M. A comprehensive mechanosensory connectome reveals a somatotopically
organized neural circuit architecture controlling stimulus-aimed grooming of the
*Drosophila* head. Preprint at https://doi.org/10.1101/2025.05.19.654894 (2025).
41.  Oleskevich, S., Descarries, L., Watkins, K. C., Se´gue´la, P. & Daszuta, A.
Ultrastructural features of the serotonin innervation in adult rat hippocampus: An

888              immunocytochemical description in single and serial thin sections. *Neuroscience*
889              42, 777–791 (1991).

42. Séguéla, P., Watkins, K. C. & Descarries, L. Ultrastructural relationships of serotonin axon terminals in the cerebral cortex of the adult rat. *Journal of Comparative Neurology* 289, 129–142 (1989).

43. Lin, A. *et al.* Network statistics of the whole-brain connectome of Drosophila. *Nature 2024 634:8032* 634, 153–165 (2024).

44. Lu, J. *et al.* Selective inhibitory control of pyramidal neuron ensembles and cortical subnetworks by chandelier cells. *Nat Neurosci* 20, (2017).

45. Woodruff, A. R., Anderson, S. A. & Yuste, R. The Enigmatic Function of Chandelier Cells. *Front Neurosci* 4, (2010).

46. Cuntz, H., Forstner, F., Borst, A. & Häusser, M. One rule to grow them all: A general theory of neuronal branching and its practical application. *PLoS Comput Biol* 6, (2010).

47. Lacin, H. & Truman, J. W. Lineage mapping identifies molecular and architectural similarities between the larval and adult Drosophila central nervous system. *Elife* 5, (2016).

48. Rajagopalan, S., Nicolas, E., Vivancos, V., Berger, J. & Dickson, B. J. Crossing the midline: Roles and regulation of Robo receptors. *Neuron* 28, (2000).

49. Harriger, L., van den Heuvel, M. P. & Sporns, O. Rich Club Organization of Macaque Cerebral Cortex and Its Role in Network Communication. *PLoS One* 7, e46497 (2012).

50. Liang, X. *et al.* The Rich-Club Organization in Rat Functional Brain Network to Balance Between Communication Cost and Efficiency. *Cerebral Cortex* 28, 924–935 (2018).

51. Card, G. & Dickinson, M. H. Visually Mediated Motor Planning in the Escape Response of Drosophila. *Current Biology* 18, (2008).

52. Dombrovski, M. *et al.* Synaptic gradients transform object location to action. *Nature* 613, (2023).

53. Kramer, P. F. *et al.* Synaptic-like axo-axonal transmission from striatal cholinergic interneurons onto dopaminergic fibers. *Neuron* 110, 2949-2960.e4 (2022).

54. Ammer, G., Vieira, R. M., Fendl, S. & Borst, A. Anatomical distribution and functional roles of electrical synapses in Drosophila. *Current Biology* 32, (2022).

55. Lopez, J., Boerner, J., Robbins, K., Pena, R. F. O. & Murphey, R. Frazzled/DCC regulates gap junction formation at a Drosophila giant synapse through transcription. Preprint at https://doi.org/10.1101/2025.04.08.647628 (2025).

56. Holcroft, C. E. *et al.* Innexins Ogre and Inx2 are required in glial cells for normal     postembryonic development of the Drosophila central nervous system. *J Cell Sci* 126, 3823–3834 (2013).

57. Phelan, P. *et al.* Molecular mechanism of rectification at identified electrical synapses in     the Drosophila giant fiber system. *Curr Biol* 18, 1955–1960 (2008).

58. Orr, B. O., Borgen, M. A., Caruccio, P. M. & Murphey, R. K. Netrin and frazzled regulate presynaptic gap junctions at a Drosophila giant synapse. *Journal of Neuroscience* 34, (2014).

59. Winding, M. *et al.* The connectome of an insect brain. *Science (1979)* 379, (2023).

933    60.    Hagberg, A. A., Schult, D. A. & Swart, P. J. Exploring Network Structure,
934            Dynamics, and Function using NetworkX. in *Proceedings of the 7th Python in*
935            *Science Conference* (2008). doi:10.25080/tcwv9851.
936    61.    Schlegel, P. *et al.* navis-org/navis: Version 1.10.0. Preprint at (2025).
937    62.    Hunter, J. D. Matplotlib: A 2D graphics environment. *Comput Sci Eng* 9, (2007).
938    63.    Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in
939            Python. *Nat Methods* 17, (2020).
940    64.    McCarthy, E. V *et al.* Synchronized bilateral synaptic inputs to Drosophila
941            melanogaster neuropeptidergic rest/arousal neurons. *Journal of Neuroscience* 31,
942            (2011).
943    65.    Shiu, P. K. *et al.* A Drosophila computational brain model reveals sensorimotor
944            processing. *Nature* 634, (2024).
945    66.    Schnizler, K. *et al.* A novel chloride channel in Drosophila melanogaster is
946            inhibited by protons. *Journal of Biological Chemistry* 280, (2005).
947    67.    Stimberg, M., Brette, R. & Goodman, D. F. M. Brian 2, an intuitive and efficient
948            neural simulator. *Elife* 8, (2019).
949    68.    Lee, D. & O'Dowd, D. K. Fast excitatory synaptic transmission mediated by
950            nicotinic acetylcholine receptors in Drosophila neurons. *Journal of Neuroscience*
951            19, (1999).
952    69.    Lee, D., Su, H. & O'Dowd, D. K. GABA receptors containing Rdl subunits mediate
953            fast inhibitory synaptic transmission in Drosophila neurons. *Journal of*
954            *Neuroscience* 23, (2003).
955    70.    Saumweber, T. *et al.* Behavioral and synaptic plasticity are impaired upon lack of
956            the synaptic protein SAP47. *Journal of Neuroscience* 31, (2011).
957    71.    Churgin, M. A. *et al.* A neural correlate of individual odor preference in
958            Drosophila. *Elife* 12, (2025).
959    72.    Clements, J. *et al.* NeuronBridge: an intuitive web application for neuronal
960            morphology search across large data sets. *BMC Bioinformatics* 25, (2024).
961    73.    Augustin, H. *et al.* Reduced insulin signaling maintains electrical transmission in a
962            neural    circuit in aging flies. *PLoS Biol* 15, e2001655–e2001655 (2017).
963    74.    Gu, H. & O'Dowd, D. K. Cholinergic Synaptic Transmission in Adult Drosophila
964            Kenyon Cells In Situ. *Journal of Neuroscience* 26, 265–272 (2006).