The Drosophila connectome reveals Axo-Axonic Synapses on Descending Neurons

Cesar Ceballos[1], Juan Lopez[1], Ty Roachford[1], Daniel Sanchez[1], Sabrina Jara[1], Kelli Robbins[1],
Casey L Spencer[2], Rodney Murphey[1], Rodrigo FO Pena[1],*

[1]Department of Biological Sciences, Florida Atlantic University, FL 33458, Jupiter, Florida, USA

[2]Harriet Wilkes Honors College, Florida Atlantic University, Jupiter, FL, 33458, USA

*penar@fau.edu

Abstract

Axo-axonic synapses can veto, amplify, or synchronize spikes, yet their circuit-scale logic is
unknown. Using the complete electron-microscopy connectome of the adult male Drosophila, we
charted every axo-axonic input onto the 1,314 descending neurons that carry brain commands to
the ventral nerve cord. By definition, any synapse connected to a descending neuron within the
cord is axo-axonic. Thus, we uncovered the ascending-descending and interneurons-descending
axo-axonic relationship. Neurons with many partners (high-degree) integrate into the network
without clustering into an interconnected 'rich-club' of hubs. We identified an octet of ascending
neurons (AN08B098) whose axo-axonic input to the Giant Fibers (DNp01) predicted modulation
of the escape circuit. Immunostaining confirms their cholinergic identity, while optogenetic
activation confirmed that this excitatory cohort increases DNp01 excitability, validating
connectome-derived rules. Our work delivers a map of axo-axonic wiring in a complete ventral
nerve cord connectome and provides constraints for models of fast motor control.

Keywords: Connectome, Drosophila, Axo-axonic, Giant Fiber escape circuit, Network topology,
Electron-microscopy reconstruction, Motor control

Introduction

Neurons communicate through a variety of synaptic motifs. In flies, the canonical arrangement is axo-dendritic, but a second configuration, axo-axonic, can exert unusually powerful control over spike initiation and transmitter release[1,2]. Axo-axonic contacts (AACs) in invertebrates have been known for some time[2–6]. Similar studies have also been conducted in vertebrates[7–9]. In the mammalian cortex, exemplified by chandelier cells, recent studies show how inhibition modulates the pyramidal axons at their initial segments[10,11]. In fish, excitatory spiral fibers and M-cells control a similar circuit[12]. Although these examples suggest broader significance, the quantitative wiring logic of axo-axonic synapses remained poorly characterized, primarily due to the absence of a fully mapped system at the necessary synaptic resolution.

This limitation has been addressed by recent electron microscopy reconstructions of the complete adult male nerve cord and female brain (MANC[13–15] and FAFB[16]) in *Drosophila melanogaster*, now publicly available through the NeuPrint interface[17]. Early graph analyses of the FAFB data revealed rich-club hubs and other mesoscale motifs[16], but systematic exploration of the ventral nerve cord (VNC), the insect analogue of the vertebrate spinal cord, is only beginning. At the heart of the VNC, a pair of descending neurons, known as Giant Fibers (GFs), function as command-like neurons that receive visual input in the brain and drive jump and flight motor neurons [18–20]Click or tap here to enter text.. These GFs receive a high number of AACs; thus, it is only natural to assume that an AAC-to-behavior modulation is in place. This system is comparable to the descending escape circuits in larger insects, which have been shown to be the target of axo-axonic connections, exemplified by the input to the descending contralateral movement detector (DCMD) neurons in the locust central nervous system[4].

Within the VNC, a single class of neurons is ideally positioned for studying axo-axonic wiring and their connection to the GFs: Descending neurons (DNs). Because DNs have their somata and dendrites in the brain yet project their axons into the VNC[21–24], any synapse made onto a DN within the cord is, by definition, axo-axonic. In fact, we have manually curated 1,000 ascending-descending neuron connections, where we observed clear presynaptic mitochondrial expression (93%), a hallmark of axonal terminals[25] (example shown in Fig. S1, and curation details in Table S1). The MANC connectome contains DN, ascending neurons (ANs), intrinsic interneurons (INs), and motor neurons (MNs), offering a comprehensive substrate for network-wide analysis of this motif. Although classic anatomical work suggested that each GF acts largely in isolation, our work using the MANC connectome reveals an unexpectedly rich set of axo-axonic inputs onto the GF axons, including direct contacts from other DNs, ANs, and INs. In the present study, we show that the GF receives 97 distinct axo-axonic partners, with some offering a crucial role for axo-axonic modulation, i.e., an octet of ascending neurons (AN08B098). Below, we will refer to these neurons of interest as Axc.

Past light-microscopic surveys catalogued individual DN types and noted scattered DN-to-DN contacts[21], yet the circuit-level role of axo-axonic synapses has remained unexplored. The outstanding issues are their prevalence, partner preference, and functional impact on fast behavior. Specifically, it is unknown how frequently axo-axonic connections occur among the 861,591 possible DN pairs, and whether such a sparse motif can nevertheless mediate rapid, system-wide control required for escape. We tackle these questions by mining the complete electron-microscopy connectome of the adult male ventral nerve cord (MANC). Our analysis (i) enumerates every axo-axonic input onto all 1,314 DNs, (ii) dissects the morphological relationship to DN connectivity, (iii) shows that the DN network is small-world but not rich-club, partitioned into thirteen functional communities, (iv) uses large-scale leaky integrate-and-fire simulations to predict which highly connected AACs modulate the Giant Fiber escape pathway, and (v) confirms those predictions with targeted optogenetic activation *in vivo*.

Our results provide the first system-level blueprint of axo-axonic wiring in the *Drosophila* adult nervous system, suggesting general design principles for descending motor control that may extend beyond insects. They also illustrate the functional power of the sparse axo-axonic network described below and place the GFs at the center of a distributed control system rather than a simple linear reflex arc.

Results

Global Distribution of Axo-Axonic Inputs to Descending Neurons.

To uncover the prevalence of axo-axonic connections made onto DNs in the VNC, we used the MANC dataset available through the NeuPrint+ interface and API[26–28]. Figure 1*A-C* shows morphological reconstructions of some exemplary DNs of interest from Neuroglancer: DNp01 (GFs, black dot), DNd03 (blue dot), and DNg02 (red dot, Fig. 1*A-C*). For each neuron, Neuroglancer overlays the 3D-EM reconstruction on the VNC volume. The GFs, whose axons are nearly unbranched (green or blue), receive one of the highest numbers of axo-axonic inputs (n = 97; see NeuPrint+) in the dataset. DNd03 is a highly arborized neuron with a similarly large input count (n=160), while DNg02 has less arborization but similar axo-axonic inputs (n=110). Interestingly, all three share similar synaptic strength (i.e. the equivalent of synapse count from the connectome MANC) despite differences in arborization (DNp01 2352, DNd03 2364, DNg02 1751; Fig. 1*H*). Neuroglancer presents transmission-EM micrographs at 500 nm resolution taken from MANC; presynaptic active zones are marked by magenta dots, postsynaptic densities by cyan dots, and partner axons are pseudo-colored to match the 3D rendering (Fig. 1*D-F*).

Figure 1. Descending neurons (DNs) and their axo-axonic inputs in the VNC. *(A)*-*(C)* Exemplary Neuroglancer reconstructions of three DNs of interest. *(D)*-*(F)* EM images illustrate axo-axonic connections formed by the three reconstructed neurons: *(D)* shows DNp01 to DNp01 connectivity, *(E)* shows DNd03 connectivity with DNp42, and *(F)* shows DNg02 to DNg02 connectivity (same colors as in A-C). Red circles identify the locations of presynaptic T-bars; blue circles show postsynaptic densities. Circles were enlarged for visibility. *(G)* Histogram of the number of axo-axonic neurons innervating one DN. Presynaptic neurons can be DN, AN, or IN. *(H)* Histogram of the cumulative strength of the synaptic connections for each DN. Presynaptic neurons can be DN, AN, or IN. *(I)* Scatter plot and linear regression of the number of axo-axonic neurons vs their synaptic strength. $R^2 = 0.92$ and $p < 0.001$.

We quantified the distribution of axo-axonic connections (i.e. presynaptic DN, AN and IN) made onto DNs reported by MANC. Figure 1*G* reveals a highly skewed distribution of partner count; the median DN is innervated by a single presynaptic neuron (inter-quartile range 1–3), and 11% of DNs receive input from only one cell with a synaptic strength of six or greater. A small set of outliers, however, attracts hundreds of partners, including DNp01 (GFs) with 97 distinct axo-axonic inputs. Strikingly, the GFs receive these connections despite possessing one of the simplest axonal arbors in the dataset (Fig. 1*A*). All other DNs exhibit more extensively branched and structurally complex axonal arbors, which are likely to account for their higher levels of innervation. Total input strength follows a similar heavy-tailed pattern (Fig. 1*H*). The modal DN receives fewer than 20 synapses, whereas the GF is again among the top of the list in terms of individual contacts. Across the population, partner count and cumulative strength are tightly coupled (Pearson r = 0.96, p < 0.001; Fig. 1*I*), indicating that neurons that recruit many partners also receive proportionally stronger innervation. Together, these findings identify the distribution of axo-axonic connections made with DNs in the VNC. We confirm that the synapses quantified in the graph analysis correspond to active zones and demonstrate how neurons with markedly different morphologies can converge on similar axo-axonic input profiles.

Using these findings, we next examine whether anatomical complexity serves as a strong, though not perfect, predictor of synaptic strength for axo-axonic inputs onto DNs (Fig. 2). Representative reconstructions illustrate the general pattern: neurons with short, unbranched axons receive few contacts, those with intermediate arborization receive proportionately more, and highly branched cells accumulate the greatest axo-axonic strength (Fig. 2*A-C*). A kernel density estimation (KDE) plot of all 1,314 DNs reinforces this observation, showing a diagonal ridge in which increasing cable length accompanies increasing total input strength (Fig. 2*F*). Linear, quadratic, and symbolic regressions all yield significant positive slopes, confirming a monotonic relationship (Fig. 2*G*). However, the shallow gradient and the presence of two striking outliers, a long axon with sparse innervation and a short axon with dense innervation, indicate that branch length alone does not determine synaptic investment (Fig. 2*D-E*). Thus, we find that a presynaptic neuron's morphology broadly predicts the degree of modulation received by a partner DN, although additional cell-type-specific factors likely refine the final allocation of axo-axonic synapses.

141
Figure 2. Analysis of morphology and synaptic strength between DN-to-DN axo-axonic connections. (*A-E*)
Examples of DNs include (legend colors indicate pre- and postsynaptic in the figure): (A) low connectivity with simple
morphology, (B) moderate connectivity and morphology, and (C) high connectivity with complex morphology.

Additionally, (D) low connectivity with complex morphology and (E) high connectivity with simple morphology
represent exceptions. These neurons generally align with the overall trend—a linear correlation between
morphological complexity and synaptic strength—while (D) and (E) are outliers. Dots indicate synapse locations. *F*)
Kernel density estimation (KDE) of the scatter data of total input strength vs cable length, *G*) Scatter plot of total input
strength vs cable length and the corresponding linear regression, polynomial regression of a quadratic function, and
symbolic regression.

Further exploration revealed a unique feature of connectivity. In the VNC, neurons will often form
synapses with multiple downstream partners (Fig. 3*A-B*). We found certain neurons, such as
members of AN08B098, had a disproportionately larger number of connections made onto a single
target type of DNs such as the GF than other neuron type. To determine whether these neurons
constitute a separable class, we quantified neuron output specialization for every presynaptic
neuron using three metrics: the fraction of total synapses made onto the dominant DN, the entropy
of the neuron type distribution, and the number of connections of the second largest neuron type
connection (off-target) relative to the largest DN connection (on-target).

For each presynaptic neuron, we first quantified the percentage of strength of its outgoing synapses
that were made onto its most strongly targeted DN. Neurons were required to exceed a threshold
value and the percentage of strength of the dominant DN was permitted to exceed the threshold,
ensuring a uniquely dominant target. Initially, we assumed threshold values that separate two
populations: multi-target vs single-type biases. The final threshold was determined by optimizing
the minimization of an error parameter. A statistical comparison of the distributions using the
Kolmogorov-Smirnov test (with a significant p-value < 0.001) revealed two distinct distributions
of presynaptic neurons, which we defined as single-type biased and multi-target, with threshold
values of 0.27 in MANC v1.2.1 and 0.1 in male-cns v0.9. In addition, single-biased cannot send
more than 15 connections to other off-target DN to exclude external influence[29].

In both connectomes, we identified several single-type biased axo-axonic connections (Fig. 3*C*) to
DNs in both MANC v1.2.1 and male-cns v0.9. DNp01 receives the highest number of single-target
AACs in MANC (n = 16) and among the highest for male-cns (n = 11). Among these neurons, five
belong to the Axc type, and the remaining are labeled as "unknown".

177
Figure 3. Neurons that provide exclusive axo-axonic connections to DNs. (*A*) Sketch of single type biased vs (*B*)
multi-target neuron. Single-type biased neurons project a significant proportion of their outgoing connections to a
single DN, while unbiased neurons form several strong connections among partners. (*C*) Number of exclusive neurons
per DN for both MANC v1.2.1 (blue histogram) and Male-cns v0.9 (orange histogram). The Giant Fibers (DNp01;
marked by an asterisk) are among a small population of DNs that receive most single-type biased connections, i.e., as
seen in (D), there are approximately 16 neurons that innervate DNp01 in MANC and 11 in Male-cns. (*D*) For each
dataset, the fraction of outputs made from neurons onto their most strongly targeted DN type was computed per
neuron, and a dataset-specific cutoff was identified from the distribution as the point separating the population of
unbiased neurons and single-type biased neurons (MANC 0.27, male-cns 0.1). Single-type biased neurons across both
datasets had similar properties, including high target preference, reduced target entropy, and reduced off-target
connection strength (Kolmogorov-Smirnov test, p < 0.001). We further refined this by removing neurons whose off-
target connections had 15 or greater synapses, suggesting non-influence over their synaptic partners[29].
190
Compared with other highly connected DNs, the GFs attain an equivalent synaptic strength with a
simple morphology, further demonstrating how morphological complexity alone does not predict
axo-axonic input. These results identify the GFs as exceptional hubs for axo-axonic modulation

due to their significant connectivity despite their relatively simple morphology compared to other
highly connected DNs. However, these findings raise several new questions: How are AAC
populations connected, and how do they act on GFs? Is the GF an outlier, or do similar axo-axonic
motifs govern other rapid motor pathways in the fly? How do these contacts alter GF excitability?
We address these questions by investigating the strongest single-target AAC population
presynaptic to the GFs, the Axc type neurons.
200
Sparse yet Globally Connected Axo-Axonic Network
To address how AAC populations are connected, and their influence on other DNs, we extracted
every DN-to-DN contact from the updated MANC dataset. The resulting adjacency matrix is
strikingly sparse: only about 1% of the possible ordered pairs carry even a single synapse (Fig.
4*A*). Moreover, partner choice is not random: homotypic pairs (same DN type) and heterotypic
pairs (different types) show distinct transmitter profiles, as shown for acetylcholine, GABA, and
glutamate, implying that neurotransmitter identity contributes to the selectivity of these long-range
axo-axonic contacts (Fig. 4*D*). In homotypic DN connections, ~90% of the connections are
cholinergic (excitatory), while heterotypic DN connections are ~60% cholinergic [the remaining
are GABA/glutamate (inhibitory)]. This suggests that AACs between DNs, while rare, are
predominantly excitatory. connections.
212
We then asked whether the same wiring principle applies to non-descending inputs, namely
ascending neurons (ANs) and intrinsic interneurons (INs). Repeating the analysis with ANs and
INs as presynaptic partners to DNs again produced very sparse matrices: about 1.2% of all possible
AN-to-DN pairs and 0.7% of the IN-to-DN pairs contain at least one synapse (Fig. 4*B-C*). Figure
4*E* shows the neurotransmitter profile for AN-to-DN and IN-to-DN pairs. Both pair profiles were
similar, with nearly half of the neurons classified as cholinergic and the other half
GABA/glutamate. This suggests that AACs between AN-to-DN and IN-to-DN follow the same
rules for neurotransmitter identity as DN-to-DN AACs.
221

Figure 4. Axo-axonic connectivity analysis. (*A*) Connectivity matrix of pairs of DN-to-DN pairs. Only 1% of all possible connections were identified, suggesting that DN-to-DN axo-axonic connections are extremely rare. (*B*) Connectivity matrix of pairs of AN-to-DN pairs. Only 1.2% of all possible connections were found, which suggests that AN-to-DN axo-axonic connections are also exceptionally rare. (*C*) Connectivity matrix of pairs of IN-to-DN pairs. Only 0.7% of all possible connections were observed, indicating that IN-to-DN axo-axonic connections are uncommon. (*D*) Pie chart of predicted neurotransmitter of DN of the different type (left, e.g., DNp01 -> DNp02) and same type (right, e.g., DNp01 ->DNp01) that make pairs of axo-axonic connections. (*E*) Pie chart of predicted neurotransmitter of AN-to-DN and IN-to-DN axo-axonic connections. (*F*) Histograms of the number of (F1) presynaptic DN (in-degree) and (F2) postsynaptic DN (out-degree) for DN-to-DN pairs. (*G*) Histograms of the number of (G1) presynaptic AN and (G2) postsynaptic DN for AN-to-DN pairs. (*H*) Histograms of the number of (H1) presynaptic IN and (H2) postsynaptic DN for IN-to-DN pairs. (*I*) Rich-club analysis of the DN-to-DN axo-axonic network. (*J*) Connectivity matrix of pairs of AN/DN/IN-to-DN connections. Filled circles follow the same color scheme as in Fig. 1*A-C*. Cyan square identifies Axc-type neurons. Dashed green lines and gray tones for dots separate ANs, DNs, and INs. (*K-P*) Examples of each category from DN-to-DN, AN-to-DN, and IN-to-DN. (From left to right: (*K*) DNx01-DNx01, (*L*) DNb04-DNb04, (*M*) AN08B099-DNp01, (*N*) AN10B019-DNp01, (*O*) IN11A001-DNp01 and (*P*) IN06B059-DNp01). (*Q*) Percentage of contralateral and ipsilateral connections between DN-to-DN, AN-to-DN, and IN-to-DN. (*R-T*) Pie chart of predicted neurotransmitter of DN to contralateral DN, AN to contralateral DN, and IN to contralateral DN of axo-axonic connections.

Lastly, we show the distribution of the number of pre-synaptic DN, AN, or IN that project axo-axonic synapses to a post-synaptic DN (Fig. 4*F-H* top row) as well as the number of postsynaptic DN that are innervated axo-axonically by the same DN, AN, or IN (Fig. 4*F-H* bottom row). The distributions are similar across neuron types, showing a decay pattern resembling an exponential (Poisson) distribution, where a few neurons are highly connected, and a few neurons are sparsely connected. When we merge all three presynaptic classes into a single connectivity matrix (Fig. 4*J*,) the same pattern from Figure 4*A-C* emerges: extreme sparseness (see dots where our exemplary neurons are located; color scheme as in Fig. 1*A*.). Note that in Fig. 4*A* there is a clear diagonal line which indicates DNs synapsing onto their contralateral counterparts. We also highlight the degree of convergence across presynaptic classes and identify connectivity hubs based on their total input. Taken together, regardless of presynaptic identity, the findings demonstrate that the network's organization combines stringent partner selectivity with complete coverage of the descending system.

Global wiring principles were identified to explore more refined AAC spatial connectivity preferences and determine whether they are repeated across DNs, ANs, and INs. Figure 4 shows examples of the contralateral axo-axonic connections between DN-to-DN (Fig. 4*K-L*), AN-to-DN (Fig. 4*M-N*), and IN-to-DN (Fig. 4*O-P*). Figure 4*Q* shows the percentage of contralateral and ipsilateral connections. AACs between contralateral DNs are rare. However, ANs and INs projecting to contralateral DNs are frequent. Interestingly, the neurotransmitter profiles from our earlier findings are preserved regardless of the type of presynaptic neuron, where approximately half of the neurons are cholinergic (excitatory), and the other half is GABAergic/glutamatergic (inhibitory) (Fig. 4*R-T*).

Collectively, these findings suggest a coherent narrative. Whether the presynaptic axon descends, ascends, or remains intrinsic to the cord, axo-axonic wiring obeys the same two rules: partner selectivity is extremely high, and midline crossing is common only for neurons that share strong

269 reciprocal ties. The result is a VNC in which local subtypes can interact tightly with their mirror
270 images, yet the great majority of neurons remain insulated from direct axo-axonic influence.
271

Graph-Theoretic Architecture of the Axo-Axonic Network.
To address if GF is an outlier or do similar axo-axonic motifs govern other rapid motor pathways
in the fly, we study connectivity matrices that are derived employed as a directed graph to examine
the system-wide arrangement of sparse axo-axonic interactions among DNs (Fig. 4*A*). The graph
contains 884 nodes and 2,757 edges, giving a density of 0.007. Despite this low density, the
average shortest-path length is only 4.03, indicating that any two DNs can be reached through a
small number of intermediate axonal contacts. In fact, we can compare with the chance level: the
formula $\langle L \rangle = \ln(n) / \ln(\bar{k})$ is a well-known and widely used approximation for the average
shortest-path length in large random networks, such as the Erdős-Rényi model. Here, n is the
number of vertices in the graph, $\bar{k}$ is the mean degree, which can be obtained by the ratio m/n
with m being the edge count. In our problem, $n = 884$ DNs and $m = 2,757$ axo-axonic contacts
leave us with $\langle L \rangle \approx 5.9$. Therefore, the average shortest path length is indeed shorter than you
would expect by chance. The clustering coefficient is likewise low at 0.013, indicating limited
local redundancy. Community detection identifies 13 separate modules, consistent with the
network's functional subdivision. Node degree ranges from 1 to 29 with a mean of 6.24, reflecting
a modest hierarchical spread.
288

The rich-club coefficient Φ(k) measures the density of connections among nodes whose total
degree (in-degree + out-degree) equals or exceeds a threshold k. Φ(k) = 1 indicates that the high-
degree subgraph is fully connected, whereas Φ(k) = 0 means that those nodes share no edges
beyond chance level. The rich-club coefficient in our data peaks at 0.076 for nodes with a degree
of at least 18, so high-degree hubs do not preferentially connect to each other; integration is
therefore distributed rather than centralized (Fig. 4*I*). The modest peak value and the absence of
a sustained plateau indicate that high-degree DNs do not preferentially interconnect to form a rich
club; instead, network integration is distributed rather than hub-dominated.
297

A slightly positive degree assortativity of 0.029, a measure that suggests a tendency of nodes to
connect with other nodes in a similar way, suggests a weak tendency for neurons of similar degree
to link. Finally, the maximum betweenness centrality of 0.049, a measure indicating how often a
node lies on the shortest path between all pairs of nodes in a network, highlights a small set of
bridge neurons that channel information between modules. Together, these metrics portray a
network that combines efficient global communication with modular specialization, an
architecture well-suited for coordinating diverse motor outputs.
305

Axo-axonic modulators of the Giant Fiber escape circuit.
The MANC connectome introduces several novel neuron types to the Giant Fiber escape circuit.
To understand and underscore the significance of AAC modulation of the circuit, it is important
to know which individual cells can exert a strong, targeted influence on a system such as the GF

system[18,19]. As shown in Figure 3, the GFs are among a small population of DNs that receive
single-target connections. A focused search within MANC revealed the Axc type neurons, a cluster
of eight neurons whose axo-axonic output is both nearly exclusive and among the strongest to the
GFs (Figs. 3-4). These Axc-type neurons are predicted to be cholinergic by the MANC
connectome. The BANC connectome demonstrates that these neurons do not connect to the GF in
the brain (see Fig. S3).
316
To test whether the connectome-predicted Axc type neurons truly form chemical AACs with the
GFs, we used split-Gal4 drivers to direct membrane-bound GFP to Axc axons, back-filled the GFs
with tetramethyl-rhodamine, and labelled presynaptic active zones with anti-Bruchpilot (nc82).
High-resolution confocal stacks revealed discrete GFP-positive boutons that colocalize with Brp
puncta on the descending GF tract in the first thoracic neuromere (Fig. 5*A*,*C*), confirming axo-
axonic contacts between Axc type neurons and GF. The split-Gal4 line being used labels
AN08B098. However, there are a few extra neurons labeled that can come from AN08B099 and
make weak connections to the GF. We have also provided (i) single-section confocal images with
optical enhancement showing the synaptic connectivity and (ii) similar EM reconstructions from
Neuprint to demonstrate that synapse locations along the axon correspond to synapse locations
identified in the MANC connectome (Fig.5*G-O*). Immunostaining for Choline Acetyltransferase
(ChAT) revealed that Axc somata, as labelled through genetic expression of mCD8::GFP, are
ChAT-positive (Fig. 5*D-G*), establishing this cohort as likely cholinergic and therefore excitatory.
330

Figure 5. Axcs make cholinergic axo-axonic synapses on the GFs as predicted by the MANC connectome.
(*A*) AN08B098-type neurons (green) form axo-axonic connections at distinct locations along Giant Fiber (red, GF)
axons. The scale bar shown is 20 µm. (*B*) The AN08B098-GF connections reconstructed in Neuroglancer mirror the
morphology seen with fluorescence in *A*. (*C*) We validated AN08B098-to-GF connectivity using anti-bruchpilot to
stain for T-bars in active zones (white) along AN08B098-type neurons and colocalized the resulting fluorescence with
the GFs (red), which were filled with tetramethylrhodamine. The scale bar shown is 20 µm. (*D*) Neuroglancer reveals
presynaptic sites at similar locations seen in colocalized fluorescent image *C*. (*E-F*) XY, and XZ plane view of the
preparation shown in *A* and *B*. The zoomed-in 6 µm inlays show AN08B098 forming a single synapse with the GF in
E. The yellow arrows detail the precise location AN08B098 forms the Brp-positive chemical synapse (white) to the
GFs in F. Scale bars shown are 5 µm. (*G-O*) EM images showing monosynaptic connections between single GF
(green) and AN08B098 neurons identified by the following MANC id: (G-H) 21041, (I) 21589, (J) 23949, (K) 152261,
(L) 16900, (M) 20444, (N) 22275, and (O) 24038. Pre-and postsynaptic sites are detected in EM slices using a 3D
convolutional neural network to identify T-bars (cyan dots) and postsynaptic densities[30] (PSDs, magenta dots). (*P-S*)
We expressed anti-choline acetyltransferase (anti-ChAT) and anti-GFP in AN08B098 neurons. Anti-ChAT

346   colocalizes to AN08B098 cells, with particularly bright staining in the cell bodies. This finding suggests acetylcholine
347   synthesis is present within these cells, validating connectome transmitter predictions for AN08B098.
348
To address how these contacts alter GF excitability, and following this imaging confirmation, we
investigated whether these AACs could measurably influence GF activity in a simulation of the
complete VNC connectome. To test this, we stimulated all VNC neurons with a Poisson
distribution of excitatory synaptic currents at 30 Hz and 5 nS. Additionally, one GF was stimulated
more strongly at 10 Hz and 40 nS to establish a baseline level of excitability (Fig. 6*A, where* spikes
are indicated by large black dots at the bottom of the raster plot and arrows underneath). When the
eight cholinergic Axc neurons were activated, the GF crossed threshold more often, indicating that
the axo-axonic inputs are net excitatory (Fig. 6*B*). Additional simulations with different scenarios
where stimulation and deletion of non-Axc (nAxc) populations demonstrate that the Axc cells are
paramount for increased excitability (Figs. 6*B-E*). GF subthreshold voltage histograms quantify
the increased excitability (Fig. 6*F*), supported by statistical significance (Fig. 6*H*). These
simulations highlight a previously unrecognized cohort of axo-axonic synaptic modulation,
providing a rapid mechanism for amplifying the escape-circuit output.

362
Figure 6. Simulations using the whole VNC network. (*A*) Stimulation of the GF and corresponding raster plot of
all the neurons in the VNC and voltage trace of GF response (spikes are shown as large black dots at the bottom of
the raster plot, also marked with an arrow). (*B*) Schematic of the GF stimulation by Axc type neurons and
corresponding raster plot of all the neurons in the VNC and voltage trace of GF response (GF spikes are represented
by large black dots at the bottom of the raster plot, while Axc spikes are represented by large red dots). (*C*) Similar to
previous plots, but now the stimulation is driving non-Axc (nAxc), and the Axc is removed from the simulation. (*D*)

Similar to previous plots, but now the stimulation is driving Axc cells, and nAxc are deleted. (*E*) Similar to *A*, but now
Axc cells are deleted. (*F*) Histogram comparison of subthreshold voltages during GF-only and GF+Axc stimulation.
(*G*) Similar to *F*, but now the histogram shows the difference between silencing nAxc and stimulation in Axc vs.
silencing Axc and stimulation in nAxc. (*H*) Violin plots showing statistical analysis of some of the comparisons. The
overall excitability of the GF increases when the Axc neurons are activated, leading to higher firing probability for
the GF.
375
As the control experiment, we also tested the possibility of silencing nAxc cells, which are defined
here as presynaptic cells to the GF that contribute to polysynaptic pathways, and Axc cells
selectively (Figs. 6*C-D*). The results from these figures demonstrate that when nAxc connections
are silenced, the increased excitability persists, and when the Axc monosynaptic connections are
silenced, but nAxc is retained, the excitability is decreased. Fig. 6*G-H* shows additional
quantification of this Axc to GF excitability, with the only non-significant comparison being Axc
stim vs. silencing nAxc + Axc stim, demonstrating that the possibility that Axc acts
polysynaptically could be neglected.
384

385   **Testing connectome predictions *in vivo* via optogenetic stimulation of axo-axonic neurons.**
386

To assess the predictions of these connectome simulations, we tested the ideas *in vivo* using classic
electrophysiology and optogenetics. We used the split-Gal4 line that labels the Axc neurons
(BDSC #89945) to drive a UAS construct containing a mVenus-tagged red-shifted
channelrhodopsin, CsChrimson (BDSC #55136). Expression of CsChrimson was verified with
immunohistochemistry (Fig. S2A). We electrically stimulated the GFs at 24 Hz in flies expressing
CsChrimson in Axc type neurons and recorded from the jump muscle (TTM) and flight muscle
(DLM) as previously described[31], (Fig. 7*A-B*). We then set the electrical stimulation of the GF just
below threshold. This near-threshold stimulation resulted in occasional (~20%) responses in both
muscles (n = 5 specimens, Fig. 7*B* and 7*E*). However, when the same preparation was illuminated
(590 nm, 0.37mW) during the stimulus train, the sub-threshold electrical stimulus elicited
responses from both muscles nearly 100% of the time (n = 5 specimens, Fig. 7*D* and 7*F*). A one-
way ANOVA found significant differences between responses before, during, and after light
application ($p < 0.001$), but there was no significant difference for the control group ($p = 0.53$)
(Fig. 7*G*). Post-hoc multiple comparisons revealed significant differences between responses at
baseline (before light application) and during and after light application. We also found significant
differences using a paired t-test for the same responses during and after light application
($p<0.001$), but not before ($p>0.26$).
404

Figure 7. Optogenetic activation of the Giant Fiber system. (*A*) Sketch of the electrophysiological recordings with optogenetic activation of axo-axonic cells (Axc). (*B*) Example of raw traces for control (no light application). A zoomed-in panel is presented, showing a constant difference in latencies of ~0.6 ms between TTM and DLM indicative of GF spiking. (*C*) Simulation of a computational model of the VNC connectome emulating the experimental results under subthreshold conditions and light stimulation. Note that GF only spikes when the light is applied and is time-locked with the electrical pulses. Action potentials are indicated with arrows. (*D*) In vivo example of raw traces when light was applied. A zoomed-in panel is presented, showing difference in latencies of ~0.6 ms between TTM and DLM responses. (*E*) Response probability per fly (n = 5) under no-light conditions. (*F*) response probability per fly (n = 5) under light conditions (light was applied during the blue shaded region). (G) Average and SEM of response probability (paired t-test, * = *p*<0.001). **Spike initiation in the VNC**. (*H*) Sketch of the electrophysiological recordings with optogenetic activation of the GF before and after decapitation. (*I*) Recordings from both TTM and DLM before decapitation. The constant latency between TTM and DLM indicates the presence of GF action potentials. A zoomed-in panel shows the constant latency between TTM and DLM ~0.77 ms. (*J*) Response of the TTM to optical stimulation in a specimen driving CsChrimson in the GF as above. The blue curve represents the response of the TTM in animals with the head on, exhibiting a long tail of post-light response. The black curve is after the head was removed to remove the spike-initiating zone in the brain. (*K*) Recordings from both TTM and DLM after decapitation. The constant latency between TTM and DLM indicates the presence of GF action potentials. A zoomed-in panel shows the constant latency between TTM and DLM ~0.76 ms.

To verify that light stimulation does not elicit effects in the absence of CsChrimson expression or function, we performed two control experiments. In the first experiment, flies expressed GFP instead of CsChrimson (Fig. S2A); in the second, CsChrimson-expressing flies were raised without dietary cis-trans-retinal, thereby preventing optogenetic activation (Fig. S2*B*). A one-way ANOVA revealed no significant differences in stimulus-evoked responses across conditions (*p* = 1). Post-hoc comparisons revealed no significant differences between the baseline, light, and post-light periods. Paired t-tests also found no significant differences between light and no-light conditions during and after stimulation. These results confirm that the observed effects are specifically attributable to CsChrimson activation.

How the Axc alters excitability depends in part on the site of spike initiation for the GF. In the present experiments, the stimulating electrodes are in the brain where there are no monosynaptic connections from Axc to GF[26]. The only monosynaptic Axc-to-GF connections are in the VNC (Fig. 5, S3). How can the optical excitation of the Axcs in the VNC, where they make synaptic contact, have a modulatory effect on the GF spike initiation zone (SIZ) in the brain, where they do not make contact? We have now demonstrated that the GF also possesses a spike-initiating region(s) in the VNC, where the Axc could influence spike initiation. We demonstrated this alternative SIZ by driving CsChrimson in the GF, removing the head to remove the SIZ in the brain, and testing headless animals with optical stimulation to reveal GF activity generated in the VNC (Fig. 7J-*K*). GF activity driven by the light stimulation is revealed by the response of both TTM and DLM with a constant latency between them indicative of GF action potentials (Fig. 7*K*). Therefore, the Axcs can directly influence this SIZ in the VNC.

Unexpectedly, we also note a prolonged response of the GF after the light was turned off (Fig. 7D, 7F and 7G). Before light application, the TTMn response probability was low (~20%, black trace,

Fig. 7*G*), during light application the probability increased to almost 100% (blue trace, Fig. 7*G*)
and after light-off activity returned to baseline very slowly (>100 msec). This prolonged response
after light-off has been observed in several other situations for CsChrimson. For instance, at the
larval neuromuscular junction (NMJ), CsChrimson drove spiking in motor neurons (monitored by
muscle synaptic potentials) and exhibited a post-light response hundreds of milliseconds in
duration[32]. In a second example, descending neurons DNp07 and DNp10, when expressing
CsChrimson, exhibit a 100-400 msec response after light-off[33]. Finally, we tested this idea in the
GF by expressing CsChrimson in the GF and illuminating the preparation at 590 nM and observed
a long tail of TTM response of approximately 100 msec (Fig. 7*J,* blue trace). In brief, a variety of
examples show that CsChrimson when activated produces a strong response with a long slow
decay after the light is turned off (Fig 7*F-G)*.
461
To confirm the parallel between the simulation and the *in vivo* experiment, we also developed a
simulation that more closely matched the *in vivo* experiments (Fig. 7*C*). A large-scale network of
the VNC, composed of leaky integrate-and-fire neuron models, was used to simulate this
experiment. The GF was electrically stimulated just below threshold at 24 Hz. In these simulations,
the GF only spikes in the presence of the light. Both the simulation and the optogenetic results on
the living animal suggest that the Axc-to-GF connection is excitatory, consistent with the
connectome and chemical predictions that the synapse is cholinergic. These results agree with our
simulation's predictions that the main effect of the optogenetic activation of the cholinergic axo-
axonic neurons (Axcs) is to increase the excitability of the GF (Fig. 7*C*).
471
472   Overall, these results support our prediction that the primary effect of optogenetic activation of the
473   cholinergic axo-axonic neurons (Axcs) is to increase the excitability of the GF. Our conclusion is
474   that the optically activated Axcs excite the GF via their synapses in the VNC and alter the
475   likelihood of GF action potentials.
476

Discussion

By analyzing all 1,314 brain-originating descending neurons, we move beyond scattered descriptions of axo-axonic contacts and derive circuit-level design rules for presynaptic modulation. The quantitative principles that emerged within this work are: (i) extreme sparseness showing that between 0.7 to 1.2% of possible DN-to-DN, AN-to-DN, and IN-to-DN pairs form axo-axonic synapses; (ii) a tight linear relation between synaptic strength and partner multiplicity; and (iii) a small-world architecture that distributes integration rather than concentrating it in a rich-club core. Together, these features constitute a wiring grammar for axo-axonic control in the adult *Drosophila* motor system.

Axo-axonic synapses have previously been identified in the *Drosophila* brain connectome [34–37]. Excitatory cholinergic synapses, like those investigated here, have been observed in the fly olfactory system, where projecting neurons (PN) form lateral PN-to-PN connections, similar to the DN-to-DN connections we identified[34]. Axo-axonic connections have also been reported between olfactory sensory neurons (OSNs)[36]. Bristle mechanosensory neurons (BMNs) are also shown to be axo-axonically connected[38]. These similarities suggest that axo-axonic connectivity principles might be conserved across the VNC and brains in flies, although the functional role of AACs remains unexplored.

Notably, axo-axonic wiring has revealed an extraordinarily selective, yet globally percolating schema, in contrast to, for example, the serotonin axo-axonic configuration in the hippocampus, which comprises 30%[10,39] or in the cortex, which comprises 19%[10,40]. In the fly VNC, only ~1 % of the possible DN-to-DN pairs form axo-axonic synapses, an order of magnitude sparser than axo-dendritic contacts in either the MANC or FAFB brain data sets[14,41]. Despite their scarcity, these connections are sufficient to incorporate every DN into a single, strongly connected network, yielding short average path lengths (~4 hops) and thus preserving the ability for any axon to gain rapid influence over remote peers. For comparison, the entire network has an average path length of 2.8 (~3 hops). This combination of stringent partner choice with network-wide reach resembles presynaptic inhibition motifs in mammalian spinal cord and chandelier-cell control of cortical pyramidal neurons, suggesting convergent architectural solutions across phyla[42,43]. What accounts for the sparsity of axo-axonic connections? One possibility is wiring economy: boutons on sister axons consume precious membrane and cytoplasmic resources[44]. Another is developmental lineage. Many high-degree bridge neurons derive from hemilineages that extend across neuromeres[45], hinting that growth-cone timing and Slit-Robo–mediated midline crossing[46] pre-pattern the opportunities for axo-axonic contact. Comparative EM work in other insects and in zebrafish larval spinal cord will clarify whether these archetypes represent a universal motor blueprint.

As previously found, rich clubs with dense interlinking of high-degree hubs are present in the fruit *Drosophila* connectome[16]. They are also widely present in vertebrates, such as in the cerebral

cortex[47,48]. In contrast, the axo-axonic fly DN network revealed only a weak, transient rich-club
coefficient, suggesting that presynaptic modulation in the nerve cord is decentralized. Control must
therefore emerge from many mid-degree "broker" neurons (with fewer connections) and with high
betweenness rather than from a minority of super-hubs. Such a layout guards against single-point
failure and allows different motor modules to be gated independently, consistent with behavioral
data showing that flies can blend local reflexes with whole-body maneuvers[49].
523
In terms of functional implications, axo-axonic contacts positioned directly on descending axons
are ideally placed to veto, amplify, or synchronize commands emanating from the brain before
they fan out onto premotor interneurons and motoneurons. We have shown that Axc is amplifying
GF activity. Experiments with headless animals demonstrate the existence of a second spike
initiation zone (SIZ) in the VNC. We believe this is the target of the Axc synaptic modulation.
Choline acetyltransferase (ChAT) immunostaining confirms that Axc type neurons are
cholinergic, and optogenetic activation of these cells reliably enhances the GF's excitability. Our
simulations also demonstrate this principle: activating a cohort of eight cholinergic Axc type
neurons, which exclusively target the Giant Fibers (GFs), enhances their firing.
533
534 Crucially, however, the GF is not unique; 17 other DNs receive >50 presynaptic partners that
535 occupy bridge positions between functional modules. These neurons may be promising targets for
536 future studies investigating the role of axo-axonic connections in precise behavior modulation.
537 Future opto-physiological experiments probing these DNs will test whether distributed control is
538 the norm rather than the exception.
539
Excitatory axo-axonic connections, exemplified by the Axc to GF projection described here, are
not unique to this system and have been reported in other organisms, including the well-
characterized Mauthner cells in zebrafish[12] and dopamine cells in mice[50]. In these examples, the
primary function of the excitatory axo-axonic synapse is to increase the release probability of the
axon terminal they innervate. This mechanism can even bypass upstream action potentials,
representing a hijacking mechanism that may have physiological roles. The findings presented in
this study may contribute to a broader understanding of axo-axonic connectivity principles across
species[51–5525,56].
In conclusion, our study presents the first comprehensive blueprint of axo-axonic connectivity
within a fully reconstructed adult ventral nerve cord system. The network revealed here offers
concrete parameters for next-generation models of rapid motor decision making in both
invertebrates and vertebrates. By formalizing direct axon-to-axon communications, the work
bridges a critical gap between connectome structure and motor function and opens new avenues
for dissecting the fastest circuits in the animal kingdom.
554
Limitations of study

One limitation of our study is that the connectome used in the analysis was derived from a single
male specimen. New datasets are now being released that are sex-specific or can address
developmental differences, such as the Female Adult Nerve Cord Connectome (FANC) and the
Brain and Nerve Cord Connectome (BANC). All functional whole-network tests are *in silico*, and
although experimental verification is provided through optogenetics, the large-scale network based
on leaky integrate-and-fire neuron model respects synapse counts but ignores conductance
dynamics and neuromodulators influences. Furthermore, EM of the MANC connectome (v1.2.1)
did not detect gap-junctional coupling, suggesting that the present map under-reports electrical
axo-axonic contacts that could further reshape information flow[51–55].
565
Lastly, dendro-axonic synapses account for only 2% of all synapses[25,56]. Thus, we cannot rule out
the possibility that some of the synapses investigated in this study are dendro-axonic, although the
vast majority are axo-axonic. Nevertheless, we curated a large number of EM images (Fig. S1) to
demonstrate the presence of mitochondria in axo-axonic synapses of descending neurons,
suggesting that this structure can be used as a marker of axo-axonic synapses. Based on this, we
found that synapses of AN08B098 onto Giant Fiber contain presynaptic mitochondria, suggesting
that these synapses are axo-axonic.
573
574 We found that the number of neurons in the Gal4 line was larger than the number of neurons found
575 in the connectome that belong to the type AN08B098. We cannot establish the source of this
576 difference. Although we characterized two other Gal4 lines (Bloomington stock #87537 and
577 #87664) and we found similar counts.
578
579 The long tail of response following activation of Axc in Fig 7 could be explained by multiple
580 underlying mechanisms; local recurrent excitatory loop between GF and Axc, unknown
581 descending neurons that feedforward to this circuit, but we are unable to identify the source of
582 these inputs.
583
584

585 **STAR Methods**
586

Detailed methods include the following:
• **KEY RESOURCES TABLE**
• **RESOURCE AVAILABILITY**
• **EXPERIMENTAL MODELS AND SUBJECT DETAILS**
• **METHODS DETAILS**
▪ Electrophysiology
▪ Solution preparations
▪ Optogenetics
▪ Dye-injection
▪ Immunostaining
▪ Confocal Microscopy
▪ Image Analysis
▪ Computational model
▪ Neuron model
▪ Synapse model
• **QUANTIFICATION AND STATISTICAL ANALYSIS**
▪ Data acquisition and analysis
▪ Connectivity analysis
▪ Network analysis
▪ Statistical and visual analysis
▪ Software and libraries

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse Anti-Drosophila choline acetyltransferase	DSHB	RRID:AB_528122
Mouse anti-Bruchpilot (nc82)	DSHB	RRID:AB_2314866
Rabbit anti-GFP	Invitrogen	RRID:AB_221569
Goat anti-rabbit secondary	Jackson ImmunoResearch	RRID:AB_2338021
Goat anti-mouse Alexa Fluor 647 secondary	Jackson ImmunoResearch	RRID:AB_2338902
Anti-mVenus antibody (conjugated)	MilliporeSigma	MABE1906
Bacterial and virus strains		
N/A	N/A	N/A
Biological samples		
N/A	N/A	N/A
Chemicals, peptides, and recombinant proteins		
All trans retinal 500mg	MilliporeSigma	R2500-500MG
Tetramethylrhodamine-labeled dextran (10,000 MW)	Invitrogen	D1817
O'Dowd's Drosophila saline	[57]	N/A
Formaldehyde, Para	Fisher Scientific	T353-500
Triton X-100	MilliporeSigma	T8787-100ML
Critical commercial assays		
N/A	N/A	N/A
Deposited data		
N/A		
Experimental models: Cell lines		
N/A	N/A	N/A
Experimental models: Organisms/strains		
SS25451: w[1118]; VT063643-p65.AD; VT059224-GAL4.DBD	Bloomington Drosophila Stock Center	RRID: BDSC_89495
w[1118]; P{y[+t7.7] w[+mC]=20XUAS-IVS-CsChrimson.mVenus}attP2	Bloomington Drosophila Stock Center	RRID: BDSC_55136
w[*]; P{y[+t7.7] w[+mC]=20XUAS-IVS-mCD8::GFP}attP2	Bloomington Drosophila Stock Center	RRID: BDSC_32194
Control-	Bloomington Drosophila Stock Center	RRID:BDSC_
Oligonucleotides		
N/A	N/A	N/A
Recombinant DNA		
N/A	N/A	N/A
Software and algorithms		
Python	http://www.python.org/	RRID:SCR_008394
Anaconda	https://docs.anaconda.com/	RRID:SCR_025572

Jupyter Notebook	https://jupyter.org/	RRID:SCR_018315
neuprint-python API	[27]	N/A
NetworkX	https://networkx.org/	RRID:SCR_016864
MATLAB	https://www.mathworks.com/products/matlab.html	RRID:SCR_001622
MANC	[14]	N/A
Inkscape	https://inkscape.org/en/	RRID:SCR_014479
Brian2	http://briansimulator.org/	RRID:SCR_002998
NeuronBridge v3.2.6	[58]	N/A
Other		
Axon Digidata 1440A Data Acquisition System	Molecular Devices	Model 1440A
Clampex software (version 10 and 11)	http://www.moleculardevices.com/products/software/pclamp.html	RRID:SCR_011323
Nikon AXR confocal microscope with NSPARC detector	Nikon	N/A
Nikon Eclipse E600FN fluorescence microscope	Nikon	N/A
590 nm optogenetic LED setup	Thorlabs	M590F3; M118L02; LEDD1B

Lead contact

Further information and requests for resources and reagents should be directed to and will be fulfilled by the Lead Contact, Rodrigo Pena (penar@fau.edu).

Materials Availability

This study did not generate new unique reagents.

EXPERIMENTAL MODELS AND SUBJECT DETAILS

Fly Genetics

Flies were raised at 25°C with a natural (12-hr) light cycle, and data were collected from both male and female adults 2-4 days old. The Gal4 drivers used in our study were selected through NeuronBridge[58]. Neurons of interest were identified through NeuPrint+, then identified using NeuronBridge software v3.2.6, and finally chosen based on similarity score and visual accuracy. A split-Gal4 driver specific to AN08B098 (Axc) was identified using NeuronBridge software v3.2.6 and chosen based on similarity score. SS25451 contains w[1118]; VT063643-p65.AD; VT059224-GAL4.DBD (BDSC #89495) (see Fig S4). Constructs used were 20XUAS-IVS-CsChrimson.mVenus (#55136), and 20XUAS-IVS-mCD8::GFP (BDSC #32194).

METHOD DETAILS

Electrophysiology

Flies were mounted dorsal side up in dental wax after CO_2 anesthesia. Giant Fibers were activated by extracellular stimuli applied using Grass stimulators with tungsten wire electrodes placed in the brain through the eyes, while a tungsten wire electrode was placed in the fly's abdomen to ground

the circuit [30]. Muscle recordings were obtained using glass microelectrodes placed directly into
both jump (TTM) and flight (DLM) muscles [18,30,55,58,73]. Recording microelectrodes were filled with
O'Dowd's Drosophila saline [74]. We do not directly monitor GF spikes but instead use TTM
responses as a reporter. Extracellular stimulation of the GF pathway activates both the TTM and
the DLM with a short and constant latency between them (~0.6 ms) as described in the early
recordings of this type in the 1980s[37,59]. Action potentials in the GF are indicated by synaptic
potentials in both the TTM and the DLM with the constant latency between them (see Figure 7) as
described in a more recent protocol for the preparation [30]. It has long been assumed that the GF has
the lowest threshold for descending neurons using this method of brain stimulation due to its large
size and the presence of a spike initiation region in the brain. Recently, a spike initiation zone
(SIZ) has been confirmed to be located in the brain at the dendrite-axon bifurcation [35]. The Axon
Digidata 1440A Data Acquisition System was used to digitize recordings which are then visualized
using Clampex software (version 10 & 11, Molecular Devices). Muscle responses were obtained
by stimulating at 24 Hz.

Solution preparations
Phosphate buffered saline was made from tablets purchased from Fisher BioReagents (BP2944-
100). One tablet was dissolved in double distilled water to yield a solution of 10 mM Phosphate
Buffer, 2.7 mM KCl, and 137 mM NaCl. Solutions were stored at the recommended 4°C.

All trans retinal (500 mg), which was purchased from Sigma-Aldrich, is dissolved in 17.6 ml
absolute ethanol to make 100 mM stock solutions. 100 µl of 100 mM stock solution is
transferred to small tubes and wrapped in aluminum foil and kept in -20°C freezer (ATR should
be kept away from light since it is sensitive to light and it would be degraded, thus becoming
ineffective if it is exposed to light for a long time). To prepare fly food supplemented with 400
µM ATR, 5 mL fly food is dissolved in microwave. The food is left to cool down, then 20 µl of
400 µM ATR is mixed well with fly food or 20 µl of absolute ethanol is mixed with food as a
control. The food vial is wrapped in aluminum foil then the food is left until it was solidified
well; otherwise, the flies get stuck in the wet food. The flies are transferred from its vial to ATR
containing vial and kept in a dark place (to keep the ATR from degradation) at room temperature
22-23°C.

O'Dowd saline was made by dissolving: NaCl (0.59 g); $NaHCO_3$ (0.174 g); KCl (0.0224 g); $MgCl_2$
(0.0814 g); $CaCl_2$ (0.0148 g); NaH_2PO_4 (0.0138 g); Trehalose (0.1892 g); and Sucrose (1.54 g) in
100 mL double distilled water. Saline was kept in 4°C for storage or in room temperature as
needed.

Optogenetics
We drove channelrhodopsin expression in neurons of interest using the UAS-Gal4 system. Flies
expressing csChrimson (BDSC #55136) were crossed with our driver line to produce: w1118;

VT063643-p65/+; 20XUAS-IVS-csChrimson.mVenus/VT059224-GAL4.DBD. Adults were placed in the dark at 50% relative humidity on standard cornmeal-molasses food, supplemented with all-trans-retinal (0.4 mM post-eclosion) for two days before testing. Light stimulation was performed for 100 ms while flies are mounted in dental wax as described above. We used a 590 nm high-power fiber-coupled LED (1.5 mW maximum light intensity, Thorlabs M590F3) with an optic fiber of diameter 200 µm and N.A. 0.22 to activate the mVenus csChrimson (M118L02). We tested various placements for the optic fiber and found the strongest activation of csChrimson to occur when the fiber was less than 1 cm from the sample and aimed at the neck. Light intensity was controlled via T-Cube LED Driver (Thorlabs LEDD1B) and kept at maximum power during testing. Each sample received electrical-only, optogenetic-only or paired opto-electrical stimulation. If Axc is stimulated only, no response is recorded, meaning that csChrimson alone is unable to elicit spikes in the GF.

We also conducted two control experiments for the optogenetic protocol (Fig. S2). The first control used flies of the same genotype as the experimental group but were not raised on food supplemented with all-trans-retinal. The second control involved flies that were raised on retinal-supplemented food but lacked the UAS-csChrimson.mVenus transgene (w1118; VT063643/+; VT059224-GAL4/VT059224-GAL4). The results are shown in Figure S2 B-C.

Dye-injection

The VNC and connective of young adult male flies were dissected and kept in cold O'Dowd's saline within a Sylgard 184-filled dissection dish. Samples were immediately transferred to a Nikon Eclipse E600FN fluorescent microscope. A tetramethylrhodamine-labeled dextran dye (10,000MW, Invitrogen, D1817) solution was backfilled into a pulled glass electrode tip (~30 MΩ resistance tip) and filled with O'Dowd's saline. Electrodes were placed into a 3D hydraulic micromanipulator. The electrode was lowered into place, penetrating the GF axons near the neck connective. A Gettings Model 5A amplifier was used to inject positive current pulses (between 1 nA and 5 nA), with a reference electrode placed into the saline-filled dish. GFs were filled with dye for approximately 5 minutes each.

Immunostaining

A mouse primary antibody against ChAT (1:20, DSHB, 4B1) was used to confirm cholinergic cells and coupled to secondary Goat anti-mouse AlexaFluor 647 (1:500, 115-605-003). A mouse primary antibody against bruchpilot (1:50, DSHB, NC82) was used to label presynaptic chemical active zones and coupled to secondary Goat anti-mouse AlexaFluor 647 (1:500, 115-605-003). Primary rabbit anti-GFP (1:500, Invitrogen A11122) and secondary goat anti-rabbit (1:500, Jackson Immuno, 111-225-144) were used to amplify the fluorescence of mCD8::GFP. mVenus-tagged csChrimson. Fluorescence was amplified using a conjugated antibody (1:500, Millipore, MABE1906). Primary antibody solutions contained 3% BSA and 0.3% Triton X-100 in PBS. The secondary antibody solutions contained PBS.

Samples were dissected and processed according to the dye-injection methods above. Samples
were then immediately fixed in PBS + 4% PFA for 25 minutes at room temperature (22°C).
Samples were washed with PBS and placed on a rotator (60RPM) for 30 minutes. Samples were
permeabilized with PBS + 0.5% Triton X-100 for 45 minutes at 22°C. Primary antibodies were
added, and samples were stored at 4°C for 48 hours. Samples were then washed on a rotator with
PBS for 4 hours at 22°C, changing the solution every 30 minutes. Secondary antibodies were
added, and samples were stored at 4°C for 24 hours. Samples were then washed on a rotator with
PBS for 4 hours at 22°C, changing the solution every 30 minutes. Samples were dehydrated with
a series of ethanol steps, 50%, 70%, 90%, and 100% for 8 minutes each step. Samples were
trimmed and embedded in 100% methylsalycylate, covered with cover slips, sealed, and imaged
within 24 hours.

Confocal Microscopy
Samples were imaged on a Nikon AXR confocal using an NSPARC detector and a 60x oil
immersion objective. Z-dimension step size was 194 μm for all results displayed. Laser channels
used were 488 nm, 561 nm, and 640 nm. Channels were imaged in series using filters to avoid
cross-excitation of close wavelengths, and 2x integration was used.

Image Analysis
Z-stack files were imported into Imaris software v10.2.0, and the fluorescence intensity of each
channel was adjusted for best fit. GF and AN08B098 (Axc) channels were surfaced using a
thresholding method that matched fluorescence in each Z dimension. Masks of each channel were
created for each surface to detect colocalization of Brp or ChAT. Images were acquired in volume
view and exported.

Computational model
To study the impact of axo-axonic projections onto DNp01 (GF), we constructed an *in-silico* model
of the ventral nerve cord (VNC) by utilizing the entire MANC v1.2.1 connectome. First, we
scrubbed the connectivity data by omitting connections with a total strength less than 6, then we
excluded all connectivity data where the neurotransmitter type, neuron type, or neuron ID was
unknown or invalid. This resulted in a total of 23,437 valid neurons and 1,152,548 connections
between them, with 619,684 excitatory (cholinergic) and 532,864 inhibitory (glutamatergic or
GABAergic) connections. In the fly, cholinergic receptors are excitatory, whereas glutamatergic
and GABAergic receptors are inhibitory[60].

Neuron model
To achieve biological plausibility while avoiding computational intractability, leaky integrate-and-
fire (LIF) neuron models were used:

$$C_m \frac{dv}{dt} = -g_L(v - V_{rest}) - g_{ach}(v - E_{ach}) - g_{gaba}(v - E_{gaba}) - g_{glut}(v - E_{glut}), \quad (1)$$

Unlike other models[61], we employ distinct parameters for the GF compared to the rest of the
neurons in the connectome, i.e., the membrane resistance. In Eq. (1), v is the neuron membrane
potential, g_L, g_{ach}, g_{GABA}, and g_{glut} are the conductances for leak channels and acetylcholine
(nicotinic), GABA (GABA$_A$), and glutamate ionotropic receptors (nAChR, GABA$_A$R, and GluCl,
respectively). Similarly, V_{rest}, E_{ach}, E_{GABA}, and E_{glut} are their respective reversal potentials.
Since GABA$_A$R and GluR are chloride-permeable channels, their reversal potentials were obtained
from chloride reversal potential[62]. C_m is the membrane capacitance. These are parameterized with
values obtained from the literature, which investigates the biophysical properties of fruit fly
synapses. Furthermore, we collected physiologically relevant values for the membrane
capacitance, resistance, and leak reversal potential. Simulations of these neurons were made using
the BRIAN2 simulator [63].
766

767 **Synapse model**
768 For each of the synaptic currents ($x = ach, gaba, glut$), the conductance obeys
769

770
$$\frac{dg_x}{\tau_x} = -g_x, \qquad (2)$$

771 which increases by J_x upon a presynaptic spike. The list of parameters is included in Table 1.

Parameter	Symbol	Value	Ref.
Acetylcholine receptor (nAChR) conductance	J_{ach}	0.27 nS	64
nAChR decay time	τ_{ach}	1.1 ms	64
nAChR reversal potential	E_{ach}	0 mV	65
GABA receptor (GABA$_A$R) conductance	J_{GABA}	0.8 nS	Estimated
GABA$_A$R decay time	τ_{GABA}	5.4 ms	66
GABA$_A$R reversal potential	E_{GABA}	-60 mV	60,62
Glutamate receptor (GluCl) conductance	J_{glut}	0.8 nS	Estimated
GluCl decay time	τ_{glut}	5 ms	67
GluCl reversal potential	E_{glut}	-60 mV	62
Membrane Resistance (GF)	R_m	25 MOhm	64
Membrane Resistance (other neurons)	R_m	300 MOhm	68

Resting membrane potential	V_{rest}	-55 mV	[61,68]
Action potential threshold	V_{th}	-40 mV	[68]
Reset potential	V_{reset}	-55 mV	Estimated
Refractory period	τ_{ref}	2 ms	Estimated
Membrane capacitance	C_m	72 pF	[68]

Table 1. List of parameters used in the connectome simulation of the VNC in this paper. For each of our simulations, we ran the entire VNC network with a step-size of 0.1 ms during 2000 ms. Furthermore, for each simulation, each neuron in the VNC network was provided with an excitatory Poisson input of 30 Hz at a synaptic conductance of 5 nS to simulate typical endogenous activity driven by neurons in the fly brain. During stimulation experiments, neurons DNp01, DNg02, and/or DNd03 were stimulated with additional Poisson input at 10 Hz at a synaptic conductance of 40 nS. In Figure 8, the same network was used, but the GF was electrically stimulated just below threshold with a Poisson-distributed stimulation at 24 Hz.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data acquisition and analysis

Data for VNC neurons used in the study came from the publicly available *Drosophila* male adult nerve cord connectome[14]. NeuPrint+ API[26,28] access is granted to registered users, and Python accessibility is maintained through the neuPrint-python package[27]. The latest *Drosophila* MANC data can be accessed here: https://www.janelia.org/project-team/flyem/manc-connectome.

Connectivity analysis

We retrieved all simple synaptic connections using neuprint.fetch_simple_connections[26–28] separately for: outputs (presynaptic AN, IN, and DN as presynaptic) and Inputs (postsynaptic DN). Connections were summed by partner neuron and filtered to those with total strength ≥ 6. For each DN, we then counted the number of distinct AN, IN, and DN partners in both the output and input directions.

Network analysis

Network analysis was performed using Python with the NetworkX[69] library to extract quantitative measures of connectivity. From the structural connectivity matrix, we calculated a comprehensive set of graph-theoretical metrics to characterize the network topology. These included basic properties (number of nodes and edges, network density), integration measures (average path length, diameter), segregation indices (clustering coefficient, modularity with community detection using the Louvain algorithm), centrality metrics (degree, betweenness, closeness, and eigenvector centrality), and organizational principles (degree assortativity, rich club coefficient).

The rich club analysis was specifically conducted across multiple degree thresholds to identify potential hierarchical organization among high-degree hub regions. All weighted graph measures

incorporated the synaptic strengths from the original connectivity matrix. This multimodal graph-theoretical approach allowed for a detailed characterization of both local and global properties of the network's structural connectome, revealing important aspects of neural communication efficiency, functional segregation, and network resilience.

Statistical and visual analysis

SWC analysis was conducted in Python using the Navis client[70]. We stored morphology and input/output counts using the Pandas library. Relationships between morphology metrics (cable length, branch points, node count) and partner counts were assessed with Pearson correlation and linear regression (scipy.stats.pearsonr, linregress). Results were visualized with Matplotlib[71]. Network-style connection diagrams were rendered interactively with ipycytoscape.

Software and libraries

NeuPrint+ analyses were performed in Python 3.12.4 using the neuprint-python API. We used the Janelia NeuPrint client (neuprint), and Navis (v1.10.0) for data access and network building; data manipulation and grouping were done with pandas v2.2.2 and NumPy v1.26.4; graph-based operations with NetworkX v3.2.1; interactive visualization with ipycytoscape v1.3.3 and ipywidgets v8.1.5; plotting and PDF export with Matplotlib v3.10.1; and statistical analyses (Pearson's r, linear regression) with SciPy[72] v1.13.1. Auxiliary parsing and utility functions relied on Python's built-in json, re, itertools, and warnings modules.

Acknowledgments

R.P. was funded by an IBRO Collaborative Research Grants. The authors thank the FAU Jupiter Life Sciences Initiative (JLSI) for funding.

Author contributions

C.C.C. and R.P. were responsible for the conceptualization of the work. C.C.C., J.L., T.R., D.S., S.J., and C.S. analyzed the data and made the figures. R.P. worked on the graph theoretical measures. D.S. performed and analyzed the electrophysiology recordings. S.J. and C.S. performed immunofluorescence. K.R. maintained fly stocks and set up experiment crosses. T.R. built and ran the network simulations. R.P., C.S., and R.M. supervised the research and secured funding. All authors participated in the writing, reviewed, and approved the manuscript.

Declaration of interests

The authors declare no competing interests.

References

1. Brette, R. Theory of axo-axonic inhibition. *PLoS Comput. Biol.* **21**, e1013047 (2025).

2. Schneider-Mizell, C. M. *et al.* Structure and function of axo-axonic inhibition. *Elife* **10**, (2021).

3. Stretton, A. O. W. *et al.* Structure and physiological activity of the motoneurons of the nematode Ascaris. *Proc. Natl. Acad. Sci. U. S. A.* **75**, (1978).

4. Pearson, K. G. & Goodman, C. S. Presynaptic inhibition of transmission from identified interneurons in locust central nervous system. *J. Neurophysiol.* **45**, (1981).

5. Atwood, H. L. & Kwan, I. Reciprocal axo-axonal synapses between excitatory and inhibitory neurons in crustaceans. *Brain Res.* **174**, (1979).

6. Altrup, U., Speckmann, E. J. & Caspers, H. Differentiation between anti- and orthodromic responses to nerve stimulation in neurons with axo-axonal synapses (Helix pomatia). *Neurosci. Lett.* **11**, (1979).

7. Andersen, P., Eccles, J. C., Schmidt, R. F. & Yokota, T. Depolarization of presynaptic fibers in the cuneate nucleus. *J. Neurophysiol.* **27**, (1964).

8. Wendell-Smith, C. P., Blunt, M. J., Baldwin, F. & Paisley, P. B. Neurone - Satellite cell relationship. *Nature* **205**, (1965).

9. Gobel, S. & Dubner, R. Fine structural studies of the main sensory trigeminal nucleus in the cat and rat. *Journal of Comparative Neurology* **137**, (1969).

10. Cover, K. K. & Mathur, B. N. Axo-axonic synapses: Diversity in neural circuit function. *Journal of Comparative Neurology* **529**, 2391–2401 (2021).

11. Dudok, B. *et al.* Recruitment and inhibitory action of hippocampal axo-axonic cells during behavior. *Neuron* **109**, (2021).

12. Lacoste, A. M. B. *et al.* A Convergent and Essential Interneuron Pathway for Mauthner-Cell-Mediated Escapes. *Current Biology* **25**, 1526–1534 (2015).

13. Takemura, S.-Y. *et al.* A Connectome of the Male \textit{Drosophila} Ventral Nerve Cord. *bioRxiv* (2023).

14. Cheong, H. S. J. *et al.* Transforming descending input into behavior: The organization of premotor circuits in the Drosophila Male Adult Nerve Cord connectome. *bioRxiv* (2023).

15. Marin, E. C. *et al.* Systematic annotation of a complete adult male Drosophila nerve cord connectome reveals principles of functional organisation. *Elife* **13**, (2024).

16. Lin, A. *et al.* Network statistics of the whole-brain connectome of Drosophila. *Nature 2024 634:8032* **634**, 153–165 (2024).

17. Clements, J. *et al.* neuPrint: Analysis Tools for EM Connectomics. *bioRxiv* 2020.01.16.909465 (2020) doi:10.1101/2020.01.16.909465.

18. Allen, M. J., Godenschwege, T. A., Tanouye, M. A. & Phelan, P. Making an escape: Development and function of the Drosophila giant fibre system. *Seminars in Cell and Developmental Biology* vol. 17 Preprint at https://doi.org/10.1016/j.semcdb.2005.11.011 (2006).

19. Asinof, S. K. & Card, G. M. Neural Control of Naturalistic Behavior Choices. *Annu. Rev. Neurosci.* **47**, 369–388 (2024).

20. Cheong, H. S. J. *et al.* Transforming descending input into behavior: The organization of premotor circuits in the Drosophila Male Adult Nerve Cord connectome. *bioRxiv* (2023).

21. Stürner, T. *et al.* Comparative connectomics of Drosophila descending and ascending neurons. *Nature 2025* 1–15 (2025) doi:10.1038/s41586-025-08925-z.

22. Takemura, S. *et al.* A Connectome of the Male Drosophila Ventral Nerve Cord. *Elife* **13**, (2024).

23. Marin, E. C. *et al.* Systematic annotation of a complete adult male Drosophila nerve cord connectome reveals principles of functional organisation. *Elife* **13**, (2024).

24. Schneider-Mizell, C. M. *et al.* Quantitative neuroanatomy for connectomics in Drosophila. *Elife* **5**, (2016).

25. Plaza, S. M. *et al.* neuPrint: An open access tool for EM connectomics. *Front. Neuroinform.* **16**, (2022).

26. Berg, S. & Schlegel, P. neuprint-python. Preprint at https://doi.org/10.5281/zenodo.7592899 (2017).

27. Clements, J. *et al.* neuPrint: Analysis Tools for EM Connectomics. *bioRxiv* 2020.01.16.909465-2020.01.16.909465 (2020) doi:10.1101/2020.01.16.909465.

28. Bates, A. S. *et al.* Distributed control circuits across a brain-and-cord connectome. Preprint at https://doi.org/10.1101/2025.07.31.667571 (2025).

29. Allen, M. J. & Godenschwege, T. A. Electrophysiological recordings from the Drosophila giant fiber system (GFS). *Cold Spring Harb. Protoc.* **5**, (2010).

30. Klapoetke, N. C. *et al.* Independent optical excitation of distinct neural populations. *Nat. Methods* **11**, (2014).

31. Ache, J. M. *et al.* Neural Basis for Looming Size and Velocity Encoding in the Drosophila Giant Fiber Escape Pathway. *Current Biology* **29**, (2019).

32. Scheffer, L. K. *et al.* A connectome and analysis of the adult drosophila central brain. *Elife* **9**, (2020).

33. Manoim, J. E., Davidson, A. M., Weiss, S., Hige, T. & Parnas, M. Lateral axonal modulation is required for stimulus-specific olfactory conditioning in Drosophila. *Current Biology* **32**, 4438-4450.e5 (2022).

34. Gruber, L. *et al.* The unique synaptic circuitry of specialized olfactory glomeruli in Drosophila melanogaster. Preprint at https://doi.org/10.7554/eLife.88824.1 (2023).

35. Calle-Schuler, S. A., Santana-Cruz, A. E., Kmecová, L., Hampel, S. & Seeds, A. M. A comprehensive mechanosensory connectome reveals a somatotopically organized neural circuit architecture controlling stimulus-aimed grooming of the *Drosophila* head. Preprint at https://doi.org/10.1101/2025.05.19.654894 (2025).

36. Oleskevich, S., Descarries, L., Watkins, K. C., Se´gue´la, P. & Daszuta, A. Ultrastructural features of the serotonin innervation in adult rat hippocampus: An immunocytochemical description in single and serial thin sections. *Neuroscience* **42**, 777–791 (1991).

935   37.   Séguéla, P., Watkins, K. C. & Descarries, L. Ultrastructural relationships of
936   serotonin axon terminals in the cerebral cortex of the adult rat. *Journal of*
937   *Comparative Neurology* **289**, 129–142 (1989).
938   38.   Lin, A. *et al.* Network statistics of the whole-brain connectome of
939   Drosophila. *Nature 2024 634:8032* **634**, 153–165 (2024).
940   39.   Lu, J. *et al.* Selective inhibitory control of pyramidal neuron ensembles and
941   cortical subnetworks by chandelier cells. *Nat. Neurosci.* **20**, (2017).
942   40.   Woodruff, A. R., Anderson, S. A. & Yuste, R. The Enigmatic Function of
943   Chandelier Cells. *Front. Neurosci.* **4**, (2010).
944   41.   Cuntz, H., Forstner, F., Borst, A. & Häusser, M. One rule to grow them all:
945   A general theory of neuronal branching and its practical application. *PLoS*
946   *Comput. Biol.* **6**, (2010).
947   42.   Lacin, H. & Truman, J. W. Lineage mapping identifies molecular and
948   architectural similarities between the larval and adult Drosophila central
949   nervous system. *Elife* **5**, (2016).
950   43.   Rajagopalan, S., Nicolas, E., Vivancos, V., Berger, J. & Dickson, B. J.
951   Crossing the midline: Roles and regulation of Robo receptors. *Neuron* **28**,
952   (2000).
953   44.   Harriger, L., van den Heuvel, M. P. & Sporns, O. Rich Club Organization of
954   Macaque Cerebral Cortex and Its Role in Network Communication. *PLoS*
955   *One* **7**, e46497 (2012).
956   45.   Liang, X. *et al.* The Rich-Club Organization in Rat Functional Brain
957   Network to Balance Between Communication Cost and Efficiency. *Cerebral*
958   *Cortex* **28**, 924–935 (2018).
959   46.   Card, G. & Dickinson, M. H. Visually Mediated Motor Planning in the
960   Escape Response of Drosophila. *Current Biology* **18**, (2008).
961   47.   Kramer, P. F. *et al.* Synaptic-like axo-axonal transmission from striatal
962   cholinergic interneurons onto dopaminergic fibers. *Neuron* **110**, 2949-
963   2960.e4 (2022).
964   48.   Ammer, G., Vieira, R. M., Fendl, S. & Borst, A. Anatomical distribution and
965   functional roles of electrical synapses in Drosophila. *Current Biology* **32**,
966   (2022).
967   49.   Lopez, J., Boerner, J., Robbins, K., Pena, R. F. O. & Murphey, R.
968   Frazzled/DCC regulates gap junction formation at a Drosophila giant
969   synapse through transcription. Preprint at
970   https://doi.org/10.1101/2025.04.08.647628 (2025).
971   50.   Holcroft, C. E. *et al.* Innexins Ogre and Inx2 are required in glial cells for
972   normal   postembryonic development of the Drosophila central nervous
973   system. *J. Cell Sci.* **126**, 3823–3834 (2013).
974   51.   Phelan, P. *et al.* Molecular mechanism of rectification at identified electrical
975   synapses in   the Drosophila giant fiber system. *Curr. Biol.* **18**, 1955–1960
976   (2008).
977   52.   Orr, B. O., Borgen, M. A., Caruccio, P. M. & Murphey, R. K. Netrin and
978   frazzled regulate presynaptic gap junctions at a Drosophila giant synapse.
979   *Journal of Neuroscience* **34**, (2014).

980   53.   Winding, M. *et al.* The connectome of an insect brain. *Science (1979).* **379**,
981   (2023).
982   54.   Gu, H. & O'Dowd, D. K. Cholinergic Synaptic Transmission in Adult
983   *Drosophila* Kenyon Cells *In Situ. The Journal of Neuroscience* **26**, 265–272
984   (2006).
985   55.   Clements, J. *et al.* NeuronBridge: an intuitive web application for neuronal
986   morphology search across large data sets. *BMC Bioinformatics* **25**, (2024).
987   56.   McCarthy, E. V *et al.* Synchronized bilateral synaptic inputs to Drosophila
988   melanogaster neuropeptidergic rest/arousal neurons. *Journal of Neuroscience*
989   **31**, (2011).
990   57.   Shiu, P. K. *et al.* A Drosophila computational brain model reveals
991   sensorimotor processing. *Nature* **634**, (2024).
992   58.   Schnizler, K. *et al.* A novel chloride channel in Drosophila melanogaster is
993   inhibited by protons. *Journal of Biological Chemistry* **280**, (2005).
994   59.   Stimberg, M., Brette, R. & Goodman, D. F. M. Brian 2, an intuitive and
995   efficient neural simulator. *Elife* **8**, (2019).
996   60.   Moreno-Sanchez, A. *et al.* Morphology and synapse topography optimize
997   linear encoding of synapse numbers in Drosophila looming responsive
998   descending neurons. Preprint at https://doi.org/10.7554/eLife.99277.1
999   (2024).
1000   61.   Lee, D. & O'Dowd, D. K. Fast excitatory synaptic transmission mediated by
1001   nicotinic acetylcholine receptors in Drosophila neurons. *Journal of*
1002   *Neuroscience* **19**, (1999).
1003   62.   Lee, D., Su, H. & O'Dowd, D. K. GABA receptors containing Rdl subunits
1004   mediate fast inhibitory synaptic transmission in Drosophila neurons. *Journal*
1005   *of Neuroscience* **23**, (2003).
1006   63.   Saumweber, T. *et al.* Behavioral and synaptic plasticity are impaired upon
1007   lack of the synaptic protein SAP47. *Journal of Neuroscience* **31**, (2011).
1008   64.   Churgin, M. A. *et al.* A neural correlate of individual odor preference in
1009   Drosophila. *Elife* **12**, (2025).
1010   65.   Hagberg, A. A., Schult, D. A. & Swart, P. J. Exploring Network Structure,
1011   Dynamics, and Function using NetworkX. in *Proceedings of the 7th Python*
1012   *in Science Conference* (2008). doi:10.25080/tcwv9851.
1013   66.   Schlegel, P. *et al.* navis-org/navis: Version 1.10.0. Preprint at (2025).
1014   67.   Hunter, J. D. Matplotlib: A 2D graphics environment. *Comput. Sci. Eng.* **9**,
1015   (2007).
1016   68.   Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific
1017   computing in Python. *Nat. Methods* **17**, (2020).
1018
1019

HIGHLIGHTS

- Axo-axonic connections made with descending neurons (DNs) are rare in the VNC.
- There is a linear correlation between the number of synapses and the number of partners a neuron has.
- Axo-axonic neurons form exclusive connections with the Giant Fibers.
- Axo-axonic neuron activity influences DN excitability *in silico* and *in vivo*.