Changes in Histotone Isoform Abundnance and Histone Post-Translational Modifications

during Anoxia Tolerance and Recovery in WS40NE cells of *Austrofundulus limnaeus*

Chelsea Hughes[1], Elizabeth A. Mojica[2], Dietmar Kültz[2], Jason E. Podrabsky[1]

1. Department of Biology, Center for Life in Extreme Environments, Portland State University,

P.O. Box 751, Portland, OR 97201, USA

2. Department of Animal Sciences and Genome Center, University of California, Davis, One

Shields Avenue, Meyer Hall, Davis, CA 95616, USA

Keywords: mass spectrometry; epigenetics; histone post-translational modifications; anoxia;
stress tolerance

Abstract

Anoxia is an often-lethal stressor to vertebrates, yet some vertebrates have adapted cellular mechanisms to survive in anoxic conditions. Embryos of the annual killifish *Austrofundulus limnaeus* have the greatest tolerance to anoxia of all vertebrates, yet the epigenetic mechanisms that support their anoxia tolerance are unknown. Using mass spectrometry, 1043 unique biologically relevant histone post-translational modifications (unimod+histone residue) were detected in WS40NE cells, representing thirteen types of biologically relevant histone post-translational modifications (hPTMs) present during normoxia, 1 d anoxia, 4 d anoxia, and aerobic recovery from anoxia. Of these 1043 hPTMs, 816 were significantly differentially expressed in at least one comparison. Thirty-six significant hPTMs were considered highly condition dependent. Additionally, at least four histone isoforms were differentially expressed, representing H2A, H2B, and H3 isoforms. Our data suggests that specific histone modifications as well as changes in histone isoform abundance in WS40NE cells may be necessary to successfully respond to extreme changes in oxygen availability.

Introduction

All organisms must respond to their environment, which offers both opportunities and threats. By changing which genes are expressed, organisms can modify their phenotype to take advantage of opportunities and neutralize threats. However, environmental stressors can often lead to aberrant gene expression in organisms that lack these adaptations. This is particularly true during anoxic stress in mammals, which can cause apoptosis and inappropriate reentry into the cell cycle in mammalian cells [1, 2]. Specifically in mammalian neurons, apoptosis is associated with cellular dysregulation that leads these terminally differentiated cells to re-enter the cell

cycle, which often leads to catastrophic cell failure [2]. Additionally, mitochondrial

dysregulation is associated with anoxia due to inhibition of the electron transport chain (ETC)

and depolarization of the inner mitochondrial membrane that can lead to apoptosis [3-5]. The

subsequent cell death causes tissue damage, which continues when oxygen is returned to the

affected area, a phenomenon known as ischemia-reperfusion injury in mammals [6]. Ischemia-

reperfusion increases reactive oxygen species (ROS) such as the superoxide anion (O^{-2}),

hydrogen peroxide (H_2O_2), and hydroxyl radical (OH^-) [7]. While low levels of ROS are a

normal byproduct of mitochondrial metabolism [8], high levels of ROS can lead to neuronal

death within 5 minutes [9, 10]. In contrast, a handful of vertebrates have adapted mechanisms to

modify their gene expression to survive environmental stressors, including osmotic, thermal,

ROS, and anoxia with little or no damage [11-19].Therefore, there is a need to understand how

gene expression is regulated in anoxia tolerant vertebrates to understand how they avoid these

fates. This study focuses on epigenetic histone modifications in response to anoxia in a cell line

isolated from anoxia-tolerant embryos of the annual killifish, *Austrofundulus limnaeus*.

Epigenetic responses to a variety of stressors are documented in multiple organisms [20-23].

However, few histone post-translational modifications (hPTMs) have been investigated in

response to hypoxia or anoxia. Additionally, these data often come from anoxia intolerant

organisms that are experiencing a disease state and are likely failing [24, 25]. Additionally, this

research often focuses only on well-known residues and specific types of hPTMs such as

methylation [26, 27]. For example, histone-3 lysine-4 trimethylation (H3K4me3) and

H3K36me3 are known to change in HeLa cells within 1 h of hypoxic exposure, but HeLa cells

begin to die within 1 d of severe hypoxia [28, 29]. Limited research exists on histone

modifications in anoxia tolerant organisms, but this work again focuses on well-known

modifications such as acetylation and methylation [30]. It is unknown how less studied hPTMs,

such as lactylation, 4-hydroxynonenal, and oxidation are impacted by anoxic stress, particularly

in anoxia tolerant species. Even better characterized hPTMs like ubiquitylation are not well

understood in anoxia. While ubiquitylation of the proteome has been studied in ischemia and

anoxia recovery [31, 32], to our knowledge, anoxia-induced histone ubiquitylation has never

been explored in a continuous cell line.

Embryos of the annual killifish *Austrofundulus limnaeus* have the greatest tolerance to anoxia of

all vertebrates, making them the ideal model to study the cellular mechanisms necessary for

anoxia tolerance. This remarkable tolerance of anoxia is supported by the ability of embryos to

arrest development and enter a state of anoxia-induced quiescence [33]. The global histone

landscape of developing embryos (Wourms' Stage 36) under normoxia and after 24 h of anoxia

suggests a stabilization of both histone isoform and hPTM abundance [34]. This is surprising, as

hPTMs are known to respond to stressors quickly [22, 28, 35, 36]. However, this stabilization

represents an averaged histone landscape for multiple cell types [34]. It is unclear how specific

cell lines of *A. limnaeus* respond to anoxia, particularly in the absence of tissue-dependent cell

signaling. Additionally, no research has investigated how the histone landscape of *A. limnaeus*

changes when exposed to longer bouts of anoxia or during recovery from anoxia.

PSU-AL-WS40NE (from now on WS40NE) is a neuroepithelial cell line that was isolated from

an embryonic (Wourms' Stage 40) *A. limnaeus* tissue explant. These cells can survive for at least

49 days without oxygen [37], while SH-SY5Y human neuroblastoma cells can survive anoxia for

only approximately 1.3 days [38], and mouse N2A cells can survive for just over 3 days [37].

Here, we describe for the first time the global histone landscape of WS40NE cells exposed to

anoxia and recovery from anoxia. We hypothesize cells undergo extensive changes to histone

isoforms and the hPTM landscape in response to anoxia and aerobic recovery from anoxia.

Results

Histone Abundance

Our methods resulted in the identification of 1,927 peptides across at least twenty-three distinct

histone proteins based on their unique protein accession number. These proteins were

representative of H1, H2A, H2B, and H3 isoforms. No H4 isoforms were detected. Sequence

coverage ranged from 17-100% (Fig. 1). At least four histones were differentially expressed

across the four experimental time points (Fig. 1).

Figure 1. Histone isoform abundance in WS40NE cells. Twenty distinct histone proteins were identified based on

their unique protein accession number, as well as three groups of histones that could not be distinguished. Bar

graphs represent the quantile normalized mean ± SEM of histone relative abundance during normoxia (orange), 1 d

anoxia (green), 4 d anoxia (blue), and 1 d of recovery (dark purple). Letters over bars represent statistically

significant isoform abundances determined by pairwise comparisons. Pie charts represent the sequence coverage

(light purple) and the percent of peptide sequences that were not detected (blue) for each isoform. Sequence

coverage is defined as the total observed protein sequence length divided by the total protein length. Note that due to

different peptide ionization efficiencies, histone isoform abundance can only be compared across conditions and not

to one another within a condition.

A histone H3.3 protein increased significantly within 1 d of anoxia, while an H2A-like protein decreased across this same time frame. Two histone H2B isoforms decreased in abundance after 4 d of anoxia. However, principal component analysis demonstrated broad sample similarity across conditions when considering the abundances of all histone isoforms (Fig. 2).

Figure 2. Principal component analysis of histone isoform abundance in WS40NE cells exposed to anoxia and

aerobic recovery from anoxia. The mean quantile normalized abundance values of all histone isoforms were used

to construct a PCA Plot. Normoxic (purple), 1 d anoxic (green), 4 d anoxic (blue), and recovery samples (pale blue)

are plotted according to PC1 and PC2 with a 95% confidence ellipse surrounding each cluster.

Histone Post-translational Modifications

Histone groups were evaluated in a residue-specific manner to determine specific sites of

differential modification. Principal component analysis (PCA) illustrates that, globally, the

hPTM landscape is most unique after 4 d of anoxia compared to normoxia (Fig. 3A). It is worth

noting the high variation in PC2 for the recovery time point (Fig. 3A). Out of 1293 unique

hPTMs detected, 1043 unique hPTMs (hPTM type+residue) were considered biologically

relevant, representing thirteen types of biologically relevant PTMs (Fig. 3B,C). Four hundred

and sixty-one residues were modified with biologically relevant hPTMs, with many residues

being modified with multiple types of hPTMs. These types of modifications included commonly

studied modifications such as acetylation, methylation, phosphorylation, and ubiquitylation, and

less studied modifications such as lactylation and 4-hydroxynonenal. All thirteen types of

modifications were statistically significant on at least one residue in at least one comparison,

with oxidation/hydroxylation and dioxidation having the greatest number of significantly

modified residues, followed closely by dehydration and phosphorylation (Fig. 3B). As expected,

the percent sequence coverage is positively correlated with the number of significant hPTMs

(Fig. 3D, $r = 0.7$, $p = 0.0003$), and thus for certain histones our ability to draw global conclusions

is somewhat limited. Of the unique biologically relevant PTMs detected, 816 were significant in

at least one comparison (Fig. 3E). Hundreds of significant differential hPTMs were present

across all four timepoints with 4 d of anoxia having the highest number compared to other time

points (Fig. 3E).

Figure 3. Quantification of global patterns of significant histone post-translational modifications in WS40NE

cells exposed to anoxia and recovery from anoxia. (A) The number of significant hPTMs across all histone

isoforms plotted by modification type. **(B)** Venn diagram illustrating the occurrence of the 297 unique and

biologically-relevant hPTMs across the experimental conditions used in this study. **(C)** PCA plot of the M-values of

all hPTMs indicate that hPTM patterns at 4 d of anoxia are unique compared to the other experimental conditions. It

is also noteworthy that variation in global patterns of hPTMs appears to be high during aerobic recovery. Individual

samples (data points) are surrounded by 95% confidence ellipses for each treatment group. **(D)** There is a positive

correlation (r = 0.7, p = 0.0003) between percent sequence coverage for each histone isoform and the number of

significant hPTMs identified. However, several isoforms do not follow this trend, suggesting these are highly

modified isoforms. **(E)** The bar graph represents the number of significantly expressed hPTMs in each t-test

comparison.

Thirty hPTMs were highly abundant (having an average relative abundance of 50% or greater)

across all conditions. Of the high abundance hPTMs, 15 displayed an average relative abundance

across all samples of exactly 100% during all conditions, meaning that each specific residue was

always modified by that PTM (Table 1). Five histones had at least one hPTM site with 100%

relative abundance, representing H1, H2A, and H3 histones. At least 8 distinct types of PTMs are

represented in this group with dehydration accounting for 30% of these stable modifications.

Table 1. Histone post-translational modifications with 100% relative abundance in WS40NE cells. Fifteen

hPTMs, across 5 histone isoforms, always had the same hPTM across all experimental conditions. At least eight

distinct types of PTMs are represented.

Protein Accession	Protein Description	Amino Acid + Position	PTM Description
XP_013861857.1	H1-gamma,_late-like	196K	Acetylation
XP_013861857.1	H1-gamma,_late-like	175K	Biotinylation
XP_013861857.1	H1-gamma,_late-like	173P	Oxidation/Hydroxylation
XP_013861857.1	H1-gamma,_late-like	191K	Acetylation
XP_013861857.1	H1-gamma,_late-like	157K	Acetylation
XP_013861857.1	H1-gamma,_late-like	228K	Dioxidation
XP_013861857.1	H1-gamma,_late-like	24R	4-hydroxynonenal
XP_013861857.1	H1-gamma,_late-like	46S	Phosphorylation
XP_013878240.1	H1-like	64S	Dehydrated
XP_013885354.1	H2B.L4-like	46S	Dehydrated
XP_013885354.1	H2B.L4-like	55S	Dehydrated
XP_013885354.1	H2B.L4-like	43T	Dehydrated
XP_013885354.1	H2B.L4-like	31Y	Dehydrated
XP_013878241.1	H3-like	24R	Oxidation/Hydroxylation
XP_013878241.1	H3-like	17K	Oxidation/Hydroxylation

Individual hPTMs at specific residues were also evaluated, and the most condition-dependent

hPTMs were identified. Significant differential modifications were detected on all groups of

histones (H1, H2A, H2B, H3), with H1-gamma late-like (XP_013861857.1) being the only

isoform that did not have any significant hPTMs detected. Thirty-seven hPTMs were considered

highly condition dependent, being significantly differentially expressed across 5-6 comparisons

(Table 2).

Table 2. Highly condition-dependent histone post-translational modifications in WS40NE cells exposed to

anoxia and aerobic recovery from anoxia. To be considered highly condition-dependent, hPTMs had to be

significant across at least 5 of 6 different comparisons. Thirty-seven hPTMs were selected as the most condition

dependent. Differential modifications of histone residues were evaluated using a two-tailed t-test with the

Benjamini-Hochberg correction applied to account for multiple hypothesis testing.

Protein Accession	Protein Description	Amino Acid + Position	PTM Description
XP_013863633.1	H1.0-B	25K	Acetylation
XP_013863633.1	H1.0-B	19K	4-hydroxynonenal
XP_013879563.1	H1-like	17K	Acetylation
XP_013879563.1	H1-like	51K	4-hydroxynonenal
XP_013879563.1	H1-like	63S	Phosphorylation
XP_013879563.1	H1-like	62R	Oxidation/Hydroxyl
XP_013878193.1	H2A-like	102S	Dehydration
XP_013879560.1	H2A-like	123K	Carboxyethyl/ lactylation
XP_013854970.1	H2A.V	116K	Oxidation/hydroxylation
XP_013854970.1	H2A.V	121K	Biotinylation

XP_013854970.1	H2A.V	122K	Dioxidation
XP_013885354.1	H2B.L4-like	74Y	Phosphorylation
XP_013885354.1	H2B.L4-like	78S	Ubiquitylation
XP_013885354.1	H2B.L4-like	79T	Dehydration
XP_013885354.1	H2B.L4-like	79T	Phosphorylation
XP_013885354.1	H2B.L4-like	82S	Dehydration
XP_013885354.1	H2B.L4-like	83R	Oxidation/hydroxylation
XP_013885354.1	H2B.L4-like	116K	Methylation
XP_013879561.1 / XP_013878239.1 / XP_013878163.1 / XP_013869445.1	H2B_1/2	41Y	Phosphorylation
XP_013879561.1 / XP_013878239.1 / XP_013878163.1 / XP_013869445.1	H2B_1/2	84K	4-hydroxynonenal
XP_013879561.1 / XP_013878239.1 / XP_013878163.1 / XP_013869445.1	H2B_1/2	89T	Phosphorylation
XP_013879561.1 / XP_013878239.1 / XP_013878163.1 / XP_013869445.1	H2B_1/2	90S	Dehydration
XP_013879561.1 / XP_013878239.1 / XP_013878163.1 / XP_013869445.1	H2B_1/2	90S	Phosphorylation

XP_013879561.1 /			
XP_013878239.1 /			
XP_013878163.1 /			
XP_013869445.1	H2B_1/2	114T	Dehydration
XP_013879561.1 /			
XP_013878239.1 /			
XP_013878163.1 /			
XP_013869445.1	H2B_1/2	124K	Methylation
XP_013857090.1	H2B_1/2-like	41Y	Phosphorylation
XP_013857090.1	H2B_1/2-like	78R	Dioxidation
XP_013857090.1	H2B_1/2-like	81H	4-hydroxynonenal
XP_013857090.1	H2B_1/2-like	84K	Acetald+28 / Dimethylation
XP_013874662.1	H2B_1/2	35S	Phosphorylation
XP_013874662.1	H2B_1/2	41Y	Phosphorylation
XP_013874662.1	H2B_1/2	55S	Phosphorylation
XP_013874662.1	H2B_1/2	63S	Ubiquitylation
XP_013874662.1	H2B_1/2	86S	Phosphorylation
XP_013874662.1	H2B_1/2	89T	Phosphorylation
XP_013874662.1	H2B_1/2	90S	Phosphorylation
XP_013874662.1	H2B_1/2	124K	Methylation

Globally, hPTMs did not cover the majority of the residues that are theoretically available.

Lysine residues were the most modified with 48.7-51.4% of the available residues modified

across the different experimental conditions (Fig. 4A). The modification of specific amino acid

residues varied based on experimental conditions (Fig. 4A). Lysines had the most types of

modifications, with 10 distinct hPTMs detected globally, and of these, oxidation/hydroxylation,

dioxidation, and acetylation were the most common (Fig. 4A). Serine (S) had the second most

types of modifications, with 5 distinct hPTMs occurring globally; dehydration and

phosphorylation dominated serine modifications (Fig. 4A). Globally, there were significant

changes in all types of hPTMs across all six comparisons (Fig. 4B,C): (a) 5 comparing normoxia

to 24 h anoxia, (b) 12 comparing normoxia to 4 d anoxia, (c) 8 comparing normoxia to 24 h

recovery,(d) 6 comparing 24 h anoxia to 4 d anoxia, (e) 4 comparing 24 h anoxia to 24 h

recovery, and (f) 2 comparing 4 d anoxia to 24 h recovery. The majority of the most common

modifications decreased significantly during anoxia and remained low even after 24 h of aerobic

recovery (Fig. 4B,C). Notably, acetylation, dehydration, oxidation, and dioxidation all increase

during anoxia. (Fig. 4B,C).

Figure 4. Global changes in the histone PTM landscape of WS40NE cells during anoxia and aerobic recovery

from anoxia. (A) Thirteen types of hPTMs were detected across nine types of amino acid residues in WS40NE

cells. While most residue types are unmodified, lysine (K) residues were the most consistently modified (54.7 -

58.3%) across the different experimental conditions. **(B)** From a global perspective, the abundance of most hPTMs

decreases in response to anoxia and fails to return to normoxic values after 1 d of aerobic recovery. Acetylation,

oxidation, dioxidation and dehydration increase modestly in response to anoxia. Fold change data were calculated

from the global mean abundance data for each hPTM.

Characterization of histone H1

Six H1 proteins were identified: (**A**) H1-like (XP_013879563.1), (**B**) H1-like

(XP_013855658.1), (**C**) H1-like (XP_013878240.1), (**D**) H1-like (XP_013888293.1), (**E**) H1-

gamma late-like (XP_013861857.1), and (**F**) H1.0-B (XP_013863633.1). Thirteen types of

PTMs were identified across these isoforms (Figs. 5A). Interestingly, H1-like (XP_013879563.1)

appears to carry more modifications compared to the other isoforms even when percent coverage

is taken into account (Figs. 1, 3D, 5C). Nine of the 15 modifications with an average relative

abundance of exactly 100% in all samples are H1 isoforms, with H1-gamma late-like

(XP_013861857.1) having 8 of the 10 (Table 1).

Figure 5. Significant differential histone post-translational modifications on histone H1 isoforms in WS40NE

cells. (**A**) The most commonly significant differential hPTM across H1 isoforms were oxidation (22 hPTMs),

dioxidation (20 hPTMs), and acetald+28/dimethylation (17 hPTMs). (**B**) Venn diagram illustrating the occurrence of

unique and biologically-relevant hPTMs on H1 histone isoforms across experimental treatment groups. (**C**) Five of

the six H1 histone isoforms had residues with significantly different hPTM modifications in response to anoxia. The

percentage above each bar represents the peptide sequence coverage of that isoform. Note that H1-gamma late-like

is modified, but those modifications are stable across experimental treatment groups.

All six H1 isoforms had residues with significant hPTM modifications, while 5 isoforms had at

least one hPTM that changed in relative abundance in response to experimental treatments. One

hundred and forty-three significant hPTMs were present on H1 isoforms in at least one condition

(Fig. 5). The most common differential modifications in response to the experimental treatments

across H1 isoforms were oxidation (22 significant unique hPTMs), dioxidation (20 significant

unique hPTMs), and acetald+28/dimethylation (17 significant unique hPTMs) (Fig. 5). Six

highly condition-dependent hPTMs were significant across five different comparisons: H1-like

(XP_013879563.1) 17K acetylation, 62R oxidation/hydroxylation, and 63S phosphorylation; H1-

like (XP_013855658.1) 51K 4-hydroxynonenal; and H1.0-B (XP_013863633.1) 19K 4-

hydroxynonenal and 25K acetylation (Fig. 6 and Table 2).

A) XP_013879563.1 H1−like

Average Relative Abundance (%)

Peptide sequence:
MSELQSDSLSVLQRQTKKRKMAEVAPAPAPAKAAKKKSTKLKK
VGPSVSELIVKAVAASKERSGVSAAALKKTLAAGGYDVDKNKA
RVKTAIKSLVTKGTLVQTKGIGASGSFKMNKQADTKAKKPAAKK
PAPKAKKPAAKKPAAAKKPKKAAAAAKKPAAAKKSPKKAKKPA
AAAKNAAKSPKKTTKSPKKVVKKAPAAKKSPAKKAAKPKAKKA
APKKK

PTM Description
- 4−hydroxynonenal
- Acetald+28 / Dimethyl
- Acetyl
- ADP−Ribosyl
- Biotin
- Carboxyethyl / Lactyl
- Dehydrated
- Dimethyl
- Dioxidation
- Methyl
- Oxidation/Hydroxyl
- Phospho
- Ubiquityl

B) XP_013855658.1 H1−like

Average Relative Abundance (%)

Peptide sequence:
MAEVAPAPAPAKAAKKKSTKPKKAGPSVSELIVKAVAASK
ERSGVSAAALKKTLAAGGYDVDKNKARVKTAIKSLVTKG
TLVQTKGIGASGSFKMNKQADSKAKNPAAKKAAPKAKK
PAAKKPAAAKKPKKPAAAAKKPAAAKKSPKKAKKPAAAA
KKAAKSPKKTTKSPKKVVKKAPAAKKSPAKKAAKPKAKK
AAPKKK

Figure 6. Characterization of post-translational modifications on Histone H1 isoforms in WS40NE cells exposed to anoxia and aerobic recovery from anoxia. Isoforms represented are (**A**) H1-like (XP_013879563.1), (**B**) H1-like (XP_013855658.1), (**C**) H1-like (XP_013878240.1), (**D**) H1-like (XP_013888293.1), (**E**) H1-gamma late-like (XP_013861857.1), and (**F**) H1.0-B (XP_013863633.1). Each panel represents a different amino acid position where at least one hPTM was detected. Amino acid residues are abbreviated using their one letter code. The x-axis displays the condition of the samples: 0 d in anoxia (0), 1 d in anoxia (1), 4 d in anoxia (4), or 1 d of aerobic recovery from 4 d of anoxia (R). The y-axis scales differ across panels to better visualize changes in low abundance hPTMs. Detected (purple) and modified (blue) peptides and the entire peptide sequences for each isoform are listed beside the panels.

Characterization of histone H2A

Nine H2A proteins were identified: (**A**) H2A (XP_013874651.1), (**B**) H2A-like

(XP_013879560.1), (**C**) H2A-like (XP_013878193.1), (**D**) macro-H2A.2 (XP_013858189.1), (**E**)

core histone macro-H2A.1 (XP_013877703.1), (**F**) H2A.V (XP_013854970.1), (**G**) H2A.Z

(XP_013858796.1), (**H**) H2AX-like (XP_013887205.1), and (**I**) H2AX-like (XP_013862107.1/

XP_013878242.1). Thirteen types of PTMs were identified across these isoforms (Fig. 7A) with

257 hPTMs significantly differentially abundant on H2A isoforms in at least one condition (Fig.

7B). All nine H2A isoforms had residues with significant hPTM modifications, with H2A

(XP_013874651.1) and H2A-like (XP_013879560.1) appearing to have a high proportion of

hPTMs for their percent coverage in the data (Fig. 7C). The top three most common hPTMs that

were significantly different in abundance in response to experimental treatments across H2A

isoforms were dioxidation (45 significant unique hPTMs), oxidation (38 significant unique

hPTMs), and dehydration (25 significant unique hPTMs) (Fig. 7A). Two highly condition-

dependent hPTMs were significantly different across six different comparisons: H2A-like

(XP_013879560.1) 123K carboxyethylation/lactylation (Figs. 7E, 8B) and H2A-like

(XP_013878193.1) 102S dehydration (Figs. 7D, 8C). Two additional highly condition-dependent

hPTMs were significantly different across five different comparisons: H2A.V

(XP_013854970.1)121K biotinylation and 122K dioxidation (Fig. 9A, Table 2).

Figure 7. Significant differential post-translational modifications on histone H2A isoforms in WS40NE cells exposed to anoxia and aerobic recovery from anoxia. (A) The top three most commonly significant differential hPTMs across histone H2A isoforms were dioxidation (45 hPTMs), oxidation (38 hPTMs), and dehydration (25 hPTMs). (**B**) Venn diagram illustrating the occurrence of unique and biologically-relevant hPTMs on H2A histone isoforms across experimental treatment groups. (**C**) At least nine H2A isoforms had residues that were significantly modified. The percentage above each bar represents the peptide sequence coverage of that isoform. Note that two isoforms appear to be highly modified based on their percent sequence coverage. Two specific hPTMs were

262   significant across all six pairwise comparisons: **(D)** H2A-like (XP_013878193.1) 102S dehydration and **(E)** H2A-

263   like (XP_013879560.1) 123K carboxyethylation/lactylation. Bars represent means ± SEM (n=11-12).

D) XP_013858189.1 macro-H2A.2

Peptide sequence:
MSARGGKKKVTKLSRSSRAGVIFPVGRMMRYLRTGTHKY
RIGMGAPVYMAAVIEYLAAEILELAGNAARDNKKGRITPRHI
KLAVANDEELNQLLRGVTISNGGVLPRIHPELLSKKRGSRA
KADTQTSTPEKQEERSKNKKVVKPFKKFKGKRGRKPKSAE
GDKESVPSSSVEDGPGDGFTILSAKSLFLGQKLSLTESEISK
IGSIKVEGIINPTNAELDLKDGVGNALEKAGGREFLDGVKEL
RKAQGPLEVASVAVSQANGMAARFVIHCNVPQWGSEKCE
DQLEKTVKNCLSAAEEKKLKSVAFPSLPAGRNGFPKQTAA
QLILKAVSNHFVSSTSSSLKNIYFVLFDSESIGIYLQEMAKLD
TK

E) XP_013877703.1 macro-H2A.1

Peptide sequence:
MSSRGGKKKSTKTSRSTKAGVIFPVGRMLRYIKRGLPKYRI
GVGAPVYLAAVLEYLTAEILELAGNAARDNKKGRVTPRHILL
AIANDEELNQLLKGVTIAAGGVLPNIHPELLAKKRGAKGKLE
TPVSPAPEKKPKTVKKTATKKMTGKKAGGKAKKQGEVSKA
ASADSTTEGSPADSFTVLSTKSLFLGQKLNLIHSEVSNLAGF
DVDGVINPTNTELELKDDLGSALEKKGGKEFTEALQELKKK
NGPLEVAGAVLTGGFGLPAKYVIHCNSPGWGSDKCEEMLE
KTVKNCLALADEKKLKSVAFPSIGSGRNGFPKQTAAQLILKA
ISSYFVATMSSTIKTVYFVLFDSESIGIYVQEMAKLETS

PTM Description
- 4-hydroxynonenal
- Acetald+28 / Dimethyl
- Acetyl
- ADP-Ribosyl
- Biotin
- Carboxyethyl / Lactyl
- Dehydrated
- Dimethyl
- Dioxidation
- Oxidation/Hydroxyl
- Phospho
- Ubiquityl

Figure 8. Characterization of the post-translational modifications on histone H2A isoforms in WS40NE cells exposed to anoxia and aerobic recovery from anoxia. Isoforms represented are (**A**) H2A (XP_013874651.1), (**B**) H2A-like (XP_013879560.1), (**C**) H2A-like (XP_013878193.1), (**D**) macro-H2A.2 (XP_013858189.1), and (**E**) macro-H2A.1 (XP_0138777031.1). Each panel represents a different amino acid position where at least one hPTM was detected. Amino acid residues are abbreviated using their one letter code. The x-axis displays the condition of the samples: 0 d in anoxia (0), 1 d in anoxia (1), 4 d in anoxia (4), or 1 d of aerobic recovery after 4 days in anoxia

272   (R). The y-axis scales differ across panels to better visualize changes in low abundance hPTMs. Detected (purple)

273   and modified (blue) peptides and the entire peptide sequences are listed beside the panels.

A) XP_013854970.1 H2A.V

Peptide sequence:
MAGGKAGKDSGKAKAKAVSRSQRAGLQFPVGRIHRHL
KTRTTSHGRVGATAAVYSAAILEYLTAEVLELAGNASKDL
KVKRITPRHLQLAIRGDEELDSLIKATIAGGGVIPHIHKSLI
GKKGQQKTA

PTM Description
- 4–hydroxynonenal
- Acetald+28 / Dimethyl
- Acetyl
- ADP–Ribosyl
- Biotin
- Carboxyethyl / Lactyl
- Dehydrated
- Dimethyl
- Dioxidation
- Methyl
- Oxidation/Hydroxyl
- Phospho
- Ubiquityl

B) XP_013858796.1 H2A.Z

Peptide sequence:
MAGGKAGKDSGKTKTKAISRSQRAGLQFPVGRIHRHLK
SRTTSHGRVGATAAVYSAAILEYLTAEVLELAGNASKDL
KVKRITPRHLQLAIRGDEELDSLIKATIAGGGVIPHIHKSLI
GKKGQQKTV

C) XP_013887205.1 H2AX–like

Peptide sequence:
MSGRGKTGGKARAKAKTRSSRAGLQFPVGRVHRLLRKG
NYAEYRVGAGAPVYLAAVLELTAEILELAGNAARDNKKTRI
IPRHLQLAVRNDEELNKLLGGVTIAQGGVLPNIQAVLLPKK
TGQSAPSSGKAGKKAFSQSQEY

D) XP_013862107.1 / XP_013878242.1 H2AX_H2AX

Peptide sequence:
MSGRGKSGAKARAKAKTRSSRAGLQFPVGRVHRLLR
KGNYAERVGAGAPVYLAAVLEYLTAEILELAGNAARDN
KKTRIIPRHLQLAVRNDEELNKLLGGVTIAQGGVLPNIQ
AVLLPKKTGQSAPSSGKAGKKASSQSQEY

Figure 9. Characterization of the post-translational modifications on histone H2A isoforms in WS40NE cells exposed to anoxia and aerobic recovery from anoxia. (**A**) H2A.V (XP_013854970.1), (**B**) H2A.Z (XP_013858796.1), (**C**) H2AX-like (XP_013887205.1), and (**D**) H2AX-like (XP_013862107.1/ XP_013878242.1). Each panel represents a different amino acid position where at least one hPTM was detected. Amino acid residues are abbreviated using their one letter code. The x-axis displays the condition of the samples: 0 d in anoxia (0), 1 d in anoxia (1), 4 d in anoxia (4), or 1 d of aerobic recovery following 4 d of anoxia (R). The y-axis scales differ across panels to better visualize changes in low abundance hPTMs. Detected (purple) and modified (blue) peptides and the entire peptide sequences are listed beside the panels.

Characterization of histone H2B

At least five H2B proteins were identified: (**A**) H2B_1/2-like (XP_013874662.1), (**B**) H2B_1/2 (XP_013879561.1 / XP_013878239.1 / XP_013878163.1 / XP_013869445.1), (**C**) H2B_1/2-like (XP_013857090.1), (**D**) H2B.L4-like (XP_013885354.1), and (**E**) H2B.3-like (XP_013886452.1). Thirteen types of PTMs were identified across these isoforms (Fig. 10A). Three hundred and twenty-seven significant hPTMs were present across H2B isoforms in at least one condition, making this the most significantly modified class of histones in this study (Fig. 10B). All identified H2B isoforms had at least one residue with a significant hPTM, while 4 of the 5 were highly modified (Fig. 10C). The top three most commonly significant differential hPTMs across H2B isoforms were phosphorylation (58 significant unique hPTMs), dehydration (49 significant unique hPTMs), and oxidation/hydroxylation (35 significant unique hPTMs) (Fig. 10A).The following hPTMs displayed an average relative abundance of exactly 100% in all samples: H2B.L4-like (XP_013885354.1) 31Y, 43T, 46S, and 55S dehydration (Fig. 11, Table 1).

Figure 10. Significant differential post-translational modifications on histone H2B isoforms in WS40NE cells exposed to anoxia and aerobic recovery from anoxia. (A) The top three most commonly significant differential

hPTMs across H2B isoforms were phosphorylation (58 hPTMs), dehydration (48 hPTMs), and

oxidation/hydroxylation (35 hPTMs). (**B**) Venn diagram illustrating the occurrence of unique and biologically-

relevant hPTMs on H2B histone isoforms across experimental treatment groups. (**C**) At least five H2B isoforms had

residues with significant hPTMs. The percentage above each bar represents the peptide sequence coverage of that

isoform. Three hPTMs on H2B.L4-like (XP_013885354.1) were highly condition dependent across all six

comparisons: (**D**) 78S ubiquitylation, (**E**) 82S dehydration, and (**F**) 83R oxidation/hydroxylation. Bars represent the

mean ± SEM (n=11-12).

Three hPTMs on H2B.L4-like (XP_013885354.1) were highly condition dependent across all six

comparisons: 78S ubiquitylation, 82S dehydration, and 83R oxidation/hydroxylation (Fig.

10D,E,F). An additional twenty-three hPTMs were highly condition dependent across five

different comparisons on four different histone isoforms: eight on H2B_1/2-like

(XP_013874662.1), seven on H2B_1/2 (XP_013879561.1 / XP_013878239.1 / XP_013878163.1

/ XP_013869445.1), four on H2B_1/2-like (XP_013857090.1), and four on H2B.L4-like

(XP_013885354.1) (Fig. 11, Table 2). Two of these highly condition-dependent modifications on

H2B.L4-like (XP_013885354.1) occur on the same residue: 79T dehydration and

phosphorylation (Fig. 11D, Table 2).

A) XP_013874662.1 H2B_1/2–like

Peptide sequence:
MPEPAKSAPKKGSKKAVTKTAGKGGKKKRKT
RKESYAIYVYKVLKQVHPDTGISSKAMSIMNSF
VNEIFERIASEASRLAHYNKRSTITSREIQTAVR
LLLPGELAKHAVSEGTKAVTKYTSSK

PTM Description
- 4–hydroxynonenal
- Acetald+28 / Dimethyl
- Acetyl
- ADP–Ribosyl
- Biotin
- Carboxyethyl / Lactyl
- Dehydrated
- Dimethyl
- Dioxidation
- Methyl
- Oxidation/Hydroxyl
- Phospho
- Ubiquityl

B) XP_013879561.1 / XP_013878239.1 / XP_013878163.1 / XP_013869445.1 H2B_1/2

Peptide sequence:
MPEPAKSAPKKGSKKAVTKTAGKGGKKKR
KTRKESYAIYVYKVLKQVHPDTGISSKAMSI
MNSFVNDIFERIASEASRLAHYNKRSTITSR
EIQTAVRLLLPGELAKHAVSEGTKAVTKYTS
SK

C) XP_013857090.1 H2B_1/2–like

Peptide sequence:
MPEPAKSAPKKGSKKAVTKTAGKGGKKKRKTKKE
SYAIYVYKVLKQVHPDTTISSKAVSFMNSFVNNIFE
RIASKASRLAHYNKRSTITSREIQTTVRLLLPGELA
KYPNN

PTM Description
- 4–hydroxynonenal
- Acetald+28 / Dimethyl
- Acetyl
- ADP–Ribosyl
- Biotin
- Carboxyethyl / Lactyl
- Dehydrated
- Dimethyl
- Dioxidation
- Methyl
- Oxidation/Hydroxyl
- Phospho
- Ubiquityl

Y-axis: Average Relative Abundance (%)

D) XP_013885354.1 H2B.L4–like

Peptide sequence:
MSNDVSKKKGKSSGDKKAKRKVKRRETYAMYIY
KVLKQVHPDTGISSRAMSIMNSFVNDLFERIATEA
SRLAQYNKRSTITSREVQTAVRLLLPGELAKHAVS
EGTKAVTKYTSSK

Y-axis: Average Relative Abundance (%)

X-axis: Condition

E) XP_013886452.1 H2B.3–like

40R

Y-axis: Average Relative Abundance (%)

Peptide sequence:
MSIMNSFINIFKRIASKASHLAHYCKRSTITSREIQT
DVRLLLPGELAKHTMSKGTKAXTKYTSSX

Figure 11. Characterization of the post-translational modifications on histone H2B isoforms in WS40NE cells

exposed to anoxia and aerobic recovery from anoxia. Isoforms represented are (**A**) H2B_1/2-like

(XP_013874662.1), (**B**) H2B_1/2 (XP_013879561.1/ XP_013878239.1 / XP_013878163.1 / XP_013869445.1), (**C**)

H2B_1/2-like (XP_013857090.1), (**D**) H2B.L4-like (XP_013885354.1) and (**E**) H2B.3-like (XP_013886452.1).

Each panel represents a different amino acid position where at least one hPTM was detected. Amino acid residues

are abbreviated using their one letter code. The x-axis displays the condition of the samples: 0 d in anoxia (0), 1 d in

anoxia (1), 4 d in anoxia (4), or 1 d of aerobic recovery after 4 d of anoxia (R). The y-axis scales differ across panels

to better visualize changes in low abundance hPTMs. Detected (purple) and modified (blue) peptides and the entire

peptide sequences are listed beside the panels.

Characterization of histone H3

At least three H3 proteins were identified: (**A**) H3.3 (XP_013887037.1 / XP_013885059.1 /

XP_013879174.1 / XP_013870443.1), (**B**) H3-like (XP_013879564.1), and (**C**) H3-like

(XP_013878241.1). All identified H3 isoforms had at least one residue with a significant hPTM,

with 12 types of hPTMs identified across these isoforms (Fig. 12A). The most common hPTMs

that differ significantly across experimental treatments on H3 isoforms were

oxidation/hydroxylation (23 significant unique hPTMs) and dioxidation (18 significant unique

hPTMs) (Fig. 12A). Eighty-nine hPTMs were present across all three H3 isoforms in at least one

condition (Fig. 12B). Two hPTMs displayed an average relative abundance of exactly 100% in

all samples: H3-like (XP_013878241.1) 17K and 24R oxidation/hydroxylation (Fig. 13C, Table

1).

Figure 12. Significant differential post-translational modifications on histone H3 isoforms in WS40NE cells

exposed to anoxia and aerobic recovery from anoxia. (A) The most commonly significant differential hPTMs

across H3 isoforms were oxidation (23 hPTMs) and dioxidation (18 hPTMs). (**B**) Venn diagram illustrating the

occurrence of unique and biologically-relevant hPTMs on H3 histone isoforms across experimental treatment

groups. (**C**) All identified H3 isoforms had residues with significant hPTM modifications. The percentage above

each bar represents the peptide sequence coverage of that isoform. Bars represent the mean ± SEM (n=11-12).

Figure 13. Characterization of the post-translational modifications on histone H3 isoforms in WS40NE cells exposed to anoxia and aerobic recovery from anoxia. Isoforms represented are (**A**) H3.3 (XP_013887037.1/

XP_013885059.1/ XP_013879174.1 / XP_013870443.1), (**B**) H3-like (XP_013879564.1), and (**C**) H3-like

(XP_013878241.1). Each panel represents a different amino acid position where at least one hPTM was detected.

Amino acid residues are abbreviated using their one letter code. The x-axis displays the condition of the samples: 0

d in anoxia (0), 1 d in anoxia (1), 4 d in anoxia (4), or 1 d aerobic recovery after 4 d of anoxia (R). The y-axis scales

differ across panels to better visualize changes in low abundance hPTMs. Detected (purple) and modified (blue)

peptides and the entire peptide sequences are listed beside the panels.

 Extracellular Lactate Accumulation

While previous studies have explored accumulation of lactate during exposure to anoxia in

WS40NE cells, this was the first study to monitor how this lactate accumulation persisted during

recovery (Fig. 14). As previously shown, anoxic WS40NE cells accumulate extracellular lactate

at a significantly higher rate than normoxic cells. After 1 d of aerobic recovery, this trend

persists.

Figure 14. Extracellular lactate accumulation in WS40NE cells exposed to anoxia and recovery from anoxia.

(**A**) Total media lactate concentration during exposure to anoxia and recovery from anoxia. Due to the recovery

media change, the data represents only 1 day of lactate accumulation in normoxic recovery conditions. (**B**) The

average rate of lactate accumulation (mM/day) in the same samples. Different letters indicate significant differences

in means within each graph (one-way ANOVA with Tukey's HSD test $p > 0.05$, $n = 6$).

Discussion

Histone isoforms and post-translational modifications change in response to anoxia and aerobic

recovery from anoxia in WS40NE cells. This is consistent with hypoxic responses in hPTMs and

histone isoforms observed in human cells [39, 40]. Significant differences in histone isoform

abundance were detected in response to anoxia. Additionally, of the 1043 unique biologically

relevant hPTMs detected, 816 hPTMs were significantly modified in at least one condition. All

histone groups showed significant post-translational modifications in response to changing

oxygen availability. H2B was the most significantly modified histone group with 327 significant

differential hPTMs present in at least one condition. H2B_1/2 (XP_013879561.1 /

XP_013878239.1 / XP_013878163.1 / XP_013869445.1) was the most significantly modified

histone, with 101 significant modifications. Interestingly, highly condition-dependent hPTMs

often occurred on adjacent residues, with two highly condition-dependent hPTMs sharing a

single residue. This suggests that histones of WS40NE cells have highly modifiable peptide

regions that are responsive to changes in oxygen availability and likely help to coordinate

changes in gene expression associated with their extreme anoxia tolerance.

Changes in histone isoform abundance

Several histone isoforms change in abundance in response to anoxia. While, to our knowledge,

no study has identified anoxia-dependent changes in histone isoform abundance, there are

several studies that document changes in histone isoforms in response to heat stress, hypoxia,

ischemia, and oxidative stress [40-43]. Of note, the increase in histone H3.3 observed in this

study is consistent with a role for "priming" the cells for a robust gene expression response to

stress as has been shown in human lung fibroblasts [40]. Importantly, histone H3.3 expression

responds within 1 day of anoxia and does not immediately return to pre-anoxic levels after 1 day

of recovery, suggesting a prolonged change in poise of the chromatin. H3.3 is typically

associated with euchromatin and active regulatory elements [44], suggesting that the genomic

location of the histone H3.3 isoforms may indicate genes that are immediately responsible for

supporting cellular functions in the absence of oxygen, or may represent genes primed and ready

to be expressed during recovery from anoxia. If future studies can identify the genomic location

of the H3.3 isoforms, these genes would be prime candidates for explaining anoxic-

preconditioning in embryos of *A. limnaeus* [45].

H2A/H2B dimers flank entry and exit portals for DNA in the nucleosome, and thus changes to

these histone isoforms are highly likely to impact gene expression [46, 47]. In WS40NE cells,

one histone H2A isoform decreases within 1 day of anoxia, while two histone H2B isoforms

decrease, but only after 4 days of anoxia, suggesting unique regulation and usage of these

isoforms in response to anoxia. Histone H2A isoforms have been shown to increase after

exposure to hypoxia and ischemia in some mammalian models [48, 49], a result that is opposite

of that in WS40NE cells. The reason for this shift and their potential for altering gene expression

is worthy of future studies.

Global patterns of histone ubiquitylation in WS40NE cells exhibit similar patterns to whole

embryos

The global decrease in histone ubiquitylation during anoxia reported here supports earlier work

on whole embryos of *A. limnaeus* [50]. WS40 embryos (the stage of embryo that WS40NE cells

were isolated from) exhibit a global decrease in ubiquitin conjugates after 3 days of anoxia that

return to pre-anoxic levels after 1 day of aerobic recovery [50]. Interestingly, a ubiquitylated 23

kDa protein, hypothesized to be a histone, in these embryos was absent during anoxia and

present only when oxygen was available. In the residue-specific data presented here, only one

residue fits this pattern: serine 63 on H2B-1/2-like (XP_013874662.1) decreases from 41.3%

ubiquitylated in normoxia to 2.3% after 4 days of anoxia. A similar pattern on the same serine

residue is observed in another H2B-1/2-like isoform (XP_013857090.1), but on this isoform the

residue has a much lower rate of modification and instead is dominated by dehydration. Thus,

antibody- and MS-based data suggest a possible role for this ubiquitylation in managing gene

expression in response to long-term anoxia. However, it should be noted that different stage

embryos illustrated different patterns, with developing embryos that exhibit extreme anoxia

tolerance (WS36 embryos) accumulating ubiquitin conjugates only after 1 d of aerobic recovery

from long-term anoxia [50]. Future work should be done to elucidate the complex nature of the

ubiquitin system in both WS40NE cells and *A. limnaeus* embryos, particularly in understanding

the role of histone ubiquitylation in anoxia.

Global decreases in multiple hPTMs deviate from expected patterns

Global changes to hPTMs have been recorded in mammalian cells in response to hypoxia; large

changes in modifications of H3 and H4 histones are seen in HEK 293T, HeLa and RKO cells

with smaller changes occurring on H2A and H2B histones [39]. However, few studies have

investigated anoxia-induced changes in hPTMs in a modification-specific way [34].

Interestingly, global methylation in WS40NE cells is stable for the first day of anoxia, but then

significantly decreases during 4 d anoxia and remains low for 1 d of aerobic recovery. As

H3K4me3 and H3K36me3 increase within 1 h during hypoxia in mammalian systems, we

expected global methylation in *A. limnaeus* to reflect similar changes [28]. Increases in

methylation in hypoxic mammalian systems is thought to be due to decreased activity of histone

demethylases when oxygen is limiting, mainly because many of these enzymes use molecular

oxygen as a substrate, and at least one histone demethylase appears to be upregulated in a

hypoxia-inducible factor dependent manner, presumably to combat the overall loss of

demethylase activity during hypoxia [51]. Thus, the large and sustained decrease in global

methylation exhibited in killifish WS40NE cells is an unexpected result when referenced to the

mammalian literature. Histone methylation, including di- and trimethylation, is associated with

transcriptional activation and repression in a residue specific manner [52, 53], so it is difficult at

this time to draw any conclusions for the role that decreased methylation might play. Of note,

only one site of methylation in our dataset was highly condition dependent (H2B_1/2,

XP_013874662.1 and XP_013869445.1, 124K). This residue does not correspond with any

known mammalian hPTMs but may serve a similar function and should be investigated further.

Given its role in the regulation of DNA repair [54], it is surprising that global levels of ADP-

ribosylation consistently decrease in anoxia and remain depressed during aerobic recovery. Most

residue-specific patterns match this global pattern with either stable or decreasing levels of ADP-

ribosylation during anoxia. However, in the cases where an increase occurs on a specific residue,

it is almost always during aerobic recovery. This suggests a conserved role for ADP-ribosylation

of histones associated with oxidative stress and the DNA damage response [54, 55]. However,

PARPs including diphtheria-toxin-like ADP-ribose transferases (ARTDs) and cholera-toxin-like

ARTCs can modify a variety of residues: lysine, aspartate, glutamate, serine, arginine, tyrosine,

and histidine [56, 57]. Additionally, ADP-ribosylation can occur on arginine, serine, threonine,

and cysteine residues via the action of sirtuins (SIRT4, SIRT6, and SIRT7) that perform mono

(ADP-ribose) (MAR) addition [56]. Histone ADP-ribosylation is associated with a variety of

functions beyond DNA damage response, including chromatin remodeling and facilitation of

gene transcription by making chromatin more accessible [58, 59]. A decrease in ADP-

ribosylation may be associated with a decrease in chromatin availability, decreasing the

accessibility of the genome to polymerases and transcription factors. Thus, the overall decrease

in histone ADP-ribosylation may support the downregulation of transcription associated with

entrance into anoxia-induced quiescence.

Recent studies suggest lactate accumulation in mammalian cells may play a role in cell signaling

and regulation through histone lactylation [60], including stimulating gene transcription in

macrophages [61]. WS40NE cells accumulate extracellular lactate throughout anoxia and during

at least the first day of aerobic recovery. Lactate is a glycolytic end-product that has classically

been considered an unavoidable consequence of vertebrate anaerobic metabolism, but in

WS40NE cells, 49 unique carboxyethylation/lactylation modifications significantly changed in

abundance in at least one comparison and one of these modifications was classified as highly

condition dependent (Table 2). Surprisingly, increased lactate accumulation in WS40NE cells is

correlated with *decreased*, not increased, global carboxyethylation/lactylation. This pattern is the

opposite of what is seen in human MCF-7, HeLa, and MDA-MB-231 cells, which demonstrate a

dose-dependent increase in histone lactylation in response to 25 mM exogenous L-lactate [61].

On a residue-specific level there are 70 residues with carboxyethylation/lactylation present, and

one of these residues is highly condition dependent. Given the extreme rarity of protein

carboxyethylation, these are most likely sites of lactylation [62]. In almost all cases, residues in

WS40NE histones experience a decrease in lactylation during anoxia. It is possible that the

slower rate of lactate accumulation in WS40NE cells (7.8 mM extracellular lactate after 4 days

of anoxia) does not produce enough lactate to cause increase lactylation. However, lactate levels

are relatively low during normoxia, and yet many histone residues still have a high percentage of

lactylation. Thus, it would seem that lactate availability is not likely to limit histone lactylation in

WS40NE cells. It is worth considering that decreased rates of lactate accumulation and decreased

histone lactylation in WS40NE cells may at least partially underlie their extreme anoxia

tolerance, as this is contrary to trends that appear in anoxia intolerant mammalian cells.

Several hPTMs suggest protection against ROS damage

Stress related hPTMs, including oxidation/hydroxylation and dioxidation, increase globally in

response to anoxic stress and aerobic recovery in WS40NE cells. This increase is expected, as

these hPTMs are known to increase during stress due to increased rates of reactive oxygen

species (ROS) production [63, 64]. While oxidation of proteins can be the result of nonenzymatic

ROS damage, several enzymes are involved in histone hydroxylation and oxidation [65-67].

Jumonji domain containing 6 (JMJD6) protein is known to hydroxylate histone lysine residues

while the lysyl oxidase (LOX) family of proteins are able to oxidize histone lysines [65, 66].

These enzymes may be interesting targets of future work in the WS40NE cell line, because a

number of specific residues defy the global pattern and decrease in response to anoxia, while

others remain stable, suggesting an enzymatic mechanism and not oxidative damage. Given the

role of prolyl-hydroxylases in regulating oxygen-dependent gene expression, investigating

histones that decrease in oxidation in response to anoxia (e.g., H3-like, XP_013879564.1, 196R)

may hold promise for finding histone oxygen sensors that could participate in regulation of gene

expression at the local level.

Unexpectedly, another stress related hPTM, 4-hydroxynonenal (4-HNE), remains stable for the

first day of anoxia and then significantly decreases after 4 days of anoxia. This decrease

continues during the initial 1 day of aerobic recovery. 4-hydroxynonenal is a product of lipid

peroxidation, and therefore its presence would be expected to increase with ROS stress [68], as

would be expected during both the transition into and out of anoxia. Fish, however, are known to

have high levels of antioxidants that mediate ROS stress and allow them to maintain low levels

of lipid peroxidation [69, 70]. In addition, embryos of *A. limnaeus* have an extremely high level

of antioxidant capacity [71]. Thus, it is possible that WS40NE cells have a mechanism to prevent

lipid peroxidation during anoxia.

Dehydration and phosphorylation – a reciprocal relationship?

Dehydration, especially of serine, threonine, and tyrosine, increases globally in response to

anoxic stress and aerobic recovery in WS40NE cells. In addition, some histone isoforms contain

stable and apparently permanent dehydrations, suggesting functional importance and a likely

enzymatic synthesis of this PTM. Amino acid dehydration is an elimination reaction that results

in a dehydroamino acid through the removal of a hydroxyl group [72] or by the direct removal of

a phosphate group [73]. On residues S and T, it is specifically a beta-elimination that results in

production of dehydroalanine [74]. Dehydroalanine, though common in bacterial and fungal

models, is considered rare in mammals [75, 76]. However, to our knowledge, dehydroamino

acids, including dehydroalanines, have not been investigated in context of fish models.

Additionally, dehydration resulting in dehydroalanines has not been well described as a hPTM,

although it has been described as an intermediary PTM in the maturation of lanthipeptide

antibiotics [72], and as a PTM in microbially-catalyzed modification of eukaryotic kinases [73].

In the case of kinase modification, microbial enzymes, such as phosphothreonine lysases, can

target specific MAP kinases and dehydrate residues that would normally be phosphorylated,

leading to inactivation and signaling dysregulation [73]. An enzyme responsible for dehydration

of these amino acids (S and T) would be needed to support this potential mechanism in WS40NE

cells, but it remains unknown. While no specific candidate enzyme has been identified, it is

interesting that eukaryotic genomes contain LanC-like proteins that share sequence and structural

similarity to bacterial proteins involved in the synthesis of lanthipeptides [73].

Dehydration of residues in WS40NE cell histones increases globally, while conversely, global

phosphorylation decreases during anoxia and aerobic recovery from anoxia. These patterns are

surprising, and the apparent reciprocal patterns of abundance are potentially very interesting.

Stress-induced phosphorylation of specific H3 residues is described in mammalian cells (10S and

11T), although these specific residues were not captured in our dataset [77]. Nevertheless, we

hypothesize that histone dehydration may be a protective feature that prevents inappropriate

phosphorylation of specific residues. Competition between dehydration and phosphorylation

appears to be particularly present on two canonical H1 isoforms (XP_013879563.1 and

XP_013855658.1) in our dataset. Phosphorylation of H1 in response to anoxia is understudied;

however, global dephosphorylation of mammalian H1 is documented to occur after induction of

apoptosis [78]. In WS40NE cells, H1 residues show various patterns of modifications. For

example, the relative abundance of XP_013855658.1 59Y phosphorylation remains stable in

anoxia but the relative abundance of dehydration increases. Conversely, XP_013855658.1 46S

phosphorylation increases in anoxia while dehydration decreases. These unique response patterns

at different residues suggest a residue-specific mechanism may lead to changes in

phosphorylation and dehydration at specific H1 sites. Other residues not captured by our

methodology should also be investigated for dehydration and phosphorylation competition. For

example, in mammalian cells, phosphorylation of H2B 14S is associated with apoptosis and is a

documented response to hypoxia [77, 79]. This residue best corresponds to 12S (13S if the first

methionine is counted as residue 1) in *A. limnaeus* [34], and was unfortunately not captured in

the dataset. Dehydration of this residue could be a possible mechanism for preventing apoptosis

and deserves future investigation.

One important site of phosphorylation was detected in WS40NE histones: H2A.X 137S.

Phosphorylation of mammalian 139S (γ–H2A.X) initiates the DNA repair response and would

therefore be expected to increase during anoxia due to an increase in ROS [80]. Surprisingly,

137S, the equivalent residue in WS40NE histone H2A.X, experiences a decrease in

phosphorylation by 1 day in anoxia and continues to decrease during 4 days of anoxia.

Interestingly, a nearby dehydrated residue, 142Y, follows the opposite pattern, increasing during

anoxia. In this dataset, phosphorylation never occurs at this tyrosine residue. This is unexpected,

as constitutive phosphorylation of mammalian 142Y by Williams–Beuren syndrome

transcription factor (WSTF) occurs during normal conditions while dephosphorylation of this

site occurs during the DNA damage response [81]. This dephosphorylation has been theorized to

enhance recruitment of protein kinases to better maintain γ–H2A.X phosphorylation [81].

Therefore, it is interesting that this site is dehydrated, as these dehydrations may prevent

phosphorylation and thus prevent the dampening of the DNA damage response that is observed

in mammalian cells [81]. These sites should be explored in the future, as they are implicated in

regulation of transcription and may have nearby sites of dehydration that are involved in their

modification status [82].

Multiple hPTMs do not return to baseline after 1 day aerobic recovery

The histone landscape of WS40NE cells remains altered after 1 day of aerobic recovery from 4

days of anoxia. Which is still considered a relatively short duration of anoxia given that the cells

can survive for 49 days without oxygen [37]. Globally, 8 of the 13 types of hPTMs detected are

significantly different in recovery compared to normoxia. Based on the $\log_2$ fold-change, global

acetylation, oxidation/hydroxylation, and dioxidation are significantly elevated during recovery

while global phosphorylation, biotinylation, 4-hydroxynonenal, methylation, and

carboxyethylation/lactylation are all decreased. Of the unique hPTMs that were significantly

modified, 42% are associated with recovery. While many of the hPTMs may simply not have

returned to baseline, other modifications may be involved in recovery mechanisms. Further work

will need to be done to determine if these hPTMs return to normoxic levels, or if they represent

potential heritable epigenetic alterations to the histone landscape and act as a cellular memory of

past exposures to anoxia.

Conclusion

WS40NE cells are unique in their ability to survive long bouts of anoxia and demonstrate

changes in their histone landscape in response to different lengths of time in anoxia and recovery

from anoxia. This response includes altering both histone isoform relative abundance and hPTM

relative abundance at multiple amino acid positions. This response likely contributes to the

anoxia tolerance displayed in this cell type. Downstream effects of specific hPTM changes

should be studied in the future to determine how specific hPTMs can impact gene expression and

anoxia tolerance. Of specific interest is the potential reciprocal relationship between dehydration

and phosphorylation as a regulatory mechanism to alter gene expression in response to anoxia.

Methods

WS40NE Cell Maintenance and Anoxic Exposure

Cells were grown in cell culture media (L-15 + 8.5% FBS, 5 mM glucose, 100 U/mL penicillin/

streptomycin) in 100×20 mm tissue culture dishes at 30°C and split when confluent using

TrypLE Express [37]. For all anoxic experiments, plates of cells were seeded at 2.5×10^6 cells

per plate and given 24 h to adhere. Cells were then introduced into an anoxic chamber (Bactron

EZ, Sheldon Manufacturing, Cornelius, OR, USA) that contains an atmosphere of 95% N_2 and

5% H_2. Media was immediately replaced with anoxic media that had been purged with N_2 gas for

30 min and then pre-equilibrated to the atmosphere of the anoxic chamber overnight. Cells were

harvested at **A)** 0 h of anoxia (normoxic), **B)** 1 d of anoxia, **C)** 4 d of anoxia, and **D)** after 1 d of

aerobic recovery following 4 d of anoxia (n = 12). To harvest, media was removed and a cell

scraper was used to gently dissociate cells from the bottom of the plate. Two 100×20 mm tissue

culture dishes were pooled for each replicate. Cell pellets were generated by subjecting the

isolated cells to centrifugation at 500 x g for 7 min at 4° C, and removing the supernatant.

Additionally, media samples were taken directly prior to the harvest of all four timepoints and

used to determine the levels of extracellular lactate accumulation. Cell pellets and spent media

were flash frozen in liquid nitrogen. Samples were stored at -80° C until use for histone acid

extraction or determination of lactate concentration.

Histone Acid Extraction and Protein Digestion

A middle-down approach to proteomics was used for liquid chromatography mass spectrometry

(LCMS) using previously described methods [34, 83]. Histone acid extracts were generated by

resuspending cell pellets in 0.4 N H_2SO_4 and incubating the samples with constant rotation at 4

°C for 4 h. Samples were then centrifuged at 16,000 x g for 10 min at 4 °C and the supernatant

was transferred into a fresh 1.5 mL tube. This process was repeated twice. Trichloroacetic acid

(TCA) was added to each sample drop-by-drop until a volume equal to 25% of the estimated

sample volume had been added. Samples were inverted to mix and incubated overnight at 4 °C.

Once the histone proteins had precipitated, samples were centrifuged at 3,400 x g for 5 min at 4

°C, and the supernatant was discarded. The remaining pellet was rinsed twice: first in ice cold

acetone with 0.1% hydrochloric acid and then with 100% acetone. For both rinses, the samples

were centrifuged at 3,400 x g for 2 min at 4 °C and the supernatant was removed. After both

rinses were complete, histone pellets were air dried in a fume hood for 5 min. Samples were

stored at -80 °C until the in-solution digestion.

To begin the digestion, each sample was brought to room temperature (RT), and the histone

pellet was resuspended in 138 μL of 8 M urea. Dithiothreitol (DTT) was added to a final

concentration of 10 mM. Each sample was then briefly vortexed, centrifuged, and incubated at

37 °C for 30 min. Iodoacetamide (IAA) was added to each sample to a final concentration of 30

mM, and samples were briefly vortexed and centrifuged before a 30 min incubation in the dark at

RT. Samples were briefly vortexed and centrifuged again after the incubation and protein

concentration was determined through a bicinchoninic acid (BCA) assay, (cat# 23250, Thermo

Scientific, Waltham, MA, USA). Each sample (50 μg of total protein) was diluted 5:6 (v:v) in

0.1M sodium phosphate buffer (pH 7.8). V8 protease beads (cat#20151, Thermo Scientific,

Waltham, MA, USA) were added at 1:50 ratio (protease:total protein). In sodium phosphate

buffer, V8 protease cleaves proteins at the carboxyl end of both glutamate and aspartate residues.

After a 20 h incubation with constant rotation at 30 °C, protease beads were removed by

centrifugation at 500 x g for 2 min and the supernatants were retained. Supernatants were

subjected to an additional centrifugation at 19,000 x g for 5 min, and those subsequent

supernatants were transferred to fresh tubes and dried using a SpeedVac (Thermo Savant,

model# ISS110, Thermo Scientific, Waltham, MA, USA). The peptide pellets were reconstituted

to a final concentration of 333 ng/μL in 0.1% formic acid diluted in LCMS-grade water. For

peptide clean-up, Pierce C18 Spin Columns (cat# 89873, Thermo Scientific, Waltham, MA,

USA) were used according to the manufacturer's instructions.

Data-dependent acquisition (DDA) was used to generate peak lists [84], and PEAKS Suite X

Plus and MSFragger to annotate histone peptides using the *A. limnaeus* reference proteome [85].

Data-independent acquisition (DIA) was then used to quantify the peptides in the samples.

Lactate Accumulation Assay

Lactate concentration in the cell culture media was measured enzymatically as previously

described [37]. Briefly, each well of a 96-well plate (black walls, clear bottom, Greiner Bio-One,

Monroe, NC, USA) contained the following enzyme assay cocktail: 218 μL of Glycine (0.6 M)-

Hydrazine (0.5 M) buffer (pH = 9.2), 25 μL 27 mM NAD^+ , 5 μL of lactate standard or sample,

and 2.5 μL lactate dehydrogenase (LDH, 14 U/mL, Sigma Aldrich). Change in absorbance was

measured at 340 nm every min for 60 min using a plate reader (Tecan Infinity M200Pro,

Männendorf, Switzerland). Final concentration of lactate in the samples was determined by

interpolation using a lactate standard curve.

Histone and hPTM Library Generation

Peptides modified with hPTMs were distinguished from unmodified sequences based on their

mass shift (Fig. S1). A list of histone PTMs was compiled according to the histone protein name,

amino acid residue/position, and modification type using Python code as previously described

[34]. All histone PTMs detected were evaluated for their biological relevance (classified as post-

translational, other, or multiple) according to their Unimod accession and amino acid residue.

Modifications classified by Unimod as known artefacts or chemical derivatives were removed

from analysis. Modifications with the same mass shift are listed together. Peptides that mapped

to multiple histones in the *A. limnaeus* genome are presented as a protein family and are reported

together.

Figure S1: Quantification of histone post-translational modifications in Skyline. A representative MS spectrum

(**A**) and peak area (**B**) of the most abundant ions for a modifed peptide, E.GTK[+42]AVTKYTSSK, and (**C,D**) the

corresponding unmodified peptide E.GTKAVTKYTSSK. A mass shift of +42 on a peptide represents acetylation.

Protein Abundance and hPTM Statistical Analyses

Histone isoform abundance was evaluated using quantile normalized raw protein abundance data

in Python as previously described [34]. Principal component analysis (PCA) and two-tailed t-

tests with the Benjamini-Hochberg correction applied to account for multiple hypothesis testing

[86] were used to make inferences about histone isoform abundance.

Peptide abundance, as determined by normalized area under the curve, and raw protein

abundance were obtained using Skyline (Version 22.2) [87]. Relative abundance values were

used to calculate beta-values and M-values. Beta-value is a transformation of relative abundance

expressed as a number between 0 and 1 calculated by adding a value of 100 to the denominator

to standardize low intensity peptides [88]. The M-value is a logit transformation of the beta-

value [89]. Additionally, the relative abundance of each hPTM was used to calculate the $\log_2$

fold change, and a two-tailed t-test was used to compare the relative abundance of hPTMs in all

conditions with the Benjamini-Hochberg correction applied to account for multiple hypothesis

testing [86]. This relative abundance analysis was repeated to analyze global hPTM levels and

calculate the $\log_2$ fold change. Additionally, an ANOVA was performed on the global relative

abundance of each hPTM using Prism (ver 10.0.0, GraphPad, Software, Boston, MA, USA). A

Dunnett's multiple comparison test was used to determine significance of all conditions

compared to normoxia. The global relative coverage of each modifiable amino acid residue was

calculated to determine what percentage of each hPTM was present at each modifiable residue.

Histone PTM maps were constructed for anoxic and normoxic samples using the mean relative

abundance of hPTMs. The M-value was used to perform a principal component analysis (PCA).

All abundance plots and PCA plots were made in R using the package ggplot2 [90]. Global

hPTM abundance, protein relative abundance, and global relative residue coverage were

visualized using Prism (ver 10.0.0, GraphPad, Software, Boston, MA, USA). All Python and R

code is accessible by the public on GitHub at

https://github.com/hughcj11/WS40NE_Histone_PTM_Analysis. Supplementary data including

average relative abundance, t-statistic, p-value, and corrected p-value is provided for each

unique, biologically relevant hPTM are provided in Supplementary Table S1 (Excel workbook)

across all aerobic conditions.

Author Contributions

706   JEP and CH conceived and designed the study. CH and EAM performed the experiments and

707   acquired the data. CH analyzed the data. DK and JEP contributed reagents/materials/analysis

708   tools. CH and JEP wrote the paper. All authors edited and approved the manuscript.

Acknowledgements

711   We would like to thank Karim Soliman for his guidance in the creation of the Python data

712   analysis pipeline and visualization of the data. We would like to thank Dr. Daniel Zajic for his

713   help performing the lactate accumulation assay.

Funding

716   Funded by National Science Foundation Grant IOS-2209383 and IOS-2025832.
717

Data and resource availability

The datasets (DDA and DIA raw data) supporting the conclusions of this article are available in

Panorama Public (https://panoramaweb.org/CH0001JP.url) and ProteomeXchange (PXD067707:

https://proteomecentral.proteomexchange.org/cgi/GetDataset?ID=PXD067707).

Reviewer login:

Email: panorama+reviewer361@proteinms.net

Password: 3dqk$TMm1AJbo%

References

1.   Lipton P. Ischemic cell death in brain neurons. Physiological Reviews. 1999;79(4):1431-
568.
2.   Katchanov J, Harms C, Gertz K, Hauck L, Waeber C, Hirt L, et al. Mild cerebral
ischemia induces loss of cyclin-dependent kinase inhibitors and activation of cell cycle
machinery before delayed neuronal cell death. The Journal of Neuroscience. 2001;21(14):5045-
53.
3.   Andersson BS, Aw TY, Jones DP. Mitochondrial transmembrane potential and pH
gradient during anoxia. American Journal of Physiology - Cell Physiology. 1987;252(4):C349-
C55.
4.   Pamenter ME, Gomez CR, Richards JG, Milsom WK. Mitochondrial responses to
prolonged anoxia in brain of red-eared slider turtles. Biology Letters. 2016;12(1):20150797. doi:
10.1098/rsbl.2015.0797. PubMed PMID: 26763217; PubMed Central PMCID:
PMCPMC4785919.
5.   Galli GLJ, Richards JG. Mitochondria from anoxia-tolerant animals reveals commons
strategies to survive without oxygen. Journal of Comparative Physiology B. 2014;184:285-302.
doi: 10.1007/s00360-014-0806-3.
6.   Hammerman C, Kaplan M. Ischemia and reperfusion injury. The ultimate
pathophysiologic paradox. Clinics in Perinatology. 1998;25(3):757-77. doi: 10.1016/S0095-
5108(18)30110-6. PubMed PMID: 9779346.
7.   Salas-Vidal E, Méndez-Cruz FJ, Ramírez-Corona A, Reza-Medina B. Oxygen, reactive
oxygen species and developmental redox networks: Evo-Devo Evil-Devils? The International
Journal of Developmental Biology. 2021;65(4-5-6):345-56. doi: 10.1387/ijdb.200170es.
8.   Scherz-Shouval R, Elazar Z. ROS, mitochondria and the regulation of autophagy. Trends
in Cell Biology. 2007;17(9):422-7.
9.   Kirino T. Delayed neuronal death in the gerbil hippocampus following ischemia. Brain
Research. 1982;239(1):57-69.
10.   Cao W, Carney J, Duchon A, Floyd R, Chevion M. Oxygen free radical involvement in
ischemia and reperfusion injury to brain. Neuroscience Letters. 1988;88(2):233-8.
11.   Feder ME, Hofmann GE. Heat-shock proteins, molecular chaperones, and the stress
response: evolutionary and ecological physiology. Annual Review of Physiology.
1999;61(1):243-82. PubMed Central PMCID: PMCHeat Shock Proteins.
12.   Crowe KM, Newton JC, Kaltenboeck B, Johnson C. Oxidative stress responses of gulf
killifish exposed to hydrocarbons from the Deepwater Horizon oil spill: Potential implications
for aquatic food resources. Environmental Toxicology and Chemistry. 2014;33(2):370-4.
13.   Kültz D. Plasticity and stressor specificity of osmotic and heat shock responses of
Gillichthys mirabilis gill cells. American Journal of Physiology. 1996;271(4):C1181-C93.
PubMed Central PMCID: PMCOsmoregulation.
14.   Carrasco-Malio A, Díaz M, Mella M, Montoya M, Miranda A, Landaeta M, et al. Are the
intertidal fish highly resistant to UV-B radiation? A study based on oxidative stress in *Girella
laevifrons* (Kyphosidae). Ecotoxicology and environmental safety. 2014;100:93-8.
15.   Baker DW, Matey V, Huynh KT, Wilson JM, Morgan JD, Brauner CJ. Complete
intracellular pH protection during extracellular pH depression is associated with hypercarbia

tolerance in white sturgeon, *Acipenser transmontanus*. American Journal of Physiology -
Regulatory, Integrative and Comparative Physiology. 2009;296(6):R1868-R80. doi:
10.1152/ajpregu.90767.2008.
16.   Storey KB. Reptile freeze tolerance: Metabolism and gene expression. Cryobiology.
2006;52(1):1-16.
17.   Churchill TA, Storey KB. Dehydration tolerance in wood frogs: a new perspective on
development of amphibian freeze tolerance. American Journal of Physiology.
1993;265(34):R1324-R32. PubMed Central PMCID: PMCDehydration.
18.   Belkin DA. Anoxia: Tolerance in Reptiles. Science. 1963;139(3554):492-3. doi:
10.1126/science.139.3554.492.
19.   Bickler PE. Cerebral anoxia tolerance in turtles: regulation of intracellular calcium and
pH. American Journal of Physiology. 1992;263(6):R1298-R302. PubMed Central PMCID:
PMCAnoxia.
20.   Mojica EA, Fu Y, Kültz D. Salinity-responsive histone PTMs identified in the gills and
gonads of Mozambique tilapia (*Oreochromis mossambicus*). BMC Genomics. 2024;25(1):586.
Epub 20240611. doi: 10.1186/s12864-024-10471-3. PubMed PMID: 38862901; PubMed Central
PMCID: PMCPMC11167857.
21.   Dyson MH, Thomson S, Mahadevan LC. Heat shock, histone H3 phosphorylation and the
cell cycle. Cell Cycle. 2005;4(1):e64-e-8. PubMed Central PMCID: PMCHeat Shock.
22.   Kim JM, To TK, Ishida J, Morosawa T, Kawashima M, Matsui A, et al. Alterations of
lysine modifications on the histone H3 N-tail under drought stress conditions in Arabidopsis
thaliana. Plant & Cell Physiology. 2008;49(10):1580-8. Epub 20080908. doi:
10.1093/pcp/pcn133. PubMed PMID: 18779215.
23.   Ma T, Keller JA, Yu X. RNF8-dependent histone ubiquitination during DNA damage
response and spermatogenesis. Acta Biochimica et Biophysica Sinica (Shanghai).
2011;43(5):339-45. Epub 20110328. doi: 10.1093/abbs/gmr016. PubMed PMID: 21444325;
PubMed Central PMCID: PMCPMC3080603.
24.   Wang Y, Lyu Y, Tu K, Xu Q, Yang Y, Salman S, et al. Histone citrullination by PADI4
is required for HIF-dependent transcriptional responses to hypoxia and tumor vascularization.
Science Advances. 2021;7(35):eabe3771.
25.   Wrann S, Kaufmann MR, Wirthner R, Stiehl DP, Wenger RH. HIF mediated and DNA
damage independent histone H2AX phosphorylation in chronic hypoxia. Biological Chemistry.
2013;394(4):519-28. doi: 10.1515/hsz-2012-0311.
26.   Batie M, Fasanya T, Kenneth NS, Rocha S. Oxygen-regulated post-translation
modifications as master signalling pathway in cells. EMBO reports. 2023;24(12). doi:
10.15252/embr.202357849.
27.   Das T, Kamle A, Kumar A, Chakravarty S. Hypoxia induced sex-difference in zebrafish
brain proteome profile reveals the crucial role of H3K9me3 in recovery from acute hypoxia.
Frontiers in Genetics. 2021;12:635904. Epub 20220131. doi: 10.3389/fgene.2021.635904.
PubMed PMID: 35173759; PubMed Central PMCID: PMCPMC8841817.
28.   Batie M, Frost J, Frost M, Wilson JW, Schofield P, Rocha S. Hypoxia induces rapid
changes to histone methylation and reprograms chromatin. Science. 2019;363(6432):1222-6. doi:
10.1126/science.aau5870.
29.   Mucci S, Isaja L, Rodriguez-Varela MS, Ferriol-Laffouillere SL, Marazita M, Videla-
Richardson GA, et al. Acute severe hypoxia induces apoptosis of human pluripotent stem cells
by a HIF-1alpha and P53 independent mechanism. Scientific Reports. 2022;12(1):18803. Epub

20221105. doi: 10.1038/s41598-022-23650-7. PubMed PMID: 36335243; PubMed Central
PMCID: PMCPMC9637190.
30.   Wijenayake S, Storey KB. Dynamic regulation of histone H3 lysine (K) acetylation and
deacetylation during prolonged oxygen deprivation in a champion anaerobe. Molecular and
Cellular Biochemistry. 2020;474(1-2):229-41. doi: 10.1007/s11010-020-03848-x.
31.   Anchordoguy TJ, Hand SC. Reactivation of ubiquitination in *Artemia franciscana*
embryos during recovery from anoxia-induced quiescence. Journal of Experimental Biology.
1995;198(6):1299-305. PubMed Central PMCID: PMCAnoxia.
32.   Liu CL, Martone ME, Hu BR. Protein ubiquitination in postsynaptic densities after
transient cerebral ischemia. Journal of Cerebral Blood Flow & Metabolism. 2004;24(11):1219-
25.
33.   Podrabsky J, Wilson N. Hypoxia and anoxia tolerance in the annual killifish
Austrofundulus limnaeus. Integrative and Comparative Biology. 2016;56(4):500-9. doi:
10.1093/icb/icw092.
34.   Hughes C, Mojica EA, Kültz D, Podrabsky JE. Global maintenance of histone post-
translational modifications during the transition into anoxia in embryos of the annual killifish
Austrofundulus limnaeus. Environmental Epigenetics. 2025;in press:1-32.
35.   Van HT, Santos MA. Histone modifications and the DNA double-strand break response.
Cell Cycle. 2018;17(21-22):2399-410. Epub 20181114. doi: 10.1080/15384101.2018.1542899.
PubMed PMID: 30394812; PubMed Central PMCID: PMCPMC6342081.
36.   Weiner A, Hsieh TH, Appleboim A, Chen HV, Rahat A, Amit I, et al. High-resolution
chromatin dynamics during a yeast stress response. Molecular Cell. 2015;58(2):371-86. Epub
20150319. doi: 10.1016/j.molcel.2015.02.002. PubMed PMID: 25801168; PubMed Central
PMCID: PMCPMC4405355.
37.   Riggs C, Le R, Kültz D, Zajic D, Summers A, Alvarez L, et al. Establishment and
characterization of an anoxia-tolerant cell line, PSU-AL-WS40NE, derived from an embryo of
the annual killifish *Austrofundulus limnaeus*. Comparative Biochemistry and Physiology - Part
B. 2019;232:11-22. doi: 10.1016/j.cbpb.2019.02.008.
38.   Fordel E, Thijs L, Martinet W, Schrijvers D, Moens L, DeWilde S. Anoxia or oxygen and
glucose deprivation in SH-SY5Y cells: A step closer to the unraveling of neuroglobin and
cytoglobin functions. Gene. 2007;398:114-22.
39.   Hsu KF, Wilkins SE, Hopkinson RJ, Sekirnik R, Flashman E, Kawamura A, et al.
Hypoxia and hypoxia mimetics differentially modulate histone post-translational modifications.
Epigenetics. 2021;16(1):14-27. Epub 20200701. doi: 10.1080/15592294.2020.1786305. PubMed
PMID: 32609604; PubMed Central PMCID: PMCPMC7889154.
40.   Nagagaki Y, Kozakura Y, Mahandaran T, Fumoto Y, Nakato R, Shirahige K, et al.
Histone H3.3 chaperone HIRA renders stress-responsive genes poised for prospective lethal
stresses in acquired tolerance. Genes to Cells. 2024;29(9):722-34. doi: 10.1111/gtc.13140.
41.   Stowell C, Wang L, Arbogast B, Lan J-q, Cioffi GA, Burgoyne CF, et al. Retinal
proteomic changes under different ischemic conditions-implication of an epigenetic regulatory
mechanism. International Journal of Physiology, Pathophysiology and Pharmacology.
2010;2(2):148.
42.   Zager RA, Johnson AC. Renal ischemia-reperfusion injury upregulates histone-
modifying enzyme systems and alters histone expression at proinflammatory/profibrotic genes.
American Journal of Physiology-Renal Physiology. 2009;296(5):F1032-F41.

43.   Zhou A, Simon RP, David L. Nascent proteomes of ischemic-injured and ischemic-
tolerant neuronal cells. International Journal of Computational Biology and Drug Design.
2011;4(1):40. doi: 10.1504/ijcbdd.2011.038656.
44.   Martire S, Banaszynski LA. The roles of histone variants in fine-tuning chromatin
organization and function. Nature Reviews Molecular Cell Biology. 2020;21(9):522-41. doi:
10.1038/s41580-020-0262-8.
45.   Podrabsky JE, Riggs CL, Duerr JM. Anoxia Tolerance During Vertebrate Development -
Insights from Studies on the Annual Killifish *Austrofundulus limnaeus*. In: Padilla P, editor.
Anoxia. Rijeka, Croatia: InTech; 2012. p. 3-24.
46.   Talbert PB, Henikoff S. Histone variants at a glance. Journal of Cell Science.
2021;134(6):jcs244749. doi: 10.1242/jcs.244749.
47.   Singh R, Bassett E, Chakravarti A, Parthun MR. Replication-dependent histone isoforms:
a new source of complexity in chromatin structure and function. Nucleic Acids Research.
2018;46(17):8665-78. doi: 10.1093/nar/gky768. PubMed PMID: 30165676; PubMed Central
PMCID: PMCPMC6158624.
48.   Boidot R, Branders S, Helleputte T, Rubio LI, Dupont P, Feron O. A generic cycling
hypoxia-derived prognostic gene signature: application to breast cancer profiling. Oncotarget.
2014;5(16):6947-63. doi: 10.18632/oncotarget.2285.
49.   Shah S, Verma T, Rashid M, Gadewal N, Gupta S. Histone H2A isoforms: Potential
implications in epigenome plasticity and diseases in eukaryotes. Journal of Biosciences.
2020;45(1). doi: 10.1007/s12038-019-9985-0.
50.   Meller CL, Meller R, Simons RP, Podrabsky JE. Patterns of ubiquitylation and
SUMOylation associated with exposure to anoxia in embryos of the annual killifish
Austrofundulus limnaeus. Journal of Comparative Physiology B. 2014;184:235-47. doi:
10.1007/s00360-013-0791-y.
51.   Wellmann S, Bettkober M, Zelmer A, Seeger K, Faigle M, Eltzschig HK, et al. Hypoxia
upregulates the histone demethylase JMJD1A via HIF-1. Biochemical and Biophysical Research
Communications. 2008;372(4):892-7. doi: https://doi.org/10.1016/j.bbrc.2008.05.150.
52.   Fabrizio P, Garvis S, Palladino F. Histone methylation and memory of environmental
stress. Cells. 2019;8(4):339. doi: 10.3390/cells8040339.
53.   Kim J, Lee H, Yi SJ, Kim K. Gene regulation by histone-modifying enzymes under
hypoxic conditions: a focus on histone methylation and acetylation. Experimental & Molecular
Medicine. 2022;54(7):878-89. Epub 20220722. doi: 10.1038/s12276-022-00812-1. PubMed
PMID: 35869366; PubMed Central PMCID: PMCPMC9355978.
54.   Brustel J, Muramoto T, Fumimoto K, Ellins J, Pears CJ, Lakin ND. Linking DNA repair
and cell cycle progression through serine ADP-ribosylation of histones. Nature Communications.
2022;13(1). doi: 10.1038/s41467-021-27867-4.
55.   Marnett LJ. Oxyradicals and DNA damage. Carcinogenesis. 2000;21(3):361-70. doi:
10.1093/carcin/21.3.361.
56.   Poltronieri P, Miwa M, Masutani M. ADP-ribosylation as post-translational modification
of proteins: Use of inhibitors in cancer control. International Journal of Molecular Sciences.
2021;22(19):10829. doi: 10.3390/ijms221910829.
57.   Bartlett E, Bonfiglio JJ, Prokhorova E, Colby T, Zobel F, Ahel I, et al. Interplay of
histone marks with serine ADP-ribosylation. Cell Reports. 2018;24(13):3488-502.e5. doi:
10.1016/j.celrep.2018.08.092.

58.   Mohapatra J, Tashiro K, Beckner RL, Sierra J, Kilgore JA, Williams NS, et al. Serine
ADP-ribosylation marks nucleosomes for ALC1-dependent chromatin remodeling. Elife.
2021;10. Epub 20211207. doi: 10.7554/eLife.71502. PubMed PMID: 34874266; PubMed
Central PMCID: PMCPMC8683085.
59.   Martinez-Zamudio R, Ha HC. Histone ADP-ribosylation facilitates gene transcription by
directly remodeling nucleosomes. Molecular and Cellular Biology. 2012;32(13):2490-502. Epub
20120430. doi: 10.1128/MCB.06667-11. PubMed PMID: 22547677; PubMed Central PMCID:
PMCPMC3434492.
60.   Bhagat TD, Von Ahrens D, Dawlaty M, Zou Y, Baddour J, Achreja A, et al. Lactate-
mediated epigenetic reprogramming regulates formation of human pancreatic cancer-associated
fibroblasts. eLife. 2019;8:e50663. Epub 20191101. doi: 10.7554/eLife.50663. PubMed PMID:
31663852; PubMed Central PMCID: PMCPMC6874475.
61.   Zhang D, Tang Z, Huang H, Zhou G, Cui C, Weng Y, et al. Metabolic regulation of gene
expression by histone lactylation. Nature. 2019;574(7779):575-80.
62.   Zhang W, Tan X, Lin S, Gou Y, Han C, Zhang C, et al. CPLM 4.0: an updated database
with rich annotations for protein lysine modifications. Nucleic Acids Research.
2022;50(D1):D451-D9. doi: 10.1093/nar/gkab849. PubMed PMID: 34581824; PubMed Central
PMCID: PMCPMC8728254.
63.   Liu X, Wang J, Boyer JA, Gong W, Zhao S, Xie L, et al. Histone H3 proline 16
hydroxylation regulates mammalian gene expression. Nature Genetics. 2022;54(11):1721-35.
doi: 10.1038/s41588-022-01212-x.
64.   Gantner BN, Palma FR, Kayzuka C, Lacchini R, Foltz DR, Backman V, et al. Histone
oxidation as a new mechanism of metabolic control over gene expression. Trends in Genetics.
2024;40:739-46.
65.   Unoki M, Masuda A, Dohmae N, Arita K, Yoshimatsu M, Iwai Y, et al. Lysyl 5-
hydroxylation, a novel histone modification, by Jumonji domain containing 6 (JMJD6). Journal
of Biological Chemistry. 2013;288(9):6053-62. Epub 20130109. doi: 10.1074/jbc.M112.433284.
PubMed PMID: 23303181; PubMed Central PMCID: PMCPMC3585044.
66.   Serra-Bardenys G, Peiro S. Enzymatic lysine oxidation as a posttranslational
modification. FEBS Journal. 2022;289(24):8020-31. Epub 20211002. doi: 10.1111/febs.16205.
PubMed PMID: 34535954; PubMed Central PMCID: PMCPMC10078733.
67.   Berlett BS, Stadtman ER. Protein oxidation in aging, disease, and oxidative stress.
Journal of Biological Chemistry. 1997;272(33):20313-6. doi: 10.1074/jbc.272.33.20313.
PubMed PMID: 9252331.
68.   Masson N, Ratcliffe PJ. HIF prolyl and asparaginyl hydroxylases in the biological
response to intracellular O_2 levels. Journal of Cell Science. 2003;116(15):3041-9.
69.   Trenzado C, Hidalgo MC, García-Gallego M, Morales AE, Furné M, Domezain A, et al.
Antioxidant enzymes and lipid peroxidation in sturgeon *Acipenser naccarii* and trout
Oncorhynchus mykiss. A comparative study. Aquaculture. 2006;254(1):758-67.
70.   Fontagné S, Lataillade E, Breque J, Kaushik S. Lipid peroxidative stress and antioxidant
defence status during ontogeny of rainbow trout (*Oncorhynchus mykiss*). British journal of
nutrition. 2008;100(01):102-11.
71.   Wagner JT, Knapp MJ, Podrabsky JE. Antioxidant capacity and anoxia tolerance in
Austrofundulus limnaeus embryos. Journal of Experimental Biology. 2019;222(12):jeb204347.
doi: 10.1242/jeb.204347.

72.   Lagedroste M, Reiners J, Knospe CV, Smits SHJ, Schmitt L. A structural view on the
maturation of lanthipeptides. Frontiers in Microbiology. 2020;11. doi:
10.3389/fmicb.2020.01183.

73.   Brennan DF, Barford D. Eliminylation: a post-translational modification catalyzed by
phosphothreonine lyases. Trends in Biochemical Sciences. 2009;34(3):108-14. doi:
10.1016/j.tibs.2008.11.005.

74.   Jones LH. Dehydroamino acid chemical biology: an example of functional group
interconversion on proteins. RSC Chemical Biology. 2020;1(5):298-304. doi:
10.1039/d0cb00174k.

75.   Lai K-Y, Galan SRG, Zeng Y, Zhou TH, He C, Raj R, et al. LanCLs add glutathione to
dehydroamino acids generated at phosphorylated sites in the proteome. Cell. 2021;184(10):2680-
95.e26. doi: 10.1016/j.cell.2021.04.001.

76.   Bogart JW, Bowers AA. Dehydroamino acids: chemical multi-tools for late-stage
diversification. Organic & Biomolecular Chemistry. 2019;17(15):3653-69. doi:
10.1039/c8ob03155j.

77.   Johnson AB, Barton MC. Hypoxia-induced and stress-specific changes in chromatin
structure and function. Mutation Research/Fundamental and Molecular Mechanisms of
Mutagenesis. 2007;618(1-2):149-62. doi: 10.1016/j.mrfmmm.2006.10.007.

78.   Kratzmeier M, Albig W, Hanecke K, Doenecke D. Rapid dephosphorylation of H1
histones after apoptosis induction. Journal of Biological Chemistry. 2000;275(39):30478-86. doi:
10.1074/jbc.M003956200. PubMed PMID: 10874037.

79.   Cheung WL, Ajiro K, Samejima K, Kloc M, Cheung P, Mizzen CA, et al. Apoptotic
phosphorylation of histone H2B is mediated by mammalian sterile twenty kinase. Cell.
2003;113(4):507-17. doi: 10.1016/s0092-8674(03)00355-6. PubMed PMID: 12757711.

80.   Fernandez-Capetillo O, Lee A, Nussenzweig M, Nussenzweig A. H2AX: the histone
guardian of the genome. DNA Repair. 2004;3(8-9):959-67. doi: 10.1016/j.dnarep.2004.03.024.

81.   Xiao A, Li H, Shechter D, Ahn SH, Fabrizio LA, Erdjument-Bromage H, et al. WSTF
regulates the H2A.X DNA damage response via a novel tyrosine kinase activity. Nature.
2009;457(7225):57-62. doi: 10.1038/nature07668.

82.   Rossetto D, Avvakumov N, Côté J. Histone phosphorylation. Epigenetics.
2012;7(10):1098-108. doi: 10.4161/epi.21975.

83.   Mojica EA, Kültz D. A strategy to characterize the global landscape of histone post-
translational modifications within tissues of nonmodel organisms. Journal of Proteome Research.
2024;23:2780−94. Epub 20230825. doi: 10.1021/acs.jproteome.3c00246. PubMed PMID:
37624673.

84.   Root L, Campo A, Macniven L, Con P, Cnaani A, Kültz D. A data-independent
acquisition (DIA) assay library for quantitation of environmental effects on the kidney proteome
of *Oreochromis niloticus*. Molecular Ecology Resources. 2021;21(7):2486-503. doi:
10.1111/1755-0998.13445.

85.   Kong AT, Leprevost FV, Avtonomov DM, Mellacheruvu D, Nesvizhskii AI. MSFragger:
ultrafast and comprehensive peptide identification in mass spectrometry-based proteomics.
Nature Methods. 2017;14(5):513-20. Epub 20170410. doi: 10.1038/nmeth.4256. PubMed PMID:
28394336; PubMed Central PMCID: PMCPMC5409104.

86.   Benjamini Y, Hochberg Y. Controlling the false discovery rate: A practical and powerful
approach to multiple testing. Journal of the Royal Statistical Society Series B. 1995;57:289-300.
doi: 10.1111/j.2517-6161.1995.tb02031.x.

87.   MacLean B, Tomazela DM, Shulman N, Chambers M, Finney GL, Frewen B, et al.
Skyline: an open source document editor for creating and analyzing targeted proteomics
experiments. Bioinformatics. 2010;26(7):966-8. Epub 20100209. doi:
10.1093/bioinformatics/btq054. PubMed PMID: 20147306; PubMed Central PMCID:
PMCPMC2844992.
88.   Du P, Zhang X, Huang C-C, Jafari N, Kibbe WA, Hou L, et al. Comparison of Beta-
value and M-value methods for quantifying methylation levels by microarray analysis. BMC
Bioinformatics. 2010;11:587. doi: 10.1186/1471-2105-11-587.
89.   Chappell K, Graw S, Washam CL, Storey AJ, Bolden C, Peterson EC, et al. PTMViz: a
tool for analyzing and visualizing histone post translational modification data. BMC
Bioinformatics. 2021;22(1):275. doi: 10.1186/s12859-021-04166-9.
90.   Wickham H. Elegant Graphics for Data Analysis. New York: Springer-Verlag; 2016.
1010