REST/NRSF phosphorylation by CaMKIV regulates its transcriptional repressor activity and half-life

Hanako Tsushima Semini[1,2] *, Beatrice Corradi[1,2] *, Dmytro Shmal[1], Martina Chiacchiaretta[1 ^], Nara Liessi[4], Andrea Di Fonzo[4], Emanuele Carminati[1,2 ^], Andrea Armirotti[4], Anna Fassio[2,3], Luca Maragliano[1,5 §], Anna Rocchi[1 ^], Fabio Benfenati[1,3 §]

[1] Center for Synaptic Neuroscience and Technology, Istituto Italiano di Tecnologia, Largo Rosanna Benzi, 10, 16132 Genova, Italy
[2] Department of Experimental Medicine, Università degli Studi di Genova, Viale Benedetto XV 3, 16132 Genova, Italy
[3] IRCCS Ospedale Policlinico San Martino, Largo Rosanna Benzi 10, 16132 Genova, Italy
[4] Analytical Chemistry Facility, Istituto Italiano di Tecnologia, Genoa, Italy
[5] Department of Life and Environmental Sciences, Polytechnic University of Marche, Via Brecce Bianche, 60131, Ancona, Italy

* Equal contribution

Running title:
Regulation of REST/NRSF by CaMKIV

^ Current address:
Martina Chiacchiaretta: Department of Neuroscience, Tufts University School of Medicine, Boston, MA
Emanuele Carminati: AGC Biologics, Bresso, Italy
Anna Rocchi: Polo Genomics Genetics Biology, Siena, Italy

§ **Corresponding authors:**

Fabio Benfenati, MD
Istituto Italiano di Tecnologia
Genova, Italy
Email: fabio.benfenati@iit.it

Luca Maragliano, PhD
Istituto Italiano di Tecnologia and Polytechnic University of Marche
Genova and Ancona, Italy
Email: l.maragliano@staff.univpm.it

ABSTRACT

REST is a repressor of a large cluster of neural genes containing RE1 motifs. In neurons, REST homeostatically regulates intrinsic excitability and synaptic transmission, However, whether REST can be regulated by Ca^{2+} and Ca^{2+}-activated kinases is unknown. Here, we investigated Ca^{2+}/calmodulin-dependent protein kinases (CaMKs) as upstream regulators of REST fate and activity. We show that REST is phosphorylated by CaMKIV at a main Ser-322 site located within the linker between the 5th and the 6th N-terminal Zn-finger domains. Phosphomimic REST mutant in Serine-322 decreased REST repressor activity and caused its transition from nucleus to cytosol, followed by degradation. Molecular dynamics simulations of the phosphomimic N-terminal REST and the DNA RE1 sequence revealed a sharp decrease in the stability of the REST-RE1 binding interface. Moreover, the homeostatic effects of CaMKIV on the amplitude of excitatory synaptic currents were inhibited by the genetic deletion of REST. The results demonstrate that REST phosphorylation by CaMKIV regulates the repressor activity of REST on neural genes and has a crucial role in the homeostatic regulation of REST levels.

Keywords: REST/NRSF degradation, nuclear/cytoplasmic translocation, Zn-finger domain, Ca^{2+}/calmodulin protein kinases

INTRODUCTION

REST (RE1-Silencing Transcription Factor), also known as NRSF (Neuron-Restrictive Silencer Factor), is a transcriptional repressor that plays a key role in gene regulation (Bruce et al., 2004; Balls & Mandel, 2005). It contains 9 Zn-finger domains, through which it binds to a specific DNA sequence of 17-21 nucleotides named Repressor Element-1 (RE1; Mori et al., 1992; Kraner et al., 1992), at which it recruits co-factors (such as Co-REST, mSin3, and HDACs) to modify chromatin structure and repress the transcription of the downstream gene. RE1 elements have been identified in over 1,800 genes in the human and mouse genomes, displaying a variable number of copies and variable affinity for REST (Bruce et al., 2004). Many of the target genes of REST are neuronal genes, while others are involved in cell cycle regulation and cytoskeleton assembly. In non-neuronal tissues, REST is active and suppresses neuronal gene expression, while during neuronal differentiation, REST is downregulated, allowing neuronal genes to be expressed (Ballas et al., 2005; Yeo et al., 2009; Rodenas-Ruano et al., 2005; Aoki et al., 2011; see Baldelli & Meldolesi for review).

In addition to its role in neuronal differentiation, residual REST expression is preserved in mature neurons, albeit at low levels, and plays post-developmental roles in the brain. In neurons, REST is an essential actor of homeostatic plasticity responses. It is promptly induced by hyperactivity and migrates into the nucleus, where it inhibits the transcription of genes governing intrinsic excitability and synaptic transmission (Baldelli & Meldolesi, 2015; Michetti & Benfenati, 2024). Under conditions of neuronal hyperactivity, the induction of REST inhibits the expression of voltage-gated Na^+ channels (Pozzi et al., 2013), blunts excitatory synaptic transmission by acting at the presynaptic level (Pecoraro-Bisogni et al., 2018), and boosts inhibitory synaptic transmission impinging onto excitatory neurons (Prestigio et al., 2021). REST is also involved in controlling changes in neuronal networks in response to early-life adversity, promoting resilience (Singh-Taylor et al., 2018; Mampay & Sheridan, 2019; Bolton et al., 2020). In addition, REST was also identified to have a key role in the regulation of lifespan and in stress resistance during aging by repressing hyperactivity (Lu et al., 2014; Zullo et al., 2019) and preventing senescence by regulating the autophagic flux (Rocchi et al., 2021). A homeostatic effect against hyperactivity was also observed in astrocytes, where REST increases glutamate uptake and potassium clearance from the synaptic environment (Centonze et al., 2023).

Beyond its canonical role in neurodevelopment and its pivotal role in regulating brain homeostasis, REST dysregulation has been implicated in a variety of pathological contexts, including epilepsy, neuroinflammation, brain ischemia, pain, Alzheimer's and Huntington's diseases, where it may contribute to neuronal vulnerability or adaptive responses (Buckley et al., 2010; Lu et al., 2014; McClelland et al., 2011, 2014; Hall et al., 2024; Carminati et al., 2021; Buffolo et al., 2021; Natali et al., 2023; Carminati et al., 2020; see Hwang & Zukin, 2018 for review).

REST shuttles between the nucleus, where it associates with a variety of interactors and actuates transcriptional repression, and the cytoplasm, where it remains inactive. Several regulators of REST favor its retention in the nucleus, such as the co-factors mSin3 and Co-REST and RILP (Shimojo & Hersh, 2006; Ballas & Mandel, 2005), and optogenetic disruption of these interactions

inhibits REST activity in the nucleus (Paonessa et al., 2016). Although no full-length experimental structure is available for REST, domain mapping suggests that it is a modular protein comprising N-terminal and C-terminal repression domains and a central Zn-finger DNA-binding region. The latter adopts the canonical C2H2 architecture, consisting of nine conserved Cys2–His2 motifs arranged in tandem, each coordinating a Zn^{2+} ion and mediating DNA binding (Palm et al., 1998; Rockowitz et al., 2014; Brenner et al., 2005). Structural modeling and sequence alignment suggest that REST's Zn-finger (ZNF) domains adopt the classical ββ-α fold, namely a β-hairpin followed by an α-helix, stabilized by Zn^{2+} coordination (Wolfe et al., 1999). Each Zn-finger module typically interacts with a tri-nucleotide segment of DNA, allowing REST to contact an extended nucleotide sequence and achieve high specificity (Klug, 2010).

REST activity is tightly regulated, not only at the transcriptional and epigenetic levels, but also through post-translational modifications, among which phosphorylation plays a central role. Indeed, phosphorylation was demonstrated to regulate REST half-life and turnover. Casein kinase 1 (CK1) was found to associate and phosphorylate REST at four serine residues (S^{1013}, S^{1024}, S^{1027}, S^{1030}) within two degron motifs in the C-terminal region, enabling recognition by the E3 ubiquitin ligase β-TrCP and targeting it for proteasomal degradation (Westbrook et al., 2008; Guardavaccaro et al., 2008; Kaneko et al., 2014). Moreover, extracellular signal-regulated kinases 1 and 2 (ERK1/2) were found to phosphorylate the proline-directed phosphorylation motifs in REST. ERK1/2 phosphorylation promotes rapid REST degradation, relieving repression of neuronal gene expression, while their dephosphorylation by the C-terminal domain small phosphatase 1 (CTDSP1) increases REST stability and half-life (Nesti et al., 2014). This ERK-mediated regulation plays an important role during neurogenesis, where ERK activation in response to growth factors leads to REST downregulation, allowing neuronal differentiation.

Despite these insights, the understanding of the full spectrum of REST phosphorylation sites and their functional consequences remains incompletely understood. While phosphorylation has been predominantly seen as a mechanism to downregulate REST in neural progenitors and differentiating neurons, it could also participate in regulating the downstream effects of REST overexpression during hyperactivity in mature neurons by negatively modulating REST transcriptional repression and returning REST levels to baseline. As calcium entry and calmodulin activation are the most likely signals triggering homeostatic plasticity and REST overexpression (Turrigiano, 2008, 2011), here we investigated whether REST can also be phosphorylated by Ca^{2+}/calmodulin-dependent kinases (CaMKs) and whether its phosphorylation has any effect on its repressor activity, nuclear/cytosolic shuttling, and half-life.

We have found that, consistent with the presence of consensus phosphorylation sites for CaMKs, REST is phosphorylated by CaMKIV in primary neurons on a main site intercalated between the 5th-6th Zn-finger domains, and its phosphorylation causes dissociation from the DNA RE1 motif, inhibition of its transcriptional repressor activity, escape from the nucleus to the cytoplasm where it undergoes degradation by extra-lysosomal proteases or pH-independent lysosomal proteases. Moreover, the homeostatic effects of CaMKIV on excitatory synaptic transmission were inhibited by the genetic deletion of REST. Overall, the effects of the CaMK-dependent phosphorylation of REST are important to blunt the homeostatic transcriptional responses of REST to hyperactivity

and to return them to the basal condition, indicating a key physiological role of the interplay of CaMKIV/REST in homeostatic plasticity responses.

RESULTS

Inhibition of Ca²⁺-calmodulin-dependent kinases increases the levels of REST protein in primary neurons

Recent evidence suggests that post-translational modifications, including phosphorylation and ubiquitination, influence the stability and turnover of the REST protein. ERK1/2 and CK1 kinases phosphorylate REST and promote its degradation by the ubiquitin proteasome system (Westbrook et al., 2008; Kaneko et al., 2014; Nesti et al., 2014). Given that REST is upregulated in mature neurons by hyperactivity, a condition in which Ca^{2+} entry occurs, we tested the possibility that phosphorylation by CaMKs regulates REST turnover and fate in neurons. We first evaluated whether Ca^{2+}/calmodulin (CaM) signaling interferes with REST stability. When primary cortical neurons were incubated with the CaM inhibitor W7, an increase in REST protein was observed (**Figure 1A**).

Figure 1. Inhibition of Ca²⁺-calmodulin-dependent activities increases the endogenous levels of REST protein in primary mouse cortical neurons. A. *Left:* Representative REST immunoblots of primary cortical neurons (DIV14) treated with the cell-permeable competitive Ca^{2+}-calmodulin antagonist W7 (20 µM). Calnexin was used as a control for equal loading. *Right:* Mean (± sem with individual experimental points) REST protein expression shown as a ratio between REST and calnexin immunoreactivities. **p=0.0071; unpaired Student's *t*-test (n=5 independent neuronal preparations). **B.** *Left:* Representative REST immunoblots of DIV 14 primary cortical neurons treated with the phosphatase inhibitor cyclosporin A (CsA; 1 µM). Calnexin was used as a control for equal loading. *Right:* Mean (± sem with individual experimental points) REST expression shown as a ratio between REST and calnexin immunoreactivities. p=0.36; unpaired Student's *t*-test (n=5 independent neuronal preparations). **C.** *Left:* Representative REST immunoblots of DIV14 primary cortical neurons treated with vehicle (Ctrl), the cell-permeable CaMK inhibitor KN93 (10 µM), or its inactive derivative KN92 (10 µM). Calnexin was used as a control for equal loading. *Right:* Mean (± sem with individual experimental points) REST protein expression shown as a ratio between REST and calnexin immunoreactivities for the three treatments. *p=0.031, Ctrl *vs* KN93; *p=0.032, KN93 *vs* KN92; one-way ANOVA/Holm-Sidak's tests (n=5 independent neuronal preparations).

Since CaM activates various kinases and phosphatases, we asked which downstream effector could be involved. Primary cortical neurons were treated with either the calcineurin inhibitor ciclosporin A (CsA) or with the competitive CaMK blocker KN93. While inhibition of calcineurin did not affect REST protein levels (**Figure 1B**), KN93, at concentrations inhibiting all CaMKs, induced a significant increase in REST protein levels that was completely absent when its inactive analogue KN92 was used (**Figure 1C**). None of the treatments was associated with any change in REST mRNA levels, as shown by RT-qPCR analysis (**Figure S1**). The same treatments, performed on primary cortical astrocytes, did not alter the levels of both REST protein and mRNA (**Figure S2**), indicating that the effects of CaMK inhibition were specific for neurons.

Expression of constitutively active CaMKI and CaMKIV decreases the expression and the transcriptional repressor activity of REST in HEK293T cells

KN93 is a broad-spectrum CaMK inhibitor with several molecular targets, and thus, the results do not identify the specific CaMKs that mediate the effects on REST protein levels. To address this point, we overexpressed CaMKI, CaMKII, and CaMKIV as the wild-type (WT) form, and as constitutively active (CA) and kinase-dead (KD) variants in HEK293T cells that constitutively express REST. Under these experimental conditions, we evaluated (i) the changes in the expression levels of the REST protein by immunoblotting as a ratio between REST and calnexin immunoreactivities; and the effects on the REST repressor activity by using a gene reporter assay in which *Renilla* luciferase is driven by the SV40 promoter fused to a triple RE1*cis* site from the promoter region of the *SCN2A* gene, a well-known REST target (**Figure 2**; Paonessa et al., 2016; see Materials and Methods).

Figure 2. Expression of wild-type, constitutively active, or kinase-dead CaM kinases in HEK293T cells implicates CaMKI and CaMKIV in the control of protein levels and transcriptional repressor activity of REST. A-C. *Left:* Representative immunoblots monitoring the expression of wild-type (WT), constitutively active (CA), and kinase-dead (KD) CaMKs, REST, and calnexin, respectively, in HEK293T cells, 48 h after transfection with the kinase constructs. The expression of CaMKI (**A**) and CaMKII (**B**) was followed using antibodies to HA and GFP fused tags, respectively, while CaMKIV (**C**) was detected with anti-CaMKIV-specific antibodies. *Middle:* Mean (± sem with individual experimental points) REST protein levels after expression of the three variants of CaMKI (**A**), CaMKII (**B**), and CaMKIV (**C**), shown as REST/calnexin immunoreactivity ratios. *Right:* A RE1-Luciferase construct, with *Renilla* Luciferase (*RLuc*) gene cloned downstream of the RE1 sequence, was used to detect the repressor activity of endogenous REST in HEK293T cells, transfected with the three variants of CaMKI (**A**), CaMKII (**B**), and CaMKIV (**C**). The *Renilla/Firefly* Luciferase chemiluminescence ratio is shown as mean (± sem with individual experimental points). Panel A - REST/calnexin: ***p=0.0007 CA *vs* KD, ****p<0.0001 CA *vs* WT; *RLuc*: ****p<0.0001 CA *vs* WT or KD. Panel B - REST/calnexin: p=0.86 CA *vs* WT, p=0.98 CA *vs* KD; *RLuc*: p=0.74 CA *vs* WT, p=0.58 CA *vs* KD. Panel C - REST/calnexin: ****p<0.0001 CA *vs* WT or KD; *RLuc*: **p=0.0018 CA *vs* WT; **p=0.0057 CA *vs* KD; one-way ANOVA/Tukey's tests (n=8-9 independent HEK293T cell preparations).

Western blotting analysis revealed that the constitutively active forms of both CaMKI and CaMKIV significantly decreased REST levels, while the constitutively active form of CaMKII was virtually ineffective (**Figure 2A-C**, *left and middle panels*). No effects with respect to the WT forms of the kinases were observed with the respective kinase-dead variants, suggesting that the endogenous activation of these kinases is not functional in this cell line under these conditions. In agreement with the observed decrease in REST levels, we observed a markedly increased luciferase signal in HEK293T cells transfected with the constitutively active forms of both CaMKI and CaMKIV and no effect with the active form of CaMKII (**Figure 2A-C**, *right panels*), indicating a corresponding drop in the REST repressor activity at RE1 sites. The data indicate that CaMKI and CaMKIV are potentially implicated in the regulation of REST protein stability, at least in HEK293Tcells.

Knockdown of CaMKIV, but not of CaMKI, in primary neurons recapitulates the effects of CaMK inhibition on REST protein levels

To get insights into the mechanism underlying the CaMKI/CaMKIV-mediated regulation of REST levels in neurons, the REST protein level and activity were analyzed after silencing of either CaMKI or CaMKIV by RNA interference using lentiviral vectors (**Figure 3** and **Figure S3**). While we used a very effective, previously reported shRNA for CaMKII (Ageta-Ishihara et al., 2009), we tested two distinct shRNAs (sh#1 and sh#2) for their efficiency in knocking down CaMKIV (**Figure S3A**). Both shRNAs effectively downregulated CaMKIV in primary neurons with >60% decrease in CaMKIV levels. We selected sh#2 for the successive experiments and first tested the specificity of the knockdown with respect to the other CaMKs and REST. The shRNA for CaMKIV was very selective and was not associated with any change in REST, CaMKI, or CaMKII mRNA levels (**Figure S3B-D**).

To ascertain whether both CaMKI and CaMKIV are active in decreasing REST levels in neurons, we silenced either CaMKI or CaMKIV by transducing neurons with lentivirus encoding the respective shRNAs. Western blotting analysis revealed that silencing CaMKIV, but not CaMKI, induced an increase in REST protein levels (**Figure 3A,B**). In conclusion, the data suggest that CaMKIV is responsible for the post-transcriptional modulation of REST protein levels in primary neurons, which is in line with the preliminary results obtained with the CaMK inhibitors. Notably, while CaMKI is expressed by neurons and astrocytes, CaMKII and CaMKIV are virtually neuron-specific (**Figure S4**), indirectly confirming the role played by CaMKIV in the regulation of REST

levels that was observed in neurons but not astrocytes under the same experimental conditions. Moreover, the distinct effects of CaMKI and CaMKIV in neurons may depend on their subcellular distribution. Whereas CaMKIV is predominantly nuclear, and therefore positioned to directly interact with nuclear REST, CaMKI is largely cytosolic, which limits its ability to efficiently target REST in neurons despite being effective in heterologous systems such as HEK293T.

Figure 3. Downregulation of CaMKIV, but not of CaMKI, increases the endogenous level of REST in mouse primary cortical neurons. A. Primary mouse cortical neurons were transduced at DIV 7 with lentiviruses encoding shRNAs to downregulate CaMKIα (shCaMKI) or a scrambled version thereof (shSCR). *Left:* Representative immunoblots monitoring the expression of endogenous CaMKIα, REST, and calnexin in the neuronal preparations at DIV14. *Middle and Right:* Mean (± sem with individual experimental points) CaMKIα (*middle*) and REST (*right*) expression shown as ratios with calnexin immunoreactivity. CaMKIα: ****p<0.0001; REST/NRSF: p=0.53; unpaired Student's *t*-test (n=5 independent preparations). **B.** Primary mouse cortical neurons were transduced at DIV 7 with lentiviruses encoding shRNAs to knock down CaMKIV (shCaMKIV) or a scrambled version thereof (shSCR). *Left:* Representative immunoblots monitoring the expression of endogenous CaMKIV, REST, and calnexin in the neuronal preparations at DIV14. *Middle and Right:* Mean (± sem with individual experimental points) CaMKIV (*middle*) and REST (*right*) expression, shown as ratio with calnexin immunoreactivity. CaMKIV: **p=0.0024; REST/NRSF: *p=0.025; unpaired Student's *t*-test (n=7 independent preparations).

REST degradation is triggered by CaMKIV phosphorylation

Phosphorylation by CK1 and ERK1/2 has been shown to target REST to the ubiquitin-proteasome degradation. Thus, we sought to investigate whether phosphorylation by CaMKIV, also decreasing the levels of REST protein, was associated with a similar mechanism. We initially considered three possible pathways of degradation, namely the proteosomal, macroautophagic, lysosomal, or cytosolic proteolytic degradation (**Figure 4A**).

To this aim, we transfected HEK293T cells with WT, constitutively active, or kinase-dead CaMKIV and treated them with cycloheximide (CHX) to block REST neo-synthesis. Then, we treated the cells with either vehicle or MG-132, a specific inhibitor of the ubiquitin-proteasome system. As reported above (see Figure 2), only cells expressing CA-CaMKIV exhibited significantly decreased REST levels. However, this downregulation of REST protein was not attributable to proteasome-mediated degradation, as MG-132 was totally ineffective in antagonizing the decreased REST levels associated with the expression of the constitutively active kinase (**Figure**

4B). The absence of effects of MG-132 in cells expressing WT- or KD-CaMKIV also indicates that, in the absence of specific activation of signal transduction pathways, REST is not constitutively targeted to the proteasome.

Next, we addressed the possibility that REST is degraded by alternative pathways, such as in autophagolysosomes or lysosomes. To ascertain this, we transfected HEK293T cells with either WT- or CA-CaMKIV and blocked REST neo-synthesis (see above) before treating the cells with vehicle, bafilomycin A1 (an inhibitor of v-ATPase responsible for lysosomal acidification), or leupeptin (an inhibitor of lysosomal and cytosolic proteases). The differential effect of these two blockers is demonstrated by the significant increase in the autophagy-related proteins LC3B-II and p62 observed after treatments with bafilomycin A1, but not with leupeptin (**Figure 4C**, *left panel*; **Figure S5A,B**). The results show that only leupeptin could reverse the drop in REST protein levels associated with expression of CA-CaMKIV (**Figure 4C**), indicating that REST degradation is likely mediated by proteases that are either cytosolic or not impaired by the loss of lysosomal acidification.

Figure 4. CaMKIV phosphorylation of REST triggers its lysosomal degradation. A. Schematics of the four possible degradation pathways that CaMKIV-phosphorylated REST can undertake and the three inhibitors that were used to dissect the fate of REST. UPS, ubiquitin proteasome system. **B.** HEK293T cells were transfected with WT, constitutively active (CA), or kinase dead (KD) CaMKIV and treated with cycloheximide (CHX) to inhibit further translation. Then, cells were treated for 16 h with either vehicle (DMSO) or MG132 (5 μM), an inhibitor of the ubiquitin-proteasome system. *Left:* Representative immunoblots monitoring the expression of wild-type (WT), constitutively active (CA), and kinase-dead (KD) CaMKs, REST, and calnexin, respectively, in HEK293T cells, 48 h after transfection with the kinase constructs. *Right:* Mean ($\pm$ sem with individual experimental points) REST protein levels, shown as ratios with calnexin immunoreactivity under the various experimental conditions. No effects of inhibition of the ubiquitin-proteasome system were observed. *p=0.017, CA *vs* WT in DMSO; **p=0.0048, CA *vs* WT in MG132); 2-way ANOVA/Tukey's tests (n=4 independent preparations). **C.** HEK293T cells were transfected with either WT or constitutively active CaMKIV (CA CaMKIV) and treated with cycloheximide to inhibit further translation. Then, cells were treated for 16 h with vehicle (DMSO), bafilomycin A1 (Baf, 10 and 100 nM; an inhibitor of lysosomal acidification and autophagy), and leupeptin (Leu, 40 μM; an inhibitor of the lysosomal and cytosolic proteases). *Left:* Representative immunoblots monitoring the expression of REST, calnexin, and the autophagy markers p62 and LC3B-II. *Right:* Mean ($\pm$ sem with individual experimental points) of REST protein levels, shown as ratios with calnexin immunoreactivity under the various experimental conditions. ***p=0.0010, CA *vs* WT in DMSO and WT/DMSO *vs* CA/Baf10 and 100; *p=0.0264 CA/Leu *vs* CA/DMSO; 2-way ANOVA/Tukey's tests (n=4 independent preparations).

REST is phosphorylated by CaMKIV on Serine322 between the 5[th] and the 6[th] Zn-finger domain

CaMKIV, together with CaMKI and CaMKII, is collectively defined as a "multifunctional CaM kinase" for its ability to phosphorylate a wide range of substrates (Xu et al., 2024; Kaiser et al., 2024). CaMKIV needs to be activated in the cytoplasm by CaMK kinase (CaMKK) before entering the nucleus, where it acts as a potent activator of Ca^{2+}-dependent transcriptional effects through phosphorylation of downstream transcriptional effectors, such as CREB, MeCP2, and HDAC4 (Lemrow et al., 2004; Matthews et al., 1994; Xu et al., 2024). We thus searched for the presence of putative consensus sequences for CaMKIV phosphorylation in the primary structure of mouse and human REST (**Figure 5A**; White et al., 1998).

We identified three putative CaMKIV phosphorylation sites within the 8 ZNF central cluster region of mouse and human REST, namely Serine177 (S177), Threonine294 (T294), and Serine322 (S322), located at 1-position of the ZNF linkers L1, L4, and L5. All three sites display the required -3 Arginine residue but have the required hydrophobic residue at -4, rather than at the canonical -5 position (where they have a basic Histidine). While all three putative sites have a hydrophobic residue at +4 that is highly preferred by CaMKI, T294, and S322 are particularly interesting since they display a +2 acidic residue (Glutamate) found in optimal consensus sequences for CaMKII. In contrast, S177 has a basic residue instead. The three putative phosphorylation sites were analyzed for their phosphorylation potential for CaMKs using the prediction software Musite and NetPhos3.1. As shown in **Table S1**, in both human (hREST) and mouse (mREST) sequences, the S322 site has the highest phosphorylation potential, while S177 displays the lowest. In addition to these structural issues, the REST peptide linkers encompassing the T294 and S322 putative sites share a strong analogy with other transcription factors containing ZNF clusters, such as the transcription factor Ikaros (Jantz & Berg, 2004), in which ZNF domains are usually organized in arrays joined by short 5-7-mer linker sequences that often start with a phosphorylatable residue. In the REST protein, the linker peptide containing the putative site 1 (S177) is the least conserved from mouse to man and does not have any resemblance to the short linker peptide sequences joining adjacent ZNF domains, being a long 36-mer peptide connecting ZNF1 with ZNF2. On the contrary, the 7-mer T294- and S322-containing linkers L4

and L5 have a strong resemblance to Icaros ZNF linkers. For all these reasons, we concentrated our attention on T294 and S322 and tested whether they can be phosphorylated by CaMKIV (**Figure 5B**).

To this aim, we expressed Myc-tagged recombinant mREST in HEK293T cells, affinity-purified it with anti-Myc magnetic beads, and phosphorylated it on beads by purified constitutively active CaMKIV. Purified phosphorylated mREST was subjected to tryptic cleavage and tandem mass spectrometry (MS-MS). Tandem mass spectra for L4 (top) and L5 (bottom) tryptic peptides revealed phosphorylation of S322, but not of T294, as clearly demonstrated by the corresponding diagnostic sequence fragment ions (**Figure 5C**).

LINKER 1 (L1): S177AKKFFVEESAEKQAKAWESGSSPAEEGEFSKGPIR
LINKER 4 (L4): T294GERPYK
LINKER 5 (L5): S322GEKPFK

Figure 5. Serine-322 in the linker between the 5th-6th Zn-finger domains is the major CaMKIV phosphorylation site in mouse REST. A. Alignment of the primary structures of human, rat, and mouse REST. Amino acids are shown in single-letter code. Sequences shaded in green refer to the Zn-finger domains (numbered 1 to 6), while three putative phosphorylation sites ($S^{177/182}$, $T^{294/299}$, and $S^{322/327}$ in mouse and human, respectively) that match the *consensus* sequence for CaMKs (S/T-XX-R/K) and are conserved between mouse and human REST are highlighted in red. **B.** The S^{177}, T^{294}, and S^{322} phosphorylation sites encompass the linker peptides between the 1st-2nd (L1), the 4th-5th (L4), and the 5th-6th (L5) in mouse REST Zn-finger domains, respectively. **C.** Tandem mass spectra for L4 (top) and L5 (bottom) tryptic peptides derived from HA-tagged recombinant mouse REST expressed in HEK293T cells, affinity-purified with anti-HA magnetic beads, and phosphorylated on beads by purified constitutively active CaMKIV. L5 shows phosphorylation of S322, as clearly demonstrated by the corresponding diagnostic sequence fragment ions (asterisk in the spectrum).

Phosphomimic mutants of REST translocate to the cytoplasm, where they undergo degradation

The location of T294 and S322 in the very short L4/L5 ZNF linkers of mREST resembles that of the transcription factor Ikaros, in which phosphorylation of the first residue in the short ZNF linkers impairs the binding of the transcription factor to DNA (Jantz & Berg, 2004). This raises the possibility that phosphorylation of S322 (and, possibly, of T294) affects the interaction between REST and the RE1 motifs in the DNA. To verify that, we replaced either or both phosphorylatable residues with negatively charged Glutamate to generate phosphomimic REST mutants and examine their activity on REST nucleus/cytoplasm shuttling, fate, and transcriptional repression (**Figure 6**). HEK293T cells were transfected with either WT mREST or with the single (T294E, TE, and S322E, SE) or the double (TESE) mREST pseudo-phosphorylated mutants, using cytosolic mCherry and transfection reporter and Hoechst staining to morphologically identify the nuclear areas (**Figure 6A**, *left panel*). Morphometric analysis of the presence of REST immunoreactivity within the nuclear area was performed by calculating the Manders' overlap coefficient, which showed a significant degree of the extent of colocalization for the SE mutant and smaller decreases for the other mutants, suggesting that the SE phosphomimic mutant of mREST has a lower affinity for its DNA target sequences and partially escapes from the nucleus (**Figure 6A**, *right panel*).

The successive analysis aimed at evaluating the total REST protein levels after transfection of HEK293T cells with either WT mREST or with the single or double mREST pseudo-phosphorylated mutants revealed that the decreased nuclear localization of REST was paralleled by an increased cytoplasmic degradation that was particularly dramatic for the SE mutant (**Figure 6B**).

Finally, we analyzed REST repressor activity after transfection of HEK293T cells with the three phosphomimic mutants and compared it with WT mREST in the absence or presence of co-transfected CA-CaMKIV by using the gene reporter assay in which *RLuc* is driven by the SV40 promoter fused to a triple RE1*cis* site. Strikingly, *RLuc* expression was significantly increased with the SE REST mutant, mimicking the decrease in transcriptional repression observed with CA-CaMKIV (**Figure 6C**). Taken together, the data on the REST phosphomimic mutant SE indicate that phospho-S322 is the phosphorylation site that mimics the effects of CA-CaMKIV overexpression on REST localization, stability, and repressor activity.

Figure 6. The phosphomimic S322E REST mutant has an impaired repressor activity and shuttles to the cytosol, where it undergoes degradation in HEK293T cells. A. HEK293T cells were transfected with WT REST or with its T294E (TE), S322E (SE), and S322E/T294E (TESE) phosphomimic mutants with soluble mCherry as transfection reporter. *Left:* Representative samples for the 4 experimental groups are shown, in which REST immunostaining, Hoechst nuclear staining, and mCherry fluorescence were used to evaluate the cytosolic/nuclear localization of REST mutants. Right: Mean (± sem with individual experimental points) of the Manders' coefficient evaluating the extent of overlap between REST immunoreactivity and Hoechst nuclear staining. *p=0.039; one-way ANOVA/Dunnett's tests. **B.** A RE1-Luciferase construct, with *Renilla* luciferase (*RLuc*) gene cloned downstream the promoter region containing triple RE1 sequence from *Scn2a*, was used to assess the extent of transcriptional repression exerted by the expression of WT REST or of its phosphomimic mutants in HEK293T cells. *Left:* The *Renilla/Firefly* Luciferase chemiluminescence ratio is shown as mean (± sem with individual experimental points). *Right:* As a positive control, either WT or catalytically active (CA) CaMKIV was transfected in the same HEK293T cells transfected with WT mREST. *p=0.032; one-way ANOVA/Dunnett's tests. *p=0.0035; unpaired Student's *t*-test (n=3 independent experiments). **C.** HEK293T cells were transfected with WT REST or with its T294E (TE), S322E (SE), and S322E/T294E (TESE) phosphomimic mutants. The levels of phosphomimic SE are strongly decreased in comparison

Phosphomimic substitutions destabilize the mREST:RE1 interaction interface and promote local structural collapse

To evaluate the impact of phosphomimic substitutions on the stability of the mREST:RE1 interaction interface, we conducted atomistic molecular dynamics (MD) simulations on both wild-type and mutant complexes. We focused on the mREST segment V130-H267. The mREST:RE1 complex was reconstructed using a multi-step protocol that included structural modeling, protein-DNA docking, complex refinement, and optimization of Zn^{2+} coordination (**Figures S6-S11**). The resulting structure was used as the starting conformation for three independent 1-µs replicate MD simulations (indicated with R1 to R3).

To monitor global, structural deviations over time, we computed the Root Mean Square Deviation (RMSD) using all the heavy atoms of the protein and of the DNA. The RMSD profiles are stable around values between 2.8 and 4 Å, indicating no significant structural deviations from the starting configuration (**Figure S12**). We quantified the integrity of the protein-DNA interface by measuring the distance between a selected phosphate atom of the DNA backbone, adjacent to the first Zn-finger module, and the center of mass (COM) of the Zn-coordinating side chain atoms (Cysteine and Histidine residues) as a distance-based descriptor. This distance was stable throughout all simulations of the WT system, with average values of 10.3 ± 2.4 Å (R1), 11.1 ± 1.5 Å (R2), and 7.6 ± 0.7 Å (R3), and no local structural deviations were observed (**Figure 7A**), indicating that the WT mREST:RE1 complex maintains a stable interface over time.

To evaluate the potential destabilizing effects of phosphorylation-mimicking mutations, two mREST variants were generated: S322E and T294E, in which S322 and T294, respectively, were substituted with Glutamate. Each mutant was simulated in nine independent 1-µs MD replicas. For the S322E mutant, three replicas (R2, R4, R5) showed an increase in the COM–phosphate distance to 11.2 ± 3.3 Å (R2), 11.8 ± 2.5 Å (R4), and 16.1 Å (R5), with maximum values reaching 24.8 Å, 24.3 Å, and 30.6 Å, respectively, indicating dissociation of the first Zn-finger domain from its DNA binding site (**Figure 7B**). Three other replicas (R7, R8, R9) exhibited structural irregularities, prompted by DNA termini fragilization, leading to complete system collapse in R9 (**Figure S14**). The remaining three trajectories (R1, R3, R6) maintained stable interactions, with average distances of 8.8 ± 1.7 Å, 7.5 ± 0.8 Å, and 9.9 Å ± 1.8 Å, and lower peak excursions (13.6–17.7 Å) (**Figure S13**). For the T294E mutant, two replicas (R1, R4) displayed interface dissociation, with average distances of 11.9 ± 2.2 Å and 10.3 ± 2.6 Å, and maximum distance values of 22.4 Å and 25.0 Å (**Figure S15A**). Two trajectories (R8, R9) showed DNA fragilization, and five replicas (R2, R3, R5, R6, R7) remained structurally stable, with average distances ranging from 7.8 ± 1.0 Å to 11.6 ± 1.6 Å and peak values between 13.0 and 20.9 Å (**Figure S15B**). Taken together, these results show that both phosphomimic substitutions at S322 and T294 introduce measurable destabilization of the mREST:RE1 interaction interface, but that the interface dissociation and structural fragilization are more probable and of higher impact in S322E.

The CaMKIV/REST interaction underlies the homeostatic effects of CaMKIV on the excitatory synaptic transmission in primary neurons

CaMKIV is known to couple increases in intracellular Ca^{2+} to compensatory changes in excitatory synaptic transmission (Joseph & Turrigiano, 2017). Having identified a mechanistic link between CaMKIV activation and REST activity and half-life, we investigated the interplay between CaMKIV and REST by studying excitatory synaptic transmission in primary cortical neurons. We knocked down CaMKIV by RNA interference in primary cortical neurons derived from conditional REST knockout mice (Rest[GTi]) in which the REST gene was deleted by transduction of Cre recombinase (Cre) or left intact by transduction with a catalytically inactive Cre deletion mutant (ΔCre) and measured miniature excitatory postsynaptic currents (mEPSCs) by patch-clamp in voltage-clamp configuration. Primary neurons in which CaMKIV was knocked down displayed an increase in mEPSC amplitude and no effects on the EPSC frequency, in agreement with previous reports (Joseph & Turrigiano, 2017). On the other hand, REST KO promoted an increase in excitatory strength with augmented mEPSC frequency and no significant effects on mEPSC amplitude (Pegoraro-Bisogni et al., 2018). Interestingly, the concomitant REST deletion virtually abolished the effects of CaMKIV silencing (interaction), indicating that the Ca^{2+}-dependent homeostatic effects of CaMKIV are dependent, at least in part, on the presence of REST (**Figure 8**).

Figure 8. The effects of CaMKIV knockdown on excitatory synaptic transmission in primary cortical neurons are abolished by the conditional knockout of REST/NRSF. A. Representative traces of mEPSCs recorded in primary cortical neurons that were knocked out for REST (*Rest*^{Cre}) or kept WT (*Rest*$^{\Delta Cre}$) and co-transduced with either *CaMKIV*^{Scr} or *CaMKIV*^{Sh} expressing lentiviruses. **B.** Mean (± sem with individual experimental points) of the mEPSC amplitude (*left*) and frequency (*right*) in *Rest*$^{\Delta Cre}$ + *CaMKIV*^{Scr} (n=20), *Rest*$^{\Delta Cre}$ + *CaMKIV*^{Sh} (n=15), *Rest*^{Cre} + *CaMKIV*^{Scr} (n=17), *Rest*^{Cre} + *CaMKIV*^{Sh} (n=21) neurons from n=3 independent neuronal preparations. Amplitude: *p=0.0172, *CaMKIV*^{Scr} *vs CaMKIV*^{Sh} with *Rest*$^{\Delta Cre}$; Frequency: **p=0.071, *Rest*$^{\Delta Cre}$ *vs Rest*^{Cre} with *CaMKIV*^{Scr}; *p=0.0126, *CaMKIV*^{Scr} *vs CaMKIV*^{Sh} with *Rest*^{Cre}; 2-way ANOVA/Tukey's tests.

DISCUSSION

Mature neurons can regulate their network activity to maintain a target level over time through homeostatic plasticity mechanisms regulating intrinsic excitability and synaptic transmission. Several distinct homeostatic transcriptional and posttranslational mechanisms have been identified (Michetti & Benfenati, 2024). They are triggered by a proxy of hyperactivity that most studies indicate as the cytosolic Ca^{2+} concentration, representing an integrated activity signal. Calcium increases activate CaM-dependent enzymes, including the array of CaMKs that, in turn, act at the posttranslational and transcriptional levels, modifying function and expression of ion channels and synaptic proteins (Turrigiano et al., 2011). A key role in the activity-regulated transcriptional mechanisms is played by transcription factors that can be directly stimulated by hyperactivity (e.g., cFos and Arc; Qiu et al., 2022) or activated by phosphorylation. Among CaMKs, CaMKIV, once activated by CaMKK, enters the nucleus and phosphorylates a series of transcription factors such as CREB and MeCP2. A prominent transcription factor that responds rapidly and directly to hyperactivity is REST, an epigenetic master regulator that acts by repressing transcription of a large cluster of neuron-specific genes by interacting with RE1 motifs in their promoters (Ballas & Mandel, 2005; Baldelli & Meldolesi, 2015). In addition to the fundamental role of REST downregulation during neuronal differentiation, which allows the full expression of the neuronal phenotype, REST acts as an important homeostatic regulator in mature neurons that preserves them from hyperactivity and senescence. Dysregulation of REST in terms of overexpression or cell mislocalization was found to associate with brain diseases, such as epilepsy, stroke, neuroinflammation, and neurodegeneration (Buckley et al., 2010; Lu et al., 2014; McClelland et al., 2011, 2014; Hall et al., 2024; Carminati et al., 2021; Buffolo et al., 2021; Natali et al., 2023; Carminati et al., 2020; see Hwang & Zukin, 2018 for review). These considerations emphasize the importance of a tight regulation of REST-DNA interactions, nuclear-cytoplasmic shuttling, and cytoplasmic fate.

REST was reported to be phosphorylated by CK1 and ERK1/2 at distinct sites in its C-terminal region within phosphodegron sequences that direct the protein to the proteasome degradation machinery (Westbrook et al., 2008; Guardavaccaro et al., 2008; Kaneko et al., 2014; Nesti et al., 2014). These mechanisms are thought to be important to decrease cell REST levels in neural progenitors to permit neuronal differentiation. As calcium is believed to be the most likely activator of the homeostatic responses mediated by REST, and REST displays several consensus sequences for CaMKs, we have analyzed the possibility that REST is phosphorylated by CaMKs and that phosphorylation modulates its transcriptional activity and/or cellular fate. Indeed, we found that (i) REST is phosphorylated by CaMKIV at a major phosphorylation site between the 5th and 6th ZNF domains (S322); (ii) phosphorylation by CaMKIV inhibits the transcriptional repressor activity of REST by decreasing its affinity for RE1 motifs; (iii) dissociation from DNA favors shuttling of phospho-REST from the nucleus to the cytoplasm where it undergoes protease-mediated degradation; (iv) the homeostatic effects of CaMKIV require the presence of REST.

Structural studies have shown that the consensus sequences for phosphorylation by CaMKs are, in large part, shared by CaMKI, II, and IV and conform to the following minimal motif: Hyd-X-R-X-X-S*/T*-X-X-X-Hyd (Lee et al., 1994; Dale et al., 1995). While the -3 Arg and -5 hydrophobic residue seem important for all kinases, CaMKI prefers a +4 hydrophobic residue, while CaMKII

prefers a +2 acidic residue. The motif for CaMKIV is more degenerate with fewer requirements than the other kinases, supporting the designation of CaMKIV as a multifunctional protein kinase. Of the three putative CaMKIV phosphorylation sites (S177, T294, S322) within the 8 ZNF central cluster region of mouse and human REST, analysis of the location of the basic, hydrophobic and acidic residues in the consensus sequences found that the sequences encompassing T294 and S322 conform better to these requirements, in addition to being the most evolutionary conserved of the three putative sites. Such observation was fully confirmed in the prediction for putative phosphorylation sites by the software Musite and NetPhos3.1, in both the mouse and human REST. Moreover, the REST peptide linkers encompassing the T294 and S322 putative sites share a strong analogy with other DNA-binding transcription factors containing ZNF clusters organized in arrays joined by short 5-7-mer linker sequences. In the 4 ZNF-containing transcription factor Ikaros, for example, ZNF domains are joined by three 5-mer peptides starting with a phosphorylatable residue that, if phosphorylated, impairs the binding of the transcription factor to DNA (Jantz & Berg, 2003). Indeed, MS analysis of in vitro phosphorylated recombinant REST revealed a detectable phosphate incorporation in S322 that thus appears to be the major CaMKIV site in REST.

To explore the mechanistic basis by which phosphorylation modulates REST-DNA binding affinity, we employed atomistic molecular dynamics (MD) simulations of the mREST:RE1 complex in both WT and phosphomimic variants. The simulations revealed that the S322E substitution, mimicking CaMKIV-mediated phosphorylation, induces a notable structural destabilization at the protein-DNA interface. In three independently simulated trajectories over nine, we observed dissociation of the first Zn-finger domain from its DNA binding site, a phenomenon that was totally absent in WT simulations. Furthermore, in several S322E trajectories, destabilization extended beyond the local interface, leading to fragilization of the DNA termini and a collapse of the full complex. Although a comparable loss of interface stability was observed also in some of the T294E simulations, dissociation events occurred less frequently and were not accompanied by complex-wide structural collapse, indicating a more limited effect of the mutation on DNA binding.

These computational findings align with prior experimental observations showing that phosphorylation within or near Zn-finger domains can impair DNA-binding affinity. In particular, phosphorylation of conserved linker sequences in Cys2-His2-type Zn-finger proteins, such as Ikaros and Sp1, has been shown to abolish DNA binding during mitosis by introducing conformational constraints or electrostatic repulsion (Dovat et al., 2002; Jantz & Berg, 2004; Rizkallah et al., 2011). Consistent with this mechanism, our biochemical experiments and MD simulations on mREST mutants demonstrate that a phosphomimic substitution between ZNF5 and ZNF6 within the Zn-finger module directly destabilizes the coordination geometry and its contact with DNA. To our knowledge, this represents the first MD-based evidence of a site-specific post-translational modification within a Zn-finger disrupting the structural stability of a DNA–protein complex at atomic resolution. The simulation results provide structural validation for the biochemical finding that CaMKIV-mediated phosphorylation of REST at Ser322 reduces its affinity for RE1 motifs, facilitating its dissociation from chromatin. Importantly, the absence of collapse or dissociation in WT simulations highlights the specificity of the phosphomimic effect and supports a model in which phosphorylation acts as a trigger for nuclear export and cytosolic degradation. The atomistic-level insight offers a mechanistically distinct layer of control of REST activity directly

acting through the DNA-binding domain, in addition to the previously reported regulation via C-terminal phosphodegrons.

Release of CaMKIV-phosphorylated REST from DNA causes its outflow to the cytoplasm, where it undergoes degradation. There are similarities and differences with respect to previous reports on REST phosphorylation. In fact, both CK1 and ERK1/2 are fundamental signaling pathways to achieve REST downregulation during neural development. Their phosphorylation creates phosphodegrons in the C-terminal region of REST, far from the DNA-interacting ZNF domains, that directly target it to the proteasome (Westbrook et al., 2008; Guardavaccaro et al., 2008; Kaneko et al., 2014; Nesti et al., 2014). Given the location of the CaMKIV phosphorylation site within the ZNF domain, REST degradation is likely the consequence of its dissociation from DNA and cytoplasmic migration, rather than the result of the generation of a phosphodegron recruiting ubiquitin ligases. Our results show that CaMKIV-phosphorylated REST is not targeted to the proteasome. Moreover, REST degradation is resistant to the inhibition of organelle acidification by bafilomycin A1, indicating that it does not depend on low pH active proteases present in lysosomes and autophagosomes. Rather, its degradation is blocked by leupeptin, which mostly inhibits lysosomal proteases, which have recently been shown to be active also in extra-lysosomal cytosolic localizations (Conesa-Bakkali et al., 2025). Thus, the available evidence suggests that REST degradation could be mediated by either extra-lysosomal proteases or lysosomal proteases that do not require acidic pH (Curevo et al., 1997).

The interaction between two recognized prominent homeostatic factors, CaMKIV and REST, that are co-activated in response to intracellular Ca^{2+} fluxes during hyperactivity is physiologically of interest. REST responds very rapidly to hyperactivity with increased expression, migration into the nucleus, and repression of target activity genes (Pozzi et al., 2013; Prestigio et al., 2021). CaMKIV is highly expressed in neurons, where it acts downstream of Ca^{2+} signaling. It is activated by the CaMKK phosphorylation, migrates into the nucleus, and modulates transcription of various genes by phosphorylating transcription factors (Xu et al., 2024). The importance of the CaMKIV/REST interplay in the homeostatic effects of CaMKIV is further underlined by the observation that the CaMKIV-induced changes in excitatory synaptic strength (Joseph & Turrigiano, 2017) are virtually abolished by the genetic deletion of REST. As calcium is believed to be the most likely activator of the homeostatic responses mediated by both REST and CaMKIV, the results suggest that the homeostatic responses of neurons to hyperactivity may require an interplay between CaMKIV and REST, which may define the temporal window of activation of the homeostatic response before the return to basal conditions.

The timing of epigenetic changes and transcriptional effects induced by REST is a key issue, since transcriptional effects can last for long times and thus require a negative regulation of REST activation and overexpression to return to baseline. Termination of REST activity may consist of a sequence of events started by phosphorylation by CaMKIV in the nucleus, dissociation of phospho-REST from the DNA, shuttling to the cytoplasm, followed by degradation. Thus, the elucidated CaMKIV-REST interplay can play an important regulatory role, bringing back the homeostatically increased REST levels to a physiological transcriptional profile after the inhibition of hyperactivity. Such a mechanistic model, shown in **Figure 9**, defines two phases of the homeostatic response to hyperactivity, namely an early phase triggering long-lasting

transcriptional effects to decrease neuronal activity, and a subsequent phase in which the transcriptional activators/repressors are switched off.

Figure 9. Proposed model for the effects of REST phosphorylation by CaMKIV. Calcium entry during hyperactivity activates both REST expression and CaMKIV. REST homeostatically downregulates intrinsic excitability and excitatory synaptic transmission by acting at the transcriptional level in the nucleus. Meanwhile, CaMKIV, activated by Ca^{2+}/calmodulin, translocates to the nucleus and phosphorylates REST at S^{322} adjacent to the 5^{th} Zn-finger domain, which participates in the interaction of the transcription factor with DNA. Phosphorylated REST has a reduced affinity for the RE1 sequences present in the promoter region of its target genes, and, upon unbinding, it is shuttled back to the cytosol, where it is eventually targeted for degradation. This way, the homeostatic plasticity response triggered by hyperactivity will terminate once activity has been normalized and the increased REST levels have returned to baseline.

In conclusion, we found a control mechanism that may contribute to closing the homeostatic transcriptional loop in response to hyperactivity and that involves two well-known transcriptional regulators. A deeper understanding of REST phosphorylation dynamics by CaMKIV could uncover novel regulatory nodes with therapeutic potential in neurological disorders.

Acknowledgements. We are grateful to Drs. Gail Mandel (Vollum Institute, Portland, OR), Thomas Floss (Helmholtz Zentrum München, Deutsches Forschungszentrum für Gesundheit und Umwelt, Neuherberg, Germany), and the German Gene Trap Consortium (GGTC-Partners) for providing us with the REST conditional knockout mice. We are also grateful to Drs Angus Nairn (Yale University School of Medicine, New Haven, MA) and Taku Kaitsuka (Kumamoto University, Kumamoto, Japan) for kindly providing the DNA constructs for WT, catalytically active, and kinase-dead CaMKs. We acknowledge ISCRA for awarding this project access to the LEONARDO supercomputer, owned by the EuroHPC Joint Undertaking, hosted by CINECA (Bologna, Italy). We also gratefully acknowledge the Data Science and Computation Facility and its Support Team for their support and assistance on the IIT High Performance Computing Infrastructure (Genova, Italy). We also thank: Drs. Maria Sabina Cerullo and Alessandra Romei for help in some initial experiments; Dr. Pietro Baldelli for critical reading of the manuscript; drs. Riccardo Navone (Istituto Italiano di Tecnologia, Genova, Italy), Laura Emionite, and Michele Cilli (IRCCS Ospedale Policlinico San Martino, Genova, Italy) for assistance in breeding the mice; Drs. Diego Moruzzo and Arta Mehilli (Istituto Italiano di Tecnologia, Genova, Italy) for help in genotyping; Drs. Ilaria Dallorto and Rossana Ciancio (Istituto Italiano di Tecnologia, Genova, Italy) for administrative assistance.

Funding. The study was supported by research grants from the Italian Ministry of Health (RF-2021-12374404 and PNRR-MR1-2022-12376528), Compagnia di San Paolo Torino (Trapezio #118787 and #124077), the Italian Ministry of University and Research (PRIN 2020-WMSNBL and PRIN 2022-XWEYBN), and by IRCCS Ospedale Policlinico San Martino (Ricerca Corrente) and #NEXTGENERATIONEU (NGEU) National Recovery and Resilience Plan (NRRP), project MNESYS (PE0000006) – A Multiscale integrated approach to the study of the nervous system in health and disease (DN. 1553 11.10.2022 to HTS and AF) and project B83C22004960002 Light4Brain (to FB).

Author Contributions. HTS, AR, DS and EC run molecular biology, biochemistry, viral vector preparation, and immunohistochemical analyses; BC and LM setup the REST-DNA model, run the MD simulations, and analyzed the data; AA, NL and ADF carried out the mass spectrometry analysis; HTS and AF carried out the REST degradation experiments; MC carried out in vitro electrophysiological experiments and analysis; FB, AR and AF conceived the study, supervised the experiments, and analyzed the experimental data; FB wrote the manuscript and supported the research; all authors critically discussed the experimental results and contributed to manuscript writing and revision.

MATERIALS AND METHODS

Materials and experimental animals

Experimental animals. Wild-type C57BL/6J male mice were purchased from Charles River (Calco, Italy). GTinvRest (RestGTi) mice (Nechiporuk et al. 2016), obtained by Gail Mandel (Portland, OR) and the German Gene Trap Consortium (GGTC-Partners), were kept on a C57BL/6J background in homozygosity under conditions of environmental enrichment. Mice were maintained on a 12:12 h light/dark cycle at constant temperature (22 ± 1 °C) and humidity (60 ± 10%), with water and pellet diet *ad libitum.* All the experiments were strictly following the guidelines of the European Community Council (Directive 2014/63/EU of May 15, 2014) and were approved by the Italian Ministry of Health (Authorization #427/2021-PR).

Materials. W7 (N-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide hydrochloride), KN92, KN93, bicuculline, CGP58845, D-APV, and tetrodotoxin were from Tocris (Bristol, UK). Cyclosporin A, cycloheximide, MG132, leupeptin, and bafilomycin A1 were from Merck Life Science (Milan, Italy). All other chemicals, unless otherwise specified, were from Sigma-Aldrich (Milano, Italy)

Primary neuron cultures

Cultures of primary cortical neurons were obtained either from wild-type C57Bl/6 or GTinvRest mice embryos, as previously described (Baldelli et al., 2007; Chiappalone et al., 2009; Valente et al., 2016). These cultures are very homogeneous, contain ~75% excitatory neurons with a recognizable pyramidal shape, ~20-25% glutamic acid decarboxylase (GAD)-positive inhibitory neurons, very low percentages of astrocytes (5-7%), and no microglial cells due to the embryonic nature of the preparation. In brief, cerebral cortices from E17–E18 embryos were dissociated by enzymatic digestion in 0.125% trypsin for 20 min at 37 °C and then triturated with a fire-polished Pasteur pipette. Primary cultures of dissociated cortical neurons were subsequently plated onto poly-L-lysine (0.1 mg/mL, Merck Life Science)-coated 18-mm glass coverslips (6 × 10^4 cells/coverslip for immunofluorescence and electrophysiology experiments) and poly-L-lysine (0.01 mg/mL)-coated 35-mm wells (5 × 10^5 cells/well for molecular biological and biochemical experiments). Neurons were maintained in a culture medium consisting of Neurobasal, supplemented with B27 (1:50 v/v), Glutamax (1% w/v), penicillin-streptomycin (1%) (ThermoFisher Scientific, Monza, Italy), and kept at 37 °C in a 5% CO_2 humidified atmosphere for 14 days before harvesting.

Manipulations of CaM-dependent pathways in primary cortical neurons

Neurons were treated for 24 h prior to harvesting with the following compounds: N-(6-aminohexyl)-5-chloro-1-naphthalenesulfonamide hydrochloride (W7; 20 µM), Cyclosporin A (CsA; 1 µM), N-[2-[N-(4-chlorocinnamyl)-N-methylaminomethyl]phenyl]-N-(2-hydroxyethyl)-4-methoxybenzene sulfonamide (KN93; 10 µM), and 2-[N-(4-methoxybenzenesulfonyl)]amino-N-(4-chlorocinnamyl)-N-methylbenzylamine phosphate (KN92; 10 µM). All drugs were obtained from Tocris. Dimethyl sulfoxide (DMSO; Cat. No. D2650, Sigma-Aldrich, Milan, Italy) was used as a vehicle control according to the experimental conditions. To specifically alter their CaMK activities, HEK293T cells were alternatively transfected with the following plasmids using Lipofectamine 2000 (ThermoFisher Scientific):

pCMV-HA CaMK1 (WT, CA or KD; kind gift of Dr Taku Kaitsuka, Kumamoto University,

Kumamoto, Japan);
pBS-EGFP-MT-CaMK2A (EMBL Database #AJ890283), pBS-EGFP-MT- CaMK2A K42R (CA), and pBS-EGFP-MT- CaMK2A T286D (KD), kind gifts from Dr. Agnes Thalhammer (Istituto Italiano di Tecnologia, Genova, Italy);
pRSV-CaMK4WT (#45062; Addgene, Watertown, MA), pRSV-CaMK4 CA (1-313) (#45063; Addgene, Watertown, MA), and pRSV-CaMK4 KD was mutagenized using pRSV-CaMKIV WT as template.

REST expression in Hek293 cells

HEK293T cells at approximately 70% confluency in 15-cm plates were transfected with 9.2 µg of pCS-myc-mREST per plate (Pozzi et al. 2013). Cells were washed and collected 48 h after transfection in phosphate-buffered saline (PBS). Cell lysates were prepared using the Pierce c-Myc-Tag Magnetic IP/Co-IP Kit (ThermoFisher Scientific) according to the manufacturer's instructions. Briefly, cells were collected by centrifugation at 100xg for 5 min. Pellets were resuspended in 6 mL of Co-IP buffer (ThermoFisher Scientific) and incubated on ice for 10 min before centrifugation at 18,500xg for 15 min at 4 °C, and the supernatant was sonicated at 25% power for 5 s on ice. The protein concentration was determined by the bicinchoninic assay using Pierce™ BCA Protein Assay Kit (ThermoFisher Scientific). The remaining total lysates were concentrated using Amicon Ultra Centrifugal Filter (50 kDa MWCO; Merck Life Science) at 18,500xg for 20 min at 4 °C. Anti-c-Myc magnetic beads were washed three times in Tris-buffered saline containing Tween-20 (TBS-T) (25 mM Tris, 0.15 M NaCl, 0.05% v/v Tween-20).

On-bead REST phosphorylation by CaMKIV

Approximately 200 µg of the concentrated total lysate was incubated with Pierce[TM] anti-c-Myc magnetic beads (ThermoFisher Scientific) that were pre-washed according to the manufacturer's instructions, diluted 1:15 with TBS-T buffer, and incubated at room temperature (RT) for 30 min under rotation. Beads, collected on the magnetic stand and washed in TBS-T three times, were resuspended in alkaline phosphatase buffer (5 mM Tris-Cl pH 8.7, 10 mM NaCl, 1 mM MgCl$_2$, 0.1 mM DTT) supplemented with protease inhibitor cocktail (Merck Life Science) and treated with 25 U of alkaline phosphatase (Merck Life Science) for 30 min at 30 °C. The reaction was stopped by removing the supernatant on the magnetic stand and washing the beads twice with 200 µL of TBS-T buffer supplemented with 10 mM EDTA and protease inhibitors. The beads were subsequently resuspended in *in vitro* phosphorylation assay buffer (50 mM HEPES at pH7.4, 10 mM MgCl$_2$, 1 mM EGTA, 0.4 mM DTT), supplemented with protease and phosphatase inhibitors, 60 µM Mg^{2+}-ATP, and 600 nM calmodulin (Merck Life Science). Either 1 µg of active recombinant CamKIV (Merck Life Science) or vehicle (negative control) was added to the samples, and the mix was incubated for 1 h at RT under rotation. The on-bead kinase assay was stopped by removing the supernatant and washing the beads with an ice-cold assay buffer (10 mM Tris-HCl at pH 7.4, 75 mM NaCl, 20 mM EDTA) supplemented with protease and phosphatase inhibitors. The bound fractions were eluted in 2 x Laemmli buffer (4% SDS, 20% glycerol, 120 mM Tris-HCl pH 6.8, 200 mM DTT).

Protein Digestion and Mass Spectrometry

The protein content resulting from the on-bead phosphorylation assay was reduced with DTT, alkylated with iodoacetamide, then digested overnight with trypsin following a well-established

protocol (Castagnola et al., 2024; Liessi et al., 2021). The proteomic analysis was conducted on a Thermo Exploris 480 orbitrap system coupled with a Dionex Ultimate 3000 nano-LC system. After trapping and desalting, the tryptic peptides were loaded on an Aurora C18 column (75 mm x 250 mm, 1.6 µm particle size) nanocolumn (Ion Opticks, Fitzroy, Australia) and separated using a linear gradient of acetonitrile in water (both added with 0.1% formic acid), from 3% to 41% in 1 h, followed by column cleaning and reconditioning. The flow rate was set to 300 nL/min. Total run time was 1.5 h. Injection volume was set to 1 µL. Peptides were analyzed in positive ESI mode, using a capillary voltage set to 2.0 kV. The RF lens was set to 40%, and the AGC target was set to 300%. Data acquisition was performed in Data Dependent mode (DDA) MS/MS spectra were acquired in HCD mode. All the collected MS/MS spectra were analyzed using Proteome Discoverer software. Cysteine carbamido-methylation was selected as a fixed modification; acetylation of protein N-Term, methionine oxidation, and phosphorylation at serine, threonine, and tyrosine were selected as variable modifications. Positive protein identifications were retained at a 1% false discovery rate (FDR) threshold and with at least two peptides matching the protein sequence. Unless otherwise specified, all reagents were purchased from Sigma Aldrich, and LC-MS solvents from ThermoFisher. All Instruments and software were purchased from ThermoFisher Scientific.

Inhibition of REST degradation pathways
HEK293T cells at approximately 70% confluency in 3.5-cm plates were transfected with 0.5 µg of either pRSV-CaMKIV (full-length, WT) or pRSV-CaMKIV (1-313; constitutively active) using Lipofectamine 2000 (ThermoFisher Scientific) and DMEM (ThermoFisher Scientific), supplemented with FBS (10%; ThermoFisher Scientific) and penicillin-streptomycin (1%; ThermoFisher Scientific), and were kept at 37 °C in a 5% CO_2 humidified atmosphere. One day after transfection, the medium was replaced with DMEM (ThermoFisher Scientific) with cycloheximide (35 µM; Merck Life Science), 5% FBS, and 1% penicillin-streptomycin. Cells were kept at 37 °C in a 5% CO_2 humidified atmosphere. After 3 h, cells were washed three times in DMEM/5% FBS/1% penicillin-streptomycin and incubated in the same medium containing vehicle (0.1 % DMSO; Merck Life Science), MG132 (5 µM; Merck Life Science), Bafilomycin A1 (10-100 nM; Merck Life Science), or Leupeptin (40 µM; Merck Life Science). Following 16 h of incubation at 37 °C in a 5% CO_2 humidified atmosphere, cells were washed three times in PBS and processed as described for Western blotting.

Site-directed mutagenesis of REST phosphorylation sites
Nucleotides encoding Threonine-294 and Serine-322 were mutated to either alanine or glutamic acid using the protocol of DpnI-mediated site-directed mutagenesis by Fisher and Pei (1997). PCR amplification was performed on the template plasmid containing wild-type mREST cDNA in pEZ-Lv229-mREST-IRES2-mCherry constructs with the following protocol: 95 °C for 30 s, 18 cycles of: 95 °C for 30 s, 55 °C for 1 min, 68 °C for 20 min using the pair of primers for each mutated site. The methylated template DNA contained in the PCR products was digested by incubation with 10 U of DpnI for 2 h at 37 °C. A fraction (1/50) of the digested PCR product was used to transform TOP10-competent cells by heat shock at 42 °C for 30 s, incubated in SOC medium for 30 min at 37 °C, and then plated on an LB-agar plate with ampicillin (100 µg/mL). Mini preps from 4 colonies were prepared for full-length sequencing to identify the correct clone with mutated sites.

Western blotting

Primary cortical neurons or HEK293T cells were used to obtain total cell lysates. Cells were collected in Lysis buffer (150 mM NaCl, 1% NP-40, 0.5% sodium deoxycholate, 0.1% SDS, 2 mM EDTA, 50 mM Tris-HCl pH 7.4) enriched with protease and phosphatase inhibitor cocktails (Merck Life Science, Milan, Italy). Following a 10-min incubation on ice, cell lysates were centrifuged at 18,500xg for 15 min at 4 °C. The supernatant was collected and sonicated at 25% power for 5 s on ice. The BCA assay (ThermoFisher Scientific) was used to determine protein concentration. Equivalent protein amounts were subjected to SDS-PAGE on 7.5% polyacrylamide gels and blotted onto nitrocellulose membranes (Whatman, Maidstone, UK). Blotted membranes were blocked for 1 h using 1% BSA (Merck Life Science) in TBS (10 mM Tris, 150 mM NaCl, pH 8.0) together with 0.1% Triton X-100, and incubated overnight at 4 °C with the following primary antibodies: anti-CaMKI (#ab68234 Abcam, Cambridge, UK), anti-CaMKII (#07-1496, Merck Life Science), anti-CaMKIV (#38 610275 BD Biosciences, Milano, Italy), anti-REST (#07-579, Merck Life Science), anti-GAPDH (#2118, Cell Signalling/Euroclone, Milano, Italy), anti-Calnexin (#ADI-SPA-860, Enzo Lifesciences, Farmingdale, NY). Membranes were washed and incubated at RT for 1 h with the following peroxidase-conjugated secondary antibodies: goat anti-mouse IgG (1:10,000; #32430; ThermoFisher Scientific) and goat anti-rabbit IgG (1:5000; #32460; ThermoFisher Scientific). Bands were revealed using the ECL chemiluminescence detection system (ThermoFisher Scientific), acquired on the iBright Imaging System (ThermoFisher Scientific), and the immunoreactivity quantification was performed by densitometric analysis with ImageJ.

Real-time qPCR

Total cellular RNA was extracted with TRIzol (ThermoFisher Scientific), and RNA concentration was quantified using the Nanodrop-1000 spectrophotometer (ThermoFisher Scientific). cDNA was synthesized starting from 0.25 µg RNA with SuperScript IV Reverse Transcriptase kit (ThermoFisher Scientific) according to the manufacturer's instructions. Gene expression was measured by real-time qPCR using SsoAdvanced Universal SYBR Green Supermix (Bio-Rad Laboratories, Milano, Italy) on a C1000 Touch Thermal Cycler (Bio-Rad) on a CFX96 Real-Time System following the manufacturer's protocol, with the following protocol: 5 min at 95 °C; 10 s at 95 °C; 20 s at 60 °C; 10 s at 72 °C for 45 cycles; melting curve (heating ramp from 55 to 95 °C) to check for amplification specificity. The following primers (final concentration 0.25 µM) were used:

REST_F: 5'- GAACCACCTCCCAGTATG -3'
REST_R: 5'- CTTCTGACAATCCTCCATAG -3'
CaMKI_F: 5'- ATCAAGGAAGTCAGGGTTT -3'
CaMKI_R: 5'- GCAGTGAAGAGTGAGAGG -3'
CaMKII_F: 5'- ACTTCCTTCCACCACTTC -3'
CaMKII_R: 5'- TGAGATACAGCATTCCATACA -3'
CaMKIV_F: 5'- CAGTTCATGTTCAGGAGAAT- 3'
CaMKIV_R: 5'- AATGTAGTCAGCCGTTTC -3'
Gapdh_F: 5'- AGGTCGGTGTGAACGGATTTG -3'
Gapdh_R: 5'- TGTAGACCATGTAGTTGAGGTCA -3'
The primers used for the mutagenesis were the following:

mRest_T294E_F: 5'-AGCACGTGCGAACTCACGAAGGAGAACGCCCGTAT-3'
mRest_T294E_R: 5'-ATACGGGCGTTCTCCTTCGTGAGTTCGCACGTGCT-3'
mRest_S322E F: 5'-GACACATGCGGACTCATGAAGGTGAGAAGCCAT-3'
mRest_S322E R: 5'-ATGGCTTCTCACCTTCATGAGTCCGCATGTGTC-3'
Relative gene expression was determined using the 2-ΔΔCT method, normalizing data on the *Gapdh* housekeeping transcript (Vandesompele et al., 2002).

Cell transfection and immunocytochemistry
HEK293T cells on 18-mm coverslips at 70% confluency were transfected with 0.5 µg of pEZ-Lv229-mREST-IRES2-mCherry WT (GeneCopoeia, Rockville MD) or mutagenized constructs therefrom using Lipofectamine 2000 (ThermoFisher Scientific). One day after transfection, cells were fixed in 4% paraformaldehyde in 0.1 M phosphate buffer pH 7.4 (PB) for 20 min at RT. Fixed cells were stained with anti-REST antibody (#MABE1982, clone 12C11) and Hoechst to visualize the nuclear area. Briefly, cells were washed in PBS three times, permeabilized and blocked, and incubated with the primary antibody overnight at 4 °C in 0.1% Triton X-100 and 5% normal goat serum (NGS) in PBS. After four washes in high-salt PBS (20 mM Na_2HPO_4/ 20 mM NaH_2PO_4/ 0.5 M NaCl) for a total of 20 min, neurons were incubated in the same buffer with Alexa-conjugated secondary antibodies (1:500, ThermoFisher Scientific) and counterstained with Hoechst33342 for nuclear detection. Following a wash in high-salt PBS for 20 min, coverslips were washed once in PBS, once in double-distilled water, and mounted using a Moviol mounting medium. Tilescan images of transfected HEK293T cells were acquired using a 60x objective with a Leica SP8 confocal microscope (Leica Microsystem, Wetzlar, Germany). Colocalization of anti-REST immunostaining signal with Hoechst or with mCherry was analyzed in binarized tilescan images using the JACoP plugin for the FIJI distribution of ImageJ. Colocalization was computed between all channels, with special focus on nuclear localization of anti-REST fluorescence. Manders' overlap coefficients were extracted as measures of reciprocal channel colocalization alongside Pearson's correlation coefficient with Costes' randomization (200 iterations) to test for overlap quality and significance.

Luciferase assay
HEK293T cells at 50% confluency in a 6-well plate were co-transfected with RE1X3-pGL3 promoter vector (Paonessa et al., 2016), with mutagenized mREST in pEZ-Lv229-mREST-IRES2-mCherry, or with pRSV-CaMKIV(1-313) (#45063, Addgene, Watertown, MA) and the pRL TKSV40 vector expressing *Renilla* luciferase (Promega, Milano, Italy), using Lipofectamine 2000 (ThermoFisher Scientific). Two days after transfection, cells were washed in PBS twice and collected in 1x Passive Lysis Buffer (PLB) from Dual-Luciferase Assay Kit (Promega), supplemented with protease inhibitor cocktail. The lysates were subjected to one free-thaw cycle and were centrifuged at 100xg at 4 °C for 5 min to remove residual cell debris. The supernatants were collected, diluted 1:5 in 1xPLB, and added to multi-wells containing the luciferase substrate LARII. After a first basal reading of luminescence, a second reading was performed after the addition of Stop & Glo Reagent (Promega). All readings of luminescence were made using a Spark Multimodal Microplate Reader (Tecan, Zurich, Switzerland).

Transduction of primary neurons

Generation of lentiviral vectors. VSV-pseudo-typed third-generation lentiviruses were obtained as previously described (De Palma and Naldini, 2002). Briefly, lentiviruses were produced by transient four-plasmid co-transfection into HEK293T cells using the Ca^{2+} phosphate transfection method. Supernatants were collected, passed through a 0.45-µm filter, and purified by ultracentrifugation (75,000xg for 2 h, at 4 °C). Viral vectors were titrated at concentrations ranging from 1×10^8 to 1×10^9 transducing units (TU)/mL and used at a multiplicity of infection (MOI) of 5-10.

REST RNA interference. For specific knockdown of CaMKs in neurons, short hairpin RNA (shRNA) was cloned by inserting the annealed oligos of shRNA for either CaMKI or CaMKIV into the pLenti-U6-shRNA-PGK-GFP vector between the XhoI and HpaI sites.

The shRNA oligonucleotides used were as follows:

shCaMKIa Sense oligo III: 5'- TGCATTGTAGCCCTGGATGACTTCAAGAGA
GTCATCCAGGGCTACAATGCTTTTTTC - 3'

shCaMKIa Antisense oligo III: 5'- CTCGAGAAAAAAGCATTGTAGCCCTGGATGA
CTCTCTTGAAGTCATCCAGGGCTACAATGCA - 3'

shCaMKIa Sense oligo IV: 5'- TGGATCAAGCACCCCAACATTTTCAAGAGA
AATGTTGGGGTGCTTGATCCTTTTTTC - 3'

shCaMKIa Antisense oligo IV: 5'- CTCGAGAAAAAAGGATCAAGCACCCCAACATT
TCTCTTGAAAATGTTGGGGTGCTTGATCCA - 3'

shCaMKIV Sense oligo II: 5'- TGAGAGAATCTTCTTTATGCATTCAAGAGA
TGCATAAAGAAGATTCTCTCTTTTTTC - 3'

shCaMKIV Antisense oligo II: 5'- CTCGAGAAAAAAGAGAGAATCTTCTTTATGCA
TCTCTTGAATGCATAAAGAAGATTCTCTCA - 3'

shCaMKIV Sense oligo III: 5'- TGGTGTTAAAGAAAACAGTGGTTCAAGAGA
CCACTGTTTTCTTTAACACCTTTTTTC - 3'

shCaMKIV Antisense oligo III: 5'- CTCGAGAAAAAAGGTGTTAAAGAAAACAGTGG
TCTCTTGAACCACTGTTTTCTTTAACACCA - 3'

scramble Sense oligo: 5'- TGAGAGAATCTTCTTTATGCATTCAAGAGA
TTGGGTTGAAGGTGGATCCCTTTTTTC - 3'

scramble Antisense oligo: 5'- CTCGAGAAAAAAGGGATCCACCTTCAACCCAA
TCTCTTGAATGCATAAAGAAGATTCTCTCA - 3

Transduction of primary neurons. For silencing CaMKs in WT primary cortical neurons, neurons were transduced with lentiviral vectors encoding the respective shRNA together with the GFP reporter (see above). For silencing CaMKIV and/or knocking out REST, primary cortical neurons from RestGTi cKO mice were alternatively transduced with sequences containing active or inactive Cre recombinase (Cre and its deletion mutant ΔCre) that were cloned into pGFP-Lenti using the PGK ubiquitous promoter (Kaeser et al., 2011) together with either scrambled or CaMKIV shRNA using the following vector: pLenti-U6-(scramble/shCaMKIV)-PGK-(ΔCRE/CRE-T2A-GFP). For all the experiments, cKO cortical neurons at 7 days in vitro (DIV) were infected at a multiplicity of infection (MOI) of 10. After 24 h, half of the medium was replaced with fresh medium. Transduction efficiency was. The efficiency of infection was estimated to range between 70% and 90% by counting neurons expressing GFP with respect to the total number of DAPI-stained cells. Primary cortical neurons were infected at 7 DIV. After 24 h, half of the medium was replaced with fresh medium. Experiments were performed 7 days after infection.

Structural Modeling of the REST:RE1 Complex

Atomistic Molecular Dynamics simulations of the wild-type REST-RE1 complex. Structures of the WT REST–RE1 assembly were generated as described in the Appendix (Section *Molecular Modeling of Wild-Type REST* and following). The solution builder plug-in of the CHARMM-GUI web server (Lee et al., 2016, 2019) was utilized for setting up the system and for generating input files to be used in all-atom molecular dynamics (MD) simulations. The DNA-protein assembly was inserted into a rectangular box of TIP3 water molecules (Jorgensen et al.,1983) with a minimum solute-to-box-edge distance of 15 Å, applied along all directions. The total system charge was neutralized by adding Na^+ and Cl^- ions reaching a physiological concentration of 0.15 M. The whole system amounts to ∽112000 atoms. Simulations were performed with NAMD 2.14 (Phillips et al., 2020), using the Amber Ff14SB (Maier et al., 2015) force field for the protein and the BSC1 (Ivani et al., 2016) force field for the nucleic acids. Tetragonal periodic boundary conditions (PBC; de Leeuw et al., 1980) were applied. The Particle Mesh Ewald method (Darden et al., 1993) was used to treat electrostatic interactions and to avoid box surface effects with a maximum grid spacing of 1 Å and sixth-order beta splines. A cutoff of 12 Å was employed for Lennard-Jones interactions, which were smoothly switched off starting at 10 Å. Chemical bond distances involving hydrogen atoms were constrained using the SHAKE algorithm (Ryckaert et al., 1977). A time step of 2 fs was utilized. Before entering the production stage, the system was energy minimized and equilibrated in the isobaric-isothermal ensemble (NPT) with P=1 atm and T=310 K. Pressure and temperature were maintained constant using the Nose-Hoover Langevin piston (Feller et al., 1995) and the Langevin thermostat (Leimkuhler et al., 2009), respectively. Afterwards, three independent, 1-µs-long unbiased all-atom MD simulations were performed in the NPT ensemble.

Clustering of wild-type REST-RE1. To generate phosphomimic mutant conformations, we extracted snapshots from the equilibrium simulations of WT REST-RE1. Trajectories were individually subjected to clustering analysis employing TTClust4.10.3 (Tubiana et al., 2018). MD frames were aligned using the backbone of both the protein and the DNA as a reference. Pairwise RMSDs were assembled into a matrix and used for hierarchical clustering with the Ward algorithm (Murtagh & Legendre, 2014) to build the linkage matrix, minimizing the increase in within-cluster variance. Structures of cluster centers were visually inspected using the PyMOL Molecular Graphics System (2002, Version 2.0, Schrödinger, LLC, New York, NY), and, for each replica, the center of the most populated cluster was taken as the representative conformation of that MD run.

Atomistic Molecular Dynamics Simulations of REST phosphomimics. Two distinct phosphomimic mutants were generated by varying the amino acid identity of positions T154 and S192 (T294 and S322 in the experimental system, respectively) into glutamic acid. All simulation procedures and parameters were the same as for the WT system. A total of 18 independent, 1 µs-long, unbiased all-atom MD simulations, 9 per mutated system, were performed in the NPT ensemble.

Patch-clamp electrophysiology

Whole patch-clamp recordings were made from primary cortical neurons as previously described (REF) using a Multiclamp 700B/Digidata 1400A system (Molecular Devices, Sunnyvale, CA). Patch pipettes, prepared from borosilicate glass, were pulled and fire-polished to a final resistance of 4-5 MΩ when filled with a standard internal solution. For all the experiments, neurons were maintained in standard extracellular Tyrode solution containing (in mM) 140 NaCl, 2 $CaCl_2$, 1

MgCl$_2$, 4 KCl, 10 glucose, and 10 HEPES (pH 7.3 with NaOH). For recording miniature excitatory postsynaptic currents (mEPSCs), bicuculline (30 µM), CGP58845 (10 µM), D-APV (50 µM), and tetrodotoxin (TTX; 300 nM) were added to the extracellular solution to block GABA$_A$, GABA$_B$, NMDA receptors and generation and propagation of spontaneous APs. The internal solution (K-gluconate) used for recording mEPSCs in voltage-clamp configuration contained (in mM): 126 K gluconate, 4 NaCl, 1 MgSO$_4$, 0.02 CaCl$_2$, 0.1 BAPTA, 15 glucose, 5 HEPES, 3 ATP and 0.1 GTP (pH 7.3 with KOH). All reagents were from Tocris, otherwise specified. mEPSCs were acquired at a 10 kHz sample frequency and filtered at 1/5 of the acquisition rate with a low-pass Bessel filter. The amplitude and frequency of the miniature excitatory events were calculated using a peak detector function with appropriate threshold amplitudes and areas using MiniAnalysis (Synaptosoft) and Prism (GraphPad Software Inc., La Jolla, CA). All experiments were performed at RT.

Statistical analysis
Data are expressed as means ± SEM with individual experimental replications, as detailed in the figure legends. Normal distribution of data was assessed using the D'Agostino-Pearson's normality test (n>6) or the Shapiro-Wilk test (n<6). The F-test was used to compare the variance between the two sample groups. The two-tailed unpaired Student's t-test or the non-parametric Mann-Whitney's U-test was used to compare two experimental groups based on data distribution. One-way or two-way ANOVA, followed by Tukey's or Dunnett's multiple comparison test, was used to compare more than two normally distributed experimental groups. The Kruskal-Wallis ANOVA was used to compare more than two non-normally distributed experimental groups, followed by Dunn's multiple comparison test. The significance level was preset to p<0.05. Statistical analysis was done using Prism (GraphPad Software, Inc.).

REFERENCES

Ageta-Ishihara N, Takemoto-Kimura S, Nonaka M, Adachi-Morishima A, Suzuki K, Kamijo S, Fujii H, Mano T, Blaeser F, Chatila TA, Mizuno H, Hirano T, Tagawa Y, Okuno H, Bito H (2009) Control of cortical axon elongation by a GABA-driven Ca^{2+}/calmodulin-dependent protein kinase cascade. *J Neurosci* **29**: 13720-13729.

Aoki H, Hara A, Era T, Kunisada T, Yamada Y (2012) Genetic ablation of Rest leads to in vitro-specific derepression of neuronal genes during neurogenesis. *Development* **139**: 667-677.

Baldelli P, Fassio A, Valtorta F, Benfenati F (2007) Lack of synapsin I reduces the readily releasable pool of synaptic vesicles at central inhibitory synapses. *J Neurosci* **27**: 13520-13531.

Baldelli P, Meldolesi J (2015) The transcription repressor REST in adult neurons: physiology, pathology, and diseases. *eNeuro* **2**: ENEURO.0010-15.2015.

Ballas N, Mandel G (2005) The many faces of REST oversee epigenetic programming of neuronal genes. *Curr Opin Neurobiol* **15**: 500-506.

Ballas N, Grunseich C, Lu DD, Speh JC, Mandel G (2005) REST and its corepressors mediate plasticity of neuronal gene chromatin throughout neurogenesis. *Cell* **121**: 645-657.

Bolton JL, Schulmann A, Garcia-Curran MM, Regev L, Chen Y, Kamei N, Shao M, Singh-Taylor A, Jiang S, Noam Y, Molet J, Mortazavi A, Baram TZ (2020) Unexpected transcriptional programs contribute to hippocampal memory deficits and neuronal stunting after early-life adversity. *Cell Rep* **33**: 108511.

Brenner C, Deplus R, Didelot C, Loriot A, Viré E, De Smet C, Gutierrez A, Danovi D, Bernard D, Boon T, Pelicci PG, Amati B, Kouzarides T, de Launoit Y, Di Croce L, Fuks F (2005) Myc represses transcription through recruitment of DNA methyltransferase corepressor. *EMBO J* **24**: 336-346.

Bruce AW, Donaldson IJ, Wood IC, Yerbury SA, Sadowski MI, Chapman M, Göttgens B, Buckley NJ (2004) Genome-wide analysis of repressor element 1 silencing transcription factor/neuron-restrictive silencing factor (REST/NRSF) target genes. *Proc Natl Acad Sci USA* **101**:10458-63.

Buckley NJ, Johnson R, Zuccato C, Bithell A, Cattaneo E (2010) The role of REST in transcriptional and epigenetic dysregulation in Huntington's disease. *Neurobiol Dis* **39**: 28-39.

Buffolo F, Petrosino V, Albini M, Moschetta M, Carlini F, Floss T, Kerlero de Rosbo N, Cesca F, Rocchi A, Uccelli A, Benfenati F (2021) Neuroinflammation induces synaptic scaling through IL-1β-mediated activation of the transcriptional repressor REST/NRSF. *Cell Death Dis* **12**: 180.

Carminati E, Buffolo F, Rocchi A, Michetti C, Cesca F, Benfenati F (2020) Mild Inactivation of RE-1 Silencing Transcription Factor (REST) Reduces susceptibility to kainic acid-induced seizures. *Front Cell Neurosci* **13**: 580.

Castagnola V, Tomati V, Boselli L, Braccia C, Decherchi S, Pompa PP, Pedemonte N, Benfenati F, Armirotti A (2024). Sources of biases in the *in vitro* testing of nanomaterials: the role of the biomolecular corona. *Nanoscale Horiz* **9**: 799-816.

Centonze E, Marte A, Albini M, Rocchi A, Cesca F, Chiacchiaretta M, Floss T, Baldelli P, Ferroni S, Benfenati F, Valente P (2023) Neuron-restrictive silencer factor/repressor element 1-silencing transcription factor (NRSF/REST) controls spatial K^+ buffering in primary cortical astrocytes. *J Neurochem* **165**: 701-721.

Chiappalone M, Casagrande S, Tedesco M, Valtorta F, Baldelli P, Martinoia S, Benfenati F (2009) Opposite changes in glutamatergic and GABAergic transmission underlie the diffuse hyperexcitability of synapsin I-deficient cortical networks. *Cereb Cortex* **19**: 1422-1439.

Conesa-Bakkali R, Morillo-Huesca M, Martínez-Fábregas J (2025) Non-canonical, extralysosomal activities of lysosomal peptidases in physiological and pathological conditions: new clinical opportunities for cancer therapy. *Cells* **14**: 68.

Cuervo AM, Dice JF, Knecht E (1997) A population of rat liver lysosomes responsible for the selective uptake and degradation of cytosolic proteins. *J Biol Chem* **272**: 5606-5615.

Dale S, Wilson WA, Edelman AM, Hardie DG (1995) Similar substrate recognition motifs for mammalian AMP-activated protein kinase, higher plant HMG-CoA reductase kinase-A, yeast SNF1, and mammalian calmodulin-dependent protein kinase I. *FEBS Lett* **361**: 191-195.

Darden T, York D, Pedersen L (1993) Particle mesh Ewald: An N·log(N) method for Ewald sums in large systems. *J Chem Phys* **98**: 10089-10092.

de Leeuw SW, Perram JW, Smith ER (1980) Simulation of electrostatic systems in periodic boundary conditions. I. Lattice sums and dielectric constants. *Proc. R. Soc. Lond A* **373**: 27-56.

Dovat S, Ronni T, Russell D, Ferrini R, Cobb BS, Smale ST (2002) A common mechanism for mitotic inactivation of C2H2 zinc finger DNA-binding domains. *Genes Dev* **16**: 2985-29890.

Feller SE, Zhang Y, Pasto, RW, Brooks BR (1995) Constant pressure molecular dynamics simulation: The Langevin piston method. *J Chem Phys* **103**: 4613-4621.

Fisher CL, Pei GK (1997) Modification of a PCR-based site-directed mutagenesis method. *Biotechniques* **23**: 570-1, 574.

Guardavaccaro D, Frescas D, Dorrello NV, Peschiaroli A, Multani AS, Cardozo T, Lasorella A, Iavarone A, Chang S, Hernando E, Pagano M (2008) Control of chromosome stability by the beta-TrCP-REST-Mad2 axis. *Nature* **452**: 365-369.

Hall AM, Kamei N, Shao M, Mun HS, Chen K, Chen Y, Baram TZ (2024) Inhibition of Neuron-Restrictive Silencing Factor (REST/NRSF) chromatin binding attenuates epileptogenesis. *eNeuro* **11**: ENEURO.0006-24.2024.

Hwang JY, Zukin RS (2018) REST, a master transcriptional regulator in neurodegenerative disease. *Curr Opin Neurobiol* **48**: 193-200.

Ivani I, Dans PD, Noy A, Pérez A, Faustino I, Hospital A, Walther J, Andrio P, Goñi R, Balaceanu A, Portella G, Battistini F, Gelpí JL, González C, Vendruscolo M, Laughton CA, Harris SA, Case DA, Orozco M (2016) Parmbsc1: a refined force field for DNA simulations. *Nat Methods* **13**: 55-58.

Jantz D, Berg JM (2004) Reduction in DNA-binding affinity of Cys2His2 zinc finger proteins by linker phosphorylation. *Proc Natl Acad Sci USA* **101**: 7589-7593.

Jorgensen WL, Chandrasekhar J, Madura JD, Impey RW, Klein ML (1983) Comparison of simple potential functions for simulating liquid water. *J Chem Phys* **79**: 926-935.

Joseph A, Turrigiano GG (2017) All for one but not one for all: excitatory synaptic scaling and intrinsic excitability are coregulated by CaMKIV, whereas inhibitory synaptic scaling is under independent control. *J Neurosci* **37**: 6778-6785.

Kaiser J, Risteska A, Muller AG, Sun H, Lei B, Nay K, Means AR, Cousin MA, Drewry DH, Oakhill JS, Kemp BE, Hannan AJ, Berk M, Febbraio MA, Gundlach AL, Hill-Yardin EL, Scott JW (2025) Convergence on CaMK4: A key modulator of autism-associated signaling pathways in neurons. *Biol Psychiatry* **97**: 439-449.

Kaneko N, Hwang JY, Gertner M, Pontarelli F, Zukin RS (2014) Casein kinase 1 suppresses activation of REST in insulted hippocampal neurons and halts ischemia-induced neuronal death. *J Neurosci* **34**: 6030-6039.

Klug A (2010) The discovery of zinc fingers and their development for practical applications in gene regulation and genome manipulation. *Q Rev Biophys* **43**: 1-21.

Kraner SD, Chong JA, Tsay HJ, Mandel G (1992) Silencing the type II sodium channel gene: a model for neural-specific gene regulation. *Neuron* **9**: 37-44.

Lee JC, Kwon YG, Lawrence DS, Edelman AM (1994) A requirement of hydrophobic and basic amino acid residues for substrate recognition by Ca^{2+}/calmodulin-dependent protein kinase Ia. *Proc Natl Acad Sci USA* **91**: 6413-6417.

Lee J, Cheng X, Swails JM, Yeom MS, Eastman PK, Lemkul JA, Wei S, Buckner J, Jeong JC, Qi Y, Jo S, Pande VS, Case DA, Brooks CL 3rd, MacKerell AD Jr, Klauda JB, Im W (2016) CHARMM-GUI input generator for NAMD, GROMACS, AMBER, OpenMM, and CHARMM/OpenMM simulations using the CHARMM36 additive force field. *J Chem Theory Comput* **12**: 405-413.

Lee J, Patel DS, Ståhle J, Park SJ, Kern NR, Kim S, Lee J, Cheng X, Valvano MA, Holst O, Knirel YA, Qi Y, Jo S, Klauda JB, Widmalm G, Im W (2019) CHARMM-GUI membrane builder for complex biological membrane simulations with glycolipids and lipoglycans. *J Chem Theory Comput* **15**: 775-786.

Leimkuhler B, Noorizadeh E, Theil F (2009) A gentle stochastic thermostat for molecular dynamics. *J Stat Phys* **135**: 261-277.

Lemrow SM, Anderson KA, Joseph JD, Ribar TJ, Noeldner PK, Means AR (2004) Catalytic activity is required for calcium/calmodulin-dependent protein kinase IV to enter the nucleus. *J Biol Chem* **279**: 11664-11671.

Liessi N, Maragliano L, Castagnola V, Bramini M, Benfenati F, Armirotti A (2021) Isobaric labeling proteomics allows a high-throughput investigation of protein corona orientation. *Anal Chem* **93**: 784-791.

Lu T, Aron L, Zullo J, Pan Y, Kim H, Chen Y, Yang TH, Kim HM, Drake D, Liu XS, Bennett DA, Colaiácovo MP, Yankner BA (2014) REST and stress resistance in ageing and Alzheimer's disease. *Nature* **507**: 448-454.

Maier JA, Martinez C, Kasavajhala K, Wickstrom L, Hauser KE, Simmerling C (2015) ff14SB: improving the accuracy of protein side chain and backbone parameters from ff99SB. *J Chem Theory Comput* **11**: 3696-3713.

Mampay M, Sheridan GK (2019) REST: An epigenetic regulator of neuronal stress responses in the young and ageing brain. *Front Neuroendocrinol* **53**: 100744.

Matthews RP, Guthrie CR, Wailes LM, Zhao X, Means AR, McKnight GS (1994) Calcium/calmodulin-dependent protein kinase types II and IV differentially regulate CREB-dependent gene expression. *Mol Cell Biol* **14**: 6107-6116.

McClelland S, Brennan GP, Dubé C, Rajpara S, Iyer S, Richichi C, Bernard C, Baram TZ (2014) The transcription factor NRSF contributes to epileptogenesis by selective repression of a subset of target genes. *Elife* **3**: e01267.

McClelland S, Flynn C, Dubé C, Richichi C, Zha Q, Ghestem A, Esclapez M, Bernard C, Baram TZ (2011) Neuron-restrictive silencer factor-mediated hyperpolarization-activated cyclic nucleotide-gated channelopathy in experimental temporal lobe epilepsy. *Ann Neurol* **70**: 454-464.

Michetti C, Benfenati F (2024) Homeostatic regulation of brain activity: from endogenous mechanisms to homeostatic nanomachines. *Am J Physiol Cell Physiol* **327**: C1384-C1399.

Mori N, Schoenherr C, Vandenbergh DJ, Anderson DJ (1992) A common silencer element in the SCG10 and type II Na$^+$ channel genes binds a factor present in nonneuronal cells but not in neuronal cells. *Neuron* **9**: 45-54.

Murtagh F, Legendre P (2014) Ward's hierarchical agglomerative clustering method: which algorithms implement ward's criterion? *J Classif* **31**: 274-295.

Natali G, Michetti C, Krawczun-Rygmaczewska A, Floss T, Cesca F, Benfenati F (2023) Conditional knockout of REST/NRSF in excitatory neurons reduces seizure susceptibility to chemical kindling. *Front Cell Neurosci* **17**: 1267609.

Nesti E, Corson GM, McCleskey M, Oyer JA, Mandel G (2014) C-terminal domain small phosphatase 1 and MAP kinase reciprocally control REST stability and neuronal differentiation. *Proc Natl Acad Sci USA* **111**: E3929-36.

Palm K, Belluardo N, Metsis M, Timmusk T (1998) Neuronal expression of zinc finger transcription factor REST/NRSF/XBR gene. *J Neurosci* **18**: 1280-1296.

Paonessa F, Criscuolo S, Sacchetti S, Amoroso D, Scarongella H, Pecoraro Bisogni F, Carminati E, Pruzzo G, Maragliano L, Cesca F, Benfenati F (2016) Regulation of neural gene transcription by optogenetic inhibition of the RE1-silencing transcription factor. *Proc Natl Acad Sci USA* **113**: E91-100.

Pecoraro-Bisogni F, Lignani G, Contestabile A, Castroflorio E, Pozzi D, Rocchi A, Prestigio C, Orlando M, Valente P, Massacesi M, Benfenati F, Baldelli P (2018) REST-dependent presynaptic homeostasis induced by chronic neuronal hyperactivity. *Mol Neurobiol* **55**: 4959-4972.

Phillips JC, Hardy DJ, Maia JDC, Stone JE, Ribeiro JV, Bernardi RC, Buch R, Fiorin G, Hénin J, Jiang W, McGreevy R, Melo MCR, Radak BK, Skeel RD, Singharoy A, Wang Y, Roux B, Aksimentiev A, Luthey-Schulten Z, Kalé LV, Schulten K, Chipot C, Tajkhorshid E (2020) Scalable molecular dynamics on CPU and GPU architectures with NAMD. *J Chem Phys* **153**: 044130.

Pozzi D, Lignani G, Ferrea E, Contestabile A, Paonessa F, D'Alessandro R, Lippiello P, Boido D, Fassio A, Meldolesi J, Valtorta F, Benfenati F, Baldelli P (2013) REST/NRSF-mediated intrinsic homeostasis protects neuronal networks from hyperexcitability. *EMBO J* **32**: 2994-3007.

Prestigio C, Ferrante D, Marte A, Romei A, Lignani G, Onofri F, Valente P, Benfenati F, Baldelli P. REST/NRSF drives homeostatic plasticity of inhibitory synapses in a target-dependent fashion. *Elife* **10**: e69058.

Qiu Y, O'Neill N, Maffei B, Zourray C, Almacellas-Barbanoj A, Carpenter JC, Jones SP, Leite M, Turner TJ, Moreira FC, Snowball A, Shekh-Ahmad T, Magloire V, Barral S, Kurian MA, Walker MC, Schorge S, Kullmann DM, Lignani G (2022) On-demand cell-autonomous gene therapy for brain circuit disorders. *Science* **378**: 523-532.

Rizkallah R, Alexander KE, Hurt MM (2011) Global mitotic phosphorylation of C2H2 zinc finger protein linker peptides. *Cell Cycle* **10**: 3327-3336.

Ryckaert J-P, Ciccotti G, Berendsen HJC (1977) Numerical integration of the Cartesian equations of motion of a system with constraints: molecular dynamics of n-alkanes. *J Comput Phys* **23**: 327-341.

Rocchi A, Carminati E, De Fusco A, Kowalska JA, Floss T, Benfenati F (2021) REST/NRSF deficiency impairs autophagy and leads to cellular senescence in neurons. *Aging Cell* **20**: e13471.

Rodenas-Ruano A, Chávez AE, Cossio MJ, Castill PE, Zukin RS (2012) REST-dependent epigenetic remodeling promotes the developmental switch in synaptic NMDA receptors. *Nat Neurosci* **15**: 1382-1390.

Rockowitz S, Lien WH, Pedrosa E, Wei G, Lin M, Zhao K, Lachman HM, Fuchs E, Zheng D (2014) Comparison of REST cistromes across human cell types reveals common and context-specific functions. *PLoS Comput Biol* **10**: e1003671.

Shimojo M, Hersh LB (2006) Characterization of the REST/NRSF-interacting LIM domain protein (RILP): localization and interaction with REST/NRSF. *J Neurochem* **96**: 1130-1138.

Singh-Taylor A, Molet J, Jiang S, Korosi A, Bolton JL, Noam Y, Simeone K, Cope J, Chen Y, Mortazavi A, Baram TZ (2018) NRSF-dependent epigenetic mechanisms contribute to programming of stress-sensitive neurons by neonatal experience, promoting resilience. *Mol Psychiatry* **23**: 648-657.

Tubiana T, Carvaillo JC, Boulard Y, Bressanelli S (2018) TTClust: a versatile molecular simulation trajectory clustering program with graphical summaries. *J Chem Inf Model* **58**: 2178-2182.

Turrigiano GG (2008) The self-tuning neuron: synaptic scaling of excitatory synapses. *Cell* **135**: 422-435.

Turrigiano G (2011) Too many cooks? Intrinsic and synaptic homeostatic mechanisms in cortical circuit refinement. *Annu Rev Neurosci* **34**: 89-103.

Valente P, Castroflorio E, Rossi P, Fadda M, Sterlini B, Cervigni RI, Prestigio C, Giovedì S, Onofri F, Mura E, Guarnieri FC, Marte A, Orlando M, Zara F, Fassio A, Valtorta F, Baldelli P, Corradi A, Benfenati F (2016) PRRT2 is a key component of the Ca^{2+}-dependent neurotransmitter release machinery. *Cell Rep* **15**: 117-131.

Vandesompele J, De Preter K, Pattyn F, Poppe B, Van Roy N, De Paepe A, Speleman F (2002) Accurate normalization of real-time quantitative RT-PCR data by geometric averaging of multiple internal control genes. *Genome Biol* **3**: RESEARCH0034.

Westbrook TF, Hu G, Ang XL, Mulligan P, Pavlova NN, Liang A, Leng Y, Maehr R, Shi Y, Harper JW, Elledge SJ (2008) SCFbeta-TRCP controls oncogenic transformation and neural differentiation through REST degradation. *Nature* **452**: 370-374.

White RR, Kwon YG, Taing M, Lawrence DS, Edelman AM (1998) Definition of optimal substrate recognition motifs of Ca^{2+}-calmodulin-dependent protein kinases IV and II reveals shared and distinctive features. *J Biol Chem* **273**: 3166-3172.

Wolfe SA, Greisman HA, Ramm EI, Pabo CO (1999) Analysis of zinc fingers optimized via phage display: evaluating the utility of a recognition code. *J Mol Biol* **285**: 1917-1934.

Xu H, Yong L, Gao X, Chen Y, Wang Y, Wang F, Hou X (2024) CaMK4: Structure, physiological functions, and therapeutic potential. *Biochem Pharmacol* **224**: 116204.

Yeo M, Berglund K, Augustine G, Liedtke W (2009) Novel repression of Kcc2 transcription by REST-RE-1 controls developmental switch in neuronal chloride. *J Neurosci* **29**: 14652-14662.

Zullo JM, Drake D, Aron L, O'Hern P, Dhamne SC, Davidsohn N, Mao CA, Klein WH, Rotenberg A, Bennett DA, Church GM, Colaiácovo MP, Yankner BA (2019) Regulation of lifespan by neural excitation and REST. *Nature* **574**: 359-364.