1

2 Manuscript type: Article

Quantifying a trait-mediated indirect effect of an environmental stressor on predator-prey dynamics

Last revision: December 11, 2025

Shota Shibasaki[1#] & Masato Yamamichi[1,2,3,4,5,6*]

1. Center for Frontier Research, National Institute of Genetics, 1111 Yata, Mishima,
Shizuoka, 411-8540 Japan

2. Genetics Program, Graduate Institute for Advanced Studies, SOKENDAI (The
Graduate University for Advanced Studies), 1111 Yata, Mishima, Shizuoka 411-
8540, Japan

3. School of the Environment, The University of Queensland, St. Lucia, Brisbane,
Queensland 4072, Australia

4. Department of International Health and Medical Anthropology, Institute of
Tropical Medicine, Nagasaki University, 1-12-4 Sakamoto, Nagasaki 852-8523,
Japan

5. Institute for Multidisciplinary Sciences, Yokohama National University, 79-5
Tokiwadai, Hodogaya, Yokohama, Kanagawa 240-8501, Japan

6. Department of Ecosystem Studies, Graduate School of Agricultural and Life
Sciences, The University of Tokyo, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657,
Japan

[*]To whom correspondence: M.Y. myamamichi@nig.ac.jp

[#]Current affiliation: Faculty of Culture and Information Science, Doshisha University,
Kyotanabe, Kyoto 610-0394, Japan

ORCID: SS 0000-0002-8196-0745, MY 0000-0003-2136-3399

Keywords: *Brachionus calyciflorus*, *Chlamydomonas reinhardtii*, *Chlamydomonas*
sphaeroides, *Chlorella vulgaris*, environmental stressor, Lotka-Volterra model,
plankton, predator-prey dynamics, salinity stress, trait-mediated indirect effect

No. of words in the main text: 4535; No. of words in Abstract: 350; No. of references:
62; No. of figures: 4; No. of tables: 2

Abstract

Trait-mediated indirect interactions have been intensively studied to understand ecological dynamics in biological communities. Yet, little is known about the relative importance of trait-mediated indirect effects of environmental stressors on population dynamics. How does an environmental stressor indirectly affect the population dynamics of a focal species by altering the traits of interacting species and the strength of interspecific interactions? Here, we quantified the direct and trait-mediated indirect effects of an environmental stressor by combining rotifer-alga experiments and Bayesian parameter estimation of a dynamic model. These days, human activities salinize freshwater lakes globally, thereby increasing the death rates and decreasing birth rates of plankton species. Salinity stress is also known to induce cell clumping in certain phytoplankton species. As larger clumps work as a defense trait against gape-limited predation by zooplankton, the salinity stress can affect zooplankton not only directly but also indirectly through phytoplankton trait changes. We first show that a green alga, *Chlamydomonas sphaeroides*, formed larger clumps than the two model species of green algae (*Chlamydomonas reinhardtii* and *Chlorella vulgaris*) under a moderate salinity stress (0.06M NaCl). Then, by tracking the clump size distributions of *Chlamydomonas sphaeroides*, we confirmed that small clumps are more vulnerable to predation by rotifers, *Brachionus calyciflorus*, as previous studies demonstrated. Finally, we co-cultured the green algae and rotifers for a week with and without salinity stress and fitted the Lotka-Volterra predator-prey model. We first estimated how salinity stress increased the rotifer mortality rate using *Chlorella vulgaris*, which seldom showed clump formation. Then, by using *Chlamydomonas sphaeroides*, we estimated how salinity stress increased the mortality rate and decreased the attack rate due to clump formation. We found that salinity stress increased the rotifer mortality rate by more than seventeen-fold, and decreased the attack rate on *Chlamydomonas sphaeroides* to approximately half of that without salinity stress. These results indicate that salinity stress can weaken the predator-prey interaction, and thus salinization can harm freshwater zooplankton species through increasing the mortality rates and decreasing the attack rates. This will be an important step for a quantitative understanding of how environmental stressors can affect community dynamics via trait modifications.

Introduction

Classic studies in ecology have tended to focus on interspecific interactions mediated by population densities, but recent studies have demonstrated the importance of considering trait dynamics driven by behavioral changes, phenotypic plasticity, or rapid evolution (Bolker et al. 2003, Miner et al. 2005, Bolnick et al. 2011, Hendry 2017). Particularly, trait-mediated indirect interactions (TMII) have been intensively studied to understand community dynamics (Werner and Peacor 2003, Schmitz et al. 2004, Utsumi et al. 2010, Ohgushi et al. 2012, Golubski et al. 2016). For example, trophic cascades were traditionally investigated through the perspective of density-mediated indirect interactions: carnivores reduce the population density of herbivores, which increases plant biomass (Hairston et al. 1960, Estes and Palmisano 1974). Studies on TMII, in contrast, emphasized the importance of behavioral change of intermediary species with the foraging-predation risk trade-off: carnivores may change the behavior of herbivores (e.g., "landscape of fear"), which can further increase or even decrease local plant biomass (Schmitz et al. 2004).

On the other hand, little is known about the relative importance of direct and trait-mediated indirect effects of environmental stressors on population dynamics. Human activities are currently affecting various species negatively, and researchers tend to focus on the loss of biodiversity thorough the direct effects of environmental stressors on death or birth rates. But if some species change their traits in response to anthropogenic disturbances, it can alter the demography of other interacting species (i.e., the trait-mediated indirect effects) (Abrams 1995, Werner and Peacor 2003, Peacor et al. 2020). For example, human activities can invoke fear in some species and reduce their foraging behaviors (Smith et al. 2024), which can increase their prey's performance (Leighton et al. 2010, Mehlhoop et al. 2022). While an increasing number of studies proposed the potential importance of trait-mediated indirect effects, it has in general been difficult to quantify the relative contributions of direct and indirect pathways of environmental stressor to focal species' population dynamics.

Here we use a rotifer-alga experimental system (Fig. 1) for quantifying trait-mediated indirect effects. The rotifer-alga system has been used in various studies to understand species coexistence (Rothhaupt 1988), predator-prey population cycles (Fussmann et al. 2000, Verschoor et al. 2004, Scheuerl and Stelzer 2019, Blasius et al. 2020), and eco-evolutionary dynamics (Boraas et al. 1998, Fussmann et al. 2003, Yoshida et al. 2003, Yoshida et al. 2007, Becks et al. 2010, Becks et al. 2012, Hiltunen et al. 2014, Kasada et al. 2014, Declerck et al. 2015, Haafke et al. 2016, Hermann and

99  Becks 2022). Particularly, many studies focused on clump formation as algal defense

100 against rotifer predation (reviewed in Yamamichi 2020). In the freshwater unicellular

101 green alga *Chlamydomonas reinhardtii*, the defense trait is palmelloid clump formation,

102 which is too big to be eaten by gape-limited predators including rotifers (Lurling and

103 Beekman 2006, Sathe and Durand 2016, Réveillon and Becks 2024), and such

104 multicellularity is heritable and can evolve in laboratory experiments (Becks et al. 2010,

105 Herron et al. 2019, Bernardes et al. 2021).

106   Importantly, rapid evolution driven by predation is not a sole cause of clump

107 formation: clump formation in green algae is also induced by phenotypic plasticity

108 triggered by environmental stressors (Cheloni and Slaveykova 2021, de Carpentier et al.

109 2022, Cornwallis et al. 2023) including salinity stress (Khona et al. 2016, Farkas et al.

110 2023). Human activities have salinized freshwater systems globally through, for

111 example, roadway de-icing (Dugan et al. 2017, Kaushal et al. 2021, Cunillera-Montcusí

112 et al. 2022), impeding population growth of freshwater phyto- and zooplankton species

113 (Sarma et al. 2006, Venâncio et al. 2019). Therefore, human activities may not only

114 directly increase the death rates of zooplankton species, but also indirectly harm them

115 by inducing palmelloid clump formation of green algae.

116   Here, to understand trait-mediated indirect effects of salinity stress, we

117 conducted laboratory experiments using four freshwater plankton species: a rotifer

118 (*Brachionus calyciflorus*) as a predator and three green algae (*Chlamydomonas*

119 *sphaeroides*, *Chlamydomonas reinhardtii*, and *Chlorella vulgaris*) as prey.

120 *Chlamydomonas sphaeroides* is not a model organism, but phylogenetically close to

121 *Chlamydomonas reinhardtii* (Yumoto et al. 2013) and was used as a food for *B.*

122 *calyciflorus* in previous experiments (Rothhaupt 1988). We first show that

123 *Chlamydomonas sphaeroides* formed remarkably larger clumps than the other two algal

124 species under a moderate salinity stress (0.06M NaCl) by phenotypic plasticity. Then,

125 we show that large clumps of *Chlamydomonas sphaeroides* were less likely to be

126 consumed by rotifers by comparing the clump size distributions before and after

127 exposure to predation under salinity stress. Finally, we show that the rotifer's attack rate

128 on the green alga became approximately half under salinity stress by Bayesian

129 parameter estimation of the Lotka-Volterra model based on co-culture time-series data

130 with and without salinity stress. These results indicate that salinity-induced clump

131 formation can weaken predator-prey interactions in freshwater ecosystems, highlighting

132 the importance of considering trait-mediated indirect effects of environmental stressors

133 to understand community dynamics in changing environments.

Materials and Methods

Strain Information

The unicellular green algae, *Chlorella vulgaris* (NIES-227), *Chlamydomonas reinhardtii* (NIES-2236), and *Chlamydomonas sphaeroides* (NIES-2242), were obtained from the National Institute for Environmental Studies (NIES), Japan. The green algae were maintained in the C medium (Ichimura 1971) at 23°C with the light cycle L:D = 14:10. The details of the culture conditions of the green algae were identical to our previous study (Shibasaki and Yamamichi 2024).

The facultative parthenogenetic rotifer *B. calyciflorus* (*elevetus*) (*sensu* Otake et al. 2025) was isolated from a water sample in Kanazawa Castle Park, Ishikawa prefecture, Japan, obtained in November 2023 by Dr. Yurie Otake at Kyoto University. The rotifer has been maintained in C medium with *Chlorella vulgaris* as prey under 25°C with the light cycle 14:10. The rotifer culture was transferred into a fresh C medium with a weekly dilution rate ca. 10 % while adding 1 ml culture of *Chlorella vulgaris.*

Before adding the rotifers to experimental flasks, the stock culture of rotifers was filtered twice with a 40 μm cell strainer, left for rotifers to starve for four hours in the dark at 25°C, and then filtered and washed again so that *Chlorella vulgaris* did not contaminate the experimental flasks. This protocol follows a previous study with a similar experimental system (Hermann and Becks 2022).

Comparing the clump sizes

We first compared the clump sizes of the three green algae in the following four conditions with three replicates in each: (i) control (C medium), (ii) with predators (adding rotifers at an initial density of ca. one female individual/ml), (iii) with salt (0.06M NaCl C medium), and (iv) with predators and salt (adding rotifers at initial density ca. one female individual/ml to 0.06M NaCl C medium). We grew the green algae (and rotifers, if applicable) in 50 ml of the media in a 300 ml Erlenmeyer flask for seven days at 25°C with the light cycle L:D = 14:10 (using daylight fluorescent lamps with light intensity 145 ± 5 μmol m^{-2}s^{-1}) while shaking (60 rpm). After cultivation, we collected 1 ml of culture from each flask and added 10 μl of Lugol's solution (20 g of potassium iodide and 10 g of iodine mixed in 200 ml of distilled water) to fix the green algae. We measured the sizes (i.e., the number of cells per clump) of 300 clumps per sample under microscopy.

Evaluating salinity-induced clump formation as a defense trait

Next, we examined whether large clumps of *Chlamydomonas sphaeroides* induced by the salinity stress reduced rotifer predation. We first grew *Chlamydomonas sphaeroides* in 50 ml of 0.06M NaCl C medium for seven days in six 300 ml Erlenmeyer flasks so that the green algae formed clumps. Then, we sampled 5 ml of culture from each flask and fixed it using 50 μl of Lugol's solution. We measured the sizes of 300 clumps in each. 5 ml of rotifer cultures (in 0.06M NaCl C medium) were added to three of the six flasks (i.e., test cases) so that the initial density of rotifers was 0.1 individual/ml, while we added 5 ml of fresh 0.06M NaCl C medium to the other three flasks (controls). The six cultures were grown for six hours under light (daylight fluorescent lamps with light intensity 145 ± 5 μmol m^{-2}s^{-1}) while shaking (60 rpm) at 25°C. We then sampled 5 ml of culture from each flask, fixed them with Lugol's solution, and measured the clump sizes as explained above.

Predator-prey dynamics and parameter estimation

To investigate how salinity stress alters predator-prey dynamics, we grew the rotifers and green algae (*Chlorella vulgaris* and *Chlamydomonas sphaeroides*) with and without salinity stress (C medium with 0.06M NaCl and 0M NaCl, respectively) for seven days. The plankton species were grown in 50 ml of the growth medium in a 300 ml-Erlenmeyer flask at 25°C with the light cycle 14:10 (light intensity145 ± 5 μmol m^{-2}s^{-1}) while shaking (60 rpm). We estimated the population densities of green algae and rotifers daily by sampling 1 ml of culture from each flask. Because it was difficult to estimate the densities of *Chlamydomonas sphaeroides* with clumps using an automated cell counter Countess II FL (Thermo Fisher Scientific Inc.), we removed clumping as follows: we centrifuged the sampled cultures (4°C, 2120 x g, 10 min), removed the supernatant, added 1 ml of C medium to each, and left them for two hours at 25°C under lights. This protocol resolved many clumps because the clump formation in this system was caused by phenotypic plasticity (Khona et al. 2016; see also Fig. S12). After two hours, we fixed the samples by adding 10 μl of Lugol's solution to each. For each sample, we measured the densities of the green algae three times by Countess II FL (using 30 μl culture) and counted female rotifers under microscopy.

To distinguish the direct (i.e., increasing the mortality rate) and indirect (i.e., decreasing attack rate due to the salinity-induced clump formation of prey) impacts of salinity stress on the rotifer's population dynamics, we fitted our experimental data to the following Lotka-Volterra predator-prey model with prey logistic growth and predator linear (Holling type I) functional response:

$$\frac{dX_i}{dt} = X_i\left[r_{ij}\left(1 - \frac{X_i}{K_{ij}}\right) - a_{ij}Y\right],$$

$$\frac{dY}{dt} = Y\left(c_i a_{ij} X_i - m_j\right)$$

209   (1)

where X_i is the scaled density of the green algae i (cells/ml divided by 10^8 for

Chlorella vulgaris and 10^7 for *Chlamydomonas sphaeroides*), Y is the scaled density of

female rotifers (female individuals/ml divided by 10^3), r_{ij} is the intrinsic growth rate of

the green algae i in the absence ($j = 1$) or presence ($j = 2$) of the 0.06M NaCl salinity

stress, K_{ij} is the carrying capacity of the green algae i in the absence ($j = 1$) or presence

($j = 2$) of the 0.06M NaCl salinity stress, a_{ij} is the rotifer's attack rate on green algae i in

the absence ($j = 1$) or presence ($j = 2$) of the 0.06M NaCl salinity stress, c_i is the

conversion efficiency of the green algae i to the rotifer's density, which we assume the

salinity stress would not affect, and m_j is the mortality rate of the rotifers in the absence

($j = 1$) or presence ($j = 2$) of the 0.06M NaCl salinity stress. We used the scaled

densities of the prey and predator so that their maximum densities were about unity,

which facilitated the following Bayesian parameter estimation.

Parameter values were estimated using Bayesian statistics (see Supporting

Information for technical details). First, we estimated posterior distributions of the

intrinsic growth rates (r_{ij}) and the carrying capacities (K_{ij}) of *Chlorella vulgaris* ($i = 1$)

and *Chlamydomonas sphaeroides* ($i = 2$) in the absence ($j = 1$) or presence ($j = 2$) of the

0.06M NaCl salinity stress, respectively, from nine-day mono-culture experiments

(Supporting Information S1.1-S1.2 and Figs. S1-S4). Next, we estimated the attack rate

(a_{1j}), conversion efficiency (c_1), and mortality rate (m_j) of rotifers in the absence ($j = 1$)

or presence ($j = 2$) of the 0.06M NaCl salinity stress based on the seven-day co-culture

experiments with *Chlorella vulgaris* (Supporting Information S1.3 and Figs. S5-S7).

We assumed that the negative impact of salinity on rotifers came through the mortality

rate only (i.e., $a_{11} = a_{12}$) because *Chlorella vulgaris* seldom formed clumps in the

presence of the 0.06M NaCl salinity stress (see Results). Finally, we estimated the

attack rates on *Chlamydomonas sphaeroides* (a_{2j}), conversion efficiency (c_2), and

mortality rate (m_j) of rotifers in the absence ($j = 1$) or presence ($j = 2$) of the 0.06M

NaCl salinity stress (Supporting Information S1.3 and Figs. S8-S10) utilizing the

posterior distributions of m_j inferred from the rotifer-*Chlorella* co-culture experiments

as prior distributions.

Experimental evolution of clumping formation

We also cultivated *Chlamydomonas sphaeroides* under salinity stress for 91 days to see whether larger clumps can evolve or not, because palmelloid clump formation might be adaptive under environmental stress (Tong et al. 2022). Previous studies showed that palmelloid clump formation of *Chlamydomonas reinhardtii* can evolve rapidly in laboratory experiments (Becks et al. 2010). Although clump formation of *Chlamydomonas sphaeroides* is phenotypic plasticity (see Results), we expected that we might be able to observe genetic assimilation, where clumps originally formed in response to salinity stress later become genetically fixed via selection (Pigliucci et al. 2006). Such evolved strains would allow us to investigate whether larger clumps were more unlikely to be consumed by rotifers or not in the absence of salinity stress. We grew the ten lines of the green algae in 20 ml of C medium in 200 ml Erlenmeyer flasks at 25°C with the light cycle 14:10 (light intensity 145 ± 5 µmol m^{-2}s^{-1}) while shaking (120 rpm). Within the ten lines, five lines were grown under salinity stress (0.06M NaCl) while the rest five were without salt (0M NaCl). The initial algal density was approximately 1×10^5 cells/ml. Every week, we collected 200 µl of culture medium from each flask, centrifuged the sampled cultures (4°C, 2120 x g, 10 min), removed the supernatant, and added each sample to 20 ml of fresh C medium with or without NaCl. Note that we accidentally sampled 1000 µl, instead of 200 µl, of cultures and transferred into fresh media on day 21.

We weekly measured algae densities and mean diameter of particles (i.e., either single cells or multi-cellular clumps) using an automatic cell counter analyzer, CASY TT (Innovatis). In addition, we also examined whether *Chlamydomonas sphaeroides* maintained palmelloid clump formation when salinity stress was removed as follows: we collected 5 ml of culture medium from each line under salinity stress, centrifuged the sampled cultures, removed the supernatant, and grew in 2 ml of fresh C medium with and without NaCl, respectively, for six hours. After six hours, we measured the mean diameter of the particles by CASY TT. If the green algae lost phenotypic plasticity, the mean diameter without salinity stress was expected to be as high as with salinity stress. In contrast, the six-hour culture under salinity stress would not affect the mean diameters. Note that a part of the data on the mean diameter is missing on days 42 and 91, because we could not use CASY TT due to its malfunctioning on December 31st, 2024, and February 18th, 2025, respectively.

Statistical analyses

All statistical analyses were performed in R version 4.3.1 (R Core Team 2024). The
effects of salt and rotifers on clump sizes of the green algae were analyzed by the non-
parametric two-way analysis of variance (ANOVA) with the aligned rank
transformation in ARTool library version 0.11.1 (Kay et al. 2025). The linear mixed
model was implemented by lmerTest library version 3.1.3 (Kuznetsova et al. 2017). The
parameter values in the predator-prey model were estimated by RStan library version
2.32.6 (Stan Development Team 2025). In the main text, we reported the mean values
and 95% credible intervals (CI) of the posterior distributions of the fold changes in the
attack rate on *Chlamydomonas sphaeroides* and the rotifer mortality rate.
285

Results

Chlamydomonas sphaeroides generates large clumps under salt stress

We first compared the clump sizes of the three freshwater green algae, *Chlorella*
vulgaris, *Chlamydomonas reinhardtii*, and *Chlamydomonas sphaeroides*, in the
presence of either or both of two stressors: salt and predators (rotifers). Although all
four taxa in this study are freshwater species, they can survive in the C medium
containing 0.06M NaCl, which is about 10% seawater salinity (see Figs. S1-S10 as
examples) (Shibasaki and Yamamichi 2024). Figure 2 shows that the median clump
sizes of the three green algae were ~1.0 in the control cases, confirming that they were
unicellular algae. While *Chlorella vulgaris* remained essentially unicellular in the
presence of salt and rotifers, the two *Chlamydomonas* species increased clump size in
the presence of salt. The median clump size of *Chlamydomonas reinhardtii* increased
from one to two cells due to the salinity stress, but their typical clump sizes were less
than four cells per clump. *Chlamydomonas sphaeroides* increased in median clump size
from one to eight or more cells under salinity stress, and we even observed more than
30 cells in a single clump (see the bottom panels of Fig. 2). Statistical analyses also
suggest that salinity had a large impact on the clump sizes of *Chlamydomonas*
sphaeroides (Table 1). Because the green algae immediately became unicellular again
when the salinity stress was removed (Fig. S12), the palmelloid clump formation in
response to salinity stress seems to be phenotypic plasticity, not evolution.
306

Larger clumps impeded the predation by the rotifers

Next, we examined whether large clumps of *Chlamydomonas sphaeroides* under
salinity stress could work as a defense trait against gape-limited predation by rotifers.
We addressed this question by comparing the clump size distribution before and after

exposure to rotifer predation for six hours. Figure 3 shows how the distributions of

clump sizes changed with (Fig. 3A) and without (Fig. 3B) rotifers. In the absence of

rotifers, the median clump sizes decreased or were little changed in six hours (Fig. 3B).

On the other hand, the median clump sizes clearly increased in the presence of rotifers

(Fig. 3A). The linear mixed model suggested that the clump sizes, on average,

decreased in six hours but increased if the green algae co-cultured with the rotifers

(Table 2). The violin plots show that the proportion of small clumps (four or fewer cells

per clump) decreased with predators. Because the doubling time of *Chlamydomonas*

sphaeroides under salinity was estimated to be $\log2/r_{22} \approx 1.33$ days (see Figs. S3-S4),

which is longer than six hours, the presence of rotifers would not induce the formation

of larger palmelloid clump in this experiment. Thus, the rotifer predation would

decrease the proportion of small clumps, but not large ones.

We then examined how salinity stress changed predator-prey dynamics by

conducting co-culture experiments of the green algae and rotifers for seven days. While

the population dynamics of the green algae *Chlamydomonas sphaeroides* did not

seemingly change as a function of salinity stress (Fig. 4A), we observed a remarkable

decrease in rotifer's population growth in the presence of salinity stress (Fig. 4B). To

disentangle the relative importance of direct and trait-mediated indirect impacts of

salinity stress, we fitted the experimental data to the Lotka-Volterra predator-prey

model (Eq. 1) by performing the Bayesian estimation (see Supporting Information and

Figs. S1-S10 for more details). Fig. 4C shows that the mortality rate of the rotifers

under salinity stress (m_2) was more than 17 times (95% CI [6.07, 58.27]) larger than that

without salinity stress (m_1). The attack rate on *Chlamydomonas sphaeroides* under

salinity stress (a_{22}) was, on the other hand, about half (95% CI [0.18, 0.98]) of that in

the absence of salinity stress (a_{21}). Owing to the low rotifer density, the posterior

distribution of a_{22} remained close to its prior and exhibited substantial uncertainty,

whereas the posterior distribution of a_{21} shifted markedly from its prior (Fig. S10).

These results indicate that the attack rates on *Chlamydomonas sphaeroides* are affected

by salinity. Taken together, we successfully quantified the direct and trait-mediated

indirect impacts of salinity stress on the rotifer population dynamics, and the direct

impact was more severe.

The salinity stress did not lead to the evolution of larger clumps or loss of

phenotypic plasticity

We also examined whether the clump size evolved or not when *Chlamydomonas*

sphaeroides were grown under salinity for 91 days. Fig. S11 shows that, while all five

lines without salinity stress (i.e., controls; orange in Fig. S11A) grew constantly during

the experiment, the lines under salinity stress (blue in Fig. S11A) gradually decreased in

growth rate from day 7 with some variation: one line (Salt-1) maintained a moderate

population density (approximately $1\text{-}2\times10^5$ cells/ml) during the experiment, whereas

another (Salt-4) continued to decrease its density from day 70 and went extinct (its

density was <100 cells/ml) at the end of the experiment. Fig. S11B shows the dynamics

of mean diameters of the particles (single cells and multicellular clumps) during the

experiment. Without salinity stress, the mean diameter remained around 10 μm,

suggesting that the green algae were unicellular. On the other hand, the mean diameter

under salinity stress was about 20 μm. This indicates that, if a single cell size did not

change, a single clump would be composed of $2^3 = 8$ cells on average, which is

consistent with our observation under microscopy (Figs. 1-2). However, there was no

evidence that clump size evolved to be larger under salinity stress.

We also investigated whether *Chlamydomonas sphaeroides* maintained clump

formation without salt during the experiment by weekly transferring the green algae into

fresh C media with and without NaCl for six hours. Fig. S12 shows the mean diameter

returned to approximately 10 μm without salinity stress, while the mean diameter

remained 20 μm with the 0.06M NaCl salinity stress. These results indicate that the

green algae returned to single cells when the salinity stress was removed, and that

phenotypic plasticity was maintained during the 91-day experiment.

Discussion

While human activities can directly threaten organisms by harvesting or destroying their

habitats, these impacts can also propagate through communities by altering species

interactions: i.e., trait-mediated indirect effects (Abrams 1995, Werner and Peacor

2003). Despite its potential importance, the relative impacts of direct and indirect effects

on population dynamics were unclear. Here, we used a rotifer-alga system to quantify a

trait-mediated indirect effect in freshwater communities. We found that the 0.06M NaCl

salinity stress induced the algal clump formation (Fig. 2), and it weakened the

interspecific interaction (predation) between the green algae and rotifers (Figs. 3-4). In

freshwater systems, salinization is a severe problem caused by human activities (Dugan

et al. 2017, Kaushal et al. 2021, Cunillera-Montcusí et al. 2022). Our results

demonstrate the importance of a quantitative understanding of how environmental

stressors can affect community dynamics via trait modifications.

Previous studies investigated the clump formation of *Chlamydomonas*
reinhardtii as a defense trait against predation (Lurling and Beekman 2006, Becks et al.
2010, Becks et al. 2012, Sathe and Durand 2016, Bernardes et al. 2021). However,
green algae form multicellular clumps under other kinds of stress (Cheloni and
Slaveykova 2021, de Carpentier et al. 2022, Cornwallis et al. 2023) including salinity
stress (Khona et al. 2016, Farkas et al. 2023). Our study shows that *Chlamydomonas*
sphaeroides was sensitive to the salinity stress and formed larger clumps than two other
model green algae, *Chlorella vulgaris* and *Chlamydomonas reinhardtii*, under 0.06M
NaCl (Fig. 2). Salinity-induced clump formation was caused by phenotypic plasticity
because the green algae immediately lost the clumps and became unicellular when the
salinity was removed from the media (Fig. S12). The green algae formed clumps
without predators (Figs. 2 and S11B), and thus their effect on the predator-prey
interaction seems a by-product of a response to salinity. Although Tong et al. (2022)
argued that the salinity-induced clump formation can be adaptive under stressful
environments, our evolution experiment did not support this claim. Nevertheless, the
salinity-induced clump formation worked as a defense trait against predation by rotifers
as shown in previous studies (Becks et al. 2010, Réveillon and Becks 2024): our
experiments showed that the rotifers were unlikely to consume large clumps (Fig. 3)
and their attack rate on *Chlamydomonas sphaeroides* decreased under salinity (Fig. 4D).
Our results, therefore, indicate that salinity stress weakened the predator-prey
interaction by inducing the algae clump formation.
We found that the trait-mediated indirect effect of salinity stress (decreasing
the attack rate on *Chlamydomonas sphaeroides* a_{2j} by half) was less intense than the
direct effect (increasing the rotifer's mortality rate m_j by 17-fold) (Fig. 4). However,
when we extrapolate the results, we found that it can have a significant effect: in the
Lotka-Volterra predator-prey model (Eq. 1), the predators cannot persist if $c_i a_{ij} K_{ij} < m_j$.
Hence, salinity stress can promote rotifer extinction by reducing a_{ij} through algal clump
formation as well as increasing m_j and decreasing K_{ij}. This is in stark contrast to the idea
of indirect evolutionary rescue, where prey defense evolution prevents predator
extinction (Yamamichi and Miner 2015, Cortez and Yamamichi 2019, Hermann and
Becks 2022). Hermann and Becks (2022) showed that increasing salinity reduces the
rotifer density through the direct effect, and the reduced predation pressure selects for
less defended *Chlamydomonas reinhardtii* due to the growth-defense trade-off (Becks et
al. 2010, Réveillon and Becks 2024). Because of the reduced frequency of defended
algae, in turn, rotifer populations may be rescued from extinction. In our system, clump
formation is induced by salinity stress, suggesting that rotifers may not be rescued from

extinction. It will be interesting to integrate these two opposite consequences of

increasing salinity for better understanding how clump formation under salinity stress

can pleiotropically affect salt tolerance and predator-prey dynamics.

Expanding our experimental system may provide deeper insights into how

human salinization activities disturb freshwater communities. For example, we will be

able to combine two or more stressors (e.g., pH, temperature) to experimentally show

how realistic environmental changes alter ecological dynamics (Kawaguchi and

Yamamichi 2025). Although we focused on interactions between two species, we can

introduce another green alga that may be less sensitive to salinity stress (e.g., *Chlorella*

vulgaris) to the rotifer-*Chlamydomonas* system to investigate how salt changes the

outcome of exploitative and apparent competition between the two algae species.

Another possibility is introducing a zooplankton species that consumes (e.g.,

Asplanchna brightwelli) (Verschoor et al. 2004) or competitor of *B. calyciflorus* (e.g.,

B. rubens) (Rothhaupt 1988). Such expansion would allow us to investigate how

salinization would change the stability and structure of complex food webs. Shibasaki

and Terui (2024) recently showed that species richness non-linearly alters food chain

length. Because salinity stress would reduce the number of freshwater predator species

by increasing mortality rates and causing the trait-mediated indirect effects, it may be

possible to observe the nonlinear relationship between species richness and food chain

length by manipulating salinity in our freshwater systems in future studies.

In conclusion, the present study demonstrates that the rotifer-alga system is

useful for investigating trait-mediated interaction effects caused by environmental

stressors. We focused on the salinization of freshwater systems and analyzed how

salinity stress changes the predator-prey interaction between green algae

Chlamydomonas sphaeroides and rotifers *B. calyciflorus*. In addition to the direct effect

of salinity stress increasing rotifer mortality rate, salinity stress induced the

multicellular clump formation of the green algae, which worked as a by-product defense

trait against gape-limited predation by rotifers. Therefore, the salinity impeded the

growth of rotifers both directly and indirectly. Future studies will be able to investigate

how freshwater salinization alters food webs by extending this experimental system.

Acknowledgment

The three strains of the green algae *Chlorella vulgaris* (NIES-227), *Chlamydomonas*

reinhardtii (NIES-2236), and *Chlamydomonas sphaeroides* (NIES-2242), were obtained

from the National Institute for Environmental Studies (NIES), Japan. We thank Yurie

Otake for sharing the water sample from which we isolated *Brachionus calyciflorus*. We
also appreciate Mayumi Aono for helping with the experiments as a technician. This
study was supported by the Japan Society for the Promotion of Science (JSPS) Grant-in-
Aid for Scientific Research (KAKENHI) JP22H02688, JP22H04983, and JP25K02340,
Japan Science and Technology Agency (JST) CREST JPMJCR23N5, Inamori Research
Grant 2024, and Research Organization for Information and Systems (ROIS) Research
Grant of Strategic Research Project 2024-SRP-10 to M. Y.
459

460 Author contribution

461 S.S. and M. Y. conceived the project, S.S. performed the experiments, conducted the
462 statistical analyses, and wrote the initial draft. M.Y. supervised the project and acquired
463 the funding. Both authors reviewed and edited the manuscript.
464

465 Conflict of interest

466 The authors declare no conflict of interest.
467

468 Data availability

469 The experimental codes and programming codes are available from the following
470 Zenodo repository: https://doi.org/10.5281/zenodo.17875555.
471

472 Reference

Abrams, P. A. 1995. Implications of dynamically variable traits for identifying,
classifying, and measuring direct and indirect effects in ecological communities.
The American Naturalist **146**:112-134.
Becks, L., S. P. Ellner, L. E. Jones, and N. G. Hairston, Jr. 2010. Reduction of adaptive
genetic diversity radically alters eco-evolutionary community dynamics.
Ecology Letters **13**:989-997.
Becks, L., S. P. Ellner, L. E. Jones, and N. G. Hairston, Jr. 2012. The functional
genomics of an eco-evolutionary feedback loop: linking gene expression, trait
evolution, and community dynamics. Ecology Letters **15**:492-501.

Bernardes, J. P., U. John, N. Woltermann, M. Valiadi, R. J. Hermann, and L. Becks.
2021. The evolution of convex trade-offs enables the transition towards
multicellularity. Nature Communications **12**:4222.

Blasius, B., L. Rudolf, G. Weithoff, U. Gaedke, and G. F. Fussmann. 2020. Long-term
cyclic persistence in an experimental predator–prey system. Nature **577**:226-
230.

Bolker, B., M. Holyoak, V. Křivan, L. Rowe, and O. Schmitz. 2003. Connecting
theoretical and empirical studies of trait-mediated interactions. Ecology
84:1101-1114.

Bolnick, D. I., P. Amarasekare, M. S. Araújo, R. Bürger, J. M. Levine, M. Novak, V. H.
W. Rudolf, S. J. Schreiber, M. C. Urban, and D. A. Vasseur. 2011. Why
intraspecific trait variation matters in community ecology. Trends in Ecology &
Evolution **26**:183-192.

Boraas, M. E., D. B. Seale, and J. E. Boxhorn. 1998. Phagotrophy by a flagellate selects
for colonial prey: A possible origin of multicellularity. Evolutionary Ecology
12:153-164.

Cheloni, G., and V. I. Slaveykova. 2021. Morphological plasticity in *Chlamydomonas*
reinhardtii and acclimation to micropollutant stress. Aquatic Toxicology
231:105711.

Cornwallis, C. K., M. Svensson-Coelho, M. Lindh, Q. Li, F. Stábile, L.-A. Hansson,
and K. Rengefors. 2023. Single-cell adaptations shape evolutionary transitions
to multicellularity in green algae. Nature Ecology & Evolution **7**:889-902.

Cortez, M. H., and M. Yamamichi. 2019. How (co)evolution alters predator responses
to increased mortality: extinction thresholds and hydra effects. Ecology
100:e02789.

Cunillera-Montcusí, D., M. Beklioğlu, M. Cañedo-Argüelles, E. Jeppesen, R. Ptacnik,
C. A. Amorim, S. E. Arnott, S. A. Berger, S. Brucet, H. A. Dugan, M. Gerhard,
Z. Horváth, S. Langenheder, J. C. Nejstgaard, M. Reinikainen, M. Striebel, P.
Urrutia-Cordero, C. F. Vad, E. Zadereev, and M. Matias. 2022. Freshwater
salinisation: a research agenda for a saltier world. Trends in Ecology &
Evolution **37**:440-453.

de Carpentier, F., A. Maes, C. H. Marchand, C. Chung, C. Durand, P. Crozet, S. D.
Lemaire, and A. Danon. 2022. How abiotic stress-induced socialization leads to
the formation of massive aggregates in *Chlamydomonas*. Plant Physiology
190:1927-1940.

Declerck, S. A. J., A. R. Malo, S. Diehl, D. Waasdorp, K. D. Lemmen, K. Proios, and S.
Papakostas. 2015. Rapid adaptation of herbivore consumers to nutrient
limitation: eco-evolutionary feedbacks to population demography and resource
control. Ecology Letters **18**:553-562.

Dugan, H. A., S. L. Bartlett, S. M. Burke, J. P. Doubek, F. E. Krivak-Tetley, N. K.
Skaff, J. C. Summers, K. J. Farrell, I. M. McCullough, A. M. Morales-Williams,
D. C. Roberts, Z. Ouyang, F. Scordo, P. C. Hanson, and K. C. Weathers. 2017.
Salting our freshwater lakes. Proceedings of the National Academy of Sciences
of the United States of America **114**:4453-4458.

Estes, J. A., and J. F. Palmisano. 1974. Sea otters: their role in structuring nearshore
communities. science **185**:1058-1060.

Farkas, A., B. Pap, O. Zsíros, R. Patai, P. Shetty, G. Garab, T. Bíró, V. Ördög, and G.
Maróti. 2023. Salinity stress provokes diverse physiological responses of
eukaryotic unicellular microalgae. Algal Research **73**:103155.

Fussmann, G. F., S. P. Ellner, and N. G. Hairston, Jr. 2003. Evolution as a critical
component of plankton dynamics. Proceedings of the Royal Society B:
Biological Sciences **270**:1015-1022.

Fussmann, G. F., S. P. Ellner, K. W. Shertzer, and N. G. Hairston, Jr. 2000. Crossing
the Hopf bifurcation in a live predator-prey system. science **290**:1358-1360.

Golubski, A. J., E. E. Westlund, J. Vandermeer, and M. Pascual. 2016. Ecological
networks over the edge: hypergraph trait-mediated indirect interaction (TMII)
structure. Trends in Ecology & Evolution **31**:344-354.

Haafke, J., M. Abou Chakra, and L. Becks. 2016. Eco-evolutionary feedback promotes
Red Queen dynamics and selects for sex in predator populations. Evolution
70:641-652.

Hairston, N. G., Sr., F. E. Smith, and L. B. Slobodkin. 1960. Community structure,
population control, and competition. The American Naturalist **94**:421-425.

Hendry, A. P. 2017. Eco-evolutionary Dynamics. Princeton University Press, Princeton,
NJ.

Hermann, R. J., and L. Becks. 2022. Change in prey genotype frequency rescues
predator from extinction. Royal Society open science **9**:220211.

Herron, M. D., J. M. Borin, J. C. Boswell, J. Walker, I.-C. K. Chen, C. A. Knox, M.
Boyd, F. Rosenzweig, and W. C. Ratcliff. 2019. *De novo* origins of
multicellularity in response to predation. Scientific Reports **9**:2328.

Hiltunen, T., S. P. Ellner, G. Hooker, L. E. Jones, and N. G. Hairston Jr. 2014. Eco-
evolutionary dynamics in a three-species food web with intraguild predation:
intriguingly complex. Advances in Ecological Research **50**:41-73.

Ichimura, T. 1971. Sexual cell division and conjugation-papilla formation in sexual
reproduction of *Closterium strigosum*. Pages 208-214 *in* Proceedings of the 7th
International Seaweed Symposium. The University of Tokyo Press, Tokyo.

Kasada, M., M. Yamamichi, and T. Yoshida. 2014. Form of an evolutionary tradeoff
affects eco-evolutionary dynamics in a predator-prey system. Proceedings of the
National Academy of Sciences of the United States of America **111**:16035-
16040.

Kaushal, S. S., G. E. Likens, M. L. Pace, J. E. Reimer, C. M. Maas, J. G. Galella, R. M.
Utz, S. Duan, J. R. Kryger, A. M. Yaculak, W. L. Boger, N. W. Bailey, S. Haq,
K. L. Wood, B. M. Wessel, C. E. Park, D. C. Collison, B. Y. a. I. Aisin, T. M.
Gedeon, S. K. Chaudhary, J. Widmer, C. R. Blackwood, C. M. Bolster, M. L.
Devilbiss, D. L. Garrison, S. Halevi, G. Q. Kese, E. K. Quach, C. M. P. Rogelio,
M. L. Tan, H. J. S. Wald, and S. A. Woglo. 2021. Freshwater salinization
syndrome: from emerging global problem to managing risks. Biogeochemistry
154:255-292.

Kawaguchi, Y. W., and M. Yamamichi. 2025. A single episode of sexual reproduction
can produce large variation in population growth rates under dual stressors.
Journal of Evolutionary Biology **38**:778-785.

Kay, M., L. A. Elkin, J. J. Higgins, and J. O. Wobbrock. 2025. ARTool: Aligned Rank
Transform for Nonparametric Factorial ANOVAs. R package version 0.11.2.

Khona, D. K., S. M. Shirolikar, K. K. Gawde, E. Hom, M. A. Deodhar, and J. S.
D'Souza. 2016. Characterization of salt stress-induced palmelloids in the green
alga, *Chlamydomonas reinhardtii*. Algal Research **16**:434-448.

Kuznetsova, A., P. B. Brockhoff, and R. H. B. Christensen. 2017. lmerTest package:
tests in linear mixed effects models. Journal of statistical software **82**:1-26.

Leighton, P. A., J. A. Horrocks, and D. L. Kramer. 2010. Conservation and the
scarecrow effect: Can human activity benefit threatened species by displacing
predators? Biological Conservation **143**:2156-2163.

Lurling, M., and W. Beekman. 2006. Palmelloids formation in *Chlamydomonas*
reinhardtii: defence against rotifer predators? International Journal of
Limnology **42**:65-72.

Mehlhoop, A. C., B. Van Moorter, C. M. Rolandsen, D. Hagen, A. Granhus, R. Eriksen,
T. H. Ringsby, and E. J. Solberg. 2022. Moose in our neighborhood: Does

perceived hunting risk have cascading effects on tree performance in vicinity of

roads and houses? Ecology and Evolution **12**:e8795.

Miner, B. G., S. E. Sultan, S. G. Morgan, D. K. Padilla, and R. A. Relyea. 2005.

Ecological consequences of phenotypic plasticity. Trends in Ecology &

Evolution **20**:685-692.

Ohgushi, T., O. Schmitz, and R. D. Holt. 2012. Trait-Mediated Indirect Interactions:

Ecological and Evolutionary Perspectives. Cambridge University Press,

Cambridge, UK.

Otake, Y., M. Kurokawa, N. Maruoka, M. Nakagawa, W. Makino, and J. Urabe. 2025.

Taxonomic re-evaluation of the *Brachionus calyciflorus* species complex

(Rotifera) in Japanese freshwaters. Limnology **26**:153-167.

Peacor, S. D., B. T. Barton, D. L. Kimbro, A. Sih, and M. J. Sheriff. 2020. A framework

and standardized terminology to facilitate the study of predation-risk effects.

Ecology **101**:e03152.

Pigliucci, M., C. J. Murren, and C. D. Schlichting. 2006. Phenotypic plasticity and

evolution by genetic assimilation. Journal of Experimental Biology **209**:2362-

2367.

R Core Team. 2024. R: A language and environment for statistical computing. R

Foundation for Statistical Computing, Vienna, Austria.

Réveillon, T., and L. Becks. 2024. Trade-offs between defense and competitive traits in

a planktonic predator–prey system. Ecology **105**:e4456.

Rothhaupt, K. O. 1988. Mechanistic resource competition theory applied to laboratory

experiments with zooplankton. Nature **333**:660-662.

Sarma, S. S. S., S. Nandini, J. Morales-Ventura, I. Delgado-Martínez, and L. González-

Valverde. 2006. Effects of NaCl salinity on the population dynamics of

freshwater zooplankton (rotifers and cladocerans). Aquatic Ecology **40**:349-360.

Sathe, S., and P. M. Durand. 2016. Cellular aggregation in *Chlamydomonas*

(Chlorophyceae) is chimaeric and depends on traits like cell size and motility.

European Journal of Phycology **51**:129-138.

Scheuerl, T., and C. P. Stelzer. 2019. Asexual reproduction changes predator population

dynamics in a life predator–prey system. Population Ecology **61**:210-216.

Schmitz, O. J., V. Krivan, and O. Ovadia. 2004. Trophic cascades: the primacy of trait-

mediated indirect interactions. Ecology Letters **7**:153-163.

Shibasaki, S., and A. Terui. 2024. Food web complexity modulates environmental

impacts on food chain length. Oikos **2024**:e10331.

Shibasaki, S., and M. Yamamichi. 2024. The double-edged effect of environmental
fluctuations on evolutionary rescue. BioRxiv.
Smith, J. A., M. E. McDaniels, S. D. Peacor, E. C. Bolas, M. J. Cherry, N. J. Dorn, O.
K. Feldman, D. L. Kimbro, E. K. Leonhardt, N. E. Peckham, M. J. Sheriff, and
K. M. Gaynor. 2024. Population and community consequences of perceived risk
from humans in wildlife. Ecology Letters **27**:e14456.
Stan Development Team. 2025. RStan: the R interface to Stan.
Tong, K., G. O. Bozdag, and W. C. Ratcliff. 2022. Selective drivers of simple
multicellularity. Current Opinion in Microbiology **67**:102141.
Utsumi, S., O. Kishida, and T. Ohgushi. 2010. Trait-mediated indirect interactions in
ecological communities. Population Ecology **52**:457-459.
Venâncio, C., B. B. Castro, R. Ribeiro, S. C. Antunes, N. Abrantes, A. M. V. M. Soares,
and I. Lopes. 2019. Sensitivity of freshwater species under single and
multigenerational exposure to seawater intrusion. Philosophical Transactions of
the Royal Society B: Biological Sciences **374**:20180252.
Verschoor, A. M., M. Vos, and I. van der Stap. 2004. Inducible defences prevent strong
population fluctuations in bi- and tritrophic food chains. Ecology Letters **7**:1143-
1148.
Werner, E. E., and S. D. Peacor. 2003. A review of trait-mediated indirect interactions
in ecological communities. Ecology **84**:1083-1100.
Yamamichi, M. 2020. Effects of Rapid Evolution on Population Cycles and Extinction
in Predator–Prey Systems. Pages 19-49 *in* A. Mougi, editor. Diversity of
Functional Traits and Interactions. Springer, Singapore.
Yamamichi, M., and B. E. Miner. 2015. Indirect evolutionary rescue: prey adapts,
predator avoids extinction. Evolutionary Applications **8**:787-795.
Yoshida, T., S. P. Ellner, L. E. Jones, B. J. M. Bohannan, R. E. Lenski, and N. G.
Hairston, Jr. 2007. Cryptic population dynamics: Rapid evolution masks trophic
interactions. PLOS Biology **5**:1868-1879.
Yoshida, T., L. E. Jones, S. P. Ellner, G. F. Fussmann, and N. G. Hairston, Jr. 2003.
Rapid evolution drives ecological dynamics in a predator-prey system. Nature
424:303-306.
Yumoto, K., F. Kasai, and M. Kawachi. 2013. Taxonomic re-examination of
Chlamydomonas strains maintained in the NIES-Collection. Microbiology and
Culture Collections **29**:1-12.
656
657

Tables

Table 1. Non-parametric two-way ANOVA and its effect size (partial η^2). Df is the degree of freedom.

Species	Stressor	Df	Df.res	*F* value	*p* value	partial η^2
Chlamydomonas sphaeroides	Salt	1	1196	1626.542	0	0.576
	Predator	1	1196	65.765	0	0.052
	Salt × Predator	1	1196	21.373	0	0.018
Chlamydomonas reinhardtii	Salt	1	1196	241.7099	0	0.168
	Predator	1	1196	14.1246	0	0.012
	Salt × Predator	1	1196	3.3138	0.069	0.003
Chlorella vulgaris	Salt	1	1196	7.1755	0.008	0.006
	Predator	1	1196	20.3152	0	0.017
	Salt × Predator	1	1196	4.5742	0.033	0.004

662   **Table 2**. Fixed effect in the linear mixed model on the clump sizes after predation. S.E.

663   is standard error and Df is the degree of freedom.

Variable	Explanation	Estimate	S.E.	Df	t value	p value
Intercept		2.054	0.081	5.157	25.307	0
Time	Six hours passed	−0.169	0.056	3592	−3.015	0.003
Predators	Presence of rotifers	−0.27	0.115	5.157	−2.354	0.064
Time × predators		0.343	0.115	5.157	2.991	0.029

665 Figures

Figure 1. Schematic representation of the system.

The green algae *Chlamydomonas sphaeroides* is consumed by rotifers *Brachionus calyciflorus* (the solid and dashed oblique arrows between them). While the salinity directly decreases the growth of both species (the three arrows from the salt), it also induces the clump formation of the green algae (the horizontal arrow). The clumps weaken the predator-prey interaction because rotifers are gape-limited predators (i.e., the trait-mediated indirect effect of salinity stress: the dashed oblique arrows). The salt icon by Servier https://smart.servier.com/ is licensed under CC-BY 3.0 Unported https://creativecommons.org/licenses/by/3.0/. Photos were taken by S. Shibasaki.

Figure 2. Large clump formation was induced by salinity stress in *Chlamydomonas*
sphaeroides.

Top: The distributions of clump sizes of *Chlamydomonas reinhardtii* (left),
Chlamydomonas sphaeroides (middle), and *Chlorella vulgaris* (right) under four
conditions (orange: control, blue: with rotifers, green: with 0.06M NaCl, and yellow:
with rotifers and 0.06M NaCl) after seven days. The black dots represent the median
clump sizes of 900 clumps. Bottom: examples of clump sizes of *Chlamydomonas*
sphaeroides in the control condition (left) and in the presence of salinity stress and
rotifers (right). The white bars represent the scale (50 μm). Photos were taken by S.
Shibasaki.

Figure 3. Large clumps were not consumed by rotifers.

The distributions of clump sizes of *Chlamydomonas sphaeroides* in the presence (A) or

absence (B) of rotifers in 0.06M NaCl C medium with three replicates (flasks). By

comparing the size distributions before (blue) and after (yellow) the exposure to rotifers

for six hours, it is clear that rotifers consumed smaller clumps (A) and salinity stress

cannot explain the difference (B). The black dots represent the median clump size of

300 clumps.

Figure 4. Predator-prey dynamics with or without salinity stress.

A and B: Population dynamics of *Chlamydomonas sphaeroides* (A) and the rotifers (B)

are shown. The shapes of markers represent the salinity stress (circle: 0M NaCl, and

triangle: 0.06M NaCl). Under each salinity stress, we performed a one-week experiment

with three replicates (shown by different colors). The error bars in panel A represent the

standard errors, while the markers show the mean values. C and D: The posterior

probability distributions of the fold changes of the rotifer's mortality rate (m_2/m_1: C) and

attack rates on *Chlamydomonas sphaeroides* (a_{22}/a_{21}: D) are shown, respectively. The

vertical blue lines show the median values, and the shaded areas show 95% CIs. The

vertical red dashed lines show one (i.e., no fold changes).