Extracellular vesicles carrying surface-anchored adiponectin prevent obesity-related metabolic complications by enhancing insulin sensitivity

Alexia Blandin[3,#], Josy Froger[1,2,#], Margot Voisin[1,2], Maëlle Lachat[3], Grégory Hilairet[2], Katarzyna Polak[3], Julie Magusto[1,2], Lisa Meslier[1,2], Mikaël Croyal[1,4], Mathilde Gourdel[4], Quentin Massiquot[2], Julien Chaigneau[5], Maxime Carpentier[1], Lionel Fizanne[5], Xavier Prieur[1], Cédric Le May[1], Jérôme Boursier[5], Bertrand Cariou[1], Robert Mamoun[3], Bernadette Trentin[3] and Soazig Le Lay[1,2,*]

[1]Nantes Université, CHU Nantes, CNRS, INSERM, l'institut du thorax, F-44000 Nantes, France.

[2]Université Angers, *SFR* ICAT, 49000 Angers, France

[3]Ciloa, 356 rue Maurice Béjart, 34080 Montpellier, France

[4]Nantes Université, CHU Nantes, Inserm, CNRS, SFR Santé, Inserm UMS 016, CNRS UMS 3556, Nantes, France

[5]HIFIH, CHU Angers, Université Angers, *SFR* ICAT, 49000 Angers, France

[#]These authors equally contributed to this work

*Corresponding author: Soazig LE LAY

L'unité de recherche de l'institut du thorax

Inserm UMR1087/CNRS UMR6291, IRS-UN – 8 quai Moncousu – BP 70721 44007 Nantes Cedex 1

e-mail: soazig.lelay@inserm.fr, Tel: +33-244-688-413

Keywords

4 Main Figures, 1 Supplementary Table**,** 3 Supplementary Figures

Total words (Introduction, Methods, Results and Discussion) : **3579**

References : 27

Abbreviations:

ACC, acetyl-CoA carboxylase; ACLY, ATP citrate lyase; AdipoQ, adiponectin gene; Col1A1, collagen type I alpha 1; ELOVL6, elongation of very long chain fatty acids protein 6; F4/80, EGF-like module-containing mucin-like hormone receptor-like 1 (macrophage marker); FAS, fatty acid synthase; IL-1β, interleukin-1 beta; IL-6, interleukin-6; MCP1, monocyte chemoattractant protein 1; PDK4, pyruvate dehydrogenase kinase 4; PGC1α, peroxisome proliferator-activated receptor gamma coactivator 1-alpha; PLIN1, perilipin 1; PLIN2, perilipin 2; PPARγ2, peroxisome proliferator-activated receptor gamma isoform 2; TGF-β1, transforming growth factor beta 1.

Abstract (197 words)

Adiponectin (Adpn) is a potent insulin-sensitizing adipokine with therapeutic promise for type 2 diabetes (T2D) and metabolic dysfunction-associated steatohepatitis (MASH). Its clinical use remains limited due to difficulties in producing stable, bioactive high–molecular weight forms. We previously demonstrated that adipose tissue-derived extracellular vesicles (EVs) naturally carry oligomeric Adpn on their surface, enhancing its stability and activity. Here, we engineered EVs displaying Adpn either anchored to the membrane (EV$^{PP-Adpn}$) or soluble (EV^{Adpn}), alongside control EVs lacking Adpn (EV^{CTL}), and evaluated their effects in a diet-induced obesity mouse model.

Mice received twice a week intraperitoneal injections of EVs or vehicle for six weeks. Only EV$^{PP-Adpn}$ significantly improved insulin sensitivity and reduced weight gain. Mechanistically, EV$^{PP-Adpn}$ enhanced adipocyte lipid storage through upregulation of insulin-responsive lipogenic genes and reduced hepatic steatosis, inflammation and fibrosis markers. It also activated AKT and AMPK pathways in liver and skeletal muscle, consistent with adiponectin signaling. In addition, EV$^{PP-Adpn}$ lowered circulating ceramides, lipids molecules promoting insulin resistance, in agreement with ceramidase activity induced by adiponectin receptor (AdipoR) signaling.

Our results demonstrate that surface-anchored Adpn EVs restore tissue-specific insulin signaling and ameliorate metabolic dysfunctions, highlighting their potential as a novel biotherapeutic strategy for T2D and MASH.

Introduction

The increasing global prevalence of type 2 diabetes (T2D) and its associated cardiometabolic complications underscores an urgent need for innovative therapeutic strategies, particularly given the limited availability of effective insulin-sensitizing molecules.

Among potential targets, adiponectin (Adpn) has emerged as a promising candidate due to its insulin-sensitizing, anti-inflammatory, and cardioprotective properties. Adpn, an adipokine, also known as adipocyte complement-related protein of 30 kDa (Acrp30), is specifically secreted by adipocytes [1]. Adpn abundantly circulates in blood in multimeric forms [2], and its plasma concentrations is decreased in patients with obesity and T2D [3]. Mechanistically, multimeric forms of Adpn exerts their effects through the activation of its receptors, AdipoR1 and AdipoR2, which are widely expressed in metabolic tissues [4]. These receptors mediate downstream signaling pathways such as AMP-activated protein kinase (AMPK) and peroxisome proliferator-activated receptor-α (PPARα), promoting glucose uptake, fatty acid oxidation, and anti-inflammatory responses [5]. However, the clinical translation of recombinant Adpn-based therapies has been hindered by significant challenges, especially the production of stable high molecular weight (HMW) Adpn oligomers.

We recently demonstrated that extracellular vesicles (EVs) derived from adipocytes are naturally coated with Adpn oligomers and act as natural cargos of Adpn accounting for 20% of the Adpn circulating pool [6]. Our results also demonstrated enhanced stability of EV-associated Adpn oligomers in the bloodstream compared to free Adpn as well as maintenance of Adpn-associated EVs' insulin-sensitizing effects [6]. EVs are lipid bilayer-enclosed structures secreted by various cell types, and they serve as carriers for bioactive molecules, including proteins, lipids, and nucleic acids [7]. EVs display natural abilities to cross biological barriers and interact with cells, modulating target cell responses and displaying immunomodulatory and regenerative activities [8]. Thus, EVs have recently emerged as promising drug delivery nano-vehicles due to their high bioavailability, excellent biocompatibility, and minimal immunogenicity, highlighting their potential as innovative therapeutic carriers [9].

This study aims to harness EVs as delivery systems for bioactive Adpn oligomers to treat metabolic disorders. To this goal, we mastered the production of customized EVs presenting oligomerized forms of Adpn at their surface and assess their therapeutic potential in mitigating insulin resistance and metabolic dysfunctions of high-fat diet fed mice. Treatment of these mice with EVs presenting Adpn at their surface enhances adipocyte lipid storage through insulin-regulated lipogenesis and lipid storage gene induction, while improving Metabolic dysfunction-Associated SteatoHepatitis (MASH) lesions (steatosis, inflammation, fibrosis) by restoring hepatic insulin sensitivity. Altogether, our results demonstrate the insulin-sensitizing potential

96   of surface Adpn-exposing EVs, paving the way for an innovative biotherapy for managing T2D
97   and MASH.
98
Methods
Animal experiments
Male and female C57BL/6J (B6) mice (10 weeks old) were fed a high-fat diet (HFD; 61% kcal
from fat, 20% protein, 20% carbohydrate; D12492, Safe Diets) *ad libitum* for 6 weeks. Mice
were housed under standard conditions (12 h light/12 h dark cycle) with *ad libitum* access to
food and water. During the HFD period, mice received biweekly intraperitoneal (i.p.) injections
of EVs, resuspended in 100 µL sterile 0.9% NaCl. Control animals received an equivalent
volume of PBS in 0.9% NaCl as vehicle. All procedures were approved by the French Ministry
of Research and the local ethics committee (CEEA - 006) and complied with EU Directive
2010/63/EU. At sacrifice, tissues were collected from either random-fed mice or after a 4 h fast
followed by insulin injection. At the end of the experiment, body weight and the weights of
organs/tissues, including liver, visceral white adipose tissue (VAT), subcutaneous white
adipose tissue (SAT), interscapular brown adipose tissue (BAT), pancreas, spleen and
quadriceps (representative of skeletal muscle) were recorded. Adiposity (%) was calculated as
the sum of VAT and SAT weights divided by total body weight. Samples were fixed in 10%
formalin for histological analysis or snap-frozen in liquid nitrogen and stored at $-80\,°C$ for
molecular analyses.
116
117   ***Hepatic Lipid Quantification***
118   Hepatic lipids were extracted from snap-frozen liver samples using a modified Bligh and Dyer
119   method[10]. Hepatic triglycerides (TG) were quantified using enzymatic kits (Roche Diagnostics)
120   and expressed relative to liver weight (µg TG per g of liver tissue).
121
122   ***Plasma FGF21 Measurement***
123   Circulating FGF21 concentrations was measured using commercial ELISA kits designed for
124   mouse measurments (R&D systems) from plasma samples according to the manufacturer's
125   protocol.
126
Metabolic Testing and Insulin Signaling Assessment
Glucose (tolerance test (GTT) and insulin (tolerance test (ITT) were conducted during the fifth
week of EV injections. Mice were fasted for 4 hours prior to testing. For GTT, glucose (1.25
g/kg body weight) was administered intraperitoneally (i.p**.**) at time 0. For ITT, mice were
injected i.p. with 0.75 IU/kg body weight human insulin (Humulin, Lilly) at time 0. Blood glucose
was measured from tail vein samples at 0, 15, 30, 45, 60, 90, and 120 minutes using

glucometer strips (Accu-Chek Guide). Plasma insulin concentrations were determined using

commercial mouse ELISA kits (Crystal Chem) from serum samples collected during the GTT

on heparin-coated microvettes (Sarstedt) following the manufacturer's instructions

To evaluate insulin signaling in target tissues, mice were fasted for 4 hours and injected i.p.

with human insulin (3 IU/kg; Humulin, Lilly). Tissues were collected 15 minutes post-injection

for analysis of insulin response.

Western blot

Insulin and AMPK signaling were assessed in mouse tissues collected 15 minutes after i.p.

insulin injection. VAT, SAT, and skeletal muscle (quadriceps) were harvested at this time point.

Tissue homogenization was performed on 20mg of snap-frozen liver or muscle tissue and 100

mg of snap-frozen VAT using CK14 soft tissue beads and the Precellys homogenizer,

according to the manufacturer's instructions. Protein lysates (15 µg) were subjected to SDS–

PAGE and Western blotting, as previously described[11]. The following primary antibodies were

used: phospho-Akt (Ser473, #4060), total Akt (pan, #4691), phospho-AMPKα (Thr172, #2535)

and total AMPKα (#2532) (Cell Signaling Technology), GAPDH (#ABC-AC002-100, ABclonal),

and β-actin (#A5316, Sigma-Aldrich). IRDye-conjugated secondary antibodies (LI-COR

Biosciences) were used, and signals were detected using an Odyssey CLx imaging system

(LI-COR). Quantification was performed using Image Studio software. Representative blots

per protein shown include samples from at least three independent animals per group.

Histology and tissue staining

Adipose and liver tissues were fixed in 10% PBS-buffered formalin for at least 24 hours,

paraffin-embedded, and sectioned at 5 µm. After deparaffinization and rehydration with specific

commercial solutions (Impath), sections were stained with with hematoxylin and eosin (HE) for

morphology and picrosirius red (PS) to assess fibrosis. Adipocyte diameter (VAT, SAT),

steatosis and fibrosis areas (liver) were quantified using dedicated ImageJ plugins.

RNA Extraction and qPCR

RNA was extracted from liquid nitrogen–frozen tissues. For VAT, RNA was isolated using

QIAzol and purified with the RNeasy Microkit (Qiagen), including DNase I treatment. Liver and

muscle RNA were extracted using the Maxwell® RSC simplyRNA Tissue Kit and instrument

(Promega). RNA quantity and purity were assessed via NanoDrop ND-2000 (Thermo Fisher).

cDNA was synthesized from 1 µg RNA using SuperScript™ II (Invitrogen) and random

hexamers, then purified with the Qiaquick PCR kit (Qiagen). qPCR was performed with 3 ng

cDNA, Maxima™ SYBR Green Master Mix (Thermo Fisher), and 0.3 µM primers using a CFX

Opus 384 system (Bio-Rad). Cycling conditions were 95°C for 10 min, then 40 cycles of 95°C

for 15 s and 60°C for 30 s. Specificity was confirmed by melting curve analysis. Gene

expression was calculated via the $2^{-\Delta\Delta Cq}$ method, normalized to 18S and 36B4, and analyzed

with CFX Maestro (Bio-Rad). Primer sequences are available upon request.

Ceramide analysis by mass spectrometry

Ceramides were quantified from 10 µL of plasma using targeted lipid liquid chromatography–

tandem mass spectrometry (LC-MS/MS), as previously described [11].

Quantification and Statistical Analysis

Data were analyzed with GraphPad Prism and presented as dot plots of independent

experiments. Statistical tests used are specified in figure legends. Significance was set at p ≤

0.05, indicated as *p ≤ 0.05, **p ≤ 0.01, ***p ≤ 0.005, and ****p ≤ 0.001.

Results

Generation and characterization of bioengineered EVs carrying Adpn.

To exploit the therapeutic potential of Adpn-associated EVs, we engineered EVs carrying native forms of Adpn according to a patented technology [12]. To this end, we used two chimeric gene constructs encoding either wild-type Adpn or Adpn anchored to the EV surface via its N-terminus, achieved by fusing the Adpn gene to a transmembrane domain and a pilot peptide (PP-Adpn) that directs it towards the EV secretion pathway (Figure S1A). Bioengineered EVs were isolated from the supernatants of HEK 293T cell lines stably transfected with Adpn (EV^{Adpn}) or with PP-Adpn (EV$^{PP-Adpn}$), using a controlled purification process combining tangential flow filtration and chromatography. In parallel, control EVs (EV^{CTL}) were produced from the supernatants of non-transfected HEK 293T cells. All EV preparations showed enrichment of the EV markers Alix and syntenin-1, as demonstrated by Wlestern blot (Figure S1B), and of CD81, as shown by ELISA (Figure S1C). Nano-flow cytometry analysis indicated that the size distribution of both control EVs and bioengineered EVs was within the expected range for EVs (50-120 nm) (Figure S1D), with no significant differences in EV median size (~70-90 nm) or concentration (~10^{10} particles/mL) among preparations (Figure S1E-F). Quantification of Adpn content per EV confirmed successful and consistent loading across bioengineered EV batches, while Adpn was undetectable in control EVs (Figure S1G). Western blot analysis of EV^{Adpn} and EV$^{PP-Adpn}$ confirmed the presence of Adpn, with a molecular weight consistent with the designed construct (Figure S1H). EV-associated Adpn was present in its high-molecular-weight proteo-oligomeric forms in both bioengineered EVs, which are recognized as the metabolically active forms (Figure S1I).

Dose-dependent metabolic effects of bioengineered EVs

As part of our investigation into the metabolic effects of bioengineered EVs carrying Adpn, we first sought to determine an effective and physiologically relevant dosing strategy. EV^{Adpn}, EV$^{PP-Adpn}$ and EV^{CTL} were administered by IP injections twice a week over a 6-week period to 10-week-old mice fed a high-fat diet (HFD). Three doses of EV-associated Adpn (0.5 ng, 2.5 ng, and 25 ng) were tested, with dosing standardized to the quantified Adpn content of each EV preparation (Figure S2A-B). The lowest dose (0.5 ng) matched the Adpn amount found in 5 µg of VAT-derived EVs previously shown to enhance insulin sensitivity[6]. The amount of EV^{CTL}, devoid of any Adpn, was matched to the highest particle number used among EV^{Adpn} and EV$^{PP-Adpn}$ to control for potential particle-related effects (Figure S2B). A PBS-injected group served as vehicle control to distinguish Adpn-associated EV specific effects.

The highest dose (25 ng) of EV^{CTL} and EV^{Adpn} (1.45 × 10^8 EVs/injection; Figure S2B) increased weight gain after 6 weeks of HFD compared to the lowest dose (0.5 ng; Figure S2C-D;F-G),

and was associated with increased BAT weight for EV^{CTL} (Figure S2K) and higher spleen weight for EV^{Adpn} (Figure S2N). Likewise, the lowest dose (0.5 ng) of $EV^{PP-Adpn}$ (1.0×10^7 EVs) significantly increased weight gain versus vehicle (Figure S2E;H), along with elevated subcutaneous AT (SAT) weight (Figure S2J) and higher glycemia in both fed and fasted states (Figure S2O-P). In contrast, 25 ng of $EV^{PP-Adpn}$ (5.0×10^8 EVs) significantly reduced both weight gain and SAT mass compared to its 0.5 ng dose (Figure S2E;H;J), despite involving more particles (Figure S2B). These findings suggest that higher EV concentrations may worsen metabolic outcomes under HFD, unless EVs are surface-coated with Adpn, as illustrated with high-dose $EV^{PP-Adpn}$. Based on these observations, the highest dose (25 ng) was selected for subsequent experiments to enable comprehensive metabolic phenotyping of bioengineered EVs and their therapeutic potential.

Surface Adpn-anchored EVs ($EV^{PP-Adpn}$) improve insulin sensitivity and glucose tolerance

Male and female HFD-fed mice were thus treated with EV^{CTL}, EV^{Adpn} or $EV^{PP-Adpn}$ (at 25 ng Adpn-equivalent doses, or equivalent particle numbers for EV^{CTL}), or with vehicle (PBS), to assess their metabolic effects. No significant differences in body weight over time (Figure 1A), weight gain (Figure 1B) or adiposity at sacrifice (Figure 1C) were observed between groups, regardless of sex. Similarly, no significant differences in individual tissue weights (including AT depots, liver, pancreas, spleen and muscle) were detected between groups at sacrifice, indicating the absence of hepatomegaly or splenomegaly and supporting the lack of EV-induced organotoxicity (Figure S3A-G). In addition, serum biochemistry analyses (Table S1) revealed no signs of hepatic or renal toxicity, further supporting the safety of repeated EV injections. Random-fed glycemia, monitored throughout the experiment, remained stable in both females and males (Figure 1D), as did fasting glycemia, measured at the end of the protocol (Figure 1E). Insulin tolerance tests (ITT) demonstrated a significant improvement in insulin sensitivity exclusively in $EV^{PP-Adpn}$ treated mice of both sexes, as indicated by increased area over the curve (AOC) (Figure 1F). Similarly, intraperitoneal glucose tolerance tests (IP-GTT) showed improved glucose tolerance in males treated with $EV^{PP-Adpn}$, while in females, both EV^{Adpn} and $EV^{PP-Adpn}$ enhanced glucose tolerance (Figure 1G). Glucose intolerance induced by HFD is often associated with impaired insulin secretion. In the $EV^{PP-Adpn}$-treated group, the improvement in glucose tolerance was accompanied by the restoration of the 15-minute insulin peak following glucose load (T15 vs. T0), which was absent in all other groups (Figure 1H). No such induction was detected in any group of females (Figure 1H). These improvements occurred without detectable changes in pancreatic β-cell mass or size, as indicated by comparable insulin immunostaining across groups on pancreatic sections (Figure S3K-N).

$EV^{PP\text{-}Adpn}$ promote healthy adipose tissue expansion

Although total AT mass remained unchanged across groups at the end of treatment (Figure 1C, Figure S3A-C), histological analysis of VAT revealed a significant shift in adipocyte size distribution toward larger diameters in both EV^{Adpn} and $EV^{PP\text{-}Adpn}$ treated mice, reflecting an increase in average adipocyte size (Figure 2A-C). This effect was specific to VAT and was not observed in SAT (Figure S3H-J). Importantly, only $EV^{PP\text{-}Adpn}$ treatment resulted in a coordinated upregulation of mRNA levels for key adipocyte markers such as PPARγ2, PLIN1, and AdipoQ (Figure 2D), as well as insulin-regulated lipogenic enzymes gene expression including ELOVL6, FAS, ACC and ACLY (Figure 2E). No significant changes were detected in the expression of inflammatory genes (IL1β, MCP1, F4/80, IL6) (Figure 2F). Of note, circulating NEFA levels were unchanged across groups (Table S1), indicating that basal lipolysis is not altered by any EV treatments. These histological and molecular changes indicate that $EV^{PP\text{-}Adpn}$ promote insulin-sensitive AT expansion, consistent with the known effects of Adpn on these pathways, which likely support more efficient fat storage and healthier adipose tissue remodeling.

$EV^{PP\text{-}Adpn}$ alleviate liver steatosis and MASH-related gene expression

Liver histology revealed a marked reduction in steatosis in $EV^{PP\text{-}Adpn}$–treated animals (Figure 3A–B), further supported by a significant decrease in hepatic triglyceride content (Figure 3D) and reduced protein expression of PPARγ2 (Figure 3E-F). In line with these findings, serum biochemistry analyses revealed significant improved liver enzyme profiles (ALAT, ASAT) in both EV^{Adpn} and $EV^{PP\text{-}Adpn}$ treated groups (Table S1), supporting the beneficial hepatic effects observed histologically. Although fibrosis area remained unchanged across liver sections (Figure 3C), consistent with the limited fibrotic response in HFD-fed mouse models, we observed a significant downregulation of MASLD-related genes in the $EV^{PP\text{-}Adpn}$ group. These included the lipid droplet marker PLIN2, pro-inflammatory genes (IL1β, MCP1, F4/80), and fibrotic markers (TGFβ1, Col1A1) (Figure 3G), indicating improved lipid handling and attenuated inflammatory and fibrotic signaling.

Analysis of hepatic metabolic gene expression showed no major changes in lipogenic (FAS, ACC) or oxidative (PDK4, PGC1α) genes (Figure 3H). Phosphoenolpyruvate Carboxykinase (PEPCK) is a gluconeogenic enzyme normally downregulated by insulin, but this repression is lost in HFD-fed mice[13]. $EV^{PP\text{-}Adpn}$ treatment restored this repression, as shown by a significant reduction in PEPCK mRNA expression, with a similar trend observed for glucose-6-phosphatase (G6Pase) (Figure 3H).

Circulating FGF21, a hepatokine elevated in response to liver fat accumulation and metabolic stress, was drastically reduced in $EV^{PP\text{-}Adpn}$ treated mice (Figure 3I). Previous reports have

shown that increased FGF21 reflects a compensatory but dysregulated response to hepatic
lipid overload and impaired metabolic signaling in MASLD [14,15]. Therefore, the reduction of
FGF21 following EV$^{PP-Adpn}$ treatment likely indicates improved hepatic lipid metabolism and
restored metabolic homeostasis.
These findings suggest that EV$^{PP-Adpn}$ selectively restores hepatic insulin responsiveness at the
transcriptional level and alleviates key features of MASH, including steatosis, inflammation,
and fibrotic gene expression, thereby highlighting its therapeutic potential in MASLD.
304
EV$^{PP-Adpn}$ promotes adiponectin-dependent insulin signaling and ceramide clearance
To further investigate the insulin-sensitizing effects of bioengineered EVs displaying surface-
anchored adiponectin (EV$^{PP-Adpn}$), we evaluated the activation of key adiponectin-responsive
signaling pathways in VAT, liver, and skeletal muscle following acute insulin stimulation (3 U,
15 min). Western blot analyses revealed an increased insulin-stimulated phosphorylation of
AKT (Ser473) in all examined tissues of EV$^{PP-Adpn}$ treated mice compared to EV^{CTL} mice,
reaching statistical significance in the liver, thereby confirming enhanced hepatic insulin
signaling (Figure 4A). Additionally, we observed a general trend toward increased AMPK
(Thr172) phosphorylation across tissues, a pathway known to mediate adiponectin's effects
on lipid metabolism and energy homeostasis (Figure 4B). This effect was significant in skeletal
muscle, suggesting enhanced metabolic flexibility in this tissue (Figure 4B). We further
investigated plasma ceramide levels, given the ceramidase activity linked to AdipoR signaling
and the well-established lipotoxic role of ceramides in insulin resistance. Notably, EV$^{PP-Adpn}$
treatment significantly decreased total ceramides (Figure 4C), dihydroceramides (Figure 4D),
and hexosylceramides (Figure 4E), particularly long-chain subspecies (C22:00 to C24:00),
consistent with the systemic metabolic improvements observed. Altogether, these findings
support that EV$^{PP-Adpn}$ enhances tissue-specific insulin signaling via engagement of the
adiponectin pathway, in line with transcriptional data indicating restored insulin
responsiveness.
324
325

Discussion

Adpn is a promising target for managing T2D-related cardiometabolic complications by enhancing insulin sensitivity, but the lack of stable, bioactive multimeric forms has so far hampered its therapeutic development. To address this, we engineered two types of Adpn-enriched EVs, either broadly distributed (EV^{Adpn}) or anchored at the surface (EV$^{PP-Adpn}$), using a HEK-derived EV bioingeneering platform. Adoptive transfer of EV$^{PP-Adpn}$ in HFD-fed mice improved insulin sensitivity, promoted insulin-driven adipose lipid storage, and reduced hepatic steatosis, inflammation, and fibrosis. In contrast, EV^{Adpn} did not display such beneficial effects. These results support the therapeutic potential of surface Adpn-enriched EVs as an innovative strategy to counteract insulin resistance-related metabolic disorders such as T2D and metabolic dysfunction-associated steatohepatitis (MASH).

In our study, mice injected intraperitoneally with increasing doses of control HEK293T-derived EVs (EV^{CTL}, devoid of any Adpn) exhibited significant weight gain, especially at doses above 1.0×10^8 particles. Cross-study comparisons remain difficult due to variability in EV quantification methods; NanoFCM, for example, tends to underestimate particle counts compare to nanoparticle tracking analysis (NTA; SLL, *unpublished data*). Dosing strategies also differ, with some studies relying on particle number and others on EV protein content. Furthermore, most toxicity and immunogenicity assessments are performed under normal diet, with limited evaluation of metabolic effects under HFD. Yet, the i.p. injection route and EV origin (e.g., HEK293, allogeneic) may also influence immune responses, as HEK-derived EVs can carry mildly immunogenic components[16,17]. Notably, the increase in neutrophils count post-HEK293-derived EV injection may be relevant in HFD-induced obesity[17], given that obesity-related inflammation involves neutrophil and macrophage recruitment in adipose tissue[18]. These observations stress the need to assess EV metabolic effects with proper controls (vehicle, empty EVs) and under dietary stress conditions such as HFD, mimicking the obesogenic nutritional environment.

EV$^{PP-Adpn}$, engineered to display Adpn specifically at the vesicle surface, exert significant insulin-sensitizing effects, unlike EVs containing non-surface-anchored Adpn. This suggests that surface-localized Adpn engages AdipoR1/R2 receptors, consistent with our previous findings using lean adipose-derived EVs [6]. While no significant changes were observed in total body weight or tissue mass, EV$^{PP-Adpn}$ improved insulin sensitivity, primarily in the liver, by suppressing gluconeogenesis and reducing hepatic steatosis, inflammation, and fibrosis. In adipose tissue, EV$^{PP-Adpn}$ induced insulin-regulated lipogenic and adipocyte remodeling genes, promoting metabolically healthy expansion. These effects were accompanied by enhanced

glucose-stimulated insulin secretion, without altering pancreatic β-cell mass. In parallel, EV[PP-]

[Adpn] reduced circulating ceramide levels, consistent with AdipoR-mediated ceramidase

activitation [19]. As ceramides are recognized as lipotoxic mediators impairing insulin signaling

and mitochondrial function, their reduction may drive the metabolic benefits observed, although

causality remains to be clarified [20]. Finally, the observed reduction in circulating FGF21 levels

following EV[PP-Adpn] treatment likely reflects a decreased need for compensatory hepatokine

secretion, consistent with restored hepatic metabolic function. This is relevant given that

elevated FGF21 levels in MASLD have been associated with impaired metabolic signaling and

FGF21 resistance, characterized by reduced receptor responsiveness despite high ligand

levels [15,21]. Its normalization may thus indicate a reversal of this dysfunctional axis and further

support the systemic metabolic benefits of EV[PP-Adpn].

Although EV[PP-Adpn] exert clear insulin-sensitizing effects, particularly in the liver and adipose

tissue, their capacity to reverse insulin resistance appears more modest than that observed

with native adipocyte-derived EVs [6]. This may reflect the combinatorial action of diverse

bioactive cargos within adipose EVs, including proteins, lipids, microRNAs or mitochondrial

components, acting synergistically to promote metabolic benefits [22]. Recent studies highlight

the beneficial effect of adipose cargos as engineering of EVs to enrich them with adipokines

reveal their beneficial metabolic potential. For instance, loading leptin into macrophage-derived

EVs thus successfully targeted the brain and restored breathing function in obese mice,

overcoming leptin resistance[23]. Similarly to our bioengineered Adpn EVs, apelin-loaded EVs

improved glucose tolerance and activated AMPK and Akt pathways in diabetic mice,

enhancing systemic insulin sensitivity [24]. In a separate study, EVs engineered for targeted

apelin delivery to pulmonary arterial hypertension lesions reversed pathological vascular

remodeling and improved cardiac function[25]. Finally, knowing that adiponectin is recognized

as one of the main downstream effectors of the metabolic effects of FGF21 on glucose

homeostasis and insulin sensitivity in mice [26], bioengineered EVs combining surface FGF21

and enclosed miR-223 demonstrated liver-preferential delivery and a marked reduction in the

MASH phenotype [27]. These examples underscore the therapeutic potential of combinatorial

EV cargoes in metabolic disease. Thus, while EV[PP-Adpn] already demonstrate insulin-sensitive

properties, their clinical optimization may require integration with complementary bioactive

payloads to fully recapitulate the potency of native adipose EVs.

References

1. Scherer, P.E., Williams, S., Fogliano, M., Baldini, G., and Lodish, H.F. (1995). A novel serum protein similar to C1q, produced exclusively in adipocytes. J Biol Chem *270*, 26746-26749. 10.1074/jbc.270.45.26746.

2. Pajvani, U.B., Du, X., Combs, T.P., Berg, A.H., Rajala, M.W., Schulthess, T., Engel, J., Brownlee, M., and Scherer, P.E. (2003). Structure-function studies of the adipocyte-secreted hormone Acrp30/adiponectin. Implications fpr metabolic regulation and bioactivity. J Biol Chem *278*, 9073-9085. 10.1074/jbc.M207198200.

3. Straub, L.G., and Scherer, P.E. (2019). Metabolic Messengers: Adiponectin. Nat Metab *1*, 334-339. 10.1038/s42255-019-0041-z.

4. Yamauchi, T., Kamon, J., Ito, Y., Tsuchida, A., Yokomizo, T., Kita, S., Sugiyama, T., Miyagishi, M., Hara, K., Tsunoda, M., et al. (2003). Cloning of adiponectin receptors that mediate antidiabetic metabolic effects. Nature *423*, 762-769. 10.1038/nature01705.

5. Li, Y., Onodera, T., and Scherer, P.E. (2024). Adiponectin. Trends Endocrinol Metab *35*, 674-675. 10.1016/j.tem.2024.05.004.

6. Blandin, A., Amosse, J., Froger, J., Hilairet, G., Durcin, M., Fizanne, L., Ghesquiere, V., Prieur, X., Chaigneau, J., Vergori, L., et al. (2023). Extracellular vesicles are carriers of adiponectin with insulin-sensitizing and anti-inflammatory properties. Cell reports *42*, 112866. 10.1016/j.celrep.2023.112866.

7. Cocozza, F., Grisard, E., Martin-Jaular, L., Mathieu, M., and Thery, C. (2020). SnapShot: Extracellular Vesicles. Cell *182*, 262-262 e261. 10.1016/j.cell.2020.04.054.

8. Salomon, C., Das, S., Erdbrugger, U., Kalluri, R., Kiang Lim, S., Olefsky, J.M., Rice, G.E., Sahoo, S., Andy Tao, W., Vader, P., et al. (2022). Extracellular Vesicles and Their Emerging Roles as Cellular Messengers in Endocrinology: An Endocrine Society Scientific Statement. Endocr Rev *43*, 441-468. 10.1210/endrev/bnac009.

9. Claridge, B., Lozano, J., Poh, Q.H., and Greening, D.W. (2021). Development of Extracellular Vesicle Therapeutics: Challenges, Considerations, and Opportunities. Front Cell Dev Biol *9*, 734720. 10.3389/fcell.2021.734720.

10. Sotin, T., Ge, X., Schonke, M., Vince, L., Thouzeau, A., Frey, S., Lorant, V., Krul, L., Pronk, A.C.M., Lalai, R., et al. (2025). A rare gain of function variant of hepatic lipase attenuates hypercholesterolemia and atherosclerosis in mice via an LDL receptor-independent mechanism. Cardiovascular research. 10.1093/cvr/cvaf097.

11. Croyal, M., Kaabia, Z., Leon, L., Ramin-Mangata, S., Baty, T., Fall, F., Billon-Crossouard, S., Aguesse, A., Hollstein, T., Sullivan, D.R., et al. (2018). Fenofibrate decreases plasma ceramide in type 2 diabetes patients: A novel marker of CVD? Diabetes Metab *44*, 143-149. 10.1016/j.diabet.2017.04.003.

12. Mamoun, R., Trentin, B., and Polak, K. (2023). Chimeric adiponectin polypeptides, extracellular vesicle comprising the same, and uses thereof. International patent WO/2023/104822, patent application PCT/EP2022/084653 06.12.2022.

13. Samuel, V.T., and Shulman, G.I. (2016). The pathogenesis of insulin resistance: integrating signaling pathways and substrate flux. J Clin Invest *126*, 12-22. 10.1172/JCI77812.

14. Rusli, F., Deelen, J., Andriyani, E., Boekschoten, M.V., Lute, C., van den Akker, E.B., Muller, M., Beekman, M., and Steegenga, W.T. (2016). Fibroblast growth factor 21 reflects liver fat accumulation and dysregulation of signalling pathways in the liver of C57BL/6J mice. Scientific reports *6*, 30484. 10.1038/srep30484.

15.   Fisher, F.M., Chui, P.C., Antonellis, P.J., Bina, H.A., Kharitonenkov, A., Flier, J.S., and
Maratos-Flier, E. (2010). Obesity is a fibroblast growth factor 21 (FGF21)-resistant
state. Diabetes *59*, 2781-2789. 10.2337/db10-0193.
16.   Chen, Z.Q., Tang, T.T., Tang, R.N., Zhang, Y., Zhang, Y.L., Yang, H.B., Song, J., Yang, Q.,
Qin, S.F., Chen, F., et al. (2025). A comprehensive evaluation of stability and safety for
HEK293F-derived extracellular vesicles as promising drug delivery vehicles. J Control
Release *382*, 113673. 10.1016/j.jconrel.2025.113673.
17.   Zhu, X., Badawi, M., Pomeroy, S., Sutaria, D.S., Xie, Z., Baek, A., Jiang, J., Elgamal, O.A.,
Mo, X., Perle, K., et al. (2017). Comprehensive toxicity and immunogenicity studies
reveal minimal effects in mice following sustained dosing of extracellular vesicles
derived from HEK293T cells. Journal of extracellular vesicles *6*, 1324730.
10.1080/20013078.2017.1324730.
18.   Uribe-Querol, E., and Rosales, C. (2022). Neutrophils Actively Contribute to Obesity-
Associated Inflammation and Pathological Complications. Cells *11*.
10.3390/cells11121883.
19.   Holland, W.L., Miller, R.A., Wang, Z.V., Sun, K., Barth, B.M., Bui, H.H., Davis, K.E.,
Bikman, B.T., Halberg, N., Rutkowski, J.M., et al. (2011). Receptor-mediated activation
of ceramidase activity initiates the pleiotropic actions of adiponectin. Nat Med *17*, 55-
63. 10.1038/nm.2277.
20.   Summers, S.A. (2018). Could Ceramides Become the New Cholesterol? Cell Metab *27*,
276-280. 10.1016/j.cmet.2017.12.003.
21.   Geng, L., Lam, K.S.L., and Xu, A. (2020). The therapeutic potential of FGF21 in
metabolic diseases: from bench to clinic. Nat Rev Endocrinol *16*, 654-667.
10.1038/s41574-020-0386-0.
22.   Le Lay, S., and Scherer, P.E. (2025). Exploring adipose tissue-derived extracellular
vesicles in inter-organ crosstalk: Implications for metabolic regulation and adipose
tissue function. Cell reports *44*, 115732. 10.1016/j.celrep.2025.115732.
23.   Freire, C., Ramsey, J.D., Pho, H., Kojima, R., Zhao, Y., Kim, L., Anokye-Danso, F., Berger,
S., Ahima, R.S., Batrakova, E.V., et al. (2022). Leptin-loaded Extracellular Vesicles Treat
Sleep-disordered Breathing in Mice with Obesity. Am J Respir Cell Mol Biol *67*, 720-
723. 10.1165/rcmb.2022-0229LE.
24.   Cui, J., Wang, M., Zhang, W., Sun, J., Zhang, Y., Zhao, L., Hong, Z., Li, D., Huang, Y.X.,
Zhang, N., and Chen, Y. (2024). Enhancing insulin sensitivity in type 2 diabetes
mellitus using apelin-loaded small extracellular vesicles from Wharton's jelly-derived
mesenchymal stem cells: a novel therapeutic approach. Diabetol Metab Syndr *16*, 84.
10.1186/s13098-024-01332-w.
25.   Kim, J., Choi, Y.S., Kim, J., Wahab, R., Lee, Y., Joo, H., Park, S., Lee, I., Kim, S.A., Hwang,
Y.H., et al. (2025). Targeted delivery of apelin using a novel extracellular vesicle
platform for pulmonary arterial hypertension treatment. Biomaterials *323*, 123438.
10.1016/j.biomaterials.2025.123438.
26.   Lin, Z., Tian, H., Lam, K.S., Lin, S., Hoo, R.C., Konishi, M., Itoh, N., Wang, Y., Bornstein,
S.R., Xu, A., and Li, X. (2013). Adiponectin mediates the metabolic effects of FGF21 on
glucose homeostasis and insulin sensitivity in mice. Cell Metab *17*, 779-789.
10.1016/j.cmet.2013.04.005.
27.   Cho, H., Ju, H., Ahn, Y., Jang, J., Cho, J., Park, E., Kang, S.M., Lee, J., Seo, D., Baek, M.C.,
and Yea, K. (2025). Engineered extracellular vesicles with surface FGF21 and enclosed

miR-223 for treating metabolic dysfunction-associated steatohepatitis. Biomaterials
321, 123321. 10.1016/j.biomaterials.2025.123321.
493
494

ACKNOWLEDGMENTS

We thank the SCAHU platform, especially Elodie Guillet, Jérôme Roux, Adeline Mourlan, Nina Meignan-Fizanne for mouse housing, care and technical assistance for animal functional exploration. We also thank the PACeM platform, especially L. Bonneau and J. Cayon, for qRT-PCR experiments.

The authors were supported the French National Research Agency (ANR-22-CE18-0026-02 EVADIPO). SLL is granted by Genavie, FHU GO NASH, Société Francophone du Diabète and the FFRD (sponsored by Fédération Française des Diabétiques, Abbott, AstraZeneca, Eli Lilly, Merck Sharp & Dohme, and Novo Nordisk).

The graphical abstract was created with BioRender.com

DECLARATION OF INTERESTS

These authors declare the following potential conflict of interests:

R.M. is a co-founder and CEO of Ciloa ; B.T is a co-founder and CSO of Ciloa.

B.T., K.P., and R.M. are inventors of patent WO2023104822A1 describing the stable anchoring of Adiponectin on EVs.

S.L.L. declare no competing interests.

All other authors declare no potential conflict of interest.

INCLUSION AND DIVERSITY

We support inclusive, diverse, and equitable conduct of research.

AUTHOR CONTRIBUTIONS

Methodology for EV production and characterization, A.B., M.L., K.P., R.M., B.T.

Methodology for animal experimentation and analysis, J.F., M.V., G.H., L.M., J.M., L.F., X.P., C.L.M., B.C., S.L.L.

Histology analysis, Q.M., J.C.

Blood biochemistry analysis, M.C.

Lipidomic analysis, M.C., M.G. ;

Conceptualization, S.L.L., B.C., R.M., B.T.

Writing – original draft, S.L.L.

Writing, review and editing, J.F., M.V., X.P., C.L.M., J.B., B.C., A.B., R.M.,S.L.L.

Supervision of the animal experimental part and project coordination, S.L.L.

All authors have read and agreed to the published version of the manuscript.

S.L.L. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of the data and the accuracy of the data analysis.

Figure 1. Surface-anchored Adpn-EVs (EV^PP-Adpn) improve insulin sensitivity and glucose tolerance in HFD-fed mice.

Figure 1

Figure 1. Surface-anchored Adpn-EVs (EV$^{PP-Adpn}$) improve insulin sensitivity and glucose tolerance in HFD-fed mice.

(A-C) Body weight trajectory **(A)**, total weight gain **(B)**, and adiposity (%) at sacrifice **(C)** in male (top) and female (bottom) HFD-fed mice after 6 weeks of i.p. injections with EV^{CTL}, EV^{Adpn} or EV$^{PP-Adpn}$ (25 ng Adpn-equivalent per dose, or particle-matched EV^{CTL}) or vehicle (PBS).

(D-E) Random-fed glycemia measured throughout the experiment, n=16-18 for males (top); n=9-10 for females (bottom) **(D)** and fasting glycemia measured at sacrifice **(E)** in males (top) and females (bottom).

(F) Insulin tolerance tests (ITT) performed in males (top) and females (bottom), with area over the curve (AOC) presented on the right.

(G) Intraperitoneal glucose tolerance tests (GTT) in males (top) and females (bottom), with area under the curve (AUC) shown on the right.

(H) Insulin levels measured at baseline (T0) and 15 minutes post-glucose injection (T15) during GTT in males (top) and females (bottom). Five animals per group were analyzed.

Data are presented as mean ± SEM. Dot plots represent the number of independent animals analyzed. Statistical differences were calculated using one-way ANOVA followed a Tukey's multiple comparisons test. $*p < 0.05$, $**p < 0.01$.

Figure 2. EV^{PP-Adpn} promote promote insulin-sensitive AT expansion inducing healthy visceral adipose tissue remodeling

Figure 2

Figure 2. EV^PP-Adpn promote promote insulin-sensitive AT expansion inducing healthy visceral adipose tissue remodeling

(A-C) Histological analysis of VAT in male HFD-fed mice after 6 weeks of i.p. injections with EV^CTL, EV^Adpn, or EV^PP-Adpn (25 ng Adpn-equivalent per dose or particle-matched EV^CTL) or vehicle (PBS). Representative hematoxylin–eosin (HE)-stained VAT sections are shown **(A)**, from which adipocyte diameters were measured. Analysis of VAT adipocyte size distribution is presented in **(B)**, and mean adipocyte area in **(C)**. n=6-8 animals per group were analyzed. Scale bar, 100 μm.

(D-F) Quantitive PCR gene expression analysis in VAT from male HFD-fed injected mice. mRNA expression was analyzed for key adipogenic markers (PDK4, PPARγ2, PLIN1, AdipoQ) **(D)**, insulin-responsive lipogenic genes (ELOVL6, FAS, ACC, ACLY) **(E)** and pro-inflammatory markers (IL1β, MCP1, F4/80, IL6) **(F)**. Bar colors: Green, Vehicle; Red, EV^CTL; Black, EV^Adpn; Blue, EV^PP-Adpn.

Data are presented as mean ± SEM. Dot plots represent the number of independent animals analyzed. Statistical differences were calculated using one-way ANOVA followed a Tukey's multiple comparisons test. $*p < 0.05$, $**p < 0.01$.

Figure 3. EV^PP-Adpn reduce liver steatosis and improve MASLD/MASH-related gene expression signatures in HFD-fed mice.

Figure 3

Figure 3. EV^{PP-Adpn} reduce liver steatosis and improve MASLD/MASH-related gene expression signatures in HFD-fed mice.

(A-C) Histological analysis of liver sections. Representative hematoxylin–eosin (HE) and picrosirius red (PS) staining (A) are shown for male mice after 6 weeks of treatment with EV^{CTL}, EV^{Adpn} or EV^{PP-Adpn} (25 ng Adpn-equivalent per dose, or particle-matched EV^{CTL}) or vehicle (PBS). Quantification of steatosis area on HE-stained sections **(B)** and fibrosis area on PS-stained sections **(C)** is presented. n=6-8 animals per group were analyzed. Scale bar, 100 µm.

(D) Biochemical measurement of hepatic triglyceride content.

(E–F) PPARγ2 protein expression in liver tissue from injected male mice. Representative image blots **(E)** and protein quantification **(F)** are presented. β-actin serve as loading control.

(G-H) Quantitative PCR analysis of hepatic gene expression in HFD-fed injected male mice. Relative mRNA levels of MASLD-associated genes **(G)** including steatotic (PLIN2), pro-inflammatory (IL1β, MCP1, F4/80) and fibrotic markers (TGFβ1, Col1A1). Expression of key metabolic genes **(H)** related to gluconeogenesis (PEPCK, G6Pase), lipogenesis (FAS, ACC) and β-oxidation (PDK4, PGC1α). Bar colors: Green, Vehicle; Red, EV^{CTL}; Black, EV^{Adpn}; Blue, EV^{PP-Adpn}.

(I) Circulating FGF-21 in injected male mice.

Data are presented as mean ± SEM. Dot plots represent the number of independent animals analyzed. Statistical significance was assessed using one-way ANOVA followed by Tukey's multiple comparisons test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure 4. EVPP-Adpn promote adiponectin-dependent insulin signaling and reduces plasma ceramide levels.

Figure 4

Figure 4. EV^{PP-Adpn} promote adiponectin-dependent insulin signaling and reduces

plasma ceramide levels.

(A-B) Western blot analysis of AKT (Ser473) **(A)** and AMPK (Thr172) **(B)** phosphorylation in

liver, VAT, and skeletal muscle following insulin injection in mice. Representative blots are

shown, with GAPDH as loading control. Quantification of phospho-AKT/total AKT **(A)** and

phospho-AMPK/total AMPK ratios **(B)** are presented below the respective blots.

(C-E) Plasma ceramide profiling for total and subspecies of ceramides **(C)**, dihydroceramides

(D) and hexosylceramides **(E)**.

Data are presented as mean ± SEM. Dot plots represent the number of independent animals

analyzed. Statistical significance was assessed by one-way ANOVA with Tukey's post hoc

test. $*p < 0.05$, $**p < 0.01$, $****p < 0.0001$.