Meta-analysis reveals the tempo of evolutionary parallelism of local adaptation between native and introduced ranges of plant species

Romane Normand[a,*], Alexis Heckley[b,c], Kathryn Hodgins[d], Samantha Grover[e], Tim Connallon[d,†], Akane Uesugi[a,†]

[a] Department of Biology, School of Science, RMIT University, Bundoora, VIC 3083, Australia
[b] Department of Biology, McGill University, Montréal, Quebec, Canada
[c] School of Biological Sciences, University of Oklahoma, Norman, OK, USA
[d] School of Biological Sciences, Monash University, Clayton, VIC 3800, Australia
[e] Applied Chemistry and Environmental Science, School of Science, RMIT University, Melbourne, VIC 3000, Australia

[*] Corresponding author (romane.r.normand@gmail.com)
[†] Equal contributions

Classification: Biological Sciences/Evolution

Keywords: clinal divergence | parallel patterns | evolutionary predictability | environmental gradients | plant traits

Abstract

Natural selection can cause evolution to be predictable despite the randomness of processes that generate genetic variation. However, the predictability of evolution can be difficult to evaluate, owing to the complexity of environmental conditions that determine selection in each species, and our inability to rewind and replay the tape of life to directly assess evolution's repeatability. Invasive species are valuable systems for evaluating evolutionary predictability, as populations in native and introduced ranges evolve separately, yet often encounter similar environmental challenges, providing potential for parallel patterns of local adaptation between ranges. However, it remains unclear how pervasive such parallelism is, or how rapidly it evolves following species' introductions. To address these questions, we carried out a meta-analysis of clinal divergence in native and introduced populations of the same plant species and evaluated the tempo of evolutionary parallelism of clines between ranges. We show that clinal divergence in introduced ranges was shallower than clinal divergence in native ranges for size, morpho-physiology, and phenology traits, but similar for reproductive and defense traits. Evolutionary parallelism of clinal divergence increased with the age of the species' introduction, with greater parallelism in relatively old introductions caused by increased alignment of the direction of clinal divergence between the ranges rather than changes in the relative magnitudes of divergence in each range. These results suggest a high predictability of local adaptation in mature species' introductions, with drift, genetic constraints and founder effects likely contributing to evolutionary lags in local adaptation within young introductions.

Significance Statement

Species with broad geographic ranges must cope with diverse environments that generate

selection for local adaptation. Invasive species are powerful systems for evaluating the rate and

ways that species locally adapt, potentially including parallel patterns of trait divergence

(clines) within native and introduced ranges. We conducted a meta-analysis of trait clines in

native and introduced populations of the same plant species to quantify the tempo and

predictability of parallelism between ranges. Introduced populations rapidly re-evolve clines

observed within the native range, with degrees of parallelism predicted by the age of the

introduction. Strong parallelism emerges within 200 generations following species

introductions, owing to increased alignment of cline directions over time. These findings

suggest a surprisingly high predictability of local adaptation.

Introduction

A general understanding of the process of adaptation must address several broad questions

about how ecology, evolution and genetics interact in natural populations. How predictable is

evolution? Do shared environmental conditions lead to reliably parallel evolutionary outcomes

(1, 2), or are there multiple genetic or phenotypic solutions to the same environmental

challenges (3)? Do populations typically adapt rapidly over ecological timescales (4, 5), or do

they experience sustained evolutionary lags in adaptation to their environments, owing to

limited genetic variation or other constraints (6, 7)? To what extent does evolution influence

ecological dynamics, including expansion, contraction, and fragmentation of species ranges

(8–10)? What are the key environmental variables—abiotic and biotic—that determine natural

selection and the direction of evolutionary change (11, 12)? Each of these questions is difficult

to answer, especially in natural systems.

Invasive species provide exceptional study systems for addressing questions about

adaptation, including its predictability, rate, and genetic basis (13, 14). For example, shared

environmental factors between the native and introduced ranges of a species can favour parallel

local adaptation between ranges ("parallelism"), resulting in geographically similar patterns of

spatial divergence along temperature, aridity, or other spatial gradients in the environment (e.g.,

(15–17)). In other cases, evolutionary contingency or environmental differences between the

ranges can promote evolutionary divergence between native versus introduced populations

(e.g., (18, 19)). In addition, the age of an introduction constrains the time available for

adaptation. In some cases, rapid evolutionary diversification across the new range may keep

pace with the range expansion itself, leading to substantial local adaptation within decades of

the introduction (20, 21). In other cases, introduced populations may experience evolutionary

lags, with protracted periods of suboptimal local adaptation across their introduced ranges (22,

23). Finally, these phenomena may play out differently among trait types (24), which may show

differences in the strength or direction of selection that they experience, or in their genetic

capacities for evolutionary change.

Comparisons of native and introduced populations of the same species offer two major

advantages for evaluating the evolutionary rate and predictability of local adaptation. First,

such studies have a natural, paired design, akin to case-control experiments, in which the

evolution of a common set of traits can be directly compared across the native and introduced

ranges (25). Such comparisons are particularly powerful when carried out using individuals

reared in common garden environments (24), which allows evolved differences among

populations to be isolated from environmental factors. One can then directly contrast the

direction and steepness of geographic clines across the introduced range, whose populations

are relatively young and potentially maladapted to their local conditions, with the native range,

whose populations have had extensive time to evolve and are presumably better adapted to

local conditions (26, 27). Second, dates of introduction are often well-documented in historical

records, which provides opportunities to quantitatively test whether indices of local adaptation

(*e.g.*, clines in introduced ranges and parallelism with native range clines) become stronger as

the time since the introduction increases (28, 29). If selection favors similar trait–environment

relationships in both ranges, older introductions should exhibit greater parallelism than younger

introductions (30).

Studies of local adaptation in native versus introduced populations of the same species

have progressed to a point where we now have many high-quality case studies spanning a

diverse range of plant taxa (24). These studies, if analysed in a common framework, provide

an opportunity to test hypotheses about local adaptation, including its degree of parallelism

between native and introduced portions of species' ranges, and the tempo with which it evolves

following introductions into new ranges. With this goal in mind, we conducted a systematic

search for studies of local adaptation in plants and identified 41 studies, spanning 31 species,

that report cline estimates for a total of 586 traits in both the native and introduced portions of the species' range. Of these studies, 34 were conducted in common garden experiments, allowing estimates of genetic clines across populations sourced from different regions of the species' range. Our dataset expands upon earlier meta-analyses of trait clines in native versus introduced plant populations by incorporating studies published over the last 15 years (Colautti and Lau (24) include studies published up to 2011).

We begin with a brief theoretical overview of quantitative predictions for the evolution of clines in native and introduced populations of a species, including the effects of time since introduction, environmental differences between ranges, and evolutionary constraints, on degrees of evolutionary parallelism. We then carry out a detailed analysis of published cline data to address three questions: (1) How predictable is local adaptation, as implied by evolutionary parallelism of trait clines between native and introduced ranges? (2) How rapid is the evolution of local adaptation, as implied by the interaction between the time since introduction and the degree of parallelism between introduced and native populations? (3) To what extent do different trait categories (e.g., morphological, phenological, and defense traits) vary in their degree of parallelism or in their evolutionary lags in introduced ranges?

Trait clines in native versus introduced ranges: Theoretical predictions

Our interpretation of the meta-analysis data relies upon predictions of cline theory from evolutionary quantitative genetics (*e.g.*, (31)), which we outline here and substantially expand upon in the Supplementary Information (SI Appendix 1). We focus here on simple scenarios in which the phenotypic optimum for each trait changes linearly across geographic gradients and local selection is strong relative to drift and gene flow, resulting in strong local adaptation at migration-selection equilibrium in each range. These simplifying assumptions should be viewed as (at best) rough approximations of clinal divergence in real populations.

Consequences of these simplifying assumptions and their violation are briefly discussed below

and more thoroughly explored in the SI Appendix 1.

With strong selection relative to gene flow, the evolution of trait clines depends upon

the rate at which trait optima change across environmental gradients within each range and

whether populations have had sufficient time to evolve to equilibrium. Let B_N and B_I,

respectively, represent the spatial rates of change (slopes) of the optima for a single trait across

the native and introduced ranges (trait optima are depicted by the dashed lines in Fig. 1A).

Native range populations are assumed to be at equilibrium, at which trait clines closely track

spatial changes in the optimum (see Fig. 1A, where the slope of the native range cline, b_N,

matches the slope of the optimum: $b_N \approx B_N$). Since introduced populations have had less time

to evolve within the introduced range, their trait clines (b_I) are more likely to deviate from

equilibrium, particularly so immediately following the introduction. The ratio of trait cline

slopes in the introduced relative to the native range, over time, can be approximated as:

151
$$\frac{b_I}{b_N} = \frac{B_I}{B_N}(1 - (1 - h^2\gamma)^t) \tag{1}$$

where h^2 is the trait's heritability, γ is the strength of stabilizing selection around the optimum

within each environment, and t is the number of generations since the introduction event.

Under this simple model, the ratio of introduced to native range cline slopes is nearly

zero shortly after introduction because populations are not yet locally adapted to the new range

(e.g., $b_I/b_N \approx 0$ after $t = 10$ generations in the example from Fig. 1A). Over time, cline slope

ratios converge to $b_I/b_N = B_I/B_N$, providing a measure of the degree of parallelism of local

adaptation between the native and introduced ranges. ***Perfect parallelism*** between ranges is

expected when $b_I/b_N = 1$ at equilibrium. ***Partial parallelism*** is expected when geographic

shifts in the optima occur in the same direction between ranges but differ in magnitude

($b_I/b_N > 0$ and $b_I/b_N \neq 1$ at equilibrium). ***Antiparallelism*** arises when trait optima shift in

opposite directions between ranges ($b_I/b_N < 0$ at equilibrium). In this model, a strong

empirical pattern of parallelism is thus expected when patterns of local selection are similar

between ranges and evolutionary lags in local adaptation are quickly resolved. Deviations from

perfect parallelism can arise from several factors, including evolutionary lags in local

adaptation (which may be strongly influenced by invasion history, including multiple

introductions and bottlenecks), divergent patterns of selection between ranges, and differences

between ranges in the effects of genetic or demographic constraints to local adaptation (such

constraints are explored in the SI Appendix 1; also see: (24, 31, 32)).

Organisms are comprised of many traits that collectively respond to selection, leading

to multivariate clinal divergence. Multivariate parallelism between the native and introduced

ranges of a species can be quantified by comparing vectors of trait cline slopes from each range.

Let $\boldsymbol{b_I} = \left(b_{I,1}, b_{I,2}, \ldots, b_{I,n}\right)$ and $\boldsymbol{b_N} = \left(b_{N,1}, b_{N,2}, \ldots, b_{N,n}\right)$ represent the introduced and native

trait cline slopes for the same set of n traits. A measure of parallelism between the ranges is:

$$R_P = \frac{\|\boldsymbol{b_I}\|}{\|\boldsymbol{b_N}\|} \cos(\theta) \tag{2}$$

(see Fig. 1B), where the Euclidean norms of the vectors ($\|\boldsymbol{b_I}\|$ and $\|\boldsymbol{b_N}\|$) represent the

magnitudes of clinal divergence in each range, θ is the angle between the vectors, and $\cos(\theta)$

represents the correlation (hereafter the "alignment") of cline slopes between native and

introduced ranges (also see (33, 34)). Greater Euclidean norms indicate stronger overall clinal

divergence across all traits combined, as steeper slopes for any individual trait increase the total

magnitude of divergence. In contrast, larger $\cos(\theta)$ indicates that the directions and relative

magnitudes of trait slopes are similar between native and introduced ranges. $R_P = 1$ when there

is perfect parallelism of local adaptation between ranges (i.e., $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\| = \cos(\theta) = 1$).

Deviations from $R_P = 1$ occur when there is a lag in local adaptation across the introduced

range or selection differs between the ranges, leading to differences in the cline magnitudes

between ranges ($\|\boldsymbol{b}_N\| \neq \|\boldsymbol{b}_I\|$) and misaligned orientations of cline vectors ($\cos(\theta) < 1$).

The cline estimates included in our meta-analysis are not time-series data, but

contemporary snapshots of a clinal divergence in taxa with different residence times within

their introduced ranges. To illustrate how evolutionary lags and differences in selection

between ranges might influence measures of parallelism in the meta-analysis data, we

simulated ratios of introduced and native trait clines (see Fig. 1C-D) across broad distributions

of introduction times among species and $h^2\gamma$ values among traits, assuming partial alignment

of local selection between ranges (B_N and B_I values were drawn from identical marginal

distributions with a modest correlation of 0.5). The simulations ignore effects of estimation

error (a feature of real data), invasion-history, genetic drift, and differences between ranges in

the strength of genetic or demographic constraints to local adaptation (see the SI Appendix 1

and Discussion). These omissions should result in faster adaptation and stronger parallelism in

the simulations than might occur in nature.

Overall, the theory predicts that clines should be shallow in young relative to old

introductions, with parallelism between the ranges initially weak ($b_I/b_N < 1$, on average, for

small t) but increasing in strength over time ($b_I/b_N = 1$, on average, with large t). Variation of

b_I/b_N among traits may, nevertheless, persist (Fig. 1C-D) if traits show variation in the ratio

of invasive to native slopes of the trait optima (i.e., if B_I/B_N varies among traits). In multivariate

space, parallelism (R_p) is predicted to be weak with young introductions because the magnitude

of clinal divergence in the introduced range and its alignment with divergence in the native

range are both initially low (Figure A1 in SI Appendix 1). Both factors are predicted to increase

with residence time in the introduced range, leading to an increase in parallelism over time,

though persistent misalignment in older introductions could indicate sustained selection

differences between ranges and/or constraints to local adaptation.

Figure 1. Outline of theoretical predictions for the meta-analysis. **Panel A:** Hypothetical predictions for equilibrium trait clines in the native (black) and introduced populations (red), relative to an introduced cline prior to equilibrium (shortly after introduction). 't' represents the number of generations. Results are based on the uniform population density model presented in the SI Appendix 1(based on (31)), which also presents extensions of the theory that relax some of the simplifying assumptions that apply to the models presented in the main text (*i.e.*, eq. (1) and the surrounding text). **Panel B:** Metrics of multivariate "parallelism" of trait cline slopes. The black arrow represents the magnitude of native cline vector, which is scaled to one. The red arrow shows the magnitude and the direction of the introduced cline vector relative to the native vector. The purple arrow shows the overall degree of parallelism. **Panels C-D:** Simulated data on cline slope ratios for different traits and species with different amounts of time since an introduction into a new range. Simulations of trait cline slopes are based on eq. (1), with $\gamma = 0.1$, h^2 values sampled from beta distribution with a mean of 0.5 and variance of 0.023 (which is compatible with heritability estimates (35), and B_N and B_I values drawn from a bivariate normal distribution with $\text{corr}(B_I, B_N) = 0.5$, and marginal means and variances of $\bar{B}_I = \bar{B}_N = 5$ and $\text{var}(B_I) = \text{var}(B_N) = 1$. **Panel C** shows the univariate parallelism metric (b_I/b_N, as in Panel B) versus time since introduction. **Panel D** shows the distributions of b_I/b_N among traits after 10 and 100 generations of evolution in the introduced range.

Methods

Meta-analysis data

We compiled the meta-analysis data following PRISMA guidelines ((36); see SI Appendix 2, Fig. S1), and included studies reporting trait clines in both the native and introduced ranges of the study species. Eligible articles were identified from searches of Web of Science, Scopus, and PubMed (initial searches were carried out in December 2022 and repeated in May 2025). Keywords were a combination of synonyms related to clines (e.g., "clinal variation"), introduced species (e.g., "introduced"), plant (e.g., "weed"), and the term "adapt*" (see SI Appendix 2, Fig. S1 for details). We included studies using common garden experiments and field surveys published in academic journals with titles and abstracts in English. Studies cited in identified articles were added to our database if they met the inclusion criteria.

When possible, we obtained raw data published online or provided by authors. When raw data were not available, we extracted mean trait values and standard errors from figures using ImageJ (37). Studies with less than four populations were excluded from the analysis. We also excluded studies focusing on 'expanded' ranges, where the native range has recently expanded within the same continent (38). A total of 41 studies met our inclusion criteria. Of those, 29 measured trait variation in a common garden, seven used field surveys, and five used both. One field study (39) examined multiple species within the same localities; we treated each species' dataset as a separate study. Some studies grew plants under multiple common garden treatments (e.g., different water or nutrient regimes) or implemented common garden experiments at different sites. These "experiments" were analysed separately because trait values could not be directly compared across experiments. When multiple introduced ranges were tested against a single native range, we considered each pair of contrasts as a separate "experiment". The final dataset includes paired (native and introduced) cline estimates for 586 traits, from 104 experiments, spanning 31 species and 12 plant families (SI Box S1 and Table

S1). Each trait was classified into one of the following categories (SI Appendix 2, Table S2):

morpho-physiology (structural features and functional processes), reproduction (investment in

offspring production), size (biomass and growth traits), phenology (timing of developmental

stages) and defense (physical and chemical traits that resist herbivory). Species were

distinguished by cultivation status (yes/no) using a published dataset (40).

Estimation of the number of generations since introduction

To estimate species' introduction times, we used the specific year of introduction to the study

area as provided by authors. When that information was not available, we used the first record

of introduction reported by Seebens et al. (41), for countries matching the source populations

of each species in our dataset. When records for the source country were unavailable, we used

records from the closest country on the same continent. For studies involving populations

sampled from multiple countries, we used the average year of the first record across the

countries. Years since introduction were calculated as [year of seed collection (or survey in

field studies) – year of first record]. When seed collection dates were not available, we used

the year when common gardens took place. The number of generations since introduction was

calculated as [years since introduction]/[generation time], with species' generation times (in

years) obtained from Compagnoni et al. (42). When generation time data were not available,

we approximated generation time as $T_z = \alpha + 0.5(\omega - \alpha)$, where α is the species' age at

reproductive maturity and ω is the typical lifespan (43), each obtained from TRY database (44).

With accurate measures of α and ω, T_z provides a good approximation of generation time (45).

Effect size calculations

All statistical analyses were performed with the R software v.4.4.1 (46) .

 Estimation of individual trait cline slopes

For each experiment, we estimated cline slopes using linear models with the population-level

trait means as the response variable and the environmental gradient the authors of the original

studies focused on as the explanatory variable. These environmental gradients included

latitude, principal components of BioClim data, temperature, elevation, aridity, altitude,

longitude and soil electrical conductivity. We also analysed the data using a shared

environmental predictor across the set of experiments (e.g., by absolute latitude, temperature,

and precipitation - the latter two predictors were calculated following Atwater et al. (47)).

Within an experiment, traits were standardised (mean = 0; standard deviation = 1) across both

native and introduced populations before model fitting. Traits with highly skewed distributions

were square root or log transformed prior to standardisation to improve residual normality. For

each trait, population means were used to obtain a native slope (b_N), an introduced slope (b_I),

and their standard errors. In the following analyses, we focus on ratios of cline slopes (i.e.,

b_I/b_N and its multivariate analogue), which facilitate comparisons of effect sizes between the

wide assortment of traits and study systems in the dataset, regardless of the original units of

measurement of the traits or the environmental gradients chosen by the authors.

Multivariate clinal divergence

We carried out multivariate analyses using the 98 experiments with more than one trait. For

each experiment, we compiled vectors of introduced and native trait clines ($\boldsymbol{b_I}$ and $\boldsymbol{b_N}$) and

calculated their cosine similarity, $\cos(\theta)$, as:

308
$$\cos(\theta) = \frac{\sum_{i=1}^{n} b_{I,i} b_{N,i}}{\|\boldsymbol{b_I}\| \cdot \|\boldsymbol{b_N}\|}$$

where $b_{I,i}$ and $b_{N,i}$ are the introduced and native slopes of the ith of n traits in the experiment,

and $\|\boldsymbol{b_I}\| = \sqrt{\sum_i^n b_{I,i}^2}$ and $\|\boldsymbol{b_N}\| = \sqrt{\sum_i^n b_{N,i}^2}$ are the total magnitudes of clinal divergence in

each range (34, 48). Values of $\cos(\theta)$ are analogous to correlation coefficients, with $\cos(\theta) =$

1 implying a perfect positive correlation, $\cos(\theta) = 0$ implying no correlation, $\cos(\theta) = -1$

implying a perfect negative correlation of cline slopes between ranges. Our measure of

multivariate parallelism is $Rp = \|\boldsymbol{b_I}\|\cos(\theta)/\|\boldsymbol{b_N}\|$, as described in eq. (2), with $R_p > 0$

indicating at least partial parallelism between ranges, and $R_p = 1$ with $\cos(\theta) = 1$ indicating

perfect parallelism between ranges.

Uncertainty around the point estimates of $\cos(\theta)$ (alignment of clines), $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$

(relative magnitudes), and R_p (parallelism), was estimated by bootstrapping the original data

by randomly selecting individual samples with replacement. For each bootstrapped sample, we

used population means of trait values to calculate the parameters using the methods described

above. In cases where only population means and standard errors were available (but not the

raw data), we simulated trait means for each population by pseudo-random sampling from a

normal distribution with mean corresponding to the point estimate of the mean for the

population and standard deviation corresponding to its standard error. Each set of simulated

means across populations within a range was used to calculate cline slopes, from which we

recalculated the multivariate parameters. We repeated this process 10,000 times to approximate

confidence intervals for these parameters.

Univariate clinal divergence

To examine cline parallelism for different trait categories, we carried out univariate analyses

using estimates of the b_I/b_N ratio, per trait, using the linear models. Our univariate analyses

focus on traits with statistically significant clines in at least one range (336 of 586 traits in the

dataset), as those without a significant cline in either range are irrelevant to tests of parallelism.

Following Friedrich et al. (49), we analysed the natural logarithm of cline slope ratios,

which facilitates parametric statistical analyses of meta-analysis effect sizes. Logarithms of

negative numbers are undefined, and a substantial proportion of b_I/b_N ratios in the dataset were

negative. We dealt with this issue by first analysing ratios of absolute values ($|b_I/b_N|$), which

quantify the relative magnitudes of clines but ignore similarities/differences in cline directions,

and separately analysing b_I/b_N ratios of traits with clines slopes of the same sign, for which

$\ln(b_I/b_N)$ has a defined value. 95% confidence intervals for the mean of $\ln(|b_I/b_N|)$ and $\ln(b_I/b_N)$

were calculated using equations presented by Friedrich et al. (49). Systematic differences

between native and introduced cline slopes correspond to cases where 95% confidence

intervals of the pooled effect sizes do not overlap with zero.

To test whether the proportion of clines going in the same direction ($b_I/b_N > 0$) changes

with the number of generations since introduction, we built a generalised linear model (*glm*)

with generations-since-introduction as the fixed effect, a binomial error distribution and a logit

link function, and a binary response variable ($0 = [b_I/b_N < 0]$ and $1 = [b_I/b_N > 0]$).

Meta-analysis overview

All meta-analyses were performed using the function *rma.mv* from the package 'metafor' (50).

Parameters in the meta-analytical models were estimated using restricted maximum likelihood

(51), with Knapp-Hartung adjustments used to calculate the confidence interval of pooled

effects (52). We performed multilevel random-effects models to account for the hierarchical

structure of non-independence in our dataset (multiple traits in multiple experiments within

single studies; multiple studies examine the same species). We used the function *var.comp* from

the package 'dmetar' (53) to calculate the total heterogeneity (I^2), and heterogeneity at each

level ($I^2_{level\ 2}$ and $I^2_{level\ 3}$). We tested whether residual heterogeneity was due to sampling error

only using the QE statistic, with significant QE indicating that residual heterogeneity can be

explained by factors not included in the model (50).

The meta-analysis of multivariate parameters estimated pooled effect sizes and 95%

confidence intervals for $\cos(\theta)$, $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$, and R_p, with random effects of study identity

nested within a species. We tested for relationships between multivariate parameters and the

number of generations since introduction by including generations since introduction,

alongside method (common garden experiment vs. field surveys), cultivation status, and

number of traits, as moderators. We generated models with all combinations of these

explanatory variables as main effects, and with pairwise interactions considered only for

generations-since-introduction using the function *glmulti* (package 'glmulti' (54)) and selected

the best model based on Akaike information criterion with small sample size correction (AICc).

Models within ΔAICc < 2 were considered competitive (55), and we compared the model-

averaged importance of each term within the competitive model set to determine the final

model structure.

For the univariate analysis, random effects of experiment identity, nested within the

study identity, accounted for within- and between-study heterogeneity. Species identity was

removed from the models because the multivariate analysis revealed little heterogeneity

explained by species identity. Following the same method as before, we examined how trait

category, method, cultivation status, generations-since-introduction and pairwise interaction

with the latter affected the extent of parallelism between native and introduced ranges. The

package 'orchaRd' was used for graphical purposes (56).

Results

Multivariate analysis of clinal divergence

Across experiments, we found partial, but significant (t_{97} = 4.2, P < 0.0001), parallelism between native and introduced clines (the mean pooled effect size of R_p was 0.28, i.e. 28% of what is expected under perfect parallelism, with 95% CI [0.15, 0.41]; Fig. 2A). The alignment of native and introduced range vectors ($\cos(\theta)$) was positive in 65% (64 of 98) of experiments, indicating that native and introduced populations tend to clinally diverge in similar directions (χ^2 = 4.1, P = 0.04). The mean alignment between native and introduced cline vectors was $\cos(\theta)$ = 0.30 ([0.15, 0.45], t_{97} = 4.1, P < 0.0001), while the ratio of the cline vector magnitudes ($\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$) was 0.73 ([0.60, 0.86], t_{97} = 11.3, P < 0.0001), indicating that introduced populations show relatively strong clines (73% of native cline magnitudes) that are weakly aligned with the native clines. Residual heterogeneity (i.e., QE) was significant, primarily due to study identity in all three metrics (SI Appendix 2, Table S3).

Variation in R_p and $\cos(\theta)$ was best explained by the additive effects of generations since introduction and study method (SI Appendix 2, Table S4). R_p increased with the number of generations since introduction (*slope* = 0.0051, t = 3.7, P = 0.0004, Fig. 2B) and was higher in common gardens than field surveys (t = 3.0, P = 0.004). Similarly, $\cos(\theta)$ increased with introduction time (*slope* = 0.0030, t = 2.7, P = 0.008, Fig. 2C) and was higher in common gardens than field surveys (t = 3.3, P = 0.001). In contrast, none of the moderators we examined, including generations since introduction, explained variation in $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$ (*slope* = 0.0013, t = 1.1, P = 0.28, Fig. 2D). These results indicate that the increase in parallelism with time since introduction is primarily due to increasing alignment of clines rather than changes in their relative magnitudes.

The above results are based on clines associated with the spatial or environmental gradients chosen by the original study's authors (most chose latitude, though some used

predictors like temperature, precipitation, or aridity). To evaluate the robustness of our results,

we repeated the multivariate analyses using re-estimated trait clines predicted by absolute

latitude, or gradients of temperature or precipitation. Results for R_p, $\cos(\theta)$, and $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$

were similar for each predictor of clinal divergence, though clines across precipitation gradients

resulted in overall quantitatively weaker degrees of parallelism relative to other gradients (SI

Appendix 2, Table S5). Associations with time since introduction remained significant for

latitudinal clines, but not for clines predicted by temperature or precipitation gradients (SI

Appendix 2, Table S6).

Figure 2. Multivariate clinal divergence between introduced and native ranges. **(A)** Cline vectors of native and introduced ranges are presented as arrows (black and red, respectively), with the angle between the vectors (θ) indicating their mean degree of alignment across studies. The length of the introduced vector represents the mean magnitude of introduced vector relative to that of the native vector ($\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$), where the native vector magnitude is scaled to one ($\|\boldsymbol{b_N}\| = 1$). Grey shading denotes the 95% CI of θ, and pink shading denotes the 95% CI of $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$. The purple arrow indicates the degree of multivariate parallelism between ranges (R_p), with shading denoting its 95% CI. An arrow length greater than 0 and less than 1 indicates partial parallelism. **Panels B-D** show the associations between the number of generations since introduction and **(B)** R_p, **(C)** $\cos(\theta)$, and **(D)** $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$, with meta-analytic data shown as bubble plots. Asterisks indicate a significant association with time (*** $P < 0.0001$).

Univariate analysis of divergence across different trait categories

Among the 336 traits with a significant cline in at least one range (Fig. 3), 197 (59%) were in the same direction ($b_I/b_N > 0$) and 139 (41%) were opposite ($b_I/b_N < 0$). Native clines were, on average, steeper than introduced clines, with the magnitude of introduced clines being 71% of their native counterparts (mean $\ln(|b_I/b_N|)$ = -0.34, 95% CI = [-0.55, -0.13]). Residual heterogeneity (i.e., QE) was significant, primarily due to study identity (SI Appendix 2, Table S3).

Figure 3. Results of the linear model tests for clinal variation in 586 traits of
the meta-analysis. Slope estimates were (left to right): non-significant in both
ranges, significant in the native range only, significant in the introduced range
only, or significant in both ranges.

Model selection revealed that variability in $\ln(|b_I/b_N|)$ was best explained by 'trait

category' alone (SI Appendix 2, Table S4). The magnitude of clinal divergence was similar

between ranges for defense and reproductive traits, with introduced clines showing 127% and

86% the magnitude of native clines, respectively ($\ln(|b_I/b_N|) = 0.24$ [-0.15, 0.63] and $\ln(|b_I/b_N|)$

$= -0.15$ [-0.43, 0.12]). In contrast, clinal divergence was systematically shallower for morpho-

physiology (51%, $\ln(|b_I/b_N|) = -0.67$ [-1.01, -0.32]), phenology (52%, $\ln(|b_I/b_N|) = -0.65$ [-0.94,

-0.35]), and size traits (55%, $\ln(|b_I/b_N|) = -0.60$ [-0.85, -0.34]).

While the relative magnitude of clinal divergence in each range was not associated with

the time since introduction (*slope* = 0.0007, t = 0.0019, *P* = 0.73; Fig. 4C), the proportion of

clines diverging in the same direction increased with time (*slope* = 0.009, z = 0.002, *P* = 0.0002;

see Fig. 4D). This increase holds qualitatively for each trait category analysed separately,

though interactions with time were only significant for the size-related traits (SI Appendix 2,

Fig. S2), possibly owing to the smaller sample sizes of other categories.

Figure 4. Univariate estimates of parallelism between native and introduced ranges. **Panels A-B** show the absolute ratio of introduced to native slopes in log scale ($\ln(|b_I/b_N|)$), analysed separately for each trait category. **(A)** includes values of all traits, and **(B)** includes only traits whose cline estimates diverge in the same direction ($b_I/b_N > 0$). The mean effect sizes (white circles) are plotted with the 95% confidence intervals (bold line), with the number of traits given in brackets (SI Appendix 2, Table S7). Asterisks indicate significant deviation from $\ln(|b_I/b_N|) = 0$ (perfect parallelism). The individual effect sizes (coloured circles) for five trait categories are plotted with circle size denoting precision (inverse variance). **Panels C-D** show the relationship between the number of generations since introduction and **(C)** cline slope ratios, or **(D)** the probability that clines in each range diverge in the same direction (*i.e., $b_I/b_N > 0$*, analysed using a generalized linear model). Asterisks indicate a significant association with time. * $P < 0.05$, ** < 0.01, *** < 0.0001.

Similar results apply for the 197 traits with cline estimates diverging in the same direction (traits where $b_I/b_N > 0$; Fig. 4B). In these cases, magnitudes of introduced clines were 76% of the magnitudes of native clines ($\ln(b_I/b_N) = $ -0.28 [-0.50, -0.06], SI Appendix 2, Table

S3). Cline slopes were similar between introduced and native ranges for defense traits (80%,

$\ln(b_I/b_N)$ = -0.22 [-0.67, 0.22]) and reproductive traits (99%, $\ln(b_I/b_N)$ = -0.01 [-0.29, 0.27]),

while clines were shallower in introduced ranges for morpho-physiology (59%, $\ln(b_I/b_N)$ = -

0.52 [-0.94, -0.09]), phenology (65%, $\ln(b_I/b_N)$ = -0.43 [-0.74, -0.12]), and size traits (64%,

$\ln(b_I/b_N)$ = -0.45 [-0.73, -0.17]). There was no association between $\ln(b_I/b_N)$ and the number of

generations since introduction (SI Appendix 2, Fig. S3).

Discussion

Our meta-analysis reveals three broad patterns of evolutionary parallelism between the trait

clines of native versus introduced ranges. First, multivariate parallelism (R_p) across the entire

dataset is relatively low, overall—on average, only 28% of the divergence predicted under

perfect parallelism. This low average R_p was due to shallower clinal divergence in introduced

relative to native ranges ($\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$ = 0.73) along with a modest alignment of cline

directions between ranges ($\cos(\theta)$ = 0.30). Second, multivariate parallelism between ranges

markedly increases with the residence time of introduced populations (generations since

introduction), resulting in strong parallelism between ranges involving relatively old

introductions (i.e., $R_p \approx 1$ at 200 generations post introduction; Fig. 2C). This increase in

parallelism over time is primarily caused by increased alignment of cline directions between

ranges rather than changes in relative cline magnitudes. Third, degrees of parallelism between

ranges vary among traits, and are stronger for reproductive and defense traits than for

phenological, morpho-physiological, and size-related traits; the latter three categories exhibit

weaker clines in introduced ranges. Below, we discuss processes that might explain these

observations before outlining opportunities for future work.

Why are trait clines shallower in introduced ranges?

Several evolutionary mechanisms could contribute to shallower clines in introduced relative to

native ranges (*i.e.*, $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\| < 1$). Under simple theoretical models that assume strong

selection and no genetic or demographic constraints (Fig. 1), shallower clines in introduced

ranges can emerge from time lags in local adaptation (57), or shallower underlying

environmental gradients in introduced range (*i.e.*, $|B_I/B_N| < 1$; recall that B_I and B_N refer to

slopes of changes in trait optima). The time-lag hypothesis predicts an increase in $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$

with the residence time of introduced populations—a pattern that we did not observe.

Alternatively, environmental gradients might be shallower in introduced ranges if species are

disproportionately introduced into geographic regions where the environmental gradients are

relatively shallow, which would facilitate range expansions in those regions (58). For example,

urban environments that many invasive plants occupy (59) may provide more homogeneous

landscapes than natural systems (60). Unfortunately, direct comparisons of environmental

gradients between native and introduced ranges are lacking, making it difficult at present to

formally evaluate this hypothesis.

A more complex picture emerges in the presence of genetic, evolutionary, and

demographic factors that constrain local adaptation, potentially leading to shallower clines in

introduced ranges. For example, weaker selection and/or lower genetic variation in introduced

ranges would enhance the swamping effects of gene flow on clinal divergence, resulting in

shallower clines at migration-selection balance. However, heritability estimates for fitness-

related traits (61–65) and linear selection gradients (24) generally have similar magnitudes in

native and introduced ranges, offering little support for this explanation, though such data are

admittedly sparse.

Demographic differences between ranges—including differences in the rate of dispersal

or the spatial distribution of population density (see SI Appendix 1)—can also lead to different

degrees of swamping of local adaptation between ranges. Invasive herbaceous plant species

tend to have higher dispersal rates than non-invasive ones (66), and increased dispersal ability

can evolve within introduced ranges (61, 67), which could lead to higher swamping by gene

flow in introduced ranges relative to native ranges. The distribution of population density

affects local adaptation by mediating the rate of gene flow from the center of a range to its

edges, with 'abundant center' distributions leading to much stronger swamping than under

relatively uniform patterns of population density across a range (31). Introduced populations

that more closely conform to an abundant-center scenario than do their native counterparts

(which might be likely in actively expanding introductions; (68)) will be more susceptible to

swamping.

Strong genetic correlations between traits can further constrain rates of local adaptation

and greatly extend time lags in local adaptation but nevertheless lead to weak associations

between the relative magnitudes of clinal divergence in each range and the time since

introduction (Fig. 5). Figure 5A illustrates one such example, where $\|\boldsymbol{b}_I\|/\|\boldsymbol{b}_N\|$ for two

strongly correlated traits is relatively flat across the first 200 generations of evolution in the

introduced range, though the increase is much more visible over longer time intervals. We

return to this point in the following section.

Figure 5. Evolution of clines for a pair of genetically correlated traits. The plots show a scenario in which the optima for the pair of traits shift in opposite directions, with $B_1 = 2$ and $B_1 = -1$, and the traits are otherwise subject to the same strength of stabilizing selection ($\gamma = 0.05$). Heritability for each trait is $h^2 = 1$ and the genetic correlation between the traits is strongly positive ($\rho = 0.95$). Parameters of selection and variation are assumed to be identical between the native and introduced ranges, so that perfect parallelism ($\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$ and $\cos(\theta) = 1$) is predicted at equilibrium. Panel **(A)** shows the evolution of $\cos(\theta)$ and $\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$ across 1000 generations since the introduction at generation $t = 0$, with the inset highlighting the predictions for the first 200 generations. Panel **(B)** shows the evolution of clines for each trait—including a counter-cline that arises early for trait 2 but ultimately resolves as introduced populations approach migration-selection equilibrium. Details of the model can be found in SI Appendix 1.

The trend towards shallower slopes in introduced ranges does not, however, apply across the board. Defense and reproductive traits have similar cline magnitudes between ranges, while clines for phenological, morpho-physiological, and size-related traits were shallower (roughly half as steep) in introduced relative to native ranges. These differences among trait categories might stem from differences in evolutionary potential for rapid local adaptation, with traits under exceptionally strong selection or with high heritabilities evolving more rapidly in introduced ranges. Reproductive traits, for example, are thought to be under strong selection because they are closely related to fitness (69, 70). Likewise, the average heritability of plant secondary metabolites (which includes defensive compounds) is estimated to be more than twice as large as the heritability of phenological, morphological, physiological, and size traits, which should facilitate responses to selection ((71); see also (61, 72)).

Why does the alignment of cline directions between ranges increase over time?

Four factors potentially contribute to the weaker alignment of cline directions in younger than older introductions (recall that $\cos(\theta)$ quantifies alignment between ranges). First, traits differ in how responsive they are to natural selection: those with high heritability and under strong

selection will rapidly adapt to local conditions across the new range, while those with low

heritability or weak selection should exhibit protracted lags in local adaptation (see the SI

Appendix 1). This heterogeneity of lag-times among traits leads to temporary misalignments

in the relative magnitudes of clines in each range, which eventually resolve as each trait evolves

to migration-selection equilibrium in the new range. However, while this scenario might

contribute to the temporal increase in alignment that we observe, it is unlikely to be the only

explanation, since this scenario also predicts a concurrent, and much larger, increase in

$\|\boldsymbol{b_I}\|/\|\boldsymbol{b_N}\|$, which we do not observe (see Fig. A1, SI Appendix 1), nor can it account for the

high proportion of counter-clines in young introductions.

Second, in the early stages of invasion, stochastic processes can play important roles in

the initial formation of clines. Founder effects and genetic drift during range expansion can

produce clines in random directions (24), many of which will not align with those in the native

range. Similarly, spatial sorting of dispersal-related traits during expansion, documented in

both plants and animals (see (73) for review), can create gradients that are potentially correlated

with environmental variables but primarily reflect colonisation history rather than local

adaptation. Furthermore, multiple introductions from genetically divergent source populations,

or hybridisation, could generate irregular or even counter-clines, particularly in younger

invasions where admixture between lineages is ongoing. If selection pressures are similar

between ranges, these stochastic and historical effects on the initial formation of clines are

expected to diminish over time, as natural selection reorients clines to align with those

maintained in the native range, leading to the observed increase in parallelism among the older

introductions.

Third, the directions of selection in each range might change over time, leading to a

greater alignment of both selection and trait clines in old relative to young introductions. For

example, selection may fundamentally differ between young and old introductions if the former

experiences a release from natural enemies, but the latter does not. Enemy release might relax

selection of defense traits within the new range and facilitate evolution of defense trait clines

caused by drift or by indirect responses to selection of genetically correlated traits (as predicted

by the "evolution of increased competitive ability" (EICA) hypothesis (74)). Moreover,

relaxation of herbivory may alter the pattern of selection on secondary metabolites with dual

functions in herbivore defense and abiotic stress tolerance (28), with metabolite concentrations

in recently introduced species evolving in response to selection for abiotic stress tolerance

rather than herbivore defense (28). Clines favoured by selection in young introductions may,

thus, differ substantially from those in the native range. However, subsequent accumulation of

natural enemies in the introduced range over time (75, 76) may gradually reshape the selective

landscapes of older introduced ranges, leading to a stronger alignment of selection and

evolutionary parallelism of clines between native and introduced ranges.

Finally, strong genetic correlations between traits can both constrain the rate of local

adaptation in introduced ranges and lead to transient counter-clines in traits subject to relatively

weak selection (Fig. 5B). Under such conditions, $\|\boldsymbol{b}_I\|/\|\boldsymbol{b}_N\|$ for a set of traits can rapidly

increase soon after the introduction and then remain evolutionarily stagnant due to genetic

constraints to further local adaptation of each trait, whereas the alignment of the clines from

each range steadily increases over the same time interval (see the inset of Fig. 5A). Strong

parallelism of clines (e.g., $\|\boldsymbol{b}_I\|/\|\boldsymbol{b}_N\| \approx \cos(\theta) \approx 1$) can ultimately evolve between the

native and introduced ranges, yet strong genetic constraints can draw out this process and,

depending on the time window of the observation, reveal a pattern in which $\|\boldsymbol{b}_I\|/\|\boldsymbol{b}_N\|$ is

relatively constant over time while $\cos(\theta)$ steadily increases, as found in our meta-analysis.

Strong genetic correlations and trade-offs between fitness-related traits may indeed be likely.

For example, both early flowering and large biomass (and seed production) are favoured at

high latitudes, yet these traits negatively covary, providing the potential for the evolution of

transient counter-clines in the more weakly selected trait (see (22, 77)).

Conclusions and future directions

A trait or allele frequency cline is not, in itself, evidence for local adaptation (24). However,

repeated evolution of parallel clines between geographically disjunct regions of a species' range

is difficult to reconcile with nonadaptive explanations and is instead indicative of repeated local

adaptation to shared environmental conditions between the regions (78, 79). We found that

geographic clines in plant traits showed strong evolutionary parallelism between native and

introduced ranges with relatively long residence times (on the order of 100 or more

generations). This result applies to a broad set of plant species and trait types and implies that

local adaptation is extensive and predictable across native ranges and mature introduced ranges.

More surprisingly, our analysis suggests that the greater parallelism of older relative to

younger introductions is primarily due to an increase in the alignment of cline directions

between ranges rather than changes to the relative magnitudes of clines in introduced versus

native ranges—a pattern that holds qualitatively across different trait categories. Although

several factors may contribute to this pattern, we suspect it is caused by rapid evolution of

maladaptive clines for a subset of traits in young introductions—owing to processes like drift,

founder effects, and spatial sorting during colonization, or to indirect responses to selection of

genetically correlated traits. This is followed, in the longer term, by more gradual evolution of

local adaptation that results in increased parallelism between clines in each range. Direct tests

for enrichment of maladaptive clines in young relative to old introductions (e.g., through

estimates of selection of clinally diverging traits) would, therefore, be a worthwhile focus for

future work. Other potentially useful extensions of our study include analyses of nonlinear

clines (which are sometimes reported in cline studies, e.g., (15, 80)) and their evolutionary

parallelism between ranges, and analyses of the joint tempo of changes in trait clines and the

geographic breadths of introduced species' ranges, both of which may increase over time, with

adaptive clines possibly facilitating expansions of range breadth (58).

Finally, we used a snapshot of clinal divergence in species that differed in their arrival

times into new ranges to indirectly infer temporal dynamics of cline evolution. However, a true

time-series analysis of evolutionary parallelism between native and introduced clines would

also be valuable, though such data are sparse. In a rare exception, Wu and Colautti (22)

analysed herbarium records to compare clinal patterns of flowering phenology in three

introduced ranges of *Lythrum salicaria* over 150 years, and found lags in the evolution of

adaptive clines in all three regions. Because herbarium specimens are collected in the field, the

contribution of phenotypic plasticity to the trait variation must be controlled (22), although

genomic analyses (30) or quantitative genetic approaches can be leveraged to overcome those

issues (81). In our meta-analysis, we found that multivariate parallelism was weaker in field

studies compared to common garden experiments (though no such differences were observed

in the univariate analysis), which suggests that herbarium-based phenotypic cline estimates

could be valuable for inferring lower bounds for the evolution of parallelism between ranges.

References

1. J. Arendt, D. Reznick, Convergence and parallelism reconsidered: what have we learned
about the genetics of adaptation? *Trends Ecol Evol* **23**, 26–32 (2008).

2. D. L. Stern, The genetic causes of convergent evolution. *Nat Rev Genet* **14**, 751–764
(2013).

3. S. J. Gould, *Wonderful life: the Burgess Shale and the nature of history* (WW Norton &
Company, 1989).

4. P. W. Messer, S. P. Ellner, N. G. Hairston, Can population genetics adapt to rapid
evolution? *Trends Genet* **32**, 408–418 (2016).

5.   A. A. Agrawal, A. P. Hastings, M. T. J. Johnson, J. L. Maron, J.-P. Salminen, Insect
676   herbivores drive real-time ecological and evolutionary change in plant populations.
677   *Science* **338**, 113–116 (2012).

678   6.   R. Gomulkiewicz, D. Houle, Demographic and genetic constraints on evolution. *Am Nat*
679   **174**, E218–E229 (2009).

680   7.   J. Maynard Smith, What determines the rate of evolution? *Am Nat* **110**, 331–338 (1976).

681   8.   J. Polechová, Evolution of species' range and niche in changing environments. *bioRxiv*
682   2025–01 (2025). https://doi.org/10.1011/2025.01.16.633367.

683   9.   J. Polechová, Is the sky the limit? On the expansion threshold of a species' range. *PLoS*
684   *Biol* **16**, e2005372 (2018).

685   10.   M. Kirkpatrick, N. H. Barton, Evolution of a species' range. *Am Nat* **150**, 1–23 (1997).

686   11.   C. M. Caruso, *et al.*, What are the environmental determinants of phenotypic selection?
687   A meta-analysis of experimental studies. *Am Nat* **190**, 363–376 (2017).

688   12.   A. M. Siepielski, *et al.*, Precipitation drives global variation in natural selection. *Science*
689   **355**, 959–962 (2017).

690   13.   P. J. Prentis, J. R. Wilson, E. E. Dormontt, D. M. Richardson, A. J. Lowe, Adaptive
691   evolution in invasive species. *Trends Plant Sci* **13**, 288–294 (2008).

692   14.   K. A. Hodgins, P. Battlay, D. G. Bock, The genomic secrets of invasive plants.
693   *New Phytol* **245**, 1846–1863 (2025).

694   15.   L. A. van Boheemen, D. Z. Atwater, K. A. Hodgins, Rapid and repeated local adaptation
695   to climate in an invasive plant. *New Phytol* **222**, 614–627 (2019).

696   16.   J. L. Maron, M. Vilà, R. Bommarco, S. Elmendorf, P. Beardsley, Rapid evolution of an
697   invasive plant. *Ecol Monogr* **74**, 261–280 (2004).

698   17.   J. R. Etterson, *et al.*, Parallel patterns of clinal variation in *Solidago altissima* in its
699   native range in central USA and its invasive range in Japan. *Botany* **86**, 91–97 (2008).

700   18.   W. Liu, X. Chen, J. Wang, Y. Zhang, Does the effect of flowering time on biomass
701   allocation across latitudes differ between invasive and native salt marsh grass *Spartina*
702   *alterniflora*? *Ecol Evol* **12**, e8681 (2022).

703   19.   W.-W. Liu, Y.-H. Zhang, Geographical variation in germination traits of the salt-marsh
704   cordgrass *Spartina alterniflora* in its invasive and native ranges. *J Plant Ecol* **14**, 348–
705   360 (2021).

706   20.   F. Colomer-Ventura, *et al.*, Contemporary evolution of an invasive plant is associated
707   with climate but not with herbivory. *Funct Ecol* **29**, 1475–1485 (2015).

708   21.   J. M. Alexander, P. J. Edwards, M. Poll, C. G. Parks, H. Dietz, Establishment of parallel
709   altitudinal clines in traits of native and introduced forbs. *Ecology* **90**, 612–622 (2009).

22.   Y. Wu, R. I. Colautti, Evidence for continent-wide convergent evolution and stasis
throughout 150 y of a biological invasion. *Proc Natl Acad Sci U S A* **119**, e2107584119
(2022).

23.   S. G. Innes, J. S. Santangelo, N. J. Kooyers, K. M. Olsen, M. T. J. Johnson, Evolution in
response to climate in the native and introduced ranges of a globally distributed plant.
Evolution **76**, 1495–1511 (2022).

24.   R. I. Colautti, J. A. Lau, Contemporary evolution during invasion: evidence for
differentiation, natural selection, and local adaptation. *Mol Ecol* **24**, 1999–2017 (2015).

25.   O. Bossdorf, *et al.*, Phenotypic and genetic differentiation between native and
introduced plant populations. *Oecologia* **144**, 1–11 (2005).

26.   E. C. Woods, S. E. Sultan, Post-introduction evolution of a rapid life-history strategy in
a newly invasive plant. *Ecology* **103**, e3803 (2022).

27.   Y. Yang, *et al.*, Rapid evolution of latitudinal clines in growth and defence of an
invasive weed. *New Phytol* **230**, 845–856 (2021).

28.   S. G. Innes, J. S. Santangelo, N. J. Kooyers, K. M. Olsen, M. T. J. Johnson, Evolution in
response to climate in the native and introduced ranges of a globally distributed plant.
Evolution **76**, 1495–1511 (2022).

29.   P. A. H. Westley, What invasive species reveal about the rate and form of contemporary
phenotypic change in nature. *Am Nat* **177**, 496–509 (2011).

30.   P. Battlay, *et al.*, Large haploblocks underlie rapid adaptation in the invasive weed
Ambrosia artemisiifolia. *Nat Commun* **14**, 1717 (2023).

31.   G. García-Ramos, M. Kirkpatrick, Genetic models of adaptation and gene flow in
peripheral populations. *Evolution* **51**, 21–28 (1997).

32.   A. Duputié, F. Massol, I. Chuine, M. Kirkpatrick, O. Ronce, How do genetic
correlations affect species range shifts in a changing environment? *Ecol Lett* **15**, 251–
259 (2012).

33.   T. F. Hansen, D. Houle, Measuring and comparing evolvability and constraint in
multivariate characters. *J Evol Biol* **21**, 1201–1219 (2008).

34.   Y. E. Stuart, *et al.*, Contrasting effects of environment and genetics generate a
continuum of parallel evolution. *Nat Ecol Evol* **1**, 158 (2017).

35.   T. A. Mousseau, D. A. Roff, Natural selection and the heritability of fitness components.
Heredity **59**, 181–197 (1987).

36.   M. J. Page, *et al.*, The PRISMA 2020 statement: an updated guideline for reporting
systematic reviews. *BMJ* **372**, n71 (2021).

37.   C. A. Schneider, W. S. Rasband, K. W. Eliceiri, NIH Image to ImageJ: 25 years of image
analysis. *Nat Methods* **9**, 671–675 (2012).

38. F. Essl, *et al.*, A conceptual framework for range-expanding species that track human-
induced environmental change. *BioScience* **69**, 908–919 (2019).

39. J. M. Alexander, P. J. Edwards, M. Poll, C. G. Parks, H. Dietz, Establishment of parallel
altitudinal clines in traits of native and introduced forbs. *Ecology* **90**, 612–622 (2009).

40. D. Z. Atwater, C. Ervine, J. N. Barney, Climatic niche shifts are common in introduced
plants. *Nat Ecol Evol* **2**, 34–43 (2018).

41. H. Seebens, *et al.*, No saturation in the accumulation of alien species worldwide. *Nat
Commun* **8**, 14435 (2017).

42. A. Compagnoni, *et al.*, Herbaceous perennial plants with short generation time have
stronger responses to climate anomalies than those with longer generation time. *Nat
Commun* **12**, 1824 (2021).

43. M. Pacifici, *et al.*, Generation length for mammals. *Nat Conserv* **5**, 89–94 (2013).

44. J. Kattge, *et al.*, TRY plant trait database – enhanced coverage and open access. *Glob
Chang Biol* **26**, 119–188 (2020).

45. J. Staerk, *et al.*, Performance of generation time approximations for extinction risk
assessments. *J Appl Ecol* **56**, 1436–1446 (2019).

46. R Core Team, R: A Language and Environment for Statistical Computing. R Foundation
for Statistical Computing, Vienna, Austria. (2024). Deposited 2024.

47. D. Z. Atwater, C. Ervine, J. N. Barney, Climatic niche shifts are common in introduced
plants. *Nat Ecol Evol* **2**, 34–43 (2018).

48. C. Carvalho, *et al.*, Multivariate selection mediated by aridity predicts divergence of
drought-resistant traits along natural aridity gradients of an invasive weed. *New Phytol*
234, 1088–1100 (2022).

49. J. O. Friedrich, N. K. Adhikari, J. Beyene, The ratio of means method as an alternative
to mean differences for analyzing continuous outcome variables in meta-analysis: a
simulation study. *BMC Med Res Methodol* **8**, 32 (2008).

50. W. Viechtbauer, Conducting meta-analyses in R with the metafor package. *J Stat Softw*
36, 1–48 (2010).

51. A. F. Zuur, E. N. Ieno, N. J. Walker, A. A. Saveliev, G. M. Smith, "Mixed Effects
Modelling for Nested Data" in *Mixed Effects Models and Extensions in Ecology with R*,
A. F. Zuur, E. N. Ieno, N. Walker, A. A. Saveliev, G. M. Smith, Eds. (Springer, 2009),
pp. 101–142.

52. G. Knapp, J. Hartung, Improved tests for a random effects meta-regression with a single
covariate. *Stat Med* **22**, 2693–2710 (2003).

53. M. Harrer, P. Cuijpers, T. Furukawa, D. D. Ebert, dmetar: companion R package for the
guide "Doing meta-analysis in R." *R package version 0.0* **9000**, 2019 (2019).

54. V. Calcagno, C. de Mazancourt, glmulti: an R package for easy automated model
selection with (generalized) linear models. *J Stat Softw* **34**, 1–29 (2010).

55. K. P. Burnham, D. R. Anderson, Multimodel inference: understanding AIC and BIC in
model selection. *Sociol Methods Res* **33**, 261–304 (2004).

56. S. Nakagawa, *et al.*, orchaRd 2.0: An R package for visualising meta-analyses with
orchard plots. *Methods Ecol Evol* **14**, 2003–2010 (2023).

57. J. A. Crooks, Lag times and exotic species: The ecology and management of biological
invasions in slow-motion1. *Ecoscience* **12**, 316–329 (2005).

58. J. Polechová, N. H. Barton, Limits to adaptation along environmental gradients. *Proc
Natl Acad Sci U S A* **112**, 6401–6406 (2015).

59. K. Štajerová, P. Šmilauer, J. Brůna, P. Pyšek, Distribution of invasive plants in urban
environment is strongly spatially structured. *Landsc Ecol* **32**, 681–692 (2017).

60. P. M. Groffman, *et al.*, Ecological homogenization of urban USA. *Front Ecol Environ*
12, 74–81 (2014).

61. L. C. Rigby, M. D. Hall, K. Monro, A. Uesugi, Evolution of "invasion syndrome" in
invasive goldenrod is not constrained by genetic trade-offs. *Evol Appl* **17**, e13734
(2024).

62. B. Calsbeek, S. Lavergne, M. Patel, J. Molofsky, Comparing the genetic architecture
and potential response to selection of invasive and native populations of reed canary
grass. *Evol Appl* **4**, 726–735 (2011).

63. S. J. Franks, G. S. Wheeler, C. Goodnight, Genetic variation and evolution of secondary
compounds in native and introduced populations of the invasive plant *Melaleuca
quinquenervia*. *Evolution* **66**, 1398–1412 (2012).

64. B. V. McGoey, J. R. Stinchcombe, Introduced populations of ragweed show as much
evolutionary potential as native populations. *Evol Appl* **14**, 1436–1449 (2021).

65. Y. Sakata, *et al.*, Environmentally triggered variability in the genetic variance–
covariance of herbivory resistance of an exotic plant *Solidago altissima*. *Ecol Evol* **10**,
3103–3111 (2020).

66. B. Zhang, A. Hastings, E. D. Grosholz, L. Zhai, The comparison of dispersal rate
between invasive and native species varied by plant life form and functional traits. *Mov
Ecol* **11**, 73 (2023).

67. B. L. Phillips, G. P. Brown, R. Shine, Evolutionarily accelerated invasions: the rate of
dispersal evolves upwards during the range advance of cane toads. *J Evol Biol* **23**,
2595–2601 (2010).

68. Q. Guo, Central-marginal population dynamics in species invasions. *Front Ecol Evol* **2**,
23 (2014).

69. R. E. Irwin, P. S. Warren, L. S. Adler, Phenotypic selection on floral traits in an urban
landscape. *Proc R Soc B Biol Sci* **285**, 20181239 (2018).

70. Ø. Totland, Environment-dependent pollen limitation and selection on floral traits in an
alpine species. *Ecology* **82**, 2233–2244 (2001).

71. M. A. Geber, L. R. Griffen, Inheritance and natural selection on functional traits. *Int J
Plant Sci* **164**, S21–S42 (2003).

72. M. T. Johnson, A. A. Agrawal, J. L. Maron, J.-P. Salminen, Heritability, covariation and
natural selection on 24 traits of common evening primrose (*Oenothera biennis*) from a
field experiment. *J Evol Biol* **22**, 1295–1307 (2009).

73. K. A. Hodgins, D. G. Bock, L. H. Rieseberg, Trait evolution in invasive species. *Annual
Plant Reviews Online* **1**, 459–496 (2018).

74. B. Blossey, R. Notzold, Evolution of increased competitive ability in invasive
nonindigenous plants: a hypothesis. *J Ecol* **83**, 887–889 (1995).

75. A. R. Zangerl, M. C. Stanley, M. R. Berenbaum, Selection for chemical trait remixing in
an invasive weed after reassociation with a coevolved specialist. *Proc Natl Acad Sci U S
A* **105**, 4547–4552 (2008).

76. Y. Sakata, M. Yamasaki, Y. Isagi, T. Ohgushi, An exotic herbivorous insect drives the
evolution of resistance in the exotic perennial herb *Solidago altissima*. *Ecology* **95**,
2569–2578 (2014).

77. R. I. Colautti, S. C. H. Barrett, Rapid adaptation to climate facilitates range expansion of
an invasive plant. *Science* **342**, 364–366 (2013).

78. J. G. Oakeshott, *et al.*, Alcohol dehydrogenase and glycerol-3-phosphate dehydrogenase
clines in *Drosophila melanogaster* on different continents. *Evolution* **36**, 86–96 (1982).

79. J. R. Adrion, M. W. Hahn, B. S. Cooper, Revisiting classic clines in *Drosophila
melanogaster* in the age of genomics. *Trends Genet* **31**, 434–444 (2015).

80. A. A. Hoffmann, A. R. Weeks, Climatic selection on genes and traits after a 100 year-old
invasion: a critical look at the temperate-tropical clines in *Drosophila melanogaster*
from eastern Australia. *Genetica* **129**, 133–147 (2007).

81. A. S. Kim, *et al.*, Temporal collections to study invasion biology. *Mol Ecol* **32**, 6729–
6742 (2023).