**Mitochondrial fission regulates ROS for ventral furrow formation in *Drosophila* gastrulation**

Somya Madan[1], Sayali Chowdhary[1,3], Sakshi Phalke[1], Harsh Mittal[1], Manos Mavrakis[2] and Richa Rikhy[1]*

1. Indian Institute of Science Education and Research Pune, Dr. Homi Bhabha Road, Pune 411008, India
2. Institut Fresnel, CNRS UMR7249, Aix Marseille Univ, Centrale Med, 13013 Marseille, France
3. Present address, Cardiovascular Research Institute, Department of Cell and Developmental Biology, Weill Cornell Medicine, 1300 York Avenue, New York, NY, 10021, USA

* To whom correspondence is addressed: richa@iiserpune.ac.in, https://orcid.org/0000-0002-4262-0238

**Highlights**

- Mitochondria migrate apically in constricting areas of ventral cells in gastrulation
- Microtubules and Dorsal/NFkB regulate apical mitochondrial migration in ventral cells
- Drp1 mutants with large, less active, basal mitochondria show ventral furrow defects
- Ventral furrow defects in Drp1 mutants are rescued by increasing mitochondrial ROS

**Summary**

Mitochondrial fusion and fission are essential for embryogenesis, but the mechanisms by which they regulate morphogenesis are not entirely understood. In this study, we elucidate the role of mitochondrial fission in *Drosophila* gastrulation. We find that fragmented mitochondria migrate apically in constricting ventral cells in a microtubule and Dorsal/NFkb dependent manner. Reduced mitochondrial fission in Drp1 mutant

embryos leads to accumulation of larger mitochondria basally and fewer apical

mitochondria. Drp1 mutants show decreased Myosin II, resulting in defective ventral

furrow formation. Transcriptomic analysis from Drp1 mutants shows no changes in the

Dorsal signaling pathway. Proteomic analysis reveals increased levels of antioxidant

enzymes and decreased respiratory complexes, resulting in low ROS levels. Depleting

SOD2 in Drp1 mutants restores ROS levels, allowing apical mitochondrial migration,

Myosin II increase and proper ventral furrow formation. Our study reveals a significant

role for mitochondrial dynamics and activity for apical constriction during gastrulation.

**Introduction**

Embryonic development is elaborate and dynamic, involving various morphogenetic

processes and cellular remodeling events such as cell division, migration, tissue folding

and invagination. These intricate cellular movements occur due to feedback between

signaling pathways and organelle function. They are energy-intensive and require

specific metabolites for appropriate activation. Mitochondria, the powerhouse of the cell,

are the source of energy and metabolites essential for supporting the dynamic

morphogenetic events during embryogenesis [1–3]. Apart from ATP production,

mitochondria have multiple other functions, such as redox balance, calcium buffering

and epigenetic regulation through which they can modulate signaling pathways and

cytoskeletal architecture. Mitochondria are well known to be integral to morphogenetic

processes such as epithelial to mesenchymal transition [4], cell division [5–7], cell migration

[8–10] and wound healing [11,12]. Mitochondrial distribution to sites of active cellular

dynamics and local release of metabolites aids in driving cell shape changes.

Mitochondria are maternally inherited in most organisms; hence, oocyte

mitochondrial health is a crucial determinant of proper embryonic growth and

development [13]. Oocyte and early embryo mitochondria are small and spherical with low

ATP production and oxygen consumption [2,14–17]. In humans, it has been shown that

oocytes with mitochondria exhibiting low electron transport chain (ETC) activity can lead

to defective embryonic development [18,19]. Disruption of mitochondrial membrane

potential and ETC complexes causes developmental arrest in zebrafish [20].

Mitochondria are dynamic and undergo regulated cycles of fusion and fission,
resulting in varied morphologies that range from globular to intricate reticular networks.
The Dynamin superfamily of large GTPases carries out mitochondrial fusion and
fragmentation. Dynamin-related protein 1 (Drp1) functions in mitochondrial fission, by
assembling around the mitochondrion, constricting it and separating it into two daughter
mitochondria via its GTPase activity [21–23]. Mitochondrial fusion is carried out by
Mitofusins 1 and 2 (Mfn1 and 2) and Optic atrophy 1 (Opa1). Mfn1 and 2 fuse the outer
mitochondrial membrane, and Opa1 is responsible for inner mitochondrial membrane
fusion [24–26]. Mitochondrial fission enables inheritance of functional mitochondria into the
oocyte[27]. Mitochondrial fission and fusion play a crucial role in embryonic development.
Mfn1/2 mutant mouse embryos are embryonic lethal [24]. Loss of Opa1 in mouse
embryos also leads to retarded growth and embryonic lethality [28]. Maternal depletion of
Drp1 and Opa1 in *Drosophila* leads to a decrease in embryo formation and lethality [15,29].
Drp1 knockout mice are also embryonic lethal, and they exhibit neurological defects
[30,31]. Knockdown of Drp1 in *C. elegans* also causes embryonic lethality [22]. Mitochondria
also contribute to embryonic development by producing metabolites that act as
regulators of gene expression [32,33]. Mouse embryos cultured in the presence of
mitochondrial metabolic inhibitor, amino-oxyacetate, show reduced fetal and placental
growth [34]. Differential mitochondrial positioning and metabolic activity, along the
embryonic axis, regulate cell fate determination in Xenopus and sea urchin embryos [35–
39]. However, the mechanism by which mitochondrial morphology, activity and
distribution modulate cell shape changes and morphogenesis during embryonic
development remains unexplored.
*Drosophila* gastrulation is a well defined model system to study mitochondrial
distribution and function in morphogenesis. It depends upon rapid cellular movements
taking place at different domains of the embryo, including internalization of the
mesoderm precursor cells by invagination called the ventral furrow (VF) formation. It
begins with a change in the shape of a few cells at the ventral midline encompassing 10
to 12 cells in width and 80 cells in length. These cells first undergo flattening of the
apical membrane, followed by apical constriction with the help of the acto-myosin
contractile network, changing from columnar to wedge shape. The neighbouring cells

stretch towards the ventral midline simultaneously. As a result of these two events, the

constricting cells invaginate, forming a U-shaped VF which is later internalised, forming

the mesoderm [40–43]. VF formation is activated by the transcription factors Twist and

Snail downstream of the Dorsal/NFkB signaling pathway [41,44].

We examine mitochondrial dynamics during VF formation in gastrulation (Figure

1A) in *Drosophila* embryos to understand the role of mitochondria in driving

morphogenesis. Mitochondria are fragmented and highly abundant in early *Drosophila*

blastoderm embryos [15,16]. We find that mitochondrial migration from basal to apical

regions occurs during gastrulation in the VF cells, regulated by microtubules and the

Dorsal/NFkB signaling pathway. Mitochondrial fission protein Drp1 mutant embryos

show accumulation of larger mitochondria basally that fail to migrate apically, and

embryos contain defects in VF formation. Mitochondria in Drp1 mutant embryos are

functionally compromised with decreased ETC components and ROS levels. Drp1

mutant embryos do not exhibit changes in the Dorsal/NFkB pathway genes, but instead

lead to increased expression of anti-oxidant genes. Increasing ROS genetically by

depletion of mitochondrial SOD2 or decreasing mitochondrial fusion protein Opa1 in

Drp1 mutant embryos results in mitochondrial fragmentation and apical migration in the

VF, also rescuing the furrow defects. Thus, we propose that fragmented mitochondria

migrate apically in the ventral cells during gastrulation to activate Myosin II by

increasing ROS locally, thereby facilitating apical constriction and enabling proper VF

formation.

**STAR Methods**

**Key resources table**

| REAGENT or RESOURCE | SOURCE | IDENTIFIER |
|---|---|---|
| Antibodies | | |
| anti-Drp1 | Boster Bio | DZ41314 |

| | | |
|---|---|---|
| anti-Dorsal | DSHB | 7A4c |
| anti-Twist | Eric F. Wieschaus | NA |
| anti-Twist | Siegfried Roth, Cologne University, Germany | NA |
| Streptavidin, Alexa Fluor® 488 Conjugate | Invitrogen | S32354 |
| Alexa Fluor™ 647 Phalloidin | Invitrogen | A22287 |
| Hoechst | Invitogen | H3570 |
| beta-actin | Cell signaling | #4967 |
| ATP beta | Abcam | ab14730 |
| Goat anti-Rabbit IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, HRP | Thermo Fisher Scientific | A16110 |
| Goat anti-mouse IgG (H+L) Highly Cross-Adsorbed Secondary Antibody, HRP | Thermo Fisher Scientific | G21040 |
| Experimental models: Organisms/strains | | |
| *Drosophila*: UAS-*dl*RNAi | Bloomington *Drosophila* Stock Centre | BDSC#32934 |
| *Drosophila*: *dl*[1] | Bloomington *Drosophila* Stock | BDSC#3236 |

| | | |
|---|---|---|
| | Centre | |
| *Drosophila*:$dl^4$ | Bloomington *Drosophila* Stock Centre | BDSC#7096 |
| *Drosophila*: UAS-SOD2i | Bloomington *Drosophila* Stock Centre | BDSC#32983 |
| *Drosophila*: UAS-*opa1*$^i$ | Bloomington *Drosophila* Stock Centre | BDSC#32358 |
| *Drosophila*: $dl^1$,UASp-Dl$^{WT}$ | Girish Ratnaparkhi, IISER Pune, India | NA |
| *Drosophila*: UASp-Drp1$^{SG}$ | Chowdhary et al, 2020 | NA |
| *Drosophila*: UASp-mito-PAGFP | Chowdhary et al, 2020 | NA |
| *Drosophila*: UASp-mito-GFP | Rachel Cox USUHS, Bethesda, MD | NA |
| *Drosophila*: SqhGFPmat67/cyo; 3xmScarletCAAX15mat/TM3SbSer | Yu-Chiun Wang | NA |
| *Drosophila*: *nanos*-Gal4::UASp-mito-GFP | Chowdhary et al, 2020 | NA |

| | | |
|---|---|---|
| *Drosophila*: *nanos*-Gal4::UASp-mito-GFP::*sqh*-Sqh-mCherry | Chowdhary et al, 2020 | NA |
| *Drosophila*: *nanos*-Gal4::*sqh*-Sqh-mCherry | This study | NA |
| *Drosophila*: UASp-Drp1$^{SG}$;;SOD2i | This study | NA |
| *Drosophila*: UASp-Drp1$^{SG}$;;*opa1*$^{i}$ | This study | NA |
| Chemicals, peptides, and recombinant proteins | | |
| Slow fade Gold Antifade | Thermo Fisher Scientific | S36937 |
| Paraformaldehyde powder | Himedia | TC703 |
| Heptane | Merck | 1.04365.2521 |
| Bovine Serum Albumin | Himedia | TC545 |
| Colchicine | Sigma | #C9754 |
| Citra Solv Cleaner Degreaser Orang | Citra-Solv | https://www.amazon.in/Citra-Solv-Cleaner-Degreaser-Fluid/dp/B007USNPCK/ref=sr_1_1?crid=5T1OR4WG6LH3&dib=eyJ2IjoiMSJ9.pXwdRYLYtNZ-iSDC4AsbzmtXT9xqk7y7p33Gl3zGnHviLw8QgchHVEss4MkP1gUuBzadq-Z-8kjBqK3u_S-heA.-dT6BtISLVpWYJku8etkiS05 |

| | | PcTrFBbmDOjuP6BbybU& dib_tag=se&keywords=citra solv&qid=1755606744&spr efix=citrasol%2Caps%2C23 5&sr=8-1 |
|---|---|---|
| Protease Inhibitor Cocktail | Roche, Sigma-Aldrich | 04693116001 |
| BCA protein estimation kit | Thermo Fisher Scientific | #23225 |
| Polyvinylidene fluoride (PVDF) membrane | Merck | Immobiolon IPVH00010 |
| Tween-20 | Sigma-Aldrich | #P9416 |
| ECL Western Blotting detection reagent | GE healthcare Amersham | #RPN2236 |
| Glutaraldehyde Grade I | Sigma | G5882 |
| Critical commercial assays | | |
| Mitochondrial isolation kit | Sigma | MITOISO-1 |
| TF-HIGH CAP CDNA REV TRANS KIT | Applied Biosystems | 4374966 |
| KAPA SYBR FAST UNI (10ML) | Sigma | KK4602 |
| Deposited data | | |
| RNA-seq data files | This study | GEO: GSE304019 |
| Mass spec data files | This study | PRIDE: PXD066604 |

| Oligonucleotides | | |
|---|---|---|
| Primers for qPCR | | |
| GstD1-Forward Primer: CGACTCCCTGTACCCTAAGT | Sigma | NA |
| GstD1-Reverse Primer: ACACCTGTGGGTAGTAGTAGTT | Sigma | NA |
| Jafrac1-Forward Primer: ACCGAGATCATTGCGTTCTC | Sigma | NA |
| Jafrac1- Reverse Primer: CTTGGCGTGTTGATCCAGGC | Sigma | NA |
| SOD3-Forward Primer: TGGATGCATCAGCATGGGTG | Sigma | NA |
| SOD3-Reverse Primer: CGTGGAGTTGGCCTCCAGGT | Sigma | NA |
| Software and algorithms | | |
| Fiji | [45] | http://fiji.sc |
| Cellpose | [46] | NA |
| Graphpad Prism v 10.4.1 | Informer Technologies, Inc. | http://graphpad-prism.software.informer.com/ |
| VolcaNoseR | NA | https://huygens.science.uva.nl/VolcaNoseR/ |

| Illustrator | Adobe | NA |
|---|---|---|
| Other | | |
| PELLET PESTLE, DISPOSABLE CORDLESS MOTO | Sigma Aldrich | Z359971-1EA |
| 2-well Chambered Coverglass w/ non-removable wells Borosilicate Glass 1.0 | Thermo Fisher Scientific | 155380 |

**Experimental model and subject details**

*Drosophila* **genetics**

All the fly crosses were maintained in the standard cornmeal agar medium. UASp-mito-PAGFP was crossed to *nanos*-Gal4::*sqh*-Sqh-mCherry at 25 $^{o}$C. $dl^{i}$ was crossed to maternal driver line *mat67*;*mat15* carrying maternal □4 *tubulin*-Gal4-VP16, homozygous for chromosome II and III, at 25 $^{o}$C. $dl^{1}$ and $dl^{4}$ mutants were crossed to each other at 25 $^{o}$C. $dl^{1}$ mutant was crossed to *nanos*-Gal4::UASp-mito-GFP::*sqh*-Sqh-mCherry at 28 $^{o}$C. $dl^{1}$,UASp-Dl$^{WT}$ was crossed to *nanos*-Gal4::UASp-mito-GFP at 28 $^{o}$C. For the apical constriction experiment, Sqh-GFP67mat/cyo; 3xmScarlet-CAAX *mat*15/TM3 Sb Ser was crossed to $w^{1118}$ and UASp-*Drp1*$^{SG}$ at 25 $^{o}$C. Embryos were obtained from the F1 generation flies using standard genetic crosses.

**Maintaining the dosage of Gal4 across single and double mutant crosses**

Comparable copies of transgenes were maintained across genotypes to ensure comparative dosage and effects of dilution of Gal4 for expression. Double mutant transgenes, *Drp1*$^{SG}$;*opa1*$^{i}$ and *Drp1*$^{SG}$;*SOD2*$^{i}$, were crossed with either nanos-Gal4 or *nanos*-Gal4::*sqh*-Sqh-mCherry at 28 $^{o}$C. Single mutant transgenes, *Drp1*$^{SG}$ and *SOD2*$^{i}$, were crossed to either *nanos*-Gal4::UASp-mito-GFP or *nanos*-Gal4::UASp-mito-GFP::*sqh*-Sqh-mCherry at 28 $^{o}$C to maintain similar Gal4 amounts as those in the double mutant transgenes.

**Method details**

**Embryonic Lethality Estimation**

3-4 hr old embryos were collected from cages. 100 embryos were aligned using a brush

on a fresh sugar agar plate in a 10x10 grid and kept at the same temperature as the

cage. Unhatched embryos were counted after 24 hrs and 48 hrs. This was done 3

times, and the average of unhatched embryos was calculated and reported as a

percentage of the total number of embryos.

**Live imaging**

2.5 hr old embryos from cages at 28 $^{o}$C were dechorionated using bleach for 60-90

seconds, washed and then placed horizontally in a lab-tek chamber. End-on imaging

was done to study the process of VF formation. For End-on imaging, after mounting

embryos in the lab-tek chamber and adding 1X PBS, each embryo was picked up gently

from one end using a single hair brush such that it stands vertically. The lab-tek

chamber was then carefully placed at the confocal microscope avoiding any jerk that

could cause embryos to fall or tilt. The lab-tek chamber was then screened for

completely vertical embryos of the desired stage and imaged using 63X/1.4 NA oil

objective on Zeiss laser scanning microscope LSM 710 or Leica laser scanning

confocal microscope SP8. The laser power and gain settings were adjusted using the

range indicator mode such that the acquired 8-bit image was within the 0-255 range.

End-on live imaging was done at 1024x1024 resolution at 2 airy units. 3-5 Z stacks

were taken every 30-60 seconds, with 1 μm interval, starting from 45-50 micron depth.

For imaging apical constriction, 2.5 hrs old embryos were taken from cages at 25 $^{o}$C.

Embryos at early cellularization were selected and dechorionated using bleach and

then washed. Embryos were then placed on an agar slab with the ventral side facing

upwards. Lab-tek chamber, with a drop of heptane glue in the centre, was then inverted

onto the agar slab gently such that the embryos stuck on their ventral side. 1X PBS was

added, and embryos were imaged. 2 color imaging was done using 60X oil objective on

the Evident SpinSR10 microscope. The laser power and other acquisition settings were

171  kept the same for control and mutant. 16 bit image was acquired with a time interval of

172  7.8 seconds.

**Immunostaining**

3 hr old embryos were collected from cages at 28 $^{\circ}$C for cellularization and gastrulation

stages. Embryos were dechorionated using bleach, washed and then fixed using 1:1

Heptane and 8% PFA or 4% PFA (for Drp1 staining) solution for 25 mins. Embryos

were then devitellinized either using an insulin needle (hand de-vitellinization) or by

shaking in 1:1 Heptane:Methanol (Methanol de-vitellinization for Drp1 staining). After

devitellinization, embryos were washed 3 times with 1X PBST (1X PBS with 0.3% Triton

X-100) for 5 mins each, blocked using 2% BSA (Sigma-Aldrich) in 1X PBST for 1 hr at

room temperature, then incubated with primary antibodies (anti-Drp1 Booster Bio

DZ41314, rabbit, 1:500; anti-Dorsal DSHB 7A4c, mouse, 1:1000; anti-Twist (Eric

Wieschaus, Princeton, USA, rabbit, 1:300), anti-Twist (Siegfried Roth, Koln University,

Germany, rabbit, 1:500) overnight at 4 °C. Excess primary antibody was removed by

giving 3 washes with 1X PBST 5 mins each. Fluorescently labelled dyes (streptavidin

1:1000, phalloidin 1:1000) and/or secondary antibodies (1:1000), were added and

incubated at room temperature for 1 hr after covering with aluminium foil. Following this,

3 washes of 1X PBST were given, and DNA stain Hoechst (1:1000) was added in the

second wash. Finally, embryos were mounted in Slow fade Gold antifade and imaged

using Plan Apochromat 63x/1.4 NA, oil immersion objective on Leica laser scanning

confocal microscope SP8. Care was taken that the acquired 8-bit image was within the

0-255 range and did not saturate. Imaging was done at 512x512 resolution with a line

averaging of 2. 0.5 µm optical sections were acquired for Z stacks at 1AU. For DHE

staining, embryos were fixed in 1:1 4% PFA:Heptane for 10 mins and then methanol de-

vitellinized. Embryos were taken in 0.1% PBST (1X PBS with 0.1% Triton X-100) and

permeabilized with 5% BSA solution made in 1% PBST (1X PBS with 1% Triton X-100)

for 90 mins. Blocking was done with 5% BSA made in 0.1% PBST for 30 mins at room

temperature. Embryos were incubated in DHE (2 µM in 1X PBS) for 10 mins. Lastly,

one wash was given with 1X PBS and then embryos were mounted in Slow fade Gold

antifade.

**Photoactivation**

Embryos expressing mito-PAGFP and Sqh-mCherry were mounted in end-on
orientation. A ring shaped ROI was drawn at the basal region at the end of
cellularization/ beginning of gastrulation. The selected ROI was photoactivated using a
405 nm laser at 10% power, using a 60×/1.4 NA Plan apochromat objective on Evident
FV4000 inverted confocal microscope. Images were obtained every 30 secs.

**Colchicine treatment via Citrasolv**

2-2.5 hr old *nanos*-Gal4::UASp-mito-GFP::*sqh*-Sqh-mCherry embryos were collected
from cages at 28 $^o$C. Embryos were treated with bleach for 3 mins and then washed
with distilled water. Further, citrasolv treatment was given (1:10 in 1X PBS) for 5 mins
on the rocker (embryos were in the sieve). Citrasolv was removed by rinsing the
embryos 2-3 times with 1X PBS. Then the embryos were transferred using a brush to a
tube containing 1X PBS, and 1mM colchicine (Sigma, #C9754) was added and
incubated for 15 mins on the rocker. Finally, the drug was removed and embryos were
rinsed with 1X PBS once and then mounted on a lab-tek chamber and imaged using
63X/1.4 NA oil objective on Zeiss laser scanning microscope LSM 710.

**TEM of control gastrulation embryos**

A 30 min collection of *yw* embryos at 25 $^o$C was taken and aged till the gastrulation
stage. Embryos were then dechorionated with bleach and fixed for 10 min in a mixture
of 1 ml glutaraldehyde 25% and 4 ml heptane. Heptane and glutaraldehyde were
removed, and embryos were immersed in phosphate-buffered saline (PBS) containing
0.1% Tween 20 and 0.1% bovine serum albumin (BSA). Embryos were hand-
devitellinized on a double-stick tape with a hypodermic needle (BD Microlance 3–
30Gx1/2″, BD 304000). De-vitellinized embryos of the desired stages were transferred
to a freshly prepared fixative solution (2.5% glutaraldehyde in 50 mM cacodylate buffer,
pH 7.2–7.4, 0.1% tannic acid) and then stored at 4°C in the same solution overnight.
Embryos were post-fixed for 2 h in 1% osmium, 2% glutaraldehyde in 50 mM
cacodylate buffer at 4°C. They were then washed in cacodylate buffer, dehydrated in a

graded ethanol series, and finally embedded in epon. Ultrathin sagittal sections (80 nm)

were prepared with an ultramicrotome, mounted onto copper grids, contrasted with 1%

uranyl acetate, and examined with a FEI Tecnai G2 200 kV electron microscope.

Images were acquired with a FEI Eagle charge-coupled device (CCD) camera (4096 ×

4096 pixels) for 1700× magnification and an Olympus Veleta CCD camera (2048 ×

2048 pixels) for 3100× and 8500× magnifications. The procedure was carried out in

Institut de Biologie du Développement de Marseille (IBDM) electron microscopy facility

(Marseille, France).

**TEM of control and Drp1$^{SG}$ cellularization embryos**

1.5 hr embryos were collected, aged for 2 hrs, dechorionated using bleach for 1 min

and then rinsed with distilled water. Embryos were then transferred to a scintillation vial

containing a 1:1 solution of 3% glutaraldehyde and heptane for 2 hr at 4°C on the

rocker for fixation. After fixation, all the solution was removed, fresh heptane was added

and then embryos were hand-devitellinized using an insulin needle. Devitellinized

embryos were collected in a 0.5ml eppendorf tube. Two washes were given with 0.1M

sodium cacodylate buffer for 10 mins each and post-fixed in 1 % Osmium tetroxide

(OsO4) for 1 hr at 4 $^{o}$C. The tissues were en-block stained in 2% uranyl acetate

followed by dehydration in an ascending alcohol series for 10 minutes each. The tissues

were infiltrated in Araldite resin (A and B) and finally embedded in Araldite resin B and

polymerized at 60 $^{o}$C for 48 hrs. Ultrathin sections of 70 nm were cut using an

ultramicrotome (LeicaUC7) and collected on copper slot grids. The sections were

contrasted with lead citrate. The images were acquired using a Transmission Electron

Microscope, JEM 1400 Plus, JEOL (Japan) at 120 kV. The sample embedding,

sectioning and imaging were carried out in the Transmission Electron Microscope

facility, ACTREC, Tata Memorial Centre, Kharghar, Navi Mumbai, India.

**Western Blotting**

3 hr old embryos were collected from cages at 28 $^{o}$C and dechorionated using bleach

for 1 min, followed by washing in distilled water. Then embryos were crushed on ice in

150-200 µl lysis buffer (1%Triton-X 100, 50 mM Tris HCl; pH 8.0, 150 mM NaCl,

Protease Inhibitor Cocktail; PIC 1:50) using a homogeniser gun. The homogenate was
then centrifuged at 15000 g for 30 mins at 4 °C. The supernatant was removed
carefully, avoiding the lipid layer at the interface, and stored at -80 °C. Protein
concentration of the supernatant was determined using BCA protein estimation kit
(Thermo Fisher Scientific #23225). A sample for SDS-PAGE was prepared by boiling
the protein sample (supernatant) in 1X Laemmli buffer at 95 $^{o}$C for 10 mins. The
denatured protein sample was then loaded into 10% or 12% polyacrylamide gel
followed by transfer onto a 0.45µm polyvinylidene fluoride (PVDF) membrane (Merck
Immobiolon IPVH00010). Post transfer, the blot was blocked for 1 hr at room
temperature with 5% skimmed milk or 5% BSA dissolved in 1X TRIS buffer saline (TBS)
with 0.1% Tween-20 (Sigma-Aldrich, #P9416) detergent. After blocking, the primary
antibodies (Drp1 1:5000; β-actin Cell signaling #4967, rabbit, 1:1000; ATP beta abcam
ab14730, mouse, 1:2000) was added and kept at 4 $^{o}$C for 12-16 hrs. Following this, the
blot was then washed, 3 washes 15 mins each, using 1X TRIS buffer saline (TBS) with
0.1% Tween-20. Then, secondary antibody is added for 1 hr at a concentration of
1:10,000 and kept at room temperature. Blot is then again washed thrice and then
developed using ECL Western Blotting detection reagent (GE Healthcare Amersham,
#RPN2236).
282

**Mitochondria Isolation**

Mitochondria are isolated from embryos using the mitochondria isolation kit (Sigma,
MITOISO1). 3 hr old embryos are taken from cages at 28 $^{o}$C and dechorionated using
bleach for 1 min. Dechorionated embryos are rinsed with a solution EB-A twice. Then,
embryos are crushed on ice in 200 µl EB-A with 2mg/ml BSA and centrifuged at 600g
for 5 mins at 4 $^{o}$C. The pellet is discarded, and the supernatant is centrifuged at
11,000g for 10 mins at 4 $^{o}$C. The supernatant was discarded. The pellet is then
resuspended in 100 µl EB A buffer and centrifuged at 600 g for 5 mins at 4 $^{o}$C.
Following this, the pellet obtained is discarded as it contains the nuclear fraction. The
supernatant is centrifuged at 11,000 g for 10 mins at 4 $^{o}$C. The pellet obtained after this
spin is enriched in mitochondria, and it is resuspended in a storage buffer and stored at
-20 $^{o}$C.

**RNA sequencing**

For sample preparation for RNA sequencing, embryos were manually sorted to avoid non-developing embryos and to select for only cellularizing and gastrulating embryos. Firstly, egg laying was synchronised by discarding the first 2 hr collection. For genotypes that laid enough embryos, 1 hr collection was taken and aged for 1.5-2 hrs, whereas for mutants that did not lay enough embryos 2.5-3 hr collection was taken directly after synchronisation from cages at 28 $^o$C. Embryos were dechorionated using bleach and washed with distilled water. Then, around 50-60 embryos were taken, using a brush, on a slide, and around 20-30 µl 1X PBS was added on top of the embryos. Embryos of desired stages were then sorted using a single-haired brush under a stereo microscope. These embryos were taken in a tube and crushed in TriZol (Qiagen) using a homogeniser gun, snap frozen in liquid nitrogen, and then stored at -80 °C. Multiple embryo collections were pooled such that there were 70-100 embryos (equal proportion of cellularizing and gastrulating embryos) in each replicate. RNA was isolated using the TRIZOL method. The concentration of RNA was measured using Qubit RNA BR Assay (Invitrogen, Cat# Q10211). RNA purity was checked using QIAxpert, and RNA integrity was checked on TapeStation using RNA ScreenTapes (Agilent, Cat# 50675579). The samples were taken for the library preparation and sequencing.

Library Prep Protocol:

KAPA RNA HyperPrep Kit (cat #0000141759) protocol was used to prepare libraries for mRNA sequencing. An initial Concentration of 200ng of total RNA was taken for the assay. mRNA molecules are captured using magnetic oligo Poly(T) beads (KAPA mRNA capture kit, Cat# 7962240001). Following purification, the enriched mRNA was fragmented using divalent cations under elevated temperatures (94 degrees for 6 minutes). The cleaved RNA fragments were copied into first-strand cDNA using reverse transcriptase and random priming. Combined 2nd strand synthesis and A-tailing is done, which converts the cDNA:RNA hybrid to double-stranded cDNA (dscDNA), incorporates dUTP into the second cDNA strand, and adds dAMP to the 3' ends of the resulting dscDNA. This is followed by Adapter Ligation, in which dsDNA indexing

adapters with 3' dTMP overhangs are ligated to library insert fragments. The adapter-

ligated products were then purified and enriched carrying appropriate adapter

sequences at both ends using high-fidelity, low-bias PCR (the strand marked with dUTP

is not amplified, allowing strand-specific sequencing) using the following thermal

conditions: initial denaturation 98 $^{\circ}$C for 45sec; 13 cycles of - 98 $^{\circ}$C for 15sec, 60 $^{\circ}$C for

30sec, 72 $^{\circ}$C for 30sec; final extension of 72 $^{\circ}$C for 1mins. PCR products are then

purified and checked for fragment size distribution on Fragment Analyzer using HS

NGS Fragment Kit (1-6000bp) (Agilent, Cat# DNF-474-1000).

Sequencing Protocol:

Prepared libraries were quantified using Qubit HS Assay (Invitrogen, Cat# Q32854).

The obtained libraries were pooled and diluted to the final optimal loading

concentration. The pooled libraries were then loaded onto Illumina NovaSeq V1.5

instrument at Medgenome, India, to generate 40M, 150bp paired-end reads/sample.

**RT-PCR**

The concentration of RNA was adjusted to 1µg using nuclease-free water. High-

capacity cDNA reverse transcription kit was used for the synthesis of cDNA using the

manufacturer's protocol. cDNA was used at a dilution of 1:5 for all reactions.

**Mass Spectrometry**

Two kinds of samples were prepared for mass spectrometry, total protein sample and

mitochondrial protein sample. Total protein sample was prepared similarly to the sample

prep for western blots, wherein embryo lysate in lysis buffer is centrifuged at 15000 g

for 30 mins at 4 $^{\circ}$C and the supernatant is used for mass spec. For the mitochondrial

protein sample, mitochondria were isolated from embryos as mentioned above. The

final mitochondrial pellet was resuspended in a buffer containing 5% SDS and 0.1M Tris

pH 8.0, and used for mass spec. The following processes were done with Vproteomics,

New Delhi, India.

Sample Preparation for Mass Spec

25 µg of the protein sample was first reduced with 5 mM TCEP and subsequently

alkylated with 50 mM iodoacetamide. The protein was then digested with Trypsin at a

1:50 Trypsin-to-lysate ratio for 16 hours at 37 $^{o}$C. After digestion, the mixture was

purified using a C18 silica cartridge and then concentrated by drying in a speed vac.

The resulting dried pellet was resuspended in buffer A, which consists of 2% acetonitrile

and 0.1% formic acid.

**Quantification and statistical analysis**

**Image Analysis and quantification**

**Mitochondrial fluorescence in the ventral furrow**

Mean mitochondrial fluorescence was measured in apical (above the nucleus) and

basal (below the nucleus) regions of cells at 3 different regions of the embryo: Ventral

(cells undergoing constriction, marked in red), neighbouring (cells adjacent to the

constricting cells, marked in green), and lateral (cells on the lateral region of the

embryo, marked in blue). 5 different time points were chosen, based on the shape and

extent of invagination of the furrow being formed, starting from the end of cellularization

till the time point when the furrow closes. The background intensity was taken from a

region within the embryo that did not contain any visible fluorescence and subtracted

from the apical and basal intensities. Ratio of background subtracted apical:basal

intensities were normalized with the ratio of the first time point i.e., end of cellularization

and plotted as Mean ± SEM using GraphPad Prism 10.4.1. Image analysis was done

using ImageJ.

**Quantification of mitochondria in TEM**

Mitochondria in the EMs were identified based on structure and electron density.

Mitochondria in each cell were counted manually, and their size and circularity were

measured using ImageJ.

**Mitochondrial and Myosin II fluorescence in photoactivation**

Fluorescence of mitochondria and Myosin II was followed in the apical and basal

regions of the ventral and lateral cells. Background normalization was done by

subtracting the mean intensity of a region within the embryo. Background subtracted

mean apical:basal intensities of mitochondria and Myosin II for the 2 regions (ventral

and lateral) were measured across time and plotted as Mean ± SEM using GraphPad

Prism 10.4.1.

**Measurement of mitochondrial area in apical and basal sections**

Single planes at the apical and basal regions were used for analysis. Firstly,

background normalization was done by subtracting the mean intensity value. The

thresholding was done, and particles greater than 0.05 $\mu m^2$ were marked using the

"particle analyser" tool in ImageJ. Mean particle size was determined for each embryo.

The data were plotted as a violin plot and analyzed with a Mann–Whitney test using

GraphPad Prism 10.4.1.

**Measurement of relative mitochondrial area in apical sections**

Apical stacks of late cellularization embryos were selected using phalloidin and Hoechst

as a reference. Particles greater than 0.05 $\mu m^2$ were selected using thresholding.

Relative mitochondrial area was obtained by dividing the total area of all thresholded

mitochondrial particles in the apical stacks by the total area of the imaging field and

expressed as a percentage. The data were plotted and analyzed using GraphPad Prism

10.4.1.

**Dorsal and Twist intensity**

Nuclei were marked using Hoechst staining. Fluorescence intensity of Dorsal and Twist

was measured in the marked nuclei. Mean ventral to dorsal fluorescence intensity was

plotted and compared using the Mann-Whitney test in GraphPad Prism 10.4.1.

**Mitochondrial fluorescence in Dorsal overexpression**

ROI was drawn, using the segmented line tool, in the apical region of the entire embryo

(end-on) in control and *dl*[1],UASp-Dorsal and the mean mitochondrial fluorescence was

measured. The perimeter of the embryos was measured and normalized to arbitrary units in percentage such that the ventral midpoint is marked as 0, left to that is 0 to -50, and right to that is 0 to 50. Mean of the normalized intensities at every 1.0 airy unit was plotted using GraphPad Prism 10.4.1.

**Quantification of apical area and Myosin II intensity during apical constriction**

For representation and quantification, a single Z-plane at 5 µm depth from the apical cortex was taken for ScarletCAAX. For Myosin II, average intensity Z-projections were taken of the planes that showed Sqh-GFP signal. The time point when the furrow reaches a depth of 2.5-2.7 µm was taken as the end, and 7.8 mins before that was taken as the first time point. Cells were segmented and tracked, and their area and Sqh-GFP intensity were measured using Cell Pose and Trackmate. Segmented cells were color-coded based on the area. Area and Myosin II intensity of 40 cells from one control and mutant embryo were plotted as a heat map in GraphPad Prism 10.4.1. Mean apical area and mean Myosin II intensity were normalized to their respective first time and plotted as Mean ± SEM using GraphPad Prism 10.4.1.

**Quantification of DHE fluorescence**

The mean fluorescence intensity of DHE was calculated for a single plane located just above the nucleus. The mean intensity values of all genotypes were normalized with the average mean intensity of the control, and plotted using GraphPad Prism 10.4.1 and compared using the Mann-Whitney test.

**Measurement of ventral furrow width**

A single time point was chosen based on the furrow shape (when the furrow was U-shaped- T4). The furrow width was calculated by drawing a line across the furrow at the point where the Myosin II signal (Sqh-mCherry) ended. The mean values were plotted as a violin plot in GraphPad Prism 10.4.1 and compared using the Mann-Whitney test.

**Myosin II fluorescence in the ventral furrow**

Mean Myosin II (Sqh-mCherry) intensity was measured in the ventral cells by drawing

an ROI around the apical Myosin II signal. Background intensity was measured from a

region within the embryo, which did not have visible fluorescence. Background

subtracted mean Myosin II intensity was normalized with the first time point and plotted

with SEM using GraphPad Prism 10.4.1.

**RNA Sequencing analysis**

Read quality check

The following parameters were checked from fastq file - Base quality score distribution,

Sequence quality score distribution, Average base content per read, GC distribution in

the reads, PCR amplification issue, Check for over-represented sequences and Adapter

trimming. Based on the quality report of fastq files, sequence reads were trimmed

where necessary to only retain high quality sequence for further analysis. In addition,

the low-quality sequence reads were excluded from the analysis. The adapter trimming

was performed using Trimmomatic (v-0.36).

Contamination removal

For the RNA-Seq analysis, the unwanted sequences were removed, especially

nonpolyA tailed RNAs from the sample (assuming that poly-A tailed RNAs are

sequenced). The unwanted sequences include - mitochondrial genome sequences,

ribosomal RNAs, transfer RNAs, adapter sequences and others. Contamination

removal was performed using Bowtie2 (2.2.4).

Read alignment

The paired-end reads were aligned to the reference *Drosophila melanogaster* genome

using HISAT2 (2.1.0) to quantify reads mapped to each transcript. Alignment

percentage of reads was in the range of 91-95 percent for all the samples.

Expression estimation

The aligned reads were used for estimating the expression of the genes. The raw read

counts were estimated using FeatureCount (1.5.2).

Differential expression analysis

The raw read counts were normalized using DESeq2. The ratio of normalized read

counts for treated over control was taken as the fold change. Genes were first filtered

based on the p-value (≤ 0.05). A distribution of these log2 (foldchange) values were

found to be normally distributed. Those genes which were found to have -1≤

log2(foldchange) ≥1 were considered statistically significant. Gene ontology of the

significantly differentially expressed gene set was performed using ShinyGO 0.81.

Volcano plot was plotted using VolcaNoseR. Heat maps were plotted in GraphPad

Prism 10.4.1 using log (CPM+1) values for genes which were unchanged and z-

normalized values for differentially expressed genes. The RNA isolation, sequencing

and analysis was done with MedGenome Labs, Bangalore, India.

**RT-PCR analysis**

The $\Delta\Delta Ct$ method was used to calculate the gene expression changes among

comparative genotypes (control and Drp1$^{SG}$)

$$\Delta Ct_{\text{sample}} = Ct_{\text{target gene}} - Ct_{\text{reference gene}}$$

$$\Delta\Delta Ct = \Delta Ct_{\text{mutant sample}} - \Delta Ct_{\text{control sample}}$$

$$\text{Fold change} = 2^{-\Delta\Delta Ct}$$

Where, Ct is the cycle threshold or cq value. The reference gene is the housekeeping

gene *rp49*. Comparisons between genotypes were made, and fold change was plotted

in the form of a bar chart to represent the difference in levels of gene expression.

**Western Blot analysis**

The image was first inverted. Background subtraction was done. The intensity of bands

was normalized by dividing it by the intensity of the loading control.

**Mass Spectrometric Analysis of Peptide Mixtures**

For mass spectrometric analysis, all the experiments were performed on an Easy-nLC-

1000 system (Thermo Fisher Scientific) coupled with an Orbitrap Exploris 240 mass

spectrometer (Thermo Fisher Scientific) and equipped with a nano electrospray ion

source. 1 µg of phosphopeptides sample was dissolved in buffer A containing 2%

acetonitrile/0.1% formic acid and resolved using Picofrit column (1.8-micron resin, 15cm

length). Gradient elution was performed with a 0–38% gradient of buffer B (80%

acetonitrile, 0.1% formic acid) at a flow rate of 500nl/min for 96 mins, followed by 90%

of buffer B for 11 min and finally column equilibration for 3 minutes. Orbitrap Exploris

240 was used to acquire MS spectra under the following conditions: Max IT = 60ms,

AGC target = 300%; RF Lens = 70%; R = 60K, mass range = 375–1500. MS2 data was

collected using the following conditions: Max IT= 60ms, R= 15K, AGC target 100%.

Lock mass option was enabled for polydimethylcyclosiloxane (PCM) ions (m/z =

445.120025) for internal recalibration during the run. MS/MS data was acquired using a

data-dependent top20 method dynamically choosing the most abundant precursor ions

from the survey scan, wherein dynamic exclusion was employed for 30s.

Data Processing

RAW files generated were analyzed with Proteome Discoverer (v2.5) against the

Uniprot *Drosophila melanogaster* database. For dual Sequest and Amanda search, the

precursor and fragment mass tolerances were set at 10 ppm and 0.02 Da, respectively.

The protease used to generate peptides, i.e. enzyme specificity, was set for trypsin/P

(cleavage at the C terminus of "K/R: unless followed by "P"). Carbamidomethyl on

cysteine as fixed modification and oxidation of methionine and N-terminal acetylation

were considered as variable modifications for database search. Both peptide spectrum

match and protein false discovery rate were set to 0.01 FDR.

Further downstream analysis was performed using in-house R-scripts. For

differential expression analysis limma package in R was used. For differential protein

abundance analysis, the abundance values were first Log2 transformed. Then the data

was filtered based on valid values in at least 70% of the replicates within each

condition. Missing values were imputated based on normal distribution. Abundance

values were median-based quantile normalised. Bioreplicates were categorized and

grouped in Control and Treated groups. Post-processing, t-test was applied which

provided the pValues and log2 fold change values. Proteins having p value<0.05 were

considered significant. log2 fold change ≥0.3 was considered significant. Abundance

values of significant proteins were Z-scale transformed. Heatmap was generated on

these proteins Z-normalized data using GraphPad Prism 10.4.1. Gene ontology of the

significantly differentially expressed proteins was performed using ShinyGO 0.81.

Volcano plot was plotted using VolcaNoseR.

**Statistics and n values**

The Mann-Whitney test was applied for statistical analysis. n values ranged from 3 to 27

and are mentioned in the respective figure legends for each experiment. Significance

levels are indicated as follows: $p \geq 0.05$ (ns), $p < 0.05$ (*), $p < 0.01$ (**), $p < 0.001$ (***),

$p < 0.0001$ (****).

**Results**

**Mitochondria are enriched apically in the ventral furrow cells during gastrulation**

Mitochondria are fragmented, dispersed and highly abundant in the early *Drosophila*

embryos [15,16]. We imaged mitochondria during cellularization and gastrulation with

fluorescently coupled streptavidin [16]. As seen previously, mitochondria were enriched

apically as cellularization proceeded, with mitochondria present above the nucleus in

late stages of cellularization (Figure 1B). We further evaluated the distribution of

mitochondria in gastrulation. Mitochondria appeared as small, punctate structures

predominantly in apical regions, above the nucleus, in VF cells as compared to the

basal regions (Figure 1C). We performed live imaging of mito-GFP expressing embryos

during gastrulation in an end-on orientation to study the dynamics of mitochondria. The

end-on orientation enabled the visualization of apico-basal mitochondrial dynamics in

the dorsal, lateral, and ventral axes of each embryo simultaneously. We observed an

apical enrichment of mitochondria in constricting VF cells compared to the neighboring

and the lateral cells. This enrichment increased as gastrulation proceeded and peaked

when cells invaginated to form the VF (Figure 1D, Video 1). We measured the mean

apical and basal mitochondrial fluorescence in 3 regions, constricting ventral cells (VF

cells, red), immediately neighbouring cells (green) and lateral cells (blue) across 5 time

points of VF formation. The apical:basal mean mitochondrial fluorescence ratio was
normalised to the first time point at the end of cellularization. This quantification showed
that the apical mitochondrial intensity increased with time in the ventral cells but
remained almost constant in the neighbouring and lateral cells (Figure 1E). We
observed a similar abundance of mitochondria in the apical regions of the VF cells
compared to the lateral cells in transmission electron micrographs (Figure 1F). The
numbers of apical mitochondria per cell were significantly increased in ventral cells
compared to lateral cells (Figure 1G). In summary, we found that small, punctate
mitochondria were enriched in apical regions in the VF cells during gastrulation.
579
**Mitochondria translocate apically in the ventral furrow cells in a microtubule-**
**dependent manner**
Mitochondria translocate in cells with the help of motor proteins in a microtubule-
dependent manner from the perinuclear region to the periphery in migrating cells and
from the cell body to the synapse in neurons and from basal to apical regions in cells in
*Drosophila* cellularization [15,47–49]. We assessed whether apical migration in VF cells
occurred from basal regions and if this was dependent on microtubules. Live embryos
expressing matrix-labelled photoactivatable mito-GFP (mito-PAGFP) and Sqh-mCherry
were imaged in the end-on orientation. A region of interest in the basal section of all
cells was photoactivated at the end of cellularization or the beginning of gastrulation
(Figure 2A, white ROI). Remarkably, we saw an increased movement of mitochondria
from the basal to the apical regions in the VF cells as compared to lateral cells
immediately after photoactivation (Figure 2A, Video 2). The mitochondrial fluorescence
in apical regions of ventral cells increased to a significantly greater extent as compared
to lateral cells (Figure 2B). This demonstrated that mitochondrial migration from the
basal region to the apical regions was enhanced explicitly in the ventral cells during
gastrulation. The increase in mitochondrial intensity was accompanied by an increase in
the Myosin II fluorescence in the ventral cells (Figure 2A and B). We observed
mitochondria migrate linearly in these cells (Figure 2A, Zoomed in, white arrowheads).
To further assess whether the migration was microtubule-dependent, we treated
embryos expressing mito-GFP with the microtubule-depolymerizing drug colchicine at

the end of cellularization and imaged them live in an end-on orientation during the

process of gastrulation. Colchicine treated embryos showed reduced VF invagination,

as has been reported in previous studies [50]. Apical mitochondrial enrichment in the

constricting cells was significantly reduced, and mitochondria were instead enriched at

the basal side (Figure 2C). We quantified the mean apical:basal mitochondrial intensity

in the constricting cells at a time point showing comparable constriction in the ventral

cells in control and colchicine treated embryos, and found that it was significantly

reduced in colchicine treated embryos compared to the control embryos (Figure 2D).

These data confirmed that mitochondria migrate in a microtubule-dependent manner

from the basal to the apical side in VF cells during gastrulation.

**Dorsal/NFkB regulates apical translocation of mitochondria in gastrulation**

Dorso-ventral patterning in *Drosophila* embryos is activated by the Dorsal/NFkB

(Dorsal) signaling pathway in ventral cells [51]. We further assessed the role of the Dorsal

pathway in regulating the migration of mitochondria in the VF cells. We used the *dl*RNAi

and transheterozygous Dorsal null mutants ($dl^A/dl^1$) to assess the mitochondrial

distribution in these embryos. In both cases, embryos showed 100% lethality and did

not proceed beyond cellularization to the formation of the VF. We therefore assessed

the mitochondrial dynamics in cellularization in Dorsal depleted embryos. *dl*RNAi and

$dl^A/dl^1$ mutants were stained with streptavidin, and mitochondrial morphology and

distribution were assessed. We found that the apical mitochondrial fluorescence in

*dl*RNAi and $dl^A/dl^1$ embryos was reduced compared to the controls (Figure 3A). The

percentage of apical area occupied by mitochondria in late cellularizing embryos was

calculated using a threshold for mitochondrial fluorescence. The area occupied by

mitochondria in apical sections in *dl*RNAi and $dl^A/dl^1$ embryos was significantly reduced

as compared to control embryos (Figure 3B). However, the mitochondrial size in the

apical and basal sections was similar in *dl*RNAi, $dl^A/dl^1$ and control embryos (Figure

3C). Loss of Dorsal decreased the apical mitochondrial migration in cellularization but

did not change the mitochondrial morphology. We further assessed haplo-insufficient

heterozygous $dl^1/+$ embryos for defects in mitochondrial migration to the apical side.

$dl^1/+$ embryos showed 74% lethality and have been previously shown to cause ventral

patterning defects due to haploinsufficiency at higher temperatures [52–55]. First, we

assessed the mitochondrial migration in cellularization in $dl^1$/+ embryos. The relative

apical area occupied by mitochondria was significantly lower than that of the controls.

The mitochondrial size in apical and basal sections was comparable to that of control

embryos (Figures 3A-3C). Further, we imaged mitochondria in live $dl^1$/+ embryos

expressing mito-GFP mounted in the end-on orientation (Video 3). 24% (4/17) $dl^1$/+

embryos did not form the VF, 47% (8/17) of embryos showed a shallow defective furrow

(Figure 3D, white arrowhead), and 29% (5/17) of embryos showed VF formation similar

to controls. In the $dl^1$/+ embryos with defective furrow, the apical to basal mitochondrial

fluorescence in the ventral cells was significantly lower compared to controls (Figure

3E).

We further tested the mitochondrial migration in embryos overexpressing Dorsal.

We overexpressed Dorsal in the background of $dl^1$ ($dl^1$,UASp-Dorsal) using *nanos*-Gal4,

UASp-mito-GFP (referred to as Dl-OE). Dorsal overexpression led to localization of

Dorsal and Twist antibodies on both the dorsal and ventral side in 88% of the embryos

in contrast to controls showing an expression on the ventral side (Figure S1A). This

increase in expression led to a decrease in the ratio of ventral:dorsal nuclear

fluorescence for Dorsal and Twist in Dorsal overexpressing embryos as compared to

controls (Figure S1B). Further, we investigated mitochondrial dynamics using mito-GFP

in live embryos overexpressing Dorsal mounted in end-on orientation (Video 4). 83%

(5/6) of the embryos did not form the VF, and the remaining 17% (1/6) of the embryos

contained invaginating cells with an abnormal furrow. We compared mitochondrial

dynamics in the early time points of gastrulation. We observed that the apical increase

in mitochondrial fluorescence in controls was restricted to approximately 8 cells. In

Dorsal overexpressing embryos, 33% (2/6) of the embryos showed increased apical

mitochondrial fluorescence throughout the embryo, 50% (3/6) showed enhanced apical

fluorescence in approximately half of the embryo, and 17% (1/6) of the embryos

showed a restricted increase in apical mitochondrial signal (Figure 3D and S1C). We

quantified apical:basal mitochondrial intensity in gastrulation when apical constriction

was visible and plotted this with respect to the distance along the dorso-ventral axis

(Figure 3F). Control embryos showed a peak of apical mitochondrial fluorescence in the

ventral region, whereas the mitochondrial fluorescence in Dorsal overexpressing

embryos was significantly spread across a greater number of cells. These data showed

that Dorsal regulates apical mitochondrial migration in the VF cells.

**Drp1 mutant embryos show accumulation of larger mitochondria in basal regions**

**and decreased apical mitochondrial migration**

Loss of mitochondrial fission leads to an increase in mitochondrial size, which affects

their migration within cells. Loss of mitochondrial fission protein, Drp1, in neurons has

been shown to cause mitochondrial accumulation in the cell body and depletion of

mitochondria from synapses and dendrites [56–58]. Drp1 depleted migrating cells show

perinuclear accumulation of mitochondria and defects in actin cytoskeletal remodelling

at the leading edge [10,47,48]. Drp1 null mutant germ line clones do not produce

embryos[29]. We have previously expressed GTPase domain mutants of Drp1 to

abrogate mitochondrial fission in embryonic development and neural stem cells [15,59].

We expressed the GTPase domain mutant of Drp1, Drp1$^{SG}$, to disrupt mitochondrial

fission in embryos using *nanos*-Gal4, UASp-mito-GFP. We found that approximately

84% of Drp1$^{SG}$ embryos did not hatch into larvae at 24 hrs. Mitochondria were present

in large, clustered spots in the basal regions and small, dispersed spots in apical

regions in Drp1$^{SG}$ embryos (78%) compared to small, dispersed mitochondria in apical

and basal regions in controls in VF cells (Figure 4A, white arrowhead). We assessed

the localization and expression levels of Drp1 in the control and Drp1$^{SG}$ embryos

(Figure 4B-D and S2A). In control embryos, Drp1 antibody fluorescence was present on

mitochondria in cellularization (Figure 4B, white arrowhead). In Drp1$^{SG}$ expressing

embryos, Drp1 antibody fluorescence was present as punctate structures on the

clustered mitochondria in the basal sections, and Drp1 punctae did not colocalize with

mitochondria in apical regions (Figure 4B, white arrowheads). Drp1 levels were found to

be higher in western blots in gastrulation, but this was not statistically significant (Figure

4C). Drp1$^{SG}$ embryos showed a significantly higher expression of Drp1 compared to

controls (Figure 4D).

We performed transmission electron microscopy of Drp1$^{SG}$ embryos to assess

the change in mitochondrial morphology. The mitochondria in late cellularizing Drp1$^{SG}$

embryos were larger compared to controls in apical and basal regions, and basal

mitochondria were larger than apical mitochondria (Figures 4E and 4G-4H). Fewer

mitochondria were found in apical regions in Drp1$^{SG}$ embryos compared to controls

(Figures 4E and 4F), consistent with immunostainings (Figure 4B). The circularity of

mitochondria was similar in Drp1$^{SG}$ expressing embryos and controls (Figures 4I and

4J). These data showed that mitochondrial fission regulated by Drp1 was essential for

regulating mitochondrial size and transport to apical regions in VF cells during

gastrulation.

**Drp1 mutant embryos show decreased apical constriction and Myosin II in the**

**ventral cells leading to a broader ventral furrow**.

VF formation and invagination are facilitated by the apical constriction of ventral cells

that follows a ratchet-like mechanism, with repeated pulses of constriction and

stabilization resulting in an incremental reduction in the apical cell area. These

contractions are driven by an intact acto-myosin network active apically in the VF cells.

Starting with discrete spots that intensify in the apical regions across VF cells, and pull

the adherence junctions together, Myosin II pulses help to constrict the VF cells [60,61].

We assessed the extent of apical constriction and Myosin II distribution in the ventral

cells in Drp1$^{SG}$ embryos expressing reporters of Myosin II light chain, Sqh-GFP and

plasma membrane, mScarletCAAX. Drp1$^{SG}$ embryos showed less apical constriction in

ventral cells compared to controls, with a similar furrow invagination (Figures 5A-C,

Videos 5 and 6). We compared the apical area and Myosin II intensity over time in 40

cells from one control and Drp1 mutant embryo. We observed a progressive decrease

in apical area and an increase in Myosin II intensity in controls (Figures 5B and D).

However, Drp1$^{SG}$ embryos showed an increase in the number of cells with reduced

apical constriction and Myosin II levels (Figure 5B and D). The overall apical area and

Myosin II intensity were significantly reduced in Drp1$^{SG}$ embryos compared to controls

(Figure 5C and E). We then visualized the overall VF morphology by imaging the

embryos in end-on orientation. We observed that Drp1$^{SG}$ embryos had a significantly

broader VF with a width of 12.4 (±2.33) µm compared to 8.6 (±0.23) µm in controls

(Figures 5F and 5G, Videos 7 and 8). The apical Myosin II intensity of VF cells in the

end-on view was also decreased in Drp1$^{SG}$ embryos compared to controls (Figures 5F

and H). These data suggested that reduced apical Myosin II in Drp1$^{SG}$ embryos leads to

an overall decrease in apical constriction in VF cells, resulting in a broader VF.

**RNA seq analysis shows no change in transcription of Dorsal/NFkB pathway**

**genes and an increased expression of anti-oxidant genes in Drp1 mutant**

**embryos**

Since we found that the Dorsal signaling pathway regulated mitochondrial dynamics in

early embryos (Figure 3), we evaluated whether the expression of Dorsal signaling

pathway genes and mesodermal genes was affected in Drp1$^{SG}$ expressing embryos, by

RNA sequencing. We sorted control and Drp1$^{SG}$ expressing embryos in cellularization

and early gastrulation for RNA extraction for sequencing. The principal component

analysis from the RNA sequencing output showed clustering of the three replicates of

control and Drp1$^{SG}$, and separation between the two genotypes (Figure S3A). In total,

1099 genes were differentially expressed in Drp1$^{SG}$ when compared with control

embryos (p value ≤0.05 and Log2FC≤-1 and ≥1). Amongst the differentially expressed

genes, 708 genes were up-regulated and 391 genes were down-regulated (Figure 6A).

Gene Ontology (GO) analysis performed on these differentially expressed genes was

categorized into Biological Process (BP) and Molecular Function (MF). BP enrichment

analysis revealed that upregulated genes were associated with 'biological regulation',

'multicellular organismal processes' and 'developmental processes'. This is in

accordance with another study in mice, where it has been shown that differentially

expressed genes (DEGs) in mitochondrial fission depleted embryonic stem cells are

enriched in 'multicellular organism development' and 'cell differentiation' [62].

Downregulated genes showed enrichment for pathways related to 'response to

stimulus', 'locomotion' and 'nervous system processes'. MF enrichment analysis

showed that upregulated genes were involved in 'catalytic activity', 'ion binding' and

'DNA-binding transcription factor activity'. Downregulated genes were enriched in

'transmembrane transporter activity', 'signaling receptor activity' and 'molecular

transducer activity' (Figure S3B). We examined the expression of Dorsal signaling

pathway genes and found no difference in their expression between control and Drp1$^{SG}$

embryos (Figure 6B). Further using immunostaining, we observed that the nuclear

localization of Dorsal and Twist signaling was unaltered in Drp1$^{SG}$ embryos (Figure 6C).

We also found that the expression of mesodermal genes, *SPE*, *T48*, *eve* and *MEF2*

was unchanged in Drp1$^{SG}$ mutant embryos as compared to controls (Figure 6D). The

expression of genes regulating the activation of Myosin II, *Rho1, Rok, RhoGEF2* and

*dia* was also similar in both controls and Drp1$^{SG}$ embryos (Figure S3C).

Mitochondria can modulate nuclear gene expression via metabolites such as

acetyl-CoA and NAD+, which contribute to epigenetic remodeling. Communication

between mitochondria and nucleus is also mediated by ATP, reactive oxygen species

(ROS) and Ca$^{2+}$ [63]. Importantly, we observed that the expression of anti-oxidant genes

such as *Sod3*, *Jafrac1*, *Trxr-2*, *GstD1*, *GstD3*, *GstE3*, *GstE6* and *GstE8* was up-

regulated in Drp1$^{SG}$ (Figures 6E and 6F). In addition, using quantitative RT-PCR, we

were able to verify an increase in the expression of antioxidant genes *Sod3*, *Jafrac1*

and *GstD1* in Drp1$^{SG}$ embryos (Figure S3D).

In summary, the Dorsal signaling pathway was upstream of mitochondrial

dynamics in regulating the VF for gastrulation. Change in mitochondrial fission led to an

increase in levels of anti-oxidant gene transcription, but no change in Dorsal signaling

pathway genes.

**Proteomic analysis shows an increased level of anti-oxidant proteins and a**

**decreased level of ETC complexes in Drp1 mutant embryos**

We performed proteome analysis using mass spectrometry of total and mitochondrial

protein extracts from control and Drp1$^{SG}$ expressing embryos to assess the change in

anti-oxidant proteins and ETC proteins. The principal component analysis showed that

the three replicates of control and Drp1$^{SG}$ clustered together, with distinct separation

between the two groups (Figure S4A). In the total proteomics, 3838 proteins were

detected, out of which 131 were found only in control embryos and 53 were detected

only in Drp1$^{SG}$ embryos. 729 proteins were differentially expressed (p value ≤0.05 and

Log2FC≤-0.3 and ≥0.3) with 312 being upregulated and 417 being downregulated

(Figure 7A). GO analysis was performed on the differentially expressed proteins. It was

categorized into Biological Process (BP) and Molecular Function (MF). BP enrichment

analysis revealed that upregulated proteins were associated with 'cellular localization'
and 'protein transport'. Downregulated proteins showed enrichment for pathways
related to 'metabolic processes', 'translation' and 'mitochondrial gene expression'. MF
enrichment analysis showed that upregulated proteins were involved in 'catalytic
activity' and various 'transmembrane transporter activity'. Downregulated proteins were
enriched in 'oxidoreductase activity' and 'NADH dehydrogenase activity' (Figure S4B).
We found that anti-oxidant proteins, Sod1, GstD3, GstE1, GstE6, GstE7 and Prx6c
were upregulated in Drp1$^{SG}$ (Figures 7B and 7C).

A total of 3303 proteins were detected from mitochondrial protein extracts, out of
which 30 were found only in control embryos and 126 were found only in Drp1$^{SG}$
embryos. Out of the 3147 common proteins, 1498 proteins were differentially expressed
with p value ≤0.05 and Log2FC≤-0.3 and ≥0.3. Amongst the differentially expressed
proteins, 783 were downregulated and 715 proteins were upregulated (Figure 7D). GO
terms were categorized into Biological Process (BP) and Molecular Function (MF)
domains. In the BP analysis, amongst the upregulated proteins, there was an
enrichment for 'biological regulation' and 'cellular localization and transport'.
Downregulated proteins were associated with 'metabolic processes' and 'mitochondrial
gene expression, translation and organization'. MF enrichment analysis revealed that
upregulated genes were enriched in 'protein binding' and 'catalytic activity'. The
downregulated proteins were involved in 'catalytic activity', 'oxidoreductase activity', and
'electron transfer activity' (Figure S4C). Interestingly, we found that many of the
subunits of the five ETC complexes were significantly downregulated in the
mitochondria of Drp1$^{SG}$ (Figures 7E and 7F), suggesting that the larger mitochondria
are metabolically less active. A similar downregulation of ETC complex subunits was
observed in the mass spectrometry of total proteins from Drp1$^{SG}$ embryos as well
(Figure 7A). Additionally, the levels of ATPbeta in mitochondria isolated from control
and Drp1$^{SG}$ expressing embryos using western blot analysis were confirmed to be
significantly reduced in Drp1$^{SG}$ (Figure S4D).

Together, the mass spectrometry analysis revealed an increase in anti-oxidant
protein levels and a decrease in mitochondrial ETC components, suggesting a
decrease in mitochondrial activity in Drp1$^{SG}$ embryos.

**Drp1 mutant embryos show reduced levels of ROS and Drp1$^{SG}$;SOD2$^i$ embryos**

**show rescue of anti-oxidant levels, ROS and mitochondrial morphology.**

ROS is produced in the mitochondria as a result of ETC activity primarily at complex I

and complex III [64]. We saw that anti-oxidant gene transcription and protein levels were

increased, and ETC subunits from many of the respiratory complexes were reduced in

Drp1$^{SG}$ embryos. We therefore estimated the levels of ROS in Drp1$^{SG}$ embryos using

the fluorescent dye dihydroethidium (DHE). DHE gets oxidised by superoxides forming

a fluorescent product that gives red fluorescence visualized using microscopy [65]. We

found that the mean fluorescence intensity of DHE was significantly lower in Drp1$^{SG}$

compared to controls (Figures 8A and 8B). Superoxides formed by the ETC get reduced

to $H_2O_2$ with the mitochondrial superoxide dismutase SOD2, and $H_2O_2$ can diffuse out

of the mitochondria. Superoxides can also transit into the cytoplasm via the voltage

gated anion channel (VDAC) from the mitochondria to the cytoplasm [66]. We checked if

there was an increase in overall ROS in embryos depleted of the anti-oxidant

mitochondrial superoxide dismutase (SOD2). We found that SOD2 RNAi (SOD2$^i$)

expressing embryos showed an increase in DHE fluorescence and ROS levels as

compared to controls. We made Drp1$^{SG}$;SOD2$^i$ combinations and assessed them for

ROS and found that the levels were restored and comparable to controls (Figures 8A

and 8B).

We then asked if combining Drp1$^{SG}$ and SOD2$^i$ could alleviate the gene

expression changes seen in Drp1$^{SG}$ embryos. Principal component analysis showed

that 2 out of 3 samples of Drp1$^{SG}$;SOD2$^i$ clustered between control and Drp1$^{SG}$,

suggesting a partial reversal of the transcriptional changes (Figure S3A). We examined

the expression of anti-oxidant genes that were upregulated in Drp1$^{SG}$ (log2FC > 1

compared to the control). We compared their log2FC values in Drp1$^{SG}$;SOD2$^i$ calculated

relative to control and found that the genes *Sod3, Jafrac1, Trxr-2, GstD1, GstD3, GstE6*

and *GstE8*, but not *GstE3*, showed a reduced log2FC compared to Drp1$^{SG}$ (Figures 8C

and 8D), indicating rescue of their expression levels upon depletion of SOD2 in Drp1

mutant embryos.

ROS has been shown to impact mitochondrial morphology. Increasing ROS

leads to mitochondrial fragmentation in *Drosophila* follicle cells and cells undergoing

delamination during *Drosophila* dorsal closure [67,68]. We therefore checked mitochondrial

distribution in Drp1$^{SG}$;SOD2$^{i}$ embryos in the VF. We observed a reduction in basal

mitochondrial clustering and a rescue of apical mitochondrial migration in the VF cells in

Drp1$^{SG}$;SOD2$^{i}$ embryos as compared to Drp1$^{SG}$ alone (Figure 8E). Oxidative stress

causes mitochondrial fragmentation and release of mitochondrial fusion protein, Opa1,

into the cytosol in mammalian HT22 cells [69]. Opa1 depletion also causes an increase in

ROS in *Drosophila* follicle cells and mitochondrial fragmentation in follicle cells and

embryos [15,67]. We therefore knocked down Opa1 in Drp1$^{SG}$ embryos and observed that

the mitochondrial migration and clustering defect was suppressed (Figure 8E).

Together, these data showed that Drp1 mutant embryos have decreased ROS and

increasing ROS supported mitochondrial fission for apical transport in VF cells during

gastrulation.

**Fragmenting mitochondria and increasing ROS rescues apical Myosin II and**

**ventral furrow formation in Drp1$^{SG}$;*opa1*$^{i}$ and Drp1$^{SG}$;SOD2$^{i}$ embryos.**

Drp1$^{SG}$ embryos showed a decrease in ROS, apical Myosin II and apical constriction

(Figures 5 and 8). The Drp1$^{SG}$;SOD2$^{i}$ combination showed increased ROS and rescue

of mitochondrial migration to apical regions (Figure 8). ROS has been shown to regulate

Myosin II in various processes such as dorsal closure [68], wound healing [70] and

*Drosophila* cellularization [15]. We therefore assessed Myosin II levels and VF

morphology in Sqh-mCherry expressing Drp1$^{SG}$;*opa1*$^{i}$ and Drp1$^{SG}$;SOD2$^{i}$ embryos

(Videos 9 and 10). Drp1$^{SG}$;*opa1*$^{i}$ showed a partial rescue in the apical Myosin II levels,

and the furrow width was 6.5 (±0.91) μm, which was restored compared to Drp1$^{SG}$ and

even smaller than controls. The apical Myosin II intensity of Drp1$^{SG}$;SOD2$^{i}$ was similar

to that of controls, and the furrow width was 7.3 (±1.69) μm, which was significantly

smaller compared to Drp1$^{SG}$. SOD2$^{i}$ embryos showed Myosin II levels and furrow width

(6.98±0.81 μm) comparable to the controls (Figures 9A-9C, Video 11). This data shows

that fragmenting mitochondria, either by decreasing fusion or increasing ROS, leads to

apical Myosin II localization and VF formation. In summary, our data collectively show

that mitochondria regulate VF furrow dynamics through ROS mediated regulation of

Myosin II contractility in VF cells.

**Discussion**

We demonstrate that small, punctate and fragmented mitochondria migrate from basal

to apical regions in ventral cells during ventral furrow formation in gastrulation in

*Drosophila* embryogenesis. Mitochondria migrate to apical regions in all cells during

cellularization [15]. However, migration in gastrulation occurs specifically at the VF in a

microtubule-dependent manner and, notably, is regulated by the Dorsal signaling

pathway. We find that maintaining mitochondrial size is essential to migrate, as larger

mitochondria in Drp1 mutant embryos are unable to migrate apically. Also interestingly,

reverting mitochondrial size with SOD2$^i$ and *opa1*$^i$ in Drp1$^{SG}$ embryos enables them to

migrate to apical regions. RNA seq analysis reveals that Drp1 mutant embryos do not

show a change in the Dorsal signaling pathway and mesodermal genes, but show an

increased expression of anti-oxidant genes. This confirms that the Dorsal signaling

pathway is upstream of mitochondrial migration. The mitochondria in Drp1 mutant

embryos have reduced activity due to a decrease in the levels of ETC components,

leading to a decrease in ROS. Since the embryos also have an increase in anti-oxidant

gene transcription, we conclude that they have an overall reduced environment. The

apical Myosin II levels are decreased in Drp1 mutant embryos during gastrulation,

resulting in reduced apical constriction, leading to larger ventral cells and broader VF.

Suppression of these defects upon fragmenting mitochondria and increasing ROS,

coincident with mitochondrial migration in the VF cells, indicates that apical

mitochondrial migration is essential for this morphogenetic process.

Mitochondria have been shown to exert their effect on the patterning of

embryonic axes in various organisms by their differential distribution and local changes

in activity. In sea urchin embryos, mitochondrial density and cytochrome oxidase activity

at the oral pole are higher, thereby establishing a respiratory gradient along the oral-

aboral (OA) axis, which is thought to activate the Nodal signaling pathway [37,39]. The

polarity of the OA axis can be altered by changing the respiratory gradient [71–73], and

dorsal activation of HIFα specifically contributes to dorso-ventral (DV) axis specification

[38]. A gradient of mitochondrial membrane potential with higher respiratory activity on the

ventral side has also been found in *Drosophila* blastoderm embryos [74]. In Xenopus

embryos, the cells of the dorsal blastopore lip in the Spemann-Mangold organizer

region undergo morphological changes in the form of apical constriction and mesoderm

internalization similar to the *Drosophila* VF. Multiple studies have reported the dorsal

region of the Xenopus embryos to have higher mitochondrial content and oxygen

consumption [35,36,75,76]. The organizer has been shown to be established by oxidative

metabolism driven by mitochondrial activity [36]. In mouse and zebrafish embryos,

mitochondrial ribosomal proteins and mitochondrial ATP synthesis have also been

shown to be critical for gastrulation, respectively [77,78]. Thus, the common theme that

appears from these studies and our findings is that there is a preferential mitochondrial

localization and/or activity in the embryonic regions of high cellular dynamics, which is

essential for driving cellular remodeling in these regions. Drp1 mutant embryos in our

study have reduced activity and metabolites, including ROS. It is also possible that

other metabolites, such as NAD+, are reduced, which not only changes the local

metabolism but also gives rise to epigenetic changes in the genome, further preserving

the reduced environment.

An interaction between mitochondria and Fog, a downstream effector in the

Dorsal signalling pathway, has been reported previously. Overexpression of Fog, in

*Drosophila* third instar larval muscles, leads to mitochondrial fission in a Drp1

dependent manner [79]. Fog is also enriched apically in the VF cells [80]. Thus, Fog activity

could maintain mitochondrial fission in ventral cells, facilitating apical mitochondrial

migration in the furrow. Downregulation of Drp1 inhibits Fog signalling and also leads to

VF defects, similar to Fog loss of function mutants [79]. Additionally, different isoforms of

Drp1 were reported in ventralized and lateralized embryos during gastrulation in

proteomic analysis. Knockdown of Drp1 using dsRNA resulted in mild to severe defects

in the VF [81]. However, the expression of Toll-Dorsal pathway genes and mesodermal

genes was unchanged in Drp1$^{SG}$, despite showing VF defects. This indicates that VF

invagination and mesoderm formation are regulated independently. It has also been

reported in a previous study that VF formation and mesoderm differentiation need not

be associated with each other. In the absence of Twist activity, Snail is sufficient to

induce VF invagination but not mesoderm differentiation [82]. It has also been shown in

sea urchin embryos that the gut and myocyte progenitors can differentiate even in the

absence of invagination in exogastrulae [83,84].

Mitochondrial shape and activity are closely associated with each other.

Changes in mitochondrial morphology have been shown to alter ROS levels [85] and

affect calcium homeostasis [86]. Increased fission can alter cell physiology and promote

cancer cell survival by increasing ROS production [87]. Mitochondrial fission, upon loss of

Opa1, leads to an increase in ROS levels [67,88–90]. It is conventionally believed that

increased mitochondrial fusion supports enhanced ETC activity [91,92]. However, some

studies have reported a decrease in ROS and mitochondrial activity upon Drp1

depletion, similar to our finding [93,94]. It is likely that the bigger mitochondria inherited

from oogenesis in Drp1$^{SG}$ embryos could be functionally compromised, resulting in

calcium imbalance and decreased ROS production from ETC complexes. ROS can act

as a signaling molecule and modify enzymes by oxidising cysteine residues [95], leading

to carbonylation of proteins [96]. Calcium can also regulate the cytoskeletal network

directly or via regulatory proteins. Myosin II light chain kinase (MLCK) is a serine-

threonine kinase that is activated by calmodulin, upon an increase in calcium levels,

and further it phosphorylates the regulatory Myosin II light chains, resulting in increased

contractility [97,98]. Intracellular calcium can drive remodeling of F-actin, thereby inducing

apical constriction leading to the formation of the neural tube in Xenopus [99]. ROS is

known to regulate kinases and phosphatases through oxidation [100,101]. ROS has been

shown to regulate two key Myosin II regulators, Rho kinase (ROCK) and Src kinase [68,70]

in *Drosophila* dorsal closure and zebrafish wound healing, respectively. It is possible

that the ROS levels in Drp1$^{SG}$;*opa1$^i$* might be increased due to mitochondrial fission and

thus the increased ROS, in Drp1$^{SG}$;SOD2$^i$ and Drp1$^{SG}$;*opa1$^i$*, could rescue the Myosin II

activity via upstream kinases and/or phosphatases. Thus, we conclude that

mitochondria migrate to the apical side in the VF cells to assist in Myosin II activation

via ROS in a spatiotemporal manner, thereby aiding VF formation. Additionally,

mitochondria could also regulate actin via ROS and/or calcium and could also be

essential for providing ATP or other metabolites for driving cell shape changes.

**Acknowledgements**

We thank IISER Pune microscopy and *Drosophila* facilities for help with imaging and *Drosophila* stocks and crosses. We thank RR lab members for constant input on this work. We thank FlyBase and Bloomington Stock Centers for the information and stocks used in this study. SM and SC thank the Council of Scientific and Industrial Research CSIR, India and the University Grants Commission UGC for the graduate student fellowship. HM thanks IISER Pune for the graduate student fellowship. We thank J-P. Chauvin, A. Aouane, and F. Richard at the Institut de Biologie du Développement de Marseille (IBDM) electron microscopy facility (Marseille, France) for help with embryo processing and image acquisition. We thank the Transmission Electron Microscope facility, ACTREC, Tata Memorial Centre, Kharghar, Navi Mumbai, India for help with sample processing, sectioning and imaging control and Drp1 mutant embryos. We thank Eric F. Wieschaus (Princeton University, USA), Yu-Chiun Wang (Riken, Kobe, Japan) and Girish Ratnaparkhi (IISER, Pune, India) for antibodies and fly stocks used in this study. We thank V Proteomics, Delhi, India for mass spectrometry analysis and Medgenome, Bangalore, India for RNA seq analysis.

**Author contribution**

SM, SC and RR designed the study, performed the experiments and analysed the data. SP and SM performed and analysed the western blots, RTPCR, genomics and proteomics. MM performed the experiment in Figure 1F-G, SC analysed the experiment. HM performed and analyzed the experiments in 4E-J. RR acquired the funding. SM and RR wrote the manuscript. SM, SC, HM, MM and RR edited the manuscript.

**Funding**

This project is funded by the Wellcome Trust-DBT India Alliance grant IA/S/22/1/506232.

**Declaration of interests**

1003    The authors declare no competing interests.

1004

1005    **Figure Legends**

1006

**Figure 1: Mitochondria are enriched apically in the ventral furrow cells during**

**gastrulation.**

(A) Schematic representing the *Drosophila* embryo in cellularization (left top sagittal

view, left bottom surface view) and gastrulation (right) and imaging orientations (shown

with box) used in fixed and live embryos to assess mitochondrial distribution and

dynamics in cellularization and gastrulation.

(B-C) Mitochondria are present as small punctae and enrich apically during

cellularization and gastrulation. Embryos in early, mid and late cellularization (B) and in

gastrulation at the VF (C) are labelled for mitochondria (fluorescent streptavidin, green),

F-actin (fluorescent phalloidin, red) and DNA (Hoechst, blue).

(D) Representative snapshots from live embryos expressing mito-GFP in late

cellularization (first time point) and during VF formation in gastrulation. Dorsal is up and

ventral is down. The images are shown in 16 color rainbow scale on an 8 bit image

where 0 is black and white is 255.

(E) The plot shows apical:basal mean fluorescence intensity of mito-GFP quantified in

three regions of the gastrulating embryos, ventral cells (red), neighbouring cells (green)

and lateral cells (blue), across 5 time points (n=3). The schematic of gastrulating

embryo shows the regions chosen for quantification. Data is represented as Mean ±

SEM.

(F) Representative TEM images from gastrulating embryos are shown in the lateral and

ventral region. Mitochondria are dark punctate structures enriched in apical regions.

Zoomed in images show mitochondria enriched in apical regions of ventral cells

compared to lateral cells.

(G) The plot shows quantification of the number of mitochondria above the nucleus per

cell in ventral and lateral cells (n= 14 and 13 cells, respectively from 4 embryos). The

violin plot shows median and quartiles. The statistics were performed by two-tailed

unpaired Mann-Whitney test, ****$p < 0.0001$ Scale Bar: 10µm (B, C, D), 5µm (F).

1034

**Figure 2. Mitochondria translocate apically from basal sections in the ventral furrow cells in a microtubule-dependent manner.**

(A) Representative embryo expressing mito-PAGFP and Sqh-mCherry was photoactivated in the white ROI at the end of cellularization or the beginning of gastrulation. Representative snapshots are shown from the movie before and after photoactivation (T1-T5). Zoomed in images show mitochondria being translocated from basal to apical regions in ventral cells on linear tracks.

(B) The plot shows mean apical:basal fluorescence intensity of mitochondria and Myosin II measured in ventral and lateral cells (n=3), as shown in the schematic. Data is represented as Mean ± SEM.

(C) Representative images of mito-GFP expressing embryos treated with colchicine (n=6) and the respective control (n=3). Colchicine treated embryos show reduced mitochondrial migration in the ventral cells and clustering of mitochondria in the basal regions (yellow arrowhead).

(D) The plot shows mean mito-GFP fluorescence intensity of the ventral cells expressed as a ratio of apical to basal regions. The violin plot shows median and quartiles. The statistics were performed by two-tailed unpaired Mann-Whitney test, *p<0.05. Scale Bar: 10μm (A,C).

**Figure 3. Dorsal regulates apical translocation of mitochondria.**

(A-C) Dorsal regulates apical translocation of mitochondria during cellularization. (A) Representative optical planes from grazing sections of apical, basal regions and sagittal sections of control, *dl*RNAi, *dl*$^4$/*dl*$^1$ and *dl*$^1$/+ embryos stained for mitochondria (fluorescent streptavidin, green) and DNA (Hoechst, blue) are shown.

(B) The plot shows quantification of the relative area occupied by mitochondria in apical sections from *dl*RNAi (n=7), *dl*$^4$/*dl*$^1$ (n=7) and *dl*$^1$/+ embryos (n=6) and control (n=7) embryos.

(C) The plot shows mean mitochondrial size quantified from a single optical plane in the apical (A) and basal (B) section. Number of embryos in control apical n=16, basal n=17;

*dl*RNAi apical n=14, basal n=16; *dl$^4$/dl$^1$* apical n=27, basal n=15 and *dl$^1$/+* apical n=6,

basal n=6.

(D) Representative snapshots from live movies of gastrulating control, *dl$^1$/+* and Dorsal

overexpression (Dl-OE) embryos expressing mito-GFP are shown in the 16 color

rainbow scale on an 8 bit image with black as 0 and white as 255. (E) The plot shows

mean fluorescence intensity from apical sections as a ratio to basal sections at the VF

in *dl$^1$/+* (n=4) and control (n=3) embryos. Data is represented as Mean ± SEM.

(F) The plot shows mean mitochondrial intensity from mito-GFP at apical regions

starting from the ventral midpoint (0) to the dorsal side in controls (n=3)  and Dl-OE

(n=6). Data is represented as Mean ± SEM. The violin plots show median and quartiles.

The statistics were performed by two-tailed unpaired Mann-Whitney test, ns p≥0.05,

*p<0.05, **p<0.01. Scale Bar: 10µm (A,D).

1076

**Figure 4. Drp1$^{SG}$ embryos show basal clustering and reduced apical translocation**

**of mitochondria.**

(A) Representative images of control and Drp1$^{SG}$ embryos with VF, stained for

mitochondria (fluorescent streptavidin, green), F-actin (fluorescent phalloidin, red) and

DNA (Hoechst, blue) are shown. The white arrowhead marks the mitochondria in the

respective embryos.

(B-D) Drp1 localization and expression in control and Drp1$^{SG}$ embryos. (B) Control and

Drp1$^{SG}$ cellularizing embryos are stained with Drp1 antibody (red) and for mitochondria

(fluorescent streptavidin, green). (C) The western blot and graph show Drp1 in control

cellularization and gastrulation embryos (N=3). β-actin is used as the loading control.

(D) The western blot and graph show Drp1 in control and Drp1$^{SG}$ embryos, with β-actin

used as a loading control (N=3).

(E) Representative TEM images of late cellularizing control and Drp1$^{SG}$ embryos are

shown in sagittal view (1000x magnification) and cross-sectional view (5000x

magnification). Mitochondria are electron-dense structures, highlighted with yellow ROI.

(F) The plot shows a quantification of mitochondria above the nucleus per cell in control

(n=4) and Drp1$^{SG}$ (n=4)embryos.

(G) The plot shows the size of mitochondria in apical regions in control (52 mitochondria

from 4 embryos) and Drp1$^{SG}$ embryos (21 mitochondria from 4 embryos).

(H) The plot shows the size of mitochondria in basal regions in control (34 mitochondria

from 4 embryos) and Drp1$^{SG}$ embryos (68 mitochondria from 4 embryos).

(I) The plot shows the circularity of mitochondria in apical regions in control (52

mitochondria from 4 embryos) and Drp1$^{SG}$ embryos (21 mitochondria from 4 embryos).

(J) The plot shows the circularity of mitochondria in basal regions in control (34

mitochondria from 4 embryos) and Drp1$^{SG}$ embryos (68 mitochondria from 4 embryos).

The violin plots show median and quartiles. The statistics were performed by two-tailed

unpaired Mann-Whitney test, ns p≥0.05, **p<0.01, ****p<0.0001. Scale Bar: 10µm

(A,B), 5µm (E, sagittal), 1µm (E, apical, basal).

1105

**Figure 5. Drp1$^{SG}$ embryos show decreased apical constriction and Myosin II in the**

**ventral cells resulting in a broader ventral furrow.**

(A) Representative images of live control and Drp1$^{SG}$ embryos, expressing

SqhGFP;ScarletCAAX, undergoing apical constriction in ventral cells are shown.

Orthogonal views show the extent of furrow ingression. Cells are color coded (right side

images) based on area.

(B) Heat map of one representative control and Drp1$^{SG}$ embryo shows the area of 40

cells (y axis: A-NA) with time.

(C) The plot shows the normalized mean apical area of control and Drp1$^{SG}$ embryos

(n=3 embryos each, 40 cells from each embryo). Data is represented as Mean ± SEM.

(D) Heat map of one representative control and Drp1$^{SG}$ embryo shows the Myosin II

intensity of 40 cells (y axis: A-NA) with time.

(E) The plot shows the normalized mean apical Myosin II intensity of control and Drp1$^{SG}$

embryos (n=3 embryos each, 40 cells from each embryo). Data is represented as Mean

± SEM.

(F) Representative images, in end-on view, of live control and Drp1$^{SG}$ embryos

expressing Sqh-mCherry during VF formation are shown. The red dotted line shows the

furrow width measurement.

(G) The plot shows the width of the VF in control (n=4) and Drp1$^{SG}$ (n=4).

(H) The plot shows the normalized mean myosin II intensity during VF formation in

control (n=3) and Drp1$^{SG}$ (n=3). Data is represented as Mean ± SEM. The violin plot

shows median and quartiles. The statistics were performed by two-tailed unpaired

Mann-Whitney test, *p<0.05. Scale bar 10µm (A,F).

1129

**Figure 6. Drp1$^{SG}$ embryos show no change in Dorsal pathway genes, but**

**increased expression of anti-oxidant genes**.

(A) Volcano plot shows the distribution of genes with the log2foldchange on x axis and

significance (-log10pvalue) on y axis. Every point in the plot represents a single gene.

All the genes with p value>0.05 and log2foldchange between -1 to +1 are similar in

expression in control (N=3) and Drp1$^{SG}$ (N=3), and are marked in grey. Genes marked

in green represent the significantly upregulated genes and those in pink indicate the

significantly downregulated genes. Top 10 differentially expressed genes (DEGs) and

some other genes of interest are highlighted. Bar plot showing the number of

significantly upregulated and downregulated genes is represented on the top left corner.

(B) Heat map shows the relative expression of Toll-Dorsal pathway genes in control and

Drp1$^{SG}$ embryos.

(C) Representative sagittal views of control and Drp1$^{SG}$ cellularizing embryos stained

for Dorsal (magenta), Twist (red), mitochondria (green) and DNA (blue) are shown.

(D) Heat map shows the relative expression of mesodermal genes in control and

Drp1$^{SG}$ embryos.

(E) Heat map shows the relative expression of anti-oxidant genes in control and Drp1$^{SG}$

embryos.

(F) Table shows the log2FoldChange of the anti-oxidant genes in Drp1$^{SG}$ compared to

control.

1150

**Figure 7. Drp1$^{SG}$ embryos show increased levels of anti-oxidant proteins and**

**downregulation of ETC complexes.**

(A) Volcano plot shows the distribution of all the proteins identified in the total protein

mass spec, with the log2foldchange on the x axis and significance (-log10pvalue) on

the y axis. Every point in the plot corresponds to a single protein. All the proteins with p

value>0.05 and log2foldchange between -0.3 to +0.3 are marked in grey and are similar

in expression in control (N=3) and Drp1$^{SG}$ (N=3) embryos. Green dots represent the

upregulated proteins, and pink dots represent the downregulated proteins. Top 10

differentially expressed proteins and some other proteins of interest are highlighted. Bar

plot showing the number of significantly upregulated and downregulated proteins is

represented on the top left corner.

(B) Heat map shows the relative expression of anti-oxidant proteins in control and

Drp1$^{SG}$ embryos.

(C) Table shows the log2FoldChange of the anti-oxidant proteins in Drp1$^{SG}$ compared

to control.

(D) Volcano plot shows the distribution of all the proteins identified in the mitochondrial

protein mass spec. log2foldchange has been plotted on the x axis and significance (-

log10pvalue) has been plotted on the y axis. Every point in the plot represents a

protein. All the proteins with p value>0.05 and log2foldchange between -0.3 to +0.3 are

similar in expression in control (N=3) and Drp1$^{SG}$ (N=3) and are marked in grey.

Upregulated proteins are marked in green and downregulated proteins are marked in

pink. Top 10 differentially expressed proteins and some other proteins of interest have

been highlighted. Bar plot showing the number of significantly upregulated and

downregulated proteins is represented on the top left corner.

(E) Heat map shows the relative expression of some of the subunits of all five

respiratory complexes in control and Drp1$^{SG}$ embryos.

(F) Table shows the log2FoldChange of the subunits of all five respiratory complexes in

Drp1$^{SG}$ compared to control.

1179

**Figure 8. Drp1$^{SG}$ embryos show reduced levels of ROS and Drp1$^{SG}$;SOD2$^{i}$**

**embryos show rescue of anti-oxidant levels, ROS and mitochondrial morphology.**

(A) Representative images of control, Drp1$^{SG}$, Drp1$^{SG}$;SOD2$^{i}$ and SOD2$^{i}$ cellularizing

embryos, at a single plane just above the nucleus, stained with DHE (red) and Hoechst

(blue) are shown. Zoomed in view of the cells is shown in rainbow scale (16 colors) as

an inset to highlight the relative difference in intensities.

(B) The plot shows the normalized mean fluorescence intensity of DHE. n=6 for control,

n=11 for Drp1$^{SG}$, n=7 for SOD2$^i$, n=7 for Drp1$^{SG}$;SOD2$^i$.

(C) Heat map shows the relative expression of anti-oxidant genes in control, Drp1$^{SG}$,

Drp1$^{SG}$;SOD2$^i$ and SOD2$^i$ embryos.

(D) Table shows the log2FoldChange values of the anti-oxidant genes, each with

respect to control.

(E) Representative images of Drp1$^{SG}$, Drp1$^{SG}$;*opa1*$^i$ and Drp1$^{SG}$;SOD2$^i$ and SOD2$^i$

gastrulating embryos with VF stained for DNA (Hoechst, blue), F-actin (fluorescent

phalloidin, red) and mitochondria (fluorescent streptavidin, green). White arrowhead

marks the mitochondria in respective embryos. The violin plot shows median and

quartiles. The statistics were performed by two-tailed unpaired Mann-Whitney test, ns

p≥0.05, *p<0.05, **p<0.01, ***p<0.001. Scale Bar 10μm (A,E).

1198

**Figure 9. Drp1$^{SG}$;*opa1*$^i$ and Drp1$^{SG}$;SOD2$^i$ embryos show a rescue of apical**

**Myosin II and ventral furrow formation.**

(A) Representative images in end-on view from live embryos (control, Drp1$^{SG}$,

Drp1$^{SG}$;*opa1*$^i$, Drp1$^{SG}$;SOD2$^i$ and SOD2$^i$) expressing Sqh-mCherry, starting from the

end of cellularization (0 min) till invagination of the VF are shown. The red dotted line

shows the measurement of the furrow width.

(B) The plot shows the furrow width, measured at the time point 4 when the VF is U

shaped. n=4 for control, n=4 for Drp1$^{SG}$, n=3 for SOD2$^i$, n=3 for Drp1$^{SG}$;SOD2$^i$, n=4 for

Drp1$^{SG}$;*opa1*$^i$ embryos

(C) The plot shows the normalized mean apical fluorescence of Sqh-mCherry in the VF

at the 5 time points. n=4 for control, n=4 for Drp1$^{SG}$, n=4 for SOD2$^i$, n=3 for

Drp1$^{SG}$;SOD2$^i$, n=4 for Drp1$^{SG}$;*opa1*$^i$ embryos. Data is represented as Mean ± SEM.

The violin plot shows median and quartiles. The statistics were performed by two-tailed

unpaired Mann-Whitney test, ns p≥0.05, *p<0.05.  Scale 10μm (A).

1213

**Figure 10. Summary Model.**

Mitochondria (green) migrate apically in the VF with the help of microtubules (red lines),

resulting in their apical enrichment. The apical mitochondrial migration is regulated by

Dorsal (blue nuclear gradient). Drp1 mutant embryos (Drp1$^{SG}$) display larger

mitochondria, which remain at the basal side (green) and are unable to migrate apically,

resulting in few smaller mitochondria in the apical regions. Expression of Dorsal

signaling pathway genes and mesodermal genes is not affected in Drp1$^{SG}$. Anti-oxidant

genes are upregulated in Drp1$^{SG}$ (shown by transcription of red triangle inside the

nucleus) in Drp1$^{SG}$. Levels of ETC complexes subunits and ROS (ROS, pink dots) are

reduced in Drp1$^{SG}$. Myosin II levels at the VF are decreased (yellow), resulting in a

broader VF. Knockdown of SOD2 and *opa1* in Drp1$^{SG}$ results in fragmentation of

mitochondria and rescue of apical mitochondrial migration in the VF. Levels of anti-

oxidants and ROS are also rescued in Drp1$^{SG}$;SOD2$^{i}$. Myosin II levels at the VF were

restored in both Drp1$^{SG}$;SOD2$^{i}$ and Drp1$^{SG}$;*opa1$^{i}$*, resulting in rescue of the VF

formation defect.

1229

**Supplementary Figure Legends**

**Figure. S1 Dorsal overexpression increases Dorsal and Twist expression**

(A) Representative images of control and Dl-OE embryos in cellularization, stained for

Dorsal, Twist and Hoechst (DNA, blue) are shown. Scale bar 10µm.

(B) The plot shows the Ventral:Dorsal intensity ratio of Dorsal and Twist in control (n=7,

6 respectively) and Dl-OE (n=4, 5 respectively). The violin plot shows median and

quartiles. The statistics were performed by two-tailed unpaired Mann-Whitney test,

**p<0.01.

(C) Representative snapshots from live embryos overexpressing Dorsal (Dl-OE) and

mito-GFP are shown in gastrulation. The 16 color rainbow scale is used to represent

fluorescence intensities of mito-GFP on a 8 bit scale where 0 is black and 255 is white.

1241

1242  **Figure S2. Western blot with anti-Drp1**

1243  (A) Western blot with anti-Drp1 in *Drosophila* embryos in cellularization (n=3)

1244

**Figure S3. RNA seq analysis and qPCR of Drp1 mutant embryos**

(A) Principal component analysis (PCA) of control, Drp1$^{SG}$, SOD2$^{i}$, Drp1$^{SG}$;SOD2$^{i}$

embryos in RNA sequencing is shown (n=3 for each genotype).

(B) Gene ontology (GO) analysis of upregulated and downregulated genes in control

and Drp1$^{SG}$ is shown.

(C) Heat map shows the relative expression of the genes that regulate Myosin II in

control and Drp1$^{SG}$ embryos.

(D) The plot shows the relative mRNA expression of 3 anti-oxidant genes- *Jafrac1*,

*Sod3* and *GstD1* (n=3). The statistics were performed by two-tailed unpaired Mann-

Whitney test, **p<0.01. ***p<0.001.

1255

**Figure S4. Total proteomics and mitochondrial proteomics analysis of Drp1**

**mutant embryos**

(A) Principal component analysis of control (C) and Drp1$^{SG}$ (D) embryos in total

proteomics and mitochondrial proteomics is shown (n=3 for each genotype).

(B) Gene ontology (GO) analysis of upregulated and downregulated proteins in total

proteomics in control and Drp1$^{SG}$ is shown.

(C) Gene ontology (GO) analysis of upregulated and downregulated proteins in

mitochondrial proteomics in control and Drp1$^{SG}$ is shown.

(D) Relative expression of ATPbeta in mitochondria extracts of control and Drp1$^{SG}$

embryos is shown using western blots (left). Coomassie staining is shown for equal

protein loading (centre). The plot shows the normalized intensity of ATPbeta in control

and Drp1$^{SG}$ embryos (n=3). The violin plot shows median and quartiles. The statistics

were performed by two-tailed unpaired Mann-Whitney test, ns p≥0.05.

1269

1270 **Video legends**

1271

1272 **Video 1.** Mitochondrial dynamics in mito-GFP expressing embryo undergoing ventral

1273 furrow formation, shown in 16 color rainbow scale. Time is shown in minutes and

1274 seconds, and the scale bar is 10 µm.

1275

1276 **Video 2.** Mitochondrial dynamics and Myosin II distribution in mito-PAGFP, Sqh-

1277 mCherry expressing embryo. Time is shown in minutes and seconds, and the scale bar

1278 is 10 µm.

1279

1280 **Video 3.** Gastrulation is shown in heterozygous $dl^1$ mutant embryos expressing mito-

1281 GFP, shown in 16 color rainbow scale. Time is shown in minutes and seconds, and the

1282 scale bar is 10 µm.

1283

1284 **Video 4.** Gastrulation is shown in UASp-Dl embryos expressing mito-GFP, shown in 16

1285 color rainbow scale. Time is shown in minutes and seconds, and the scale bar is 10 µm.

1286

1287 **Video 5.** Apical constriction at the VF in Sqh-GFP;mScarletCAAX expressing embryos.

1288 Time is shown in minutes and seconds, and the scale bar is 10 µm.

1289

1290 **Video 6.** Apical constriction at the VF in Drp1$^{SG}$ embryos expressing Sqh-

1291 GFP;mScarletCAAX. Time is shown in minutes and seconds, and the scale bar is 10

1292 µm.

1293

1294 **Video 7.** VF formation is shown in end-on view in Sqh-mCherry expressing embryos.

1295 Time is shown in minutes and seconds, and the scale bar is 10 µm.

1296

1297 **Video 8.** VF formation is shown in end-on view in Drp1$^{SG}$ embryos expressing Sqh-

1298 mCherry. Time is shown in minutes and seconds, and the scale bar is 10 µm.

1299

1300 **Video 9.** VF formation is shown in end-on view in Drp1$^{SG}$;*opa1*$^i$ embryos expressing

1301 Sqh-mCherry. Time is shown in minutes and seconds, and the scale bar is 10 µm.

1302

1303 **Video 10.** VF formation is shown in end-on view in Drp1$^{SG}$;*SOD2*$^i$ embryos expressing

1304 Sqh-mCherry. Time is shown in minutes and seconds, and the scale bar is 10 µm.

1305

1306 **Video 11.** VF formation is shown in end-on view in *SOD2*$^i$ embryos expressing Sqh-

1307 mCherry. Time is shown in minutes and seconds, and the scale bar is 10 µm.

1308

**References**

1.    Madan, S., Uttekar, B., Chowdhary, S., and Rikhy, R. (2021). Mitochondria lead the
way: Mitochondrial dynamics and function in cellular movements in development
and disease. Front. Cell Dev. Biol. *9*, 781933.

2.    Dumollard, R., Carroll, J., Duchen, M.R., Campbell, K., and Swann, K. (2009).
Mitochondrial function and redox state in mammalian embryos. Semin. Cell Dev.
Biol. *20*, 346–353.

3.    May-Panloup, P., Boguenet, M., Hachem, H.E., Bouet, P.-E., and Reynier, P.
(2021). Embryo and its mitochondria. Antioxidants (Basel) *10*, 139.

4.    Zhang, L., Zhang, Y., Chang, X., and Zhang, X. (2020). Imbalance in mitochondrial
dynamics induced by low PGC-1α expression contributes to hepatocyte EMT and
liver fibrosis. Cell Death Dis. *11*, 226.

5.    Finkel, T., and Hwang, P.M. (2009). The Krebs cycle meets the cell cycle:
mitochondria and the G1-S transition. Proc. Natl. Acad. Sci. U. S. A. *106*, 11825–
11826.

6.    Mitra, K., Wunder, C., Roysam, B., Lin, G., and Lippincott-Schwartz, J. (2009). A
hyperfused mitochondrial state achieved at G1-S regulates cyclin E buildup and
entry into S phase. Proc. Natl. Acad. Sci. U. S. A. *106*, 11960–11965.

7.    Martínez-Diez, M., Santamaría, G., Ortega, A.D., and Cuezva, J.M. (2006).
Biogenesis and dynamics of mitochondria during the cell cycle: significance of
3'UTRs. PLoS One *1*, e107.

8.    Kim, H.J., Shaker, M.R., Cho, B., Cho, H.M., Kim, H., Kim, J.Y., and Sun, W.
(2015). Dynamin-related protein 1 controls the migration and neuronal
differentiation of subventricular zone-derived neural progenitor cells. Sci. Rep. *5*,
15962.

9.    Lim, S., Lee, S.-Y., Seo, H.-H., Ham, O., Lee, C., Park, J.-H., Lee, J., Seung, M.,
Yun, I., Han, S.M., et al. (2015). Regulation of mitochondrial morphology by
positive feedback interaction between PKCδ and Drp1 in vascular smooth muscle

cell. J. Cell. Biochem. *116*, 648–660.

10.  Wang, L., Yu, T., Lee, H., O'Brien, D.K., Sesaki, H., and Yoon, Y. (2015).
Decreasing mitochondrial fission diminishes vascular smooth muscle cell migration
and ameliorates intimal hyperplasia. Cardiovasc. Res. *106*, 272–283.

11.  Fu, H., Zhou, H., Yu, X., Xu, J., Zhou, J., Meng, X., Zhao, J., Zhou, Y., Chisholm,
A.D., and Xu, S. (2020). Wounding triggers MIRO-1 dependent mitochondrial
fragmentation that accelerates epidermal wound closure through oxidative
signaling. Nat. Commun. *11*, 1050.

12.  Xu, S., and Chisholm, A.D. (2014). C. elegans epidermal wounding induces a
mitochondrial ROS burst that promotes wound repair. Dev. Cell *31*, 48–60.

13.  Giles, R.E., Blanc, H., Cann, H.M., and Wallace, D.C. (1980). Maternal inheritance
of human mitochondrial DNA. Proc. Natl. Acad. Sci. U. S. A. *77*, 6715–6719.

14.  Jonathan, P.W., and Davis, J. (1995). Fertilization and Early Embryolgoy: ATP
Content of Human Oocytes and Developmental Potential and Outcome after in-
Vitro Fertilization and Embryo Transfer. Human Reproduction.

15.  Chowdhary, S., Madan, S., Tomer, D., Mavrakis, M., and Rikhy, R. (2020).
Mitochondrial morphology and activity regulate furrow ingression and contractile
ring dynamics in Drosophila cellularization. Mol. Biol. Cell, mbcE20030177.

16.  Chowdhary, S., Tomer, D., Dubal, D., Sambre, D., and Rikhy, R. (2017). Analysis
of mitochondrial organization and function in the Drosophila blastoderm embryo.
Sci. Rep. *7*, 5502.

17.  Cox, R.T., and Spradling, A.C. (2003). A Balbiani body and the fusome mediate
mitochondrial inheritance during Drosophila oogenesis. Development *130*, 1579–
1590.

18.  Wilding, M., Dale, B., Marino, M., di Matteo, L., Alviggi, C., Pisaturo, M.L.,
Lombardi, L., and De Placido, G. (2001). Mitochondrial aggregation patterns and

activity in human oocytes and preimplantation embryos. Hum. Reprod. *16,* 909–

917.

19.  Heng-Kien, T.-S., Yeh, S.-H., Kao, C.-R., and Tzeng, R.-H. (2005). Abnormal

Mitochondrial Structure in Human Unfertilized Oocytes and Arrested Embryos.

Annals of the New York Academy of Sciences *1042*.

20.  Arribat, Y., Grepper, D., Lagarrigue, S., Richard, J., Gachet, M., Gut, P., and Amati,

F. (2019). Mitochondria in embryogenesis: An organellogenesis perspective. Front.

Cell Dev. Biol. *7,* 282.

21.  Bleazard, W., McCaffery, J.M., King, E.J., Bale, S., Mozdy, A., Tieu, Q., Nunnari,

J., and Shaw, J.M. (1999). The dynamin-related GTPase Dnm1 regulates

mitochondrial fission in yeast. Nat. Cell Biol. *1*, 298–304.

22.  Labrousse, A.M., Zappaterra, M.D., Rube, D.A., and van der Bliek, A.M. (1999). C.

elegans dynamin-related protein DRP-1 controls severing of the mitochondrial

outer membrane. Mol. Cell *4*, 815–826.

23.  Sesaki, H., and Jensen, R.E. (1999). Division versus fusion: Dnm1p and Fzo1p

antagonistically regulate mitochondrial shape. J. Cell Biol. *147*, 699–706.

24.  Hsiuchen, S.A., Detmer, A.J., Ewald, E.E., Griffin, S.E., and Fraser, D.C. (2003).

Mitofusins Mfn1 and Mfn2 Coordinately Regulate Mitochondrial Fusion and Are

Essential for Embryonic Development. The Journal of Cell Biology *160*, 189–200.

25.  Ban, T., Kohno, H., Ishihara, T., and Ishihara, N. (2018). Relationship between

OPA1 and cardiolipin in mitochondrial inner-membrane fusion. Biochim. Biophys.

Acta Bioenerg. *1859,* 951–957.

26.  Olichon, A., Baricault, L., Gas, N., Guillou, E., Valette, A., Belenguer, P., and

Lenaers, G. (2003). Loss of OPA1 perturbates the mitochondrial inner membrane

structure and integrity, leading to cytochrome c release and apoptosis. J. Biol.

Chem. *278,* 7743–7746.

27.  Lieber, T., Jeedigunta, S.P., Palozzi, J.M., Lehmann, R., and Hurd, T.R. (2019).
Mitochondrial fragmentation drives selective removal of deleterious mtDNA in the
germline. Nature *570*, 380–384.

28.  Moore, B.A., Gonzalez Aviles, G.D., Larkins, C.E., Hillman, M.J., and Caspary, T.
(2010). Mitochondrial retention of Opa1 is required for mouse embryogenesis.
Mamm. Genome *21*, 350–360.

29.  Mitra, K., Rikhy, R., Lilly, M., and Lippincott-Schwartz, J. (2012). DRP1-dependent
mitochondrial fission initiates follicle cell differentiation during Drosophila
oogenesis. J. Cell Biol. *197*, 487–497.

30.  Ishihara, N., Nomura, M., Jofuku, A., Kato, H., Suzuki, S.O., Masuda, K., Otera, H.,
Nakanishi, Y., Nonaka, I., Goto, Y.-I., et al. (2009). Mitochondrial fission factor Drp1
is essential for embryonic development and synapse formation in mice. Nat. Cell
Biol. *11*, 958–966.

31.  Wakabayashi, J., Zhang, Z., Wakabayashi, N., Tamura, Y., Fukaya, M., Kensler,
T.W., Iijima, M., and Sesaki, H. (2009). The dynamin-related GTPase Drp1 is
required for embryonic and brain development in mice. J. Exp. Med. *206*, i23–i23.

32.  Lees, J.G., Gardner, D.K., and Harvey, A.J. (2017). Pluripotent stem cell
metabolism and mitochondria: Beyond ATP. Stem Cells Int. *2017*, 2874283.

33.  Gut, P., and Verdin, E. (2013). The nexus of chromatin regulation and intermediary
metabolism. Nature *502*, 489–498.

34.  Wakefield, S.L., Lane, M., and Mitchell, M. (2011). Impaired mitochondrial function
in the preimplantation embryo perturbs fetal and placental development in the
mouse. Biol. Reprod. *84*, 572–580.

35.  Yost, H.J., Phillips, C.R., Boore, J.L., Bertman, J., Whalon, B., and Danilchik, M.V.
(1995). Relocation of mitochondria to the prospective dorsal marginal zone during
Xenopus embryogenesis. Dev. Biol. *170*, 83–90.

36.   Garfinkel, M., Mnatsakanyan, N., Patel, J.H., Wills, A.E., Shteyman, A., Peter, J.S.,
Kambiz, N., Alavian, E.A., and Jonas, M.K. (2023). Mitochondrial Leak Metabolism
Induces the Spemann-Mangold Organizer via Hif-1α in Xenopus. Developmental
Cell *58*.

37.   Czihak, G. (1963). 272-292 10.1007/BF00582031 Czihak Gerhard G Stazione
Zoologica, Napoli. Max-Planck-Institut für Meeresbiologie Abt. Bauer, Tübingen.
ger Journal Article Entwicklungsphysiologische Untersuchungen an Echiniden :
Verteilung und Bedeutung der Cytochromoxydase. Wilhelm Roux Arch. Entwickl.
Mech. Org. *154*, 272–292.

38.   Chang, W.-L., Chang, Y.-C., Lin, K.-T., Li, H.-R., Pai, C.-Y., Chen, J.-H., and Su,
Y.-H. (2017). Asymmetric distribution of hypoxia-inducible factor α regulates
dorsoventral axis establishment in the early sea urchin embryo. Development *144*,
2940–2950.

39.   Coffman, J.A. (2009). Mitochondria and metazoan epigenesis. Semin. Cell Dev.
Biol. *20*, 321–329.

40.   Turner, F.R., and Mahowald, A.P. (1977). Scanning electron microscopy of
Drosophila melanogaster embryogenesis. II. Gastrulation and segmentation. Dev.
Biol. *57*, 403–416.

41.   Leptin, M., and Grunewald, B. (1990). Cell shape changes during gastrulation in
Drosophila. Development *110*, 73–84.

42.   Gheisari, E., Aakhte, M., and Müller, H.-A.J. (2020). Gastrulation in Drosophila
melanogaster: Genetic control, cellular basis and biomechanics. Mech. Dev. *163*,
103629.

43.   Martin, A.C. (2020). The physical mechanisms of Drosophila gastrulation:
Mesoderm and endoderm invagination. Genetics *214*, 543–560.

44.   Ko, C.S., and Martin, A.C. (2020). The cellular and molecular mechanisms that
establish the mechanics of Drosophila gastrulation. Curr. Top. Dev. Biol. *136*, 141–

165.

45. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T.,
Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-
source platform for biological-image analysis. Nat. Methods *9*, 676–682.

46. Stringer, C., Wang, T., Michaelos, M., and Pachitariu, M. (2021). Cellpose: a
generalist algorithm for cellular segmentation. Nat. Methods *18*, 100–106.

47. Zhao, J., Zhang, J., Yu, M., Xie, Y., Huang, Y., Wolff, D.W., Abel, P.W., and Tu, Y.
(2013). Mitochondrial dynamics regulates migration and invasion of breast cancer
cells. Oncogene *32*, 4814–4824.

48. Desai, S.P., Bhatia, S.N., Toner, M., and Irimia, D. (2013). Mitochondrial
localization and the persistent migration of epithelial cancer cells. Biophys. J. *104*,
2077–2088.

49. Pilling, A.D., Horiuchi, D., Lively, C.M., and Saxton, W.M. (2006). Kinesin-1 and
Dynein are the primary motors for fast transport of mitochondria in Drosophila
motor axons. Mol. Biol. Cell *17*, 2057–2068.

50. Ko, C.S., Tserunyan, V., and Martin, A.C. (2019). Microtubules promote
intercellular contractile force transmission during tissue folding. J. Cell Biol. *218*,
2726–2742.

51. Belvin, M.P., and Anderson, K.V. (1996). A conserved signaling pathway: the
Drosophila toll-dorsal pathway. Annu. Rev. Cell Dev. Biol. *12*, 393–416.

52. Nüsslein-Volhard, C., Lohs-Schardin, M., Sander, K., and Cremer, C. (1980). A
dorso-ventral shift of embryonic primordia in a new maternal-effect mutant of
Drosophila. Nature *283*, 474–476.

53. Simpson, P. (1983). Maternal-zygotic gene interactions during formation of the
dorsoventral pattern in Drosophila embryos. Genetics *105*, 615–632.

54.  Isoda, K., Roth, S., and Nüsslein-Volhard, C. (1992). The functional domains of the
Drosophila morphogen dorsal: evidence from the analysis of mutants. Genes Dev.
*6*, 619–630.

55.  Hegde, S., Sreejan, A., Gadgil, C.J., and Ratnaparkhi, G.S. (2022). SUMOylation of
Dorsal attenuates Toll/NF-κB signaling. Genetics *221*.
https://doi.org/10.1093/genetics/iyac081.

56.  Verstreken, P., Ly, C.V., Venken, K.J.T., Koh, T.-W., Zhou, Y., and Bellen, H.J.
(2005). Synaptic mitochondria are critical for mobilization of reserve pool vesicles
at Drosophila neuromuscular junctions. Neuron *47*, 365–378.

57.  Li, Z., Okamoto, K.-I., Hayashi, Y., and Sheng, M. (2004). The importance of
dendritic mitochondria in the morphogenesis and plasticity of spines and synapses.
Cell *119*, 873–887.

58.  Rikhy, R., Kamat, S., Ramagiri, S., Sriram, V., and Krishnan, K.S. (2007).
Mutations in dynamin-related protein result in gross changes in mitochondrial
morphology and affect synaptic vesicle recycling at the Drosophila neuromuscular
junction. Genes Brain Behav. *6*, 42–53.

59.  Dubal, D., Moghe, P., Verma, R.K., Uttekar, B., and Rikhy, R. (2022). Mitochondrial
fusion regulates proliferation and differentiation in the type II neuroblast lineage in
Drosophila. PLoS Genet. *18*, e1010055.

60.  Martin, A.C., Kaschube, M., and Wieschaus, E.F. (2009). Pulsed contractions of an
actin-myosin network drive apical constriction. Nature *457*, 495–499.

61.  Martin, A.C., Gelbart, M., Fernandez-Gonzalez, R., Kaschube, M., and Wieschaus,
E.F. (2010). Integration of contractile forces during tissue invagination. J. Cell Biol.
*188*, 735–749.

62.  Seo, B.J., Choi, J., La, H., Habib, O., Choi, Y., Hong, K., and Do, J.T. (2020). Role
of mitochondrial fission-related genes in mitochondrial morphology and energy
metabolism in mouse embryonic stem cells. Redox Biol. *36*, 101599.

63.   Seto, E., and Yoshida, M. (2014). Erasers of histone acetylation: the histone
deacetylase enzymes. Cold Spring Harb. Perspect. Biol. *6*, a018713.

64.   Murphy, M.P. (2009). How mitochondria produce reactive oxygen species.
Biochem. J. *417*, 1–13.

65.   Zhao, H., Kalivendi, S., Zhang, H., Joseph, J., Nithipatikom, K., Vásquez-Vivar, J.,
and Kalyanaraman, B. (2003). Superoxide reacts with hydroethidine but forms a
fluorescent product that is distinctly different from ethidium: potential implications in
intracellular fluorescence detection of superoxide. Free Radic. Biol. Med. *34*, 1359–
1368.

66.   Han, D., Antunes, F., Canali, R., Rettori, D., and Cadenas, E. (2003). Voltage-
dependent anion channels control the release of the superoxide anion from
mitochondria to cytosol. J. Biol. Chem. *278*, 5557–5563.

67.   Uttekar, B., Verma, R.K., Tomer, D., and Rikhy, R. (2024). Mitochondrial
morphology dynamics and ROS regulate apical polarity and differentiation in
Drosophila follicle cells. Development *151*. https://doi.org/10.1242/dev.201732.

68.   Muliyil, S., and Narasimha, M. (2014). Mitochondrial ROS regulates cytoskeletal
and mitochondrial remodeling to tune cell and tissue dynamics in a model for
wound healing. Dev. Cell *28*, 239–252.

69.   Sanderson, T.H., Raghunayakula, S., and Kumar, R. (2015). Release of
mitochondrial Opa1 following oxidative stress in HT22 cells. Mol. Cell. Neurosci.
*64*, 116–122.

70.   Hunter, M.V., Willoughby, P.M., Bruce, A.E.E., and Fernandez-Gonzalez, R.
(2018). Oxidative stress orchestrates cell polarity to promote embryonic wound
healing. Dev. Cell *47*, 377–387.e4.

71.   Pease, D.C. (1941). Echinoderm bilateral determination in chemical concentration
gradients. I. The effects of cyanide, ferricyanide, iodoacetate, picrate, dinitrophenol,
urethane, iodine, malonate, etc. J. Exp. Zool. *86*, 381–404.

72. Pilocarpine, P., and Cysteine, G. Echinoderm Bilateral Determination in Chemical
Concentration Gradients. II. The Effects of Azide. The Journal of Experimental
Zoology *89*, 329–345.

73. Pease, D.C. (1942). Echinoderm bilateral determination in chemical concentration
gradients. III. The effects of carbon monoxide and other gases. J. Exp. Zool. *89*,
347–356.

74. Schiffmann, D.A. (1997). Dorsoventral asymmetry in mitochondrial membrane
potential in early Drosophila embryos. Biochem. Soc. Trans. *25*, S653.

75. Landström, U., and Lovtrup, S. (1975). On the determination of the dorso-ventral
polarity in Xenopus laevis embryos. J. Embryol. Exp. Morphol. *33*, 879–895.

76. Løvtrup, S., and Pigon, A. (1958). Inversion of the dorso-ventral axis in amphibian
embryos by unilateral restriction of oxygen supply. Development *6*, 486–490.

77. Cheong, A., Archambault, D., Degani, R., Iverson, E., Tremblay, K.D., and Mager,
J. (2020). Nuclear-encoded mitochondrial ribosomal proteins are required to initiate
gastrulation. Development *147*, dev188714.

78. Pinho, B.R., Santos, M.M., Fonseca-Silva, A., Valentão, P., Andrade, P.B., and
Oliveira, J.M.A. (2013). How mitochondrial dysfunction affects zebrafish
development and cardiovascular function: an in vivo model for testing
mitochondria-targeted drugs. Br. J. Pharmacol. *169*, 1072–1090.

79. Ratnaparkhi, A. (2013). Signaling by Folded gastrulation is modulated by
mitochondrial fusion and fission. J. Cell Sci. *126*, 5369–5376.

80. Dawes-Hoang, R.E., Parmar, K.M., Christiansen, A.E., Phelps, C.B., Brand, A.H.,
and Wieschaus, E.F. (2007). Folded gastrulation, cell shape change and the
control of myosin localization. Development *134*, 4507–4507.

81. Gong, L., Puri, M., Unlü, M., Young, M., Robertson, K., Viswanathan, S.,
Krishnaswamy, A., Dowd, S.R., and Minden, J.S. (2004). Drosophila ventral furrow

1547    morphogenesis: a proteomic analysis. Development *131*, 643–656.

1548 82. Ip, Y.T., Maggert, K., and Levine, M. (1994). Uncoupling gastrulation and
1549    mesoderm differentiation in the Drosophila embryo. EMBO J. *13*, 5826–5834.

1550 83. Ransick, A., and Davidson, E.H. (1993). A complete second gut induced by
1551    transplanted micromeres in the sea urchin embryo. Science *259*, 1134–1138.

1552 84. Venuti, J.M., Gan, L., Kozlowski, M.T., and Klein, W.H. (1993). Developmental
1553    potential of muscle cell progenitors and the myogenic factor SUM-1 in the sea
1554    urchin embryo. Mech. Dev. *41*, 3–14.

1555 85. Ježek, J., Cooper, K., and Strich, R. (2018). Reactive oxygen species and
1556    mitochondrial dynamics: The yin and yang of mitochondrial dysfunction and cancer
1557    progression. Antioxidants (Basel) *7*, 13.

1558 86. Kowaltowski, A.J., Menezes-Filho, S.L., Assali, E.A., Gonçalves, I.G., Cabral-
1559    Costa, J.V., Abreu, P., Miller, N., Nolasco, P., Laurindo, F.R.M., Bruni-Cardoso, A.,
1560    et al. (2019). Mitochondrial morphology regulates organellar Ca2+ uptake and
1561    changes cellular Ca2+ homeostasis. FASEB J. *33*, 13176–13188.

1562 87. Huang, Q., Zhan, L., Cao, H., Li, J., Lyu, Y., Guo, X., Zhang, J., Ji, L., Ren, T., An,
1563    J., et al. (2016). Increased mitochondrial fission promotes autophagy and
1564    hepatocellular carcinoma cell survival through the ROS-modulated coordinated
1565    regulation of the NFKB and TP53 pathways. Autophagy *12*, 999–1014.

1566 88. Yarosh, W., Monserrate, J., Tong, J.J., Tse, S., Le, P.K., Nguyen, K., Brachmann,
1567    C.B., Wallace, D.C., and Huang, T. (2008). The molecular mechanisms of OPA1-
1568    mediated optic atrophy in Drosophila model and prospects for antioxidant
1569    treatment. PLoS Genet. *4*, e6.

1570 89. Tang, S., Le, P.K., Tse, S., Wallace, D.C., and Huang, T. (2009). Heterozygous
1571    mutation of Opa1 in Drosophila shortens lifespan mediated through increased
1572    reactive oxygen species production. PLoS One *4*, e4492.

90.  Rubén, I., Manjarrés-Raza, C., Vicente-Gutiérrez, M., Corrado, J.P., and Bolaños,
L. (2021). Opa1 Relies on Cristae Preservation and ATP Synthase to Curtail
Reactive Oxygen Species Accumulation in Mitochondria. Redox Biology.

91.  Yao, C.-H., Wang, R., Wang, Y., Kung, C.-P., Weber, J.D., and Patti, G.J. (2019).
Mitochondrial fusion supports increased oxidative phosphorylation during cell
proliferation. Elife *8*. https://doi.org/10.7554/elife.41351.

92.  Chen, W., Zhao, H., and Li, Y. (2023). Mitochondrial dynamics in health and
disease: mechanisms and potential targets. Signal Transduct. Target. Ther. *8*, 333.

93.  Ponte, S., Carvalho, L., Gagliardi, M., Campos, I., Oliveira, P.J., and Jacinto, A.
(2020). Drp1-mediated mitochondrial fission regulates calcium and F-actin
dynamics during wound healing. Biol. Open *9*, bio.048629.

94.  Parone, P.A., Da Cruz, S., Tondera, D., Mattenberger, Y., James, D.I., Maechler,
P., Barja, F., and Martinou, J.-C. (2008). Preventing mitochondrial fission impairs
mitochondrial function and leads to loss of mitochondrial DNA. PLoS One *3*, e3257.

95.  Moldogazieva, N.T., Mokhosoev, I.M., Feldman, N.B., and Lutsenko, S.V. (2018).
ROS and RNS signalling: adaptive redox switches through oxidative/nitrosative
protein modifications. Free Radic. Res. *52*, 507–543.

96.  Fedorova, M., Kuleva, N., and Hoffmann, R. (2009). Reversible and irreversible
modifications of skeletal muscle proteins in a rat model of acute oxidative stress.
Biochim. Biophys. Acta *1792*, 1185–1193.

97.  Robison, A.J., and Colbran, R.J. (2013). Calcium | Calcium/Calmodulin-Dependent
Protein Kinases. In Encyclopedia of Biological Chemistry III (Elsevier), pp. 655–
660.

98.  Scholey, J.M., Taylor, K.A., and Kendrick-Jones, J. (1980). Regulation of non-
muscle myosin assembly by calmodulin-dependent light chain kinase. Nature *287*,
233–235.

99. Suzuki, M., Sato, M., Koyama, H., Hara, Y., Hayashi, K., Yasue, N., Imamura, H.,

Fujimori, T., Nagai, T., Campbell, R.E., et al. (2017). Distinct intracellular Ca2+

dynamics regulate apical constriction and differentially contribute to neural tube

closure. Development *144*, 1307–1316.

100. Corcoran, A., and Cotter, T.G. (2013). Redox regulation of protein kinases. FEBS

J. *280*, 1944–1965.

101. Ray, P.D., Huang, B.-W., and Tsuji, Y. (2012). Reactive oxygen species (ROS)

homeostasis and redox regulation in cellular signaling. Cell. Signal. *24*, 981–990.

1607

**A**

Dorsal

Ventral

Sagittal

Surface view

End-on

End-on

Cellularization

Gastrulation

**B**

Hoechst   Streptavidin   Phalloidin

**C**

**D**   mito-GFP

0 min          6.5 min          7.4 min          7.9 min          8.3 min

**E**

Normalized Apical:Basal mitochondrial intensity

1   2   3   4   5

Time Point

Ventral
Neighbouring
Lateral

Dorsal

Ventral

**F**          Zoomed

Lateral

Ventral

**G**

Number of apical mitochondria/cell

****

Lateral   Ventral

Fig 1

**A**

mito-PAGFP / Sqh-mCherry

Preactivation | T1 | T2 | T3 | T4 | T5

Zoomed In

Preactivation | T1 | T2 | T3 | T4 | T5

**B**

Apical:Basal mitoGFP intensity

Apical:Basal Myosin II Intensity

Time point

- - - Myosin II-Ventral
- - - Myosin II-Lateral
—•— Mito-Ventral
—•— Mito-Lateral

Lateral

Ventral

**C**

Control

Colchicine

mito-GFP

**D**

Normalized Apical:Basal mitochondrial intensity

Control | Colchicine

*

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

## C

| Proteins | log2FC |
|----------|--------|
| Sod1 | 0.3313313292 |
| GstD3 | 0.3677227784 |
| GstE1 | 0.5297582967 |
| GstE6 | 0.4620588666 |
| GstE7 | 0.7760282215 |
| Prx6c | 1.23698529 |

## F

| Proteins | log2FC | Proteins | log2FC |
|----------|--------|----------|--------|
| ND-75 | -1.292618773 | UQCR-Q | -0.97470137 |
| ND-13A | -1.484030494 | COX6B | -2.494907874 |
| ND-39 | -1.439414726 | COX7A | -2.249872269 |
| mt:ND1 | -1.478197331 | COX5A | -1.789639136 |
| SdhA | -1.055428856 | mt:CoII | -1.587067991 |
| SdhB | -0.98432467 | ATPsynD | -0.978086075 |
| Sdhaf3 | -3.262787014 | ATPsynB | -0.92933435 |
| UQCR-11 | -1.289831246 | ATPsynG | -1.777576053 |
| UQCR-14 | -1.196573863 | mt:ATPase8 | -1.804977808 |

Fig 7

Fig 8

Fig 9

Control    Drp1 mutant

Drp1 mut
Less ROS

Drp1 mut
+
SOD2 mut
More ROS

⊚ Mitochondria    · ROS    ▬ Myosin II

Fig 10