TITLE

Cytokinin receptor AHK3 modulates root-to-shoot transport of *trans*-zeatin type cytokinin

AUTHORS

Kota Monden[a†], Takamasa Suzuki[b], Mikiko Kojima[c], Yumiko Takebayashi[c], Takehiro Kamiya[d], Takatoshi Kiba[e,f], Hitoshi Sakakibara[c,f], Tsuyoshi Nakagawa[a], and Takushi Hachiya[a]

[a]Department of Molecular and Functional Genomics, Interdisciplinary Center for Science Research, Shimane University, 1060 Nishikawatsu-cho, Matsue, Shimane 690-8504, Japan

[b]College of Bioscience and Biotechnology, Chubu University, 1200 Matsumoto-cho, Kasugai, Aichi 487-8501, Japan

[c]RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro-cho, Tsurumi-ku, Yokohama, Kanagawa 230-0045, Japan

[d]Graduate School of Agricultural and Life Sciences, Tokyo University, 1-1-1 Yayoi, Bunkyo-ku, Tokyo 113-8657, Japan

[e]Institute of Plant Science and Resources, Okayama University, 2-20-1 Chuo, Kurashiki, Okayama 710-0046, Japan

[f]Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8601, Japan

[†]Corresponding author: Kota Monden (n23d203@matsu.shimane-u.ac.jp)

AUTHOR STATEMENT

The author responsible for distribution of materials integral to the findings presented in this article in accordance with the policy described in the Instructions for Authors (https://academic.oup.com/plphys/pages/General-Instructions) is Kota Monden.

RUNNING HEAD

AHK3 modulates long-distance cytokinin transport.

ABSTRACT

Trans-zeatin (*t*Z)-type cytokinins are synthesized in roots in response to nitrogen

availability, transported to shoot via xylem, and coordinate diverse physiological

processes in aerial organs. Within this mechanism, the regulation of cytokinin

biosynthesis by nitrate signaling via NIN-like protein 7 as well as the loading of *t*Z-type

cytokinins into xylem by ATP-binding cassette transporter G14 have been well studied.

However, the roles of other components remain unclear. Here, we show that cytokinin

perception and degradation in roots, as mediated by Arabidopsis histidine kinase 3

(AHK3) and cytokinin oxidase/dehydrogenase 4 (CKX4), modulate root-to-shoot

transport of *t*Z-type cytokinins. Grafting experiments demonstrated that root-specific

AHK3 deficiency systemically increased leaf blade area through long-distance signals of

root-derived *t*Z-type cytokinin, perceived by shoot-expressed AHK3. Transcriptome,

hormonome, and reporter analyses revealed that root-specific *AHK3* deficiency reduced

CKX4 expression in roots, elevating *t*Z-type cytokinin levels in roots and xylem sap and

thereby enhancing leaf cytokinin response. Transfer experiments manipulating root

nitrate levels showed that root-specific *AHK3* deficiency promoted leaf blade area in a

manner dependent on both nitrate and root-derived *t*Z-type cytokinin signaling.

Moreover, both nitrate signals and root-expressed AHK3 are required for maximal

CKX4 induction in roots, and root-specific *CKX4* deficiency enhanced leaf blade area in

a nitrate-dependent manner. These findings reveal a novel mechanism in which an

AHK3–CKX4 module governs root-to-shoot transport of *t*Z-type cytokinins, fine-tuning

leaf size according to nitrogen availability in roots.

INTRODUCTION

Cytokinins (CKs) are essential plant hormones that coordinate diverse developmental

and physiological processes, including cell division, shoot and root growth, vascular

formation, and responses to stress and nutrients (Werner et al., 2001; Riefler et al.,

2006; Matsumoto-Kitano et al., 2008; Sakakibara, 2021). In *Arabidopsis*, the

biologically active CK forms include N^6-(Δ^2-isopentenyl) adenine (iP) and *trans*-zeatin

(*t*Z). The initial and rate-limiting step of CK biosynthesis involves N^6-prenylation of

adenosine 5′-phosphates catalyzed by adenosine phosphate-isopentenyltransferases

(IPTs), producing iP-riboside 5′-phosphates (iPRPs) (Takei et al., 2001a; Miyawaki et

al., 2006). These intermediates are subsequently hydroxylated by cytochrome P450 monooxygenases CYP735A1 and CYP735A2 to form tZ-riboside 5′-phosphates ($tZRPs$) (Takei et al., 2004a). Ribotide precursors are then converted into iP and tZ by CK-activating enzyme members of the lonely guy (LOG) family (Kurakawa et al., 2007; Kuroha et al., 2009). CK homeostasis is further modulated by reversible *O*-glucosylation and tentatively reversible *N*-glucosylation catalyzed by glucosyltransferases (UGTs), and by irreversible degradation by cytokinin oxidase/dehydrogenases (CKXs) (Werner et al., 2006; Wang et al., 2011; Jin et al., 2013; Wang et al., 2013; Hošek et al., 2020). CK perception and signaling rely on a multistep His-to-Asp phosphorelay system (Hutchison and Kieber, 2002; Müller and Sheen, 2007). Three membrane-spanning histidine kinases, Arabidopsis histidine kinase (AHK)2, AHK3, and CRE1/AHK4, act as CK receptors (Inoue et al., 2001). Upon ligand binding, the phosphoryl group is transferred from AHKs to histidine phosphotransfer proteins, which shuttle into the nucleus and activate type-B response regulators (Hutchison and Kieber, 2002; Müller and Sheen, 2007). These transcription factors induce CK-responsive genes, including type-A response regulators that function as negative feedback elements (To et al., 2004).

In addition to local roles, CKs act as long-distance signaling molecules that coordinate developmental and physiological processes between roots and shoots (Sakakibara, 2021). In *Arabidopsis*, tZ-type CKs predominate in xylem sap, whereas iP-type CKs are more abundant in phloem sap (Hirose et al., 2008; Osugi et al., 2017). This distribution reflects tissue-specific expression of CK biosynthetic genes: *CYP735A2* is strongly expressed in root vasculature, while *IPT3* is mainly expressed in phloem tissues of both roots and shoots (Miyawaki et al., 2004; Takei et al., 2004a,b; Kiba et al., 2013). These activities establish a directional flow of CK species, with root-derived tZ-type CKs transported to shoots via xylem and shoot-derived iP-type CKs transported to roots via phloem tissue (Hirose et al., 2008; Matsumoto-Kitano et al., 2008; Kiba et al., 2013). A central component of the root-to-shoot CK transport system is ATP-binding cassette transporter G14 (ABCG14), which loads CKs into xylem in root stele and also contributes to phloem unloading in shoots (Ko et al., 2014; Zhang et al., 2014; Zhao et al., 2021). Heterologous transport assays in yeast have shown that

ABCG14 transports multiple CK species, including both iP- and *t*Z-types, underscoring

its pivotal role in systemic CK distribution and signaling (Zhao et al., 2023).

Root-to-shoot CK transport has been found to be tightly linked to plant

nitrogen status. For example, supplying inorganic nitrogen to root activates CK

signaling in leaves and increases xylem transport of *t*Z-riboside (*t*ZR) following nitrate

application (Sakakibara et al., 1998; Takei et al., 2001b). Moreover, nitrate-induced

changes in leaf growth rate correlate with *t*Z and *t*ZR levels in xylem sap and leaves

(Rahayu et al., 2005). In *Arabidopsis*, nitrate signaling regulates expression of *IPT3* and

CYP735A2 through NIN-like protein 7 (NLP7) and Nitrate-inducible GARP-type

transcriptional repressor 1s (NIGT1s), these transcription factors that are central to the

primary nitrate response (Miyawaki et al., 2004; Takei et al., 2004a,b; Kiba et al., 2018;

Maeda et al., 2018; Liu et al., 2022). It has been proposed that nitrate signaling

mediated by NLP7 coordinates CK metabolism and root-to-shoot CK transport to

modulate shoot CK responses (Abualia et al., 2022). In addition, long-distance signals

of root-derived *t*Z and *t*ZR differentially affect leaf growth traits, and their ratios in

xylem sap are known to be modulated by nitrogen supply (Osugi et al., 2017). Together,

these findings highlight that regulation of root-to-shoot CK transport is critical for

fine-tuning leaf growth in response to nitrogen availability in roots.

Our previous study proposed that root-expressed CK receptors are involved

in the regulation of root-to-shoot CK transport. Using grafting, we showed that

root-specific deficiencies of *AHK2* and *AHK3* increased iP- and *t*Z-type CK levels in

roots by suppressing CK degradation and inducing CK biosynthesis. This rise in root

CK content was accompanied by higher *t*Z-type CK levels and stronger expression of

CK-inducible genes in shoots (Monden et al., 2022). Notably, among the three CK

receptors, *AHK3* was identified as one of the top genes co-expressed with *NLP7*

(Supplementary Table S1), suggesting that AHK3 may act in association with nitrate

responses via NLP7. In addition, AHK3 has been reported to exhibit a higher affinity

for *t*Z than for iP (Stolz et al., 2011), implying a preferential role in *t*Z-type CK

signaling. Together, these findings suggest that root-expressed AHK3 may modulate

root-to-shoot transport of *t*Z-type CKs in response to nitrogen availability in roots,

although the underlying mechanism and physiological relevance remain unclear.

In this study, we show that CK perception and degradation in roots, mediated by AHK3 and CKX4, modulate root-to-shoot transport of *t*Z-type CKs. Growth analyses of grafted plants demonstrated that root-specific *AHK3* deficiency enlarged leaf blade area through long-distance signals of root-derived *t*Z-type CK, perceived by shoot-expressed AHK3. Transcriptome, hormonome, and reporter assays further revealed that root-specific *AHK3* deficiency reduced *CKX4* expression in roots, elevating *t*Z-type CK levels in roots and xylem sap and thereby enhancing CK responses in leaves. Transfer experiments manipulating root nitrate levels showed that nitrate-induced leaf growth promotion depends on long-distance signals of root-derived *t*Z-type CK, which is in turn controlled by root-expressed AHK3 and CKX4. We propose a mechanism in which the AHK3–CKX4 module governs root-to-shoot transport of *t*Z-type CK in response to nitrogen availability in roots, thereby fine-tuning leaf size to maintain balanced plant growth.

RESULTS

1. Root-expressed AHK3 modulates shoot and root growth

To identify which root-expressed CK receptor(s) influence shoot and root growth, we grafted Col-0 (WT) scions onto rootstocks derived from WT, single mutant (i.e., *ahk2-5*; *ahk2*, *ahk3-7*; *ahk3*, *cre1-2*; *ahk4*) or double mutant (i.e., *ahk2-5 ahk3-7*; *ahk23*, *ahk2-5 cre1-2*; *ahk24*, *ahk3-7 cre1-2*; *ahk34*) five□day□old seedlings (Riefler et al., 2006). Five□days after grafting, we selected plants (scion/rootstock: WT/WT, WT/*ahk2*, WT/*ahk3*, WT/*ahk4*, WT/*ahk23*, WT/*ahk24*, WT/*ahk34*) for further analysis if they showed vigorous growth without adventitious roots. Grafted plants were then cultivated for 7 days on either half-strength MS (1/2 MS) medium or nutrient-rich soil. We measured shoot fresh weight, leaf blade area, and leaf number. Overall, WT/*ahk3*, WT/*ahk23*, and WT/*ahk34* displayed greater shoot growth than WT/WT (Fig. 1A, Supplementary Fig. S1A). Relative to the WT/WT control, these lines showed an increase in shoot fresh weight of 26%, 35%, and 19%, and in leaf blade area of 26%, 32%, and 17%, respectively (Fig. 1, B and C). Under nutrient-rich soil conditions, shoot fresh weight increased by 34%, 25%, and 22%, and leaf blade area by 32%, 21%, and 17% in WT/*ahk3*, WT/*ahk23*, and WT/*ahk34*, respectively, compared with WT/WT (Supplementary Fig. S1, B and C). Leaf number was generally similar across grafted

plants in both conditions, although several statistically significant differences were noted (Fig. 1D, Supplementary Fig. S1D). We next monitored primary root length in grafted plants transferred to 1/2 MS medium over a five-day period. Primary root length was comparable at day 0, but WT/*ahk3*, WT/*ahk23*, and WT/*ahk34* developed significantly shorter roots than WT/WT from day 3 onward (Supplementary Fig. S2). In contrast, WT/*ahk2*, WT/*ahk4*, and WT/*ahk24* showed no significant differences in shoot or root growth relative to WT/WT (Fig. 1, Supplementary Figs. S1 and S2). Notably, the relative growth of WT/*ahk3* was comparable to WT/*ahk23* and WT/*ahk34*, indicating that loss of AHK3 alone is sufficient to alter overall plant growth. We therefore focused subsequent analyses on WT/*ahk3*, using WT/WT, WT/*ahk2*, and WT/*ahk4* as comparisons.

2. Root-expressed AHK3 modulates root CK levels

Our previous work showed that root specific deficiencies of *AHK2* and *AHK3* markedly increased iP and *t*Z type CK levels in roots, but the receptor responsible for governing root CK levels was unclear (Monden et al., 2022). To clarify this, we measured iP- and *t*Z-type CK concentrations in WT/WT, WT/*ahk2*, WT/*ahk3*, and WT/*ahk4* roots. Root concentrations of both iP- and *t*Z-type CKs were significantly elevated in WT/*ahk3* compared with WT/WT; moreover, this was true for active, precursor, and inactive forms (Fig. 2, Supplementary Fig. S3). In contrast, root iP and *t*Z type CKs were largely unchanged in WT/*ahk2* and WT/*ahk4* relative to WT/WT, except for iP7G and iP9G in WT/*ahk4* (Fig. 2, Supplementary Fig. S3). These findings indicate that AHK3 expressed in roots modulates CK homeostasis.

3. Root-expressed AHK3 influences the root transcriptome

Despite the fact that *AHK2*, *AHK3*, and *AHK4* are all expressed in roots (Higuchi et al., 2004), only AHK3 strongly affected root CK levels. This suggests that each CK receptor may regulate a distinct set of downstream genes. To test this hypothesis, we performed an RNA-seq analysis of root samples taken from WT/WT, WT/*ahk2*, WT/*ahk3*, and WT/*ahk4*. All data, including raw read counts, reads per million mapped reads (RPM), and normalized transcript levels are presented in Supplementary Tables S2–4. First, we examined differentially expressed genes (DEGs) between the roots of

the WT/WT control and three grafted plants, i.e., WT/*ahk2*, WT/*ahk3*, and WT/*ahk4*

(Supplementary Table S6). An analysis of the DEGs with a fold change of 1.5 or greater

revealed that 62 genes were significantly upregulated in WT/*ahk3* relative to WT/WT

roots, whereas 91 genes were downregulated (Fig. 3A, Supplementary Figs. S4 and S5).

Notably, relatively few DEGs were detectable in the roots of WT/*ahk2* and WT/*ahk4*

(Fig. 3A, Supplementary Figs. S4 and S5). Next, we conducted a Gene Ontology (GO)

term enrichment analysis of the 62 upregulated genes and the 87 downregulated genes

(excluding *AHK3*) identified in WT/*ahk3* roots. This revealed that multiple

hormone-related terms were enriched in both sets (Supplementary Fig. S6). To further

examine the extent to which CK receptor regulates CK signaling-related genes in roots,

we analyzed the expression of CK-inducible and -repressible genes in the roots of

grafted plants; to do so we used a curated gene list (Bhargava et al., 2013). We found

that the expression levels of both CK-inducible and -repressible genes in the roots of

WT/*ahk4* were comparable to the levels detected in WT/WT (Fig. 3B). In contrast, the

expression of CK-inducible genes in root tissue samples was significantly lower in both

WT/*ahk2* and WT/*ahk3* relative to the WT/WT (Fig. 3B). Moreover, CK-repressible

genes were significantly increased in WT/*ahk3* and showed an increasing trend in

WT/*ahk2* (Fig. 3B). Notably, the magnitude of these changes was more pronounced in

WT/*ahk3* than in WT/*ahk2* roots, suggesting that root-expressed AHK3 plays a major

role in regulating root CK signaling.

Next, to identify candidate gene(s) that may be responsible for increased

CK levels in WT/*ahk3* roots, we examined whether the DEGs observed in WT/*ahk3*

roots included genes that are known to be involved in CK biosynthesis and inactivation.

Of these genes, only *CKX4*, which is involved in CK degradation, was among the genes

that were downregulated genes in WT/*ahk3* roots relative to the WT/WT (Fig. 3, A and

C). Quantitative RT-PCR (qRT-PCR) analysis confirmed that expression of *CKX4* in

roots was significantly decreased in WT/*ahk3* relative to WT/WT (Fig. 3D). Taken

together, these results suggest that root-expressed AHK3 modulates root CK levels,

potentially via CKX4-mediated CK degradation.

4. Root-expressed AHK3 modulates *t*Z-type CKs in xylem sap

CKs synthesized in roots are transported to shoots mainly via xylem pathway (Ko et al.,

2014; Osugi et al., 2017). Given that we observed relatively high CK levels in WT/*ahk3*

roots, we hypothesized that the CK levels of xylem sap may also be altered. To verify

this, we measured the CK concentrations of xylem sap samples from WT/WT, WT/*ahk2*,

WT/*ahk3*, and WT/*ahk4*. We found that *t*Z, *t*ZR, and *t*ZRPs were relatively more

abundant than inactive forms of *t*Z-type CK and iP-type CKs in WT/WT xylem sap

(Supplementary Fig. S7A). This suggests that there is preferential root-to-shoot

transport of *t*Z and its precursors. Notably, the concentrations of *t*Z-type CKs were

significantly higher in WT/*ahk3* than in WT/WT xylem sap, while the concentrations of

iP-type CKs remained largely unchanged (Fig. 4, Supplementary Fig. S7B). Among the

major CK species detected in xylem sap, the concentration of *t*Z showed a 2.22-fold

increase in WT/*ahk3* relative to WT/WT xylem sap (Fig. 4B). In contrast, the

concentrations of *t*ZR and *t*ZRPs were both slightly higher in WT/*ahk3* than in WT/WT

xylem sap, showing 1.20-fold and 1.37-fold increases, respectively (Fig. 4B). Finally,

the concentrations of iP☐ and *t*Z☐type CKs were largely unchanged in WT/*ahk2* and

WT/*ahk4* relative to WT/WT xylem sap, except for *t*Z and *t*ZRPs in WT/*ahk4* (Fig. 4,

Supplementary Fig. S7B). Overall, these results suggest that root-expressed AHK3

modulates the root-to-shoot transport of *t*Z-type CKs, especially *t*Z.

5. Root-expressed AHK3 systemically influences the shoot transcriptome

Next, to test whether elevated *t*Z-type CK levels in WT/*ahk3* xylem sap affected shoot

CK levels, we measured CK concentrations in WT/WT, WT/*ahk2*, WT/*ahk3*, and

WT/*ahk4* shoots. Unexpectedly, *t*Z-type CK concentrations were not higher in WT/*ahk3*

than in WT/WT shoots, except for *t*Z9G (Supplementary Fig. S8). Similarly, almost all

CK species in WT/*ahk2* and WT/*ahk4* showed no marked differences compared with

WT/WT (Supplementary Fig. S8). We next analyzed whether root-specific CK receptor

deficiency may alter the shoot transcriptomes. To do so, we performed RNA-seq on

WT/WT, WT/*ahk2*, WT/*ahk3*, and WT/*ahk4* shoots. Raw read counts, RPM, and

normalized transcript levels are provided in Supplementary Tables S2, S3, and S5.

Differential expression analysis with a 1.5-fold cutoff revealed 163 genes significantly

upregulated and 35 genes downregulated in WT/*ahk3* relative to WT/WT shoots (Fig.

5A, Supplementary Table S7). No DEGs were detected in WT/*ahk2* or WT/*ahk4* shoots

(Fig. 5A, Supplementary Table S7). Moreover, the number of upregulated genes was about five times greater than the number of downregulated genes in WT/*ahk3* shoots (Fig. 5A, Supplementary Table S7). This finding indicates that root-specific *AHK3* deficiency mainly induces gene expression in shoots.

To gain insight into the functional characteristics of the upregulated set, we performed GO enrichment analysis using the set of 163 upregulated genes. Enriched categories included "Translation," "Response to inorganic substance," and "Cellular nitrogen compound biosynthetic process" (Fig. 5B), indicating activation of genes related to protein synthesis, nutrient responses, and nitrogen metabolism in WT/*ahk3* shoots. We then examined whether these transcriptional changes correlated with altered shoot element concentrations. We found that the shoot concentrations of calcium, sulfur, manganese, and molybdenum were significantly higher in WT/*ahk3* than in WT/WT (Supplementary Fig. S9), suggesting a possible association between these nutrients and altered gene expression in WT/*ahk3* shoots. In contrast, although several nitrogen-related GO terms were enriched, shoot concentrations of nitrogen and nitrate did not differ between WT/WT and WT/*ahk3* (Supplementary Fig. S9). Thus, nitrogen status in WT/*ahk3* shoots did not drive the transcriptional changes observed in protein synthesis and nitrogen metabolism genes. Because nitrogen is a major protein component and both nitrate and CK signaling promote expressions of protein synthesis-related genes (Sakakibara et al., 2006), altered CK response in WT/*ahk3* shoots may underlie these transcriptional effects.

To further assess CK response, we compared the 163 upregulated genes in WT/*ahk3* shoots with CK-inducible genes (Bhargava et al., 2013). *Arabidopsis response regulator 5* (*ARR5*) and *Glutaredoxin-S6* (*GrxS6*) were among the overlapping genes (Fig. 5A). *ARR5*, one of ten type-A *ARRs*, is a representative CK-responsive gene functioning as a negative feedback regulator of CK signaling (To et al., 2004). Expression of several other type-A *ARRs* also tended to increase in WT/*ahk3* relative to WT/WT (Fig. 5C). Consistent with the RNA-seq results, qRT-PCR confirmed that *ARR5* expression was significantly higher in WT/*ahk3* shoots than in WT/WT (Fig. 5D). Taken together, these results indicate that root-expressed AHK3 modulates shoot gene expression, at least partly through CK signaling.

6. Root-derived CK signals are required for enhanced shoot growth via root-expressed AHK3

Elevated *t*Z-type CKs in WT/*ahk3* xylem sap suggested that long-distance signals of root-derived *t*Z-type CK contribute to enhanced shoot growth. To test this, we generated *ahk3 abcg14* and *ahk3 cyp735a1 cyp735a2* (*ahk3 cypDM*) mutants by traditional crossing (Supplementary Fig. S10A). Since both *abcg14* and *cypDM* shoots display lack proper root-derived *t*Z-type CK signaling (Kiba et al., 2013; Ko et al., 2014), these mutants allow assessment of its contribution. Disruption of the respective genes in *ahk3 abcg14* and *ahk3 cypDM* was confirmed by RT-PCR (Supplementary Fig. S10B). Grafted plants (scion/rootstock: WT/WT, WT/*ahk3*, WT/*abcg14*, WT/*ahk3 abcg14*, WT/*cypDM*, WT/*ahk3 cypDM*) were cultivated on 1/2 MS medium, and shoot growth was measured. Shoot fresh weight and leaf blade area were significantly reduced in WT/*abcg14* and WT/*ahk3 abcg14*, while WT/*cypDM* and WT/*ahk3 cypDM* showed no decrease compared with WT/WT (Fig. 6, A and B). These results agree with previous findings that root-specific deficiency of *ABCG14* impairs shoot growth, whereas deficiencies of *CYP735A1* and *CYP735A2* do not (Kiba et al., 2013; Ko et al., 2014). Notably, the increases in shoot fresh weight and leaf blade area seen in WT/*ahk3* compared with WT/WT were largely suppressed by disrupting *ABCG14* and completely abolished by disrupting *CYP735A1* and *CYP735A2* in roots (Fig. 6, A and B). These findings indicate that long-distance signals of root-derived *t*Z-type CKs are required for enhanced shoot growth in WT/*ahk3*.

To determine which CK receptor(s) mediate CK signaling underlying this growth effect, we analyzed grafted plants carrying single or double mutations in *AHK2*, *AHK3*, or *AHK4* specifically in WT/*ahk3* shoots. Grafted plants (scion/rootstock: WT/WT, WT/*ahk3*, *ahk2*/*ahk3*, *ahk3*/*ahk3*, *ahk4*/*ahk3*, *ahk23*/*ahk3*, *ahk24*/*ahk3*, *ahk34*/*ahk3*) were cultivated under 1/2 MS medium, and shoot growth was measured. Increases in shoot fresh weight and leaf blade area observed in WT/*ahk3* relative to WT/WT were similarly evident in *ahk2*/*ahk3*, *ahk4*/*ahk3*, and *ahk24*/*ahk3* (Fig. 6, C and D). In contrast, these enhancements were suppressed in *ahk3*/*ahk3* and *ahk34*/*ahk3*, and markedly reduced in *ahk23*/*ahk3*, where values were significantly lower than WT/WT (Fig. 6, C and D). These results indicate that enhanced shoot growth in WT/*ahk3* depends primarily on CK signaling through shoot-expressed AHK3, with partial

contribution from AHK2. Next, to visualize sites of enhanced CK response in WT/*ahk3*

shoots, we introduced the synthetic CK reporter *TCSn:GFP* into WT/WT and WT/*ahk3*

shoots (Zürcher et al., 2013). Grafted plants (scion/rootstock: *TCSn:GFP*/WT,

TCSn:GFP/*ahk3*) were grown on 1/2 MS medium. *TCSn:GFP*/*ahk3* exhibited greater

shoot fresh weight and leaf blade area than *TCSn:GFP*/WT (Fig. 6E). Notably, GFP

fluorescence intensity in true leaf was also significantly higher in *TCSn:GFP*/*ahk3* than

in *TCSn:GFP*/WT (Fig. 6F). Finally, to examine how root-specific *AHK3* deficiency

influences leaf micro-morphology, we measured epidermal cell size and number in true

leaves of WT/WT and WT/*ahk3*. Cell size was comparable between genotypes, but cell

number was significantly higher in WT/*ahk3* than in WT/WT (Fig. 6G). These results

suggest that root-specific *AHK3* deficiency enhances leaf CK response, promoting

epidermal cell division and thereby increasing leaf size.

7. Root-expressed AHK3 modulates leaf size in a nitrate-dependent manner

AHK3 was identified as one of the top genes co-expressed with the nitrate sensor NLP7

(Supplementary Table S1), suggesting that AHK3 may play a role in plant nitrate

responses. To test this, we manipulated the nitrate levels of WT/WT and WT/*ahk3* roots.

Ten-day-old WT/WT and WT/*ahk3* were first transferred to 2.5 mM ammonium

succinate (5 mM NH_4^+) medium and incubated for 48 hours to deplete nitrate

(Supplementary Fig. S11A). Grafted plants were then transferred to medium containing

5 mM NH_4^+ supplemented with either 0.5 mM KNO_3 (KNO_3) or 0.5 mM KCl (KCl)

and grown for 5 days to assess the presence or absence of nitrate signaling in roots.

Nitrate accumulated only in WT/WT and WT/*ahk3* roots grown under KNO_3

(Supplementary Fig. S11B). Moreover, WT/WT and WT/*ahk3* appeared healthy, with

comparable shoot chlorophyll concentrations under both conditions (Supplementary Fig.

S12), indicating that nitrogen deficiency or ammonium toxicity did not occur. In

WT/WT, shoot fresh weight and leaf blade area were found to be significantly greater

under KNO_3 than KCl (Fig. 7A), showing that nitrate signaling promotes leaf growth.

Notably, shoot fresh weight and leaf blade area were similar between WT/WT and

WT/*ahk3* under KCl, but were significantly higher in WT/*ahk3* than in WT/WT under

KNO_3 (Fig. 7A).

Next, to test whether nitrate-dependent growth promotion in WT/*ahk3*
requires root-derived *t*Z-type CK signals, we analyzed WT/*abcg14* and WT/*ahk3*
abcg14 grown under KNO_3 or KCl. Interestingly, the increases in shoot fresh weight
and leaf blade area observed in WT/WT and WT/*ahk3* under KNO_3 were strongly
suppressed in WT/*abcg14* and WT/*ahk3 abcg14* (Fig. 7B). These results indicate that
long-distance signals of root-derived *t*Z-type CKs are required for nitrate-induced leaf
growth promotion. Moreover, shoot fresh weight and leaf blade area were comparable
between WT/*abcg14* and WT/*ahk3 abcg14* under KNO_3 (Fig. 7B), suggesting that
root-derived *t*Z-type CK signals may contribute to the enhanced leaf growth of
WT/*ahk3*.

To further clarify the role of AHK3 in nitrate signaling, we analyzed
expression of CK biosynthesis and inactivation genes in roots 4 hours after transfer to
KNO_3 or KCl. Nitrate accumulation was detected only in WT/WT and WT/*ahk3* roots
after KNO_3 treatment (Supplementary Fig. S11C). In WT/WT, root expression of *IPT3*,
CYP735A1, *CYP735A2*, and *CKX4* was significantly higher under KNO_3 than KCl (Fig.
7C, Supplementary Fig. S13), indicating induction by nitrate signaling. Importantly,
expression of *IPT3*, *CYP735A1*, and *CYP735A2* was similar in WT/WT and WT/*ahk3*
under KNO_3, whereas *CKX4* expression was significantly reduced in WT/*ahk3*
compared with WT/WT (Fig. 7C, Supplementary Fig. S13). These results suggest that
both nitrate signals and root-expressed AHK3 are required for maximal *CKX4*
induction.

To investigate the role of CKX4 in nitrate signaling, we obtained two
independent T-DNA alleles, *ckx4-1* (SALK_055204) and *ckx4-2* (SALK_132516), from
the Arabidopsis Biological Resource Center. *Ckx4-1* carries an intron insertion (Bartrina
et al., 2011), while *ckx4-2* carries an exon insertion (Supplementary Fig. S14). RT-PCR
confirmed reduced *CKX4* transcript in *ckx4-1* and absence in *ckx4-2* (Fig. 7D),
indicating that *ckx4-1* is a knockdown and *ckx4-2* a knockout. Grafted plants
(scion/rootstock: WT/WT, WT/*ckx4-1*, WT/*ckx4-2*) were cultivated under KNO_3 or KCl,
and shoot growth was measured. Under KCl, shoot fresh weight and leaf blade area
were similar among genotypes (Fig. 7E). Under KNO_3, however, both traits were
significantly greater in WT/*ckx4-1* and WT/*ckx4-2* compared to WT/WT (Fig. 7E).

To further clarify relationship between AHK3 and CKX4, we generated

ahk3 ckx4-1 and *ahk3 ckx4-2* mutants by traditional crossing (Supplementary Fig. S15,

A and B). Grafted plants (scion/rootstock: WT/WT, WT/*ahk3*, WT/*ckx4-1*, WT/*ckx4-2*,

WT/*ahk3 ckx4-1*, WT/*ahk3 ckx4-2*) were grown under KNO_3 or KCl, and shoot growth

was measured. All genotypes showed comparable shoot growth under KCl

(Supplementary Fig. S15C). Under KNO_3, however, shoot fresh weight and leaf blade

area were significantly greater in WT/*ahk3*, WT/*ckx4-1*, WT/*ckx4-2*, WT/*ahk3 ckx4-1*,

and WT/*ahk3 ckx4-2* compared with WT/WT (Supplementary Fig. S15C). These traits

were higher in WT/*ahk3* than in WT/*ckx4-1* and WT/*ckx4-2* (Supplementary Fig. S15C),

indicating that CKX4 acts downstream of AHK3. Importantly, WT/*ahk3 ckx4-1* and

WT/*ahk3 ckx4-2* showed no further growth increase relative to WT/*ahk3* under KNO_3

(Supplementary Fig. S15C). Together, these results suggest that nitrate-induced leaf

growth promotion occurs through a pathway mediated jointly by AHK3 and CKX4.

DISCUSSION

Recent studies have shown that long-distance signals of root-derived *t*Z-type CK

coordinate multiple physiological processes in aerial organs, including leaf growth,

photoperiod stress response, shade avoidance, and flowering time (Osugi et al., 2017;

Landrein et al., 2018; Frank et al., 2020; Gautrat et al., 2024; Bartrina et al., 2025).

These findings highlight that proper regulation of root-to-shoot *t*Z-type CK transport is

essential for plants to adapt to environmental fluctuations.

Based on our findings and prior studies (Miyawaki et al., 2004; Kiba et al.,

2013; Ko et al., 2014; Zhang et al., 2014; Sakakibara, 2021, 2025), we provide new

insight into the specific mechanisms responsible for modulating root-to-shoot *t*Z-type

CK transport, as mediated by CK perception and degradation through an AHK3–CKX4

module. Nitrate signaling promotes *t*Z-type CK biosynthesis in pericycle and phloem

companion cells via IPT3, CYP735A2, and related enzymes. *t*Z-type CKs are then

transported to the extracellular and endoplasmic reticulum (ER) space by transporters

such as ABCG14. Active form *t*Z are perceived by AHK3 on plasma membrane (PM)

and ER membrane. Activation of CK signaling through AHK3 induces expression of

CKX4, predicted to localize in apoplast, where it modulates apoplastic *t*Z-type CK

levels by degradation. These mechanisms likely influence the apoplastic *t*Z-type CK

pool and, consequently, the amount of *t*Z-type CK loaded into xylem (Supplementary Fig. S16). To further substantiate this model, it will likely be necessary to analyze the spatial expression patterns of AHK3 and CKX4 at the tissue and cellular level in future studies. Interestingly, *IPT3* expression and *t*Z-type CK levels in roots also respond to other nutrients, including phosphate, sulfate, and potassium (Hirose et al., 2008; Nam et al., 2012). Thus, AHK3 may modulates *t*Z-type CK levels in response to diverse nutrient signals. Collectively, the AHK3–CKX4 module could serve as a feedback regulator that attenuates the nutrient-induced root-to-shoot transport of *t*Z-type CKs, thereby fine-tuning leaf CK response to maintain balanced plant growth.

CKXs are known to mediate irreversible CK degradation, producing adenine (or its derivatives) and aldehyde (Schmülling et al., 2003). In *Arabidopsis*, seven CKXs differ in substrate specificity, spatial and temporal expression, and subcellular localization, targeting either vacuoles (CKX1, CKX3), ER (CKX1), apoplasts (CKX2, CKX4, CKX5, CKX6), or the cytosol (CKX7) (Werner et al., 2003; Galuszka et al., 2007; Frébort et al., 2011; Niemann et al., 2018). Expression of *CKX3*, *CKX4*, *CKX5*, and *CKX6* is strongly induced by CK treatment (Werner et al., 2006), suggesting sensitivity to internal CK levels and roles in maintaining homeostasis. Moreover, CKX2 and CKX4 show higher *in vitro* CKX activity than other CKXs, preferentially degrading active form CKs at neutral or slightly basic pH (Galuszka et al., 2007). Our data showed that root-specific *AHK3* deficiency greatly reduced *CKX4* expression in roots, without significantly altering other *CKX*s (Figs. 3 and 7). These findings suggest that root-expressed AHK3 modulates *t*Z degradation by inducing *CKX4* expression in response to the internal *t*Z level. This interpretation is supported by our observation that *t*Z concentration was markedly higher in WT/*ahk3* xylem sap than in WT/WT (Fig. 4). Furthermore, Osugi *et al.* (2017) demonstrated that root-derived *t*ZR controls both leaf number and area, whereas root-derived *t*Z regulates only leaf area. In WT/*ahk3*, the increase in xylem sap *t*Z relative to WT/WT was greater than that of *t*ZR (Fig. 4). Thus, the enlarged leaf blade area in WT/*ahk3*, without a change in leaf number, likely reflects long-distance signaling by root-derived *t*Z (Figs. 1 and 4).

AHK2, AHK3, and AHK4 localize to the PM and ER membrane, where they play central roles in CK signal transduction (Antoniadi et al., 2020; Kubiasová et al., 2020). AHK3 is considered to be primarily localized on the ER membrane and

partially on the PM (Caesar et al., 2011; Wulfetange et al., 2011). Therefore,

ER-localized AHK3 may contribute substantially to regulation of root-to-shoot

transport of *t*Z-type CKs. AHK3 is also expressed more strongly in leaves than in roots,

and its leaf expression exceeds that of AHK2 and AHK4, supporting its hypothesized

role as a major regulator of leaf CK response (Higuchi et al., 2004). This hypothesis is

consistent with our finding that leaf growth promotion, driven by increased shoot CK

response, was diminished when AHK3 was specifically disrupted in WT/*ahk3* shoots

(Fig. 6). Strikingly, root-expressed AHK3 exerted a stronger impact on the root

transcriptome and on CK levels than root-expressed AHK2 and AHK4 (Figs. 2 and 3).

Taken together, these results suggest that AHK3 also plays a major role in roots.

Previous reports indicate that rosette leaf growth, CK metabolism, leaf senescence, and

stomatal opening depend particularly on AHK3, as disruption of AHK3, but not other

CK receptors, impairs these processes (Riefler et al., 2006; Kim et al., 2006; Arnaud et

al., 2017). The distinct functions of these receptors may arise from their expression

patterns, recognition of different CK species, and specific interactions with other

proteins (Higuchi et al., 2004; Spíchal et al., 2004; Romanov et al., 2006; Stolz et al.,

2011). A key feature of AHK3 is its ligand preference: while AHK2 and AHK4 bind iP

and *t*Z with similar affinity, AHK3 shows higher affinity for *t*Z than iP (Stolz et al.,

2011). In addition, AHK3 specifically mediates CK-triggered phosphorylation of ARR2

(Kim et al., 2006). These differences in ligand preference and downstream targets may

underlie the unique functions of AHK3.

Overall, this study uncovered a comprehensive mechanism by which the

CK receptor AHK3 systemically modulates the shoot growth via exerting control of

root-to-shoot *t*Z-type CK transport. Furthermore, we clarified the role of AHK3 in

modulating leaf size in response to root nitrogen availability, a key environmental factor.

Overall, these findings provide new insight into how plants integrate nutritional cues to

coordinate systemic communication between roots and shoots.

MATERIALS AND METHODS

Plant materials

The *Arabidopsis thaliana* accession Col-0 was used as wild type. T-DNA insertion lines

ahk2-5, *ahk3-7*, *cre1-2*, *ahk2-5 ahk3-7*, *ahk2-5 cre1-2*, *ahk3-7 cre1-2* (Riefler et al.,

2006), *abcg14* (Ko et al., 2014), *cyp735a1-2 cyp735a2-2* (Kiba et al., 2013), and *ckx4-1*

(Bartrina et al., 2011) have been described previously. The *ckx4-2* (SALK_132516) line

was obtained from the Arabidopsis Biological Resource Center (https://abrc.osu.edu/).

The transgenic line *TCSn:GFP* (Zürcher et al., 2013) was described previously. Finally,

the homozygous multiple mutants *ahk3-7 abcg14*, *ahk3-7 cyp735a1-2 cyp735a2-2*,

ahk3-7 ckx4-1, and *ahk3-7 ckx4-2* were generated by crossing, and genotypes were

determined by genomic PCR using primers listed in Supplementary Table S8.

Growth conditions of grafted plants

Seeds were surface-sterilized and kept in the dark at 4°C for 3 days to break dormancy.

Approximately 120 seeds per dish were sown in rectangular dishes ($140\square \times 100 \times 20$

mm^3; Eiken Chemical Co. Ltd., Tokyo, Japan) containing 50 mL of half-strength MS

salts including vitamins (Code: M0222; Duchefa Biochemie), supplemented with 0.05%

(w/v) MES-KOH (pH 5.7), 1% (w/v) sucrose, and 0.5% (w/v) gellan gum (Fujifilm

Wako, VA, USA). Plants were grown under a 16/8-h light/dark cycle at 22°C with a

photosynthetic photon flux density (PPFD) of 25 μmol m^{-2} s^{-1}. Five-day-old seedlings

were cut perpendicularly at the hypocotyl using an injection needle tip (Code:

NN-2613S; TERUMO, Tokyo, Japan). Excised scions were connected to corresponding

rootstocks using silicone microtubes (inner $\times$ outer diameter: φ0.4 $\times$ φ0.5 mm; Code:

1-8194-04; AS ONE, Osaka, Japan). Grafted plants were incubated for 5 days under

continuous light at 27°C with a PPFD of 50 μmol m^{-2} s^{-1}. Successfully grafted plants

without adventitious roots were selected, transferred to solid media or soil pots, then

grown for further experiments.

For *in vitro* cultures, grafted plants were grown vertically for five days or

horizontally for seven days under a PPFD of 70 μmol m^{-2} s^{-1} (16/8-h light/dark cycle) at

22°C in 50 mL (vertical) or 30 mL (horizontal) of half-strength MS salts without

vitamins (Code: M0221; Duchefa Biochemie), supplemented with 0.05% (w/v)

MES-KOH (pH 5.7), 1% (w/v) sucrose, and 0.8% (w/v) purified agar (Code: 01162-15;

Nacalai Tesque, Kyoto, Japan). To facilitate non-destructive root sampling during

horizontal growth, media surfaces were covered with cellophane sheets (Hachiya et al.,
2021).

For soil cultivation, grafted plants were grown for 7 days under a PPFD of
70 μmol m^{-2} s^{-1} (16/8-h light/dark cycle) at 22°C in pots containing ~150 mL of
nutrient-rich soil (Supermix A; Sakata Seed, Kyoto, Japan).

For transfer experiments with varying nitrogen concentrations, grafted
plants were placed on cellophane sheets in round dishes (diameter 90 mm; depth
20 mm; AS ONE) containing 30 mL of half-strength MS modified basal salt mixture
without nitrogen (M531; PhytoTech Labs, Lenexa, KS, USA), supplemented with 2.5
mM $(NH_4)_2C_4H_4O_4$, 0.05% (w/v) MES-KOH (pH 5.7), 1% (w/v) sucrose, and 0.8%
(w/v) purified agar (Code: 01162-15; Nacalai Tesque) (5 mM NH_4^+ medium). Plants
were grown for 48 hours under a PPFD of 70 μmol m^{-2} s^{-1} (16/8-h light/dark cycle) at
22°C, then transferred to 5 mM NH_4^+ medium supplemented with either 0.5 mM
KNO$_3$ or 0.5 mM KCl and grown for five days under the same conditions.

520 Growth analysis of grafted plants
Shoots were weighed on a precision balance (HR-202i; A&D, Tokyo, Japan). Leaf
blades were harvested and scanned at 600 dpi (GT-X980; EPSON, Tokyo, Japan) to
measure true leaf blade area. Roots of vertically grown plants were scanned at 600 dpi
to measure primary root length. To monitor root elongation, root tip positions were
marked every 24 hours. Leaf blade area and primary root length were quantified using
ImageJ version 1.53t (US National Institutes of Health, Bethesda, MD, USA).

528 Xylem sap collection
Grafted plants were cut perpendicularly at the hypocotyl using an injection needle tip
(Code: NN-2613S; TERUMO). A silicone microtube (inner × outer diameter: φ0.5 ×
φ0.6 mm; Code: 1-8194-05; AS ONE) was attached to the cut end, and plants were
incubated for 2 hours to allow exudation. After incubation, the silicone microtube was
detached, and xylem sap collected in the tube was recovered by centrifugation into a
collection tube. This was then and stored at –80°C until further analysis. We used
12 μL aliquots of sap from each grafted plant for CK analyses.

Determination of CK species

CK extraction and quantification followed the method of Kojima *et al.* (2009), using an ultra-performance liquid chromatography–tandem quadrupole mass spectrometry system (AQUITY UPLC System/XEVO-TQS; Waters Corp., Milford, MA, USA) equipped with an octadecylsilyl column (ACQUITY UPLC HSS T3, 1.8 μm, 2.1 mm × 100 mm; Waters Corp.).

Extraction of total RNA

To extract total RNA, frozen plant samples were first homogenized using a Multi-Beads Shocker and 5 mm zirconia beads. Total RNA was then isolated with the RNeasy Plant Mini Kit (Qiagen, Hilden, Germany) with all protocols performed as per the manufacturer's protocol. For RNA-seq library preparation, RNA was further purified by on-column DNase digestion (Qiagen).

RNA-seq analysis

RNA quality was evaluated using a Qubit RNA IQ Assay Kit with a Qubit 4 Fluorometer (Thermo Fisher Scientific, Tokyo, Japan). Samples with RNA IQ scores of 9.5–10.0 were used for RNA-seq. cDNA libraries were prepared with the NEBNext Ultra II Directional RNA Library Prep Kit (E7760S, New England Biolabs, Ipswich, MA, USA), SPRIselect (B23317, Beckman Coulter), the NEBNext Poly(A) mRNA Magnetic Isolation Module (E7490S, New England Biolabs), and NEBNext Multiplex Oligos for Illumina (E7710S, New England Biolabs), with all steps following the manufacturer's instructions. Libraries were sequenced on a NextSeq 500 (Illumina, San Diego, CA, USA), and bcl files were converted to fastq using bcl2fastq (Illumina). Reads were analyzed as described by Notaguchi *et al.* (2014) and mapped to the *Arabidopsis* reference (TAIR10) using Bowtie (Langmead et al., 2009) with the options "--all --best --strata." For normalization, read count data were transformed with rlog and adjusted by subtracting the average expression for each gene using default settings of iDEP v0.96 (Ge et al., 2018). DEGs were extracted using edgeR with a false discovery rate cutoff of 0.15 under the default settings of RNAseqChef v1.1.4 (Etoh et al., 2023). Gene set enrichment was performed using ShinyGO v0.82 (Ge et al., 2020) using the

568 default parameters. During analysis we focused on the Biological Process annotation

569 category.

Reverse transcription and quantitative RT-PCR

Reverse transcription was performed with the ReverTraAce qPCR RT Master Mix with

gDNA Remover (Toyobo, Osaka, Japan). For each reaction, 0.5 µg of RNA was used

as input. Synthesized cDNA was diluted five-fold with sterile water and used as

template for qRT-PCR. Reactions were run on a QuantStudio 1 system (Thermo Fisher

Scientific) using a KOD SYBR qPCR Mix (Toyobo). Relative expression was

determined using the comparative Ct method. *TIP41* was used as an internal reference

gene. Primer sequences are listed in Supplementary Table S9.

Determination of nutrient elements

Shoots were harvested and oven-dried at 80°C for 2 days, ground with a spoon, then

stored in a desiccator with silica gel. For carbon and nitrogen quantification, we

weighed portions of dried samples, packed them into tin capsules, and subjected them to

quantitative analysis using a vario MICRO cube elemental analyzer (Elementar

Analysensysteme GmbH, Hanau, Germany). For other elements (e.g., potassium,

calcium, sulfur, phosphorus, magnesium, sodium, manganese, zinc, iron, boron, copper,

molybdenum, and cobalt), weighed portions of dried samples were digested with HNO_3

followed by H_2O_2. Digests were suspended in 1 mL of 0.08 M HNO_3, diluted tenfold

with 0.08 M HNO_3, and analyzed using inductively coupled plasma mass spectrometry

(Agilent 7800 ICP-MS; Agilent Technologies, Santa Clara, CA, USA). Here, indium

was used as an internal standard.

Nitrate level determination

Nitrate was extracted using 10 volumes of sterile water at 100°C for 20 min. A 5.0 µL

aliquot of supernatant was mixed with 40 µL of 5% (w/v) salicylic acid in concentrated

sulfuric acid and incubated at room temperature for 20 min. A mock treatment with 40

µL concentrated sulfuric acid was prepared in parallel. Finally, 1 mL of 8% (w/v)

NaOH was added to the mixture, and nitrate concentrations was determined by

recording sample absorbance at 410 nm.

Fluorescence microscopy

Fluorescence imaging was performed using an all-in-one fluorescence microscope (BZ-X710, KEYENCE, Osaka, Japan) equipped with a 4× objective (CFI Plan Fluor DL 4×/0.13, Nikon, Tokyo, Japan) and a metal halide lamp (No. 91056, KEYENCE). A narrow-band GFP filter (Ex. 480/20; Em. 510/20; 49020, Chroma, Brattleboro, VT, USA) was used for GFP detection, and a Cy5 filter (Ex. 620/60; Em. 700/75; 49006, Chroma) was used for chlorophyll autofluorescence.

Leaf cellular analysis

The fifth true leaf was excised from the shoot and scanned at 600 dpi (GT-X980; EPSON) to measure total leaf area. The abaxial surface was pressed onto Scotch transparent adhesive tape (Code: 500-3-15-10P; 3M Japan, Tokyo, Japan) and carefully peeled to isolate epidermal cells. Epidermal cell images were captured using a fluorescence microscope (BZ-X710; KEYENCE) equipped with a 20× objective (CFI Plan Apo 20×/0.75; Nikon) and a metal halide lamp (No. 91056, KEYENCE). Thirty epidermal cells were randomly selected, and their areas were measured using ImageJ v1.53t (US National Institutes of Health, Bethesda). The mean area of 30 cells was defined as epidermal cell size. The total epidermal cell number was estimated by dividing the total leaf area by the mean cell size.

Statistical analyses

Data were analyzed using unpaired two-tailed Welch's t-tests and Dunnett's multiple comparisons tests. All statistical analyses were conducted in R v4.3.0 (R Foundation for Statistical Computing, Vienna, Austria).

Data Availability Statement

RNA-seq raw data are deposited in ArrayExpress under accession number E-MTAB-15416.

Acknowledgements

We thank Dr. Ryo Tabata and Dr. Tomoko Niwa for technical advice, Ms. Maki Saiki

for ICP-MS analysis, and Prof. Thomas Schmülling for important input. We also

acknowledge the use of ChatGPT, an AI-based language tool, for support in clarifying

the manuscript, and Enago (www.enago.jp) for English language review.

Author contributions

K.M. and T.H. conceived and designed experiments. K.M., T.S., M.K., Y.T., T.Ka.,

H.S., and T.H. performed the experiments and data analysis. T.Ki., T.N., and T.H.

supervised the study. K.M. wrote the manuscript with contributions from all authors.

All authors approved the final manuscript.

Conflicts of interest

The authors declare that they have no competing interests.

Funding

This study was supported by grants from JSPS KAKENHI [No. JP24KJ1709 to K.M.,

No. 21H02087 to T.Ka., and Nos. 20K05771 and 23K04978 to T.H.] and from JST

SPRING [No. JPMJSP2155 to K.M.].

REFERENCES

Abualia R, Ötvös K, Novák O, Bouguyon E, Domanegg K, Krapp A, Nacry P, Gojon A,

Lacombe B, Benková E. Molecular framework integrating nitrate sensing in root and

auxin-guided shoot adaptive responses. *Proc Natl Acad Sci U S A.*

2022:119(31):e2122460119. 10.1073/pnas.2122460119

Antoniadi I, Novák O, Gelová Z, Johnson A, Plíhal O, Simerský R, Mik V, Vain T,

Mateo-Bonmatí E, Karady M, et al. Cell-surface receptors enable perception of

extracellular cytokinins. *Nat Commun.* 2020:11(1):4284. 10.1038/s41467-020-17700-9

Arnaud D, Lee S, Takebayashi Y, Choi D, Choi J, Sakakibara H, Hwang I.

Cytokinin-mediated regulation of reactive oxygen species homeostasis modulates

stomatal immunity in Arabidopsis. *Plant Cell.* 2017:29(3):543–559.
10.1105/tpc.16.00583

Bartrina I, Otto E, Strnad M, Werner T, Schmülling T. Cytokinin regulates the activity
of reproductive meristems, flower organ size, ovule formation, and thus seed yield in
Arabidopsis thaliana. *Plant Cell.* 2011:23(1):69–80. 10.1105/tpc.110.079079

Bartrina I, Werner S, Schenke A, Gasperini D, Werner T, Schmülling T. Root-derived
cytokinin regulates Arabidopsis flowering time through components of the age pathway.
Plant Physiol. 2025:198(3). 10.1093/plphys/kiaf204

Bhargava A, Clabaugh I, To JP, Maxwell BB, Chiang YH, Schaller GE, Loraine A,
Kieber JJ. Identification of cytokinin-responsive genes using microarray meta-analysis
and RNA-Seq in Arabidopsis. *Plant Physiol.* 2013:162(1):272–294.
10.1104/pp.113.217026

Caesar K, Thamm AMK, Witthöft J, Elgass K, Huppenberger P, Grefen C, Horak J,
Harter K. Evidence for the localization of the Arabidopsis cytokinin receptors AHK3
and AHK4 in the endoplasmic reticulum. *J Exp Bot.* 2011:62(15):5571–5580.
10.1093/jxb/err238

Etoh K, Nakao M. A web-based integrative transcriptome analysis, RNAseqChef,
uncovers the cell/tissue type-dependent action of sulforaphane. *J. Biol. Chem.*
2023:299(6):104810. 10.1016/j.jbc.2023.104810

Frank M, Cortleven A, Novák O, Schmülling T. Root-derived trans-zeatin cytokinin
protects Arabidopsis plants against photoperiod stress. *Plant Cell Environ.*
2020:43:2637–2649. 10.1111/pce.13860

Frébort I, Kowalska M, Hluska T, Frébortová J, Galuszka P. Evolution of cytokinin
biosynthesis and degradation. *J Exp Bot.* 2011:62(8):2431–2452. 10.1093/jxb/err004

Galuszka P, Popelková H, Werner T, Frébortová J, Pospíšilová H, Mik V, Köllmer I,
Schmülling T, Frébort I. Biochemical characterization of cytokinin
oxidases/dehydrogenases from Arabidopsis thaliana expressed in Nicotiana tabacum L.
J Plant Growth Regul. 2007:26(3):255–267. 10.1007/s00344-007-9008-5
698

699 Gautrat P, Buti S, Romanowski A, Lammers M, Matton SEA, Buijs G, Pierik R.
700 Phytochrome-dependent responsiveness to root-derived cytokinins enables coordinated
701 elongation responses to combined light and nitrate cues. *Nat Commun.* 2024:15:8489.
702 10.1038/s41467-024-52828-y
703

704 Ge SX, Son EW, Yao R. iDEP: an integrated web application for differential expression
705 and pathway analysis of RNA-Seq data. *BMC Bioinformatics.* 2018:19:534.
706 10.1186/s12859-018-2486-6
707

708 Ge SX, Jung D, Yao R. ShinyGO: a graphical gene-set enrichment tool for animals and
709 plants. *Bioinformatics.* 2020:36(8):2628–2629. 10.1093/bioinformatics/btz931
710

711 Hachiya T, Oya T, Monden K, Nagae A, Nakagawa T. A cellophane-supported
712 Arabidopsis culture for seamless transfer between different media is useful for studying
713 various nitrogen responses. *Soil Sci. Plant Nutr.* 2021:67(3):277–282.
714 10.1080/00380768.2021.1908094
715

716 Higuchi M, Pischke MS, Mähönen AP, Miyawaki K, Hashimoto Y, Seki M, Kobayashi
717 M, Shinozaki K, Kato T, Tabata S, et al. In planta functions of the Arabidopsis
718 cytokinin receptor family. *Proc Natl Acad Sci U S A.* 2004:101(23):8821–8826.
719 10.1073/pnas.0402887101
720

721 Hirose N, Takei K, Kuroha T, Kamada-Nobusada T, Hayashi H, Sakakibara H.
722 Regulation of cytokinin biosynthesis, compartmentalization and translocation. *J Exp Bot.*
723 2008:59(1):75–83. 10.1093/jxb/erm157
724

Hošek P, Hoyerová K, Kiran NS, Dobrev PI, Zahajská L, Filepová R, Motyka V,

Müller K, Kamínek M. Distinct metabolism of N-glucosides of isopentenyladenine and

trans-zeatin determines cytokinin metabolic spectrum in Arabidopsis. *New Phytol.*

2020:225(6):2423−2438. 10.1111/nph.16310

Hutchison CE, Kieber JJ. Cytokinin signaling in Arabidopsis. *Plant Cell.*

2002:14(suppl_1):S47–S59. 10.1105/tpc.010444

Inoue T, Higuchi M, Hashimoto Y, Seki M, Kobayashi M, Kato T, Tabata S, Shinozaki

K, Kakimoto T. Identification of CRE1 as a cytokinin receptor from Arabidopsis.

Nature. 2001:409(6823):1060–1063. 10.1038/35059117

Jin SH, Ma XM, Kojima M, Sakakibara H, Wang YW, Hou BK. Overexpression of

glucosyltransferase UGT85A1 influences trans-zeatin homeostasis and trans-zeatin

responses likely through O-glucosylation. *Planta.* 2013:237(4):991–999.

10.1007/s00425-012-1818-4

Kiba T, Takei K, Kojima M, Sakakibara H. Side-chain modification of cytokinins

controls shoot growth in Arabidopsis. *Dev Cell.* 2013:27(4):452–461.

10.1016/j.devcel.2013.10.004

Kiba T, Inaba J, Kudo T, Ueda N, Konishi M, Mitsuda N, Takiguchi Y, Kondou Y,

Yoshizumi T, Ohme-Takagi M, et al. Repression of nitrogen starvation responses by

members of the Arabidopsis GARP-type transcription factor NIGT1/HRS1 subfamily.

Plant Cell. 2018:30(4):925–945. 10.1105/tpc.17.00810

Kim HJ, Ryu H, Hong SH, Woo HR, Lim PO, Lee IC, Sheen J, Nam HG, Hwang I.

Cytokinin-mediated control of leaf longevity by AHK3 through phosphorylation of

ARR2 in Arabidopsis. *Proc Natl Acad Sci U S A.* 2006:103(3):814–819.

10.1073/pnas.0505150103

Ko D, Kang J, Kiba T, Park J, Kojima M, Do J, Kim KY, Kwon M, Endler A, Song
W-Y, et al. Arabidopsis ABCG14 is essential for the root-to-shoot translocation of
cytokinin. *Proc Natl Acad Sci USA.* 2014:111(19):7150–7155.
10.1073/pnas.1321519111

Kojima M, Kamada-Nobusada T, Komatsu H, Takei K, Kuroha T, Mizutani M,
Ashikari M, Ueguchi-Tanaka M, Matsuoka M, Suzuki K, et al. Highly sensitive and
high-throughput analysis of plant hormones using MS-probe modification and liquid
chromatographytandem mass spectrometry: an application for hormone profiling in
Oryza sativa. *Plant Cell Physiol.* 2009:50(7):1201–1214. 10.1093/pcp/pcp057

Kubiasová K, Montesinos JC, Šamajová O, Nisler J, Mik V, Semerádová H, Plíhalová
L, Novák O, Marhavý P, Cavallari N, et al. Cytokinin fluoroprobe reveals multiple sites
of cytokinin perception at plasma membrane and endoplasmic reticulum. *Nat Commun.*
2020:11(1):4285. 10.1038/s41467-020-17949-0

Kurakawa T, Ueda N, Maekawa M, Kobayashi K, Kojima M, Nagato Y, Sakakibara H,
Kyozuka J. Direct control of shoot meristem activity by a cytokinin-activating enzyme.
Nature. 2007:445(7128):652–655. 10.1038/nature05504

Kuroha T, Tokunaga H, Kojima M, Ueda N, Ishida T, Nagawa S, Fukuda H, Sugimoto
K, Sakakibara H. Functional analyses of LONELY GUY cytokinin-activating enzymes
reveal the importance of the direct activation pathway in Arabidopsis. *Plant Cell.*
2009:21(10):3152–3169. 10.1105/tpc.109.068676

Landrein B, Formosa-Jordan P, Malivert A, Schuster C, Melnyk CW, Yang W,
Turnbull C, Meyerowitz EM, Locke JCW, Jönsson H. Nitrate modulates stem cell
dynamics in Arabidopsis shoot meristems through cytokinins. *Proc Natl Acad Sci U S A.*
2018:115(6):1382–1387. 10.1073/pnas.1718670115

Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient
alignment of short DNA sequences to the human genome. *Genome Biol.*
2009:10(3):R25. 10.1186/gb-2009-10-3-r25

Liu KH, Liu M, Lin Z, Wang ZF, Chen B, Liu C, Guo A, Konishi M, Yanagisawa S,
Wagner G, et al. NIN-like protein 7 transcription factor is a plant nitrate sensor. *Science.*
2022:377(6613):1419–1425. 10.1126/science.add1104

Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H,
Yanagisawa S. A NIGT1-centred transcriptional cascade regulates nitrate signalling and
incorporates phosphorus starvation signals in Arabidopsis. *Nat Commun.*
2018:9(1):1376. 10.1038/s41467-018-03832-6

Matsumoto-Kitano M, Kusumoto T, Tarkowski P, Kinoshita-Tsujimura K, Václavíková
K, Miyawaki K, Kakimoto T. Cytokinins are central regulators of cambial activity. *Proc*
Natl Acad Sci USA. 2008:105(50):20027–20031. 10.1073/pnas.0805619105

Miyawaki K, Matsumoto-Kitano M, Kakimoto T. Expression of cytokinin biosynthetic
isopentyltransferase genes in Arabidopsis: tissue specificity and regulation by auxin,
cytokinin, and nitrate. *Plant J.* 2004:37(1):128–138.
10.1046/j.1365-313X.2003.01945.x

Miyawaki K, Tarkowski P, Matsumoto-Kitano M, Kato T, Sato S, Tarkowska D, Tabata
S, Sandberg G, Kakimoto T. Roles of Arabidopsis ATP/ADP isopentenyltransferases
and tRNA isopentenyltransferases in cytokinin biosynthesis. *Proc Natl Acad Sci USA.*
2006:103(44):16598–16603. 10.1073/pnas.0603522103

Monden K, Kojima M, Takebayashi Y, Suzuki T, Nakagawa T, Sakakibara H, Hachiya
T. Root-specific Reduction of Cytokinin Perception Enhances Shoot Growth in
Arabidopsis thaliana. *Plant Cell Physiol.* 2022:63(4):484–493. 10.1093/pcp/pcac013

Müller B, Sheen J. Arabidopsis Cytokinin Signaling Pathway. *Sci. STKE.*
2007:407:cm5. 10.1126/stke.4072007cm5

Nam YJ, Tran LS, Kojima M, Sakakibara H, Nishiyama R, Shin R. Regulatory roles of
cytokinins and cytokinin signaling in response to potassium deficiency in Arabidopsis.
PLoS One. 2012:7(10): e47797. 10.1371/journal.pone.0047797

Niemann MCE, Weber H, Hluska T, Leonte G, Anderson SM, Novak O, Senes A,
Werner T. The cytokinin oxidase/dehydrogenase CKX1 is a membrane-bound protein
requiring homooligomerization in the endoplasmic reticulum for its cellular activity.
Plant Physiol. 2018:176(3):2024–2039. 10.1104/pp.17.00925

Notaguchi M, Higashiyama T, Suzuki T. Identification of mRNAs that move over long
distances using an RNA-Seq analysis of Arabidopsis/Nicotiana benthamiana
heterografts. *Plant Cell Physiol.* 2015:56(2):311–321. 10.1093/pcp/pcu210

Osugi A, Kojima M, Takebayashi Y, Ueda N, Kiba T, Sakakibara H. Systemic transport
of trans-zeatin and its precursor have differing roles in Arabidopsis shoots. *Nat Plants.*
2017:3:17112. 10.1038/nplants.2017.112

Rahayu YS, Walch-Liu P, Neumann G, Römheld V, von Wirén N, Bangerth F.
Root-derived cytokinins as long-distance signals for NO3⁻-induced stimulation of leaf
growth. *J Exp Bot.* 2005:56(414):1143–1152. 10.1093/jxb/eri107

Riefler M, Novak O, Strnad M, Schmu☐lling T. Arabidopsis cytokinin receptor mutants
reveal functions in shoot growth, leaf senescence, seed size, germination, root
development, and cytokinin metabolism. *Plant Cell.* 2006:18(1):40–54.
10.1105/tpc.105.037796

Romanov GA, Lomin SN, Schmülling T. Biochemical characteristics and
ligand-binding properties of Arabidopsis cytokinin receptor AHK3 compared to

CRE1/AHK4 as revealed by a direct binding assay. *J Exp Bot.* 2006:57(15):4051–4058.

10.1093/jxb/erl179

Sakakibara H, Suzuki M, Takei K, Deji A, Taniguchi M, Sugiyama T. A response‐

regulator homologue possibly involved in nitrogen signal transduction mediated by

cytokinin in maize. *Plant J.* 1998:14:337–344. 10.1046/j.1365-313X.1998.00134.x

Sakakibara H, Takei K, Hirose N. Interactions between nitrogen and cytokinin in the

regulation of metabolism and development. *Trends Plant Sci.* 2006:11: 440–448.

10.1016/j.tplants.2006.07.004

Sakakibara H. Cytokinin biosynthesis and transport for systemic nitrogen signaling.

Plant J. 2021:105(2):421–430. 10.1111/tpj.15011

Sakakibara H. Five unaddressed questions about cytokinin biosynthesis. *J. Exp. Bot.*

2025:76(7):1941–1949. 10.1093/jxb/erae348

Schmülling T, Werner T, Riefler M, Krupková E, Bartrina y Manns I. Structure and

function of cytokinin oxidase/dehydrogenase genes of maize, rice, Arabidopsis and

other species. *J Plant Res.* 2003:116:241–252. 10.1007/s10265-003-0096-4

Spichal L, Rakova NY, Riefler M, Mizuno T, Romanov GA, Strnad M, Schmülling T.

Two cytokinin receptors of Arabidopsis thaliana, CRE1/AHK4 and AHK3, differ in

their ligand specificity in a bacterial assay. *Plant Cell Physiol.* 2004:45(9):1299–1305.

10.1093/pcp/pch132

Stolz A, Riefler M, Lomin SN, Achazi K, Romanov GA, Schmülling T. The specificity

of cytokinin signalling in Arabidopsis thaliana is mediated by differing ligand affinities

and expression profiles of the receptors. *Plant J.* 2011:67(1):157–168.

10.1111/j.1365-313X.2011.04584.x

Takei K, Sakakibara H, Sugiyama T. Identification of genes encoding adenylate
isopentenyltransferase, a cytokinin biosynthesis enzyme, in Arabidopsis thaliana. *J Biol*
Chem. 2001a:276(28):26405–26410. 10.1074/jbc.M102130200
882

883   Takei K, Sakakibara H, Taniguchi M, Sugiyama T. Nitrogen-dependent accumulation
884   of cytokinins in root and the translocation to leaf: Implication of cytokinin species that
885   induces gene expression of maize response regulator. *Plant Cell Physiol.*
886   2001b:42(1):85–93. 10.1093/pcp/pce009
887

888   Takei K, Yamaya T, Sakakibara H. Arabidopsis CYP735A1 and CYP735A2 encode
889   cytokinin hydroxylases that catalyze the biosynthesis of trans-zeatin. *J Biol Chem.*
890   2004a:279(40):41866–41872. 10.1074/jbc.M406337200
891

892   Takei K, Ueda N, Aoki K, Kuromori T, Hirayama T, Shinozaki K, Yamaya T,
893   Sakakibara H. AtIPT3 is a key determinant of nitrate-dependent cytokinin biosynthesis
894   in Arabidopsis. *Plant Cell Physiol.* 2004b:45(8):1053–1062. 10.1093/pcp/pch119
895

896   To JPC, Haberer G, Ferreira FJ, Deruère J, Mason MG, Schaller GE, Alonso JM, Ecker
897   JR, Kieber JJ. Type-A ARRs are partially redundant negative regulators of cytokinin
898   signaling in Arabidopsis. *Plant Cell.* 2004:16(3):658–671. 10.1105/tpc.018978
899

900   Wang J, Ma XM, Kojima M, Sakakibara H, Hou BK. N-Glucosyltransferase UGT76C2
901   is involved in cytokinin homeostasis and cytokinin response in Arabidopsis thaliana.
902   *Plant Cell Physiol.* 2011:52 (12):2200–2213. 10.1093/pcp/pcr152
903

904   Wang J, Ma XM, Kojima M, Sakakibara H, Hou BK. Glucosyltransferase UGT76C1
905   finely modulates cytokinin responses via cytokinin N-glucosylation in Arabidopsis
906   thaliana. *Plant Physiol. Biochem.* 2013:65:9–16. 10.1016/j.plaphy.2013.01.012
907

908   Werner T, Motyka V, Strnad M, Schmülling T. Regulation of plant growth by cytokinin.
909   *Proc. Natl. Acad. Sci. USA.* 2001:98(18):10487–10492. 10.1073/pnas.171304098
910

Werner T, Motyka V, Laucou V, Smets R, Van Onckelen H, Schmülling T.
Cytokinin-deficient transgenic Arabidopsis plants show multiple developmental
alterations indicating opposite functions of cytokinins in the regulation of shoot and root
meristem activity. *Plant Cell.* 2003:15(11):2532–2550. 10.1105/tpc.014928

Werner T, Köllmer I, Bartrina I, Holst K, Schmülling T. New insights into the biology
of cytokinin degradation. *Plant Biol.* 2006:8(3):371–381. 10.1055/s-2006-923928

Wulfetange K, Lomin SN, Romanov GA, Stolz A, Heyl A, Schmülling T. The
cytokinin receptors of Arabidopsis are located mainly to the endoplasmic reticulum.
Plant Physiol. 2011:156(4):1808–1818. 10.1104/pp.111.180539

Zhang K, Novak O, Wei Z, Gou M, Zhang X, Yu Y, Yang H, Cai Y, Strnad M, Liu CJ.
Arabidopsis ABCG14 protein controls the acropetal translocation of root-synthesized
cytokinins. *Nat Commun.* 2014:5(1):3274. 10.1038/ncomms4274

Zhao J, Ding B, Zhu E, Deng X, Zhang M, Zhang P, Wang L, Dai Y, Xiao S, Zhang C,
et al. Phloem unloading via the apoplastic pathway is essential for shoot distribution of
root-synthesized   cytokinins.   *Plant   Physiol.*   2021:186(4):2111–2123.
10.1093/plphys/kiab188

Zhao J, Deng X, Qian J, Liu T, Ju M, Li J, Yang Q, Zhu X, Li W, Liu C-J, et al.
Arabidopsis ABCG14 forms a homodimeric transporter for multiple cytokinins and
mediates long-distance transport of isopentenyladenine-type cytokinins. *Plant Commun.*
2023:4(2):100468. 10.1016/j.xplc.2022.100468

Zürcher E, Tavor-Deslex D, Lituiev D, Enkerli K, Tarr PT, Müller B. A robust and
sensitive synthetic sensor to monitor the transcriptional output of the cytokinin
signaling   network   in   planta.   *Plant   Physiol.*   2013:161(3):1066–1075.
10.1104/pp.112.211763

FIGURE LEGENDS

Figure 1. Effects of root-specific CK receptor deficiency on shoot growth in
medium. (A) Representative shoots of grafted plants. Scale bars = 5 mm. (B–D) Shoot
fresh weight (B), leaf blade area (C), and leaf number (D) of grafted plants. Data pooled
from two independent grafting experiments are shown as mean ± SD (n = 12). Asterisks
indicate significant differences relative to WT/WT (*$P < 0.05$, **$P < 0.01$, ***$P <$
0.001; Dunnett's test). Numbers on graphs show mean values relative to WT/WT (set =
1). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*,
ahk3-7; *a4*, *cre1-2*; *a23*, *ahk2-5 ahk3-7*; *a24*, *ahk2-5 cre1-2*; *a34*, *ahk3-7 cre1-2*.

Figure 2. Effects of root-specific CK receptor deficiency on root CK concentration.
(A–B) Root concentrations of active form iP and *t*Z and their precursors (A), and
inactive form iP- and *t*Z-type CKs (B) in grafted plants. Data are pooled from three
independent grafting experiments and are reported as mean ± SD (n = 3). Asterisks
indicate significant differences relative to WT/WT (**$P < 0.01$, ***$P < 0.001$;
Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*,
ahk2-5; *a3*, *ahk3-7*; *a4*, *cre1-2*; FW, fresh weight; *t*Z, *trans*-zeatin; iP,
N^6-(Δ^2-isopentenyl) adenine; iP and their precursors = sum of iP, iPR, and iPRPs;
inactive form iP-type CKs = sum of iP7G and iP9G; *t*Z and their precursors = sum of *t*Z,
*t*ZR, and *t*ZRPs; inactive form *t*Z-type CKs = sum of *t*Z7G, *t*Z9G, *t*ZOG, and *t*ZROG.

Figure 3. Effects of root-specific CK receptor deficiency on the root transcriptome.
(A) Volcano plots of DEGs between WT/WT and WT/*ahk2*, WT/*ahk3*, or WT/*ahk4*
roots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023) using edgeR,
applying a 1.5-fold change threshold and FDR cutoff of 0.15 (all other parameters
default). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted p-value (padj). Each
dot represents a single gene. Red, blue, and grey indicate upregulated, downregulated,
and non-significant genes, respectively. $\log_2$ FC was limited to –4 to 4 and padj to $\leqq$
50. Full-range volcano plots are provided in Supplementary Fig. S4. (B) Mean
normalized transcript levels of CK-inducible and CK-repressible genes (mean ± SD).
Gene list obtained from Bhargava *et al.* (2013). (C) Transcriptional changes in CK
biosynthesis and inactivation genes, visualized as heatmaps with MORPHEUS

(https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each

graft to WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per

row. (D) Root expression of *CKX4* in grafted plants by qRT-PCR. Relative transcript

levels were normalized to the WT/WT root mean (set = 1) and $\log_2$-transformed. Data

pooled from two independent grafting experiments are shown as mean ± SD (n = 6).

Asterisks indicate significant differences relative to WT/WT (**$P < 0.01$, ***$P <$

0.001; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT,

Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FC, fold change.

Figure 4. Effects of root-specific CK receptor deficiency on CK concentration of

xylem sap. (A–B) Xylem sap concentrations of iP- and *t*Z-type CKs (A) and of *t*Z, *t*ZR,

and *t*ZRPs (B) in grafted plants. Data pooled from three independent grafting

experiments are reported as mean ± SD (n = 3). Asterisks indicate significant

differences relative to WT/WT (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$; Dunnett's test).

Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*,

ahk3-7; *a4*, *cre1-2*; *t*Z, *trans*-zeatin; *t*ZR, *t*Z-riboside; *t*ZRPs, *t*ZR 5′-phosphates; iP-type

CKs = sum of iPR, iPRPs, and iP7G; *t*Z-type CKs = sum of *t*Z, *t*ZR, *t*ZRPs, *t*Z7G, *t*Z9G,

and *t*ZOG.

Figure 5. Effects of root-specific CK receptor deficiency on the shoot

transcriptome. (A) Volcano plots of DEGs between WT/WT and WT/*ahk2*, WT/*ahk3*,

or WT/*ahk4* shoots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023)

using edgeR, applying a 1.5-fold change threshold and FDR cutoff of 0.15 (default

parameters otherwise). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted

p-value (padj). Each dot represents one gene; red, blue, and grey indicate upregulated,

downregulated, and non-significant genes, respectively. $\log_2$ FC values were limited to

–4 to 4 and padj values to $\leqq 5$. (B) Lollipop plots showing GO enrichment analysis

from the 163 genes upregulated in WT/*ahk3* shoots, generated with ShinyGO (Ge et al.,

2020) using default settings for biological process terms. (C) Heatmaps of

transcriptional changes in type-A *ARR* genes, visualized with MORPHEUS

(https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each

graft to the WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per

row. (D) Shoot expression of *ARR5* determined by qRT-PCR. Values were normalized

to the WT/WT shoot mean (set = 1) and $\log_2$-transformed. Data pooled from two

independent grafting experiments are reported as mean ± SD (n = 6). Asterisks indicate

significant differences relative to WT/WT (*$P < 0.05$; Dunnett's test). Grafted plants

are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*,

cre1-2; FC, fold change.

1012

Figure 6. Effects of root-specific *AHK3* deficiency on shoot growth via CK

signaling. (A–E) Shoot fresh weight and leaf blade area in grafted plants: (A) WT/WT,

WT/*ahk3*, WT/*abcg14*, and WT/*ahk3 abcg14*; (B) WT/WT, WT/*ahk3*, WT/*cypDM*, and

WT/*ahk3 cypDM*; (C) WT/WT, WT/*ahk3*, *ahk2*/*ahk3*, *ahk3*/*ahk3*, and *ahk4*/*ahk3*; (D)

WT/WT, WT/*ahk3*, *ahk23*/*ahk3*, *ahk24*/*ahk3*, and *ahk34*/*ahk3*; (E) *TCSn:GFP*/WT and

TCSn:GFP/*ahk3*. (F) Representative fluorescent image of the adaxial side of the sixth

true leaf in grafted plants. Green = GFP; red = autofluorescence. Scale bars = 1 mm.

GFP intensity was quantified with ImageJ as the mean signal in the region marked by a

white dashed box. (G) Epidermal cell size and number on the abaxial side of the fifth

true leaf. Data pooled from two independent grafting experiments are reported as mean

± SD (n = 12 (A–E), 11 (F), or 6 (G)). Asterisks (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$)

indicate significant differences relative to WT/WT by Dunnett's test. Pairwise

differences by Welch's *t*-test are marked with asterisks above horizontal lines. Graph

numbers show mean values relative to WT/WT (set = 1). Grafted plants are denoted as

"scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; *a23*, *ahk2-5*

ahk3-7; *a24*, *ahk2-5 cre1-2*; *a34*, *ahk3-7 cre1-2*; *g14*, *abcg14*; *DM*, *cyp735a1-2*

cyp735a2-2; *a3g14*, *ahk3-7 abcg14*; *a3DM*, *ahk3-7 cyp735a1-2 cyp735a2-2*; *TCS*,

TCSn:GFP; ns, not significant.

1031

Figure 7. Effects of root-specific *AHK3* or *CKX4* deficiency on shoot growth in

response to nitrate signaling. (A, B, E) Shoot fresh weight and leaf blade area of

grafted plants 5 days after transfer from 5 mM NH_4^+ medium to either 5 mM NH_4^+ +

0.5 mM KNO_3 or 5 mM NH_4^+ + 0.5 mM KCl: (A) WT/WT and WT/*ahk3*; (B) WT/WT,

WT/*ahk3*, WT/*abcg14*, and WT/*ahk3 abcg14*; (E) WT/WT, WT/*ckx4-1*, and WT/*ckx4-2*.

(C) Root expression of *IPT3*, *IPT7*, *CYP735A1*, *CYP735A2*, *CKX1*, *CKX3*, *CKX4*,

CKX5, *CKX6*, *CKX7*, *UGT76C1*, *UGT76C2*, and *UGT85A1* 4 hours after transfer.

Heatmaps were generated with MORPHEUS

(https://software.broadinstitute.org/morpheus) from $\log_2$ fold change values relative to

WT/WT under 5 mM NH_4^+ + 0.5 mM KCl (set = 1). Color scale: blue (minimum) to

yellow (maximum) per row. (D) RT-PCR analysis of *CKX4* in WT, *ckx4-1*, and *ckx4-2*

seedlings grown 7 days on 1/2 MS medium. *TIP41* was the control. Primers are listed in

Supplementary Table S8. Data are shown as mean ± SD with replicates as follows: (A)

n = 18, two replicates; (B) n = 18, three replicates; (C) n = 6, two replicates; (E) n = 24,

four replicates. Asterisks (*P < 0.05, **P < 0.01, ***P < 0.001) mark significant

differences relative to WT/WT under 5 mM NH_4^+ + 0.5 mM KCl (Dunnett's test).

Pairwise comparisons used Welch's *t*-tests (A). For multiple treatments vs. one control,

we used Dunnett's tests (B, E). Significant differences are shown by asterisks above

horizontal lines. Graph numbers indicate mean values relative to WT/WT under 5 mM

NH_4^+ + 0.5 mM KCl (set = 1). Grafted plants are denoted as "scion line/rootstock line."

WT, Col-0; *a3*, *ahk3-7*; *g14*, *abcg14*; *a3g14*, *ahk3-7 abcg14*; *c4-1*, *ckx4-1*; *c4-2*, *ckx4-2*;

ns, not significant; KCl, 5 mM NH_4^+ + 0.5 mM KCl; KNO_3, 5 mM NH_4^+ + 0.5 mM

KNO_3.

Figure 1. Effects of root-specific CK receptor deficiency on shoot growth in medium. (A) Representative shoots of grafted plants. Scale bars = 5 mm. (B–D) Shoot fresh weight (B), leaf blade area (C), and leaf number (D) of grafted plants. Data pooled from two independent grafting experiments are shown as mean ± SD (n = 12). Asterisks indicate significant differences relative to WT/WT (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$; Dunnett's test). Numbers on graphs show mean values relative to WT/WT (set = 1). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; *a23*, *ahk2-5 ahk3-7*; *a24*, *ahk2-5 cre1-2*; *a34*, *ahk3-7 cre1-2*.

Figure 2. Effects of root-specific CK receptor deficiency on root CK concentration. (A–B) Root concentrations of active form iP and *t*Z and their precursors (A), and inactive form iP- and *t*Z-type CKs (B) in grafted plants. Data are pooled from three independent grafting experiments and are reported as mean $\pm$ SD (n = 3). Asterisks indicate significant differences relative to WT/WT (**P < 0.01, ***P < 0.001; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FW, fresh weight; *t*Z, *trans*-zeatin; iP, N^6-(Δ^2-isopentenyl) adenine; iP and their precursors = sum of iP, iPR, and iPRPs; inactive form iP-type CKs = sum of iP7G and iP9G; *t*Z and their precursors = sum of *t*Z, *t*ZR, and *t*ZRPs; inactive form *t*Z-type CKs = sum of *t*Z7G, *t*Z9G, *t*ZOG, and *t*ZROG.

Figure 3. Effects of root-specific CK receptor deficiency on the root transcriptome. (A) Volcano plots of DEGs between WT/WT and WT/*ahk2*, WT/*ahk3*, or WT/*ahk4* roots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023) using edgeR, applying a 1.5-fold change threshold and FDR cutoff of 0.15 (all other parameters default). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted p-value (padj). Each dot represents a single gene. Red, blue, and grey indicate upregulated, downregulated, and non-significant genes, respectively. $\log_2$ FC was limited to –4 to 4 and padj to $\leqq$50. Full-range volcano plots are provided in Supplementary Fig. S4. (B) Mean normalized transcript levels of CK-inducible and CK-repressible genes (mean $\pm$ SD). Gene list obtained from Bhargava *et al.* (2013). (C) Transcriptional changes in CK biosynthesis and inactivation genes, visualized as heatmaps with MORPHEUS (https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each graft to WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per row. (D) Root expression of *CKX4* in grafted plants by qRT-PCR. Relative transcript levels were normalized to the WT/WT root mean (set = 1) and $\log_2$-transformed. Data pooled from two independent grafting experiments are shown as mean $\pm$ SD (n = 6). Asterisks indicate significant differences relative to WT/WT (**$P < 0.01$, ***$P < 0.001$; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FC, fold change.

Figure 4. Effects of root-specific CK receptor deficiency on CK concentration of xylem sap. (A–B) Xylem sap concentrations of iP- and *t*Z-type CKs (A) and of *t*Z, *t*ZR, and *t*ZRPs (B) in grafted plants. Data pooled from three independent grafting experiments are reported as mean ± SD (n = 3). Asterisks indicate significant differences relative to WT/WT (**P* < 0.05, ***P* < 0.01, ****P* < 0.001; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; *t*Z, *trans*-zeatin; *t*ZR, *t*Z-riboside; *t*ZRPs, *t*ZR 5′-phosphates; iP-type CKs = sum of iPR, iPRPs, and iP7G; *t*Z-type CKs = sum of *t*Z, *t*ZR, *t*ZRPs, *t*Z7G, *t*Z9G, and *t*ZOG.

Figure 5. Effects of root-specific CK receptor deficiency on the shoot transcriptome. (A) Volcano plots of DEGs between WT/WT and WT/*ahk2*, WT/*ahk3*, or WT/*ahk4* shoots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023) using edgeR, applying a 1.5-fold change threshold and FDR cutoff of 0.15 (default parameters otherwise). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted p-value (padj). Each dot represents one gene; red, blue, and grey indicate upregulated, downregulated, and non-significant genes, respectively. (B) Lollipop plots showing GO enrichment analysis from the 163 genes upregulated in WT/*ahk3* shoots, generated with ShinyGO (Ge et al., 2020) using default settings for biological process terms. (C) Heatmaps of transcriptional changes in type-A *ARR* genes, visualized with MORPHEUS (https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each graft to the WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per row. (D) Shoot expression of *ARR5* determined by qRT-PCR. Values were normalized to the WT/WT shoot mean (set = 1) and $\log_2$-transformed. Data pooled from two independent grafting experiments are reported as mean $\pm$ SD (n = 6). Asterisks indicate significant differences relative to WT/WT (*P < 0.05; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FC, fold change.

Figure 6. Effects of root-specific *AHK3* deficiency on shoot growth via CK signaling. (A–E) Shoot fresh weight and leaf blade area in grafted plants: (A) WT/WT, WT/*ahk3*, WT/*abcg14*, and WT/*ahk3 abcg14*; (B) WT/WT, WT/*ahk3*, WT/*cypDM*, and WT/*ahk3 cypDM*; (C) WT/WT, WT/*ahk3*, *ahk2*/*ahk3*, *ahk3*/*ahk3*, and *ahk4*/*ahk3*; (D) WT/WT, WT/*ahk3*, *ahk23*/*ahk3*, *ahk24*/*ahk3*, and *ahk34*/*ahk3*; (E) *TCSn:GFP*/WT and *TCSn:GFP*/*ahk3*. (F) Representative fluorescent image of the adaxial side of the sixth true leaf in grafted plants. Green = GFP; red = autofluorescence. Scale bars = 1 mm. GFP intensity was quantified with ImageJ as the mean signal in the region marked by a white dashed box. (G) Epidermal cell size and number on the abaxial side of the fifth true leaf. Data pooled from two independent grafting experiments are reported as mean ± SD (n = 12 (A–E), 11 (F), or 6 (G)). Asterisks (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$) indicate significant differences relative to WT/WT by Dunnett's test. Pairwise differences by Welch's *t*-test are marked with asterisks above horizontal lines. Graph numbers show mean values relative to WT/WT (set = 1). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; *a23*, *ahk2-5 ahk3-7*; *a24*, *ahk2-5 cre1-2*; *a34*, *ahk3-7 cre1-2*; *g14*, *abcg14*; *DM*, *cyp735a1-2 cyp735a2-2*; *a3g14*, *ahk3-7 abcg14*; *a3DM*, *ahk3-7 cyp735a1-2 cyp735a2-2*; *TCS*, *TCSn:GFP*; ns, not significant.

Figure 7. Effects of root-specific *AHK3* or *CKX4* deficiency on shoot growth in response to nitrate signaling. (A, B, E) Shoot fresh weight and leaf blade area of grafted plants 5 days after transfer from 5 mM NH_4^+ medium to either 5 mM NH_4^+ + 0.5 mM KNO_3 or 5 mM NH_4^+ + 0.5 mM KCl: (A) WT/WT and WT/*ahk3*; (B) WT/WT, WT/*ahk3*, WT/*abcg14*, and WT/*ahk3 abcg14*; (E) WT/WT, WT/*ckx4-1*, and WT/*ckx4-2*. (C) Root expression of *IPT3*, *IPT7*, *CYP735A1*, *CYP735A2, CKX1*, *CKX3*, *CKX4*, *CKX5*, *CKX6*, *CKX7*, *UGT76C1*, *UGT76C2*, and *UGT85A1* 4 hours after transfer. Heatmaps were generated with MORPHEUS (https://software.broadinstitute.org/morpheus) from $\log_2$ fold change values relative to WT/WT under 5 mM NH_4^+ + 0.5 mM KCl (set = 1). Color scale: blue (minimum) to yellow (maximum) per row. (D) RT-PCR analysis of *CKX4* in WT, *ckx4-1*, and *ckx4-2* seedlings grown 7 days on 1/2 MS medium. *TIP41* was the control. Primers are listed in Supplementary Table S8. Data are shown as mean ± SD with replicates as follows: (A) n = 18, two replicates; (B) n = 18, three replicates; (C) n = 6, two replicates; (E) n = 24, four replicates. Asterisks (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$) mark significant differences relative to WT/WT under 5 mM NH_4^+ + 0.5 mM KCl (Dunnett's test). Pairwise comparisons used Welch's *t*-tests (A). For multiple treatments vs. one control, we used Dunnett's tests (B, E). Significant differences are shown by asterisks above horizontal lines. Graph numbers indicate mean values relative to WT/WT under 5 mM NH_4^+ + 0.5 mM KCl (set = 1). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a3*, *ahk3-7*; *g14*, *abcg14*; *a3g14*, *ahk3-7 abcg14*; *c4-1*, *ckx4-1*; *c4-2*, *ckx4-2*; ns, not significant; KCl, 5 mM NH_4^+ + 0.5 mM KCl; KNO$_3$, 5 mM NH_4^+ + 0.5 mM KNO_3.