TITLE

Cytokinin receptor AHK3 modulates root-to-shoot transport of *trans*-zeatin-type

cytokinin

AUTHORS

Kota Monden[a†], Takamasa Suzuki[b], Mikiko Kojima[c], Yumiko Takebayashi[c], Takehiro

Kamiya[d], Takatoshi Kiba[e,f], Hitoshi Sakakibara[c,f], Tsuyoshi Nakagawa[a], and Takushi

Hachiya[a]

[a]Department of Molecular and Functional Genomics, Interdisciplinary Center for Science

Research, Shimane University, 1060 Nishikawatsu-cho, Matsue, Shimane 690-8504,

Japan

[b]College of Bioscience and Biotechnology, Chubu University, 1200 Matsumoto-cho,

Kasugai, Aichi 487-8501, Japan

[c]RIKEN Center for Sustainable Resource Science, 1-7-22 Suehiro-cho, Tsurumi-ku,

Yokohama, Kanagawa 230-0045, Japan

[d]Graduate School of Agricultural and Life Sciences, Tokyo University, 1-1-1 Yayoi,

Bunkyo-ku, Tokyo 113-8657, Japan

[e]Institute of Plant Science and Resources, Okayama University, 2-20-1 Chuo, Kurashiki,

Okayama 710-0046, Japan

[f]Graduate School of Bioagricultural Sciences, Nagoya University, Furo-cho, Chikusa-ku,

Nagoya, Aichi 464-8601, Japan

[†]Corresponding author: Kota Monden (n23d203@matsu.shimane-u.ac.jp)

30 **RUNNING HEAD**

31 AHK3 modulates long-distance cytokinin transport.

ABSTRACT

Trans-zeatin (*t*Z)-type cytokinins are synthesized in roots in response to nitrogen

availability, transported to shoot via xylem, and coordinate diverse physiological

processes in aerial organs. Within this mechanism, the regulation of cytokinin

biosynthesis by nitrate signaling via NIN-like protein 7 as well as the loading of *t*Z-type

cytokinins into xylem by ATP-binding cassette transporter G14 have been well studied.

However, the roles of other components remain unclear. Here, we show that cytokinin

perception and degradation in roots, as mediated by Arabidopsis histidine kinase 3

(AHK3) and cytokinin oxidase/dehydrogenase 4 (CKX4), modulate root-to-shoot

transport of *t*Z-type cytokinins. Grafting experiments demonstrated that root-specific

AHK3 deficiency systemically increased leaf blade area through long-distance signals of

root-derived *t*Z-type cytokinin, perceived by shoot-expressed AHK3. Transcriptome,

hormonome, and reporter analyses revealed that root-specific *AHK3* deficiency reduced

CKX4 expression in roots, elevating *t*Z-type cytokinin levels in roots and xylem sap and

thereby enhancing leaf cytokinin response. Transfer experiments manipulating root

nitrate levels showed that root-specific *AHK3* deficiency promoted leaf blade area in a

manner dependent on both nitrate and root-derived *t*Z-type cytokinin signaling. Moreover,

both nitrate signals and root-expressed AHK3 are required for maximal *CKX4* induction

in roots, and root-specific *CKX4* deficiency enhanced leaf blade area in a nitrate-

dependent manner. These findings reveal a novel mechanism in which an AHK3–CKX4

module governs root-to-shoot transport of *t*Z-type cytokinins, fine-tuning leaf size

according to nitrogen availability in roots.

INTRODUCTION

Cytokinins (CKs) are essential plant hormones that coordinate diverse developmental and

physiological processes, including cell division, shoot and root growth, vascular

formation, and responses to stress and nutrients (Werner et al., 2001; Riefler et al., 2006;

Matsumoto-Kitano et al., 2008; Sakakibara, 2021). In *Arabidopsis*, the biologically active

CK forms include N^6-(Δ^2-isopentenyl) adenine (iP) and *trans*-zeatin (*t*Z). The initial and

rate-limiting step of CK biosynthesis involves N^6-prenylation of adenosine 5′-phosphates

catalyzed by adenosine phosphate-isopentenyltransferases (IPTs), producing iP-riboside

5′-phosphates (iPRPs) (Takei et al., 2001a; Miyawaki et al., 2006). These intermediates

are subsequently hydroxylated by cytochrome P450 monooxygenases CYP735A1 and CYP735A2 to form tZ-riboside 5′-phosphates (tZRPs) (Takei et al., 2004a). Ribotide precursors are then converted into iP and tZ by CK-activating enzyme members of the lonely guy (LOG) family (Kurakawa et al., 2007; Kuroha et al., 2009). CK homeostasis is further modulated by reversible O-glucosylation and tentatively reversible N-glucosylation catalyzed by glucosyltransferases (UGTs), and by irreversible degradation by cytokinin oxidase/dehydrogenases (CKXs) (Werner et al., 2006; Wang et al., 2011; Jin et al., 2013; Wang et al., 2013; Hošek et al., 2020). CK perception and signaling rely on a multistep His-to-Asp phosphorelay system (Hutchison and Kieber, 2002; Müller and Sheen, 2007). Three membrane-spanning histidine kinases, Arabidopsis histidine kinase (AHK)2, AHK3, and CRE1/AHK4, act as CK receptors (Inoue et al., 2001). Upon ligand binding, the phosphoryl group is transferred from AHKs to histidine phosphotransfer proteins, which shuttle into the nucleus and activate type-B response regulators (Hutchison and Kieber, 2002; Müller and Sheen, 2007). These transcription factors induce CK-responsive genes, including type-A response regulators that function as negative feedback elements (To et al., 2004).

In addition to local roles, CKs act as long-distance signaling molecules that coordinate developmental and physiological processes between roots and shoots (Sakakibara, 2021). In *Arabidopsis*, tZ-type CKs predominate in xylem sap, whereas iP-type CKs are more abundant in phloem sap (Hirose et al., 2008; Osugi et al., 2017). This distribution reflects tissue-specific expression of CK biosynthetic genes: *CYP735A2* is strongly expressed in root vasculature, while *IPT3* is mainly expressed in phloem tissues of both roots and shoots (Miyawaki et al., 2004; Takei et al., 2004a,b; Kiba et al., 2013). These activities establish a directional flow of CK species, with root-derived tZ-type CKs transported to shoots via xylem and shoot-derived iP-type CKs transported to roots via phloem tissue (Hirose et al., 2008; Matsumoto-Kitano et al., 2008; Kiba et al., 2013). A central component of the root-to-shoot CK transport system is ATP-binding cassette transporter G14 (ABCG14), which loads CKs into xylem in root stele and also contributes to phloem unloading in shoots (Ko et al., 2014; Zhang et al., 2014; Zhao et al., 2021). Heterologous transport assays in yeast have shown that ABCG14 transports multiple CK species, including both iP- and tZ-types, underscoring its pivotal role in systemic CK distribution and signaling (Zhao et al., 2023).

Root-to-shoot CK transport has been found to be tightly linked to plant

nitrogen status. For example, supplying inorganic nitrogen to root activates CK signaling

in leaves and increases xylem transport of *t*Z-riboside (*t*ZR) following nitrate application

(Sakakibara et al., 1998; Takei et al., 2001b). Moreover, nitrate-induced changes in leaf

growth rate correlate with *t*Z and *t*ZR levels in xylem sap and leaves (Rahayu et al., 2005).

In *Arabidopsis*, nitrate signaling regulates expression of *IPT3* and *CYP735A2* through

NIN-like protein 7 (NLP7) and Nitrate-inducible GARP-type transcriptional repressor 1s

(NIGT1s), these transcription factors that are central to the primary nitrate response

(Miyawaki et al., 2004; Takei et al., 2004a,b; Kiba et al., 2018; Maeda et al., 2018; Liu

et al., 2022). It has been proposed that nitrate signaling mediated by NLP7 coordinates

CK metabolism and root-to-shoot CK transport to modulate shoot CK responses (Abualia

et al., 2022). In addition, long-distance signals of root-derived *t*Z and *t*ZR differentially

affect leaf growth traits, and their ratios in xylem sap are known to be modulated by

nitrogen supply (Osugi et al., 2017). Together, these findings highlight that regulation of

root-to-shoot CK transport is critical for fine-tuning leaf growth in response to nitrogen

availability in roots.

Our previous study proposed that root-expressed CK receptors are involved

in the regulation of root-to-shoot CK transport. Using grafting, we showed that root-

specific deficiencies of *AHK2* and *AHK3* increased iP- and *t*Z-type CK levels in roots by

suppressing CK degradation and inducing CK biosynthesis. This rise in root CK content

was accompanied by higher *t*Z-type CK levels and stronger expression of CK-inducible

genes in shoots (Monden et al., 2022). Notably, among the three CK receptors, *AHK3*

was identified as one of the top genes co-expressed with *NLP7* (Supplementary Table S1),

suggesting that AHK3 may act in association with nitrate responses via NLP7. In addition,

AHK3 has been reported to exhibit a higher affinity for *t*Z than for iP (Stolz et al., 2011),

implying a preferential role in *t*Z-type CK signaling. Together, these findings suggest that

root-expressed AHK3 may modulate root-to-shoot transport of *t*Z-type CKs in response

to nitrogen availability in roots, although the underlying mechanism and physiological

relevance remain unclear.

In this study, we show that CK perception and degradation in roots, mediated

by AHK3 and CKX4, modulate root-to-shoot transport of *t*Z-type CKs. Growth analyses

of grafted plants demonstrated that root-specific *AHK3* deficiency enlarged leaf blade

area through long-distance signals of root-derived *t*Z-type CK, perceived by shoot-expressed AHK3. Transcriptome, hormonome, and reporter assays further revealed that root-specific *AHK3* deficiency reduced *CKX4* expression in roots, elevating *t*Z-type CK levels in roots and xylem sap and thereby enhancing CK responses in leaves. Transfer experiments manipulating root nitrate levels showed that nitrate-induced leaf growth promotion depends on long-distance signals of root-derived *t*Z-type CK, which is in turn controlled by root-expressed AHK3 and CKX4. We propose a mechanism in which the AHK3–CKX4 module governs root-to-shoot transport of *t*Z-type CK in response to nitrogen availability in roots, thereby fine-tuning leaf size to maintain balanced plant growth.

RESULTS

1. Root-expressed AHK3 modulates shoot and root growth

To identify which root-expressed CK receptor(s) influence shoot and root growth, we grafted Col-0 (WT) scions onto rootstocks derived from WT, single mutant (i.e., *ahk2-5*; *ahk2*, *ahk3-7*; *ahk3*, *cre1-2*; *ahk4*) or double mutant (i.e., *ahk2-5 ahk3-7*; *ahk23*, *ahk2-5 cre1-2*; *ahk24*, *ahk3-7 cre1-2*; *ahk34*) five-day-old seedlings (Riefler et al., 2006). Five days after grafting, we selected plants (scion/rootstock: WT/WT, WT/*ahk2*, WT/*ahk3*, WT/*ahk4*, WT/*ahk23*, WT/*ahk24*, WT/*ahk34*) for further analysis if they showed vigorous growth without adventitious roots. Grafted plants were then cultivated for 7 days on either half-strength MS (1/2 MS) medium or nutrient-rich soil. We measured shoot fresh weight, leaf blade area, and leaf number. Overall, WT/*ahk3*, WT/*ahk23*, and WT/*ahk34* displayed greater shoot growth than WT/WT (Fig. 1A, Supplementary Fig. S1A). Relative to the WT/WT control, these lines showed an increase in shoot fresh weight of 26%, 35%, and 19%, and in leaf blade area of 26%, 32%, and 17%, respectively (Fig. 1, B and C). Under nutrient-rich soil conditions, shoot fresh weight increased by 34%, 25%, and 22%, and leaf blade area by 32%, 21%, and 17% in WT/*ahk3*, WT/*ahk23*, and WT/*ahk34*, respectively, compared with WT/WT (Supplementary Fig. S1, B and C). Leaf number was generally similar across grafted plants in both conditions, although several statistically significant differences were noted (Fig. 1D, Supplementary Fig. S1D). We next monitored primary root length in grafted plants transferred to 1/2 MS medium over a five-day period. Primary root length was comparable at day 0, but WT/*ahk3*,

WT/*ahk23*, and WT/*ahk34* developed significantly shorter roots than WT/WT from day
3 onward (Supplementary Fig. S2). In contrast, WT/*ahk2*, WT/*ahk4*, and WT/*ahk24*
showed no significant differences in shoot or root growth relative to WT/WT (Fig. 1,
Supplementary Figs. S1 and S2). Notably, the relative growth of WT/*ahk3* was
comparable to WT/*ahk23* and WT/*ahk34*, indicating that loss of AHK3 alone is sufficient
to alter overall plant growth. We therefore focused subsequent analyses on WT/*ahk3*,
using WT/WT, WT/*ahk2*, and WT/*ahk4* as comparisons.
168
2. Root-expressed AHK3 modulates root CK levels
Our previous work showed that root-specific deficiencies of *AHK2* and *AHK3* markedly
increased iP- and *t*Z-type CK levels in roots, but the receptor responsible for governing
root CK levels was unclear (Monden et al., 2022). To clarify this, we measured iP- and
*t*Z-type CK concentrations in WT/WT, WT/*ahk2*, WT/*ahk3*, and WT/*ahk4* roots. Root
concentrations of both iP- and *t*Z-type CKs were significantly elevated in WT/*ahk3*
compared with WT/WT; moreover, this was true for active, precursor, and inactive forms
(Fig. 2, Supplementary Fig. S3). In contrast, root iP- and *t*Z-type CKs were largely
unchanged in WT/*ahk2* and WT/*ahk4* relative to WT/WT, except for iP7G and iP9G in
WT/*ahk4* (Fig. 2, Supplementary Fig. S3). These findings indicate that AHK3 expressed
in roots modulates CK homeostasis.
180
3. Root-expressed AHK3 influences the root transcriptome
Despite the fact that *AHK2*, *AHK3*, and *AHK4* are all expressed in roots (Higuchi et al.,
2004), only AHK3 strongly affected root CK levels. This suggests that each CK receptor
may regulate a distinct set of downstream genes. To test this hypothesis, we performed
an RNA-seq analysis of root samples taken from WT/WT, WT/*ahk2*, WT/*ahk3*, and
WT/*ahk4*. All data, including raw read counts, reads per million mapped reads (RPM),
and normalized transcript levels are presented in Supplementary Tables S2–4. First, we
examined differentially expressed genes (DEGs) between the roots of the WT/WT control
and three grafted plants, i.e., WT/*ahk2*, WT/*ahk3*, and WT/*ahk4* (Supplementary Table
S6). An analysis of the DEGs with a fold change of 1.5 or greater revealed that 62 genes
were significantly upregulated in WT/*ahk3* relative to WT/WT roots, whereas 91 genes
were downregulated (Fig. 3A, Supplementary Figs. S4 and S5). Notably, relatively few

DEGs were detectable in the roots of WT/*ahk2* and WT/*ahk4* (Fig. 3A, Supplementary

Figs. S4 and S5). Next, we conducted a Gene Ontology (GO) term enrichment analysis

of the 62 upregulated genes and the 87 downregulated genes (excluding *AHK3*) identified

in WT/*ahk3* roots. This revealed that multiple hormone-related terms were enriched in

both sets (Supplementary Fig. S6). To further examine the extent to which CK receptor

regulates CK signaling-related genes in roots, we analyzed the expression of CK-

inducible and -repressible genes in the roots of grafted plants; to do so we used a curated

gene list (Bhargava et al., 2013). We found that the expression levels of both CK-

inducible and -repressible genes in the roots of WT/*ahk4* were comparable to the levels

detected in WT/WT (Fig. 3B). In contrast, the expression of CK-inducible genes in root

tissue samples was significantly lower in both WT/*ahk2* and WT/*ahk3* relative to the

WT/WT (Fig. 3B). Moreover, CK-repressible genes were significantly increased in

WT/*ahk3* and showed an increasing trend in WT/*ahk2* (Fig. 3B). Notably, the magnitude

of these changes was more pronounced in WT/*ahk3* than in WT/*ahk2* roots, suggesting

that root-expressed AHK3 plays a major role in regulating root CK signaling.

Next, to identify candidate gene(s) that may be responsible for increased CK

levels in WT/*ahk3* roots, we examined whether the DEGs observed in WT/*ahk3* roots

included genes that are known to be involved in CK biosynthesis and inactivation. Of

these genes, only *CKX4*, which is involved in CK degradation, was among the genes that

were downregulated genes in WT/*ahk3* roots relative to the WT/WT (Fig. 3, A and C).

Quantitative RT-PCR (qRT-PCR) analysis confirmed that expression of *CKX4* in roots

was significantly decreased in WT/*ahk3* relative to WT/WT (Fig. 3D). Taken together,

these results suggest that root-expressed AHK3 modulates root CK levels, potentially via

CKX4-mediated CK degradation.

4. Root-expressed AHK3 modulates *t*Z-type CKs in xylem sap

CKs synthesized in roots are transported to shoots mainly via xylem pathway (Ko et al.,

2014; Osugi et al., 2017). Given that we observed relatively high CK levels in WT/*ahk3*

roots, we hypothesized that the CK levels of xylem sap may also be altered. To verify

this, we measured the CK concentrations of xylem sap samples from WT/WT, WT/*ahk2*,

WT/*ahk3*, and WT/*ahk4*. We found that *t*Z, *t*ZR, and *t*ZRPs were relatively more

abundant than inactive forms of *t*Z-type CK and iP-type CKs in WT/WT xylem sap

(Supplementary Fig. S7A). This suggests that there is preferential root-to-shoot transport

of *tZ* and its precursors. Notably, the concentrations of *tZ*-type CKs were significantly

higher in WT/*ahk3* than in WT/WT xylem sap, while the concentrations of iP-type CKs

remained largely unchanged (Fig. 4, Supplementary Fig. S7B). Among the major CK

species detected in xylem sap, the concentration of *tZ* showed a 2.22-fold increase in

WT/*ahk3* relative to WT/WT xylem sap (Fig. 4B). In contrast, the concentrations of *tZR*

and *tZRPs* were both slightly higher in WT/*ahk3* than in WT/WT xylem sap, showing

1.20-fold and 1.37-fold increases, respectively (Fig. 4B). Finally, the concentrations of

iP- and *tZ*-type CKs were largely unchanged in WT/*ahk2* and WT/*ahk4* relative to

WT/WT xylem sap, except for *tZ* and *tZRPs* in WT/*ahk4* (Fig. 4, Supplementary Fig.

S7B). Overall, these results suggest that root-expressed AHK3 modulates the root-to-

shoot transport of *tZ*-type CKs, especially *tZ*.

5. Root-expressed AHK3 systemically influences the shoot transcriptome

Next, to test whether elevated *tZ*-type CK levels in WT/*ahk3* xylem sap affected shoot

CK levels, we measured CK concentrations in WT/WT, WT/*ahk2*, WT/*ahk3*, and

WT/*ahk4* shoots. Unexpectedly, *tZ*-type CK concentrations were not higher in WT/*ahk3*

than in WT/WT shoots, except for *tZ9G* (Supplementary Fig. S8). Similarly, almost all

CK species in WT/*ahk2* and WT/*ahk4* showed no marked differences compared with

WT/WT (Supplementary Fig. S8). We next analyzed whether root-specific CK receptor

deficiency may alter the shoot transcriptomes. To do so, we performed RNA-seq on

WT/WT, WT/*ahk2*, WT/*ahk3*, and WT/*ahk4* shoots. Raw read counts, RPM, and

normalized transcript levels are provided in Supplementary Tables S2, S3, and S5.

Differential expression analysis with a 1.5-fold cutoff revealed 163 genes significantly

upregulated and 35 genes downregulated in WT/*ahk3* relative to WT/WT shoots (Fig. 5A,

Supplementary Table S7). No DEGs were detected in WT/*ahk2* or WT/*ahk4* shoots (Fig.

5A, Supplementary Table S7). Moreover, the number of upregulated genes was about

five times greater than the number of downregulated genes in WT/*ahk3* shoots (Fig. 5A,

Supplementary Table S7). This finding indicates that root-specific *AHK3* deficiency

mainly induces gene expression in shoots.

To gain insight into the functional characteristics of the upregulated set, we

performed GO enrichment analysis using the set of 163 upregulated genes. Enriched

categories included "Translation," "Response to inorganic substance," and "Cellular
nitrogen compound biosynthetic process" (Fig. 5B), indicating activation of genes related
to protein synthesis, nutrient responses, and nitrogen metabolism in WT/*ahk3* shoots. We
then examined whether these transcriptional changes correlated with altered shoot
element concentrations. We found that the shoot concentrations of calcium, sulfur,
manganese, and molybdenum were significantly higher in WT/*ahk3* than in WT/WT
(Supplementary Fig. S9), suggesting a possible association between these nutrients and
altered gene expression in WT/*ahk3* shoots. In contrast, although several nitrogen-related
GO terms were enriched, shoot concentrations of nitrogen and nitrate did not differ
between WT/WT and WT/*ahk3* (Supplementary Fig. S9). Thus, nitrogen status in
WT/*ahk3* shoots did not drive the transcriptional changes observed in protein synthesis
and nitrogen metabolism genes. Because nitrogen is a major protein component and both
nitrate and CK signaling promote expressions of protein synthesis-related genes
(Sakakibara et al., 2006), altered CK response in WT/*ahk3* shoots may underlie these
transcriptional effects.

To further assess CK response, we compared the 163 upregulated genes in
WT/*ahk3* shoots with CK-inducible genes (Bhargava et al., 2013). *Arabidopsis response*
regulator 5 (*ARR5*) and *Glutaredoxin-S6* (*GrxS6*) were among the overlapping genes (Fig.
5A). *ARR5*, one of ten type-A *ARRs*, is a representative CK-responsive gene functioning
as a negative feedback regulator of CK signaling (To et al., 2004). Expression of several
other type-A *ARRs* also tended to increase in WT/*ahk3* relative to WT/WT (Fig. 5C).
Consistent with the RNA-seq results, qRT-PCR confirmed that *ARR5* expression was
significantly higher in WT/*ahk3* shoots than in WT/WT (Fig. 5D). Taken together, these
results indicate that root-expressed AHK3 modulates shoot gene expression, at least
partly through CK signaling.

6. Root-derived CK signals are required for enhanced shoot growth via root-
expressed AHK3
Elevated *t*Z-type CKs in WT/*ahk3* xylem sap suggested that long-distance signals of root-
derived *t*Z-type CK contribute to enhanced shoot growth. To test this, we generated *ahk3*
abcg14 and *ahk3 cyp735a1 cyp735a2* (*ahk3 cypDM*) mutants by traditional crossing
(Supplementary Fig. S10A). Since both *abcg14* and *cypDM* shoots display lack proper

root-derived *t*Z-type CK signaling (Kiba et al., 2013; Ko et al., 2014), these mutants allow assessment of its contribution. Disruption of the respective genes in *ahk3 abcg14* and *ahk3 cypDM* was confirmed by RT-PCR (Supplementary Fig. S10B). Grafted plants (scion/rootstock: WT/WT, WT/*ahk3*, WT/*abcg14*, WT/*ahk3 abcg14*, WT/*cypDM*, WT/*ahk3 cypDM*) were cultivated on 1/2 MS medium, and shoot growth was measured. Shoot fresh weight and leaf blade area were significantly reduced in WT/*abcg14* and WT/*ahk3 abcg14*, while WT/*cypDM* and WT/*ahk3 cypDM* showed no decrease compared with WT/WT (Fig. 6, A and B). These results agree with previous findings that root-specific deficiency of *ABCG14* impairs shoot growth, whereas deficiencies of *CYP735A1* and *CYP735A2* do not (Kiba et al., 2013; Ko et al., 2014). Notably, the increases in shoot fresh weight and leaf blade area seen in WT/*ahk3* compared with WT/WT were largely suppressed by disrupting *ABCG14* and completely abolished by disrupting *CYP735A1* and *CYP735A2* in roots (Fig. 6, A and B). These findings indicate that long-distance signals of root-derived *t*Z-type CKs are required for enhanced shoot growth in WT/*ahk3*.

To determine which CK receptor(s) mediate CK signaling underlying this growth effect, we analyzed grafted plants carrying single or double mutations in *AHK2*, *AHK3*, or *AHK4* specifically in WT/*ahk3* shoots. Grafted plants (scion/rootstock: WT/WT, WT/*ahk3*, *ahk2*/*ahk3*, *ahk3*/*ahk3*, *ahk4*/*ahk3*, *ahk23*/*ahk3*, *ahk24*/*ahk3*, *ahk34*/*ahk3*) were cultivated under 1/2 MS medium, and shoot growth was measured. Increases in shoot fresh weight and leaf blade area observed in WT/*ahk3* relative to WT/WT were similarly evident in *ahk2*/*ahk3*, *ahk4*/*ahk3*, and *ahk24*/*ahk3* (Fig. 6, C and D). In contrast, these enhancements were suppressed in *ahk3*/*ahk3* and *ahk34*/*ahk3*, and markedly reduced in *ahk23*/*ahk3*, where values were significantly lower than WT/WT (Fig. 6, C and D). These results indicate that enhanced shoot growth in WT/*ahk3* depends primarily on CK signaling through shoot-expressed AHK3, with partial contribution from AHK2. Next, to visualize sites of enhanced CK response in WT/*ahk3* shoots, we introduced the synthetic CK reporter *TCSn:GFP* into WT/WT and WT/*ahk3* shoots (Zürcher et al., 2013). Grafted plants (scion/rootstock: *TCSn:GFP*/WT, *TCSn:GFP*/*ahk3*) were grown on 1/2 MS medium. *TCSn:GFP*/*ahk3* exhibited greater shoot fresh weight and leaf blade area than *TCSn:GFP*/WT (Fig. 6E). Notably, GFP fluorescence intensity in true leaf was also significantly higher in *TCSn:GFP*/*ahk3* than in *TCSn:GFP*/WT (Fig.

6F). Finally, to examine how root-specific *AHK3* deficiency influences leaf micro-morphology, we measured epidermal cell size and number in true leaves of WT/WT and WT/*ahk3*. Cell size was comparable between genotypes, but cell number was significantly higher in WT/*ahk3* than in WT/WT (Fig. 6G). These results suggest that root-specific *AHK3* deficiency enhances leaf CK response, promoting epidermal cell division and thereby increasing leaf size.

7. Root-expressed AHK3 modulates leaf size in a nitrate-dependent manner

AHK3 was identified as one of the top genes co-expressed with the nitrate sensor NLP7 (Supplementary Table S1), suggesting that AHK3 may play a role in plant nitrate responses. To test this, we manipulated the nitrate levels of WT/WT and WT/*ahk3* roots. Ten-day-old WT/WT and WT/*ahk3* were first transferred to 2.5 mM ammonium succinate (5 mM NH_4^+) medium and incubated for 48 hours to deplete nitrate (Supplementary Fig. S11A). Grafted plants were then transferred to medium containing 5 mM NH_4^+ supplemented with either 0.5 mM KNO_3 (KNO_3) or 0.5 mM KCl (KCl) and grown for 5 days to assess the presence or absence of nitrate signaling in roots. Nitrate accumulated only in WT/WT and WT/*ahk3* roots grown under KNO_3 (Supplementary Fig. S11B). Moreover, WT/WT and WT/*ahk3* appeared healthy, with comparable shoot chlorophyll concentrations under both conditions (Supplementary Fig. S12), indicating that nitrogen deficiency or ammonium toxicity did not occur. In WT/WT, shoot fresh weight and leaf blade area were found to be significantly greater under KNO_3 than KCl (Fig. 7A), showing that nitrate signaling promotes leaf growth. Notably, shoot fresh weight and leaf blade area were similar between WT/WT and WT/*ahk3* under KCl, but were significantly higher in WT/*ahk3* than in WT/WT under KNO_3 (Fig. 7A).

Next, to test whether nitrate-dependent growth promotion in WT/*ahk3* requires root-derived *t*Z-type CK signals, we analyzed WT/*abcg14* and WT/*ahk3 abcg14* grown under KNO_3 or KCl. Interestingly, the increases in shoot fresh weight and leaf blade area observed in WT/WT and WT/*ahk3* under KNO_3 were strongly suppressed in WT/*abcg14* and WT/*ahk3 abcg14* (Fig. 7B). These results indicate that long-distance signals of root-derived *t*Z-type CKs are required for nitrate-induced leaf growth promotion. Moreover, shoot fresh weight and leaf blade area were comparable between

WT/*abcg14* and WT/*ahk3 abcg14* under KNO$_3$ (Fig. 7B), suggesting that root-derived

*t*Z-type CK signals may contribute to the enhanced leaf growth of WT/*ahk3*.

To further clarify the role of AHK3 in nitrate signaling, we analyzed

expression of CK biosynthesis and inactivation genes in roots 4 hours after transfer to

KNO$_3$ or KCl. Nitrate accumulation was detected only in WT/WT and WT/*ahk3* roots

after KNO$_3$ treatment (Supplementary Fig. S11C). In WT/WT, root expression of *IPT3*,

CYP735A1, *CYP735A2*, and *CKX4* was significantly higher under KNO$_3$ than KCl (Fig.

7C, Supplementary Fig. S13), indicating induction by nitrate signaling. Importantly,

expression of *IPT3*, *CYP735A1*, and *CYP735A2* was similar in WT/WT and WT/*ahk3*

under KNO$_3$, whereas *CKX4* expression was significantly reduced in WT/*ahk3* compared

with WT/WT (Fig. 7C, Supplementary Fig. S13). These results suggest that both nitrate

signals and root-expressed AHK3 are required for maximal *CKX4* induction.

To investigate the role of CKX4 in nitrate signaling, we obtained two

independent T-DNA alleles, *ckx4-1* (SALK_055204) and *ckx4-2* (SALK_132516), from

the Arabidopsis Biological Resource Center. *Ckx4-1* carries an intron insertion (Bartrina

et al., 2011), while *ckx4-2* carries an exon insertion (Supplementary Fig. S14). RT-PCR

confirmed reduced *CKX4* transcript in *ckx4-1* and absence in *ckx4-2* (Fig. 7D), indicating

that *ckx4-1* is a knockdown and *ckx4-2* a knockout. Grafted plants (scion/rootstock:

WT/WT, WT/*ckx4-1*, WT/*ckx4-2*) were cultivated under KNO$_3$ or KCl, and shoot growth

was measured. Under KCl, shoot fresh weight and leaf blade area were similar among

genotypes (Fig. 7E). Under KNO$_3$, however, both traits were significantly greater in

WT/*ckx4-1* and WT/*ckx4-2* compared to WT/WT (Fig. 7E).

To further clarify relationship between AHK3 and CKX4, we generated *ahk3*

ckx4-1 and *ahk3 ckx4-2* mutants by traditional crossing (Supplementary Fig. S15, A and

B). Grafted plants (scion/rootstock: WT/WT, WT/*ahk3*, WT/*ckx4-1*, WT/*ckx4-2*,

WT/*ahk3 ckx4-1*, WT/*ahk3 ckx4-2*) were grown under KNO$_3$ or KCl, and shoot growth

was measured. All genotypes showed comparable shoot growth under KCl

(Supplementary Fig. S15C). Under KNO$_3$, however, shoot fresh weight and leaf blade

area were significantly greater in WT/*ahk3*, WT/*ckx4-1*, WT/*ckx4-2*, WT/*ahk3 ckx4-1*,

and WT/*ahk3 ckx4-2* compared with WT/WT (Supplementary Fig. S15C). These traits

were higher in WT/*ahk3* than in WT/*ckx4-1* and WT/*ckx4-2* (Supplementary Fig. S15C),

indicating that CKX4 acts downstream of AHK3. Importantly, WT/*ahk3 ckx4-1* and

WT/*ahk3 ckx4-2* showed no further growth increase relative to WT/*ahk3* under KNO_3

(Supplementary Fig. S15C). Together, these results suggest that nitrate-induced leaf

growth promotion occurs through a pathway mediated jointly by AHK3 and CKX4.

DISCUSSION

Recent studies have shown that long-distance signals of root-derived *t*Z-type CK

coordinate multiple physiological processes in aerial organs, including leaf growth,

photoperiod stress response, shade avoidance, and flowering time (Osugi et al., 2017;

Landrein et al., 2018; Frank et al., 2020; Gautrat et al., 2024; Bartrina et al., 2025). These

findings highlight that proper regulation of root-to-shoot *t*Z-type CK transport is essential

for plants to adapt to environmental fluctuations.

Based on our findings and prior studies (Miyawaki et al., 2004; Kiba et al.,

2013; Ko et al., 2014; Zhang et al., 2014; Sakakibara, 2021, 2025), we provide new

insight into the specific mechanisms responsible for modulating root-to-shoot *t*Z-type CK

transport, as mediated by CK perception and degradation through an AHK3–CKX4

module. Nitrate signaling promotes *t*Z-type CK biosynthesis in pericycle and phloem

companion cells via IPT3, CYP735A2, and related enzymes. *t*Z-type CKs are then

transported to the extracellular and endoplasmic reticulum (ER) space by transporters

such as ABCG14. Active form *t*Z are perceived by AHK3 on plasma membrane (PM)

and ER membrane. Activation of CK signaling through AHK3 induces expression of

CKX4, predicted to localize in apoplast, where it modulates apoplastic *t*Z-type CK levels

by degradation. These mechanisms likely influence the apoplastic *t*Z-type CK pool and,

consequently, the amount of *t*Z-type CK loaded into xylem (Supplementary Fig. S16). To

further substantiate this model, it will likely be necessary to analyze the spatial expression

patterns of AHK3 and CKX4 at the tissue and cellular level in future studies. Interestingly,

IPT3 expression and *t*Z-type CK levels in roots also respond to other nutrients, including

phosphate, sulfate, and potassium (Hirose et al., 2008; Nam et al., 2012). Thus, AHK3

may modulates *t*Z-type CK levels in response to diverse nutrient signals. Collectively,

the AHK3–CKX4 module could serve as a feedback regulator that attenuates the nutrient-

induced root-to-shoot transport of *t*Z-type CKs, thereby fine-tuning leaf CK response to

maintain balanced plant growth.

CKXs are known to mediate irreversible CK degradation, producing adenine

(or its derivatives) and aldehyde (Schmülling et al., 2003). In *Arabidopsis*, seven CKXs

differ in substrate specificity, spatial and temporal expression, and subcellular

localization, targeting either vacuoles (CKX1, CKX3), ER (CKX1), apoplasts (CKX2,

CKX4, CKX5, CKX6), or the cytosol (CKX7) (Werner et al., 2003; Galuszka et al., 2007;

Frébort et al., 2011; Niemann et al., 2018). Expression of *CKX3*, *CKX4*, *CKX5*, and *CKX6*

is strongly induced by CK treatment (Werner et al., 2006), suggesting sensitivity to

internal CK levels and roles in maintaining homeostasis. Moreover, CKX2 and CKX4

show higher *in vitro* CKX activity than other CKXs, preferentially degrading active form

CKs at neutral or slightly basic pH (Galuszka et al., 2007). Our data showed that root-

specific *AHK3* deficiency greatly reduced *CKX4* expression in roots, without significantly

altering other *CKX*s (Figs. 3 and 7). These findings suggest that root-expressed AHK3

modulates *t*Z degradation by inducing *CKX4* expression in response to the internal *t*Z

level. This interpretation is supported by our observation that *t*Z concentration was

markedly higher in WT/*ahk3* xylem sap than in WT/WT (Fig. 4). Furthermore, Osugi *et*

al. (2017) demonstrated that root-derived *t*ZR controls both leaf number and area,

whereas root-derived *t*Z regulates only leaf area. In WT/*ahk3*, the increase in xylem sap

*t*Z relative to WT/WT was greater than that of *t*ZR (Fig. 4). Thus, the enlarged leaf blade

area in WT/*ahk3*, without a change in leaf number, likely reflects long-distance signaling

by root-derived *t*Z (Figs. 1 and 4).

AHK2, AHK3, and AHK4 localize to the PM and ER membrane, where they

play central roles in CK signal transduction (Antoniadi et al., 2020; Kubiasová et al.,

2020). AHK3 is considered to be primarily localized on the ER membrane and partially

on the PM (Caesar et al., 2011; Wulfetange et al., 2011). Therefore, ER-localized AHK3

may contribute substantially to regulation of root-to-shoot transport of *t*Z-type CKs.

AHK3 is also expressed more strongly in leaves than in roots, and its leaf expression

exceeds that of AHK2 and AHK4, supporting its hypothesized role as a major regulator

of leaf CK response (Higuchi et al., 2004). This hypothesis is consistent with our finding

that leaf growth promotion, driven by increased shoot CK response, was diminished when

AHK3 was specifically disrupted in WT/*ahk3* shoots (Fig. 6). Strikingly, root-expressed

AHK3 exerted a stronger impact on the root transcriptome and on CK levels than root-

expressed AHK2 and AHK4 (Figs. 2 and 3). Taken together, these results suggest that

AHK3 also plays a major role in roots. Previous reports indicate that rosette leaf growth,
CK metabolism, leaf senescence, and stomatal opening depend particularly on AHK3, as
disruption of AHK3, but not other CK receptors, impairs these processes (Riefler et al.,
2006; Kim et al., 2006; Arnaud et al., 2017). The distinct functions of these receptors may
arise from their expression patterns, recognition of different CK species, and specific
interactions with other proteins (Higuchi et al., 2004; Spíchal et al., 2004; Romanov et
al., 2006; Stolz et al., 2011). A key feature of AHK3 is its ligand preference: while AHK2
and AHK4 bind iP and *t*Z with similar affinity, AHK3 shows higher affinity for *t*Z than
iP (Stolz et al., 2011). In addition, AHK3 specifically mediates CK-triggered
phosphorylation of ARR2 (Kim et al., 2006). These differences in ligand preference and
downstream targets may underlie the unique functions of AHK3.

Overall, this study uncovered a comprehensive mechanism by which the CK
receptor AHK3 systemically modulates the shoot growth via exerting control of root-to-
shoot *t*Z-type CK transport. Furthermore, we clarified the role of AHK3 in modulating
leaf size in response to root nitrogen availability, a key environmental factor. Overall,
these findings provide new insight into how plants integrate nutritional cues to coordinate
systemic communication between roots and shoots.

MATERIALS AND METHODS

Plant materials

The *Arabidopsis thaliana* accession Col-0 was used as wild type. T-DNA insertion lines
ahk2-5, *ahk3-7*, *cre1-2*, *ahk2-5 ahk3-7*, *ahk2-5 cre1-2*, *ahk3-7 cre1-2* (Riefler et al., 2006),
abcg14 (Ko et al., 2014), *cyp735a1-2 cyp735a2-2* (Kiba et al., 2013), and *ckx4-1*
(Bartrina et al., 2011) have been described previously. The *ckx4-2* (SALK_132516) line
was obtained from the Arabidopsis Biological Resource Center (https://abrc.osu.edu/).
The transgenic line *TCSn:GFP* (Zürcher et al., 2013) was described previously. Finally,
the homozygous multiple mutants *ahk3-7 abcg14*, *ahk3-7 cyp735a1-2 cyp735a2-2*, *ahk3-*
7 ckx4-1, and *ahk3-7 ckx4-2* were generated by crossing, and genotypes were determined
by genomic PCR using primers listed in Supplementary Table S8.

Growth conditions of grafted plants

Seeds were surface-sterilized and kept in the dark at 4°C for 3 days to break dormancy.

Approximately 120 seeds per dish were sown in rectangular dishes ($140 \times 100 \times 20$ mm^3;

Eiken Chemical Co. Ltd., Tokyo, Japan) containing 50 mL of half-strength MS salts

including vitamins (Code: M0222; Duchefa Biochemie), supplemented with 0.05% (w/v)

MES-KOH (pH 5.7), 1% (w/v) sucrose, and 0.5% (w/v) gellan gum (Fujifilm Wako, VA,

USA). Plants were grown under a 16/8-h light/dark cycle at 22°C with a photosynthetic

photon flux density (PPFD) of 25 μmol m^{-2} s^{-1}. Five-day-old seedlings were cut

perpendicularly at the hypocotyl using an injection needle tip (Code: NN-2613S;

TERUMO, Tokyo, Japan). Excised scions were connected to corresponding rootstocks

using silicone microtubes (inner × outer diameter: φ0.4 × φ0.5 mm; Code: 1-8194-04; AS

ONE, Osaka, Japan). Grafted plants were incubated for 5 days under continuous light at

27°C with a PPFD of 50 μmol m^{-2} s^{-1}. Successfully grafted plants without adventitious

roots were selected, transferred to solid media or soil pots, then grown for further

experiments.

For *in vitro* cultures, grafted plants were grown vertically for five days or

horizontally for seven days under a PPFD of 70 μmol m^{-2} s^{-1} (16/8-h light/dark cycle) at

22°C in 50 mL (vertical) or 30 mL (horizontal) of half-strength MS salts without vitamins

(Code: M0221; Duchefa Biochemie), supplemented with 0.05% (w/v) MES-KOH (pH

5.7), 1% (w/v) sucrose, and 0.8% (w/v) purified agar (Code: 01162-15; Nacalai Tesque,

Kyoto, Japan). To facilitate non-destructive root sampling during horizontal growth,

media surfaces were covered with cellophane sheets (Hachiya et al., 2021).

For soil cultivation, grafted plants were grown for 7 days under a PPFD of

70 μmol m^{-2} s^{-1} (16/8-h light/dark cycle) at 22°C in pots containing ~150 mL of nutrient-

rich soil (Supermix A; Sakata Seed, Kyoto, Japan).

For transfer experiments with varying nitrogen concentrations, grafted plants

were placed on cellophane sheets in round dishes (diameter 90 mm; depth 20 mm; AS

ONE) containing 30 mL of half-strength MS modified basal salt mixture without nitrogen

(M531; PhytoTech Labs, Lenexa, KS, USA), supplemented with 2.5 mM $(NH_4)_2C_4H_4O_4$,

0.05% (w/v) MES-KOH (pH 5.7), 1% (w/v) sucrose, and 0.8% (w/v) purified agar (Code:

01162-15; Nacalai Tesque) (5 mM NH_4^+ medium). Plants were grown for 48 hours under

a PPFD of 70 μmol m^{-2} s^{-1} (16/8-h light/dark cycle) at 22°C, then transferred to 5 mM

NH_4^+ medium supplemented with either 0.5 mM KNO_3 or 0.5 mM KCl and grown for

five days under the same conditions.

Growth analysis of grafted plants

Shoots were weighed on a precision balance (HR-202i; A&D, Tokyo, Japan). Leaf blades

were harvested and scanned at 600 dpi (GT-X980; EPSON, Tokyo, Japan) to measure

true leaf blade area. Roots of vertically grown plants were scanned at 600 dpi to measure

primary root length. To monitor root elongation, root tip positions were marked every 24

hours. Leaf blade area and primary root length were quantified using ImageJ version 1.53t

(US National Institutes of Health, Bethesda, MD, USA).

Xylem sap collection

Grafted plants were cut perpendicularly at the hypocotyl using an injection needle tip

(Code: NN-2613S; TERUMO). A silicone microtube (inner × outer diameter: φ0.5 × φ0.6

mm; Code: 1-8194-05; AS ONE) was attached to the cut end, and plants were incubated

for 2 hours to allow exudation. After incubation, the silicone microtube was detached,

and xylem sap collected in the tube was recovered by centrifugation into a collection tube.

This was then and stored at –80°C until further analysis. We used 12 µL aliquots of sap

from each grafted plant for CK analyses.

Determination of CK species

CK extraction and quantification followed the method of Kojima *et al*. (2009), using an

ultra-performance liquid chromatography–tandem quadrupole mass spectrometry system

(AQUITY UPLC System/XEVO-TQS; Waters Corp., Milford, MA, USA) equipped with

an octadecylsilyl column (ACQUITY UPLC HSS T3, 1.8 µm, 2.1 mm × 100 mm; Waters

Corp.).

Extraction of total RNA

To extract total RNA, frozen plant samples were first homogenized using a Multi-Beads

Shocker and 5 mm zirconia beads. Total RNA was then isolated with the RNeasy Plant

Mini Kit (Qiagen, Hilden, Germany) with all protocols performed as per the

manufacturer's protocol. For RNA-seq library preparation, RNA was further purified by

on-column DNase digestion (Qiagen).

RNA-seq analysis

RNA quality was evaluated using a Qubit RNA IQ Assay Kit with a Qubit 4 Fluorometer

(Thermo Fisher Scientific, Tokyo, Japan). Samples with RNA IQ scores of 9.5–10.0 were

used for RNA-seq. cDNA libraries were prepared with the NEBNext Ultra II Directional

RNA Library Prep Kit (E7760S, New England Biolabs, Ipswich, MA, USA), SPRIselect

(B23317, Beckman Coulter), the NEBNext Poly(A) mRNA Magnetic Isolation Module

(E7490S, New England Biolabs), and NEBNext Multiplex Oligos for Illumina (E7710S,

New England Biolabs), with all steps following the manufacturer's instructions. Libraries

were sequenced on a NextSeq 500 (Illumina, San Diego, CA, USA), and bcl files were

converted to fastq using bcl2fastq (Illumina). Reads were analyzed as described by

Notaguchi *et al*. (2014) and mapped to the *Arabidopsis* reference (TAIR10) using Bowtie

(Langmead et al., 2009) with the options "--all --best --strata." For normalization, read

count data were transformed with rlog and adjusted by subtracting the average expression

for each gene using default settings of iDEP v0.96 (Ge et al., 2018). DEGs were extracted

using edgeR with a false discovery rate cutoff of 0.15 under the default settings of

RNAseqChef v1.1.4 (Etoh et al., 2023). Gene set enrichment was performed using

ShinyGO v0.82 (Ge et al., 2020) using the default parameters. During analysis we focused

on the Biological Process annotation category.

Reverse transcription and quantitative RT-PCR

Reverse transcription was performed with the ReverTraAce qPCR RT Master Mix with

gDNA Remover (Toyobo, Osaka, Japan). For each reaction, 0.5 μg of RNA was used as

input. Synthesized cDNA was diluted five-fold with sterile water and used as template

for qRT-PCR. Reactions were run on a QuantStudio 1 system (Thermo Fisher Scientific)

using a KOD SYBR qPCR Mix (Toyobo). Relative expression was determined using the

comparative Ct method. *TIP41* was used as an internal reference gene. Primer sequences

are listed in Supplementary Table S9.

Determination of nutrient elements

Shoots were harvested and oven-dried at 80°C for 2 days, ground with a spoon, then

stored in a desiccator with silica gel. For carbon and nitrogen quantification, we weighed

portions of dried samples, packed them into tin capsules, and subjected them to

quantitative analysis using a vario MICRO cube elemental analyzer (Elementar

Analysensysteme GmbH, Hanau, Germany). For other elements (e.g., potassium, calcium,

sulfur, phosphorus, magnesium, sodium, manganese, zinc, iron, boron, copper,

molybdenum, and cobalt), weighed portions of dried samples were digested with HNO_3

followed by H_2O_2. Digests were suspended in 1 mL of 0.08 M HNO_3, diluted tenfold with

0.08 M HNO_3, and analyzed using inductively coupled plasma mass spectrometry

(Agilent 7800 ICP-MS; Agilent Technologies, Santa Clara, CA, USA). Here, indium was

used as an internal standard.

Nitrate level determination

Nitrate was extracted using 10 volumes of sterile water at 100°C for 20 min. A 5.0 μL

aliquot of supernatant was mixed with 40 µL of 5% (w/v) salicylic acid in concentrated

sulfuric acid and incubated at room temperature for 20 min. A mock treatment with 40

µL concentrated sulfuric acid was prepared in parallel. Finally, 1 mL of 8% (w/v) NaOH

was added to the mixture, and nitrate concentrations was determined by recording sample

absorbance at 410 nm.

Fluorescence microscopy

Fluorescence imaging was performed using an all-in-one fluorescence microscope (BZ-

X710, KEYENCE, Osaka, Japan) equipped with a 4× objective (CFI Plan Fluor DL

4×/0.13, Nikon, Tokyo, Japan) and a metal halide lamp (No. 91056, KEYENCE). A

narrow-band GFP filter (Ex. 480/20; Em. 510/20; 49020, Chroma, Brattleboro, VT, USA)

was used for GFP detection, and a Cy5 filter (Ex. 620/60; Em. 700/75; 49006, Chroma)

was used for chlorophyll autofluorescence.

Leaf cellular analysis

The fifth true leaf was excised from the shoot and scanned at 600 dpi (GT-X980; EPSON)

to measure total leaf area. The abaxial surface was pressed onto Scotch transparent

adhesive tape (Code: 500-3-15-10P; 3M Japan, Tokyo, Japan) and carefully peeled to

isolate epidermal cells. Epidermal cell images were captured using a fluorescence
microscope (BZ-X710; KEYENCE) equipped with a 20× objective (CFI Plan Apo
20×/0.75; Nikon) and a metal halide lamp (No. 91056, KEYENCE). Thirty epidermal
cells were randomly selected, and their areas were measured using ImageJ v1.53t (US
National Institutes of Health, Bethesda). The mean area of 30 cells was defined as
epidermal cell size. The total epidermal cell number was estimated by dividing the total
leaf area by the mean cell size.
611

612 **Statistical analyses**
613 Data were analyzed using unpaired two-tailed Welch's t-tests and Dunnett's multiple
614 comparisons tests. All statistical analyses were conducted in R v4.3.0 (R Foundation for
615 Statistical Computing, Vienna, Austria).
616

617 **Data Availability Statement**
618 RNA-seq raw data are deposited in ArrayExpress under accession number E-MTAB-
619 15416.
620

621 **Acknowledgements**
622 We thank Dr. Ryo Tabata and Dr. Tomoko Niwa for technical advice, Ms. Maki Saiki for
623 ICP-MS analysis, and Prof. Thomas Schmülling for important input. We also
624 acknowledge the use of ChatGPT, an AI-based language tool, for support in clarifying
625 the manuscript, and Enago (www.enago.jp) for English language review.
626

627 **Author contributions**
628 K.M. and T.H. conceived and designed experiments. K.M., T.S., M.K., Y.T., T.Ka., H.S.,
629 and T.H. performed the experiments and data analysis. T.Ki., T.N., and T.H. supervised
630 the study. K.M. wrote the manuscript with contributions from all authors. All authors
631 approved the final manuscript.
632

633 **Conflicts of interest**
634 The authors declare that they have no competing interests.
635

Funding

This study was supported by grants from JSPS KAKENHI [No. JP24KJ1709 to K.M.,

No. 21H02087 to T.Ka., and Nos. 20K05771 and 23K04978 to T.H.] and from JST

SPRING [No. JPMJSP2155 to K.M.].

REFERENCES

Abualia R, Ötvös K, Novák O, Bouguyon E, Domanegg K, Krapp A, Nacry P, Gojon A,

Lacombe B, Benková E. Molecular framework integrating nitrate sensing in root and

auxin-guided shoot adaptive responses. *Proc Natl Acad Sci U S A.*

2022:119(31):e2122460119. 10.1073/pnas.2122460119

Antoniadi I, Novák O, Gelová Z, Johnson A, Plíhal O, Simerský R, Mik V, Vain T,

Mateo-Bonmatí E, Karady M, et al. Cell-surface receptors enable perception of

extracellular cytokinins. *Nat Commun.* 2020:11(1):4284. 10.1038/s41467-020-17700-9

Arnaud D, Lee S, Takebayashi Y, Choi D, Choi J, Sakakibara H, Hwang I. Cytokinin-

mediated regulation of reactive oxygen species homeostasis modulates stomatal

immunity in Arabidopsis. *Plant Cell.* 2017:29(3):543–559. 10.1105/tpc.16.00583

Bartrina I, Otto E, Strnad M, Werner T, Schmülling T. Cytokinin regulates the activity of

reproductive meristems, flower organ size, ovule formation, and thus seed yield in

Arabidopsis thaliana. *Plant Cell.* 2011:23(1):69–80. 10.1105/tpc.110.079079

Bartrina I, Werner S, Schenke A, Gasperini D, Werner T, Schmülling T. Root-derived

cytokinin regulates Arabidopsis flowering time through components of the age pathway.

Plant Physiol. 2025:198(3). 10.1093/plphys/kiaf204

Bhargava A, Clabaugh I, To JP, Maxwell BB, Chiang YH, Schaller GE, Loraine A,

Kieber JJ. Identification of cytokinin-responsive genes using microarray meta-analysis

and RNA-Seq in Arabidopsis. *Plant Physiol.* 2013:162(1):272–294.

10.1104/pp.113.217026

Caesar K, Thamm AMK, Witthöft J, Elgass K, Huppenberger P, Grefen C, Horak J,

Harter K. Evidence for the localization of the Arabidopsis cytokinin receptors AHK3 and

AHK4 in the endoplasmic reticulum. *J Exp Bot.* 2011:62(15):5571–5580.

10.1093/jxb/err238

Etoh K, Nakao M. A web-based integrative transcriptome analysis, RNAseqChef,

uncovers the cell/tissue type-dependent action of sulforaphane. *J. Biol. Chem.*

2023:299(6):104810. 10.1016/j.jbc.2023.104810

Frank M, Cortleven A, Novák O, Schmülling T. Root-derived trans-zeatin cytokinin

protects Arabidopsis plants against photoperiod stress. *Plant Cell Environ.*

2020:43:2637–2649. 10.1111/pce.13860

Frébort I, Kowalska M, Hluska T, Frébortová J, Galuszka P. Evolution of cytokinin

biosynthesis and degradation. *J Exp Bot.* 2011:62(8):2431–2452. 10.1093/jxb/err004

Galuszka P, Popelková H, Werner T, Frébortová J, Pospíšilová H, Mik V, Köllmer I,

Schmülling T, Frébort I. Biochemical characterization of cytokinin

oxidases/dehydrogenases from Arabidopsis thaliana expressed in Nicotiana tabacum L. *J*

Plant Growth Regul. 2007:26(3):255–267. 10.1007/s00344-007-9008-5

Gautrat P, Buti S, Romanowski A, Lammers M, Matton SEA, Buijs G, Pierik R.

Phytochrome-dependent responsiveness to root-derived cytokinins enables coordinated

elongation responses to combined light and nitrate cues. *Nat Commun.* 2024:15:8489.

10.1038/s41467-024-52828-y

Ge SX, Son EW, Yao R. iDEP: an integrated web application for differential expression

and pathway analysis of RNA-Seq data. *BMC Bioinformatics.* 2018:19:534.

10.1186/s12859-018-2486-6

Ge SX, Jung D, Yao R. ShinyGO: a graphical gene-set enrichment tool for animals and

plants. *Bioinformatics.* 2020:36(8):2628–2629. 10.1093/bioinformatics/btz931

Hachiya T, Oya T, Monden K, Nagae A, Nakagawa T. A cellophane-supported
Arabidopsis culture for seamless transfer between different media is useful for studying
various nitrogen responses. *Soil Sci. Plant Nutr.* 2021:67(3):277–282.
10.1080/00380768.2021.1908094

Higuchi M, Pischke MS, Mähönen AP, Miyawaki K, Hashimoto Y, Seki M, Kobayashi
M, Shinozaki K, Kato T, Tabata S, et al. In planta functions of the Arabidopsis cytokinin
receptor family. *Proc Natl Acad Sci U S A.* 2004:101(23):8821–8826.
10.1073/pnas.0402887101

Hirose N, Takei K, Kuroha T, Kamada-Nobusada T, Hayashi H, Sakakibara H.
Regulation of cytokinin biosynthesis, compartmentalization and translocation. *J Exp Bot.*
2008:59(1):75–83. 10.1093/jxb/erm157

Hošek P, Hoyerová K, Kiran NS, Dobrev PI, Zahajská L, Filepová R, Motyka V, Müller
K, Kamínek M. Distinct metabolism of N-glucosides of isopentenyladenine and trans-
zeatin determines cytokinin metabolic spectrum in Arabidopsis. *New Phytol.*
2020:225(6):2423‑2438. 10.1111/nph.16310

Hutchison CE, Kieber JJ. Cytokinin signaling in Arabidopsis. *Plant Cell.*
2002:14(suppl_1):S47–S59. 10.1105/tpc.010444

Inoue T, Higuchi M, Hashimoto Y, Seki M, Kobayashi M, Kato T, Tabata S, Shinozaki
K, Kakimoto T. Identification of CRE1 as a cytokinin receptor from Arabidopsis. *Nature.*
2001:409(6823):1060–1063. 10.1038/35059117

Jin SH, Ma XM, Kojima M, Sakakibara H, Wang YW, Hou BK. Overexpression of
glucosyltransferase UGT85A1 influences trans-zeatin homeostasis and trans-zeatin
responses likely through O-glucosylation. *Planta.* 2013:237(4):991–999.
10.1007/s00425-012-1818-4

Kiba T, Takei K, Kojima M, Sakakibara H. Side-chain modification of cytokinins
controls shoot growth in Arabidopsis. *Dev Cell.* 2013:27(4):452–461.
10.1016/j.devcel.2013.10.004

Kiba T, Inaba J, Kudo T, Ueda N, Konishi M, Mitsuda N, Takiguchi Y, Kondou Y,
Yoshizumi T, Ohme-Takagi M, et al. Repression of nitrogen starvation responses by
members of the Arabidopsis GARP-type transcription factor NIGT1/HRS1 subfamily.
Plant Cell. 2018:30(4):925–945. 10.1105/tpc.17.00810

Kim HJ, Ryu H, Hong SH, Woo HR, Lim PO, Lee IC, Sheen J, Nam HG, Hwang I.
Cytokinin-mediated control of leaf longevity by AHK3 through phosphorylation of
ARR2 in Arabidopsis. *Proc Natl Acad Sci U S A.* 2006:103(3):814–819.
10.1073/pnas.0505150103

Ko D, Kang J, Kiba T, Park J, Kojima M, Do J, Kim KY, Kwon M, Endler A, Song W-
Y, et al. Arabidopsis ABCG14 is essential for the root-to-shoot translocation of cytokinin.
Proc Natl Acad Sci USA. 2014:111(19):7150–7155. 10.1073/pnas.1321519111

Kojima M, Kamada-Nobusada T, Komatsu H, Takei K, Kuroha T, Mizutani M, Ashikari
M, Ueguchi-Tanaka M, Matsuoka M, Suzuki K, et al. Highly sensitive and high-
throughput analysis of plant hormones using MS-probe modification and liquid
chromatographytandem mass spectrometry: an application for hormone profiling in
Oryza sativa. *Plant Cell Physiol.* 2009:50(7):1201–1214. 10.1093/pcp/pcp057

Kubiasová K, Montesinos JC, Šamajová O, Nisler J, Mik V, Semerádová H, Plíhalová L,
Novák O, Marhavý P, Cavallari N, et al. Cytokinin fluoroprobe reveals multiple sites of
cytokinin perception at plasma membrane and endoplasmic reticulum. *Nat Commun.*
2020:11(1):4285. 10.1038/s41467-020-17949-0

Kurakawa T, Ueda N, Maekawa M, Kobayashi K, Kojima M, Nagato Y, Sakakibara H,
Kyozuka J. Direct control of shoot meristem activity by a cytokinin-activating enzyme.
Nature. 2007:445(7128):652–655. 10.1038/nature05504
764
765 Kuroha T, Tokunaga H, Kojima M, Ueda N, Ishida T, Nagawa S, Fukuda H, Sugimoto
766 K, Sakakibara H. Functional analyses of LONELY GUY cytokinin-activating enzymes
767 reveal the importance of the direct activation pathway in Arabidopsis. *Plant Cell.*
768 2009:21(10):3152–3169. 10.1105/tpc.109.068676
769
770 Landrein B, Formosa-Jordan P, Malivert A, Schuster C, Melnyk CW, Yang W, Turnbull
771 C, Meyerowitz EM, Locke JCW, Jönsson H. Nitrate modulates stem cell dynamics in
772 Arabidopsis shoot meristems through cytokinins. *Proc Natl Acad Sci U S A.*
773 2018:115(6):1382–1387. 10.1073/pnas.1718670115
774
775 Langmead B, Trapnell C, Pop M, Salzberg SL. Ultrafast and memory-efficient alignment
776 of short DNA sequences to the human genome. *Genome Biol.* 2009:10(3):R25.
777 10.1186/gb-2009-10-3-r25
778
779 Liu KH, Liu M, Lin Z, Wang ZF, Chen B, Liu C, Guo A, Konishi M, Yanagisawa S,
780 Wagner G, et al. NIN-like protein 7 transcription factor is a plant nitrate sensor. *Science.*
781 2022:377(6613):1419–1425. 10.1126/science.add1104
782
783 Maeda Y, Konishi M, Kiba T, Sakuraba Y, Sawaki N, Kurai T, Ueda Y, Sakakibara H,
784 Yanagisawa S. A NIGT1-centred transcriptional cascade regulates nitrate signalling and
785 incorporates phosphorus starvation signals in Arabidopsis. *Nat Commun.* 2018:9(1):1376.
786 10.1038/s41467-018-03832-6
787
788 Matsumoto-Kitano M, Kusumoto T, Tarkowski P, Kinoshita-Tsujimura K, Václavíková
789 K, Miyawaki K, Kakimoto T. Cytokinins are central regulators of cambial activity. *Proc*
790 *Natl Acad Sci USA.* 2008:105(50):20027–20031. 10.1073/pnas.0805619105
791

792 Miyawaki K, Matsumoto-Kitano M, Kakimoto T. Expression of cytokinin biosynthetic
793 isopentyltransferase genes in Arabidopsis: tissue specificity and regulation by auxin,
794 cytokinin, and nitrate. *Plant J.* 2004:37(1):128–138. 10.1046/j.1365-313X.2003.01945.x
795

796 Miyawaki K, Tarkowski P, Matsumoto-Kitano M, Kato T, Sato S, Tarkowska D, Tabata
797 S, Sandberg G, Kakimoto T. Roles of Arabidopsis ATP/ADP isopentenyltransferases and
798 tRNA isopentenyltransferases in cytokinin biosynthesis. *Proc Natl Acad Sci USA.*
799 2006:103(44):16598–16603. 10.1073/pnas.0603522103
800

801 Monden K, Kojima M, Takebayashi Y, Suzuki T, Nakagawa T, Sakakibara H, Hachiya
802 T. Root-specific Reduction of Cytokinin Perception Enhances Shoot Growth in
803 Arabidopsis thaliana. *Plant Cell Physiol.* 2022:63(4):484–493. 10.1093/pcp/pcac013
804

805 Müller B, Sheen J. Arabidopsis Cytokinin Signaling Pathway. *Sci. STKE.* 2007:407:cm5.
806 10.1126/stke.4072007cm5
807

808 Nam YJ, Tran LS, Kojima M, Sakakibara H, Nishiyama R, Shin R. Regulatory roles of
809 cytokinins and cytokinin signaling in response to potassium deficiency in Arabidopsis.
810 *PLoS One.* 2012:7(10): e47797. 10.1371/journal.pone.0047797
811

812 Niemann MCE, Weber H, Hluska T, Leonte G, Anderson SM, Novak O, Senes A, Werner
813 T. The cytokinin oxidase/dehydrogenase CKX1 is a membrane-bound protein requiring
814 homooligomerization in the endoplasmic reticulum for its cellular activity. *Plant Physiol.*
815 2018:176(3):2024–2039. 10.1104/pp.17.00925
816

817 Notaguchi M, Higashiyama T, Suzuki T. Identification of mRNAs that move over long
818 distances using an RNA-Seq analysis of Arabidopsis/Nicotiana benthamiana heterografts.
819 *Plant Cell Physiol.* 2015:56(2):311–321. 10.1093/pcp/pcu210
820

Osugi A, Kojima M, Takebayashi Y, Ueda N, Kiba T, Sakakibara H. Systemic transport
of trans-zeatin and its precursor have differing roles in Arabidopsis shoots. *Nat Plants.*
2017:3:17112. 10.1038/nplants.2017.112

Rahayu YS, Walch-Liu P, Neumann G, Römheld V, von Wirén N, Bangerth F. Root-
derived cytokinins as long-distance signals for NO3−-induced stimulation of leaf growth.
J Exp Bot. 2005:56(414):1143–1152. 10.1093/jxb/eri107

Riefler M, Novak O, Strnad M, Schmülling T. Arabidopsis cytokinin receptor mutants
reveal functions in shoot growth, leaf senescence, seed size, germination, root
development, and cytokinin metabolism. *Plant Cell.* 2006:18(1):40–54.
10.1105/tpc.105.037796

Romanov GA, Lomin SN, Schmülling T. Biochemical characteristics and ligand-binding
properties of Arabidopsis cytokinin receptor AHK3 compared to CRE1/AHK4 as
revealed by a direct binding assay. *J Exp Bot.* 2006:57(15):4051–4058.
10.1093/jxb/erl179

Sakakibara H, Suzuki M, Takei K, Deji A, Taniguchi M, Sugiyama T. A response‑
regulator homologue possibly involved in nitrogen signal transduction mediated by
cytokinin in maize. *Plant J.* 1998:14:337‑344. 10.1046/j.1365-313X.1998.00134.x

Sakakibara H, Takei K, Hirose N. Interactions between nitrogen and cytokinin in the
regulation of metabolism and development. *Trends Plant Sci.* 2006:11: 440–448.
10.1016/j.tplants.2006.07.004

Sakakibara H. Cytokinin biosynthesis and transport for systemic nitrogen signaling. *Plant
J.* 2021:105(2):421–430. 10.1111/tpj.15011

Sakakibara H. Five unaddressed questions about cytokinin biosynthesis. *J. Exp. Bot.*
2025:76(7):1941–1949. 10.1093/jxb/erae348

Schmülling T, Werner T, Riefler M, Krupková E, Bartrina y Manns I. Structure and
function of cytokinin oxidase/dehydrogenase genes of maize, rice, Arabidopsis and other
species. *J Plant Res.* 2003:116:241–252. 10.1007/s10265-003-0096-4
856
857 Spichal L, Rakova NY, Riefler M, Mizuno T, Romanov GA, Strnad M, Schmülling T.
858 Two cytokinin receptors of Arabidopsis thaliana, CRE1/AHK4 and AHK3, differ in their
859 ligand specificity in a bacterial assay. *Plant Cell Physiol.* 2004:45(9):1299–1305.
860 10.1093/pcp/pch132
861
862 Stolz A, Riefler M, Lomin SN, Achazi K, Romanov GA, Schmülling T. The specificity
863 of cytokinin signalling in Arabidopsis thaliana is mediated by differing ligand affinities
864 and expression profiles of the receptors. *Plant J.* 2011:67(1):157–168. 10.1111/j.1365-
865 313X.2011.04584.x
866
867 Takei K, Sakakibara H, Sugiyama T. Identification of genes encoding adenylate
868 isopentenyltransferase, a cytokinin biosynthesis enzyme, in Arabidopsis thaliana. *J Biol
869 Chem.* 2001a:276(28):26405–26410. 10.1074/jbc.M102130200
870
871 Takei K, Sakakibara H, Taniguchi M, Sugiyama T. Nitrogen-dependent accumulation of
872 cytokinins in root and the translocation to leaf: Implication of cytokinin species that
873 induces gene expression of maize response regulator. *Plant Cell Physiol.*
874 2001b:42(1):85–93. 10.1093/pcp/pce009
875
876 Takei K, Yamaya T, Sakakibara H. Arabidopsis CYP735A1 and CYP735A2 encode
877 cytokinin hydroxylases that catalyze the biosynthesis of trans-zeatin. *J Biol Chem.*
878 2004a:279(40):41866–41872. 10.1074/jbc.M406337200
879
880 Takei K, Ueda N, Aoki K, Kuromori T, Hirayama T, Shinozaki K, Yamaya T, Sakakibara
881 H. AtIPT3 is a key determinant of nitrate-dependent cytokinin biosynthesis in
882 Arabidopsis. *Plant Cell Physiol.* 2004b:45(8):1053–1062. 10.1093/pcp/pch119
883

884 To JPC, Haberer G, Ferreira FJ, Deruère J, Mason MG, Schaller GE, Alonso JM, Ecker
885 JR, Kieber JJ. Type-A ARRs are partially redundant negative regulators of cytokinin
886 signaling in Arabidopsis. *Plant Cell.* 2004:16(3):658–671. 10.1105/tpc.018978
887

888 Wang J, Ma XM, Kojima M, Sakakibara H, Hou BK. N-Glucosyltransferase UGT76C2
889 is involved in cytokinin homeostasis and cytokinin response in Arabidopsis thaliana.
890 *Plant Cell Physiol*. 2011:52 (12):2200–2213. 10.1093/pcp/pcr152
891

892 Wang J, Ma XM, Kojima M, Sakakibara H, Hou BK. Glucosyltransferase UGT76C1
893 finely modulates cytokinin responses via cytokinin N-glucosylation in Arabidopsis
894 thaliana. *Plant Physiol. Biochem*. 2013:65:9–16. 10.1016/j.plaphy.2013.01.012
895

896 Werner T, Motyka V, Strnad M, Schmülling T. Regulation of plant growth by cytokinin.
897 *Proc. Natl. Acad. Sci. USA*. 2001:98(18):10487–10492. 10.1073/pnas.171304098
898

899 Werner T, Motyka V, Laucou V, Smets R, Van Onckelen H, Schmülling T. Cytokinin-
900 deficient transgenic Arabidopsis plants show multiple developmental alterations
901 indicating opposite functions of cytokinins in the regulation of shoot and root meristem
902 activity. *Plant Cell.* 2003:15(11):2532–2550. 10.1105/tpc.014928
903

904 Werner T, Köllmer I, Bartrina I, Holst K, Schmülling T. New insights into the biology of
905 cytokinin degradation. *Plant Biol.* 2006:8(3):371–381. 10.1055/s-2006-923928
906

907 Wulfetange K, Lomin SN, Romanov GA, Stolz A, Heyl A, Schmülling T. The cytokinin
908 receptors of Arabidopsis are located mainly to the endoplasmic reticulum. *Plant Physiol.*
909 2011:156(4):1808–1818. 10.1104/pp.111.180539
910

911 Zhang K, Novak O, Wei Z, Gou M, Zhang X, Yu Y, Yang H, Cai Y, Strnad M, Liu CJ.
912 Arabidopsis ABCG14 protein controls the acropetal translocation of root-synthesized
913 cytokinins. *Nat Commun.* 2014:5(1):3274. 10.1038/ncomms4274
914

Zhao J, Ding B, Zhu E, Deng X, Zhang M, Zhang P, Wang L, Dai Y, Xiao S, Zhang C,

et al. Phloem unloading via the apoplastic pathway is essential for shoot distribution of

root-synthesized cytokinins. *Plant Physiol.* 2021:186(4):2111–2123.

10.1093/plphys/kiab188

Zhao J, Deng X, Qian J, Liu T, Ju M, Li J, Yang Q, Zhu X, Li W, Liu C-J, et al.

Arabidopsis ABCG14 forms a homodimeric transporter for multiple cytokinins and

mediates long-distance transport of isopentenyladenine-type cytokinins. *Plant Commun.*

2023:4(2):100468. 10.1016/j.xplc.2022.100468

Zürcher E, Tavor-Deslex D, Lituiev D, Enkerli K, Tarr PT, Müller B. A robust and

sensitive synthetic sensor to monitor the transcriptional output of the cytokinin signaling

network in planta. *Plant Physiol.* 2013:161(3):1066–1075. 10.1104/pp.112.211763

FIGURE LEGENDS

Figure 1. Effects of root-specific CK receptor deficiency on shoot growth in medium.

(A) Representative shoots of grafted plants. Scale bars = 5 mm. (B–D) Shoot fresh weight

(B), leaf blade area (C), and leaf number (D) of grafted plants. Data pooled from two

independent grafting experiments are shown as mean ± SD (n = 12). Asterisks indicate

significant differences relative to WT/WT (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$;

Dunnett's test). Numbers on graphs show mean values relative to WT/WT (set = 1).

Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*,

ahk3-7; *a4*, *cre1-2*; *a23*, *ahk2-5 ahk3-7*; *a24*, *ahk2-5 cre1-2*; *a34*, *ahk3-7 cre1-2*.

Figure 2. Effects of root-specific CK receptor deficiency on root CK concentration.

(A–B) Root concentrations of active form iP and *t*Z and their precursors (A), and inactive

form iP- and *t*Z-type CKs (B) in grafted plants. Data are pooled from three independent

grafting experiments and are reported as mean ± SD (n = 3). Asterisks indicate significant

differences relative to WT/WT (**$P < 0.01$, ***$P < 0.001$; Dunnett's test). Grafted plants

are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-*

2; FW, fresh weight; *t*Z, *trans*-zeatin; iP, N^6-(Δ^2-isopentenyl) adenine; iP and their

precursors = sum of iP, iPR, and iPRPs; inactive form iP-type CKs = sum of iP7G and

iP9G; tZ and their precursors = sum of tZ, tZR, and $tZRPs$; inactive form tZ-type CKs =

sum of $tZ7G$, $tZ9G$, $tZOG$, and $tZROG$.

Figure 3. Effects of root-specific CK receptor deficiency on the root transcriptome.
(A) Volcano plots of DEGs between WT/WT and WT/$ahk2$, WT/$ahk3$, or WT/$ahk4$ roots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023) using edgeR, applying a 1.5-fold change threshold and FDR cutoff of 0.15 (all other parameters default). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted p-value (padj). Each dot represents a single gene. Red, blue, and grey indicate upregulated, downregulated, and non-significant genes, respectively. $\log_2$ FC was limited to -4 to 4 and padj to $\leqq 50$. Full-range volcano plots are provided in Supplementary Fig. S4. (B) Mean normalized transcript levels of CK-inducible and CK-repressible genes (mean ± SD). Gene list obtained from Bhargava *et al.* (2013). (C) Transcriptional changes in CK biosynthesis and inactivation genes, visualized as heatmaps with MORPHEUS (https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each graft to WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per row. (D) Root expression of *CKX4* in grafted plants by qRT-PCR. Relative transcript levels were normalized to the WT/WT root mean (set = 1) and $\log_2$-transformed. Data pooled from two independent grafting experiments are shown as mean ± SD (n = 6). Asterisks indicate significant differences relative to WT/WT (**$P < 0.01$, ***$P < 0.001$; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FC, fold change.

Figure 4. Effects of root-specific CK receptor deficiency on CK concentration of xylem sap. (A–B) Xylem sap concentrations of iP- and tZ-type CKs (A) and of tZ, tZR, and $tZRPs$ (B) in grafted plants. Data pooled from three independent grafting experiments are reported as mean ± SD (n = 3). Asterisks indicate significant differences relative to WT/WT (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; *tZ*, *trans*-zeatin; *tZR*, *tZ*-riboside; *tZRPs*, *tZR* 5′-phosphates; iP-type CKs = sum of iPR, iPRPs, and iP7G; tZ-type CKs = sum of tZ, tZR, $tZRPs$, $tZ7G$, $tZ9G$, and $tZOG$.

Figure 5. Effects of root-specific CK receptor deficiency on the shoot transcriptome.
(A) Volcano plots of DEGs between WT/WT and WT/*ahk2*, WT/*ahk3*, or WT/*ahk4* shoots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023) using edgeR, applying a 1.5-fold change threshold and FDR cutoff of 0.15 (default parameters otherwise). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted p-value (padj). Each dot represents one gene; red, blue, and grey indicate upregulated, downregulated, and non-significant genes, respectively. $\log_2$ FC values were limited to -4 to 4 and padj values to $\leqq 5$. (B) Lollipop plots showing GO enrichment analysis from the 163 genes upregulated in WT/*ahk3* shoots, generated with ShinyGO (Ge et al., 2020) using default settings for biological process terms. (C) Heatmaps of transcriptional changes in type-A *ARR* genes, visualized with MORPHEUS (https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each graft to the WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per row. (D) Shoot expression of *ARR5* determined by qRT-PCR. Values were normalized to the WT/WT shoot mean (set = 1) and $\log_2$-transformed. Data pooled from two independent grafting experiments are reported as mean $\pm$ SD (n = 6). Asterisks indicate significant differences relative to WT/WT (*$P < 0.05$; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FC, fold change.

Figure 6. Effects of root-specific *AHK3* deficiency on shoot growth via CK signaling.
(A–E) Shoot fresh weight and leaf blade area in grafted plants: (A) WT/WT, WT/*ahk3*, WT/*abcg14*, and WT/*ahk3 abcg14*; (B) WT/WT, WT/*ahk3*, WT/*cypDM*, and WT/*ahk3 cypDM*; (C) WT/WT, WT/*ahk3*, *ahk2*/*ahk3*, *ahk3*/*ahk3*, and *ahk4*/*ahk3*; (D) WT/WT, WT/*ahk3*, *ahk23*/*ahk3*, *ahk24*/*ahk3*, and *ahk34*/*ahk3*; (E) *TCSn:GFP*/WT and *TCSn:GFP*/*ahk3*. (F) Representative fluorescent image of the adaxial side of the sixth true leaf in grafted plants. Green = GFP; red = autofluorescence. Scale bars = 1 mm. GFP intensity was quantified with ImageJ as the mean signal in the region marked by a white dashed box. (G) Epidermal cell size and number on the abaxial side of the fifth true leaf. Data pooled from two independent grafting experiments are reported as mean $\pm$ SD (n = 12 (A–E), 11 (F), or 6 (G)). Asterisks (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$) indicate significant differences relative to WT/WT by Dunnett's test. Pairwise differences by

Welch's *t*-test are marked with asterisks above horizontal lines. Graph numbers show

mean values relative to WT/WT (set = 1). Grafted plants are denoted as "scion

line/rootstock line." WT, Col-0; *a2, ahk2-5*; *a3, ahk3-7*; *a4, cre1-2*; *a23, ahk2-5 ahk3-7*;

a24, ahk2-5 cre1-2; *a34, ahk3-7 cre1-2*; *g14, abcg14*; *DM, cyp735a1-2 cyp735a2-2*;

a3g14, ahk3-7 abcg14; *a3DM, ahk3-7 cyp735a1-2 cyp735a2-2*; *TCS, TCSn:GFP*; ns, not

significant.

1015

Figure 7. Effects of root-specific *AHK3* or *CKX4* deficiency on shoot growth in

response to nitrate signaling. (A, B, E) Shoot fresh weight and leaf blade area of grafted

plants 5 days after transfer from 5 mM NH_4^+ medium to either 5 mM NH_4^+ + 0.5 mM

KNO_3 or 5 mM NH_4^+ + 0.5 mM KCl: (A) WT/WT and WT/*ahk3*; (B) WT/WT, WT/*ahk3*,

WT/*abcg14*, and WT/*ahk3 abcg14*; (E) WT/WT, WT/*ckx4-1*, and WT/*ckx4-2*. (C) Root

expression of *IPT3, IPT7, CYP735A1, CYP735A2, CKX1, CKX3, CKX4, CKX5, CKX6,*

CKX7, UGT76C1, UGT76C2, and *UGT85A1* 4 hours after transfer. Heatmaps were

generated with MORPHEUS (https://software.broadinstitute.org/morpheus) from $\log_2$

fold change values relative to WT/WT under 5 mM NH_4^+ + 0.5 mM KCl (set = 1). Color

scale: blue (minimum) to yellow (maximum) per row. (D) RT-PCR analysis of *CKX4* in

WT, *ckx4-1*, and *ckx4-2* seedlings grown 7 days on 1/2 MS medium. *TIP41* was the

control. Primers are listed in Supplementary Table S8. Data are shown as mean ± SD with

replicates as follows: (A) n = 18, two replicates; (B) n = 18, three replicates; (C) n = 6,

two replicates; (E) n = 24, four replicates. Asterisks (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$)

mark significant differences relative to WT/WT under 5 mM NH_4^+ + 0.5 mM KCl

(Dunnett's test). Pairwise comparisons used Welch's *t*-tests (A). For multiple treatments

vs. one control, we used Dunnett's tests (B, E). Significant differences are shown by

asterisks above horizontal lines. Graph numbers indicate mean values relative to WT/WT

under 5 mM NH_4^+ + 0.5 mM KCl (set = 1). Grafted plants are denoted as "scion

line/rootstock line." WT, Col-0; *a3, ahk3-7*; *g14, abcg14*; *a3g14, ahk3-7 abcg14*; *c4-1,*

ckx4-1; *c4-2, ckx4-2*; ns, not significant; KCl, 5 mM NH_4^+ + 0.5 mM KCl; KNO_3, 5 mM

NH_4^+ + 0.5 mM KNO_3.

Figure 1. Effects of root-specific CK receptor deficiency on shoot growth in medium. (A) Representative shoots of grafted plants. Scale bars = 5 mm. (B–D) Shoot fresh weight (B), leaf blade area (C), and leaf number (D) of grafted plants. Data pooled from two independent grafting experiments are shown as mean $\pm$ SD (n = 12). Asterisks indicate significant differences relative to WT/WT (*P < 0.05, **P < 0.01, ***P < 0.001; Dunnett's test). Numbers on graphs show mean values relative to WT/WT (set = 1). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; *a23*, *ahk2-5 ahk3-7*; *a24*, *ahk2-5 cre1-2*; *a34*, *ahk3-7 cre1-2*.

Figure 2. Effects of root-specific CK receptor deficiency on root CK concentration. (A–B) Root concentrations of active form iP and *t*Z and their precursors (A), and inactive form iP- and *t*Z-type CKs (B) in grafted plants. Data are pooled from three independent grafting experiments and are reported as mean ± SD (n = 3). Asterisks indicate significant differences relative to WT/WT (**$P < 0.01$, ***$P < 0.001$; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FW, fresh weight; *t*Z, *trans*-zeatin; iP, N^6-(Δ^2-isopentenyl) adenine; iP and their precursors = sum of iP, iPR, and iPRPs; inactive form iP-type CKs = sum of iP7G and iP9G; *t*Z and their precursors = sum of *t*Z, *t*ZR, and *t*ZRPs; inactive form *t*Z-type CKs = sum of *t*Z7G, *t*Z9G, *t*ZOG, and *t*ZROG.

Figure 3. Effects of root-specific CK receptor deficiency on the root transcriptome. (A) Volcano plots of DEGs between WT/WT and WT/*ahk2*, WT/*ahk3*, or WT/*ahk4* roots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023) using edgeR, applying a 1.5-fold change threshold and FDR cutoff of 0.15 (all other parameters default). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted p-value (padj). Each dot represents a single gene. Red, blue, and grey indicate upregulated, downregulated, and non-significant genes, respectively. $\log_2$ FC was limited to –4 to 4 and padj to $\leqq$50. Full-range volcano plots are provided in Supplementary Fig. S4. (B) Mean normalized transcript levels of CK-inducible and CK-repressible genes (mean $\pm$ SD). Gene list obtained from Bhargava *et al.* (2013). (C) Transcriptional changes in CK biosynthesis and inactivation genes, visualized as heatmaps with MORPHEUS (https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each graft to WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per row. (D) Root expression of *CKX4* in grafted plants by qRT-PCR. Relative transcript levels were normalized to the WT/WT root mean (set = 1) and $\log_2$-transformed. Data pooled from two independent grafting experiments are shown as mean $\pm$ SD (n = 6). Asterisks indicate significant differences relative to WT/WT (**$P < 0.01$, ***$P < 0.001$; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FC, fold change.

Figure 4. Effects of root-specific CK receptor deficiency on CK concentration of xylem sap. (A–B) Xylem sap concentrations of iP- and tZ-type CKs (A) and of tZ, tZR, and tZRPs (B) in grafted plants. Data pooled from three independent grafting experiments are reported as mean ± SD (n = 3). Asterisks indicate significant differences relative to WT/WT (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; tZ, *trans*-zeatin; tZR, tZ-riboside; tZRPs, tZ 5′-phosphates; iP-type CKs = sum of iPR, iPRPs, and iP7G; tZ-type CKs = sum of tZ, tZR, tZRPs, tZ7G, tZ9G, and tZOG.

Figure 5. Effects of root-specific CK receptor deficiency on the shoot transcriptome. (A) Volcano plots of DEGs between WT/WT and WT/*ahk2*, WT/*ahk3*, or WT/*ahk4* shoots. DEG analysis was performed with RNAseqChef (Etoh et al., 2023) using edgeR, applying a 1.5-fold change threshold and FDR cutoff of 0.15 (default parameters otherwise). x-axis: $\log_2$ fold change ($\log_2$ FC); y-axis: $-\log_{10}$ adjusted p-value (padj). Each dot represents one gene; red, blue, and grey indicate upregulated, downregulated, and non-significant genes, respectively. (B) Lollipop plots showing GO enrichment analysis from the 163 genes upregulated in WT/*ahk3* shoots, generated with ShinyGO (Ge et al., 2020) using default settings for biological process terms. (C) Heatmaps of transcriptional changes in type-A *ARR* genes, visualized with MORPHEUS (https://software.broadinstitute.org/morpheus) based on $\log_2$-transformed ratios of each graft to the WT/WT mean RPM. Color scale: blue (minimum) to yellow (maximum) per row. (D) Shoot expression of *ARR5* determined by qRT-PCR. Values were normalized to the WT/WT shoot mean (set = 1) and $\log_2$-transformed. Data pooled from two independent grafting experiments are reported as mean ± SD (n = 6). Asterisks indicate significant differences relative to WT/WT (*P < 0.05; Dunnett's test). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; FC, fold change.

Figure 6. Effects of root-specific *AHK3* deficiency on shoot growth via CK signaling. (A–E) Shoot fresh weight and leaf blade area in grafted plants: (A) WT/WT, WT/*ahk3*, WT/*abcg14*, and WT/*ahk3 abcg14*; (B) WT/WT, WT/*ahk3*, WT/*cypDM*, and WT/*ahk3 cypDM*; (C) WT/WT, WT/*ahk3*, *ahk2*/*ahk3*, *ahk3*/*ahk3*, and *ahk4*/*ahk3*; (D) WT/WT, WT/*ahk3*, *ahk23*/*ahk3*, *ahk24*/*ahk3*, and *ahk34*/*ahk3*; (E) *TCSn:GFP*/WT and *TCSn:GFP*/*ahk3*. (F) Representative fluorescent image of the adaxial side of the sixth true leaf in grafted plants. Green = GFP; red = autofluorescence. Scale bars = 1 mm. GFP intensity was quantified with ImageJ as the mean signal in the region marked by a white dashed box. (G) Epidermal cell size and number on the abaxial side of the fifth true leaf. Data pooled from two independent grafting experiments are reported as mean ± SD (n = 12 (A–E), 11 (F), or 6 (G)). Asterisks (*$P < 0.05$, **$P < 0.01$, ***$P < 0.001$) indicate significant differences relative to WT/WT by Dunnett's test. Pairwise differences by Welch's *t*-test are marked with asterisks above horizontal lines. Graph numbers show mean values relative to WT/WT (set = 1). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a2*, *ahk2-5*; *a3*, *ahk3-7*; *a4*, *cre1-2*; *a23*, *ahk2-5 ahk3-7*; *a24*, *ahk2-5 cre1-2*; *a34*, *ahk3-7 cre1-2*; *g14*, *abcg14*; *DM*, *cyp735a1-2 cyp735a2-2*; *a3g14*, *ahk3-7 abcg14*; *a3DM*, *ahk3-7 cyp735a1-2 cyp735a2-2*; *TCS*, *TCSn:GFP*; ns, not significant.

Figure 7. Effects of root-specific *AHK3* or *CKX4* deficiency on shoot growth in response to nitrate signaling. (A, B, E) Shoot fresh weight and leaf blade area of grafted plants 5 days after transfer from 5 mM NH₄⁺ medium to either 5 mM NH₄⁺ + 0.5 mM KNO₃ or 5 mM NH₄⁺ + 0.5 mM KCl: (A) WT/WT and WT/*ahk3*; (B) WT/WT, WT/*ahk3*, WT/*abcg14*, and WT/*ahk3 abcg14*; (E) WT/WT, WT/*ckx4-1*, and WT/*ckx4-2*. (C) Root expression of *IPT3, IPT7, CYP735A1, CYP735A2, CKX1, CKX3, CKX4, CKX5, CKX6, CKX7, UGT76C1, UGT76C2,* and *UGT85A1* 4 hours after transfer. Heatmaps were generated with MORPHEUS (https://software.broadinstitute.org/morpheus) from log₂ fold change values relative to WT/WT under 5 mM NH₄⁺ + 0.5 mM KCl (set = 1). Color scale: blue (minimum) to yellow (maximum) per row. (D) RT-PCR analysis of *CKX4* in WT, *ckx4-1*, and *ckx4-2* seedlings grown 7 days on 1/2 MS medium. *TIP41* was the control. Primers are listed in Supplementary Table S8. Data are shown as mean ± SD with replicates as follows: (A) n = 18, two replicates; (B) n = 18, three replicates; (C) n = 6, two replicates; (E) n = 24, four replicates. Asterisks (*P < 0.05, **P < 0.01, ***P < 0.001) mark significant differences relative to WT/WT under 5 mM NH₄⁺ + 0.5 mM KCl (Dunnett's test). Pairwise comparisons used Welch's *t*-tests (A). For multiple treatments vs. one control, we used Dunnett's tests (B, E). Significant differences are shown by asterisks above horizontal lines. Graph numbers indicate mean values relative to WT/WT under 5 mM NH₄⁺ + 0.5 mM KCl (set = 1). Grafted plants are denoted as "scion line/rootstock line." WT, Col-0; *a3, ahk3-7; g14, abcg14; a3g14, ahk3-7 abcg14; c4-1, ckx4-1; c4-2, ckx4-2;* ns, not significant; KCl, 5 mM NH₄⁺ + 0.5 mM KCl; KNO₃, 5 mM NH₄⁺ + 0.5 mM KNO₃.