Experimental Study of Social Signaling through Delayed Plumage Maturation in a Colony-Nesting Seabird

Molly M. Hill[1,2], Sarah L. Dobney[1,3], Liliane K. Fanburg[1,4], Daniel J. Mennill[1,3], Liam U. Taylor[1,2,4]

[1]Bowdoin Scientific Station, Kent Island, NB, Canada

[2]Department of Ecology and Evolution, Yale University, New Haven, CT, United States

[3]Department of Integrative Biology, University of Windsor, Windsor, ON, Canada

[4]Biology Department, Bowdoin College, Brunswick, ME, United States

Target journal: ANIMAL BEHAVIOUR

Abstract

Delayed development is a widespread evolutionary strategy that can reduce competition

among highly social animals. Many seabirds exhibit delayed plumage maturation, in which

young birds spend years in visually distinct predefinitive plumages before attaining the

definitive plumage of adults. Previous work hypothesized that predefinitive plumages may

function to reduce aggression towards young, predefinitive seabirds at breeding colonies,

an idea known from other lineages of birds as the status signaling hypothesis. We tested

this hypothesis with visual stimulus experiments at a breeding colony of American Herring

Gulls (*Larus smithsonianus*). We presented painted models of four different plumage

classes (first-cycle predefinitive gull plumage, third-cycle predefinitive gull plumage,

definitive gull plumage, and a Canada Goose as a control) and measured the aggressive

responses of breeding adults at their nests. Breeding gulls responded with significantly less

frequent, lower, and slower aggression toward the first-cycle and control plumages

compared to the definitive plumage. There were no significant differences in response

towards the third-cycle plumage. These results offer support for a status signaling

hypothesis, indicating that substantial, mottled brown plumage—as worn by first-cycle

gulls—reduces aggression from breeding adults in colonies or foraging flocks. Future

research can investigate how immature seabirds—including third-cycle gulls—may

combine plumage, posture, and behavior to shift the dynamics of social behavior at

breeding colonies.

Keywords: delayed plumage maturation; delayed maturity; Herring Gull; model

presentation; plumage signals; social signals; visual ornaments.

Introduction

One evolutionary consequence of sociality is a delayed developmental form that reduces competition. For example, orangutans reduce the threat of conflict by suppressing the development of male face flanges (Kralick et al. 2023) while some highly social wrasses delay developing male sexual characteristics until potential competitors are absent (Warner and Swearer 1991). In birds, several lineages have evolved delayed plumage maturation, in which individuals spend multiple years in distinctive predefinitive plumages before acquiring definitive plumages with adult coloration (Hawkins et al. 2012). Predefinitive plumages can act as social signals that reduce aggression towards young birds (Lyon and Montgomerie 1986, Hawkins et al. 2012), such as for Long-tailed Manakins (*Chiroxiphia linearis*) at display leks (McDonald 1993) and Lazuli Buntings (*Passerina amoena*) at breeding territories (Muehter et al. 1997). Through behavioral studies of animals' reactions to different developmental stages, we can better understand the dynamics of animals' lives in complex social environments.

Here, we investigate the social signaling function of delayed plumage maturation in seabirds at breeding colonies. Prolonged delayed plumage maturation, lasting two years or longer, is present in multiple, independent lineages of seabirds including gulls (Charadriiformes: Laridae), albatrosses (Procellariiformes: Diomedeidae), and gannets (Suliformes: Sulidae). Many species in these lineages have predefinitive plumages that are distinctly darker or drabber than definitive adult plumages (Grant 1982, Tickell 2000, Nelson 1978). The ecological, evolutionary, and behavioral properties of delayed plumage maturation in seabirds are not yet clear.

45   Behavioral observations of seabirds have raised a functional hypothesis for delayed

46 plumage maturation in these lineages. In many seabird species, immature birds return to

47 nesting colonies while still in a distinctly predefinitive plumage, despite not breeding for

48 multiple additional years (e.g., Pickering 1989, Nelson 1966). Young birds in predefinitive

49 plumages also occur alongside adult birds in foraging flocks away from breeding colonies

50 (Wakefield et al. 2019). Accordingly, a functional hypothesis is that predefinitive plumages

51 are social signals that reduce the risk of conflict for young seabirds when they start visiting

52 colonies (Taylor 2024) or competing in foraging flocks. This idea corresponds to the status

53 signaling hypothesis first presented in relation to songbirds (Lyon & Montgomerie 1986).

54 Alternatively, delayed plumage maturation may be socially neutral (i.e., it may serve no

55 social signaling function) if it is merely the byproduct of delayed hormonal expression in

56 young, non-breeding birds (Schaedler et al. 2021).

57   We tested the status signaling hypothesis with a model presentation experiment at

58 a breeding colony of American Herring Gulls (*Larus smithsonianus*). We simulated

59 conspecific birds in different plumage stages near the nests of breeding gulls, using plastic

60 models painted to match different predefinitive and definitive gull plumages, as well as a

61 heterospecific control stimulus. We measured the aggressive responses of breeding gulls

62 to presentations of (1) a young, first-cycle predefinitive gull plumage, (2) an adolescent,

63 third-cycle predefinitive gull plumage, (3) a definitive gull plumage, and (4) a Canada Goose

64 as a control. We predicted that nesting gulls would respond most aggressively to the

65 definitive models and less aggressively to the first-cycle, third-cycle, and control models

66 based on the status signaling hypothesis. Alternatively, if predefinitive plumages are

socially neutral, we predicted that breeding gulls would respond with equal aggression

levels to all gull plumage models. This experiment provides an opportunity to investigate

how predefinitive plumages may function as social signals to mediate aggression at

seabird colonies.

Methods

Study species and site

American Herring Gulls (*Larus smithsonianus*) are seabirds that breed in coastal,

urban, and island colonies across much of northern North America. The species exhibits

delayed reproduction and delayed plumage maturation, taking approximately four years to

begin breeding (as described for the closely-related European Herring Gull *Larus*

argentatus; Chabrzyk and Coulson 1976) in definitive adult plumage (Grant 1982).

The definitive plumage in the American Herring Gull has white body feathers with

pale gray wings, black wing tips with white "mirror" patches, a bright yellow bill, and a red

gonydeal spot near the tip of the bill (Fig. 1A). Before reaching this definitive plumage,

young birds spend approximately one year in each of three predefinitive plumage stages:

(1) a mottled brown plumage in the first molt cycle, ~12 months old (Fig. 1B); (2) a highly

variable, mottled brown plumage with a gray saddle in the second molt cycle, ~24 months

old; and (3) a near-definitive plumage featuring gray wings with brown coverts, dark brown

primaries, a dark tail band, and a yellow bill with merged black and red spots near the tip in

the third molt cycle, ~36 months old (Grant 1982, Nisbet et al. 2017, Taylor 2024; Fig. 1C).

We conducted our study at the American Herring Gull breeding colony on Kent

Island in the Bay of Fundy, New Brunswick, Canada (latitude 44.583, longitude -66.756; Fig.

1E). This 80-hectare island consists of a large grass field along with patches of spruce and

deciduous forest. Roughly 6,000 gulls breed on the island each year (Bennett et al. 2017).

Gulls begin arriving to Kent Island from wintering grounds in the eastern and southeastern

United States in the early spring (Gross 1940, Anderson et al. 2019). They establish nesting

territories by early May. The incubation period at the site generally runs from mid-May

through mid-June, during which time birds are highly territorial and defend the areas

around their nests (Tinbergen 1953, Nisbet et al. 2017). Both parents assist with

incubation, with partners alternating between attending the nest and leaving the territory to

forage (Shlepr et al. 2021). Most gulls on Kent Island are breeding birds in definitive

plumage, but a small number of first- and second-cycle birds (<1% of the total population)

along with a larger number of third-cycle birds (~4% of the total population) are regularly

found within the colony as prebreeders (Taylor 2024).

Primary visual stimuli

Our experiment used four visual stimuli as different plumage treatments. Three

were plastic gull models (model: SG 1100; Sport Plast Decoy Co., Medelana, Italy). The

models were hand-painted with acrylic paint to match different American Herring Gull

plumage stages: first-cycle, third-cycle, and definitive (Fig. 1A–C). The control stimulus was

a plastic Canada Goose (*Branta canadensis*; model: AVX-LGCF-4PK; Avian-X Decoys,

Irving, TX, USA), which is a common breeding species on Kent Island that does not regularly

exhibit aggressive interactions with gulls (Fig. 1D). There were three exemplars of each gull

model and two exemplars of the goose model, for a total of 11 visual stimuli (Fig. S1). To

minimize the effects of pseudoreplication, exemplars were randomized across trials.

Secondary acoustic stimuli

We used acoustic stimuli to help draw attention to the visual stimuli. Acoustic

stimuli consisted of three call types presented in an order corresponding to increasing

levels of agitation: *kek* indicating mild alarm, *yeow* indicating moderate alarm, and *trumpet*

indicating territoriality and social aggression (Tinbergen 1953; Nisbet et al. 2017). Each

acoustic stimulus had the following structure: 20.0 s of silence; three repetitions of a

cluster of three *kek* calls, each separated by 3.0 s of silence; 10.0 s of silence; three

repetitions of a *yeow* call, each separated by 3.0 s of silence; 10.0 s of silence; three

repetitions of a *trumpet* call, each separated by 3.0 s of silence (Fig. 1F). We created these

stimuli from recordings we collected on Kent Island in August 2022. Recordings were

collected with an ME62/K6 microphone mounted in a Telinga parabola connected to a

Marantz PMD-661 digital recorder (44.1 kHz sampling rate; 16-bit accuracy; WAV format).

We selected high signal-to-noise ratio recordings of different individual birds, recorded in

different locations throughout the island, producing *kek*, *yeow*, and *trumpet* calls.

We prepared playback stimuli using Adobe Audition using three exemplars for each

of the three call types. *Kek* calls are often uttered in clusters of three notes spaced closely

together (Nisbet et al. 2017). For *kek* calls, we ensured that each recording had three notes,

spaced approximately 0.1 s apart, for a total duration of approximately 0.5 s. For *yeow*

calls, we used individual calls of approximately 0.3 s in duration. For *trumpet* calls, we

used individual calls, each approximately 3.0 s in duration and containing 9 notes. We

filtered background sounds out of each recording using FFT filters and the LASSO tool in

Audition. We then normalized the amplitude of each recording to -1 dB. We created 27

different audio tracks using all possible combinations of the three recordings of each call

type. To minimize pseudoreplication, tracks were randomized across trials. We broadcast

audio using a camouflage JBL Flip 5 speaker at a consistent volume that we determined to

be natural in the field based on aural assessment. Later, using a calibrated Casella CEL-

24X sound level meter, we measured the amplitude of each call type, and found *kek* calls

were broadcast at 72-76 dB SPL, *yeow* calls were 75-77 dB SPL, and *trumpet* calls were 82-

86 dB SPL, measured at 1m distance from the loudspeaker.

Nest selection

We selected nesting territories for each trial in easily accessible field and beach

habitat (Fig. 1E), choosing only nests with a parent in attendance. Most selected nests had

complete, three-egg clutches (113 / 120 nests) while the remainder had two eggs. To

account for potential variation in parental aggression due to differential investment in the

developing clutch (Pierotti and Annett 1994), we selected only nests that were predicted to

be within one week of hatching based on egg floatation (Schreiber 1970). Although some

eggs began hatching across multiple trials at a single nest, no chicks left the nest area

across the days when we presented the stimuli to the nesting gulls.

Experimental procedure

We tested the aggressive response toward gull plumage signals by presenting

models near the subjects' nests, using the four plumage treatments at each nest in a

randomized order. Each trial at a nest was separated by 24–36 hr to minimize disturbance

and habituation effects. With 120 nests and four plumage treatments per nest, the total

number of trials was 480.

For each trial, one observer approached the nest with the visual stimulus and the

loudspeaker. Models were set upright on a freely spinning, plastic spike ~0.75 m from the

nest and the loudspeaker was placed underneath (Fig. S2). That observer started the

playback stimulus and began briskly walking away from the nest, at which point a second

observer started a timer. These two observers helped categorize the behavioral responses

of parents along with the time at which each behavioral response first occurred. Nest

defense behaviors can vary when both parents are present at the nest (Shah et al. 2025), so

we also recorded the number of parents present at each trial (one, or two, or zero if adults

left the area at the start of the trial). Because many trials featured only one parent, we did

not analyze the sex of these unmarked birds using visual comparisons (the sexes are

monochromatic, and although males are larger than females, an observer can only make

such a differentiation if both parents are near one another). There were no cases where

parents from neighboring nests directly interfered with the stimuli. Each trial lasted 2 min,

or until a bird made physical contact with the visual stimulus. To prevent destruction of the

model, we stopped a trial early and removed the stimulus when a bird made physical

contact.

Behavioral scoring

We categorized behavioral responses into five aggression levels (Table 1). Level 0

(Relaxed) was reserved for gulls that exhibited relaxed behaviors such as preening or

incubating. Level 1 (Alarm) included agitation or alarm responses (*e.g.*, *kek* calls, *yeow*

calls, rapid pacing, short fluttering flights). Level 2 (Social Aggression) included territorial or

social conflict displays such as *mew* calls, *trumpet* calls, and grass-pulling displays

directed at the model (Tinbergen 1953). Level 3 (Physical Threat) included lunging or

swooping at the model without making direct contact. Level 4 (Physical Attacks) included

direct contact with the model.

Each trial was assigned a single aggression score, corresponding to the maximum

aggression level observed during that trial. We excluded behavioral responses that were

clearly directed towards the observers or towards neighboring birds, retaining other

responses for that trial. We excluded trials where birds flushed at the start of the trial and

never returned. After these filters, there was a total of 442 trials retained for analysis: 109

trials for the goose control treatment, 110 trials for first-cycle gull plumage, 113 trials for

the third-cycle gull plumage, and 110 trials for the definitive gull plumage.

Data analysis

All analyses were conducted using the *tidyverse* packages in R version

4.4.1(Wickham et al. 2019; R Core Team 2024). First, we used a binomial logistic model to

test whether the rate of non-aggressive responses (maximum aggression levels 0–1) versus

aggressive responses (maximum aggression levels 2–4) differed by plumage treatment. The

model included treatment order (1, 2, 3, or 4) and number of parents present as fixed

effects, and nest ID as a random effect (Fig. S3). We fit the model using the *glmer* function

from the *lme4* package (Bates et al. 2015). The intercept categories were the definitive

plumage treatment, treatment order 1, and a single parent in attendance for all models.

To investigate different levels of aggression in more detail, we also fit an ordinal

mixed effects model. The ordinal model allowed our five aggression levels to be increasing

in relative intensity without assigning them precise quantitative values. As in the previous

model, we included treatment order and number of parents as fixed effects along with nest

ID as a random effect. We fit the model using the *clmm* function from the *ordinal* package

(Christensen 2023).

Finally, we tested for differences in speed of aggressive responses using a linear

mixed effects model. The response variable in this model was the minimum time needed to

reach the maximum aggression level within each trial. We only included trials where there

was an aggressive response (i.e., maximum aggression levels 2–4). In addition to treatment

order (fixed effect), number of parents (fixed effect), and nest ID (random effect), this

model also included aggression level itself (2, 3, or 4) as a fixed effect. We fit the model

using the *lme* function from the *nlme* package (Pinheiro and Bates 2024)

Ethical Note

All experimental procedures were approved by the Bowdoin Scientific Station,

Bowdoin College (Protocol #2023-0008), and Yale University (Protocol #2024-20453). Birds

were not captured or handled for this experiment. We designed the experiment to minimize

the length of exposure of the animals to signals that could be perceived as aggressive.

Results

Nesting American Herring Gulls responded with variation in their rate of aggressive

response to visual stimuli of different gull plumage stages (Table 2; Fig. 2). Across all

plumage treatments, 67% of trials resulted in a non-aggressive response (298 / 442 trials;

maximum aggression levels 0–1) while 33% of trials resulted in an aggressive response (144

/ 442 trials; maximum aggression levels 2–4). Based on binomial mixed effect model

estimates, gulls were 48% less likely to respond aggressively to the first-cycle gull plumage

treatment (p < 0.05) and 86% less likely to respond aggressively to the goose control (p <

0.001) when compared to definitive gull plumage (Table 2). We found no evidence

differences in the rate of aggression towards the third-cycle plumage treatment, nor were

there effects of treatment order or number of parents present during the trial (Table 2).

Similarly, the ordinal model revealed lower aggression towards the first-cycle gull

plumage, along with the goose control, relative to the definitive gull plumage (Table 3; Fig.

3). In terms of the maximum aggression level, across all trials, 21.0% peaked at no

aggression (level 0) while 46.4% peaked at alarm response (level 1), 4.5% at social

aggression (level 2), 12.2% at physical threat (level 3), and 15.8% at physical attack (level 4;

Fig. S4). We found no evidence for relationships between aggression level and third-cycle

gull plumage, treatment order, or number of parents (Table 3, Fig. 3).

When analyzing aggressive responses only (i.e., response levels 2 and higher), gulls

took significantly longer to reach maximum aggression towards the first-cycle gull plumage

(mean response time: 43.9 ± 36.1 s) compared to the definitive gull plumage (36.7 ± 27.7 s;

Table 4; Fig. 4). We found no evidence for effects on timing from the third-cycle gull

plumage (34.9 ± 30.2 s), and a large but statistically insignificant effect from the goose

plumage (50.8 ± 37.6 s), although the power of the this comparison is likely limited by the

small number of aggressive responses towards the goose.

Aside from plumage, gulls reached peak aggression most slowly in the first trial

(47.4 ± 35.2 s), slightly faster in the second (42.8 ± 30.5 s), and much faster at the third

(33.4 ± 31.4 s) and fourth trials (34.3 ± 29.1 s; Table 4; Fig. S5), with the increases for third

and fourth trials being statistically significant. Gulls also responded more slowly when two

parents were present (53.2 ± 34.6 s) compared to one parent (36.1 ± 30.2 s; Table 4).

Response times did not differ based on level of aggression (Table 4).

Discussion

Multiple orders of seabirds have evolved prolonged delayed plumage maturation

(Hawkins et al. 2012), but few studies have investigated the evolutionary significance of

predefinitive plumages in these species. Breeding American Herring Gulls responded

toward first-cycle predefinitive plumage stimuli with less frequent aggression (Fig. 2; Table

2), lower aggression levels (Fig. 3; Table 3), and slower aggressive reaction times (Fig. 4;

Table 4) compared to definitive adult plumage stimuli. Our experiment with American

Herring Gulls supports the hypothesis that substantial brown plumage elements,

resembling those of first-cycle birds (~12 months old), act as social signals that can reduce

the risk of conflict with older, breeding birds (Taylor 2024).

These results are consistent with previous studies of delayed plumage maturation

that find predefinitive plumages decrease aggression toward young birds, such as for

lekking manakins and territorial songbirds (Lyon and Montgomerie 1986; McDonald 1993;

Hawkins et al. 2012; Schaedler et al. 2021). These studies have focused on sexually

dimorphic species, where female predefinitive plumages generally reduce competition

among young males that are beginning to compete for territories or display opportunities.

In contrast, delayed plumage maturation in seabirds is sexually monomorphic, with

predefinitive plumages that are unique to immature birds rather than resembling a

particular sex. This pattern suggests that delayed plumage maturation in seabirds has

evolved in line with their social systems, which typically involve social monogamy with

biparental territoriality and offspring care (Cockburn 2006). A promising opportunity for

future studies of delayed development would involve comparing the dynamics of delayed

plumage maturation across sexually dimorphic lineages (e.g., some songbirds or lekking

birds) versus sexually monomorphic lineages (e.g., some seabirds or large raptors).

In many gull species, predefinitive plumages have mottled brown and black

elements that resemble the natal and juvenile plumages of chicks and recent fledglings.

Thus, our results indicate that immature gulls have co-opted signals from chick plumages,

which may aid in crypsis (Tinbergen 1961, Hill and McGraw 2006) or may possibly help to

reduce aggression toward vulnerable chicks themselves (Pierotti and Murphy 1987, Nelson

1978). A plumage that reduces the risk of aggression may help young birds navigate the

social environment of the breeding colony and allow young animals to develop the

breeding and territorial skills necessary to begin breeding themselves (Taylor 2024). Young

birds of many seabird species return to colonies for years before breeding (Pickering 1966,

Nelson 1978), despite colonies being dense social environments with frequent competition

and physical aggression (Coulson 2002). Delayed plumage maturation, and the reduction

of aggression toward young gulls, may allow young birds to engage with the social contexts

without suffering injury.

Notably, our experiments found no difference in aggression toward the third-cycle

plumage treatment compared to the definitive plumage treatment. Third-cycle gulls are the

most common predefinitive birds in American Herring Gull breeding colonies, whereas

first-cycle and second-cycle birds are less frequent (Taylor 2024). There are at least three

potential explanations for this disparity. First, third-cycle birds may genuinely lack the

plumage signals that reduce aggression in younger predefinitive birds, instead developing

enough definitive traits that their plumages no longer function as signals of immaturity. For

example, third-cycle birds exhibit a red gonydeal spot, which is used in adult signaling to

chicks (Fig. 1A vs. B; Tinbergen and Perdeck 1950; Tinbergen 1953). From this perspective,

predefinitive plumages may not serve as social signals for the third-cycle birds that return

to colonies, but rather function as signals for younger birds that do not return to colonies

and instead navigate other social environments such as foraging flocks or non-breeding

grounds (Ulfstrand 1979, Wakefield et al. 2019). A second possibility is that our approach to

quantifying aggression, using five broad categories, obscured key differences in reaction to

the third-cycle plumage treatment. Our methods focused on obvious signals of aggression

but did not consider postural cues (such as extending or contracting the neck) that play

important roles in gull social interactions (Tinbergen 1961; Stout and Brass 1969, MacLean

and Bonter 2013). Thus, our results may overlook more subtle variations in the response of

breeding gulls towards all our visual stimuli. Finally, a third possibility is that our artificial

stimuli may have failed to represent salient features of third-cycle plumages. For example,

much of the brown plumage retained on third-cycle gulls is visible only on the spread wing

(Fig. S6) and was consequently not visible on our folded-wing models. Thus, gulls may

manipulate the signaling properties of predefinitive plumage through opening their wings or

other postural cues.

The relationships between physical appearance, posture, and behavior are difficult

to disentangle. Indeed, these relationships have confounded our understanding of delayed

plumage maturation in seabirds. Previous observers noted that young gulls with brown

plumages are attacked more frequently by breeding adults, not less (Huntington 1954;

Ulfstrand 1979; Taylor 2024). However, they were unable to distinguish the effects of

plumage and behavior using observational methods alone. Following classic work on visual

signaling (Tinbergen and Perdeck 1950, Jones 1990), posture (Stout and Brass 1969,

Galusha and Stout 1977), and nest defense (Shah et al. 2025) our study suggests that

visual stimulus experiments in gulls can offer new insights into how social aggression

evolves in the complicated context of a breeding colony. Our experimental investigation

supports the idea that predefinitive plumages reduce aggression in the complex,

competitive social context of a breeding colony.

References

Anderson, C. M., Gilchrist, H. G., Ronconi, R. A., Shlepr, K. R., Clark, D. E., Weseloh, D. V.

C., Robertson, G. J., & Mallory, M. L. (2019). Winter home range and habitat selection differs

among breeding populations of herring gulls in eastern North America. *Movement Ecology*,

7, 8. https://doi.org/10.1186/s40462-019-0152-x.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models

using lme4. *Journal of Statistical Software,* 67(1), 1-48.

https://doi.org/10.18637/jss.v067.i01.

Bennett, J., Jamieson, E., Ronconi, R., & Wong, S. (2017). Variability in egg size and

population declines of Herring Gulls in relation to fisheries and climate conditions. *Avian*

Conservation and Ecology, 12(2), 16. https://doi.org/10.5751/ACE-01118-120216.

Chabrzyk, G. & Coulson, J. C. (1976). Survival and recruitment in the herring gull *Larus*

argentatus. *Journal of Animal Ecology,* 45(1), 187-203.

Christensen, R (2023). Ordinal-regression models for ordinal data. R package version

2023.12-4.1, <https://CRAN.R-project.org/package=ordinal>.

Cockburn, A. (2006). Prevalence of different models of parental care in birds. *Proceedings*

of the Royal Society B, 273, 1375-1383. https://doi.org/10.1098/rspb.2005.3458.

Coulson, J. C. (2002). Colonial breeding in seabirds. *Biology of marine birds*, 87-113.

Galusha, J. G. & Stout, J. F. (1977). Aggressive communication by *Larus glaucescens* Part IV:

Experiments on visual communication. *Behavior,* 62(3), 222-235.

Grant, P. J. (1982). *Gulls: A guide to identification*. Academic Press.

Gross, A. O. (1940). The migration of Kent Island herring gulls. *Bird-Banding*, 11(4), 129-155.

https://doi.org/10.2307/4509634.

Hawkins, G. L., Hill, G. E., & Mercadante, A. (2012). Delayed plumage maturation and

delayed reproductive investment in birds. *Biological Reviews*, 87(2), 257-274.

Hill, G. E., & McGraw, K. J. (Eds.) (2006). *Bird coloration* (vol. 2; pp. 201–232). Harvard

University Press.

Huntington, C. (1954). Age discrimination in a breeding colony of the herring gull *Larus*

argentatus. *Acta XI Congressus Internationalis Ornithologici,* 467-469.

Jones, I. L. (1990). Plumage variability functions for status signalling in least auklets.

Animal Behaviour, 39, 967-975.

Kralick, A. E., O'Connell, C. A., Bastian, M. L., Hoke, M. K., Zemel, B. S., Schurr, T. G., &

Tocheri, M. W. (2023). Beyond dimorphism: Body size variation among adult orangutans is

not dichotomous by sex. *Integrative and Comparative Biology*, 63(4), 907-921.

Lyon, B. E. & Montgomerie, R. D. (1986). Delayed plumage maturation in passerine birds:

Reliable signaling by subordinate males? *Evolution*, 40(3), 605-615.

MacLean, S. A. & Bonter, D. N. (2013). The sound of danger: Threat sensitivity to predator

vocalizations, alarm calls, and novelty in gulls. *PLOS One, 8*(12), e82384.

McDonald, D. B. (1993). Delayed plumage maturation and orderly queues for status: A

manakin mannequin experiment. *Ethology,* 94(1): 31-45.

Muehter, V. R. Greene, E., & Ratcliffe, L. (1997). Delayed plumage maturation in Lazuli

buntings: tests of the female mimicry and status signalling hypotheses. *Behavioral Ecology*

and Sociobiology, 41, 281-290.

Nelson, J. B. (1966). The breeding biology of the gannet *Sula bassana* on the Bass Rock,

Scotland. *Ibis*, 108, 584-626.

Nelson, J. B. (1978). *The Gannet*. T. & A. D. Poyser Limited. p.24-25.

Nisbet, I. C. T., Weseloh, D. V., Hebert, C. E., Mallory, M. L., Poole, A. F., Ellis, J. C., Pyle, P. &

Patten, M. A. (2017). Herring gull (Larus argentatus), version 3.0. In *The Birds of North*

America (P. G. Rodewald, Ed.). Cornell Lab of Ornithology, Ithaca, NY, USA.

https://doi.org/10.2173/bna.hergul.03

Pickering, S. P. C. (1989). Attendance patterns and behaviour in relation to experience and

pair-bond formation in the Wandering Albatross *Diomedea exulans* at South Georgia. *Ibis*,

131(2), 183-195. https://onlinelibrary.wiley.com/doi/abs/10.1111/j.1474-

919X.1989.tb02761.x

Pierotti, R. & Annett, C. (1994). Patterns of aggression in gulls: asymmetris and tactics in

different social categories. *The Condor: Ornithological Applications,* 96(3), 590-599.

https://doi.org/10.2307/1369461.

Pierotti, R. & Murphy, E. C. (1987). Intergenerational conflicts in gulls. *Animal Behaviour*,

35(2), 435-444. https://doi.org/10.1016/S0003-3472(87)80268-3.

Pinheiro, J. & Bates, D., R Core Team (2024). _nlme: Linear and Nonlinear Mixed Effects

Models_. R package version 3.1-166, <https://CRAN.R-project.org/package=nlme>.

Schaedler, L. M., Taylor, L. U., Prum, R. O., & Aciães, M. (2021). Constraint and function in

the predefinitive plumages of manakins (Aves: Pipridae). *Integrative and Comparative*

Biology, 61(4), 1363-1377.

Schreiber, R.W. (1970). Breeding Biology of Western Gulls (Larus occidentalis) on San

Nicolas Island, California, 1968. *The Condor*, 72(2), 133-140.

Shah, S. S., Ellms, M. E., Dygean, F., Gunasekera, S., Hart, E., Ivanyi, R. E., Lane, M., Liu, D.,

Pitcher, D., Saucier, A., Schroeder, K. F., Triquet, J., Williams, K. M., Covino, K. M. (2025).

Investigating diurnal effects and joint nest defense behavior in Herring Gulls. *bioRxiv.*

https://doi.org/10.1101/2025.06.13.659635.

Stout, J. F., and Brass, M. E. (1969). Aggressive Communication by *Larus glaucescens:* Part

II. Visual Communication. *Behavior,* 34(1/2), 42-54.

Taylor, L. U. (2024). Young American Herring Gulls (*Larus argentatus* subsp. *Smithsonianus*)

have the opportunity for social development at the breeding colony. *Waterbirds*, 47(2): 1-

12.

Tickell, W. L. N. (2000). *Albatrosses.* Yale University Press.

Tinbergen, N. (1953). *The herring gull's world: a study of the social behavior of birds*.

Frederick A. Praeger, Inc.

Tinbergen, N. (1961). *The Herring Gull's world: Study of the social behavior of birds.* Basic

Books.

Tinbergen, N., & Perdeck, A. C. (1950). On the stimulus situation releasing the begging

response in the newly hatched Herring Gull chick (*Larus Argentatus Argentatus Pont.*).

Behavior, 3(1), 1-39.

Ulfstrand, S. (1979). Age and plumage associated differences of behaviour among

blackheaded gulls *Larus ridibundus*: foraging success, conflict victoriousness and reaction

to disturbance. *Oikos*, 33(2), 160-166.

Wakefield, E. D., Furness, R. W., Lane, J. V., Jeglinski, J. W. E., & Pinder, S. J. (2019).

Immature gannets follow adults in commuting flocks providing a potential mechanism for

social learning. *Journal of Avian Biology*, 50(10). https://doi.org/10.1111/jav.02164.

Warner, R. R., & Swearer, S. E. (1991). Social control of sex change in the bluehead wrase,

Thalassoma bifasciatum (Pisces: Labridae). *Biol. Bull.* 181, 199-204.

Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L.D., François, R., Grolemnd, G.,

Hayes, A., Henry, L., Hester, J., Kuhn, M., Pedersen, T.L., Miller, E., Bache, S.M., Müller, K.,

Ooms, J., Robinson, D., Aeidel, D.P., Spinu, V., Takahashi, K., Vaughan, D., Wilke, C., Woo,

K., & Yutani, H. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, 4(43),

1686.

Figures

Figure 1. Experimental stimuli and study site for an investigation of plumage signals in

American Herring Gulls (*Larus smithsonianus*). Photographs of (A) first-cycle gull plumage

(~12 months old); (B) third-cycle gull plumage (~36 months old); (C) definitive gull plumage

(46+ months old); and (D) Canada Goose (control), including live birds photographed on

Kent Island (left) and photographs of one model of each type. (E) A map of the American

Herring Gull breeding colony on Kent Island, New Brunswick, Canada, showing the location

of nests used in experimental trials. (F) Sound spectrograms showing an example of each

of the calls used in the acoustic stimuli.

Figure 2. American Herring Gulls were less likely to respond aggressively toward first-cycle

gull plumage, and Canada Goose control models, compared to the definitive gull plumage.

Aggression rate was calculated as the percentage of trials reaching a maximum aggression

level of 2–4 for each treatment. Asterisks indicate $p < 0.05$ (*) and $p < 0.001$ (**) compared

to the intercept category of definitive plumage, from a binomial mixed effects model.

Figure 3. American Herring Gulls responded with lower aggression towards the first-cycle gull plumage, and Canada Goose control, compared to the definitive gull plumage. Vertical lines indicate the mean aggression level for each plumage treatment. Asterisks indicate p < 0.05 (*) and p < 0.001 (***) compared to the intercept category of definitive plumage, from an ordinal mixed effects model.

Figure 4. Aggressive American Herring Gulls responded more slowly toward first-cycle gull

plumage compared to definitive gull plumage. Only trials with an aggressive response

score of 2 or higher were included, limiting the sample size for the goose control. Asterisks

indicate p <0.05 (*) compared to the intercept category of definitive plumage, from a linear

mixed effects model.

Table 1. Behavioral aggression levels used for analysis

Level	Category	Description
0	Relaxed	Incubation; preening; standing; regurgitating; eating
1	Alarm	Pacing; rapidly approaching stimulus; fluttering flight; *kek* or *yeow* vocalizations
2	Social aggression	Choking displays; grass-pulling displays; *trumpet* or *mew* vocalizations
3	Physical threat	Lunging or swooping at stimulus
4	Physical attack	Contact with stimulus

456   **Table 2.** Results of binomial mixed effects model for aggressive response rate. Trials were

457   considered non-aggressive if they reached maximum aggression levels 0–1 and aggressive

458   if they reached levels 2–4. Definitive gull plumage treatment is the intercept category.

Model term	Estimate	Odds ratio	S.E.	z	p
(Intercept)	-0.35	0.70	0.30	-1.17	0.24
Plumage – Third-cycle	-0.08	0.92	0.29	-0.28	0.78
Plumage – First-cycle	-0.66	0.52	0.30	-2.20	**0.03**
Plumage – Goose	-1.96	0.14	0.38	-5.20	**<0.001**
Trial order – 2	-0.15	0.86	0.32	-0.46	0.65
Trial order – 3	-0.05	0.95	0.33	-0.15	0.88
Trial order – 4	0.46	1.58	0.32	1.45	0.15
Num. parents – 2	0.15	1.16	0.32	0.47	0.64

460 **Table 3.** Results of ordinal model for maximum aggression level (0, 1, 2, 3, or 4). Definitive

461 gull plumage treatment is the intercept category.

Model term	Estimate	S.E.	z	p
Plumage – Third-cycle	-0.02	0.26	-0.09	0.93
Plumage – First-cycle	-0.57	0.27	-2.16	**0.03**
Plumage – Goose	-1.33	0.27	-4.95	**< 0.001**
Trial order – 2	-0.35	0.26	-1.34	0.18
Trial order – 3	-0.41	0.26	-1.58	0.12
Trial order – 4	0.03	0.26	0.13	0.89
Num. parents – 2	0.43	0.26	1.65	0.10

463   **Table 4**: Results of linear model for time until maximum aggression (s). Definitive gull

464   plumage treatment is the intercept category.

Term	Estimate	S.E.	DF	t	p
(Intercept)	49.22	9.44	85	5.22	0.00
Plumage – Third-cycle	0.92	6.03	49	0.15	0.88
Plumage – First-cycle	14.32	6.79	49	2.11	**0.04**
Plumage – Goose	18.28	9.87	49	1.85	0.07
Trial order – 2	-10.29	7.32	49	-1.41	0.17
Trial order – 3	-16.23	7.20	49	-2.26	**0.03**
Trial order – 4	-20.86	7.03	49	-2.97	**0.01**
Num. parents – 2	18.15	6.73	49	2.70	**0.01**
Max. aggression – 3	-3.50	8.37	49	-0.42	0.68
Max. aggression – 4	-9.73	8.09	49	-1.20	0.23

465