Consequences of dietary amino acid imbalance on growth and gas exchange in black soldier fly (*Hermetia illucens*) larvae

Running header: METABOLIC CONSEQUENCES OF DIETARY AMINO ACID BALANCE IN *H. ILLUCENS* LARVAE

M. Schow-Madsen[1#], K. Jensen[2#]*, M.M. Toribio[1], M.L Schøn[2,3], T.M. Schou[4], J.V. Nørgaard[2] and Johannes Overgaard[1]*

[1] Department of Biology, Zoophysiology, Aarhus University, C.F. Møllers Allé 3, 8000 Aarhus C, Denmark

[2] Department of Animal and Veterinary Sciences, Monogastric Nutrition, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark

[3] Department for Health Sciences and Technology, Sustainable Food Processing Laboratory, ETH Zurich, Schmelzbergstrasse 9, 8092 Zürich, Switzerland

[4] ENORM Biofactory A/S, Hedelundvej 15, 8762 Flemming, Denmark

[#] First authors, *Corresponding authors: kj@anivet.au.dk and johannes.overgaard@bio.au.dk

Abstract: Black soldier fly larvae (*Hermetia illucens* (L.); BSFL) production relies on efficient nutrient conversion into protein-rich biomass to ensure sustainability and economic viability. While the effects of dietary protein content and macronutrient ratios are well described, the metabolic costs of providing an imbalanced essential amino acid (EAA) supply in dietary proteins remain poorly understood. We examined how dietary EAA imbalance affects growth, metabolism, and energy allocation in BSFL. Larvae were reared on a reference diet (chicken feed) and three isonitrogenous, semi-synthetic diets: one balanced diet matching the EAA profile of the reference, and two unbalanced diets selectively deficient in either leucine, phenylalanine, and threonine or isoleucine, lysine, and tryptophan. Using open flow respirometry, daily growth monitoring, and biochemical analysis, we quantified gas exchange, body composition, and the energetic costs of biomass and protein deposition. EAA imbalance delayed development, reduced growth and protein content, and promoted lipid accumulation, while O_2 consumption and CO_2 production were reduced and respiratory exchange ratios exceeded 1.0, indicating elevated lipogenesis. Metabolic costs per unit protein deposited were higher on EAA-deficient diets (up to 3.99 kJ per kJ protein), reflecting inefficient amino acid utilisation. In contrast, the balanced diet supported more efficient protein deposition at a lower energetic cost (2.92 kJ per kJ protein). Our findings demonstrate that suboptimal dietary EAA profiles reduce metabolic efficiency in BSFL. Quantifying the physiological costs of amino acid imbalance provides benchmarks to guide feed formulations in insect production systems to minimise input costs and maximise the nutritional quality and quantity of larval output.

Keywords: essential amino acids, metabolism, physiological costs, protein utilization, respiratory exchange ratio

Orcids: M. Schow-Madsen: 0009-0004-1289-3579; K. Jensen: 0000-0003-0261-3831; M.L. Schøn: 0009-0007-6251-9034; T.M. Schou: 0000-0001-7545-7572; J.V. Nørgaard: 0000-0002-0580-0184; J. Overgaard: 0000-0002-2851-4551

Acknowledgements: This study was supported by the Green Development and Demonstration Programme (GUDP), Projects EntoFeed, 34009-21-1837, and FlyCloud, 34009-20-1652.

42	**Conflict of interest:** The authors declare no conflict of interest.

Introduction

Optimising insect nutrition is crucial in insects produced for food and feed to maximize feed conversion and utilization efficiency to minimize production costs (Oonincx *et al*., 2025). Black soldier fly larvae (*Hermetia illucens* (L.); BSFL) are increasingly applied to support circular waste management as they convert low-value organic side streams into protein- and lipid-rich biomass for animal feed (Liu *et al*., 2019; Purkayastha and Sarkar, 2022; van der Heide *et al*., 2021) and produce organic plant fertiliser from the frass (Lomonaco *et al*., 2024). Rapid growth, high feed conversion efficiency, and tolerance to a wide range of low-quality substrates make BSFL well-suited for large-scale production systems (Liu *et al*., 2019; Siddiqui *et al*., 2024). However, production efficiency - including biomass conversion rate, larval yield, growth rate, survival, and body composition - is strongly modulated by the nutritional quality of the feed (Gold *et al*., 2018; Liu *et al*., 2019). Research on BSFL has explored diverse feedstocks and underutilised organic side streams (Broeckx *et al*., 2021; Gold *et al*., 2018; Oonincx *et al*., 2015), and substantial progress has been made in understanding the effects of dietary protein and carbohydrate content (Barragan-Fonseca *et al*., 2019; Cammack and Tomberlin, 2017; Schøn *et al*., 2025a). As BSFL production scales up, it is increasingly important to evaluate how suboptimal diets affect growth, metabolism, and body composition, to develop cost-effective feeds that optimise larval yield and nutritional value while maintaining economic viability.

Despite increasing attention to BSFL nutrition, the role of micronutrient balance, especially essential amino acids (EAAs), remains insufficiently understood (Tomberlin *et al*., 2023). EAAs are required for protein synthesis and cannot be synthesised *de novo* (Wu *et al*., 2014). In BSFL, few studies have tested individual EAA limitations, showing that deficiencies or imbalances constrain somatic growth and delay development (Berggreen *et al*., 2025; Koethe *et al*., 2022; Spranghers *et al*., 2024). However, the broader metabolic consequences of imbalanced dietary EAA profiles, including effects on energy allocation and protein assimilation, remain unexplored. When EAAs are limiting, excess non-limiting amino acids cannot support protein synthesis and are instead deaminated with the resulting carbon redirected into lipid synthesis (Berggreen *et al*., 2025; Stryer, 1995), potentially increasing the metabolic cost of growth.

Growth is metabolically costly, requiring energy for biomass synthesis, tissue maintenance, and cellular turnover (Eriksen, 2022; Laganaro *et al*., 2021; Schmidt-Nilsen, 1997). It is well-described that the energetic cost of digesting and assimilating protein exceeds that of carbohydrates and lipids (Goodrich *et al*., 2024; Hawkins, 1991; Secor, 2009), and when protein is present but poorly utilisable, a greater proportion of total respiration is allocated to maintenance rather than growth (Laganaro *et al*., 2021). As feed accounts for a major part of the BSFL production costs, efficient conversion of diverse, low-value substrates into high-protein biomass is key to economic viability. High-quality substrates accelerate growth and improve larval protein content and nutrient composition (Barragan-Fonseca *et al*., 2019; Schøn *et al*., 2025a), increasing their nutritional value. In contrast, suboptimal feeds prolong development to pupation and reduce both biomass yield and quality (Barragan-Fonseca *et al*., 2019; Gold *et al*., 2018), though such diets may be cheaper and easier to formulate from low-cost side-stream. Whether the aim is to minimise feed input costs or to maximise the nutritional quality of the larval biomass, a clear understanding of the metabolic costs associated with nutrient assimilation is essential for evaluating feed efficiency. As BSFL are primarily valued for their protein content, reduced protein deposition directly lowers both the nutritional value of the larvae and the economic return. Identifying how suboptimal dietary EAA profiles influence the energetic cost of protein deposition is therefore important to guide feed formulation strategies in large-scale production.

The objectives were to evaluate how amino acid balance affected growth, survival, and body composition, how respiratory activity changes during the last 7 days of production, and to

91 estimate the energetic cost of lipid and protein deposition using oxygen-based calorimetry. This
92 study evaluates the gross cost of growth, defined as the energy expended per unit of biomass
93 gained, providing an integrative measure of metabolic efficiency during growth.

Materials and methods

Experimental animals and general design

Black soldier fly larvae were provided by the commercial BSFL facility ENORM Biofactory A/S
(Flemming, Denmark) and transported 1-hour to the laboratory facilities shortly before each
experiment. The larvae used in the experiments were 5-6 days old with a mass of ~5 mg,
corresponding to the 4^{th}-5^{th} larval stage (Gligorescu *et al.* 2019).

The experiments were designed to examine how dietary deficiencies of EAAs affect
the gross cost of growth in BSFL. To examine this, growth, gas exchange, body composition, and
energy expenditure were compared in BSFL reared on four different diets (see below) for seven
days at 27°C. Each replicate in each diet group contained 600 larvae, housed in 1 L respirometer
chambers with ~340 g of feed. Over four weeks, measurements were performed using a blocked
design, testing one or two replicates per diet weekly, resulting in seven replicates per diet. The
experimental system recorded CO_2 production and O_2 consumption on an hourly basis, and each
day, 15 larvae from each replicate were randomly selected, weighed and then returned to the
chamber. At the end of the experiment, all larvae were weighed, counted, and stored at -20°C for
later analysis of water, crude lipid, and crude protein content.

Experimental diets

This experiment included a reference diet (REF), and three treatment diets - a balanced (B), and two
unbalanced diets (UA and UB). The REF diet consisted solely of commercial chicken feed
(KylleKræs 1, Danish Agro; Table S1), which contained amino acids, minerals, and vitamins to
meet the nutrient requirements of growing chickens (23.0% protein, 65.4% carbohydrates, and 5.2%
lipids). The REF diet was included to ensure that our rearing conditions supported growth and
metabolism comparable to previous BSFL studies using high-productivity feeds (Schøn *et al.*,
2025b). The three treatment diets (B, UA, UB) were formulated to explore the impact of EAA
deficiencies (Table 1). All treatment diets contained 35.1% chicken feed to ensure a base supply of
essential macro- and micronutrients. Compared to the REF diet, the semi-synthetic treatment diets
contained less total protein to promote high protein utilisation and induce a growth-limiting effect
of EAAs, whose relative concentrations were below the levels required for maximal growth and
protein deposition (Berggreen *et al.*, 2025). Importantly, the B diet formulation has previously been
validated for BSFL growth and metabolism in Berggreen *et al.* (2025), confirming its suitability for
controlled nutritional experiments. This chicken-feed fraction in the treatment diets provided
approximately 8.1% protein, 23% carbohydrates, and 1.8% lipids. The remaining 64.9% of the diet
consisted of a carbohydrate mixture (sucrose and dextrin, 1:1), a free amino acid mixture, cellulose,
and other components including Vanderzant vitamin mixture, Wesson's salt mixture, linoleic acid
(80% linoleic acid, 20% oleic acid), cholesterol, and nipagin (1:6:1:1:1). Cellulose served as a
structural element, while nipagin functioned as a preservative. While chicken feed had a protein
content of 20.3% of diet dry mass, the semi-synthetic diets had a protein plus free amino acid
content of 15.2 % of diet dry mass to ensure a high requirement for balanced amino acids for
growth (Berggreen *et al.*, 2025).

Table 1. Ingredient and nutritional composition of the three semi-synthetic diets based on dry mass content. The three diets differed only in the specific free amino acid mix used.

Ingredients (%)	
Chicken feed[1]	35.1
Carbohydrate mix[2]	39.3
Amino acid mix[3]	7.1
Cellulose	14.3
Agar	1.7
Other additives[4]	2.5
Nutrients (%)	
Carbohydrate	78.3
Digestible carbohydrates	61.1
Protein	8.1
Free amino acid mix	7.1
Lipids	1.8
Other components[5]	4.7
Energy (kJ/g)[6]	16.3
Metabolizable energy (kJ/g)	13.6

[1]See Table S1

[2]1:1 sucrose and dextrin

[3]See Table 2

[4]Other additives include Vanderzant vitamin mixture (0.25%), Wessons salt mixture (1.50%), nipagin (0.25%) and the lipids; linoleic acid (0.20%), oleic acid (0.05%) and cholesterol (0.25%).

[5]Micronutrients plus other additives

[6]Energy is calculated using energy contents of 17 kJ/g for carbohydrates, 17 kJ/g for protein, and 37 kJ/g for lipids (Capuano *et al.*, 2018)

In diet B, the amino acid profile was matched to that of REF, enabling a direct comparison of amino acids derived exclusively from chicken feed versus those provided as free crystalline amino acids. Diet UA was selectively deficient in leucine, phenylalanine, and threonine, whereas diet UB was deficient in isoleucine, lysine, and tryptophan (Table 2). These six amino acids each limited BSFL growth significantly when not included in the free amino acid mix (Berggreen et al. 2025), and their deficiencies were here combined in two random groups in the two diets. To maintain a constant amount of free amino acid of 7.1% of the dry diet (Table 1), non-essential amino acids were adjusted to compensate for the removed EAA. All diets were prepared by mixing 320 g of heated agar-water solution with 80 g of the designated dry ingredients. After solidification in the respiration chambers, some water evaporated, yielding 351.6 ± 5.5 g of wet diet (~77% water). Each chamber contained 600 BSFL, corresponding to ~1.71 larvae per gram of wet diet or 0.133 g of dry feed per larva.

Table 2. Calculated amounts of essential and non-essential amino acids per kg dry diet in the chicken feed (REF) diet and the three treatment diets varying in amino acid composition. The B diet contains all amino acids in similar proportions as in chicken feed (Table S1), but at lower concentration. The amino acids in bold stem only from the chicken feed part of the diet, as they were not added in the mix of free amino acids. The total free amino acid mixes for the UA and UB diets were adjusted to 7.1% of the total dry diet (Table 1) by adding more non-essential amino acids.

	Amino acid (g/kg)	Diet			
		REF	**B**	**UA**	**UB**
Essential	Arginine	12.87	9.64	9.64	9.64
	Histidine	5.02	3.76	3.76	3.76
	Isoleucine	9.05	6.78	6.78	**3.61**
	Leucine	15.56	11.65	**6.21**	11.65
	Lysine	11.06	8.28	8.28	**4.41**
	Methionine	4.41	3.3	3.3	3.3
	Phenylalanine	9.79	7.33	**3.91**	7.33
	Threonine	7.49	5.61	**2.99**	5.61
	Tryptophan	2.32	1.74	1.74	**0.93**
	Valine	10.18	7.62	7.62	7.62
Non-essential	Alanine	9.22	6.9	7.82	7.53
	Asparagine	5.30	3.97	4.5	4.33
	Aspartic acid	18.90	14.15	16.03	15.44
	Cysteine	3.23	2.42	2.74	2.64
	Glutamine	5.30	3.97	4.5	4.33
	Glutamic acid	36.53	27.35	30.99	29.84
	Glycine	8.53	6.39	7.23	6.97
	Proline	11.53	8.63	9.78	9.41
	Serine	10.26	7.68	8.7	8.38
	Tyrosine	6.57	4.92	5.57	5.37

Growth analysis

To follow the daily growth, 15 larvae were each day randomly selected and weighed to determine the average mass per replicate, after which they were returned to the feed. At the end of the experiment, surviving larvae were counted and weighed, and mortality was calculated. Daily total larval mass was estimated by multiplying the mean larval mass by the estimated number of live larvae each day, under the assumption that mortality was gradual over the seven‐day period. These estimates facilitated the tracking of daily growth and its relation to daily gas exchange and metabolic rate. The growth progression is also evaluated using a specific growth rate (SGR), calculated as follows:

$$SGR = \frac{\ln(End_{body\,mass}) - \ln(Start_{body\,mass})}{Time_{days}}$$

Water, lipid, and protein content

Following the experimental period, all larvae from each replicate were oven‐dried at 60°C for 3-4 days until a stable mass was reached. Water content was determined by subtracting dry mass from

wet mass, and from each replicate, 40 larvae were subsampled to assess lipid and protein contents.
For lipid extraction, dried larvae were weighed and then washed five times with petroleum ether,
where each wash was followed by a 48 hour soak to ensure thorough lipid dissolution. The
defatted samples were then re dried in the oven at 60°C until they had a stable mass, and crude
lipid content was calculated as the difference between initial and lean dry masses. Finely ground
defatted sub-samples (4-7 mg) underwent nitrogen analysis using a combustion analyser (Na 2000,
Italy). Crude protein content was then estimated as the product of the nitrogen content (%) and 4.76
g protein/g nitrogen, which is an established conversion factor for the protein-to-nitrogen ratio in
BSFL (Janssen *et al*., 2017).

Gas exchange measurements
An open-flow respirometry setup followed the protocol described in Schøn *et al*. (2025a, 2025b)
with custom-built, airtight respiration chambers (15 cm diameter, 6 cm height, 1 L volume) placed
in a temperature-controlled cabinet set at 27 °C. A continuous flow of room air (~600 mL/min)
passed through each chamber, ensuring a stable O_2supply while removing water vapour and CO_2.
Over the 7-day experiment, each chamber lost an average of 28.7 ± 2.5 g per day, mainly due to
evaporation and CO_2 production. To measure gas exchange, a sub-sampler pump (SS-4 Sub-
Sampler Pump, Sable Systems) and an 8-channel multiplexer (RM-8 Respirometry Flow
Multiplexer, Sable Systems) sequentially sampled air (~200 mL/min) from each of the eight
chambers (seven experimental and one empty control). The sampled air passed through a calcium
chloride column to remove water vapour and a syringe filter (Acrodisc syringe filter, 25 mm, 0.2
µm Supor membrane, Pall Corporation) before analysis. The O_2 and CO_2 concentrations were
measured using a differential O_2analyser (FC-2 Differential Oxygen Analyser (Oxzilla), Sable
Systems) and a CO_2 gas analyser (LI-850 Gas Analyzer, LI-COR Biosciences). The multiplexer
cycled between chambers every 7.5 minutes, enabling hourly measurements of CO_2 production and
O_2 consumption throughout the 7-day experiment. The rate of oxygen consumption (VO_2) and
carbon dioxide production (VCO_2) (mL/min) were found by multiplying airflow rate (mL/min)
with the difference in gas concentrations between experimental and control chambers. Thus, total
gas exchange (VO$_2$ and VCO$_2$) was calculated for each replicate across the 7-day growth period.
These values were used to determine the temporal (VCO_2/VO_2) and cumulative (VCO$_2$/VO$_2$)
respiratory exchange ratio (RER).

Data Analysis
Unstable measurements and outliers were removed and interpolated following the data quality
assessment protocol described by Schøn *et al*. (2025b). One-way ANOVA was used to assess
differences in BSFL growth, gas exchange, and body composition across treatment groups,
following confirmation of normality and variance homogeneity. Tukey's Honest Significant
Difference (HSD) test was applied for post hoc comparisons, with significance set at $\alpha = 0.05$. Data
processing and visualisation were conducted in RStudio (version 2024.04.2, R version 4.3.2) (Posit
Team, 2023; R Core Team, 2023).

Results
Growth and body composition
All experimental diets supported larval growth and development with mean survival rates ranging
from 88–99%, although larvae on the UB diet exhibited a marginal but statistically significant
increase in mortality (Table 3). The REF diet resulted in the highest total dry matter yield per
replicate (32.4 g from ~ 600 larvae; Fig. 1A; Table 3). Total larvae yield in the B diet (24.7 g)
was lower than that of larvae reared on the REF diet, but higher than for larvae on the two
unbalanced diets, UA (16.9 g) and UB (12.1 g) (Tukey HSD, $\alpha = 0.05$; insert in Fig. 1A). In

accordance with the higher total growth, the average SGR was higher on the REF and B diets than
in UA and UB (Tukey HSD, α = 0.05; insert in Fig. 1B, Table 3).

Figure 1. Dietary effects on growth of black soldier fly larvae during a 7-day rearing period. A) Temporal
changes in average individual larval mass (mg wet mass), with insert depicting final total larval biomass (dry
mass) per treatment. B) Daily estimates of specific growth rate (SGR), based on the change in estimated
larval wet mass in a replicate between consecutive days. The B) insert shows an average SGR over the full 7-
day period for each treatment. Bold lines represent treatment means and semi-transparent lines indicate
individual replicates (n = 7). Insert bar graphs show means ± SE, where distinct letters denote statistically
significant differences among treatments (one-way ANOVA with Tukey's HSD, α = 0.05).

Analysis of larval body composition revealed pronounced diet-dependent differences in lipid and
protein content of the larvae after the 7-day rearing period. Crude protein content in the dry mass
was highest in larvae reared on the REF diet (34.9%), intermediate in larvae reared on the B diet
(29.0%), and lowest in larvae reared on the two unbalanced diets, UA (24.7%) and
UB (23.4%) (Table 3). Conversely, crude lipid content was lowest in larvae from the REF diet
(10.7%), more than twice as high in larvae from the B diet (26.1%), and almost four times as high
in larvae from the unbalanced diets, UB (37.4%) and UA (40.5%) (Table 3).

Gas exchange rates (V̇O₂ and V̇CO₂)
Oxygen consumption rate ($\dot{V}O_2$) and carbon dioxide production rate ($\dot{V}CO_2$) were measured
hourly over the 7-day experimental period. All treatment groups were initially characterised by
parallel increases in $\dot{V}O_2$ and $\dot{V}CO_2$ (Fig. 2) with distinct peaks over the experiment, but the
absolute rates and temporal developments of gas exchange differed markedly among treatments
(Fig. 2). Larvae reared on the REF diet exhibited $\dot{V}O_2$ peaks around day 3.5 (4.5 mL O_2/min) and
5.5 (10 mL O_2/min) followed by a marked reduction in gas exchange rate on day 6 and 7. Larvae
on the B diet followed a similar pattern, with slightly delayed peaks (day 4 and day 6.5) and more
modest absolute gas exchange rates. The two mixed, unbalanced diets showed even lower gas
exchange rates with further deductions in temporal development and peak amplitudes. Thus, in the

two unbalanced treatments, gas exchange peaked around day 5 (1.69 mL O₂/min) with a second rise emerging near the end of the experiment on day 7. When integrating the oxygen consumption over the entire period (insert in Fig. 2A), the total VO_2 was higher on the REF diet (34.5 L O₂) compared to the B diet (20.7 L O₂). Likewise, the total VO_2 in the B diet was higher than both in UA (15.9 L O₂) and UB (10.9 L O₂) (Tukey HSD, α = 0.05).

Table 3: Survival, growth, body composition, and gas exchange of black soldier fly larvae reared on diets with variable amino acid profiles (Table 2).

	Chicken feed (REF)	Balanced (B)	Unbalanced A (UA)	Unbalanced B (UB)
Growth performance				
Survival rate (%)	99.0 ± 1.5% [a]	93.9 ± 4.0% [a]	92.7 ± 3.9% [a]	88.1 ± 7.6% [b]
Larvae dry mass (DM, g per replicate)	32.4 ± 1.0 g [a]	24.7 ± 3.9 g [b]	16.9 ± 3.1 g [c]	12.1 ± 4.0 g [c]
Specific growth rate (per day)	0.50 ± 0.06 [a]	0.47 ± 0.05 [a]	0.39 ± 0.04 [b]	0.36 ± 0.04 [b]
Larvae dry mass (% DM)	30.8 ± 1.1% [a]	31.2 ± 1.4% [a]	33.5 ± 1.6% [b]	33.3 ± 1.1% [b]
Larvae crude protein (% of DM)	34.9 ± 1.5% [a]	29.0 ± 1.7% [b]	24.7 ± 1.3% [c]	23.4 ± 1.4% [c]
Larvae crude lipid (% of DM)	10.7 ± 2.5 % [a]	26.1 ± 2.9 % [b]	40.5 ± 3.1% [c]	37.4 ± 3.4% [c]
Gas exchange measurements				
VCO₂ (L, total for 7 days)	34.5 ± 2.4 L [a]	20.7 ± 3.1 L [b]	15.9 ± 3.2 L [c]	10.9 ± 3.7 L [d]
VO₂ (L, total for 7 days)	33.1 ± 2.9 L [a]	16.4 ± 2.6 L [b]	11.6 ± 2.0 L [c]	8.1 ± 2.7 L [c]
RER	1.04 ± 0.05 [a]	1.27 ± 0.09 [b]	1.38 ± 0.12 [b]	1.35 ± 0.08 [b]

RER = respiratory exchange ratio. RER is calculated from total VCO₂/VO₂. Values are mean ± standard error. Different letters indicate significant differences (one-way ANOVA followed by Tukey HSD test, α = 0.05, n = 7). DM = dry matter, V = volume.

Figure 2. Dietary effects on gas exchange in black soldier fly larvae (BSFL) during a 7-day rearing period.
Temporal trajectory of A) oxygen consumption rates ($\dot{V}O_2$, mL min⁻¹) and B) carbon dioxide production
rates ($\dot{V}CO_2$, mL min⁻¹), with insert summarising total gas exchange (VO_2 and VCO_2, L) across the full
experimental duration for each diet treatment. Bold lines represent treatment means and semi-transparent
lines indicate individual replicates (n=7). The insert bar graphs show means ± SE, where distinct letters
denote statistically significant differences among treatments (one-way ANOVA with Tukey's HSD, α = 0.05).

In the present study, all diet treatments were initially characterised by a RER close to 1.5, which
declined gradually over the 7-day period. The gradual decrease in RER was faster and more
pronounced in the REF diet (Fig. 3) and there was a tendency for the mixed balanced diet (B) to
have slightly lower RER than the two mixed unbalanced diets (UA and UB). When cumulative gas
exchange was integrated over the entire 7-day growth period, we found that larvae on the REF diet
had a lower RER (1.04). The B diet had a cumulative RER of 1.27 which was slightly lower, but not
significantly different from the RER values of the two unbalanced diets (UA: 1.38; UB: 1.34)
(Tukey HSD, α = 0.05; Fig. 3 insert).

Figure 3. Respiratory exchange ratio (RER; $V CO_2/V O_2$) for black soldier fly larvae (BSFL) during a 7-day rearing period plotted over time. Insert bar showing RER derived from cumulative O_2 consumption and CO_2 production per treatment (VCO_2/VO_2). Bold lines represent treatment means and semi-transparent lines indicate individual replicates (n = 7). The insert bar graph shows means ± SE, where distinct letters denote statistically significant differences among treatments (one-way ANOVA with Tukey's HSD, α = 0.05).

Energetic cost of growth and protein assimilation

Integrating growth, body composition, and metabolic rate allows evaluation of the total energetic cost of growth by relating oxygen consumption to energy stored in biomass, with protein and lipid deposition expressed as energy equivalents. When analysing all treatment diets, we found a strong correlation ($R^2 = 0.84$) between the total larval dry mass and total O_2 consumption across individual replicates. Thus, on average each g of dry biomass assimilation was associated with the use of $1.1 L O_2$ for aerobic metabolism (Fig. 4A). When this analysis was performed for each treatment, we found the apparent "gross cost of growth" to be higher for larvae on the REF diet (1.03 L/g), while larvae on the three treatment diets all had lower cost of growth around 0.68 L/g (Tukey HSD, α = 0.05; insert in Fig. 4A).

To evaluate a more direct ratio of metabolic energy expenditure to assimilated energy, we converted the cumulative O_2 consumption to metabolic energy using a standard oxy-calorific coefficient of 20.5 kJ L O_2^{-1} (Schmidt-Nilsen, 1997). Likewise, we calculated assimilated energy from the increment of lipid and protein by multiplying the measured fractions by their respective energy densities (37 kJ g^{-1} for lipid and 17 kJ g^{-1} for protein) (Schmidt-Nilsen, 1997). As seen in Fig. 4B, this analysis revealed marked differences in the energetic cost of energy assimilation among treatments and accordingly the general association between energy expenditure and energy assimilation had a weak correlation among all replicates ($R^2 = 0.18$). When analysed within replicates, we found that the costs of energy assimilation were more than twice as high in the REF treatment compared to the 3 mixed diets (Fig. 4B, insert). Accordingly, larvae fed on the REF diet expended 2.17 kJ per kJ lipid and protein assimilated. In contrast, larvae fed on the mixed diets showed no significant differences, with the balanced diet yielding an expenditure of 0.96 kJ, and the unbalanced diets 0.75 kJ (UA) and 0.81 kJ (UB) per kJ lipid and protein assimilated (Fig. 4B, insert) (Tukey HSD, α = 0.05).

Restricting the analysis to consider only the cost of protein assimilation we found a strong and

general correlation (Fig 4C; R^2 = 0.94) with a slope of 3.48 kJ energy expended per kJ protein
assimilated. Despite the strong general correlation, the energetic cost of protein assimilation also
differed among diets (Fig. 4C, insert). Larvae fed the mixed balanced diet (B) showed the lowest
cost (2.92 kJ), while REF (3.64 kJ) and UA (3.71 kJ) showed intermediate costs, and UB was the
least efficient, with a cost of 3.99 kJ per kJ protein assimilated) (Tukey HSD, α = 0.05).

Figure 4. Metabolic costs of growth during a 7-day rearing period in black soldier fly larvae. A) Linear
relationship between total oxygen consumption (L) and final larval dry biomass (g). B) and C) Energy
expenditure of the system, calculated by multiplying VVO$_2$ by the caloric equivalent of oxygen (20.5 kJ
L^{-1}), plotted against assimilated energy in the biomass (kJ), represented as gain in lipid plus protein (B) or
protein alone (C). Bold points represent treatment means and semi-transparent points show individual
replicates (n = 7). Black lines representing linear regression models for all replicates (n = 28). The insert bar
graphs show means ± standard error for the ratio between the expended energy (Exp.) and the assimilated
energy (Ass.), where distinct letters denote statistically significant differences among treatments (one-way
ANOVA with Tukey's HSD, α = 0.05).

Discussion

By combining respirometry with chemical analysis, we quantified how suboptimal dietary EAA profiles affects growth and metabolic expense in BSFL. The primary aim was to compare a semi-synthetic diet with balanced EAA composition with two other semi-synthetic diets with the same macronutrient composition, but with two different unbalanced EAA profiles.

Growth, body composition, and survival

Survival was slightly but significantly reduced on the UB diet compared to the REF diet (Table 3), indicating that nutritional imbalance can compromise viability. The BSFL generally tolerate wide ranges of dietary protein (Berggreen *et al.*, 2025; Oonincx *et al.*, 2015; Schneider *et al.*, 2025; Schøn *et al.*, 2025a), and high survival is also reported under isonitrogenous diets with imbalanced EAA profiles (Berggreen *et al.*, 2025). Nonetheless, severe EAA deficiencies or excesses can elevate mortality in BSFL (Koethe *et al.*, 2022; Spranghers *et al.*, 2024), as also seen in medflies (Davis, 1975).

We found that unbalanced EAA diets markedly reduced growth of BSFL compared to the balanced diet (Table 3, Fig.1). Biomass accumulation is known to depend on the total dietary protein content, and optimal macronutrient ratios have been widely studied (Barragan-Fonseca *et al.*, 2019; Berggreen *et al.*, 2025; Broeckx *et al.*, 2021, Schøn *et al.* 2025, Eggink *et al.* 2023). However, micronutrient requirements, particularly requirements for individual EAAs, remain poorly defined for BSFL (Tomberlin *et al.*, 2023). In the present study, combined restriction of three EAAs resulted in impaired growth, in accordance with recent findings showing that growth was also suppressed when individual EAAs were deficient (Berggreen *et al.*, 2025). In contrast, Lemme and Klüber (2024) reported no effects on growth following individual EAA reductions. Such discrepancies may reflect masked treatment effects due to uncontrolled EAA or micronutrient levels in the basal diets.

Larvae reared on unbalanced diets were characterised by lower fractional protein content and higher fractional lipid content than those fed on the B or REF diet (Table 3). This is consistent with effects of diets deficient in only a single EAA (Berggreen *et al.*, 2025) and is likely reflecting a metabolic prioritisation of the growing larvae. Specifically, protein synthesis is prioritised over fat storage (Lee, 2007), especially in early life stages (Talal *et al.*, 2024), when amino acids are adequate and balanced (Lee, 2007). However, when the amino acid composition is unbalanced, excess non-limiting amino acids are likely deaminated and stored as lipids (Stryer, 1995), which may explain why larvae on unbalanced diets show the highest fractional lipid content. Lipid accumulation in larvae on unbalanced diets may also result from compensatory feeding under nutrient limitation (Raubenheimer and Simpson, 1997) leading to excess intake of protein and carbohydrate subsequently converted to lipid. Similar mechanisms are reported in other insects (Berner *et al.*, 2005; Lee, 2007) and have been proposed for BSFL (Banks *et al.*, 2014). This is further supported by Pimentel *et al.* (2017) who reported that, in BSFL, protein-poor diets downregulate genes involved in protein storage and promote lipid accumulation via de novo lipogenesis.

Gas exchange

Gas exchange patterns revealed distinct metabolic responses to dietary amino acid balance (Fig. 2). Both oxygen consumption (VO_2) and carbon dioxide production (VCO_2) were reduced in larvae fed the UA and UB diets (Fig 2 inserts) with suppressed respiratory peaks and delayed development relative to the B diet (Fig 2). Similar patterns were reported by Schøn *et al.* (2025a) under low protein-to-carbohydrate conditions, which may also be considered as low-quality diets. The reduced gas exchange likely reflects lower ATP demand due to suppressed anabolic activity associated with protein synthesis and tissue growth. In BSFL, respiratory metabolism scales with growth

performance (Parodi *et al.*, 2020), and the observed metabolic downregulation is consistent with
patterns seen under other growth-limiting conditions, such as low dietary water content in *Manduca*
sexta (Van't Hof and Martin, 1989) or thermal stress in BSFL (Schøn *et al.*, 2025b). The
developmental delay may reflect prolonged larval instars, where moulting phases following
respiratory peaks are associated with reduced feeding (Gligorescu *et al.*, 2019). The substantial
elevated gas exchange in REF fed larvae, also reported by Schøn *et al.* (2025a), likely reflects
greater availability of growth-promoting micronutrients in the REF diet that enhance anabolic
processes.
382   When calculated from the total CO_2 and O_2 exchange over the 7-day period, the RER
exceeded 1.00 across treatments. The highest RER values were observed in larvae fed the B, UA,
and UB diets, with an apparent positive association with crude lipid content across treatments
(Table 3). While CO_2 was not used to estimate cost of growth (see below), it was used to calculate
RER, to get information about the combined impact of metabolic respiratory activity and the
surplus CO_2 produced during lipogenesis (Talal *et al.*, 2021). Accordingly, elevated RER values
suggest that CO_2 production arose not only from catabolism (RER: 0.7–1.0, depending on lipid or
carbohydrate utilisation, respectively (Schmidt-Nilsen, 1997)), but also from anabolic lipogenesis
(Schøn *et al.*, 2025b; Talal *et al.*, 2021) and possibly microbial metabolism during early
development (Parodi et al., 2020). This supports the suggestion by Berggreen *et al.* (2025) that
impaired protein accretion redirects excess amino acids toward lipid synthesis (Berggreen *et al.*,
2025), increasing lipid synthesis and elevating CO_2 production relative to O_2 consumption.
However, this finding was expected, as RER values above 1 are common among fast-growing
animals (Jakobsen and Thorbekt, 1993; Talal *et al.*, 2021) including BSFL (Parodi *et al.*, 2020;
Schøn *et al.*, 2025a, 2025b).

Energetic cost of growth and protein assimilation
The analysis of gross cost of growth was evaluated by relating total oxygen consumption to biomass
gain. Cumulative oxygen consumption over the 7-day period was converted to metabolic energy
using a standard oxycaloric coefficient, assuming metabolism primarily fuelled by mixed nutrients.
Biomass gain was expressed as the energy stored in newly deposited protein and lipid, calculated
from their respective energy densities. This method integrates metabolic costs as cumulative output
relative to biochemical gain and avoids potential biases associated with CO_2-based estimates,
which, as described above, can be confounded by lipogenesis and microbial activity (Gold *et al.*,
2018; Schøn *et al.*, 2025b; Talal *et al.*, 2021). The gross cost of growth reflects the combined
demands of tissue synthesis, maintenance metabolism, and microbial processes (Eriksen, 2022;
Laganaro *et al.*, 2021; Fuhrmann *et al.*, 2025), and while our whole-system measurements do not
separate these components, relative differences in cost per unit growth still allow meaningful
comparisons of substrate efficiency and nutrient limitations.
410   As an indirect measurement of energy metabolism, the total oxygen consumption
scaled linearly with final larval biomass (Fig. 4A), reflecting the well-established link between total
feed assimilation (growth) and energy consumption in insects (Laganaro *et al.*, 2021; Von
Bertalanffy, 1957). The gross cost of growth (volume oxygen used per gram dry body mass gained)
remained consistent across all three treatment diets (Fig. 4A, insert) demonstrating that the
metabolic energy used to produce a g of biomass on these three diets did not differ. Larvae fed on
the REF diet achieved the greatest biomass accumulation but also exhibited higher gross cost of
growth per gram of biomass assimilated than those on semi-synthetic diets. Notably, larvae on the
REF diet showed a distinct temporal growth pattern, characterised by ~1.5 days of biomass loss and
reduced metabolic rate near the end of the experiment, presumably because the average larvae had
reached the non-feeding prepupal larval stage. Oxygen consumed during this period was primarily
allocated to catabolic or maintenance processes rather than net growth, potentially overestimating

the true cost of tissue synthesis. A more precise definition of larval stages would allow for a more
detailed analysis of growth costs. Nonetheless, the three mixed diet groups remained in the growth
phase throughout the 7-day experiment and are therefore evaluated on more equal terms.
To understand why mass-specific metabolic costs remained unchanged between the
balanced (B) and unbalanced diets (UA and UB), we assessed energy expenditure relative to total
energy gain (lipid + protein energy) (Fig. 4B). Again, energetic costs appeared consistent across
diets, likely due to differences in biomass composition. Protein synthesis is more metabolically
demanding than lipid accumulation (Hawkins, 1991; Secor, 2009) and likely constitutes a larger
component of energy expenditure in larvae fed the B and REF diets, which best supported protein
growth. In contrast, larvae on EAA-deficient diets face suboptimal conditions for growth due to
limited substrates for protein synthesis. As a result, assimilated nutrients are redirected toward lipid
storage, and the relative allocation of energy shifts so that growth is constrained while maintenance
becomes relatively more dominant (Laganaro *et al*., 2021). Although total protein synthesis is
reduced in larvae fed unbalanced diets, the metabolic costs of lipid storage, maintenance, and
deamination of excess amino acids (Hawkins, 1991) may raise energy expenditure to levels
comparable to protein-synthesising larvae.
The relatively high energetic cost of protein assimilation is supported by comparison
of the weak association between total energy gain (lipid + protein) and energy expenditure (Fig.
4B), in contrast to the stronger correlation observed for protein gain versus energy expenditure (Fig.
4C). These results suggest that, while lipid accumulation incurs some energetic cost, protein
synthesis is the primary driver of energy expenditure during growth. This aligns with the prediction
of Eriksen (2022), who modelled BSFL energy budgets using CO_2-based fluxes and partitioned
respiration into maintenance and biosynthesis. Despite methodological differences, both approaches
converge on the conclusion that protein synthesis is substantially more energetically costly than
lipid deposition and represents the dominant metabolic driver in BSFL. Thus, when focusing solely
on protein gain, the energetic inefficiency of EAA-deficient diets became evident. Larvae on UA
and UB expended more energy per kJ of protein deposited than those on the B diet (Fig. 4C), and
the same was observed for larvae fed the REF diet, likely due to the inclusion of energy expenditure
during post-feeding stages. As diets with unbalanced EAA composition caused lower protein
deposition and higher energetic cost of protein growth, our results indicate that suboptimal dietary
profiles may constrain the efficiency of insect production systems, highlighting the need for further
research to improve the sustainability and economic viability of insect feeds.

Conclusions

This study demonstrates that unbalanced EAA profiles impair growth performance and metabolic
efficiency in BSFL. By integrating growth, body composition, and gas exchange data, we show that
suboptimal EAA profiles limit protein synthesis, promote lipid accumulation, and increase the gross
energetic cost of amino acid metabolism and protein deposition. Metabolic costs per unit of protein
deposited were substantially higher under EAA-deficient diets (up to 3.99 kJ per kJ protein) than
under the balanced diet (2.92 kJ per kJ protein), reflecting inefficient amino acid utilization.
Whether unbalanced diets are disadvantageous from a feed-cost perspective depends on the trade-
off between reduced larval growth and protein assimilation versus the potentially lower cost of low-
quality feed ingredients. The use of crystalline amino acids offers precise control of dietary
composition, but their viability depends on whether they support sufficient biomass yield and
quality. Our results suggest that protein from complex feed sources (e.g. chicken feed), may be
more efficiently assimilated than crystalline amino acids, even when EAA profiles are matched,
which is also previously reported (Berggreen *et al*., 2025). By quantifying the physiological costs of
EAA imbalance, this study provides benchmarks to support the evaluation of suboptimal feeds in

BSFL production. Further research should assess whether targeted EAA supplementation can improve protein yield and energy efficiency on low-quality substrates, thereby contributing to the development of nutritionally and economically optimized insect feeds.

References

Banks, I.J., Gibson, W.T. and Cameron, M.M., 2014. Growth rates of black soldier fly larvae fed on fresh human faeces and their implication for improving sanitation. Tropical Medicine and International Health 19: 14–22. https://doi.org/10.1111/tmi.12228

Barragan-Fonseca, K.B., Gort, G., Dicke, M., & van Loon, J.J.A., 2019. Effects of dietary protein and carbohydrate on life-history traits and body protein and fat contents of the black soldier fly *Hermetia illucens*. Physiological Entomology 44: 148–159. https://doi.org/10.1111/phen.12285

Berggreen, I.E., Schøn, M.L., Nørgaard, J.V. and Jensen, K., 2025. Optimal dietary protein content and essential amino acid limitation in larvae of the black soldier fly (*Hermetia illucens*). Journal of Insects as Food and Feed. In Press. https://doi.org/10.1163/23524588-bja10221

Berner, D., Blanckenhorn, W.U. and Körner, C., 2005. Grasshoppers cope with low host plant quality by compensatory feeding and food selection: N limitation challenged. Oikos 111: 525–533. https://doi.org/10.1111/j.1600-0706.2005.14144.x

Broeckx, L., Frooninckx, L., Slegers, L., Berrens, S., Noyens, I., Goossens, S., Verheyen, G., Wuyts, A. and Van Miert, S., 2021. Growth of black soldier fly larvae reared on organic side-streams. Sustainability 13: 12953. https://doi.org/10.3390/su132312953

Cammack, J. A. and Tomberlin, J. K., 2017. The impact of diet protein and carbohydrate on select life-history traits of the black soldier fly *Hermetia illucens* (L.) (Diptera: Stratiomyidae). Insects 8: 56. https://doi.org/10.3390/insects8020056

Capuano, E., Oliviero, T., Fogliano, V. and Pellegrini, N., 2018. Role of the food matrix and digestion on calculation of the actual energy content of food. Nutrition Reviews 76: 274-289. https://doi.org/10.1093/nutrit/nux072

Davis, G.R.F., 1975. Essential dietary amino acids for growth of larvae of the yellow mealworm, *Tenebrio molitor* L. Journal of Nutrition 105: 1071–1075. https://doi.org/10.1093/jn/105.8.1071

Eggink, K.M., Donoso, I.G. and Dalsgaard, J., 2023. Optimal dietary protein to carbohydrate ratio for black soldier fly (*Hermetia illucens*) larvae. Journal of Insects as Food and Feed 9: 789–798. https://doi.org/10.3920/JIFF2022.0102

Eriksen, N.T., 2022. Dynamic modelling of feed assimilation, growth, lipid accumulation, and CO_2 production in black soldier fly larvae. PLoS ONE 17: e0276605. https://doi.org/10.1371/journal.pone.0276605

Gligorescu, A., Toft, S., Hauggaard-Nielsen, H., Axelsen, J.A. and Nielsen, S.A., 2019. Development, growth and metabolic rate of *Hermetia illucens* larvae. Journal of Applied Entomology 143: 875-881. https://doi.org/10.1111/jen.12653

Gold, M., Tomberlin, J.K., Diener, S., Zurbrügg, C. and Mathys, A., 2018. Decomposition of biowaste macronutrients, microbes, and chemicals in black soldier fly larval treatment: a review. Waste Management 82: 302–318. https://doi.org/10.1016/j.wasman.2018.10.022

Goodrich, H.R., Wood, C.M., Wilson, R.W., Clark, T.D., Last, K.B. and Wang, T., 2024. Specific
dynamic action: the energy cost of digestion or growth? Journal of Experimental Biology 227:
jeb246722. https://doi.org/10.1242/jeb.246722

Hawkins, A.J.S., 1991. Protein turnover: a functional appraisal. Functional Ecology 5: 222–233.
https://doi.org/10.2307/2389260

Jakobsen, K. and Thorbekt, G., 1993. The respiratory quotient in relation to fat deposition in
fattening–growing pigs. British Journal of Nutrition 69: 333–343.
https://doi.org/10.1079/bjn19930037

Janssen, R.H., Vincken, J.P., Van Den Broek, L.A.M., Fogliano, V. and Lakemond, C.M.M., 2017.
Nitrogen-to-protein conversion factors for three edible insects: *Tenebrio molitor*, *Alphitobius*
diaperinus, and *Hermetia illucens*. Journal of Agricultural and Food Chemistry 65: 2275–
2278. https://doi.org/10.1021/acs.jafc.7b00471

Koethe, M., Taubert, J. and Vervuert, I., 2022. Impact of lysine supplementation on growth and
development of *Hermetia illucens* larvae. Journal of Insects as Food and Feed 8: 35–44.
https://doi.org/10.3920/JIFF2020.0059

Laganaro, M., Bahrndorff, S. and Eriksen, N.T., 2021. Growth and metabolic performance of black
soldier fly larvae grown on low and high-quality substrates. Waste Management 121: 198–205.
https://doi.org/10.1016/j.wasman.2020.12.009

Lee, K.P., 2007. The interactive effects of protein quality and macronutrient imbalance on nutrient
balancing in an insect herbivore. Journal of Experimental Biology 210: 3236–3244.
https://doi.org/10.1242/jeb.008060

Lemme, A. and Klüber, P., 2024. Rethinking amino acid nutrition of black soldier fly larvae
(*Hermetia illucens*) based on insights from an amino acid reduction trial. Insects 15: 862.
https://doi.org/10.3390/insects15110862

Liu, C., Wang, C. and Yao, H., 2019. Comprehensive resource utilization of waste using the black
soldier fly (*Hermetia illucens* (L.)) (diptera: Stratiomyidae). Animals 9: 349.
https://doi.org/10.3390/ani9060349

Lomonaco, G., Franco, A., De Smet, J., Scieuzo, C., Salvia, R. and Falabella, P., 2024. Larval frass
of *Hermetia illucens* as organic fertilizer: composition and beneficial effects on different crops.
Insects 15: 293. https://doi.org/10.3390/insects15040293

Oonincx, D.G.A.B., Gold, M., Bosch, G., Guillaume, J.B., Rumbos, C.I., Naser El Deen, S.,
Sandrock, C., Bellezza Oddon, S., Athanassiou, C.G., Cambra-López, M., Pascual, J.J., Parodi,
A.P., Spranghers, T. and Yakti, W., 2025. Bugbook: nutritional requirements for edible insect
rearing. Journal of Insects as Food and Feed. In Press. https://doi.org/10.1163/23524588-
bja10226

Oonincx, D.G.A.B., van Broekhoven, S., van Huis, A. and van Loon, J.J.A., 2015. Feed
conversion, survival and development, and composition of four insect species on diets
composed of food by-products. PLoS ONE 10: e0144601.
https://doi.org/10.1371/journal.pone.0144601

Parodi, A., De Boer, I.J.M., Gerrits, W.J.J., Van Loon, J.J.A., Heetkamp, M.J.W., Van Schelt, J.,
Bolhuis, J.E. and Van Zanten, H.H.E., 2020. Bioconversion efficiencies, greenhouse gas and

ammonia emissions during black soldier fly rearing – a mass balance approach. Journal of
Cleaner Production 271: 122488. https://doi.org/10.1016/J.JCLEPRO.2020.122488

Pimentel, A.C., Montali, A., Bruno, D. and Tettamanti, G., 2017. Metabolic adjustment of the larval
fat body in *Hermetia illucens* to dietary conditions. Journal of Asia-Pacific Entomology 20:
1307–1313. https://doi.org/10.1016/j.aspen.2017.09.017

Posit Team, 2023. RStudio: Integrated development environment for R (2023.9.1.494). Posit
Software (PBC). http://www.posit.co/

Purkayastha, D. and Sarkar, S., 2022. Sustainable waste management using black soldier fly larva: a
review. International Journal of Environmental Science and Technology 19: 12701–12726.
https://doi.org/10.1007/s13762-021-03524-7

R Core Team, 2023. A language and environment for statistical computing. R Foundation for
Statistical Computing. https://www.R-project.org.

Raubenheimer, D. and Simpson, S.J., 1997. Integrative models of nutrient balancing: application to
insects and vertebrates. Nutrition Research Reviews 10: 151–179.
https://doi.org/10.1079/nrr19970009

Schmidt-Nilsen, K., 1997. Animal physiology: adaption and environment. Cambridge University
Press, Cambridge, UK.

Schneider, L., Kisinga, B., Stoehr, N., Cord-Landwehr, S., Schulte-Geldermann, E., Moerschbacher,
B.M., Eder, K., Jha, R. and Dusel, G., 2025. Dietary protein levels in isoenergetic diets affect
the performance, nutrient utilization and retention of nitrogen and amino acids of *Hermetia*
illucens (L.) (Diptera: Stratiomyidae) larvae. Insects 16: 240.
https://doi.org/10.3390/insects16030240

Schøn, M.L., Jensen, K., Schow-Madsen, M., Gold, M., Mathys, A., Berggreen, I.E., Schou, T.M.,
Nørgaard, J.V. and Overgaard, J., 2025a. Energy expenditure and nutrient assimilation in
Hermetia illucens larvae; monitoring growth, metabolism, and body composition via gas
exchange. In preparation

Schøn, M.L., Mikkelsen, M.V.N., Jensen, K., Poulsen, J.M., Berggreen, I.E., Schou, T.M.,
Nørgaard, J.V. and Overgaard, J., 2025b. Effect of temperature on growth, metabolism, and gas
exchange in *Hermetia illucens* larvae reared under commercial and laboratory conditions.
Journal of Insects as Food and Feed 11: 1059–1074. https://doi.org/10.1163/23524588-
00001268

Secor, S.M., 2009. Specific dynamic action: a review of the postprandial metabolic response.
Journal of Comparative Physiology B 179: 1–56. https://doi.org/10.1007/s00360-008-0283-7

Siddiqui, S.A., Süfer, Ö., Çalışkan Koç, G., Lutuf, H., Rahayu, T., Castro-Muñoz, R. and Fernando,
I., 2024. Enhancing the bioconversion rate and end products of black soldier fly (BSF)
treatment – a comprehensive review. Environment, Development and Sustainability 27: 9673–
9741. https://doi.org/10.1007/s10668-023-04306-6

Spranghers, T., Moradei, A., Vynckier, K., Boudrez, M., Pinotti, L. and Ottoboni, M., 2024. Amino
acid requirements of yellow mealworm and black soldier fly larvae. Journal of Insects as Food
and Feed 6: 1047–1058. https://doi.org/10.1163/23524588-00001271

Stryer, L., 1995. Biochemistry. W.H. Freeman. New York, NY.

Talal, S., Cease, A., Farington, R., Medina, H.E., Rojas, J. and Harrison, J., 2021. High
carbohydrate diet ingestion increases post-meal lipid synthesis and drives respiratory exchange
ratios above 1. Journal of Experimental Biology 224: jeb240010.
https://doi.org/10.1242/jeb.240010

Talal, S., Harrison, J.F., Farington, R., Youngblood, J.P., Medina, H.E., Overson, R. and Cease, A.J.,
2024. Body mass and growth rates predict protein intake across animals. eLife 12: e88933.
https://doi.org/10.7554/eLife.88933

Tomberlin, J.K., Miranda, C., Flint, C., Harris, E. and Wu, G., 2023. Nutrients limit production of
insects for food and feed: an emphasis on nutritionally essential amino acids. Animal Frontiers
13: 64–70. https://doi.org/10.1093/af/vfad032

van der Heide, M.E., Stødkilde, L., Nørgaard, J.V. and Studnitz, M., 2021. The potential of locally-
sourced european protein sources for organic monogastric production: a review of forage crop
extracts, seaweed, starfish, mussel, and insects. Sustainability 13: 2303.
https://doi.org/10.3390/su13042303

Van't Hof, H.M. and Martin, M.M., 1989. The effect of diet water content on energy expenditure by
third-instar *Manduca sexta* larvae (Lepidoptera: Sphingidae). Journal of Insect Physiology 35:
433–436. https://doi.org/10.1016/0022-1910(89)90118-2

Von Bertalanffy, L., 1957. Quantitative laws in metabolism and growth. Quarterly Review of
Biology 32: 217–231. https://doi.org/10.1086/401873

Wu, G., Bazer, F.W., Dai, Z., Li, D., Wang, J. and Wu, Z., 2014. Amino acid nutrition in animals:
protein synthesis and beyond. Annual Review of Animal Biosciences 2: 387–417.
https://doi.org/10.1146/annurev-animal-022513-114113

618 **Supplementary Information**

619 Table S1. Ingredient and nutrient compositions in the chicken feed (Kyllekræs 1, Danish Agro) used
620 in the experiment. Ingredient and nutrient composition were given by the producer, and amino acid
621 contents were analysed.

Chicken feed

Ingredients (% as fed)

Wheat	32.0
Soybean meal, dehulled, toasted	30.0
Corn	25.0
Rapeseeds	4.0
Oat	2.0
Fish meal	1.6
Limestone	1.3
Sunflower seed meal	1.1
Soybean oil	1.0
Monocalcium phosphate	0.9
Vitamin and mineral mix	0.4
Sodium bicarbonate	0.3
Sodium chloride	0.2

Nutrient composition (% of dry matter)

Crude protein	20.3
Crude lipids	5.2
Carbohydrates	68.3
Digestible carbohydrates	64.9
Micronutrients (ash)	6.2
Energy (kJ/g dry matter)	17.0
Metabolizable energy (kJ/g dry matter)	16.4

Amino acids (% of protein mass)

Arginine	6.3
Histidine	2.5
Isoleucine	4.5
Leucine	7.7
Lysine	5.4
Methionine	2.2
Phenylalanine	4.8
Threonine	3.7
Tryptophan	1.1
Valine	5.0
Alanine	4.5
Asparagine	2.6
Aspartic acid	9.3
Cysteine	1.6
Glutamine	2.6
Glutamic acid	18.0
Glycine	4.2
Proline	5.7
Serine	5.1
Tyrosine	3.2