No effect of continuous transcutaneous auricular vagus nerve stimulation on the P3 and the P600 in an oddball and sentence comprehension task

Friederike Contier[1], Isabell Wartenburger[2], Mathias Weymar[1], & Milena Rabovsky[1]

[1]*Cognitive Sciences, Department of Psychology, University of Potsdam, Potsdam, Germany*

[2]*Cognitive Sciences, Department of Linguistics, University of Potsdam, Potsdam, Germany*

Corresponding author: Friederike Contier (Friederike.contier@uni-potsdam.de)

Conflict of interest statement

The authors declare no competing financial interests.

Acknowledgements

We thank L. Y. Lam, A. Sterzik, J. Kirschner, and J. Barthelmeß for their help with data collection. Funding was provided by the Research Focus Cognitive Sciences of the University of Potsdam and by the Deutsche Forschungsgemeinschaft (DFG, German Research Foundation, Project ID 317633480, SFB1287).

Abstract

The ERP components P3 and P600 have been proposed to reflect phasic activity of the locus coeruleus norepinephrine (LC/NE) system in response to deviant and task-relevant stimuli across cognitive domains. Yet, causal evidence for this link remains limited. Here, we used continuous transcutaneous auricular vagus nerve stimulation (taVNS), a non-invasive method proposed to modulate LC/NE activity, to test whether these components are indeed sensitive to noradrenergic manipulation. Forty participants completed both an active visual oddball task and a sentence processing task including both syntactic and semantic violations, while receiving continuous taVNS at the cymba conchae in one session and sham stimulation at the earlobe in another session. We observed robust P3 and P600 effects. Crucially though, taVNS had no effect on P3 or P600 amplitude. The physiological NE markers salivary alpha amylase level and baseline pupil size were also unaffected by the stimulation, suggesting that the taVNS protocol and/or task may not have been sufficient to successfully engage the norepinephrine system. Beyond the stimulation, however, exploratory analyses revealed correlations between the syntactic P600 and both the P3 and salivary alpha amylase levels, supporting the idea that the P600 might be related to both the P3 and NE. Overall, our findings do not allow for theoretical implications concerning a potential causal link between the two components and NE but highlight the need for more standardized taVNS protocols.

Key words: ERP, P3, P600, LC/NE system, taVNS

Introduction

Components of the event-related potential (ERPs) have been widely studied as neural correlates of specific cognitive processes. They have traditionally been examined within domain-specific contexts – such as emotion regulation, attention in the visual oddball paradigm, or structural processing in language tasks – leading to a fragmented understanding of their functional significance. To move beyond functional description of single ERPs and toward a mechanistic understanding of these components that may cross the boundaries of cognitive subdomains, it is crucial to consider not only their cortical generators but also the neurochemical systems that support them. ERPs are assumed to reflect postsynaptic potentials from synchronized neural populations (Luck, 2014; Mitzdorf, 1994) and are modulated by neurotransmitter activity that governs neuronal excitability and synaptic transmission (Frodl-Bauch et al., 1999). However, the specific roles that different neuromodulatory systems play in shaping ERP components remain largely unexplored with some initial exceptions. For example, the locus coeruleus norepinephrine (LC/NE) system has been proposed as a key neuromodulatory mechanism underlying a family of late positive ERP components, including the P3 and P600. Here we test this hypothesis using transcutaneous auricular vagus nerve stimulation (taVNS), a noninvasive stimulation method that has been linked to release of norepinephrine from the locus coeruleus (Badran et al., 2018; Farmer et al., 2021; Frangos et al., 2015; Schneider et al., 2025). This approach allows us to probe the neurochemical underpinnings and potential overlap of these ERP components, possibly suggesting a shared, domain-general mechanism.

The P3 is a robust and widely studied ERP component, a positive going deflection over centro-parietal electrodes, peaking around 300-600 ms post stimulus onset (Polich, 2007). It is typically elicited by so-called 'oddballs', that is, rare and relevant target stimuli among frequent non-target standard stimuli across domains (Donchin, 1981). The P3 has been

extensively used as a neural index of stimulus evaluation, attentional resource allocation, saliency processing, and context updating (e.g., Donchin, 1981; Donchin & Coles, 1988; Kok, 2001; Nieuwenhuis et al., 2005; Polich, 2007). However, its precise functional significance remains under debate. For instance, the P3's broad sensitivity to stimulus salience, expectancy violation, and behavioral relevance has initiated a discussion about whether it might reflect a specific cognitive operation or rather a domain-general integration process (O'Connell et al., 2012; Verleger, 2020). Further, although several cortical and subcortical neural generators have been proposed – mostly focusing on parietal and frontal cortices (e.g., Bledowski et al., 2004; Halgren et al., 1998; Linden, 2005) – the neurochemical mechanisms underlying the P3 are not fully understood.

The P600 is a late positive ERP component typically peaking between 500 and 900 ms after target word onset and is widely used in research on language processing. It is well known to be sensitive to various (morpho-)syntactic violations as well as ambiguities in sentence processing (e.g., Friederici et al., 1993, 1996, 2001; Gouvea et al., 2010; Hagoort et al., 1993; Münte et al., 1998; Osterhout et al., 1994; Osterhout & Mobley, 1995). For example, it is observed in response to number agreement mismatches between the subject and verb of a sentence, as in "The spoilt child _throw_ the toys on the ground." (transl. from Dutch, Hagoort et al., 1993). It is traditionally interpreted as specifically reflecting structural processing in language, such as syntactic re-analyses, integration, or effort required to build a coherent sentence structure (e.g., Friederici, 1995, 2002; Gouvea et al., 2010; Hagoort, 2003; Kaan et al., 2000; Osterhout et al., 1994). The component is thus commonly used as a neural marker for the presence of such syntactic operations in experimental paradigms. However, more recent research indicates that the P600 is elicited by a much wider range of linguistic anomalies, for instance also on the semantic, pragmatic, discourse, and even orthographic level (e.g., Brothers et al., 2020, 2021; Burkhardt, 2006; DeLong et al., 2014; Hoeks et al., 2004; Kim & Osterhout, 2005; Kuperberg et al., 2003; Münte et al., 1998; Regel et al., 2014;

Sassenhagen & Fiebach, 2019; Van Herten et al., 2005). The P600 has thus also been proposed as a general signal of error monitoring or correction in language comprehension (e.g., Ryskin et al., 2021; van de Meerendonk et al., 2009, 2010). Alternatively, it could reflect the difficulty to establish and keep a single, coherent representation of the linguistic utterance (e.g., Brouwer et al., 2012; Kuperberg et al., 2020; Li & Ettinger, 2023). Yet another alternative is that the P600 is not even specific to language at all, but is a more domain-general response to conflict or unexpected stimuli. Indeed, the anomalies eliciting a P600 are not limited to language: For example, incongruencies in music, arithmetic, as well as narrative structures in visual events also elicit parietal responses that strongly resemble the typical P600 (e.g., Christiansen et al., 2012; Cohn & Kutas, 2015; McLean et al., 2020; Núñez-Peña & Honrubia-Serrano, 2004; Patel et al., 1998; Sitnikova et al., 2008; Võ & Wolfe, 2013).

Thus, it has been proposed that the P600 might actually be a late variant of the P3 (Coulson et al., 1998a, 1998b; Leckey & Federmeier, 2019; Sassenhagen et al., 2014; Sassenhagen & Bornkessel-Schlesewsky, 2015; Sassenhagen & Fiebach, 2019). There are indeed striking similarities between the P600 in linguistic contexts and the P3 in other domains: both are late, centro-parietal positivities with indistinguishable morphology of waveforms (Sassenhagen & Fiebach, 2019). They are both remarkably sensitive to stimulus probability, salience and task relevance (e.g., Leckey & Federmeier, 2019; Pritchard, 1981) and their underlying processes require attention and cognitive control (e.g., Brothers et al., 2021; Contier et al., 2022; Gratton et al., 2018; Xu et al., 2021). However, this proposed relationship of positivities across domains is not without challenge. For example, Osterhout et al. (1996) reported that when syntactic violations occurred in a physical oddball context, the resulting ERP waveform reflected an additive effect of a P3 and a P600, which has been taken to suggest that the two may be functionally and neurophysiologically distinct. Further, in contrast to the P3, the P600 can be elicited by linguistic anomalies that do not require overt

decisions**,** and it often shows less sensitivity to stimulus probability than is typical for the P3 (Vissers et al., 2007). Taken together, while the hypothesis that the P600 reflects a domain-general signal akin to the P3 is supported by several lines of converging evidence, it remains an open question whether the P600 should be interpreted as a P3 variant, a language-specific component, or a functionally distinct – but partially overlapping – late positivity.

One prominent theory proposes that the P3 and P600 share a neural generator: norepinephrine (NE, also called noradrenaline) release from the locus coeruleus (LC) in the brain stem which plays a central role in modulating attention, arousal, and behavioral adaptation (Aston-Jones & Cohen, 2005). Such late positivities, including the P3, the late positive potential, and the P600, are assumed to reflect such phasic NE response to salient and motivationally significant events across domains (Bornkessel-Schlesewsky & Schlesewsky, 2019; Hajcak & Foti, 2020; Murphy et al., 2011; Nieuwenhuis et al., 2005; Sassenhagen et al., 2014; Sassenhagen & Bornkessel-Schlesewsky, 2015; Vazey et al., 2018). The LC-NE projects to relevant target cortices, where it promotes focused attention to relevant stimuli (neural gain, Nieuwenhuis et al., 2005) or acts as a neural interrupt signal (Bouret & Sara, 2005; Dayan & Yu, 2006).

There are indeed several functional parallels between the positivities and the LC/NE system: First, NE responses are related to motivational significance and salience as well as sensitive to stimulus probability and task relevance (Nieuwenhuis et al., 2005, 2011). Second, the LC/NE system is strongly oddball sensitive (e.g., Aston-Jones et al., 1997). Third, just like LC/NE response itself, the latency of both the P3 and P600 have been shown to align more closely with reaction times than with stimulus onsets (Bouret & Sara, 2004; Kutas et al., 1977; Sassenhagen et al., 2014; Sassenhagen & Bornkessel-Schlesewsky, 2015). This temporal alignment supports the notion that both components index similar cognitive control or salience-driven processes, potentially mediated by phasic norepinephrine. In addition, all three – the LC/NE system**,** the P3**,** and the P600 – have been implicated in long-term memory

facilitation. By activating attentional networks and involvement of limbic structures, phasic LC/NE activation has been shown to promote explicit memory formation, especially under conditions of heightened salience or arousal (e.g., Aston-Jones & Cohen, 2005; Gibbs et al., 2010; Izumi & Zorumski, 1999; Mather et al., 2016; Sara, 2009). Analogously, both P3 and P600 amplitudes have been found to predict subsequent memory performance, suggesting that these components may reflect neurocognitive processes that prioritize information for encoding (e.g., Contier et al., 2025; Dunn et al., 1998; Fabiani et al., 1986, 1990; Karis et al., 1984; Neville et al., 1986; Paller et al., 1988).

There is also ongoing research attempting to directly link the positivities to activity of the LC and its correlates. For example, unit recordings suggest simultaneous firing of LC neurons and P3 during oddball tasks in macaques (Swick et al., 1994). As measuring the LC directly remains challenging in humans, much research focuses on the task-evoked pupillary response, a common non-invasive marker of NE (e.g., Joshi et al., 2016; Liu et al., 2017; Murphy et al., 2014; Rajkowski et al., 1993; Sterpenich et al., 2006). While earlier studies failed to find covariation between the pupil and P3 (Hong et al., 2014; Kamp & Donchin, 2015; LoTemplio et al., 2020; Murphy et al., 2011), recent research suggests that it can be uncovered if response-related activity is taken into account (Chang et al., 2024; Menicucci et al., 2024). Contier et al. (2024) also found first evidence that the amplitude of the P600 covaries with the task-evoked pupillary response, particularly its temporal derivative. However, it remains difficult to unambiguously uncover the potential relationship between the pupillary response and both ERP components.

Beyond correlation, it is crucial to establish a causal relationship between the two positivities and the LC/NE system. Previous research suggests that the P3 can indeed be modulated by interventions that alter NE release, such as pharmacological manipulations (e.g., Joseph & Sitaram, 1989; Pineda & Westerfield, 1993). Together with lesion studies in rodents and monkeys (e.g., Ehlers & Chaplin, 1992; Pineda et al., 1989), these studies indeed

suggest a causal link between the P3 and norepinephrine, while such intervention research regarding the P600 is lacking thus far.

TaVNS has emerged as a promising non-invasive technique to engage central neuromodulatory pathways, with a particular emphasis on the LC/NE system (Badran et al., 2018; Farmer et al., 2021; Frangos et al., 2015; Szeska et al., 2025). The method involves electrical stimulation of the auricular branch of the vagus nerve, typically at the cymba conchae or tragus, which are known to be innervated by afferent vagal fibers (Badran et al., 2018; Peuker & Filler, 2002). These fibers project to the nucleus tractus solitarius in the brainstem, which relays signals to key neuromodulatory centers, such as the LC (e.g., Frangos et al., 2015; Kawai, 2018). Converging evidence from animal models and neuroimaging supports the notion that (ta)VNS can engage the LC and modulate central noradrenergic activity (e.g., Dorr & Debonnel, 2006; Ludwig et al., 2023; Schneider et al., 2025; Yakunina et al., 2017). In humans, taVNS has been found to affect task-evoked pupil responses (e.g., D'Agostini et al. 2023; Lloyd et al., 2023; Ludwig et al., 2023; Pervaz et al., 2025; Sharon et al., 2021; Skora et al., 2024; Villani et al., 2022; but see e.g., Keute et al., 2019; C. M. Warren et al., 2019; Burger et al., 2020), salivary alpha-amylase levels (e.g., Giraudier et al., 2022; Schneider et al., 2025; Ventura-Bort et al., 2018; C. M. Warren et al., 2019), heart rate variability (Kaduk et al., 2025) as well as electrophysiological processing in attention and memory tasks (e.g., Collins et al., 2021; Giraudier et al., 2020; Ventura-Bort et al., 2021; Ventura☐Bort et al., 2025), all of which are influenced by noradrenergic activity.

Several studies have tested whether taVNS can modulate the amplitude of the P3 but findings to date have been mixed. Some studies have reported increased P3 amplitudes under taVNS, particularly in auditory or visual oddball paradigms, although sometimes restricted to specific types of oddball stimuli (Giraudier et al., 2024; Gurtubay et al., 2022; Rufener et al., 2018; Ventura-Bort et al., 2018; C. V. Warren et al., 2020). For instance, Ventura-Bort et al. (2018) found that while there was no overall effect of taVNS on the P3, modulations were

observed, particularly for easy targets (c.f., Giraudier et al., 2024). Others have failed to observe any reliable effects (D'Agostini et al., 2022; Fischer et al., 2018; Gadeyne et al., 2022; C. M. Warren et al., 2019). Thus, the literature remains inconclusive, motivating further replication, especially when combined with physiological validation measures. Importantly, evidence of potential taVNS effects on the P600 are completely lacking. If the P600 also reflects a phasic LC/NE response to salient stimuli, then increasing LC/NE activity via taVNS should amplify or modulate the P600, similar to how it might affect the classic oddball P3.

Here we thus investigated whether the P3 and P600 are causally linked to the LC/NE system using taVNS. While receiving continuous taVNS (one session) and sham stimulation (another session), participants completed an active oddball paradigm aimed to elicit the P3 as well as a sentence processing task including both syntactic and semantic violations aimed to elicit the P600. Since taVNS is assumed to innervate the LC/NE system, the amplitude of both the P3 and P600 should be enhanced under taVNS compared to sham if they reflect LC/NE activity. This modulatory taVNS effect should be specific to the two late positivities. We also recorded the N400 in response to semantic violations in the sentence processing task, which should not be affected by taVNS, as it might reflect a rather automatic semantic prediction error and learning signal with no putative link to the LC/NE system (e.g., Rabovsky & McClelland, 2020). To confirm that taVNS (but not sham) indeed affected the LC/NE system, we assessed salivary alpha amylase levels, a putative autonomic NE correlate (e.g., Bosch et al., 2011; Nater & Rohleder, 2009) that has previously been shown to be modulated by taVNS (e.g., Giraudier et al., 2022; Schneider et al., 2025; Ventura-Bort et al., 2018; C. M. Warren et al., 2019). To contribute further to evidence for potential taVNS markers, we additionally measured baseline pupil size at several points in the study, as this NE-associated marker (e.g., Joshi et al., 2016; Laeng et al., 2012; Reimer et al., 2016) has to the best of our knowledge not been successfully manipulated via taVNS (e.g., Burger et al., 2020a, 2020b; D'Agostini et al., 2022; Keute et al., 2019; Skora et al., 2024).

Method

The study including methods, hypotheses, and statistical models was pre-registered on the
Open Science Framework
(https://osf.io/fgbxs/?view_only=743d5d402e5249de8d4f2ed753dd5bb4). Any deviations
from this plan are explicitly stated. The linked OSF project
(https://osf.io/qfhab/?view_only=510598a63fe542fe8e4f6758585841b9) also hosts all pre-
processing and analyses scripts as well as the pre-processed data.

Participants

Forty naïve volunteers participated in the experiment (35 female, 5 male). Three
participants had to be excluded and replaced since they did not complete both sessions
(preregistered data exclusion criteria). Mean age was 24.1 years (*SD* = 5.5, range = 18-38).
All participants were native speakers of German and right-handed according to the Edinburgh
Handedness Inventory (Oldfield, 1971; 12-item version; mean laterality quotient: 86, range:
41-100). Participation was compensated monetarily or with course credit. Each individual
provided written informed consent for a protocol approved by the ethics committee of the host
institution (No. 85/2020 amendment). Participants were pre-screened online for the following
exclusion criteria: Impaired and uncorrected vision, nystagmus, pregnancy, metal close to the
left ear, chronic or acute medication, neurological or mental disorders, chronic illnesses
(endocrine, metabolic, cardio-vascular, pulmonary, renal), metal implants, pumps, or
pacemakers, history of migraine, epilepsy or brain surgery, and/or speech/language
impairments.

Procedure

The study consisted of two experimental sessions, which took place seven days apart, at the same time of the day. In one session, participants received the vagus stimulation and in the other session the sham stimulation (order counterbalanced across participants). Participants were not aware of the manipulation (single-blind study). Apart from the stimulation condition, the two sessions were identical (Figure 1A below). Testing took place in a sound-proof and electrically shielded booth with dimmed light. After study information, consent, and preparation of the EEG cap, the NE markers salivary alpha amylase levels and baseline pupil size were measured ("pre"). Subsequently, the stimulation electrode was applied, intensity adjusted, and device turned on (see below). Then, the two tasks (oddball task, sentence processing task) proceeded, in counterbalanced order (but kept constant across sessions for each participant). Between the tasks, there was a short break in which salivary alpha amylase levels and baseline pupil size were measured again ("post1"). As soon as the second task was finished, the stimulation device was turned off and salivary alpha amylase levels and baseline pupil size were measured one last time ("post2"). Timing of both tasks (both stimulus presentation times and breaks) were controlled as much as possible in order to keep the duration of the experimental session under stimulation as constant as possible. Each experimental session lasted approximately 40 minutes. At the end of each session, participants filled out a questionnaire about potential adverse effects of the vagus nerve stimulation: On a 7-point scale ranging from 1 (not at all) to 7 (very much), they rated how much they felt the following sensations during the stimulation: headache, nausea, dizziness, neck pain, neck muscle contractions, stinging sensation, skin irritation on the ear, fluctuations of concentration, fluctuations of feelings, or other unpleasant feelings (Giraudier et al., 2020, 2025; Ventura-Bort et al., 2018). At debriefing after the second session, participants learned about the broader theoretical goal of the study as well as that it involved two stimulation

settings for comparison more generally. None of the participants indicated that they knew or

guessed the true nature of sham vs. active stimulation during the experiment.

Figure 1.

(A) Experimental procedure, identical across the two sessions and stimulation conditions.

sAA = Salivary alpha amylase sample. PM = Pupillometry Measurement (baseline pupil size

during 5 seconds of fixation cross). Stimulation was either vagus (one session) or sham

(another session). (B) Stimulus presentation in the oddball task, with an example stimulus

sequence in the block in which "D" was the target letter. (C) Word-by-word stimulus

presentation of one trial in the sentence processing task.

sAA = Salivary alpha amylase sample

PM = Pupillometry (Baseline pupil size during 5s fixation cross)

Transcutaneous auricular vagus nerve stimulation (taVNS)

The taVNS stimulation device consisted of two titan electrodes attached to a stabilizing mount and wired to a stimulation unit (CMO2, Cerbomed, Erlangen, Germany). The mount was always placed in the left auricle, but the location of the electrodes differed between conditions. In the vagus condition, the electrodes were positioned at the cymba conchae, which is exclusively innervated by the vagus nerve (Peuker & Filler, 2002). In the sham condition, the electrodes were rotated so that they touched the center of the earlobe (e.g., Frangos et al., 2015; Ventura-Bort et al., 2018). Thus, in the sham condition, participants had the same tingling sensation as in the vagus condition, but the stimulation did not affect the vagus nerve, since the earlobe is assumed not to be innervated by the vagus nerve (Peuker & Filler, 2002). Conservatively, stimulation was applied on the left ear in order to avoid potential cardiac side effects, although recent research does not suggest any risks even when using the right ear (e.g., Gentile et al., 2025; Tan et al., 2025). The device emitted continuous stimulation at a pulse width of 200-300 μs at 25 Hz for the duration of both tasks (appr. 40 minutes). Since the stimulation sensation varies across individuals, we adjusted the intensity of the stimulation to ensure that all participants had the same sensation during the stimulation, above the detection, but below the pain threshold (c.f., Ventura-Bort et al., 2018). In an adjustment procedure using a scale from 1 (no sensation) to 10 (pain), we determined the intensity with which the participants judged the sensation as an 8 (maximum tingling, without discomfort). The mean intensity in the vagus condition was 0.99 mA ($SD = 0.541$, range = 0.4-2.4) and in the sham condition 1.088 mA ($SD = 0.649$, range = 0.4-3.1). Statistically, intensities did not significantly differ between stimulation conditions ($t(39) = -1.563$, $p = .126$, d = 0.163).

Tasks & materials

Stimuli of both tasks were presented visually on a light grey computer screen in black Geneva font, at a distance of 60 cm from the participants.

Oddball task

To elicit the P3, we used an active, visual oddball task specifically designed to elicit a strong ERP effect while diminishing physical confounds (Kappenman et al., 2021). The task is to identify rare target stimuli ('oddballs') across many frequent non-targets ('standards').

In each instance of the task, five letters served as stimuli, each presented with a probability of .2. There were five blocks of trials and for each block, another of the five letters was designated to be the target letter (oddball) for that block. Thus, in each block, the oddball probability was .2 and the standard probability was .8, while each letter served both as a target and non-target across the whole task. Within a block, letters were presented in random order, with the restriction that targets were never followed by targets. The order of blocks was also randomized. In each session and thus, stimulation condition, we presented a total of 200 trials (40 targets and 160 non-targets), divided into five blocks of 40 trials each.

In one session, the stimulus letters were A, B, C, D, and E and in the other, K, L, M, N, and O (counterbalanced across stimulation conditions and sessions). Stimulus letters were presented in the center of the screen (height: 6.676 degrees of visual angle) for 200 ms (Fig. 1B). During the inter-stimulus interval (ISI), a fixation cross was shown for 1200-1400 ms (jittered in steps of the screen refresh interval of 16.667 ms using a rectangular distribution).

Participants' task was to respond as quickly and accurately as possible whether the presented letter is the target or non-target for that particular block. Two marked (control) keys served as response buttons for 'left' and 'right' responses, each pressed by the index finger of the respective hand. Stimulus-response mapping (right key target & left key non-target, or

vice versa) was counterbalanced across participants, but held constant within participants across tasks and sessions (see below). Written instructions were presented at the beginning of the task and reminders after each break within the task. Before the experimental blocks, participants practiced the task using a set of letters different from the experimental letters (27 trials) and were only allowed to proceed with above-chance accuracy. Between blocks, participants had a short break to rest their eyes (20 seconds). The task lasted approximately 12 minutes.

Sentence processing task

This task was a classic sentence processing task in which sentences are presented word by word and participants are prompted to make an acceptability judgement after each sentence. We used German sentences by Sassenhagen & Fiebach (2019, Exp. 2), which contain 50% correct sentences, 25% syntactic violations and 25% semantic violations.

Sentences were simple, plausible subject-verb-object (S-V-O) or object-verb-subject (O-V-S) sentences. For the syntactic violation, a second version of each sentence with an agreement violation had been constructed by exchanging the verb or pronominal subject (example 1 below) for a morphosyntactic ill-fitting one from another sentence (thereby creating an agreement violation). Although this constitutes a morphosyntactic violation, we will henceforth call it "syntactic" violation for simplicity. Semantic violations had been created by either exchanging a noun (example 2 below) or verb between two experimental sentences rendering each of the sentences highly implausible.

(1) „Den Zoo besuchen[pl] **sie**[pl] **/ *er**[sg] am Montag" [The zoo visit **they / *he** on Monday]

(2) "Der Chirurg operiert **den Leistenbruch/*die Butter** mit Vorsicht" [The surgeon operates the **rupture/*butter** carefully]

For more details on the construction of these sentences, see Sassenhagen & Fiebach (2019, Exp. 2). We also added 30 correct filler sentences in the same fashion as correct control sentences to increase the proportion of correct sentences. The final set comprised 350 sentences: 320 experimental sentences (80 syntactic violations and 80 correct control counterparts; 80 semantic violations and 80 correct control counterparts) as well as 30 correct filler sentences. These sentences were divided into 2 lists, which were counterbalanced across participants. This resulted in a classic latin-square design where each participant sees only one version of each sentence, but across participants, each sentence is presented in both its correct control and violation version equally often. These lists were then further divided into two parts, one for each experimental session (order counterbalanced across sessions and stimulation conditions). Thus, in each session (and thus, stimulation condition), participants were presented with 175 sentences of which 46% contained some kind of violation (23% syntactic, 23% semantic).

Within each session, sentences were presented in random order, with the only constraint that violation sentences of a certain type were never directly followed by another violation sentence of the same type. Each trial (see Figure 1C) started with a fixation cross, shown for two seconds. Single words of the sentence were then presented consecutively, each for 300 ms with a 300 ms blank inter-word interval. If a non-target word's length exceeded 12 letters, 16.667 ms (one screen refresh interval) were added to its presentation duration for each additional letter (for comparable approaches, see e.g., Contier et al., 2024; Hodapp & Rabovsky, 2021; Nicenboim et al., 2020). Words had a height of 1.41° degrees of visual angle. 300 ms after the offset of the final word, a question mark appeared, prompting the participant to decide as fast and accurately as possible whether the presented sentence was syntactically or semantically anomalous. Two marked (control) keys on the computer keyboard served as response buttons for 'left' and 'right' responses, each pressed by the index finger of the respective hand. Stimulus-response mapping was counterbalanced across

participants but held constant in both tasks and sessions within each participant. Thus, the same key, either the left or right control key, was assigned for targets in the oddball task and violations in the sentence processing task and vice versa. The question mark was shown until button press (but max. for 2500 ms). Participants received no feedback. The next trials started 300 ms after the response.

Short rest breaks were provided every 29 trials (approx. 20 seconds). Prior to the start of the experimental sentences, participants were provided with written instructions. Then, participants practiced the task with nine additional sentences including correct control, syntactic violation and semantic violation sentences and were only allowed to proceed with above-chance accuracy (with one chance of repetition). The task lasted approximately 27 minutes.

Data acquisition & processing

EEG data

In both tasks and sessions, we recorded participants' electroencephalogram (EEG) using 32 passive Ag/AgCl electrodes (actiCHamp Plus system, Brain Products GmbH, Gilching, Germany), spaced according to the international 10-20 system. Impedances were kept below 5 kOhm if possible. The ground electrode was positioned at FCz and the reference electrode at the right mastoid. Right-side VEOG as well as bilateral HEOG electrodes were positioned to capture eye movements and blinks. The EEG was recorded at a sampling rate of 1000 Hz (with a bandpass filter of .016-250 Hz and time constant of 10s).

Offline, we preprocessed the data in *MATLAB* R2020a (The MathWorks Inc., 2020), using the *EEGLAB* toolbox (Delorme & Makeig, 2004). Data were first downsampled to 500 Hz and re-referenced to the average of the left and right mastoid. We then used independent component analysis (Infomax ICA, Jung et al., 2001) to correct ocular artifacts. ICA was

trained on epochs spanning -0.5 to +2s (oddball task) and -1 to +2s (sentence processing task) relative to stimulus/target word onset after artifact rejection (removal of epochs exceeding +150/-150 mV). These epochs were extracted from continuous data downsampled to 125 Hz and filtered with a 1-30 Hz Butterworth filter. We removed independent components that the *IClabel* plug-in (Pion-Tonachini et al., 2019) identified to be eye-related with a probability equal to or larger than 30%. We additionally removed (max. three) channel noise components or components clearly originating from the taVNS device (predominantly 25 Hz activity on left side) based on visual inspection.

Corrected, continuous data were high-pass filtered (0.1 Hz, two-pass Butterworth with a 12 dB/oct roll-off) and low-pass filtered (30 Hz, two-pass Butterworth with a 24 dB/oct roll-off). Residual noise caused by the taVNS device was filtered out using EEGLAB's CleanLine plug-in (at 25Hz + harmonics). Bad channels exceeding an absolute z-score of 3 regarding voltage across channels were interpolated using a spherical spline function. Data were then epoched from -200 to 800 ms (oddball task) and from -200 to 1100 ms (sentence processing task) time-locked to stimulus/target word onset and baseline corrected relative to the 200 ms interval preceding this onset. Artifactual trials exceeding +/-75 mV were automatically removed.

As preregistered, error trials (i.e., trials in which participants responded incorrectly) were excluded. Table 1 displays the number of trials per task and condition combination after data preprocessing. In none of the stimulus conditions did the number of trials significantly differ between stimulation conditions (all p's > .13). Epoched data were averaged over preregistered electrode clusters and time windows and resulting trial-wise ERP values were used in linear mixed effects models. Data for both late, positive components were analyzed in a parietal region of interest (ROI: CP1, CPz, CP2, P3, Pz, P4, PO3, POz, PO4) within a 300-600 ms (P3) and 600-900 ms (P600) time window. These spatial ROIs and time windows had been chosen based on the topography and time where each of the components had been shown

to be largest in the respective task in previous studies (Contier et al., 2024; Kappenman et al., 2021; Sassenhagen & Fiebach, 2019). N400 amplitudes (in response to semantic violations and controls) were computed within a 300-500 ms time window. As preregistered, we computed N400 amplitudes within two different ROIs: For comparability, the parietal region identical to the one for the P3 and P600 mentioned above and a centro-parietal channel cluster more commonly chosen for the N400 (Cz, CP1, CPz, CP2, Pz; see e.g., Hodapp & Rabovsky, 2021; Kuperberg et al., 2020).

Table 1.

Mean (SD) number of trials per participant in each task, stimulus condition, and stimulation condition after data preprocessing. Number of presented trials in parentheses behind the stimulus condition.

	Stimulus condition	**Vagus condition**	**Sham condition**
Oddball task	Oddball condition (40)	39.5 (0.72)	39.52 (0.6)
	Standard condition (160)	152.55 (3.71)	153.73 (3.3)
Sentence processing task	Syntactic violation condition (40)	36.925 (3.13)	36.8 (2.79)
	Semantic violation condition (40)	36.475 (2.87)	36.33 (3.03)
	Control condition (80)	75 (3.71)	74.43 (3.79)

Salivary alpha amylase levels

As our main manipulation check, salivary alpha amylase (sAA) levels were collected that are proposed to correlate with NE activity (Bosch et al., 2011; Nater & Rohleder, 2009). Saliva was collected at three points during each session to investigate potential effects of the stimulation: Before the stimulation was turned on (pre), during the stimulation in the break between the two tasks (post1) and right after the stimulation was turned off (post2). Saliva

was collected using the spitting method (Hodgson & Granger, 2013): Participants were instructed to pool saliva in their mouth for about one minute and then spit it into a tube using a plastic straw. Samples were stored at -18 degrees celsius and alpha amylase levels determined via immuno-assay by *Dresden Lab Services GmbH*. Data from three participants had to be excluded since they had at least one sample in which sAA levels could not be determined. Alpha amylase levels were not normally distributed (skewness = 1.549, *SE* = 0.162) and were thus log-transformed.

sAA levels were also used to explore whether the effect of stimulation on ERP amplitudes depends on whether the stimulation actually affected the NE system in a particular participant. For these analyses, we calculated sAA "change" values that quantify how much sAA levels changed from baseline (pre). For each participant and session (i.e., stimulation condition), sAA values before stimulation onset (pre) were subtracted from values during (post1) and after (post2) the stimulation. Positive change values thus indicate an increase of sAA levels and negative values a decrease of sAA values compared to the time before the stimulation was started.

Baseline pupil size

As a second potential manipulation check, pupil baseline was measured since pupil size has been proposed to be influenced by the LC/NE system. We defined "baseline" as opposed to the pupillary response evoked by the task or stimulus (as in e.g., Contier et al., 2024). In particular, pupil size was measured for 5000 ms while participants sat quietly and looked at a fixation cross. Just like saliva, we were mainly interested in pupil size at three timepoints in each session: Before the stimulation was turned on (pre), during the stimulation in the break between the two tasks (post1) and right after the stimulation was turned off (post2). In addition, we aimed to capture any potential more fine-grained or short-lived effects of the stimulation on pupil size by measuring the pupil size during every short break (i.e.,

every 3-4 minutes) during the tasks starting right before the stimulation was turned on and

finishing right after it was turned off, resulting in 15 measurements including the three main

timepoints of interest (pre, post1, post2). Pupil size was measured using a Tobii Nano at 60

Hz at both eyes, sitting 60 cm away from the screen. The pre and post1 measurement were

preceded by a 5-point calibration procedure (see e.g., Mckinnon et al., 2020; Trueswell et al.,

2013). Offline, pupil size for each timepoint was pre-processed using the pipeline by Kret and

Sjak-Shie (2019) which involves artifact correction by removing dilation speed outliers,

trend-line deviations, and secluded samples (e.g., due to noise of erroneous pupil detection

during shut eyes). Pupil size was then averaged over the left and right pupil, low-pass filtered

(cutoff at 4Hz) and interpolated. Before interpolation, epochs were excluded that had < 50%

data points left. Our final measure for statistical analyses was the mean pupil size at each

timepoint.

Statistical analyses

For the primary, preregistered analyses of ERPs and the potential effect of the

stimulation, we performed linear mixed-effects model (LMM) analyses using the *R* package

lme4 (R core Team, 2018). This approach accounts for subject-specific and item-specific

variability and utilizes all trials, increasing statistical power (Baayen et al., 2008). These

models had the following fixed effects structure:

*ERP amplitude ~ stimulus condition * stimulation*

All models were built using trial-by-trial data. In these models, both categorical

regressors were sum coded[1] (Schad et al., 2018). In addition, these models included a random

effects structure. In all models, we included random intercepts and slopes for the main effects

---

[1] Coding of relevant regressors: Stimulation: sham (-.5) vs. vagus (.5); Oddball task: standard

(-.5) vs. oddball (.5); Sentence processing task: violation (-.5) vs. control (.5).

and interaction between the fixed effects per subject. In models with sentence processing task data, we additionally included random intercepts and slopes for the main effects and interaction between the fixed effects per sentence. Following the recommendations by Barr et al. (2013), we tried to fit this maximal random effect structure but reduced its complexity successively until the model converged. As pre-registered, we first increased the number of iterations to 10000, then introduced an optimizer ("bobyca"), and then removed correlations between random intercepts and slopes and if necessary, entire random slopes. The significance of fixed effects was determined via likelihood ratio tests to compare the fit of the model to that of a reduced model lacking the respective fixed effect but including all remaining fixed effects as well as the same random effect structure. To further investigate potential interaction effects, we conducted sub-group contrasts using the *emmeans* package (Lenth et al., 2023) with Bonferroni correction for multiple comparisons. We preregistered that if no effect of stimulation can be found, we would explore any shorter lived differences between vagus and sham stimulation on ERP amplitude. These tests compared subject-wise averaged ERP amplitudes between vagus and sham conditions in response to oddball/violations across all timepoints of the respective epoch. Clusters of consecutive timepoints exceeding a two-sided threshold ($p < .05$, Bonferroni-corrected) were tested against a null distribution of maximum cluster masses (5000 permutations, paired-sample t-tests).

To analyze potential effects of the stimulation on the NE markers, we conducted 2 (stimulation) x 3 (timepoint) repeated measures ANOVAs. As preregistered, we generally used the standard $p < .05$ criterion for determining if the likelihood ratio tests suggest that the results are significantly different from those expected if the null hypothesis were correct.

Preregistered hypotheses

Norepinephrine markers

If taVNS affects NE activity, two assessed NE markers (salivary alpha amylase and baseline pupil size) should increase more strongly or decrease less (from pre-stimulation to post-stimulation) under vagus than sham stimulation. This effect could arise regarding either of the two post-stimulation timepoints (in between tasks and after offset of the stimulation). If no such interaction effect on pupil baseline can be detected at these two timepoints, its development over the session should increase more under vagus than under sham stimulation.

ERPs

We expect a main effect of stimulus condition on the amplitude of the ERP components: more positive amplitudes on oddballs than standards (P3 effect) and on target words in sentences with syntactic violations than on control sentences (P600 effect), as well as more negative amplitudes on target words in semantic violation than control sentences (N400 effect).

Further, no effect of stimulation on the P3/P600 amplitude would be taken as evidence against the LC/NE-P3-P600 hypothesis. Conversely, the following effects would suggest that the respective component (P3, P600) is linked to LC/NE activity: 1) Main effect of stimulation on the ERP amplitude (larger amplitude under vagus than sham stimulation), 2) interaction effect between stimulus type (oddballs/syntactic violations vs standards/control sentences) and stimulation (vagus vs sham) in that amplitudes on oddballs/syntactic violations are larger under vagus than under sham, and/or 3) significant differences in group comparison tests regarding the interaction between stimulus condition and stimulation.

Since conversely, the N400 has not been proposed to be linked to LC/NE activity, its amplitude should not be affected by stimulation, meaning neither an interaction between stimulus condition (semantic violation vs control) and stimulation (vagus vs sham) nor a main effect of stimulation on amplitudes in the N400 time window.

Results

Subjective ratings on stimulation effects

In line with the previous literature (see Giraudier et al., 2025), subjective ratings on the 7-point scale suggested that participants experienced little to no side effects during the stimulation ($M = 1.35$, see Table 2 below). The Wilcoxon signed rank test (ratings were not normally distributed) revealed almost no differences between stimulation conditions (all ps > .332). Only stinging was experienced significantly stronger under vagus than under sham stimulation ($p = .025$).

Table 2.

Mean subjective ratings (standard deviation) for the stimulation side effects in the sham and vagus condition on the scale ranging from 1 (not at all) to 7 (very much).

	Sham	**Vagus**
Headache	1.35 (0.66)	1.35 (0.89)
Nausea	1.1 (0.5)	1.03 (0.16)
Dizziness	1.33 (0.88)	1.25 (0.63)
Neck pain	1.23 (0.48)	1.23 (0.58)
Neck contraction	1.3 (0.56)	1.4 (0.59)
Stinging sensation	1.45 (0.9)	1.83 (1.17)
Ear irritation	1.15 (0.43)	1.3 (1.02)
Concentration	2.5 (1.17)	2.4 (1.22)
Fluctuation of feelings	1.2 (0.72)	1.2 (0.61)
Unpleasant feelings	1.23 (0.62)	1.5 (0.91)

Norepinephrine markers

Salivary alpha amylase levels

sAA levels in each condition over all three timepoints are displayed in Figure 2A below. There was no significant main effect of stimulation on sAA levels (F(1,216) = [0.005], p = .944). There was also no main effect of timepoint on sAA levels (F(2,216) = [0. 565], p = .569), meaning that overall sAA did not systematically rise or fall across the session (see Fig. 2A). This suggests that neither the stimulation nor time-on-task significantly altered sAA. Crucially, there was also no significant interaction between timepoint and stimulation (F(2,216) = [0.050], p = .951, all Tukey HSD comparison p's > .95). Similarly, there was no significant difference in our exploratory measure of sAA change values between stimulation conditions regarding post1 (t(72) = -0.185, p = .854) or post2 (t(72) = -0.549, p = .585). Together, this suggests that the stimulation did not have any effect on the NE marker sAA level.

Baseline pupil size

Baseline pupil size in each stimulation condition over all three main timepoints is displayed in Figure 2B below. There was a significant main effect of timepoint ((F(2,234) = [4.476], p = .944) in that pupil size decreased from baseline to post2 (p = .015), indicating general habituation over the experiment. However, the stimulation did not affect baseline pupil size, neither as a main effect (F(1,234) = [0.061], p = .806) nor in interaction with timepoint (F(2,234) = [0.016], p = .984). As preregistered, we also explored potential more fine-grained effects of stimulation on pupil size. Figure 2C displays pupil size in all 15 measurements across the course of the entire experiment. However, we failed to find any shorter-lived effects, as paired t-tests revealed no significant differences between vagus and

sham condition at any timepoint (all Bonferroni-corrected p's > .551). Thus, the stimulation

did not have any measurable effect on the NE marker baseline pupil size.

Figure 2.

*Results of NE markers. (**A**) sAA Levels and (**B**) baseline pupil size in each of the two*

stimulation conditions over the course of the three main timepoints: "pre" = baseline

measurement before onset of stimulation, "Post1" = during stimulation, in break between

tasks. "Post2" = right after tasks were finished and stimulation was turned off again.

Raincloud plots depict individual data points (dots), their distribution (half–violin density),

*and summary statistics (boxplots). (**C**) depicts mean pupil size over the entire course of the*

experiment in each of the stimulation conditions. The grey box signifies the time window in

which the stimulation was on and error bars indicate the standard error of the mean.

Oddball task

Behavioral performance

Overall accuracy was very high ($M = 96\%$; $SD = 2\%$) and was significantly better in the standard condition ($M = 99\%$, $SD = 2\%$) than in the oddball condition ($M = 84\%$, $SD =$

8%; β = -2.867, SE = 0.164, z = -17.471, $\chi2$ = 91.34, p < .001). The stimulation did not affect

accuracy, neither as a main effect (β = 0.115, SE = 0.124, z = 0.928, $\chi2$ = 0.84, p = .359), nor

in interaction with stimulus condition (β = -0.230, SE = 0.2, z = -1.150, $\chi2$ = 1.249, p = .263).

Within correct trials, responses to standard stimuli were significantly faster than to oddball

stimuli (β = 0.155, SE = 0.009, t = 17.306, $\chi2$ = 86.578, p < .001). However, there was no

significant main effect of the stimulation on RTs (β = -0.002, SE = 0.011, t = -0.163, $\chi2$ =

0.027, p = .869). Lastly, there was no significant interaction between stimulus condition and

stimulation on RTs in the oddball task (β = -0.001, SE = 0.011, t = -0.117, $\chi2$ = 0.014, p =

.904). Note that all reported analyses regarding behavior were exploratory.

Effect of stimulus condition and stimulation on P3 amplitude

Stimulus condition means of the ERP amplitude in the oddball task are shown in

Figure 3. We find the classic P3 effect: ERP amplitudes (300-600 ms) over posterior regions

were significantly larger for oddballs (M = 9.96 mV, SD = 8.37) than standards (M = 3.93

mV, SD = 7.22; β = 6.041, SE = 0.512, t = 11.81, $\chi2$ = 60.829, p < .001). However, the

stimulation did not significantly affect ERP amplitudes, neither as a main effect (β = -0.36, SE

= 0.26, t = -1.382, $\chi2$ = 1.913, p = .167; mean condition difference = 0.33 mV) nor in

interaction with stimulus condition (β = -0.002, SE = 0.27, t = -0.008, $\chi2$ = 5.877, p = .994).

None of the preregistered posthoc group contrasts (stimulus condition x stimulation) revealed

any significant effects (all p's > .531). The additional, exploratory cluster-based permutation

test also revealed no significant ERP amplitude differences between the vagus and sham

condition (max cluster mass = 53.18; p = .212). Thus, taVNS did not measurably influence

the P3.

Figure 3.

Grand averaged ERP amplitude for the four conditions in the oddball task. P3 amplitudes were calculated as the mean in the time interval marked by the grey box (300-600 ms) within the posterior channel cluster marked in light green in the topographic map. The topographic map shows the mean amplitude difference between oddball and standard stimuli within the same time window. Error bands indicate the standard error of the mean (SEM).

Influence of sAA change on stimulation effect on P3

　　　　Although the stimulation did not significantly affect sAA levels on a group level, we explored whether a potential effect of the stimulation on the P3 amplitude might be restricted to participants for which the stimulation increased sAA levels (as an index of LC/NE engagement). We quantified this by using sAA change values (see Method). The rationale was the following: If the stimulation in a session would successfully affect the LC/NE system, NE markers such as sAA levels should increase. We thus explored whether an (individual) increase of sAA level, possibly due to the stimulation, affected P3 amplitude. We did so by including the continuous measure of sAA change (two values per participant, one for sham

and one for stimulation) as a predictor in the original P3 model, that is, in addition to an interaction with stimulus condition and stimulation. However, sAA change was not related to any effect of the stimulation on the P3. Regarding the change to post1 (during the stimulation), there was no significant main effect of sAA change on P3 amplitude (β = -0.232, SE = 0.294, t = -0.789, $\chi2$ = 0.526, p = .468) and no interaction between sAA and stimulus condition (β = 0.277, SE = 0.242, t = 1.140, $\chi2$ = 1.258, p = .262), or sAA and stimulation (β = -0.152, SE = 0.555, t = -0.273, $\chi2$ = 3.351, p = .341). Lastly, there was no 3-way interaction between sAA change, stimulus condition, and stimulation (β = -0.095, SE = 0.571, t = -0.167, $\chi2$ = 0.026, p = .871). The same pattern was found regarding the change to post2 (after the stimulation): There was no significant main effect of sAA change on P3 amplitude (β = 0.218, SE = 0.28, t = 0.778, $\chi2$ = 0.579, p = .447) and no significant interaction of sAA change with stimulus condition (β = 0.219, SE = 0.282, t = 0.777, $\chi2$ = 0.585, p = . 445) or stimulation (β = 0.306, SE = 0.523, t = 0.586, $\chi2$ = 0.318, p = .573). Again, there was no 3-way interaction of sAA change with stimulus condition and stimulation (β = 1.076, SE = 0.542, t = 1.986, $\chi2$ = 3.823, p = .051). In sum, change of sAA level over the course of the experiment did not affect the amplitude of the P3, neither by itself not mediated through the stimulation.

Influence of task order on stimulation effect on P3

Lastly, we explored whether the stimulation effect on the P3 might have been dependent on stimulation duration (Bömmer et al., 2024; Giraudier et al., 2024). We thus investigated whether task order (i.e., whether the oddball task came earlier in the session or not) affected the stimulation effect on the P3 by including it as an additional predictor in the model. However, task order did not influence how the stimulation affected ERP amplitudes as there was no significant 3-way interaction between task order, stimulus condition, and stimulation (β = -0.906, SE = 0.541, t = -1.676, $\chi2$ = 2.81, p = .1) and no 2-way interaction between task order and stimulation (β = -0.670, SE = 0.516, t = -1.298, $\chi2$ = 1.735, p = .188).

Sentence processing task

Behavioral performance

Overall accuracy in the sentence acceptability judgement was high ($M = 93\%$; $SD = 4\%$), indicating that participants attentively read the sentences. Accuracy was significantly better regarding correct control sentences ($M = 94\%$, $SD = 4\%$) than sentences including violations ($M = 92\%$, $SD = 5\%$; $\beta = -0.362$, $SE = 0.133$, $z = -2.730$, $\chi2 = 6.81$, $p = .009$). However, the stimulation did not affect accuracy, neither as a main effect ($\beta = 0.138$, $SE = 0.103$, $z = 1.337$, $\chi2 = 1.688$, $p = .194$), nor in interaction with stimulus condition ($\beta = -0.124$, $SE = 0.152$, $z = -0.815$, $\chi2 = 0.626$, $p = .429$). Within correct trials, response times were neither affected by stimulus condition ($\beta = -0.116$, $SE = 0.060$, $t = -1.92$, $\chi2 = 3.614$, $p = .057$), nor stimulation ($\beta = 0.028$, $SE = 0.062$, $t = 0.445$, $\chi2 = 0.203$, $p = .653$), nor their interaction ($\beta = -0.030$, $SE = 0.045$, $t = -0.678$, $\chi2 = 0.461$, $p = .497$). Note that all reported analyses regarding behavior were exploratory and collapsed across violation types.

Effect of stimulus condition and stimulation on P600 amplitude

Mean ERP amplitudes over time in each of the stimulus conditions in the sentence processing task are shown in Figure 4. Overall, we found a syntactic as well as semantic[2] P600 effect. Amplitudes between 600-900 ms at parietal sites were significantly larger in response to syntactic violations ($M = 4.91$ mV, $SD = 8.48$) than respective correct controls ($M = 0.677$ mV, $SD = 7.55$; $\beta = 4.198$, $SE = 0.479$, $t = 8.755$, $\chi2 = 43.42$, $p < .001$). Analogously, P600 amplitudes were significantly larger on semantic violations ($M = 2.81$ mV, $SD = 7.89$)

---

[2] Note that our preregistration focused on the syntactic P600 and how it is potentially influenced by the stimulation. However, since we also observed a P600 regarding semantic violations, we also explored whether the semantic P600 is influenced by the stimulation.

than respective correct controls ($M = 1.8$ mV, $SD = 7.16$; $\beta = 0.977$, $SE = 0.23$, $t = 4.252$, $\chi2 = 15.196$, $p < .001$).

Regarding the syntactic P600, the stimulation did not significantly affect ERP amplitudes, neither as a main effect ($\beta = 0.013$, $SE = 0.244$, $t = 0.051$, $\chi2 = 0.003$, $p = .959$, mean condition difference = 0.01 mV) nor in interaction with stimulus condition ($\beta = -0.586$, $SE = 0.395$, $t = -1.484$, $\chi2 = 2.201$, $p = .138$). None of the preregistered posthoc group contrasts (condition x stimulation) revealed any significant effects (all p's > .760). Likewise, the additional, exploratory cluster-based permutation test also did not reveal any significant ERP amplitude differences between the vagus and sham condition (max cluster mass = 20.88; $p = .67$).

The same results pattern was obtained regarding the semantic P600: There was no significant main effect of stimulation ($\beta = 0.192$, $SE = 0.190$, $t = 1.007$, $\chi2 = 1.015$, $p = .314$, mean condition difference = 0.18 mV), no interaction between stimulus condition and stimulation ($\beta = 0.275$, $SE = 0.381$, $t = 0.723$, $\chi2 = 0.522$, $p = .47$), and no significant posthoc group contrasts (all p's > .659). Again, the additional, exploratory cluster-based permutation test did not reveal any significant ERP amplitude differences between the vagus and sham condition (max cluster mass = 20.19; $p = .692$). Thus, taVNS did not measurably influence the amplitude of the syntactic or semantic P600.

Figure 4.

*Grand averaged ERP amplitudes in the sentence processing task aligned to target word onset for the (**A**) syntactic and (**B**) semantic sentence type, each divided by stimulation condition. P600 amplitudes were calculated as the mean in the time interval marked by the grey box (600-900 ms) within the posterior channel cluster marked in light green in the topographic maps. Topographic maps show the mean amplitude difference between violation and control stimuli within the same time window. Error bands indicate the standard error of the mean.*

Influence of sAA change on stimulation effect on P600

As in the oddball task, we explored whether a potential effect of the stimulation on the P600 amplitude might be restricted to participants for which the stimulation increased sAA levels (indicating LC/NE engagement). To that end, we included the continuous measure of sAA change as an interacting predictor in the original P600 models in addition to stimulus condition and stimulation.

Regarding the syntactic violations, sAA change to post1 (during the stimulation) did not significantly affect P600 amplitudes ($\beta = 0.055$, $SE = 0.235$, $t = 0.234$, $\chi2 = 0.106$, $p = .744$). There were also no significant interaction between sAA change and stimulation ($\beta = -0.786$, $SE = 0.481$, $t = -1.637$, $\chi2 = 2.745$, $p = .098$), or stimulus condition ($\beta = -0.406$, $SE = 0.337$, $t = -1.203$, $\chi2 = 1.42$, $p = .233$). Lastly, there was no significant 3-way interaction between sAA change, stimulus condition, and stimulation ($\beta = -0.253$, $SE = 0.775$, $t = -0.326$, $\chi2 = 0.116$, $p = .734$). For the change to post2 (after the stimulation), again, there was no significant main effect of sAA change ($\beta = 0.362$, $SE = 0.205$, $t = 1.770$, $\chi2 = 3.127$, $p = .077$) or interaction between sAA change and stimulation ($\beta = -0.397$, $SE = 0.457$, $t = -0.869$, $\chi2 = 0.71$, $p = .4$). Notably though, sAA change to post2 did significantly interact with stimulus

condition (β = -1.012, *SE* = 0.390, *t* = -2.594, $\chi2$ = 6.665, *p* = .01), reflecting the fact that

P600 amplitudes were larger the more sAA levels increased, but only in response to violations

(β = 1.017, *SE* = 0.321, *p* = .002), not controls (β = 0.005, *SE* = 0.293, *p* = .987). Lastly, there

was no significant 3-way interaction between sAA change, stimulation, and stimulus

condition (β = -0.369, *SE* = 0.756, *t* = -0.488, $\chi2$ = 0.248, *p* = .051)

Regarding semantic violations and post1 (during the stimulation) there was no

significant main effect of sAA change (β = -0.265, *SE* = 0.194, *t* = -1.368, $\chi2$ = 1.945, *p* =

.163), no interaction of sAA and stimulation (β = -0.105, *SE* = 0.396, *t* = -0.265, $\chi2$ = 0.073, *p*

= .787) and no interaction of sAA change and stimulus condition (β = 0.202, *SE* = 0.277, *t* =

0.727, $\chi2$ = 0.548, *p* = .459) on P600 amplitude. There was also no significant 3-way

interaction of sAA change with stimulation and stimulus condition (β = 0.058, *SE* = 0.610, *t* =

0.095, $\chi2$ = 0.008, *p* = .926). In the same vein, sAA change to post2 (after the stimulation) did

not significantly affect P600 amplitude (β = -0.027, *SE* = 0.261, *t* = -0.104, $\chi2$ = 0.009, *p* =

.923) and again did not significantly interact with stimulation (β = 0.110, *SE* = 0.378, *t* =

0.292, $\chi2$ = 0.076, *p* = .783), or with stimulus condition (β = 0.236, *SE* = 0.311, *t* = 0.759, $\chi2$

= 0.534, *p* = .465). Lastly, there was also no significant 3-way interaction between sAA

change, stimulus condition, and stimulation (β = 0.751, *SE* = 0.612, *t* = 1.228, $\chi2$ = 1.492, *p* =

.222).

Thus, sAA change did not influence any effect of the stimulation on the syntactic or

semantic P600. However, the observed exploratory interaction between sAA change and P600

amplitude indicates a relationship between the syntactic P600 and a marker of NE

independently of the stimulation.

Influence of task order on stimulation effect on P600

We again explored the possibility of time-dependent effects of stimulation. We thus

included task order as an additional (interacting) predictor in the P600 models. However, task

order did not influence how the stimulation affected P600 amplitudes as there was no significant 3-way interaction between task order, stimulus condition, and stimulation (syntactic P600: $\beta = 0.147$, $SE = 0.78$, $t = 0.186$, $\chi2 = 5.358$, $p = .253$; semantic P600: $\beta = 1.136$, $SE = 0.760$, $t = 1.494$, $\chi2 = 2.226$, $p = .136$). Likewise, there was no 2-way interaction between task order and stimulation (syntactic P600: $\beta = -0.261$, $SE = 0.493$, $t = -0.529$, $\chi2 = 0.294$, $p = .588$; semantic P600: $\beta = -0.072$, $SE = 0.380$, $t = -0.189$, $\chi2 = 0.035$, $p = .851$).

Effect of condition and stimulation on N400 amplitude

As a control component that is not presumed to be linked to NE release, we preregistered to test whether the N400 is affected by the stimulation. At the parietal channel cluster (same ROI as P3 and P600, see methods) we replicated the classic N400 effect: Amplitudes between 300-500 ms were significantly more negative on semantic violations ($M = -0.061$ mV, $SD = 7.45$) than respective correct controls ($M = 0.612$ mV, $SD = 7.32$; $\beta = -0.680$, $SE = 0.214$, $t = -3.182$, $\chi2 = 9.241$, $p = .002$, see Fig. 4B). As expected, the stimulation did not significantly affect N400 amplitudes, neither as a main effect ($\beta = 0.132$, $SE = 0.18$, $t = 0.695$, $\chi2 = 0.483$, $p = .487$, mean condition difference: 0.136 mV) nor in interaction with stimulus condition ($\beta = 0.398$, $SE = 0.38$, $t = 1.047$, $\chi2 = 1.098$, $p = .295$). None of the preregistered posthoc group contrasts (stimulus condition x stimulation) revealed any significant effects (all p's > .601).

We found the identical pattern of results at the centro-parietal channel cluster (common N400 ROI, see methods): There was an N400 effect between 300-500 ms in that amplitudes were significantly more negative on semantic violations ($M = -0.268$ mV, $SD = 8.44$) than correct controls ($M = 0.517$ mV, $SD = 8.34$; $\beta = -0.8$, $SE = 0.246$, $t = -3.255$, $\chi2 = 9.631$, $p = .002$). Again, the stimulation did not significantly affect N400 amplitudes, neither as a main effect ($\beta = 0.3$, $SE = 0.216$, $t = 0.695$, $\chi2 = 1.385$, $p = .174$, mean condition difference: 0.299 mV) nor in interaction with stimulus condition ($\beta = 0.602$, $SE = 0.433$, $t =$

1.390, $\chi2$ = 1.935, p = .164). None of the preregistered posthoc group contrasts (stimulus condition x stimulation) revealed any significant effects (all p's > . 206).

Relationship between ERP components across tasks

Lastly, we explored a potential relationship between the P600 and P3 effect (i.e., the mean amplitude difference between experimental and control condition) within participants, which would be predicted if both relied on the same neural generator.

Indeed, the syntactic P600 effect was positively predicted by the P3 effect (β = 0.412, SE = 0.141, t = 2.916, p = .005, Figure 5). This relationship was independent of the stimulation, as there was no interaction with stimulation (β = 0.105, SE = 0.202, t = 0.518, p = .606). Further, the relationship was unlikely due to more general cognitive state or mood, as the correlation persisted across both sessions (i.e., P600 effect in session 1 correlated with P3 effect in session 2 and vice versa, both p's < .002). Thus, we found that participants with larger P3 effects in the oddball task also displayed larger syntactic P600 effects in the sentence processing task independent of the stimulation, session, and order of tasks, further linking the two late, positive components.

Notably, this relationship of the P3 and P600 effect was specific to the syntactic P600 effect, as we did not find such an effect for the semantic P600 effect (main effect: (β = 0.085, SE = 0.098, t = 0.873, p = .385; interaction with stimulation: β = -0.088, SE = 0.14, t = -0.630, p = .530, Figure 5 below) or the N400 effect in response to semantic violations (main effect: β = -0.935, SE = 0.817, t = -1.144, p = .256; interaction with stimulation: β = 0.087, SE = 0.119, t = 0.734, p = .465, Figure 5).

Figure 5.

*Relationship (i.e., slope of fitted linear models) between P3 effect and (**A**) syntactic P600*

*effect, (**B**) semantic P600 effect, and (**C**) N400 effect in response to semantic violations.*

Shadows represent 95% confidence intervals.

Discussion

The present study investigated a modulatory effect of continuous transcutaneous auricular vagus stimulation on the ERP components P3 and P600, both of which have been proposed to reflect phasic noradrenergic activity in the locus coeruleus (e.g., Nieuwenhuis et al., 2005; Sassenhagen & Bornkessel-Schlesewsky, 2015). Moving beyond correlational evidence (e.g., Contier et al., 2024), we set out to test the hypothesis that phasic NE release causally contributes to the generation of the two late, positive ERP components. Using a within-subject, between-session design, we applied continuous taVNS to the cymba conchae, which has been linked to activation of the locus coeruleus-norepinephrine system, and compared it to sham stimulation at the earlobe. During stimulation in both sessions, participants completed an active visual oddball task (to elicit the P3) and an active sentence processing task including both syntactic and semantic violations (to elicit the P600). We observed robust main effects of stimulus condition: Oddball stimuli elicited larger P3 amplitudes than standard stimuli (P3 effect), and both syntactic and semantic violations elicited larger P600 amplitudes than target words in corresponding control sentences (P600 effect). In contrast, taVNS had no measurable effect on the amplitude of either component, neither by itself nor in interaction with stimulus condition. Notably, these null effects must be understood in the context of a possibly ineffective neuromodulatory manipulation. To assess whether taVNS affected NE as intended, we additionally measured two well-established physiological markers of NE activity (e.g., Joshi et al., 2016; Laeng et al., 2012; Nater & Rohleder, 2009): salivary alpha-amylase (sAA) and baseline pupil size at three timepoints (before, during, and after the stimulation). However, neither of the two measures showed evidence of change specifically due to the stimulation in the current study. Thus, our manipulation checks do not allow us to draw any major conclusions about a potential causal link between the two ERP components and NE.

The absence of taVNS effects on the ERP components may also be related to the mode of stimulation. In particular, the link between norepinephrine and both ERP components rests on the assumption that these components reflect phasic – not tonic – LC/NE activity. Phasic NE activity consists of large, transient, stimulus-bound bursts, while tonic activity is the continuous, baseline activity. Together, they jointly balance out exploratory and goal-directed behavior (Aston-Jones & Cohen, 2005). The P3 has been conceptualized as a neural signature of stimulus-driven salience detection via these phasic LC bursts (e.g., Nieuwenhuis et al., 2005) and the P600 may similarly reflect phasic arousal responses when encountering linguistically salient stimuli (e.g., Sassenhagen & Bornkessel-Schlesewsky, 2015). The continuous taVNS protocol in our study may have been suboptimal for engaging the targeted mechanism as continuous stimulation may primarily increase tonic LC activity or induce habituation. According to adaptive gain theory (Aston-Jones & Cohen, 2005), phasic LC responses are most effective when timed to salient events and embedded within an intermediate tonic state. Thus, even if taVNS elevated tonic NE levels, which may lead to optimal states for phasic responses for some participants, this may have also reduced the likelihood of observing noradrenergic responses in other participants – and, by extension, modulations of ERP components potentially sensitive to phasic NE activity, such as the P3 and P600. This mechanism could account for the heterogeneous effects on the P3 reported with continuous taVNS (e.g., d'Agostini et al., 2022; Fischer et al., 2018; Giraudier et al., 2024; Ventura-Bort et al., 2018). Whether continuous taVNS impacts phasic ERP components may further depend on the experimental context (e.g., properties of stimuli, stimulus context, task-relevance). For instance, continuous stimulation might exert stronger effects on late positive ERPs when phasic responses are elicited by emotionally salient (i.e., arousing) stimuli (e.g., Ventura-Bort et al., 2025) or target stimuli are presented in the context of arousing stimuli (e.g., Ventura-Bort et al., 2018; Giraudier et al., 2024; for discussion on the sAA, see Giraudier et al., 2022). Thus, the modulation of these ERP components by taVNS

may be shaped by the interaction between stimulation mode and the arousing properties of the stimuli or context (which also likely includes the saliency of stimuli given by task relevance).

The present findings thus highlight crucial methodological considerations but also the importance of the stimulation pattern, which may include settings to target phasic responses specifically. Studies using a 30s on / 30s off intermittent taVNS protocol have indeed also found enhanced P3 amplitudes (e.g., Lewine et al., 2019; C. V. Warren et al., 2020). Moreover, recent advances in short-term taVNS protocols that deliver brief bursts of stimulation have been proposed as a more precise and temporally targeted method for specifically affecting the phasic LC/NE response and have repeatedly been shown to enhance pupil dilation (e.g., D'Agostini et al. 2023; Lloyd et al., 2023; Ludwig et al., 2023; Pervaz et al., 2025; Sharon et al., 2021; Skora et al., 2024; Villani et al., 2022). Importantly, this can be used time-locked to cognitive events, specifically targeting the processing of task-relevant stimuli. For example, short bursts of taVNS during an encoding task led to better memory and larger pupil dilation for negative events (Ludwig et al., 2025). This method has also been shown to enhance target detection in rodents (Mridha et al., 2021). Interestingly, Mridha and colleagues (2021) observed the strongest phasic, stimulation evoked pupillary and neural responses during intermediate tonic NE activity, supporting the notion that providing baseline arousal conditions is crucial to successfully manipulate phasic NE activity. This phasic, event-locked method may recruit the LC/NE system more reliably than continuous, tonic stimulation approaches, offering promising methods to further investigate noradrenergic contributions to ERP components such as the P3 and P600. Ultimately, these considerations underscore the need for standardized, non-invasive stimulation protocols to target phasic NE activity specifically.

Given the likely complex relationship between positive ERP components and the neuromodulatory brain stem system, taVNS studies should furthermore be complemented by other modes of intervention. One avenue for future research might be to use alternative

methods to stimulate involved peripheral nerves such as the greater occipital nerve and the trigeminal nerve (e.g., Luckey et al., 2023; van Boekholdt et al., 2021) as well as pharmacological manipulations using NE agonists that have already been shown to modulate the P3 (e.g., Joseph & Sitaram, 1989). Another promising method may be isometric handgrip, a form of static muscular contraction in which participants constantly squeeze a handgrip device over several seconds (e.g., Nielsen & Mather, 2015). This transiently elevates sympathetic arousal and has been shown to increase phasic LC responsivity as well as pupil dilation in response to salient events and improve subsequent cognitive performance (Mather et al., 2020). As a brief, portable, and non-invasive technique, the handgrip method may serve as a valuable complementary tool here.

Although our taVNS protocol did not modulate ERP amplitudes or physiological norepinephrine markers, we did find two notable pieces of exploratory evidence supporting the P600's link to both the P3 and norepinephrine, respectively. First, we observed a within-subject correlation between the P3 effect and syntactic P600 effect that was independent of taVNS stimulation and session. Together with previous evidence on their shared link to the pupillary response (Contier et al., 2024), reaction time (Sassenhagen & Bornkessel-Schlesewsky, 2015) and the similarity of their waveforms (Sassenhagen & Fiebach, 2019), this correlation of ERP effects supports the possibility of shared or overlapping neurocognitive mechanisms, such as salience detection or context updating, potentially modulated by neuromodulators like norepinephrine. Secondly, larger sAA increase over the course of the experiment was associated with larger syntactic P600 amplitudes, suggesting that endogenous noradrenergic activity, as indexed by sAA, may enhance the neural processes underlying cognitive processing in linguistic contexts. Importantly, this exploratorily analysed relationship was selective to violation trials, suggesting that phasic LC/NE activity may specifically support processing in situations of conflict and/or uncertainty. Although correlational, this result supports the notion that the P600 is sensitive to norepinephrine and

suggests that naturally occurring variation in NE activity may be an important factor in late, positive ERP amplitude variation, even in the absence of experimental modulation. Lastly, these two links (i.e., the link of the P600 to the P3 and to the NE marker sAA level) were both found specifically for the P600 in response to syntactic, but not semantic violations. This may be partly attributable to the comparatively smaller semantic P600 effect (compared to the syntactic P600 effect), which likely constrained amplitude variability necessary to reveal reliable correlations with other neural or physiological measures. Nevertheless, it is an open question whether the P600s in response to syntactic and semantic violations reflect the same process and whether both or only the former are related to the P3 (Leckey & Federmeier, 2019). Thus, both findings certainly stimulate more research into the possible differences between the semantic and syntactic P600 and encourage incorporating a more diverse set of linguistic anomalies into single P600 studies to potentially uncover unknown processing differences and potential differential relationships to the P3 and NE.

In conclusion, our study finds no influence of continuous taVNS on either the P3 or P600 ERP component nor on physiological markers of norepinephrine, suggesting that the stimulation protocol may not have effectively modulated the LC/NE system. Thus, our results cannot provide evidence for a causal role of norepinephrine in the neural generation of the P3 or P600, but they also do not falsify this hypothesis. Instead, they highlight the need for more standardized taVNS protocols. Importantly, they emphasize the need for better-targeted manipulations of phasic norepinephrine activity, such as phasic, stimulus-aligned taVNS to advance our understanding of potential links between the late, positive ERP components and the LC/NE system. Lastly, a link between the P600 and P3 as well as increased sAA observed in exploratory analyses supports the view that the two positivities may be related and phasic norepinephrine release may contribute to late-stage language processing, particularly in response to incongruencies.

References

Aston-Jones, G., & Cohen, J. D. (2005). An Integrative Theory of Locus Coeruleus-Norepinephrine Function: Adaptive Gain and Optimal Performance. *Annual Review of Neuroscience*, *28*(1), 403–450.

https://doi.org/10.1146/annurev.neuro.28.061604.135709

Aston-Jones, G., Rajkowski, J., & Kubiak, P. (1997). Conditioned responses of monkey locus coeruleus neurons anticipate acquisition of discriminative behavior in a vigilance task. *Neuroscience*, *80*(3), 697–715. https://doi.org/10.1016/S0306-4522(97)00060-2

Baayen, R. H., Davidson, D. J., & Bates, D. M. (2008). Mixed-effects modeling with crossed random effects for subjects and items. *Journal of Memory and Language*, *59*(4), 390–412. https://doi.org/10.1016/j.jml.2007.12.005

Badran, B. W., Dowdle, L. T., Mithoefer, O. J., LaBate, N. T., Coatsworth, J., Brown, J. C., DeVries, W. H., Austelle, C. W., McTeague, L. M., & George, M. S. (2018). Neurophysiologic effects of transcutaneous auricular vagus nerve stimulation (taVNS) via electrical stimulation of the tragus: A concurrent taVNS/fMRI study and review. *Brain Stimulation*, *11*(3), 492–500. https://doi.org/10.1016/j.brs.2017.12.009

Barr, D. J., Levy, R., Scheepers, C., & Tily, H. J. (2013). Random effects structure for confirmatory hypothesis testing: Keep it maximal. *Journal of Memory and Language*, *68*(3), 255–278. https://doi.org/10.1016/j.jml.2012.11.001

Bledowski, C., Prvulovic, D., Hoechstetter, K., Scherg, M., Wibral, M., Goebel, R., & Linden, D. E. J. (2004). Localizing P300 Generators in Visual Target and Distractor Processing: A Combined Event-Related Potential and Functional Magnetic Resonance Imaging Study. *Journal of Neuroscience*, *24*(42), 9353–9360.

https://doi.org/10.1523/JNEUROSCI.1897-04.2004

Bömmer, T., Schmidt, L. M., Meier, K., Kricheldorff, J., Stecher, H., Herrmann, C. S., Thiel, C. M., Janitzky, K., & Witt, K. (2024). Impact of Stimulation Duration in taVNS— Exploring Multiple Physiological and Cognitive Outcomes. *Brain Sciences*, *14*(9), Article 9. https://doi.org/10.3390/brainsci14090875

Bornkessel-Schlesewsky, I., & Schlesewsky, M. (2019). Toward a neurobiologically plausible model of language-related, negative event-related potentials. *Frontiers in Psychology*, *10*, 1–17. https://doi.org/10.3389/fpsyg.2019.00298

Bosch, J. A., Veerman, E. C. I., de Geus, E. J., & Proctor, G. B. (2011). A-Amylase As A Reliable And Convenient Measure Of Sympathetic Activity: Don't start salivating just yet! *Psychoneuroendocrinology*, *36*(4), 449–453. https://doi.org/10.1016/j.psyneuen.2010.12.019

Bouret, S., & Sara, S. J. (2004). Reward expectation, orientation of attention and locus coeruleus-medial frontal cortex interplay during learning. *European Journal of Neuroscience*, *20*(3), 791–802. https://doi.org/10.1111/j.1460-9568.2004.03526.x

Bouret, S., & Sara, S. J. (2005). Network reset: A simplified overarching theory of locus coeruleus noradrenaline function. *Trends in Neurosciences*, *28*(11), 574–582. https://doi.org/10.1016/j.tins.2005.09.002

Brothers, T., Wlotko, E. W., Warnke, L., & Kuperberg, G. R. (2020). Going the Extra Mile: Effects of Discourse Context on Two Late Positivities During Language Comprehension. *Neurobiology of Language*, *1*(1), 135–160. https://doi.org/10.1162/nol_a_00006

Brothers, T., Zeitlin, M., Perrachione, A. C., Choi, C., & Kuperberg, G. R. (2021). Domain-general conflict monitoring predicts neural and behavioral indices of linguistic error processing during reading comprehension. *Journal of Experimental Psychology: General*. https://doi.org/10.1037/xge0001130

Brouwer, H., Fitz, H., & Hoeks, J. (2012). Getting real about Semantic Illusions: Rethinking

the functional role of the P600 in language comprehension. *Brain Research*, *1446*,

127–143. https://doi.org/10.1016/j.brainres.2012.01.055

Burger, A. M., Van der Does, W., Brosschot, J. F., & Verkuil, B. (2020). From ear to eye? No

effect of transcutaneous vagus nerve stimulation on human pupil dilation: A report of

three studies. *Biological Psychology*, *152*(December 2019), 107863.

https://doi.org/10.1016/j.biopsycho.2020.107863

Burkhardt, P. (2006). Inferential bridging relations reveal distinct neural mechanisms:

Evidence from event-related brain potentials. *Brain and Language*, *98*(2), 159–168.

https://doi.org/10.1016/j.bandl.2006.04.005

Chang, Y.-H., Chen, H.-J., Barquero, C., Tsai, H. J., Liang, W.-K., Hsu, C.-H., Muggleton, N.

G., & Wang, C.-A. (2024). Linking tonic and phasic pupil responses to P300

amplitude in an emotional face-word Stroop task. *Psychophysiology*, *61*(4), e14479.

https://doi.org/10.1111/psyp.14479

Christiansen, M. H., Conway, C. M., & Onnis, L. (2012). Similar neural correlates for

language and sequential learning: Evidence from event-related brain potentials.

Language and Cognitive Processes, *27*(2), 231–256.

https://doi.org/10.1080/01690965.2011.606666

Cohn, N., & Kutas, M. (2015). Getting a cue before getting a clue: Event-related potentials to

inference in visual narrative comprehension. *Neuropsychologia*, *77*, 267–278.

https://doi.org/10.1016/j.neuropsychologia.2015.08.026

Collins, L., Boddington, L., Steffan, P. J., & McCormick, D. (2021). Vagus nerve stimulation

induces widespread cortical and behavioral activation. *Current Biology*, *31*(10), 2088-

2098.e3. https://doi.org/10.1016/j.cub.2021.02.049

Contier, F., Höger, M., & Rabovsky, M. (2025). The P600 during sentence reading predicts behavioral and neural markers of recognition memory. *Journal of Cognitive Neuroscience*, 1-22. https://doi.org/ 10.1162/jocn.a.68

Contier, F., Wartenburger, I., Weymar, M., & Rabovsky, M. (2024). Are the P600 and P3 components linked to the task☐evoked pupillary response as a correlate of norepinephrine activity? *Psychophysiology*, *61*(7), 1–51. https://doi.org/10.1111/psyp.14565

Contier, F., Weymar, M., Wartenburger, I., & Rabovsky, M. (2022). Sustained Attention as Measured by Reaction Time Variability Is a Strong Modulator for the P600, but Not the N400. *Journal of Cognitive Neuroscience*, *34*(12), 2297–2310. https://doi.org/10.1162/jocn_a_01918

Coulson, S., King, J. W., & Kutas, M. (1998a). ERPs and Domain Specificity: Beating a Straw Horse. *Language and Cognitive Processes*, *13*(6), 653–672. https://doi.org/10.1080/016909698386410

Coulson, S., King, J. W., & Kutas, M. (1998b). Expect the Unexpected: Event-related Brain Response to Morphosyntactic Violations. *Language and Cognitive Processes*, *41*(1), 21–58. https://doi.org/10.1080/016909698386582

D'Agostini, M., Burger, A. M., Jelinčić, V., von Leupoldt, A., & Van Diest, I. (2023). Effects of transcutaneous auricular vagus nerve stimulation on P300 magnitudes and salivary alpha-amylase during an auditory oddball task. *Biological Psychology*, *182*, 108646. https://doi.org/10.1016/j.biopsycho.2023.108646

D'Agostini, M., Burger, A. M., Villca Ponce, G., Claes, S., Leupoldt, A., & Van Diest, I. (2022). No evidence for a modulating effect of continuous transcutaneous auricular vagus nerve stimulation on markers of noradrenergic activity. *Psychophysiology*, *59*(4), 88–100. https://doi.org/10.1111/psyp.13984

Dayan, P., & Yu, A. J. (2006). Phasic norepinephrine: A neural interrupt signal for unexpected events. *Network: Computation in Neural Systems*, *17*(4), 335–350. https://doi.org/10.1080/09548980601004024

DeLong, K. A., Quante, L., & Kutas, M. (2014). Predictability, plausibility, and two late ERP positivities during written sentence comprehension. *Neuropsychologia*, *61*(1), 150–162. https://doi.org/10.1016/j.neuropsychologia.2014.06.016

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Donchin, E. (1981). Surprise!? Surprise? *Psychophysiology*, *18*(5), 493–513. https://doi.org/10.1111/j.1469-8986.1981.tb01815.x

Donchin, E., & Coles, M. G. (1988). Is the P300 component a manifestation of context updating? *Behavioral and Brain Sciences*, *11*(3), 357–427. https://doi.org/10.1017/S0140525X00058027

Dorr, A. E., & Debonnel, G. (2006). Effect of vagus nerve stimulation on serotonergic and noradrenergic transmission. *Journal of Pharmacology and Experimental Therapeutics*, *318*(2), 890–898. https://doi.org/10.1124/jpet.106.104166

Dunn, B. R., Dunn, D. A., Languis, M., & Andrews, D. (1998). The relation of ERP components to complex memory processing. *Brain and Cognition*, *36*(3), 355–376. https://doi.org/10.1006/brcg.1998.0998

Ehlers, C. L., & Chaplin, R. I. (1992). Long latency event related potentials in rats: The effects of changes in stimulus parameters and neurochemical lesions. *Journal of Neural Transmission*, *88*(1), 61–75. https://doi.org/10.1007/BF01245037

Fabiani, M., Karis, D., & Donchin, E. (1986). P300 and Recall in an Incidental Memory Paradigm. *Psychophysiology*, *23*(3), 298–308. https://doi.org/10.1111/j.1469-8986.1986.tb00636.x

Fabiani, M., Karis, D., & Donchin, E. (1990). Effects of mnemonic strategy manipulation in a Von Restorff paradigm. *Electroencephalography and Clinical Neurophysiology*, *75*(1–2), 22–35. https://doi.org/10.1016/0013-4694(90)90149-E

Farmer, A. D., Strzelczyk, A., Finisguerra, A., Gourine, A. V., Gharabaghi, A., Hasan, A., Burger, A. M., Jaramillo, A. M., Mertens, A., Majid, A., Verkuil, B., Badran, B. W., Ventura-Bort, C., Gaul, C., Beste, C., Warren, C. M., Quintana, D. S., Hämmerer, D., Freri, E., … Koenig, J. (2021). International Consensus Based Review and Recommendations for Minimum Reporting Standards in Research on Transcutaneous Vagus Nerve Stimulation. *Frontiers in Human Neuroscience*, *14*. https://doi.org/10.3389/fnhum.2020.568051

Fischer, R., Ventura-Bort, C., Hamm, A., & Weymar, M. (2018). Transcutaneous vagus nerve stimulation (tVNS) enhances conflict-triggered adjustment of cognitive control. *Cognitive, Affective and Behavioral Neuroscience*, *18*(4), 680–693. https://doi.org/10.3758/s13415-018-0596-2

Frangos, E., Ellrich, J., & Komisaruk, B. R. (2015). Non-invasive Access to the Vagus Nerve Central Projections via Electrical Stimulation of the External Ear: fMRI Evidence in Humans. *Brain Stimulation*, *8*(3), 624–636. https://doi.org/10.1016/j.brs.2014.11.018

Friederici, A. D. (1995). The time course of syntactic activation during language processing: A model based on neuropsychological and neurophysiological data. *Brain and Language*, *50*(3), 259–281. https://doi.org/10.1006/brln.1995.1048

Friederici, A. D. (2002). Towards a neural basis of auditory sentence processing. *Trends in Cognitive Sciences*, *6*(2), 78–84. https://doi.org/10.1016/S1364-6613(00)01839-8

Friederici, A. D., Hahne, A., & Mecklinger, A. (1996). Temporal structure of syntactic parsing: Early and late event-related brain potential effects. *Journal of Experimental Psychology: Learning Memory and Cognition*, *22*(5), 1219–1248. https://doi.org/10.1037/0278-7393.22.5.1219

Friederici, A. D., Mecklinger, A., Spencer, K. M., Steinhauer, K., & Donchin, E. (2001).

Syntactic parsing preferences and their on-line revisions: A spatio-temporal analysis

of event-related brain potentials. *Cognitive Brain Research*, *11*(2), 305–323.

https://doi.org/10.1016/S0926-6410(00)00065-3

Friederici, A. D., Pfeifer, E., & Hahne, A. (1993). Event-related brain potentials during

natural speech processing: Effects of semantic, morphological and syntactic violations.

Cognitive Brain Research, *1*(3), 183–192. https://doi.org/10.1016/0926-

6410(93)90026-2

Frodl-Bauch, T., Bottlender, R., & Hegerl, U. (1999). Neurochemical substrates and

neuroanatomical generators of the event-related P300. *Neuropsychobiology*, *40*(2),

86–94. https://doi.org/10.1159/000026603

Gadeyne, S., Mertens, A., Carrette, E., Van den Bossche, F., Boon, P., Raedt, R., & Vonck,

K. (2022). Transcutaneous auricular vagus nerve stimulation cannot modulate the P3b

event-related potential in healthy volunteers. *Clinical Neurophysiology: Official*

Journal of the International Federation of Clinical Neurophysiology, *135*, 22–29.

https://doi.org/10.1016/j.clinph.2021.11.079

Gentile, F., Giannoni, A., Navari, A., Degl'Innocenti, E., Emdin, M., & Passino, C. (2025).

Acute right-sided transcutaneous vagus nerve stimulation improves cardio-vagal

baroreflex gain in patients with chronic heart failure. *Clinical Autonomic Research*,

35(1), 75–85. https://doi.org/10.1007/s10286-024-01074-9

Gibbs, M. E., Hutchinson, D. S., & Summers, R. J. (2010). Noradrenaline release in the locus

coeruleus modulates memory formation and consolidation; roles for α- and β-

adrenergic receptors. *Neuroscience*, *170*(4), 1209–1222.

https://doi.org/10.1016/j.neuroscience.2010.07.052

Giraudier, M., Ventura-Bort, C., Burger, A. M., Claes, N., D'Agostini, M., Fischer, R.,

Franssen, M., Kaess, M., Koenig, J., Liepelt, R., Nieuwenhuis, S., Sommer, A.,

Usichenko, T., Van Diest, I., von Leupoldt, A., Warren, C. M., & Weymar, M. (2022). Evidence for a modulating effect of transcutaneous auricular vagus nerve stimulation (taVNS) on salivary alpha-amylase as indirect noradrenergic marker: A pooled mega-analysis. *Brain Stimulation*, *15*(6), 1378–1388. https://doi.org/10.1016/j.brs.2022.09.009

Giraudier, M., Ventura-Bort, C., Szeska, C., & Weymar, M. (2025). A Pooled Analysis of the Side Effects of Non-Invasive Transcutaneous Auricular Vagus Nerve Stimulation (taVNS). *Frontiers in Human Neuroscience, 19*, 1539416. https://doi.org/10.3389/fnhum.2025.1539416

Giraudier, M., Ventura-Bort, C., & Weymar, M. (2020). Transcutaneous Vagus Nerve Stimulation (tVNS) Improves High-Confidence Recognition Memory but Not Emotional Word Processing. *Frontiers in Psychology*, *11*. https://doi.org/10.3389/fpsyg.2020.01276

Giraudier, M., Ventura-Bort, C., & Weymar, M. (2024). Effects of Transcutaneous Auricular Vagus Nerve Stimulation on the P300: Do Stimulation Duration and Stimulation Type Matter? *Brain Sciences*, *14*(7). https://doi.org/10.3390/brainsci14070690

Gouvea, A. C., Phillips, C., Kazanina, N., & Poeppel, D. (2010). The linguistic processes underlying the P600. *Language and Cognitive Processes*, *25*(2), 149–188. https://doi.org/10.1080/01690960902965951

Gratton, G., Cooper, P., Fabiani, M., Carter, C. S., & Karayanidis, F. (2018). Dynamics of cognitive control: Theoretical bases, paradigms, and a view for the future. *Psychophysiology*, *55*(3), 1–29. https://doi.org/10.1111/psyp.13016

Grunwald, T., Beck, H., Lehnertz, K., Blümcke, I., Pezer, N., Kurthen, M., ... & Elger, C. E. (1999). Evidence relating human verbal memory to hippocampal N-methyl-D-aspartate receptors. *Proceedings of the National Academy of Sciences*, *96*(21), 12085-12089. https://doi.org/10.1073/pnas.96.21.12085

Gurtubay, I. G., Perez-Rodriguez, D., Fernandez-López, E., Librero, J., Calvo, D., Bermejo, P., Pinin-osorio, C., & Lopez, M. (2022). Immediate effects and duration of a short and single application of transcutaneous auricular vagus nerve stimulation on P300 event related potential. *Frontiers in Neuroscience*, *Under cons*(March), 1–14. https://doi.org/10.3389/fnins.2023.1096865

Hagoort, P. (2003). How the brain solves the binding problem for language. *International Journal of Psychology*, *39*(5–6), 349–349. https://doi.org/10.1016/j.neuroimage.2003.1796

Hagoort, P., Brown, C., & Groothusen, J. (1993). The syntactic positive shift (sps) as an erp measure of syntactic processing. *Language and Cognitive Processes*, *8*(4), 439–483. https://doi.org/10.1080/01690969308407585

Hajcak, G. & Foti, D. (2020). Significance? Significance! Empirical, methodological, and theoretical connections between the late positive potential and P300 as neural responses to stimulus significance: An integrative review". *Psychophysiology* , *57*(7), 1–15. Httos://doi.org/10.1111/psyp.1357

Halgren, E., Marinkovic, K., & Chauvel, P. (1998). Generators of the late cognitive potentials in auditory and visual oddball tasks. *Electroencephalography and Clinical Neurophysiology*, *106*(2), 156–164. https://doi.org/10.1016/S0013-4694(97)00119-3

Hodapp, A., & Rabovsky, M. (2021). The N400 ERP component reflects an error-based implicit learning signal during language comprehension. *European Journal of Neuroscience*, *54*(9), 1–16. https://doi.org/10.1111/ejn.15462

Hodgson, N. A., & Granger, D. A. (2013). Collecting Saliva and Measuring Salivary Cortisol and Alpha-amylase in Frail Community Residing Older Adults via Family Caregivers. *Journal of Visualized Experiments*, *82*, 50815. https://doi.org/10.3791/50815

Hoeks, J. C. J., Stowe, L. A., & Doedens, G. (2004). Seeing words in context: The interaction of lexical and sentence level information during reading. *Cognitive Brain Research*, *19*(1), 59–73. https://doi.org/10.1016/j.cogbrainres.2003.10.022

Hong, L., Walz, J. M., & Sajda, P. (2014). Your eyes give you away: Prestimulus changes in pupil diameter correlate with poststimulus task-related EEG dynamics. *PLoS ONE*, *9*(3), e91321. https://doi.org/10.1371/journal.pone.0091321

Izumi, Y., & Zorumski, C. F. (1999). Norepinephrine promotes long-term potentiation in the adult rat hippocampus in vitro. *Synapse*, *31*(3), 196–202. https://doi.org/10.1002/(SICI)1098-2396(19990301)31:3<196::AID-SYN4>3.0.CO;2-K

Joseph, K. C., & Sitaram, N. (1989). The effect of clonidine on auditory P300. *Psychiatry Research*, *28*(3), 255–262. https://doi.org/10.1016/0165-1781(89)90206-0

Joshi, S., Li, Y., Kalwani, R. M., & Gold, J. I. (2016). Relationships between Pupil Diameter and Neuronal Activity in the Locus Coeruleus, Colliculi, and Cingulate Cortex. *Neuron*, *89*(1), 221–234. https://doi.org/10.1016/j.neuron.2015.11.028

Jung, T.-P., Makeig, S., McKeown, M. J., Bell, A. J., Lee, T.-W., & Sejnowski, T. J. (2001). Imaging brain dynamics using independent component analysis. *Proceedings of the IEEE*, *89*(7), 1107–1122. https://doi.org/10.1109/5.939827

Kaan, E., Harris, A., Gibson, E., & Holcomb, P. (2000). The P600 as an index of syntactic integration difficulty. *Language and Cognitive Processes*, *15*(2), 159–201. https://doi.org/10.1080/016909600386084

Kaduk, K., Petrella, A., Müller, S. J., Koenig, J., & Kroemer, N. B. (2025). Non-Invasive Auricular Vagus Nerve Stimulation Decreases Heart Rate Variability Independent of Caloric Load. *Psychophysiology*, *62*(2), e70017. https://doi.org/10.1111/psyp.70017

Kamp, S. M., & Donchin, E. (2015). ERP and pupil responses to deviance in an oddball paradigm. *Psychophysiology*, *52*(4), 460–471. https://doi.org/10.1111/psyp.12378

Kappenman, E. S., Farrens, J. L., Zhang, W., Stewart, A. X., & Luck, S. J. (2021). ERP

CORE: An open resource for human event-related potential research. *NeuroImage*,

225, 117465. https://doi.org/10.1016/j.neuroimage.2020.117465

Karis, D., Fabiani, M., & Donchin, E. (1984). "P300" and memory: Individual differences in

the von Restorff effect. *Cognitive Psychology*, *16*(2), 177–216.

https://doi.org/10.1016/0010-0285(84)90007-0

Kawai, Y. (2018). Differential Ascending Projections From the Male Rat Caudal Nucleus of

the Tractus Solitarius: An Interface Between Local Microcircuits and Global

Macrocircuits. *Frontiers in Neuroanatomy*, *12*, 63.

https://doi.org/10.3389/fnana.2018.00063

Keute, M., Demirezen, M., Graf, A., Mueller, N. G., & Zaehle, T. (2019). No modulation of

pupil size and event-related pupil response by transcutaneous auricular vagus nerve

stimulation (taVNS). *Scientific Reports*, *9*(1), 1–10. https://doi.org/10.1038/s41598-

019-47961-4

Kim, A., & Osterhout, L. (2005). The independence of combinatory semantic processing:

Evidence from event-related potentials. *Journal of Memory and Language*, *52*(2),

205–225. https://doi.org/10.1016/j.jml.2004.10.002

Kok, A. (2001). On the utility of P3 amplitude as a measure of processing capacity.

Psychophysiology, *38*(3), 557–577. https://doi.org/10.1017/s0048577201990559

Kret, M. E., & Sjak-Shie, E. E. (2019). Preprocessing pupil size data: Guidelines and code.

Behavior Research Methods, *51*(3), 1336–1342. https://doi.org/10.3758/s13428-018-

1075-y

Kuperberg, G. R., Brothers, T., & Wlotko, E. (2020). A Tale of Two Positivities and the

N400: Distinct neural signatures are evoked by confirmed and violated predictions at

different levels of representation. *Journal of Cognitive Neuroscience*, *32*(1), 12–35.

https://doi.org/10.1101/404780

Kuperberg, G. R., Sitnikova, T., Caplan, D., & Holcomb, P. J. (2003). Electrophysiological

distinctions in processing conceptual relationships within simple sentences. *Cognitive

Brain Research*, *17*(1), 117–129. https://doi.org/10.1016/S0926-6410(03)00086-7

Kutas, M., McCarthy, G., & Donchin, E. (1977). Augmenting mental chronometry: The P300

as a measure of stimulus evaluation time. *Science*, *197*(4305), 792–795.

https://doi.org/10.1126/science.887923

Laeng, B., Sirois, S., & Gredebäck, G. (2012). Pupillometry: A window to the preconscious?

Perspectives on Psychological Science, *7*(1), 18–27.

https://doi.org/10.1177/1745691611427305

Leckey, M., & Federmeier, K. D. (2019). The P3b and P600(s): Positive contributions to

language comprehension. *Psychophysiology*, *57*(7), 1–15.

https://doi.org/10.1111/psyp.13351

Lenth, R. V., Bolker, B., Buerkner, P., Giné-Vazquez, I., Herve, M., Jung, M., Love, J.,

Miguez, F., Riebl, H., & Singmann, H. (2023). *emmeans: Estimated Marginal Means,

aka Least-Squares Means* [Computer software]. https://github.com/rvlenth/emmeans

Lewine, J. D., Paulson, K., Bangera, N., & Simon, B. J. (2019). Exploration of the Impact of

Brief Noninvasive Vagal Nerve Stimulation on EEG and Event-Related Potentials.

Neuromodulation, *22*(5), 564–572. https://doi.org/10.1111/ner.12864

Li, J., & Ettinger, A. (2023). Heuristic interpretation as rational inference: A computational

model of the N400 and P600 in language processing. *Cognition*, *233*, 105359.

https://doi.org/10.1016/j.cognition.2022.105359

Linden, D. E. J. (2005). The P300: Where in the brain is it produced and what does it tell us?

Neuroscientist, *11*(6), 563–576. https://doi.org/10.1177/1073858405280524

Liu, Y., Rodenkirch, C., Moskowitz, N., Schriver, B., & Wang, Q. (2017). Dynamic

Lateralization of Pupil Dilation Evoked by Locus Coeruleus Activation Results from

Sympathetic, Not Parasympathetic, Contributions. *Cell Reports*, *20*(13), 3099–3112. https://doi.org/10.1016/j.celrep.2017.08.094

Lloyd, B., Wurm, F., De Kleijn, R., & Nieuwenhuis, S. (2023). Short-term transcutaneous vagus nerve stimulation increases pupil size but does not affect EEG alpha power: A replication of Sharon et al. (2021, Journal of Neuroscience). Brain Stimulation, 16(4), 1001-1008. https://doi.org/10.1016/j.brs.2023.06.010

LoTemplio, S., Silcox, J., Federmeier, K. D., & Payne, B. R. (2020). Inter- and intra-individual coupling between pupillary, electrophysiological, and behavioral responses in a visual oddball task. *Psychophysiology*, *June*, 1–14. https://doi.org/10.1111/psyp.13758

Luck, S. J. (2014). *An Introduction to the Event-Related Potential Technique* (2nd edition). The MIT Press.

Luckey, A.M., Adcock, K., & Vanneste, S. (2023). Peripheral nerve stimulation: A neuromodulation-based approach. *Neuroscience & Biobehavioral Reviews, 149*, 105180. https://doi.org/ 10.1016/j.neubiorev.2023.105180

Ludwig, M., Düzel, E., Betts, M. J., & Hämmerer, D. (2023). Higher pupillary dilation during phasic transcutaneous vagus nerve stimulation (taVNS) and its relevance to the noradrenergic system of the locus coeruleus (LC☐NE). *Alzheimer's & Dementia*, *19*. https://doi.org/10.1002/alz.082818

Ludwig, M., Betts, M.J., & Hämmerer, D. (2025). Stimulate to Remember? The Effects of Short Burst of Transcutaneous Vagus Nerve Stimulation (taVNS) on Memory Performance and Pupil Dilation. Psychophysiology, 62(1), e14753. https://doi.org/10.1111/psyp.14753

Mather, M., Clewett, D., Sakaki, M., & Harley, C. W. (2016). Norepinephrine ignites local hotspots of neuronal excitation: How arousal amplifies selectivity in perception and

memory. *Behavioral and Brain Sciences*, *39*(2016).

https://doi.org/10.1017/S0140525X15000667

Mather, M., Huang, R., Clewett, D., Nielsen, S. E., Velasco, R., Tu, K., Han, S., & Kennedy,

B. L. (2020). Isometric exercise facilitates attention to salient events in women via the

noradrenergic system. *NeuroImage*, *210*, 116560.

https://doi.org/10.1016/j.neuroimage.2020.116560

Mckinnon, A. I., Gray, N. S., & Snowden, R. J. (2020). Enhanced emotional response to both

negative and positive images in post-traumatic stress disorder: Evidence from

pupillometry. *Biological Psychology*, *154*, 107922.

https://doi.org/10.1016/j.biopsycho.2020.107922

McLean, D., Renoult, L., & Malcolm, G. L. (2020). Predictability facilitates rapid scene

processing: A behavioral and ERP study. *Vision Sciences Society Annual Meeting*,

20(11). https://doi.org/10.1167/jov.20.11.1189

Menicucci, D., Animali, S., Malloggi, E., Gemignani, A., Bonanni, E., Fornai, F., Giorgi, F.,

& Binda, P. (2024). Correlated P300b and phasic pupil-dilation responses to

motivationally significant stimuli. *Psychophysiology*, *61*(6), 1–14.

https://doi.org/10.1111/psyp.14550

Mitzdorf, U. (1994). Properties of Cortical Generators of Event-Related Potentials.

Pharmacopsychiatry, *27*(02), 49–51. https://doi.org/10.1055/s-2007-1014274

Mridha, Z., de Gee, J. W., Shi, Y., Alkashgari, R., Williams, J., Suminski, A., Ward, M. P.,

Zhang, W., & McGinley, M. J. (2021). Graded recruitment of pupil-linked

neuromodulation by parametric stimulation of the vagus nerve. *Nature

Communications*, *12*(1), 1–14. https://doi.org/10.1038/s41467-021-21730-2

Münte, T. F., Heinze, H. J., Matzke, M., Wieringa, B. M., & Johannes, S. (1998). Brain

potentials and syntactic violations revisited: No evidence for specificity of the

syntactic positive shift. *Neuropsychologia*, *36*(3), 217–226.

https://doi.org/10.1016/S0028-3932(97)00119-X

Murphy, P. R., O'Connell, R. G., O'Sullivan, M., Robertson, I. H., & Balsters, J. H. (2014).

Pupil diameter covaries with BOLD activity in human locus coeruleus. *Human Brain*

Mapping, *35*(8), 4140–4154. https://doi.org/10.1002/hbm.22466

Murphy, P. R., Robertson, I. H., Balsters, J. H., & O'Connell, R. G. (2011). Pupillometry and

P3 index the locus coeruleus-noradrenergic arousal function in humans.

Psychophysiology, *48*(11), 1532–1543. https://doi.org/10.1111/j.1469-

8986.2011.01226.x

Nater, U. M., & Rohleder, N. (2009). Salivary alpha-amylase as a non-invasive biomarker for

the sympathetic nervous system: Current state of research.

Psychoneuroendocrinology, *34*(4), 486–496.

https://doi.org/10.1016/j.psyneuen.2009.01.014

Neville, H. J., Kutas, M., Chesney, G., & Schmidt, A. L. (1986). Event-related brain

potentials during initial encoding and recognition memory of congruous and

incongruous words. *Journal of Memory and Language*, *25*(1), 75–92.

https://doi.org/10.1016/0749-596X(86)90022-7

Nicenboim, B., Vasishth, S., & Rösler, F. (2020). Are words pre-activated probabilistically

during sentence comprehension? Evidence from new data and a bayesian random-

effects meta-analysis using publicly available data. *Neuropsychologia*, *142*, 107427.

https://doi.org/10.1016/j.neuropsychologia.2020.107427

Nielsen, S. E., & Mather, M. (2015). Comparison of two isometric handgrip protocols on

sympathetic arousal in women. *Physiology & Behavior*, *142*, 5–13.

https://doi.org/10.1016/j.physbeh.2015.01.031

Nieuwenhuis, S., Aston-Jones, G., & Cohen, J. D. (2005). Decision making, the P3, and the locus coeruleus-norepinephrine system. *Psychological Bulletin*, *131*(4), 510–532. https://doi.org/10.1037/0033-2909.131.4.510

Nieuwenhuis, S., De Geus, E. J., & Aston-Jones, G. (2011). The anatomical and functional relationship between the P3 and autonomic components of the orienting response. *Psychophysiology*, *48*(2), 162–175. https://doi.org/10.1111/j.1469-8986.2010.01057.x

Núñez-Peña, M. I., & Honrubia-Serrano, M. L. (2004). P600 related to rule violation in an arithmetic task. *Cognitive Brain Research*, *18*(2), 130–141. https://doi.org/10.1016/j.cogbrainres.2003.09.010

O'Connell, R. G., Dockree, P. M., & Kelly, S. P. (2012). A supramodal accumulation-to-bound signal that determines perceptual decisions in humans. *Nature Neuroscience*, *15*(12), 1729–1735. https://doi.org/10.1038/nn.3248

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Osterhout, L., Holcomb, P. J., & Swinney, D. A. (1994). Brain Potentials Elicited by Garden-Path Sentences: Evidence of the Application of Verb Information During Parsing. *Journal of Experimental Psychology: Learning, Memory, and Cognition*, *20*(4), 786–803. https://doi.org/10.1037/0278-7393.20.4.786

Osterhout, L., McKinnon, R., Bersick, M., & Corey, V. (1996). On the language specificity of the brain response to syntactic anomalies: Is the syntactic positive shift a member of the P300 family? *Journal of Cognitive Neuroscience*, *8*(6), 507–526. https://doi.org/10.1162/jocn.1996.8.6.507

Osterhout, L., & Mobley, L. A. (1995). Event-Related Brain Potentials Elicited by Failure to Agree. *Journal of Memory and Language*, *34*(6), 739–773. https://doi.org/10.1006/jmla.1995.1033

Paller, K. A., McCarthy, G., & Wood, C. C. (1988). ERPs predictive of subsequent recall and recognition performance. *Biological Psychology*, *26*(1–3), 269–276. https://doi.org/10.1016/0301-0511(88)90023-3

Patel, A. D., Gibson, E., Ratner, J., Besson, M., & Holcomb, P. J. (1998). Processing Syntactic Relations in Language and Music: An Event-Related Potential Study. *Journal of Cognitive Neuroscience*, *10*(6), 717–733. https://doi.org/10.1162/089892998563121

Pervaz, I., Thurn, L., Vezzani, C., Kaluza, L., Kühnel, A., & Kroemer, N. B. (2025). Does transcutaneous auricular vagus nerve stimulation alter pupil dilation? A living Bayesian meta-analysis. *Brain Stimulation*, *18*(2), 148–157. https://doi.org/10.1016/j.brs.2025.01.022

Peuker, E. T., & Filler, T. J. (2002). The nerve supply of the human auricle. *Clinical Anatomy (New York, N.Y.)*, *15*(1), 35–37. https://doi.org/10.1002/ca.1089

Pineda, J. A., Foote, S. L., & Neville, H. J. (1989). Effects of locus coeruleus lesions on auditory, long-latency, event-related potentials in monkey. *Journal of Neuroscience*, *9*(1), 81–93. https://doi.org/10.1523/jneurosci.09-01-00081.1989

Pineda, J. A., & Westerfield, M. (1993). Monkey P3 in an "oddball" paradigm: Pharmacological support for multiple neural sources. *Brain Research Bulletin*, *31*(6), 689–696. https://doi.org/10.1016/0361-9230(93)90142-X

Pion-Tonachini, L., Kreutz-Delgado, K., & Makeig, S. (2019). ICLabel: An automated electroencephalographic independent component classifier, dataset, and website. *NeuroImage*, *198*(3), 181–197. https://doi.org/10.1016/j.neuroimage.2019.05.026

Polich, J. (2007). Updating P300: An integrative theory of P3a and P3b. *Clinical Neurophysiology*, *118*(10), 2128–2148. https://doi.org/10.1016/j.clinph.2007.04.019

Pritchard, W. S. (1981). Psychophysiology of P300. *Psychological Bulletin*, *89*(3), 506–540. https://doi.org/10.1037/0033-2909.89.3.506

Rabovsky, M., & McClelland, J. L. (2020). Quasi-compositional mapping from form to

    meaning: A neural network-based approach to capturing neural responses during

    human language comprehension. *Philosophical Transactions of the Royal Society of*

    London, Series B, Biological Sciences, 375, 20190313.

    https://doi.org/10.1098/rstb.2019.0313,

Rajkowski, J., Kubiak, P., & Aston-Jones, G. (1993). Correlations between locus coeruleus

    (LC) neural activity, pupil diameter and behavior in monkey support a role of LC in

    attention. *Proc. Soc. Neurosci.*, *19*, 974.

Regel, S., Meyer, L., & Gunter, T. C. (2014). Distinguishing neurocognitive processes

    reflected by P600 effects: Evidence from ERPs and neural oscillations. *PLoS ONE*,

    9(5), 1–11. https://doi.org/10.1371/journal.pone.0096840

Reimer, J., McGinley, M. J., Liu, Y., Rodenkirch, C., Wang, Q., McCormick, D. A., & Tolias,

    A. S. (2016). Pupil fluctuations track rapid changes in adrenergic and cholinergic

    activity in cortex. *Nature Communications*, *7*(May), 1–7.

    https://doi.org/10.1038/ncomms13289

Rufener, K. S., Geyer, U., Janitzky, K., Heinze, H. J., & Zaehle, T. (2018). Modulating

    auditory selective attention by non-invasive brain stimulation: Differential effects of

    transcutaneous vagal nerve stimulation and transcranial random noise stimulation.

    European Journal of Neuroscience, *48*(6), 2301–2309.

    https://doi.org/10.1111/ejn.14128

Ryskin, R., Stearns, L., Bergen, L., Eddy, M., Fedorenko, E., & Gibson, E. (2021). An ERP

    index of real-time error correction within a noisy-channel framework of human

    communication. *Neuropsychologia*, *158*, 107855.

    https://doi.org/10.1016/j.neuropsychologia.2021.107855

Sara, S. J. (2009). The locus coeruleus and noradrenergic modulation of cognition. *Nature*

    Reviews Neuroscience, *10*(3), 211–223. https://doi.org/10.1038/nrn2573

Sassenhagen, J., & Bornkessel-Schlesewsky, I. (2015). The P600 as a correlate of ventral attention network reorientation. *Cortex*, *66*, A3–A20. https://doi.org/10.1016/j.cortex.2014.12.019

Sassenhagen, J., & Fiebach, C. J. (2019). Finding the P3 in the P600: Decoding shared neural mechanisms of responses to syntactic violations and oddball targets. *NeuroImage*, *200*, 425–436. https://doi.org/10.1016/j.neuroimage.2019.06.048

Sassenhagen, J., Schlesewsky, M., & Bornkessel-Schlesewsky, I. (2014). The P600-as-P3 hypothesis revisited: Single-trial analyses reveal that the late EEG positivity following linguistically deviant material is reaction time aligned. *Brain and Language*, *137*, 29–39. https://doi.org/10.1016/j.bandl.2014.07.010

Schad, D. J., Vasishth, S., Hohenstein, S., & Kliegl, R. (2018). How to capitalize on a priori contrasts in linear (mixed) models: A tutorial. *Journal of Memory and Language*, *110*, 104038. https://doi.org/10.1016/j.jml.2019.104038

Schneider, C., Priovoulos, N., Prokopiou, P. C., Verhey, F. R., Poser, B. A., Ivanov, D., ... & Jacobs, H. I. (2025). Locus coeruleus fMRI response to transcutaneous auricular vagus nerve stimulation is coupled to changes in salivary alpha amylase. *Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation*, *18*(4), 1205 - 1207. https://doi.org/10.1016/j.brs.2025.06.014

Sharon, O., Fahoum, F., & Nir, Y. (2021). Transcutaneous Vagus Nerve Stimulation in Humans Induces Pupil Dilation and Attenuates Alpha Oscillations. *The Journal of Neuroscience*, *41*(2), 320–330. https://doi.org/10.1523/JNEUROSCI.1361-20.2020

Sitnikova, T., Holcomb, P. J., Kiyonaga, K. A., & Kuperberg, G. R. (2008). Two neurocognitive mechanisms of semantic integration during the comprehension of visual real-world events. *Journal of Cognitive Neuroscience*, *20*(11), 2037–2057. https://doi.org/10.1162/jocn.2008.20143

Skora, L., Marzecová, A., & Gerhard, J. (2024). Tonic and phasic transcutaneous auricular vagus nerve stimulation (taVNS) both evoke rapid and transient pupil dilation. *Brain Stimulation, 17*(2), 233-244. https://doi.org/10.1016/j.brs.2024.02.013

Sterpenich, V., D'Argembeau, A., Desseilles, M., Balteau, E., Albouy, G., Vandewalle, G., Degueldre, C., Luxen, A., Collette, F., & Maquet, P. (2006). The locus ceruleus is involved in the successful retrieval of emotional memories in humans. *Journal of Neuroscience*, *26*(28), 7416–7423. https://doi.org/10.1523/JNEUROSCI.1001-06.2006

Swick, D., Pineda, J. A., Schacher, S., & Foote, S. L. (1994). Locus coeruleus neuronal activity in awake monkeys: Relationship to auditory P300-like potentials and spontaneous EEG. *Experimental Brain Research*, *101*(1). https://doi.org/10.1007/BF00243219

Szeska, C., Klepzig, K., Hamm, A. O., & Weymar, M. (2025). Ready for translation: non-invasive auricular vagus nerve stimulation inhibits psychophysiological indices of stimulus-specific fear and facilitates responding to repeated exposure in phobic individuals. *Translational Psychiatry*, *15*(1), 135. https://doi.org/10.1038/s41398-025-03352-0

Tan, G., Huguenard, A. L., Donovan, K. M., Demarest, P., Liu, X., Li, Z., Adamek, M., Lavine, K., Vellimana, A. K., Kummer, T. T., Osbun, J. W., Zipfel, G. J., Brunner, P., & Leuthardt, E. C. (2025). The effect of transcutaneous auricular vagus nerve stimulation on cardiovascular function in subarachnoid hemorrhage patients: A randomized trial. *eLife*, *13*, RP100088. https://doi.org/10.7554/eLife.100088

The MathWorks Inc. (2022). MATLAB (R2020a), Natick, Massachusetts: The MathWorks Inc. https://www.mathworks.com

Trueswell, J. C., Medina, T. N., Hafri, A., & Gleitman, L. R. (2013). Propose but verify: Fast mapping meets cross-situational word learning. *Cognitive Psychology*, *66*(1), 126–156. https://doi.org/10.1016/j.cogpsych.2012.10.001

Van Boekholdt, L., Kerstens, S., Khatoun, A., Asamoah, B., & Mc Laughlin, M. (2021). tDCS peripheral nerve stimulation: a neglected mode of action? *Molecular Psychiatry, 26*(29), 456-461. https://doi.org/ 0.1038/s41380-020-00962-6

van de Meerendonk, N., Kolk, H. H. J., Chwilla, D. J., & Vissers, C. T. W. M. (2009). Monitoring in language perception. *Linguistics and Language Compass*, *3*(5), 1211–1224. https://doi.org/10.1111/j.1749-818X.2009.00163.x

van de Meerendonk, N., Kolk, H. H. J., Vissers, C. T. W. M., & Chwilla, D. J. (2010). Monitoring in language perception: Mild and strong conflicts elicit different ERP patterns. *Journal of Cognitive Neuroscience*, *22*(1), 67–82. https://doi.org/10.1162/jocn.2008.21170

Van Herten, M., Kolk, H. H. J., & Chwilla, D. J. (2005). An ERP study of P600 effects elicited by semantic anomalies. *Cognitive Brain Research*, *22*(2), 241–255. https://doi.org/10.1016/j.cogbrainres.2004.09.002

Vazey, E. M., Moorman, D. E., & Aston-Jones, G. (2018). Phasic locus coeruleus activity regulates cortical encoding of salience information. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(40), E9439–E9448. https://doi.org/10.1073/pnas.1803716115

Ventura□Bort, C., Giraudier, M., & Weymar, M. (2025). Transcutaneous Auricular Vagus Nerve Stimulation Enhances Emotional Processing and Long□Term Recognition Memory: Electrophysiological Evidence Across Two Studies. *Psychophysiology*, *62*(3), e70034. https://doi.org/10.1111/psyp.70034

Ventura-Bort, C., Wirkner, J., Genheimer, H., Wendt, J., Hamm, A. O., & Weymar, M. (2018). Effects of Transcutaneous Vagus Nerve Stimulation (tVNS) on the P300 and Alpha-Amylase Level: A Pilot Study. *Frontiers in Human Neuroscience*, *12*(202), 1–12. https://doi.org/10.3389/fnhum.2018.00202

Ventura-Bort, C., Wirkner, J., Wendt, J., Hamm, A. O., & Weymar, M. (2021). Establishment of Emotional Memories Is Mediated by Vagal Nerve Activation: Evidence from Noninvasive taVNS. *The Journal of Neuroscience: The Official Journal of the Society for Neuroscience*, *41*(36), 7636–7648. https://doi.org/10.1523/JNEUROSCI.2329-20.2021

Verleger, R. (2020). Effects of relevance and response frequency on P3b amplitudes: Review of findings and comparison of hypotheses about the process reflected by P3b. *Psychophysiology*, *57*(7), 1–22. https://doi.org/10.1111/psyp.13542

Villani, V., Finotti, G., Di Lernia, D., Tsakiris, M., & Azevedo, R. T. (2022). Event-related transcutaneous vagus nerve stimulation modulates behaviour and pupillary responses during an auditory oddball task. *Psychoneuroendocrinology*, *140*, 105719. https://doi.org/10.1016/j.psyneuen.2022.105719

Vissers, C. T. W. M., Chwilla, D. J., & Kolk, H. H. J. (2007). The interplay of heuristics and parsing routines in sentence comprehension: Evidence from ERPs and reaction times. *Biological Psychology*, *75*(1), 8–18. https://doi.org/10.1016/j.biopsycho.2006.10.004

Võ, M. L. H., & Wolfe, J. M. (2013). Differential Electrophysiological Signatures of Semantic and Syntactic Scene Processing. *Psychological Science*, *24*(9), 1816–1823. https://doi.org/10.1177/0956797613476955

Warren, C. M., Tona, K. D., Ouwerkerk, L., van Paridon, J., Poletiek, F., van Steenbergen, H., Bosch, J. A., & Nieuwenhuis, S. (2019). The neuromodulatory and hormonal effects of transcutaneous vagus nerve stimulation as evidenced by salivary alpha amylase, salivary cortisol, pupil diameter, and the P3 event-related potential. *Brain Stimulation*, *12*(3), 635–642. https://doi.org/10.1016/j.brs.2018.12.224

Warren, C. V., Maraver, M. J., de Luca, A., & Kopp, B. (2020). The effect of transcutaneous auricular vagal nerve stimulation (taVNS) on P3 event□related potentials during a

Bayesian oddball task. *Brain Sciences*, *10*(6), 1–20.

https://doi.org/10.3390/brainsci10060404

Xu, J., Abdel Rahman, R., & Sommer, W. (2021). Sequential adaptation effects reveal

proactive control in processing spoken sentences: Evidence from event-related

potentials. *Brain and Language*, *214*, 104904.

https://doi.org/10.1016/j.bandl.2020.104904

Yakunina, N., Kim, S. S., & Nam, E.-C. (2017). Optimization of Transcutaneous Vagus

Nerve Stimulation Using Functional MRI. *Neuromodulation: Journal of the

International Neuromodulation Society*, *20*(3), 290–300.

https://doi.org/10.1111/ner.12541