Constraints on paternal care capacity favor a monogamous mating
system in the group-living gobiid fish *Trimma marinae*

Kota Kambe[1,2#] · Kazuya Fukuda[3,4#*] · Tomoki Sunobe[3]

[1]School of Marine Biosciences, Kitasato University, Sagamihara, Kanagawa 252-0373,

Japan

[2]Current address: Laboratory of Animal Sociology, Department of Biology, Graduate

School of Science, Osaka Metropolitan University, Osaka 558-0022, Japan.

[3]Laboratory of Fish Behavioral Ecology, Tateyama Station, Field Science Center, Tokyo

University of Marine Science and Technology, Tateyama, Chiba 294-0308, Japan

[4]Current address: School of Marine Biosciences, Kitasato University, Sagamihara,

Kanagawa 252-0373, Japan

[#]Equal contribution

*Correspondence: Kazuya Fukuda (E-mail: fukuda.kazuya@kitasato-u.ac.jp, Tel:

+81-42-778-8009). 1-15-1 Kitazato, Minami-ku, Sagamihara, Kanagawa 252-0373,

Japan

ORCiD ID

Kota Kambe: 0009-0006-1931-3631

Kazuya Fukuda: 0000-0002-4355-1545

Tomoki Sunobe: 0000-0001-7137-3726

Total text pages: 8

Numbers of figures: 1

Abstract

We examined why the group-living gobiid fish *Trimma marinae* is monogamous despite

a high potential for polygamy. Our observations revealed that in 75.5% of spawning

events within established pairs, females spawned before the previous clutch hatched,

forcing males to care for two clutches simultaneously. Previous findings showed males

can care for up to three clutches, while mating with a second female would likely require

them to care for four. Our results indicate that female mate guarding is a strategy to

monopolize a male's limited paternal capacity by preventing her partner from mating with

additional females.

Keywords

Mating system, Monogamy, Mate guarding, paternal care, gobiid fish, *Trimma marinae*

Introduction

Monogamy is a mating system in which males and females typically mate with a single

partner, often maintaining this relationship beyond one reproductive cycle; given the

difficulty in defining monogamy unambiguously across animal lineages (Kvarnemo

2018; Bales et al. 2021), this study adopts the definition traditionally used in fish

behavioral ecology, following Barlow (1988) and Whiteman and Côté (2004). The

ecological conditions under which monogamy is evolutionarily stable have long been

investigated (e.g., Emlen and Oring 1977; Shuster and Wade 2003). Even when

focusing on marine fish, at least six ecological factors have been proposed to facilitate

the evolution of monogamy: biparental care, habitat limitations, low population

density/low mate availability/low mobility, increased reproductive efficiency, territorial

defense, and the net benefit of sequestering a single mate (Whiteman and Côté 2004).

Because there is a great diversity of marine fish, accumulating knowledge of the

ecological features of monogamous species is expected to strengthen existing theories

and generate new hypotheses.

Trimma marinae is a small (approx. 25 mm TL), group-living gobiid fish

found on coral reef slopes in enclosed bays of the Western Pacific (Shibukawa 2004).

The framework of Emlen and Oring (1977) predicts that species in environments with

a high potential for acquiring additional and monopolizable mates will exhibit a high

environmental potential for polygamy. Based on this framework, *T. marinae* is
predicted to favor the evolution of a polygamous mating system because it inhabits
groups where potential mates are spatially clustered. However, contrary to this
prediction, Fukuda et al. (2017a) revealed that *T. marinae* has a monogamous mating
system. They reported that this mating system primarily results from intrasexual
competition among females (i.e., mate guarding) and suggested that its adaptive
significance lies in monopolizing the limited paternal care capacity of males. By
preventing her partner from mating with other females, a female should be able to avoid
sharing the male's limited parental investment, thereby increasing her reproductive
success. This hypothesis is supported by occasional observations from Fukuda et al.
(2017a) showing that while males could care for up to three clutches, they failed to care
for all of them successfully when simultaneously guarding four clutches, with some
eggs decomposing or being cannibalized. These observations suggest that the paternal
care capacity of *T. marinae* males is constrained. However, that study did not
quantitatively investigate whether a male's care capacity is already saturated by the
reproductive output of a single female partner during continuous reproduction within
typical monogamous pairs.
To address this question, the present study investigated the spawning interval
of continuous pairs under laboratory conditions. Our aim was to determine the number
of clutches a male must typically care for simultaneously, and thereby examine the
hypothesis that females obtain benefits from mate guarding.
82
Materials and methods
The data used in this study were obtained from a rearing observation previously reported
by Fukuda et al. (2017a). The materials and methods for specimen collection and rearing
conditions, which are identical to the original report, are briefly summarized below. A
single group of 25 individuals (8 males, 17 females) was collected by SCUBA at Amami
Oshima, Kagoshima Prefecture, Japan, and transported to the laboratory. Each fish was
subcutaneously tagged for individual identification and housed in a $120 \times 45 \times 45$ cm
tank at 25°C. Ten polyvinyl chloride pipes were halved as spawning nests. Reproductive
behavior was observed between 04:00 and 12:00 for 37 consecutive days. At the rearing
temperature of 25°C, the eggs of *T. marinae* hatch on the third day post-fertilization, with

hatching triggered by it becoming dark (sunset).

Following the criteria of Fukuda et al. (2017a), a male–female pair was

considered an established monogamous bond if pairing behavior was observed for more

than 2 days. As *T. marinae* typically reproduces in monogamous pairs, we recorded

spawning intervals exclusively for females in these established pairs. To clarify

representative spawning intervals, we analyzed only females that spawned at least twice

and recorded the number of days between their successive spawning events.

Results

A total of 49 spawning intervals were recorded from established pairs, with a median

interval of 2 days; this means that females typically spawned approximately 48 hours

after the previous spawning (IQR = 1–3 days). As eggs of *T. marinae* hatch on the third

day post-fertilization, this reproductive pace indicates that males may be required to care

for two clutches simultaneously. The full frequency distribution of the spawning intervals,

which ranged from 1 to 13 days, is shown in Fig. 1. Of the 49 total recorded intervals, 37

(75.5%) were 3 days or less (i.e., intervals of 1, 2, or 3 days); this reproductive pace

requires males to care for at least two clutches simultaneously. Similarly, 26 of these

intervals (53.1%) were 2 days or less (requiring care for three or more clutches), and 12

(24.5%) were just 1 day (requiring care for four clutches).

Discussion

This study demonstrated that in 75.5% of reproductive events within monogamous *T.*

marinae pairs, females spawn their next clutch before the previous one hatches. This

indicates that males routinely face opportunities to care for two clutches simultaneously

during a typical reproductive cycle, even when paired with a single partner. This

situation suggests that mating with a second female would create a high probability of

the male being required to care for four clutches, a number exceeding the male's

paternal care capacity. Furthermore, in 53.1% of reproductive events, the spawning

pace required males to care for three clutches simultaneously, indicating that a single

female's reproductive cycle is already sufficient to saturate a male's paternal care

capacity. Therefore, to ensure that their own eggs receive sufficient care until hatching,

it would be adaptive for a female to prevent her partner from acquiring additional mates.

This female behavior may explain the evolution of a monogamous mating system in *T. marinae,* despite their high environmental potential for polygamy. The present findings offer support for the hypothesis proposed by Fukuda et al. (2017a). This species forms aggregations in which the sex ratio is biased toward females (Fukuda et al. 2017). Such a female-biased sex ratio, and the resulting female-skewed operational sex ratio (OSR), may have led to conditions that favored the evolution of female mate guarding and subsequent monogamy in *T. marinae*. Indeed, in the monogamous pipefish, *Corythoichthys haematopterus*, which also exhibits a female-biased sex ratio, female mate guarding is suggested to contribute to the maintenance of monogamy (Matsumoto and Yanagisawa 2001). This behavior is considered to be promoted by the skewed OSR (Sogabe and Yanagisawa 2007).

136   One possible factor contributing to the limited paternal care capacity of *T. marinae* is that males are smaller than females (Fukuda et al. 2017a). This pattern contrasts with that of the polygynous *Trimma* species (e.g., *T. okinawae* [Sunobe and Nakazono 1990], *T. grammistes* [Fukuda et al. 2017b], *T. caudomaculatum* [Tomatsu et al. 2018]). In these species, the larger male body size likely enables the care of multiple clutches from several females simultaneously, thus leaving females with little need to compete for paternal care. Indeed, in *T. caudomaculatum*, it has been observed that three females spawned simultaneously in a single male's nest without competition (Tomatsu et al. 2018). In *T. marinae*, however, the smaller male body size likely constrains their capacity to care for the clutches produced by the relatively larger females. This constraint is thought to drive intrasexual competition among females to monopolize a male's parental care, favoring the evolution of monogamy. The case of *T. taylori*, the only other *Trimma* species reported with a similar sexual size dimorphism to *T. marinae*, lends further support to this hypothesis; it is also suggested to be monogamous due to female competition over limited paternal care (Oyama et al. 2023). This pattern suggests that a small male body size relative to females has been a potential factor in the evolution of monogamy, at least within the *Trimma* lineage. Cases suggesting that constraints on paternal care induce female mate guarding, ultimately leading to the evolution of monogamous mating systems, have been previously reported. For example, in the coral-dwelling goby *Paragobiodon xanthosomus*, female intrasexual competition driven by limitations in paternal care is suggested to be a key

factor in the establishment of its monogamous mating system (Wong et al. 2008). This
process is similar to the one proposed for *T. marinae*. Given that parental care is
commonly provided only by males in gobiid fishes, further studies may identify
additional species exhibiting monogamy due to similar selective pressures, even in
environments with a high potential for polygyny.

Acknowledgments
We are grateful to the members of the Laboratory of Fish Behavioral Ecology, Tokyo
University of Marine Science and Technology for their advice and support during our
study. Thanks are also due to S. Yokoyama for their assistance during the study.

Author contributions
Conceptualization, Methodology: KF, TS; Investigation: KF; Formal analysis: KK;
Writing - original draft preparation: KK; Writing - review and editing: KF, TS; Funding
acquisition: TS; Resources: KF, TS; Supervision: TS.

Funding
This work was supported by JSPS KAKENHI Grant Numbers 19570016, 24370006.

Data availability
The datasets are available from the corresponding author on reasonable request.

Declarations
Conflict of interest
The authors declare that they have no conflicts of interest.

Ethical approval
At Tokyo University of Marine Science and Technology, there are no institutional ethical
regulations for fish. Therefore, all procedures adhered to the ASAB/ABS guidelines for
the treatment of animals in behavioral research.

Reference

Barlow GW (1988) Monogamy in relation to resource. In: Slobodchikoff CN (ed) The

ecology of social behavior. Academic Press, New York, pp 55–79

Bales KL, Ardekani CS, Baxter A, Karaskiewicz CL, Kuske JX, Lau AR, Savidge LE,

Sayler KR, Witczak LR (2021) What is a pair bond?. Horm Behav 136:105062.

Emlen ST, Oring LW (1977) Ecology, sexual selection, and the evolution of mating

system. Science 197:215–223

Fukuda K, Manabe H, Sakurai M, Dewa S, Shinomiya A, Sunobe T (2017a) Monogamous

mating system and sexuality in the gobiid fish, *Trimma marinae* (Actinopterygii:

Gobiidae). J Ethol 35:121–130

Fukuda K, Tanazawa T, Sunobe T (2017b) Polygynous mating system and field evidence

for bidirectional sex change in the gobiid fish *Trimma grammistes*. Inter J Pure Appl

Zool 5:92–99

Kvarnemo C (2018) Why do some animals mate with one partner rather than many? A

review of causes and consequences of monogamy. Biol Rev 93:1795–1812

Matsumoto K, Yanagisawa Y (2001) Monogamy and sex role reversal in the pipefish

Corythoichthys haematopterus. Anim Behav 61:163–170

Oyama T, Tomatsu S, Manabe H, Sakurai M, Matsuoka M, Shinomiya A, Dewa S, Sunobe

T (2023) Monogamous mating system and protandrous-like sexuality in the goby

Trimma taylori. Ichthyol Res 70:287–292

Shibukawa K (2004) Genus Trimma. In: Senou H (ed) A photographic guide to the gobiid

fishes of Japan. Heibonsha, Tokyo, pp 96–117

Shuster SM, Wade MJ (2003) Mating systems and strategies. Princeton University Press.

Sogabe A, Yanagisawa Y (2007) Sex-role reversal of a monogamous pipefish without

higher potential reproductive rate in females. Proc R Soc B 274:2959–2963

Sunobe T, Nakazono A (1990) Polygynous mating system of *Trimma okinawae* (Pisces:

Gobiidae) at Kagoshima, Japan with a note on sex change. Ethology 84:133–143

Tomatsu S, Ogiso K, Fukuda K, Deki M, Dewa S, Manabe H, Sakurai M, Shinomiya A,

Sunobe T (2018) Multi-male group and bidirectional sex change in the gobiid fish,

Trimma caudomaculatum. Ichthyol Res 65:502–506

Whiteman EA, Côté IM (2004) Monogamy in marine fishes. Biol Rev 79:351–375

Wong MYL, Munday PL, Buston PM, Jones GP (2008) Monogamy when there is

potential for polygyny: tests of multiple hypotheses in a group-living fish. Behav

Ecol 19:353–361

Figure legend

Fig1. Histogram of the spawning intervals distribution (days). The x-axis indicates the

number of days between spawning events within monogamous pairs