Evolutionary origin and functional mechanism of Lhcx in the diatom photoprotection

Minoru Kumazawa[1,2], Seiji Akimoto[3], Atsushi Takabayashi[2], Ko Imaizumi[1], Shoko Tsuji[1], Hazuki

Hasegawa[4], Atsushi Sakurai[4], Sousuke Imamura[4], Noriko Ishikawa[1], Natsuko Inoue-Kashino[5],

Yasuhiro Kashino[5] & Kentaro Ifuku[1]

[1]Graduate School of Agriculture, Kyoto University, Japan

[2]Institute of Low Temperature Science, Hokkaido University, Japan

[3]Graduate School of Science, Kobe University, Japan

[4]Space Environment and Energy Laboratories, NTT, Inc., Japan

[5]Graduate School of Science, University of Hyogo, Japan

Author for correspondence: Kentaro Ifuku (ifuku.kentaro.2m@kyoto-u.ac.jp)

Summary

•   Diatoms, a group of red-lineage algae that possess plastids derived from red algae through

secondary endosymbiosis, utilize the light-harvesting complex (LHC) subfamily Lhcx for

photoprotection; however, its evolutionary origin and functional mechanism remain unclear.

•   To address these questions, we performed molecular phylogenetic analysis of Lhcx/Lhcsr

subfamilies and produced the *Lhcx1* knockout mutants of the diatom *Chaetoceros gracilis*, which

has low Lhcx redundancy.

•   Phylogenetic analysis revealed that diatom Lhcxs and the green algal Lhcsrs evolved from a

common ancestor, with green plants acquiring them through horizontal gene transfer. The *Lhcx1*

mutants of *C. gracilis* mostly abolished non-photochemical quenching (NPQ), and the time-

resolved fluorescence measurements clarified that Lhcx1-mediated quenching occurs in the

energetically detached antenna complexes. Consistently, clear-native PAGE using Amphipol

suggested that CgLhcx1 would interact with FCP L-dimer, functioning as a peripheral antenna for

$C_2S_2M_2$ PSII–FCPII. Notably, the *lhcx1* mutant exhibited a higher PSII effective quantum yield

than wild-type (WT) under high-light acclimation, attributed to reduced antenna size and

enhanced carbon fixation capacity. The lack of NPQ accelerated high-light acclimation, with

increased xanthophylls in *lhcx1*, indicating that compensatory mechanisms can lead to enhanced

photosynthetic efficiency.

•   Our findings provide novel insights into the origin and molecular mechanisms of photoprotection

in diatoms.

Introduction

Photosynthesis consists of photochemical reactions in which two photosystems connected in series convert light energy into chemical energy. Among photosynthetic organisms, red-lineage secondary endosymbiotic algae, which possess plastids derived from red algae acquired through cellular endosymbiosis, are widely adapted to aquatic environments on Earth (Pierella Karlusich *et al.*, 2020, 2025). Light conditions in these environments are as dynamic as those in terrestrial environments, making it extremely important for photosynthetic organisms to protect their photosystems from excess light to suppress the generation of harmful reactive oxygen species caused by the absorption of excessive light energy. Non-Photochemical Quenching (NPQ), which dissipates excess light energy as heat, plays a crucial role in this protective mechanism (Niyogi, 2000).

Multiple NPQ mechanisms with different induction times exist, and these exhibit distinct characteristics depending on the taxonomic group (Goss & Lepetit, 2015; Malnoë, 2018). In particular, energy-dependent quenching (qE) is a major component of NPQ and is carried out by Light-Harvesting Complex (LHC) proteins (Peers *et al.*, 2009; Goss & Lepetit, 2015; Buck *et al.*, 2019). In red-lineage secondary endosymbiotic algae, qE, dependent on the Lhcx subfamily of the LHC family, serves as the primary NPQ mechanism (Buck *et al.*, 2019; Croteau *et al.*, 2025). Green algae, on the other hand, utilize qE dependent on the Lhcsr subfamily, which is closely related to Lhcx, as well as state transitions (qT), where LHCII detaches from photosystem II (PSII) and binds to photosystem I (PSI) through phosphorylation control to adjust the excitation balance between the two photosystems (Allorent *et al.*, 2013; Minagawa & Tokutsu, 2015). However, the evolutionary origin of Lhcx/Lhcsr has not been extensively discussed due to a lack of comprehensive phylogenetic analyses of LHC subfamilies.

The LHC of diatoms, representative of red-lineage algae, is called FCP (Fucoxanthin-Chlorophyll (Chl) *a*/*c*-binding Protein) because it binds pigments such as Chl *c*, fucoxanthin, and diadinoxanthin

in addition to Chl *a*. Diatom NPQ primarily consists of qE, which depends on the proton gradient

(ΔpH) across the thylakoid membrane formed by light (Lavaud & Kroth, 2006; Goss & Lepetit, 2015).

This qE in diatoms depends on Lhcx, which is homologous to the Lhcsr family essential for qE in

green algae (Koziol *et al.*, 2007; Peers *et al.*, 2009). However, PsbS, which is necessary for qE

induction in land plants, is absent in diatoms. While the homology between Lhcx and Lhcsr amino

acid sequences is well known, differences in the mechanisms that regulate gene expression have been

reported (Petroutsos *et al.*, 2016; Allorent *et al.*, 2016; Tokutsu *et al.*, 2019a,b; Im *et al.*, 2024; Zhang

et al., 2024).

While Lhcx is required for qE induction, xanthophyll pigments are also important for qE induction.

In diatoms, the diadinoxanthin (Ddx) to diatoxanthin (Dtx) conversion cycle (Ddx-Dtx cycle) is the

major xanthophyll cycle, with a strong positive correlation between Dtx amount and qE (Lavaud &

Lepetit, 2013). Additionally, the violaxanthin-antheraxanthin-zeaxanthin (VAZ) cycle, known in land

plants, also functions in diatoms, complementing the Ddx-Dtx cycle (Lohr & Wilhelm, 1999; Giossi

et al., 2025). Furthermore, it has been reported that mitochondria in diatoms function as electron sinks

for the photosynthetic electron transport chain, contributing to the maintenance of normal chloroplast

function (Bailleul *et al.*, 2015).

NPQ in red-lineage algae has been studied in diatoms and *Nannochloropsis* belonging to heterokont

algae (Bailleul *et al.*, 2010; Park *et al.*, 2019; Buck *et al.*, 2019, 2021; Giovagnetti *et al.*, 2022). In

model diatom species, multiple Lhcx isoforms are expressed under different conditions and show

functional redundancy. Therefore, almost complete Lhcx-deficient strains have not been created in

diatoms to date, and the physiological responses and details of excitation energy transfer in states with

significantly deficient NPQ have not been fully elucidated. In contrast, *Chaetoceros gracilis* has low

redundancy of Lhcxs, with primarily only CgLhcx1 being expressed (Kumazawa *et al.*, 2022),

providing the possibility that an almost complete Lhcx-deficient strain can be obtained by creating a

knockout of this CgLhcx1. Furthermore, the three-dimensional structures of PSII–FCPII (Wang *et al.*,

2019; Nagao *et al.*, 2019, 2022) and PSI–FCPI (Nagao *et al.*, 2020a; Xu *et al.*, 2020) supercomplexes

have already been elucidated in this species, establishing a foundation for structural biology research.

This study aims to investigate the origin of Lhcx phylogenetically and to elucidate the mechanism of

its excitation energy quenching for high light (HL) protection by creating Lhcx1-deficient mutants in

C. gracilis using CRISPR/Cas9 genome editing technology.

Results

Origin of LHCs in the Lhcx/Lhcsr Subfamily

To elucidate the origins of the diatom Lhcx and green algal Lhcsr subfamilies, which perform NPQ,

we conducted a comprehensive molecular phylogenetic analysis of light-harvesting complexes

(LHCs) from red algae, green algae, cryptophytes, and diatoms, and represented the estimated

phylogenetic tree as an unrooted tree (Figure 1). Comprehensive retrieval of LHC sequences by

BLAST confirmed that, as in our previous reports, red algae possess only the Lhcr subfamily,

cryptophytes have Lhcr and Lhcz subfamilies, and green algae contain Lhca, Lhcb, and Lhcsr

subfamilies, while diatoms possess Lhcr, Lhcz, Lhcq, CgLhcr9 homolog, Lhcx, and Lhcf subfamilies

(Kumazawa *et al.*, 2022; Kumazawa & Ifuku, 2024). In Figure 1, each LHC subfamily in the red

lineage is separated. However, within the Lhcr subfamily, there exists diversity among the Lhcr

proteins from red algae, cryptophytes, and diatoms, encompassing gene groups with distant

evolutionary relationships.

Lhcx and Lhcsr formed a monophyletic clade with an ultrafast bootstrap value of 100% (Figure 1,

inset). Within this clade, *Thalassiosira pseudonana* Lhcx6_1 and *Cyclotella cryptica* Lhcx6_1,

derived from diatoms belonging to the order Thalassiosirales, clearly branched at the most basal

position and were evolutionarily distant from typical Lhcx and Lhcsr proteins. All Lhcx1–3 proteins

from the diatom *Chaetoceros gracilis* were positioned within the typical Lhcx clade. Additionally, the

Lhcx and Lhcsr clade showed monophyly with the Lhcq, CgLhcr9 homolog, and Lhcf subfamilies,

supported by a high bootstrap value (100%). Within that clade, Lhcq was located at the most basal

position, while the Lhcx/Lhcsr clade was positioned internally. In contrast, the Lhcx/Lhcsr clade did

not show monophyly with the Lhca/Lhcb clade. Furthermore, the Lhcr of red algae and the Lhca and

Lhcb subfamilies, which are responsible for light harvesting in red and green algae possessing primary

plastids, did not form a monophyletic group; between them existed cryptophyte Lhcr, diatom Lhcr,

cryptophyte and diatom Lhcz, as well as other subfamilies of the red evolutionary lineage. These data

suggest that Lhcsr and Lhcx subfamilies are derived from the red-lineage-specific subfamilies.

Creating *Lhcx1* Knockout Strains with Ribozyme-Based Genome Editing

C. gracilis has three typical Lhcx genes (*CgLhcx1–3*): according to the genome database of *C.*

gracilis (ChaetoBase v1.1, https://chaetoceros.nibb.ac.jp/), *CgLhcx1* showed the highest transcription

level by a significant margin (Kumazawa *et al.*, 2022). Therefore, we decided to create an *Lhcx1*-

deficient mutant in *C. gracilis* using the CRISPR/Cas9 system. To facilitate the comprehensive

analysis of LHC genes using genome editing in *C. gracilis,* we have developed the inducible

CRISPR/Cas9 system, in which both Cas9 and guide RNA (gRNA) expression is regulated by the

inducible Pol-II-type promoter. First, we introduced the *Cas9* gene under the nitrate reductase (NR)

promoter into wild-type *C. gracilis* with a vector having the nourseothricin resistance gene (*nat*) via

electroporation (Supplemental Figure 1A). Colonies obtained through antibiotic-resistance selection

were further selected for high Cas9 expression in the presence of nitrate by immunoblotting

(Supplemental Figure 1B). The selected hC9_5 strain showed morphology and growth characteristics

equivalent to the wild-type strain (WT) and was established as the host strain for the genome editing

(Supplemental Figure 1C). Second, to express mature gRNA in the host hC9_5 under the NR promoter,

we employed a method utilizing ribozymes (Gao & Zhao, 2014; Zhang *et al.*, 2017). This method

employed a ribozyme-gRNA-ribozyme (RGR) construct in which a hammerhead ribozyme with self-

cleavage activity at the 3' end was attached to the 5' end of the gRNA, and an HDV ribozyme with

self-cleavage activity at the 5' end was attached to the 3' end of the gRNA. This RGR cassette for the

gene of interest can be easily constructed by overlap PCR and inserted into the pCgNRp[ble] vector,

having an antibiotic-resistant gene against zeocin (*ble*) for second selection.

The transformation of the host hC9_5 strain with the linearized pCgNRp[ble]-Lhcx1-RGR vector

produced 252 zeocin-resistant colonies, from which 23 strains were randomly selected, and 13 strains

had various mutations in the target sequence of the *CgLhcx1* gene. Among the strains confirmed for

target sequence editing, *lhcx1-7*, *15*, and *234* strains were used for subsequent analysis. When the

nucleotide sequences of the *Lhcx1* region were confirmed by Sanger sequencing, heterogeneity was

observed (Supplemental Figure 2A). Since *C. gracilis* is a diploid organism like other diatoms,

mutagenic events can occur in both alleles to completely inactivate the target gene. However,

simultaneous mutations in both alleles are rare, and in most cases, only one allele is mutated, or neutral

mutations such as in-frame deletions/insertions may occur, resulting in the formation of mixed cell

populations with different mutation status and types through subsequent cell divisions (Huang &

Daboussi, 2017). Therefore, we examined the accumulation of Lhcx1 protein in these strains by

western blotting using an anti-Lhcx1 antibody (Supplemental Figure 2B). Almost no accumulation of

Lhcx1 protein was observed in the *lhcx1-7* and *lhcx1-15* strains. On the other hand, the *lhcx1-234*

strain showed reduced accumulation of Lhcx1 compared to WT, but it was not a complete deficiency

since *lhcx1-234* may include neutral mutations such as in-frame deletions/insertions.

Phenotypic Analysis of *Lhcx1* Deficient Strains

WT and *lhcx1-7*, *15*, and *234* strains were cultured with air bubbling for 7–8 days under low light

(LL) at 30 μmol photons m^{-2} s^{-1} and high light (HL) at 300 μmol photons m^{-2} s^{-1}. No significant

differences in growth were observed between WT and genome-edited strains (Supplemental Figure 3).

To investigate the photosynthetic characteristics of WT and *lhcx1* strains, pulse-amplified modulated

(PAM) fluorescence measurements were performed on day 3 and day 7–8 using cells cultured under

LL and HL conditions (Figure 2, Supplemental Figure 4). As a result, *lhcx1-7*, *15*, and *234* were hardly

able to induce NPQ, an indicator of thermal dissipation of excess excitation energy around PSII, under

both LL and HL conditions (Figure 2A, Supplemental Figure 4A). The *lhcx1* strains cultured under

LL showed lower photochemical quenching qP under light exposure compared to WT, indicating a

more reduced state of the plastquinone pool. However, this trend was not clear in the *lhcx1* strains

cultured under HL, because non-regulated energy dissipation due to photoinhibition affects the qP

level (Figure 2B). The effective quantum yield of PSII, Y(II), showed higher values in all *lhcx1* strains

compared to the WT strain in cells cultured under HL for 7–8 days (Figure 2C). In cells cultured under

HL for 3 days, *lhcx1* strains showed a trend of higher Y(II) compared to WT, but the difference was

not significant (Supplemental Figure 4C). The *lhcx1-7, 15* strains exhibited significantly higher

oxygen-evolving activity compared to WT cells at the same total Chl *a* content of 5 μg (Supplemental

Figure 5). The *lhcx1-234* strains showed the tendency towards higher oxygen-evolving activity, though

it was not statistically significant ($p = 0.157$). This may be due to the mosaic genetic background of

Lhcx1 mutations in *lhcx1-234*.

Whole-Cell Absorption Spectra and Pigment Analysis

The whole-cell absorption spectra showed no significant differences between WT and *lhcx1*

mutants in cells cultured under LL, while in the cells cultured under HL, the *lhcx1* mutants exhibited

greater absorption in the 400–500 nm region compared to the WT (Supplemental Figure 6). Since this

region corresponds to carotenoid absorption, pigment analysis using HPLC (High Performance Liquid

Chromatography) was performed to examine carotenoid contents. The amount of Chl *a* per cell was

decreased in the mutants compared to WT under LL conditions (Figure 3A). Additionally, under HL

conditions, WT also showed a decrease in Chl *a* content compared to LL conditions, but the decrease

tends to be greater in the *lhcx1* mutants. Since the total absorption spectra were normalized at the Qy

peak of Chl *a*, carotenoid content was quantified per Chl *a* amount. Fucoxanthin, the major carotenoid

in *C. gracilis* FCP, showed no significant differences between WT and mutants under both light

conditions, except for *lhcx1-15* under HL conditions (Figure 3B). The *lhcx1-15* under HL conditions

showed slightly lower fucoxanthin content compared to WT under the same conditions. On the other

hand, for diadinoxanthin (Ddx) and diatoxanthin (Dtx), the major pigments of the diatom xanthophyll

cycle, no significant differences in accumulation were observed between WT and *lhcx1* mutants under

LL conditions (Figure 3C, D). In contrast, under HL conditions, WT showed greater accumulation of

Dtx compared to LL conditions. The mutant strains accumulated significantly more Dtx compared to

the WT under the same conditions. Regarding Ddx, both WT and mutants showed greater

accumulation under HL conditions compared to LL conditions, but significant increases in

accumulation in mutants compared to WT were not observed, except for *lhcx1-15*. The de-epoxidation

state (DEPS = [Dtx]/([Dtx]+[Ddx])), an indicator of the xanthophyll cycle, was calculated from the

accumulation of Dtx and Ddx (Figure 3E). Under LL, there was no significant difference between WT

and mutants. Under HL, the WT showed a higher DEPS (0.34) compared to LL conditions. The

mutants showed significantly higher DEPS (0.47–0.52) compared to WT under the same conditions.

Steady-State Low-Temperature Fluorescence and Antenna Size Measurements

WT and *lhcx1* cultured under HL were subjected to 30-minute treatments under either dark

conditions (dark-treated) or HL conditions (HL-treated), then were immediately frozen in liquid

nitrogen, and used for the low-temperature (77K) fluorescence spectra measurements (Figure 4A, B).

WT was measured as two independent samples cultured separately (solid and dashed lines in Figure

4A, B). The 77K steady-state fluorescence spectra well represent the distribution of excitation energy

between photosystems and light-harvesting pigment proteins. The excitation wavelength was 459 nm,

which efficiently excites Chl *c* and fucoxanthin bound to the peripheral antenna FCP in diatoms. The

77K fluorescence spectra of *C. gracilis* showed two peaks derived from CP43 and CP47 of

photosystem II, observed around 686 nm and 694 nm, respectively (Andrizhiyevskaya *et al.*, 2005;

Nagao *et al.*, 2010). In all cells cultured under HL, the mutants showed lower CP43 peaks compared

to the WT in spectra normalized at the CP47 peak (Figure 4A). The fluorescence intensity around 680

nm, which is considered to be the fluorescence from free FCP, was smaller in *lhcx1* mutants under

dark conditions compared to the WT. These indicate that energy transfer from the peripheral antenna

FCP via CP43 is prominent in the dark, especially in WT. This also suggests that the effective antenna

size of PSII differed between WT and *lhcx1* in the dark. Therefore, the antenna size after dark

adaptation of cells cultured under HL was measured by measuring the induction curve of Chl

fluorescence with a Fluorometer FL 3500 (PSI, Czech Republic) (Nedbal *et al.*, 1999). As a result, it

was revealed that *lhcx1* had a significantly smaller effective antenna size of PSII compared to WT in

all strains (Supplemental Figure 7).

Time-Resolved Low-Temperature Fluorescence Spectra

To elucidate the process of excitation energy transfer from the peripheral antenna FCP to the

photosystems, time-resolved fluorescence (TRF) spectra were measured under the same conditions as

the samples subjected to low-temperature steady-state fluorescence spectrum measurements. TRF

fluorescence was measured on samples of whole cells that were frozen in liquid nitrogen immediately

after dark treatment and HL treatment (300 μmol photons m^{-2} s^{-1}). This allows observation of

excitation energy transfer and quenching in states where NPQ is not induced and where it is induced

(Chukhutsina *et al.*, 2017). Fluorescence decay-associated spectra (FDAS) were obtained by global

analysis that decomposed the obtained TRF spectra into multiple exponential components with time

as a variable (Figure 5, Supplemental Figure 8). In FDAS, positive peaks (decay) indicate the decay

of fluorescence, while negative peaks (rise) indicate the rise of fluorescence. From the TRF spectra of

the WT, FDAS were constructed with six-time constants. The assignment of each peak followed

previous literature (Andrizhiyevskaya *et al.*, 2005; Nagao *et al.*, 2010). In the dark-adapted WT, the

FDAS with a lifetime of 40 ps showed positive peaks of FCP at <680 nm and a large negative peak

around 687 nm corresponding to CP43 (black solid line in Figure 5 and Supplemental Figure 8). This

reflects excitation energy transfer from FCP to CP43. In the spectrum of the second lifetime of 140 ps,

positive peaks were observed not only at ~680 nm but also around 687 nm corresponding to CP43,

while the waveform around 694 nm corresponding to CP47 showed a convex downward shape. This

is thought to be due to the negative peak of CP47 overlapping with the shoulder of the nearby positive

peak, and is considered to indicate excitation energy transfer from FCP/CP43 to CP47. The third

lifetime of 640 ps continued to show the same trend as 140 ps, suggesting that excitation energy

transfer from FCP/CP43 to CP47 was ongoing. The fourth and fifth lifetimes, at 1.6 ns and 3.8 ns,

showed positive peaks corresponding to CP43 and CP47, indicating simple fluorescence decay. The

longest lifetime of 26 ns did not exhibit a prominent positive peak around 710 nm, which is

characteristic of PSI fluorescence, but instead showed positive peaks corresponding to CP43 and CP47,

associated with charge recombination in photosystem II. Therefore, excitation energy transfer from

PSII to PSI, so-called spillover, did not occur.

On the other hand, WT subjected to HL (black dashed line in Figure 5 and Supplemental Figure 8)

showed a similar waveform at 40 ps as the dark-adapted WT. At 140 ps, although the positive peak

around 687 nm was much larger compared to the dark-adapted WT, the value at the dip around 694

nm was larger than that of the dark-adapted WT. In other words, WT acclimated to HL had a smaller

fluorescence rise at CP47 compared to the dark-adapted sample, while the energy decay around 687

nm was larger. The *lhcx1* mutant, which does not induce NPQ, showed spectra similar to the dark-

adapted WT in the first and second time constants when dark-adapted (blue, green, and red solid lines

in Figure 5 and Supplemental Figure 8A, B). However, in the third time constant (640 ps), while the

dark-adapted WT showed excitation energy transfer from FCP/CP43 to CP47, *lhcx1* already showed

positive peaks at both 687 nm and 694 nm, indicating fluorescence decay of CP43 and CP47. In

subsequent time constants, 687 nm and 694 nm also showed only positive peaks. The HL-acclimated

lhcx1 (blue, green, and red dashed lines in Figure 5 and Supplemental Figure 8A, B) also showed

spectra similar to the dark-adapted *lhcx1* at all time constants and did not show the large positive peak

around 687 nm at 140 ps observed in the HL-acclimated WT. Furthermore, since energy transfer from

FCPII to CP43 and CP47, and from CP43 to CP47, was completed by 140 ps, similar to the HL-treated

WT and dark-treated *lhcx1*, it was concluded that *lhcx1* has a smaller antenna size compared to the

WT even under HL conditions. From the above, while the decay of fluorescence around 687 nm is

normally attributed to energy transfer from CP43, when looking at the energy transfer of *lhcx1* at the

same time constant, such decay is not observed. Therefore, it is concluded that the decay of

fluorescence around 687 nm in the HL-acclimated WT is not decay of CP43 but quenching attributed

to Lhcx1.

Molecular interaction of Lhcx1 in thylakoid membranes

To determine the localization of Lhcx, we performed native-PAGE followed by second-dimension

SDS-PAGE and subsequent immunoblotting using anti-Lhcx1 antibody. Since conventional Clear

Native (CN)-PAGE with deoxycholate made it difficult to separate diatom $C_2S_2M_2$ (Core$_2$–S-

tetramer$_2$–M-tetramer$_2$) PSII–FCPII while keeping FCP-A (which contains CgLhcf1 as a major

component) bound to PSII (Nagao *et al.*, 2012), we applied a modified CN-PAGE using Amphipol

A8-35 (Kameo *et al.*, 2021), which enhances the stability of solubilized protein complexes, instead of

deoxycholate as the surfactant that imparts negative charge to proteins (Figure 6A). In the diatom *C.*

gracilis, a dense band was observed at a higher molecular weight position than spinach $C_2S_2M_2$

(Core$_2$–S-trimer$_2$–M-trimer$_2$) PSII–LHCII. Additionally, a broad brown band, which was observed at

a higher molecular weight position than the LHC/FCP monomer, extended to a lower molecular weight

position than the spinach LHCII trimer. This CN-PAGE gel strip was subjected to SDS-PAGE with a

gel containing 6 mol L^{-1} urea to separate each subunit (Figure 6B). Annotation of each spot was based

on previous reports (Nagao *et al.*, 2007; Zhou *et al.*, 2024). The amino-acid sequence of

HGRIAQLAFLGN shared with CgLhcf2 and CgLhcf12 was identified from the major FCP fraction

(Ishihara *et al.*, 2015). These reports indicate that FCP-B/C is one of the major FCPs in *C. gracilis*.

Furthermore, since the FCP-B/C heterodimer is presumed to be loosely bound to $C_2S_2M_2$ PSII–FCPII

and transfer excitation energy, it was designated as L (Loose)-dimer (Figure 6A, B). Additionally,

immunoblotting was performed to identify the Lhcx1 spot from the spots in the two-dimensional SDS-

PAGE (Figure 6C). The Lhcx1 spot, indicated by the blue arrowhead in Figure 6C, was detected at a

higher molecular weight than FCP-B/C in 2D-SDS-PAGE. In the 1D-CN-PAGE direction, the spot of

Lhcx1 was detected across a wide range of multimers from monomers to higher-order multimeric

complexes, co-migrating with FCP-B/C (Figure 6B). This result suggests that Lhcx1 may interact with

the FCP-B/C dimers, functioning as the loosely associated antenna of $C_2S_2M_2$ PSII–FCPII.

Phenotype of *lhcx1* Mutants in High CO_2 Environment

We investigated whether the *lhcx1* mutants, which showed a higher effective quantum yield of PSII

when acclimated to HL under normal atmospheric aeration conditions, would further improve their

PSII effective quantum yield by enhancing downstream electron sink capacity with 3% CO_2 addition.

First, even when cultured with CO_2 aeration, *lhcx1* showed almost no NPQ regardless of light intensity

(Figure 7A). The effective quantum yield of PSII, Y(II), increased in *lhcx1* when cultured with CO_2

aeration even under LL conditions, where no difference was observed with atmospheric aeration (solid

line in Figure 7C). This Y(II) under light irradiation increased with time during the initial irradiation

period and reached a constant value after about 2 minutes, which is thought to be due to the activation

of the carbon assimilation pathway. Additionally, an increase in Y(II) was also observed under HL

conditions. Photochemical quenching qP, which indicates the redox state of plastoquinone, was similar

between WT and *lhcx1* under LL conditions. As observed in Figure 2B, qP of *lhcx1* varied among

strains, with strains having lower Y(II) showing lower qP under HL conditions (Figure 7B).

To understand the general physiological response of *lhcx1*, we performed RNA sequencing analysis

under LL/HL and ambient/3%-CO_2-supplemented air conditions. Under ambient air conditions, HL

treatment induced 1,494 genes in WT and 1,530 genes in *lhcx1* mutant, with 458 genes commonly

upregulated in both genotypes. Under CO_2-enriched conditions, HL induced substantially more genes:

2,411 in WT and 2,402 in *lhcx1* mutant. To identify responses compensating the absence of NPQ in

lhcx1, we compared the magnitude of HL response (fold-change from LL to HL) between genotypes.

lhcx1 showed enhanced HL response compared to WT in 978 genes under ambient air and 311 genes

under CO_2-enriched conditions, with 105 genes showing this enhanced response under both ambient

and CO_2 conditions. KEGG pathway enrichment analysis reveals that while both WT and *lhcx1*

respond to HL through porphyrin metabolism and photosynthesis pathways (Figure 8A), *lhcx1* mutant

specifically enhances carbon fixation and central carbon metabolism pathways under both ambient air

and CO_2 conditions (Figure 8B).

Discussions

The deficiency of Lhcx, while Lhcx belongs to the LHC family primarily responsible for light

harvesting, almost completely suppresses the induction of non-photochemical quenching (NPQ). The

NPQ-deficient *lhcx1* strains acclimated by inducing various HL defense systems possessed by *C.*

gracilis and showed higher PSII activity compared to the WT. Since the increase in PSII activity was

also induced in *lhcx1* with the addition of CO_2, it was revealed that acclimation of downstream electron

sinks in the electron transport chain, along with acclimation of the upstream light-harvesting system,

plays an important role in HL acclimation of *C. gracilis*. This study suggests that the multiple

photoprotection mechanisms can be interrelated and flexibly regulated in response to environmental

changes in the marine diatom *C. gracilis*.

Origin of Lhcx

The Lhcx subfamily, which is responsible for non-photochemical quenching (NPQ) in red-lineage

secondary endosymbiotic algae such as heterokonts including diatoms, and presumably haptophytes,

and the Lhcsr subfamily, which similarly mediates NPQ in green algae, streptophyte algae, and

bryophytes, have been considered to share a common ancestral gene as they form a monophyletic

group (Koziol *et al.*, 2007). Although the Lhcx/Lhcsr subfamily is suggested to be derived from the

red lineage, detailed molecular phylogenetic relationships with other subfamilies have not been

thoroughly investigated, likely due to insufficient sequence information (Dittami *et al.*, 2010). In this

study as well, the diatom Lhcx subfamily and the basal streptophyte algae and core green algae Lhcsr

subfamily formed a monophyletic group (Figure 1). The Lhcx/Lhcsr clade formed by these showed

strong support for monophyly with the diatom Lhcq, CgLhcr9 homolog, and Lhcf subfamilies, and

was in a different clade from the green algae Lhca/Lhcb subfamily, clearly indicating that it is derived

from red-lineage algae. In particular, since the Lhcx/Lhcsr clade showed monophyly with the CgLhcr9

homolog and Lhcf subfamily, it was demonstrated that the Lhcx subfamily is one of the new LHC

subfamily groups that arose in heterokont algae along with Lhcq, CgLhcr9 homolog, and Lhcf

subfamilies. Furthermore, since Lhcsr is conserved in both core green algae and streptophyte algae,

this supports the hypothesis proposed in Dittami et al. that the ancestor of extant green plants acquired

the Lhcx/Lhcsr subfamily through horizontal gene transfer from a red-lineage alga, presumably a

heterokont alga or haptophyte, before the last common ancestor of extant green plants. The estimated

divergence time of extant green plants into Streptophyta and Chlorophyta overlaps with the emergence

of heterokonts (Strassert *et al.*, 2021), indicating that gene transfer of Lhcx/Lhcsr from heterokonts to

ancestral green plants could have occurred prior to this divergence event. It is difficult to determine

whether this gene transfer was merely a horizontal gene transfer or involved the acquisition of a

temporary red-lineage plastid, such as a kleptoplast. Additionally, looking at Figure 1, the

monophyletic clade of the green algae Lhca and Lhcb subfamilies does not form a monophyletic group

with the red algae Lhcr subfamily, and between them exist cryptophyte Lhcr and diatom Lhcr. This

does not negate the possibility that not only Lhcsr but also the Lhca and Lhcb subfamilies were

acquired through horizontal gene transfer. This result calls for reconsideration of the formation and

establishment of early chloroplasts.

Within Lhcx, Lhcx6_1, which exists at the most basal position, is present in *T. pseudonana* and

Cyclotella cryptica but not in *C. gracilis* or *Phaeodactylum tricornutum* (Figure 1, inset). This

Lhcx6_1 not only has an evolutionary distance from other typical Lhcx in the molecular phylogenetic

tree but also does not show a low NPQ phenotype when deficient (Nakayasu *et al.*, 2024). Furthermore,

Lhcx6_1 directly binds to PSII and has a completely different assembly from *C. gracilis* PSII–FCPII

(Feng *et al.*, 2023; Zhao *et al.*, 2023). Based on these observations, it is considered that Lhcx6_1 is an

atypical Lhcx and does not mediate NPQ, and we argue that its classification within the Lhcx

subfamily should also be reconsidered.

Spectroscopic Characteristics of Lhcx Quenching

In this study, we successfully obtained NPQ signals at 140 ps around 687 nm by measuring time-

resolved fluorescence comparing frozen whole cells immediately after dark adaptation and HL

irradiation, reflecting the *in vivo* state (Figure 5, Supplemental Figure 7). By using the *lhcx1* mutant

that shows a clear phenotype with almost no NPQ induction, and by optimizing the dark adaptation

and HL exposure time to 30 minutes, we were able to clearly distinguish signals attributable to Lhcx-

mediated NPQ.

Our results contrast with several previous studies on diatoms. Similar analyses were performed for

the diatoms *C. meneghinia* and *P. tricornutum* (Miloslavina *et al.*, 2009). A different approach was

taken regarding DAS fitting compared to this study, with strong assumptions placed on the

fluorescence signals of PSII and PSI, and quenching signals like those observed in this study were not

obtained. More detailed observations of quenching using time-resolved fluorescence have been

attempted in the diatom *C. meneghiniana*, but the ~200 ps signal around 685 nm observed in this study

was not observed; instead, a ~600 ps quenching signal at 695 nm was observed and attributed to PSII

(Chukhutsina *et al.*, 2014). In a similar approach using WT, a 214 ps signal at 685 nm in

Nannochloropsis gaditana was attributed to LHCII quenching, but it was not clear whether this was a

quenching signal due to the presence of Lhcx (Chukhutsina *et al.*, 2017). A similar approach has also

been used with the *lhcx1* knockdown mutant of *P. tricornutum*, but this mutant showed a limited

decrease in NPQ, and no clear NPQ signal was observed in time-resolved fluorescence (Taddei *et al.*,

2018).

The spectroscopic characteristics of NPQ in diatoms differ notably from those in green algae. In

green alga *Chlamydomonas reinhardtii stt7-9* state-transition-less mutants, NPQ signals were

observed at 150 ps in time-resolved fluorescence and ~100 ps in transient absorption at physiological

temperature, similar to our time constant, but the NPQ signal from fluorescence was detected at 682

nm and attributed to LHCII fluorescence (Tian *et al.*, 2019; Zheng *et al.*, 2024). This wavelength

difference (682 nm vs. 687 nm) likely reflects differences in measurement temperatures and distinct

pigment compositions between the two algal groups, rather than fundamental mechanistic differences.

Time-resolved fluorescence measurements in whole cells revealed longer excitation energy transfer

times compared to isolated complexes. While excitation energy transfer from FCPII to the PSII core

in $C_2S_2M_2$ PSII–FCPII occurs up to a time constant of 210 ps (Nagao *et al.*, 2022), our measurements

in dark-adapted whole cells showed a time constant of 640 ps (Figure 5, Supplemental Figure 8). This

threefold increase suggests that PSII–FCPII complexes *in vivo* possess larger antenna systems than the

isolated $C_2S_2M_2$ form, indicating the presence of additional FCPs connected to the excitation energy

network. These peripheral antenna complexes, likely including L-FCPII dimers, extend the effective

antenna size and modify energy transfer kinetics.

The mechanism of Lhcx1-mediated NPQ

The localization of diatom Lhcx has been debated for many years. Except for Lhcx6_1 associated

with PSII core (Feng *et al.*, 2023; Zhao *et al.*, 2023), the localization of Lhcx has been detected in free

FCP or bulk FCP (Lepetit *et al.*, 2010; Grouneva *et al.*, 2011), except for reports that *P. tricornutum*

Lhcx1/Lhcx2 bind to PSI (Grouneva *et al.*, 2011). Additionally, in SDS-PAGE following CN-PAGE,

T. pseudonana Lhcx1 was detected as a free monomer (Zhou *et al.*, 2024). In the 2D-CN/SDS-PAGE

of the diatom *C. gracilis* by Zhou et al. (2024), Lhcx1 was not detected. In this study, we utilized

Amphipol A8-35, a surfactant that enhances the stability of membrane proteins, to enable the

separation of diatom $C_2S_2M_2$ PSII–FCPII, which had been previously challenging with CN-PAGE

methods (Figure 6A). Under this condition, the dimer (L-dimer) consisting of CgLhcf12/CgLhcf2

(FCP-B/C) showed a broad band in the first dimension (Figure 6B), and a similar band pattern was

observed in the immunoblot for CgLhcx1 (Figure 6C). These findings suggest that CgLhcx1 may

interact with the FCP L-dimer.

Assuming the interaction between L-FCPII dimers and Lhcx1, we propose that Lhcx1-mediated

NPQ operates through two possible mechanisms: either Lhcx1 directly functions as the quenching

center, or quenching occurs through the L-dimer via its interaction with Lhcx1. This mechanism can

be classified within the framework of Q1 and Q2 quenching, where Q1 occurs in detached antenna

complexes and Q2 occurs in minor LHCs bound to the PSII core (Miloslavina *et al.*, 2009; Holzwarth

et al., 2009; Chukhutsina *et al.*, 2014). Given that L-FCPII dimers bind loosely to S/M-FCPII, Lhcx1-

mediated quenching likely represents Q1-type quenching occurring in the peripheral light-harvesting

antenna, although it is unclear whether the L-FCPII dimer detaches from PSII-S/M-FCPII.

The 140 ps quenching time constant observed in HL-treated whole cells is notably faster than

previously reported values for isolated FCP complexes. Isolated FCP-A tetramers and FCP-B/C dimers

have been reported to show characteristic quenching with time constants of 270 ps and 200 ps,

respectively (Nagao *et al.*, 2013). This acceleration in whole cells may result from two factors: first,

isolated FCP-B/C complexes lack Lhcx proteins, which appear crucial for efficient quenching; second,

isolated complexes lack the native membrane environment, including the ΔpH gradient and three-

dimensional protein organization necessary for optimal quenching function. These results highlight

the importance of studying NPQ mechanisms in intact cellular systems rather than relying solely on

isolated protein complexes.

Compensatory mechanisms to the Lhcx1 Deficiency

The *lhcx1* strain, which lacks the most highly expressed Lhcx1 in *C. gracilis* created using genome

editing, induced almost no NPQ and showed higher PSII effective quantum yield and oxygen-evolving

activity compared to the WT when acclimated to HL. Since NPQ mutants in land plants show

decreased PSII electron transport activity, i.e., decreased effective quantum yield, under HL

acclimation (Yang *et al.*, 2022), it is unusual that *lhcx1*, an NPQ mutant in this study, shows a higher

PSII effective quantum yield. Since *lhcx1* has a smaller antenna size even when dark-adapted after HL

acclimation (Figure 5, Supplemental Figure 8), the decrease in excitation energy acquisition per PSII

core is thought to contribute to a higher PSII effective quantum yield. Additionally, since the PSII

effective quantum yield increased even under LL conditions with the addition of CO_2 (Figure 7), and

transcriptomic analysis revealed that *lhcx1* specifically enhances carbon fixation and central carbon

metabolism pathways under both ambient air and CO_2 conditions (Figure 8B), it is likely that the

increase in carbon fixation capacity in the downstream electron transport chain also contributed to the

improvement of PSII effective quantum yield. The activation of downstream electron sinks can be

observed under HL conditions (Figure 2C).

HL acclimation in *lhcx1* was also suggested by the pigment analysis (Figure 3C–E). Chl content per

cell decreased in *lhcx1*, and while the amount of Fx per Chl did not change, the amounts of

xanthophylls Dtx and Ddx increased. The de-epoxidation state (DEPS) calculated from Dtx and Ddx

also increased in *lhcx1* under HL, indicating that acclimation responses under HL are enhanced from

the perspective of pigment composition as well. Generally, in WT diatoms, the amount of Dtx is very

well correlated with NPQ intensity (Lavaud & Lepetit, 2013). However, since *lhcx1* strains show

almost no induction of NPQ despite increased Dtx amounts, the correlation between Dtx amount and

NPQ requires the presence of Lhcx. Although it is not clarified whether the increased xanthophylls in

lhcx1 are incorporated in LHC/FCP, they may function as antioxidants in thylakoid membranes and

contribute to photoprotection. The above multiple photoprotection mechanisms that became apparent

in the absence of Lhcx1 would contribute to the adaptation in variable marine environments.

Materials and Methods

Phylogenetic Analysis of LHCs Across Various Photosynthetic Eukaryotic Lineages

LHC sequences were collected from genomes and transcriptomes included in the Phycocosm and

EukProt databases, in addition to our previous dataset (Kumazawa *et al.*, 2022; Kumazawa & Ifuku,

2024), following the previous method (Kumazawa & Ifuku, 2024). The LHCs in the dataset were

obtained from red algae, early-diverging streptophyte green algae (Leliaert *et al.*, 2012): *Mesostigma*

viride NIES-296 (Liang et al., 2020) and *Klebsormidium nitens* NIES-2285 (Hori *et al.*, 2014),

diatoms: *C. gracilis*, *Thalassiosira pseudonana*, *Cyclotella cryptica*, *Phaeodactylum tricornutum*, and

cryptophytes: *Guillardia theta* CCMP2712 (Curtis *et al.*, 2012) and Cryptophyceae sp. (*Baffinella* sp.)

CCMP2293 (Dorrell *et al.*, 2023). As reference sequences for Lhcsr, *Chlamydomonas reinhardtii*

Lhcsr1 (UniProt: P93664) and Lhcsr3.1 (UniProt: P0DO19), which is identical to Lhcsr3.2 (UniProt:

P0DO18), were added. Redundant sequences were removed from the LHC protein sequences using

CD-HIT (Fu *et al.*, 2012), and alignment was performed with mafft-einsi v7.525, followed by manual

curation (Katoh & Standley, 2013). The curated dataset was realigned, and trimming was performed

using ClipKit v2.3.0 in smart-gap mode to remove unaligned sites (Steenwyk *et al.*, 2020). The

phylogenetic tree was estimated using IQ-TREE v2.3.6 with the Q.pfam+I+R7 model selected by

ModelFinder based on LG and Q.pfam models (Kalyaanamoorthy *et al.*, 2017; Minh *et al.*, 2020,

2021), and visualized with iTOL v6 (Letunic & Bork, 2024). Ultra-fast bootstrap approximation

(UFBoot) and SH-aLRT support (%) tests were performed with 1000 replicates each (Guindon *et al.*,

2010; Hoang *et al.*, 2018). The aBayes support test was also conducted (Anisimova *et al.*, 2011). The

annotation of LHC subfamilies in the phylogenetic tree was based on our previous studies (Kumazawa

et al., 2022; Kumazawa & Ifuku, 2024) and drawn on the tree using InkScape software.

Cultivation

The marine centric diatom *C. gracilis* (UTEX LB 2658) was used for all analyses. Pre-cultivation was performed in IMK medium —4% (w/v) sea salt (SIGMA S9883, US) or artificial seawater made with Daigo's Artificial Seawater SP (Nihon Pharmaceutical MS, Japan) supplemented with 1x Daigo's IMK medium (Nihon Pharmaceutical MS, Japan) and 0.2 mmol L^{-1} Na_2SiO_3— at 20ºC under fluorescent lights at approximately 30 µmol photons m^{-2} s^{-1} with continuous shaking at 100 rpm. Seven–eight-day or three-day cultivation was performed in IMK medium at 25ºC under warm white LEDs at 30 µmol photons m^{-2} s^{-1} (LL) or 300 µmol photons m^{-2} s^{-1} (HL), with continuous aeration of air or 3% (v/v) CO_2.

Preparation of Thylakoid Membranes from *C. gracilis* and Spinach

Diatom WT cells cultured under HL at 25ºC, 300 µmol photons m^{-2} s^{-1} for three days were centrifuged at 1,200×*g*, 20ºC for 5 minutes to collect the cells, then resuspended in artificial seawater, washed, and centrifuged again. After removing the supernatant, the pellet was resuspended in 1 mL of artificial seawater, transferred to a 1.5 mL microcentrifuge tube, and centrifuged at 21,500×*g*, 4ºC for 5 minutes to obtain a pellet. The pellet was resuspended in 1 mL of cell disruption buffer (20 mM Tricine-KOH (pH 7.6), 0.4 mol L^{-1} sorbitol, 10 (w/v) % PEG-6000, 10 mM EDTA), mixed with 0.5 g of 0.1 mm diameter glass beads, and processed with a bead beater for 10 seconds. The sample was then well-chilled on ice, and the disruption process was repeated a total of three times. The supernatant was centrifuged at 21,500×*g*, 4ºC for 5 minutes to obtain a pellet. The pellet was resuspended in 1 mL of washing buffer (20 mmol L^{-1} Tricine-KOH (pH 7.6), 0.4 mol L^{-1} Sorbitol, 5 mmol L^{-1} $MgCl_2$, 2.5 mmol L^{-1} EDTA) and centrifuged at 50×*g*, 4ºC for 5 minutes. The portion containing thylakoid membranes was carefully collected, avoiding the upper and lower contaminants, and centrifuged at 21,500×*g*, 4ºC for 5 minutes to obtain a thylakoid membrane pellet. The pellet was resuspended in

solubilization buffer (50 m mol L^{-1} imidazole-HCl (pH 7.0 at 4℃), 20 (v/v) % glycerol), and a portion

was taken to measure the concentrations of Chl *a* and *c* (Jeffrey & Humphrey, 1975). The thylakoid

membrane of spinach (*Spinacia oleracea*) was isolated from market spinach based on previously

reported methods (Yamamoto *et al.*, 2011) using the modified buffer (50 mmol L^{-1} tricine-NaOH (pH

7.8 at room temperature), 0.4 L^{-1} sucrose, 10 mmol L^{-1} NaCl, 5 mmol L^{-1} MgCl$_2$). The thylakoid

membranes were adjusted to 1 μg μL^{-1} of Chl *a*+*c* with solubilization buffer (50 mM imidazole-HCl

(pH 7.0 at 4℃), 20 (v/v) % glycerol).

2D-CN/SDS-PAGE

The 4–13% gradient gel used for CN-PAGE was prepared according to existing methods (Järvi et

al., 2011; Kameo et al., 2021), and the entire CN-PAGE procedure followed the modified CN-PAGE

method using Amphipol A8-35 (Kameo et al., 2021). However, 75 mmol L^{-1} imidazole-HCl (pH 7.0

at 4ºC) was used in the 3x gel buffer, and 40 (w/v) % acrylamide/bis solution (37.5:1) (019-25655,

Fujifilm Wako, Japan) was used for acrylamide. The anode buffer contained 25 mmol L^{-1} imidazole-

HCl (pH 7.0 at 4ºC), and the cathode buffer contained 50 mmol L^{-1} Tricine, 7.5 mmol L^{-1} imidazole.

The Chl concentrations in resuspended thylakoids were measured (Jeffrey & Humphrey, 1975), and

thylakoid membranes were adjusted to 1 μg μL^{-1} of Chl *a*+*c* and Chl *a*+*b* with solubilization buffer

(50 mmol L^{-1} imidazole-HCl (pH 7.0 at 4℃), 20 (v/v) % glycerol) and resuspended. An equal volume

of 2 (w/v) % dodecyl-α-D-maltoside (α-DM, Anatrace, OH, USA) was added for solubilization, placed

on ice for 2 minutes, then centrifuged at 21,500×*g* for 2 minutes to remove insoluble residues. The

supernatant was mixed with half the volume of 2 (w/v) % amphipol A8-35 (Anatrace, OH, USA), and

5 μg of Chl in 15 μL was applied to each well. Electrophoresis was performed using MyPowerII 300

power supply (AE-8135, ATTO, Japan) at 40 V, 1 mA for 660 minutes at 4ºC with two mini gels

simultaneously. After electrophoresis, the gels were scanned with the scanner ES2200 (EPSON, Japan).

The CN-PAGE gels were immersed in 50 (v/v) % glycerol on ice, cut into gel strips, and stored at −

80ºC in 50 (v/v) % glycerol until use.

Gel strips were solubilized by immersing them in solubilization buffer (1% SDS and 50 mM

dithiothreitol (DTT) for 30 minutes at room temperature, and then applied to gels with a stacking gel

(pH 6.8) and separation gel (14% acrylamide (37.5:1), 6 mol L^{-1} Urea, pH 8.6). The Laemmli system

was used for electrophoresis. When the dye had electrophoresed to the bottom of the marker well at

250 V, 100 mA, the process was temporarily stopped, and Precision Plus Protein Dual Color Standards

(#1610374, BioRad, CA, USA) were applied. Electrophoresis was then continued at 250 V for 90

minutes. After electrophoresis, gels not used for immunoblotting were fluorescently stained with

Oriole fluorescent gel stain (#1610496, BioRad, CA, USA) for 100 minutes, gently rinsed with

distilled water, and then fluorescence was detected using the ChemiDoc Touch Imaging System (Bio-

Rad).

Construction of hCas9 Vector

The hCas9 sequence was amplified from the hCas9 vector (Addgene, plasmid #41815) using the

InF-hCas9-fw1 primer (TTTATAAAGCGGATCATGGACAAGAAGTACTCCATTGG) and InF-

hCas9-rv1 primer (AACAGCTTGCATGCCTCACACCTTCCTCTTCTTCTTG). The PCR product

was cloned into the linearized pCgNRp vector digested with BamHI and PstI using In-Fusion reaction

according to the manufacturer's protocol. The obtained clone vector was named pCgNRphCas9.

Construction of pCgNRpble-Lhcx1-RGR Vector

A 20-mer genome editing target sequence containing the PAM (protospacer adjacent motif)

sequence (NGG) for gRNA in the CRISPR/Cas9 system, GTTGCTATGCTTGCCGTTAT**TGG** (bold

indicates the PAM sequence), was selected. A system that expresses complete gRNA in cells via a

mRNA-type Pol II promoter by connecting self-processing ribozymes (Gao & Zhao, 2014; Zhang *et*

al., 2017) was used. This method was adapted to the genome editing system of *C. gracilis*. The genome

editing vector containing gRNA was constructed using three primers: Lhcx1-RGR-F1;

GACGAAACGAGTAAGCTCGTC**GTTGCTATGCTTGCCGTTAT**GTTTTAGAGCTAGAAATA

GCAAG (bold indicates the 20-mer sequence for gRNA target recognition excluding the PAM

sequence),   Lhcx1-RGR-F2;

GTCAGGATCC**AGCAAC**CTGATGAGTCCGTGAGGACGAAACGAGTAAGCTCGTC (BamHI

restriction enzyme site is underlined, and bold indicates the complementary sequence of the first six

bases of the 20-mer for gRNA target recognition including the PAM sequence), and PstI-RGR-Rv;

ATATCTGCAGGTCCCATTCGCCATGCCGAAGC (PstI restriction enzyme site is underlined). The

PstI-RGR-Rv primer is a universal primer in this protocol. The first fragment (199 bp) was amplified

by PCR from pRS316-RGR-GFP (Addgene, plasmid #51056) using the Lhcx1-RGR-F1 and PstI-

RGR-Rv primers. The second fragment (231 bp) was amplified by PCR from the first purified PCR

product using the Lhcx1-RGR-F2 and PstI-RGR-Rv primers. The purified second product was

digested with BamHI (NEB) and PstI (NEB), and ligated with the pCgNRpble vector linearized with

BamHI and PstI. The final product was named pCgNRp[ble]-Lhcx1-RGR.

Transformation

The electroporation transformation procedure using the Super electroporator NEPA21 Type-II

(NEPAGENE, Japan) followed previous reports (Miyahara *et al.*, 2013; Ifuku *et al.*, 2015; Ifuku &

Yan, 2020). The initial transformation of the *hCas9* gene into *C. gracilis* was performed using the

pCgNRphCas9 vector linearized with DraIII and SapI. Colonies were selected on IMK plates

containing the antibiotic Nourseothricin at 400 μg mL^{-1} (clonNAT; Hans Knoell Institute, Jena,

Germany). The introduced *hCas9* gene was confirmed by PCR using the first set: InF-hCas9-fw1

primer and hCas9-2088-Rv primer: CCTTAAAGGTGAGAGAGTCATCATGGATC, and another set:

hCas9-1965n-Fw primer: ATGAAACAGCTCAAGAGGCGCCGA and InF-hCas9-rv1 primer. The

hCas9 strain (hC9 strain) used for further genome editing was selected from eight transformants

screened by PCR and immunoblotting using anti-Cas9 antibody (described later).

The linearized pCgNRpble-Lhcx1-RGR was introduced into the hC9 strain by electroporation. This

was essentially the same as the transformation of hCas9. Before introducing the RGR vector, the vector

was linearized with HindIII (NEB) and concentrated to 850–1000 ng μL^{-1} by ethanol precipitation.

The amount of linearized vector was 5 μg per electroporation. Colonies were selected on IMK plates

containing the antibiotic Zeocin at 100 $\mu g\ mL^{-1}$ (InvivoGen #ant-an-1, US CA). Mutations were

confirmed by Sanger sequencing of the products amplified by PCR using Lhcx1_seq_FW primer:

AGAGACTGTAAGGATTCGGA and Lhcx1_seq_RV primer: GAAGGAAGATCCCTCTACCT.

Immunoblotting

Diatom cells were collected and solubilized in sample buffer containing 62.5 mmol L^{-1} Tris–HCl

(pH 6.8), 2.5% (w/v) SDS, 10% (w/v) glycerol, 2.5% (v/v) 2-mercaptoethanol, and a trace amount of

bromophenol blue (BPB). Solubilization was performed by incubating at 98ºC for 6 minutes, followed

by 30 minutes at room temperature, and then centrifugation. The protein concentration was adjusted

to 30 μg/20 μL. Thirty micrograms of protein was applied to each lane of an 8% acrylamide separation

gel with a 3% acrylamide stacking gel and separated by SDS-PAGE. Proteins were transferred to an

Immobilon-P PVDF membrane (Millipore, MA, USA) and incubated with anti-Cas9 polyclonal

antibody (632607, Takara, Japan) at a 1:2000 dilution.

For the detection of Lhcx1, SDS-PAGE gels were prepared with a 15% acrylamide separation gel

and a 4.5% acrylamide stacking gel. Two μg of protein was applied to each lane and separated by SDS-

PAGE. The anti-Lhcx1 peptide antibody was produced against the synthetic peptide

"KEEYAPGDLRFDPFGLMP". After electrophoresis, the gel was desalted and equilibrated in

Electron transfer buffer (25 mmol L^{-1} Tris, 192 mmol L^{-1} glycine, 20 (v/v) % methanol, 0.05 (w/v) %

SDS). Subsequently, proteins were transferred from the gel to an Immobilon-P PVDF membrane

(Millipore, Billerica, MA, USA) and incubated with anti-Lhcx1 antibody at a 1:5000 dilution.

Immunoreactive proteins were detected using the Amersham ECL Prime Western Blotting Detection

Reagent (Cytiva RPN2236) with the ChemiDoc Touch Imaging System (Bio-Rad).

PAM Fluorescence Measurement

Chl fluorescence by pulse-amplitude modulation was detected using an actinic light of 450 or 455

nm (blue) at 300 μmol photons m^{-2} s^{-1} with the NPQ2 protocol (200 seconds of light period and 390

seconds of dark recovery, with the first saturating pulse light in the light period at 10 seconds after the

start of illumination) of AquaPen AP-100 or AP-110 (PSI, Czech Republic) using a bandpass filter

(667 to 750 nm). The measuring pulse was set to 10%, and the super pulse (saturating pulse light) was

set to 30% (equivalent to 900 μmol photons m^{-2} s^{-1}). Cells used in the NPQ2 protocol were dark-

adapted for more than 10 minutes before measurement.

Oxygen-Evolving Activity Measurements

The diatom cells grown under HL conditions (200 μmol photons m^{-2} s^{-1} with rotational shaking at

~22ºC) were adjusted to a concentration of 5 μg Chl *a* mL^{-1} (Jeffrey & Humphrey, 1975), following

the methods described in Nakayasu et al. (Nakayasu *et al.*, 2024). Using a Clark-type oxygen electrode

(Hansatech, UK) at 25°C, oxygen evolution was measured in the medium supplemented with 10 mM

NaHCO$_3$. Cells were illuminated with white light at 300 μmol photons m^{-2} s^{-1}, and the cell

suspension in the reaction chamber was continuously stirred with a magnetic stirrer during

measurements.

HPLC Pigment Analysis

Cells cultured for 7–8 days under LL or HL conditions were collected by centrifugation at 1,500 x

g for 5 minutes, and the supernatant was removed. The pellet was immediately frozen in liquid nitrogen

and stored at -80 ºC until extraction. The frozen pellet was extracted using 100% acetone (Wako,

Japan) with sonication for 6 minutes in an ice-water bath. Subsequently, the sample was centrifuged

at 15,000 rpm for 5 minutes at 4 ºC. The resulting supernatant was filtered through a 0.45 nm

hydrophilic PTFE filter (SLPT0445NL, Hawach Scientific, IL, USA) and subjected to HPLC. The

HPLC method is based on Nagao, Ueno, et al. (2020) with modifications derived from Manuel Zapata

et al. (2000). The 0.25 mol L^{-1} pyridine contained in solution A was adjusted to pH 5.0 using acetic

acid. A Shimadzu HPLC system equipped with LC-20AD pumps and an SPD-M20A detector was

used for pigment composition analysis. The system was equipped with an Inertsil C8 reverse-phase

column (5020-01228, GL Sciences, Japan). The sample injection volume was 20 μL for each analysis.

Standard pigments of Chl a, fucoxanthin, diadinoxanthin, and diatoxanthin were purchased from DHI

(Denmark), analyzed similarly, and used to create calibration curves.

77K Steady-State Fluorescence Spectra

Cells for measurement were cultured under HL at 300 μmol photons m^{-2} s^{-1} at 25ºC for 3 days.

Samples were diluted with IMK medium and adjusted to 10 μg Chl $a+c$/150 μL. Chl concentration

was measured after extraction with 90% acetone (Jeffrey & Humphrey, 1975). The diluted samples

were treated in the dark or under HL at 300 μmol photons m^{-2} s^{-1} for 30 minutes and immediately

frozen in liquid nitrogen. Low-temperature steady-state fluorescence emission spectra of whole cells

were measured at 77K with excitation at 459 nm using a JASCO FP-8500 equipped with a PMU-130

liquid nitrogen cooling unit. The spectra were integrated from 5 consecutive measurements with a 2.5

nm sampling pitch. The spectra were normalized at the PSII CP47 peak (693–695 nm)

(Andrizhiyevskaya *et al.*, 2005; Nagao *et al.*, 2010).

Time-Resolved Fluorescence and Fluorescence Decay-Associated Spectra

Samples were prepared for time-resolved fluorescence (TRF) spectral measurements using the same

method as for steady-state measurements. TRF spectra were measured using a time-correlated single-

photon counting system in the range of 670 nm to 750 nm with wavelength intervals of 1 nm and time

intervals of 18.3 ps (Hamada *et al.*, 2017). A picosecond pulsed diode laser (PiL047X; Advanced Laser

Diode Systems) operating at 459 nm with a repetition frequency of 3 MHz was used as the excitation

light source to excite Chl *c* and fucoxanthin, which are the main light-harvesting pigments in diatoms.

Detailed conditions for TRF measurements were described in Nagao et al. (2020). Fluorescence decay-

associated spectrum (spectra) (FDAS) was constructed by global analysis (Ye *et al.*, 2025). The

analysis of fluorescence decay $I_F(\lambda, t)$ at wavelength λ with the instrument response function $I_{IRF}(t)$

followed the function:

665
$$I_F(\lambda, t) = C \sum_{i=1}^{6} A_i(\lambda) \int_{-\infty}^{t} I_{IRF}\left(t'\right) exp\left(-\frac{t - t'}{\tau_i}\right) dt'$$

The time constants were set to τ_{1-6} = 40 ps, 140 ps, 640 ps, 1.6 ns, 3.8 ns, and 26 ns based on the

analysis results of WT cells cultured under white LED light. $A_i(\lambda)$ represents each FDAS. The rise and

decay components of fluorescence are indicated by negative and positive peaks, respectively. Positive

and negative amplitudes in the same FDAS reflect energy transfer from pigments with positive

amplitude to pigments with negative amplitude (Yokono *et al.*, 2008; Akimoto *et al.*, 2012). Positive

amplitude without accompanying negative amplitude signifies energy quenching. For Figure 5 and

Figure S8, FDAS of each sample were normalized as,

673
$$\frac{A_i(\lambda)}{\int \sum_{i=1}^{6} A_i(\lambda) \, d\lambda} \times 100.$$

For easy viewing, a magnification factor 100 was used.

Measurement of PSII Antenna Size

Chl fluorescence induction curves of wild-type (WT) and three *lhcx1* mutants (*lhcx1-7*, *lhcx1-15*,

lhcx1-234) cultured under HL at 300 μmol photons $m^{-2}\,s^{-1}$ at 25ºC for 3 days were measured using

a Fluorometer FL 3500 (PSI, Czech Republic) in the presence of 40 μM DCMU after 10 minutes of

dark adaptation (Nedbal et al., 1999). The cultures were grown in two independent batches. In the

measurement, Flash intensity was set to 50%, and Flash duration was set to 100 μs. The relative

antenna size was determined by measuring the initial slope of fluorescence rise during the first 1 μs

relative to the maximum fluorescence intensity. Four replicate measurements were performed for each

strain.

For data analysis, R statistical software (R version 4.4.0 (2024-04-24)) was used, utilizing the

tidyverse, readr, lme4, emmeans, and ggplot2 packages. To account for batch-specific variations,

antenna size values were normalized by the WT mean value within each batch. A linear mixed model

(LMM) was applied to the normalized data. Estimated marginal means were calculated, and multiple

comparisons were performed using the Tukey method. The threshold for statistical significance was

set at adjusted $p < 0.05$.

RNA sequencing

Cells were harvested during the exponential growth phase after three days of treatment exposure to

LL/HL and ambient/3%-CO_2-supplemented air aeration, frozen with liquid nitrogen, and stored at −

80ºC until use. The frozen cells were crushed by bead-beating with 3-mm diameter zirconia beads.

Total RNA was extracted using the RNeasy Plant Mini Kit (Qiagen). RNA-seq libraries were

constructed using the NEBNext Ultra II RNA Library Prep Kit for Illumina (Cat No. 7770) according

to the manufacturer's instructions. Paired-end sequencing (2 × 150 bp) was performed on an Illumina

NovaSeq   6000   platform.   Adapter   sequences   (Read   1:

AGATCGGAAGAGCACACGTCTGAACTCCAGTCA,   Read   2:

AGATCGGAAGAGCGTCGTGTAGGGAAAGAGTGT) were trimmed from raw sequencing reads.

Prior to mapping, the gene structure annotation provided in ChaetoBase v1.1 (predicted_all.gff3) was

modified using the following procedures: (1) replacement of geneID attributes with Parent attributes

for transcripts lacking gene correspondence, (2) correction of transcripts incorrectly defined as genes,

and (3) addition of missing gene definitions using AGAT v1.0.0.

Clean reads were aligned to the *C. gracilis* reference genome (ChaetoBase v1.1) using STAR

v2.7.11b with the following parameters: --alignIntronMax 10000. Gene expression levels were

quantified using RSEM v1.3.3 with --strandedness reverse to account for Illumina stranded RNA-seq

protocol, and expected count matrices were generated for downstream analysis.

Differential Gene Expression Analysis

Statistical analysis of gene expression was performed using the edgeR package (v4.6.2) in R

(v4.5.0). Genes with low expression levels were filtered out by retaining only those with CPM (counts

per million) values exceeding the 20th percentile in at least one condition. A DGEList object was

created, and normalization factors were calculated using the TMM (trimmed mean of M-values)

method. Dispersion parameters were estimated using the estimateDisp function, and generalized linear

models (GLM) were fitted using glmQLFit.

Statistical Contrasts and Hypothesis Testing

To address the research objectives, the following statistical contrasts were performed:

1) Primary Analysis: HL Response Comparison

722    WT HL response: Air.HL.WT vs Air.LL.WT

723    lhcx1 HL response: Air.HL.lhcx1 vs Air.LL.lhcx1

724    Enhanced HL response in lhcx1: (Air.HL.lhcx1 - Air.LL.lhcx1) - (Air.HL.WT - Air.LL.WT)

725  2) Secondary Analysis: CO_2 Effect Validation

726    WT HL response under CO_2: CO_2.HL.WT vs CO_2.LL.WT

727    lhcx1 HL response under CO_2: CO_2.HL.lhcx1 vs CO_2.LL.lhcx1

728    Enhanced HL response in *lhcx1* under CO2: (CO2.HL.lhcx$_1$ - CO_2.LL.lhcx1) - (CO_2.HL.WT

729 - CO_2.LL.WT)

730 Differential expression was determined using quasi-likelihood F-tests implemented in glmQLFTest.

731 P-values were adjusted for multiple testing using the Benjamini-Hochberg method, with an FDR

732 threshold of 0.05.

734 **Functional Enrichment Analysis**

735 KEGG pathway enrichment analysis was conducted using the enrichKEGG function from the

736 clusterProfiler package (v4.16.0). KEGG Orthology (KO) annotations for *C. gracilis* genes were

737 obtained from BLASTP searches against the KEGG database using BLASTKOALA. Only pathways

738 with q-value < 0.05 were considered significantly enriched.

Acknowledgment

This work was supported in part by JSPS KAKENHI grant nos. JP22KJ2017 (M.K.), JP24H02086 (S.A.), JP23KJ1361 (Ko.I.), JP23H02347 (Ke.I. and S.A.), JP24H02081 (Ke.I.), and a grant no. L-2023-3-007 from the Institute for Fermentation, Osaka, Japan (Ke.I.).

Competing Interests

Authors HH, AS, and SI are employed by the NTT, Inc.

Author Contributions

M.K. and Ke.I. conceived the project; M.K. performed phylogenetic analysis, 2D-CN/SDS-PAGE, growth experiments, PAM analyses, absorbance spectrum analyses, 77K emission fluorescence analysis, antenna size measurements, and RNA-seq *in silico* analysis, and prepared samples for ime-resolved fluorescence analyses; S.A. performed time-resolved fluorescence analyses; A.T. conceived the methodology of 2D-CN/SDS-PAGE; Ko.I. measured and analyzed oxygen-evolving activity; S.T. measured pigments; H.H., A.S., and S.I. performed RNA-seq and provided funding; N.I. conceived methodology of SDS-PAGE and immunoblotting; N.I.-K. and Y.K. provided anti-Lhcx1 antibody. M.K. drafted the original manuscript, and Ke.I. revised and edited the final manuscript. All authors contributed to the discussion of the results and improvement of the manuscript.

References

References

Akimoto S, Yokono M, Hamada F, Teshigahara A, Aikawa S, Kondo A. **2012**. Adaptation of

light-harvesting systems of Arthrospira platensis to light conditions, probed by time-resolved

fluorescence spectroscopy. In: *Biochimica et Biophysica Acta - Bioenergetics*. 1483–1489.

Allorent G, Lefebvre-Legendre L, Chappuis R, Kuntz M, Truong TB, Niyogi KK, Ulm R,

Goldschmidt-Clermont M. **2016**. UV-B photoreceptor-mediated protection of the photosynthetic

machinery in *Chlamydomonas reinhardtii*. *Proceedings of the National Academy of Sciences of the*

United States of America **113**: 14864–14869.

Allorent G, Tokutsu R, Roach T, Peers G, Cardol P, Girard-Bascou J, Seigneurin-Berny D,

Petroutsos D, Kuntz M, Breyton C, *et al*. **2013**. A Dual Strategy to Cope with High Light in

Chlamydomonas reinhardtii. *The Plant Cell* **25**: 545–557.

Andrizhiyevskaya EG, Chojnicka A, Bautista JA, Diner BA, Van Grondelle R, Dekker JP.

2005. Origin of the F685 and F695 fluorescence in Photosystem II. In: *Photosynthesis Research*. 173–

180.

Anisimova M, Gil M, Dufayard JF, Dessimoz C, Gascuel O. **2011**. Survey of branch support

methods demonstrates accuracy, power, and robustness of fast likelihood-based approximation

schemes. *Systematic Biology* **60**: 685–699.

Bailleul B, Berne N, Murik O, Petroutsos D, Prihoda J, Tanaka A, Villanova V, Bligny R, Flori

S, Falconet D, *et al*. **2015**. Energetic coupling between plastids and mitochondria drives CO_2

assimilation in diatoms. *Nature* **524**: 366–369.

Bailleul B, Rogato A, De Martino A, Coesel S, Cardol P, Bowler C, Falciatore A, Finazzi G.

2010. An atypical member of the light-harvesting complex stress-related protein family modulates

diatom responses to light. *Proceedings of the National Academy of Sciences of the United States of*

America **107**: 18214–18219.

Buck JM, Kroth PG, Lepetit B. **2021**. Identification of sequence motifs in Lhcx proteins that

confer qE-based photoprotection in the diatom *Phaeodactylum tricornutum*. *Plant Journal* **108**: 1721–

1734.

Buck JM, Sherman J, Bártulos CR, Serif M, Halder M, Henkel J, Falciatore A, Lavaud J,

Gorbunov MY, Kroth PG, *et al.* 2019. Lhcx proteins provide photoprotection via thermal dissipation

of absorbed light in the diatom *Phaeodactylum tricornutum*. *Nature Communications* **10**.

Chukhutsina VU, Büchel C, Van Amerongen H. **2014**. Disentangling two non-photochemical

quenching processes in *Cyclotella meneghiniana* by spectrally-resolved picosecond fluorescence at 77

K. *Biochimica et Biophysica Acta - Bioenergetics* **1837**: 899–907.

Chukhutsina VU, Fristedt R, Morosinotto T, Croce R. **2017**. Photoprotection strategies of the

alga *Nannochloropsis gaditana*. *Biochimica et Biophysica Acta - Bioenergetics* **1858**: 544–552.

Croteau D, Jaubert M, Falciatore A, Bailleul B. **2025**. Pennate diatoms make non-photochemical

quenching as simple as possible but not simpler. *Nature Communications* **16**.

Curtis BA, Tanifuji G, Maruyama S, Gile GH, Hopkins JF, Eveleigh RJM, Nakayama T,

Malik SB, Onodera NT, Slamovits CH, *et al.* 2012. Algal genomes reveal evolutionary mosaicism

and the fate of nucleomorphs. *Nature* **492**: 59–65.

Dittami SM, Michel G, Collén J, Boyen C, Tonon T. **2010**. Chlorophyll-binding proteins revisited

- a multigenic family of light-harvesting and stress proteins from a brown algal perspective. *BMC*

Evolutionary Biology **10**: 365.

Dorrell RG, Kuo A, Füssy Z, Richardson EH, Salamov A, Zarevski N, Freyria NJ, Ibarbalz

FM, Jenkins J, Karlusich JJP, *et al.* 2023. Convergent evolution and horizontal gene transfer in

Arctic Ocean microalgae. *Life Science Alliance* **6**.

Feng Y, Li Z, Li X, Shen L, Liu X, Zhou C, Zhang J, Sang M, Han G, Yang W, *et al.* 2023.

Structure of a diatom photosystem II supercomplex containing a member of Lhcx family and dimeric

FCPII. *Science Advances* **9**.

Fu L, Niu B, Zhu Z, Wu S, Li W. **2012**. CD-HIT: Accelerated for clustering the next-generation

sequencing data. *Bioinformatics* **28**: 3150–3152.

Gao Y, Zhao Y. **2014**. Self-processing of ribozyme-flanked RNAs into guide RNAs in vitro and in

vivo for CRISPR-mediated genome editing. *Journal of Integrative Plant Biology* **56**: 343–349.

Giossi CE, Wünsch MA, Dautermann O, Schober AF, Buck JM, Kroth PG, Lohr M, Lepetit

B. **2025**. Both major xanthophyll cycles present in nature promote non-photochemical quenching in a

model diatom. *Plant Physiology*: kiaf371.

Giovagnetti V, Jaubert M, Shukla MK, Ungerer P, Bouly JP, Falciatore A, Ruban AV. 2022.

Biochemical and molecular properties of LHCX1, the essential regulator of dynamic photoprotection

in diatoms. *Plant Physiology* **188**: 509–525.

Goss R, Lepetit B. **2015**. Biodiversity of NPQ. *Journal of Plant Physiology* **172**: 13–32.

Grouneva I, Rokka A, Aro EM. **2011**. The thylakoid membrane proteome of two marine diatoms

outlines both diatom-specific and species-specific features of the photosynthetic machinery. *Journal*

of Proteome Research **10**: 5338–5353.

Guindon S, Dufayard JF, Lefort V, Anisimova M, Hordijk W, Gascuel O. **2010**. New algorithms

and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0.

Systematic Biology **59**: 307–321.

Hamada F, Murakami A, Akimoto S. **2017**. Adaptation of Divinyl Chlorophyll *a*/*b*-Containing

Cyanobacterium to Different Light Conditions: Three Strains of Prochlorococcus marinus. *Journal of*

Physical Chemistry B **121**: 9081–9090.

Hoang DT, Chernomor O, von Haeseler A, Minh BQ, Vinh LS. **2018**. UFBoot2: Improving the

Ultrafast Bootstrap Approximation. Molecular biology and evolution. *Molecular Biology and*

Evolution **35**: 518–522.

Holzwarth AR, Miloslavina Y, Nilkens M, Jahns P. **2009**. Identification of two quenching sites

active in the regulation of photosynthetic light-harvesting studied by time-resolved fluorescence.

Chemical Physics Letters **483**: 262–267.

Hori K, Maruyama F, Fujisawa T, Togashi T, Yamamoto N, Seo M, Sato S, Yamada T, Mori

H, Tajima N, *et al.* 2014. Klebsormidium flaccidum genome reveals primary factors for plant

terrestrial adaptation. *Nature Communications* **5**.

Huang W, Daboussi F. **2017**. Genetic and metabolic engineering in diatoms. *Philosophical*

Transactions of the Royal Society B: Biological Sciences **372**.

Ifuku K, Yan D. **2020**. *Efficient Transformation of the Diatoms Phaeodactylum tricornutum by*

Multipulse Electroporation. Humana, New York, NY.

Ifuku K, Yan D, Miyahara M, Inoue-Kashino N, Yamamoto YY, Kashino Y. **2015**. A stable and

efficient nuclear transformation system for the diatom *Chaetoceros gracilis*. *Photosynthesis Research*

123: 203–211.

Im SH, Lepetit B, Mosesso N, Shrestha S, Weiss L, Nymark M, Roellig R, Wilhelm C, Isono

E, Kroth PG. 2024. Identification of promoter targets by Aureochrome 1a in the diatom

Phaeodactylum tricornutum. *Journal of Experimental Botany* **75**: 1834–1851.

Ishihara T, Ifuku K, Yamashita E, Fukunaga Y, Nishino Y, Miyazawa A, Kashino Y, Inoue-

Kashino N. **2015**. Utilization of light by fucoxanthin–chlorophyll-binding protein in a marine centric

diatom, *Chaetoceros gracilis*. *Photosynthesis Research* **126**: 437–447.

Järvi S, Suorsa M, Paakkarinen V, Aro EM. **2011**. Optimized native gel systems for separation

of thylakoid protein complexes: Novel super- and mega-complexes. *Biochemical Journal* **439**: 207–

214.

Jeffrey SW, Humphrey GF. **1975**. New spectrophotometric equations for determining

chlorophylls a, b, c_1 and c_2 in higher plants, algae and natural phytoplankton. *Biochemie und*

Physiologie der Pflanzen **167**: 191–194.

Kalyaanamoorthy S, Minh BQ, Wong TKF, von Haeseler A, Jermiin LS. **2017**. ModelFinder:

fast model selection for accurate phylogenetic estimates. *Nature Methods* **14**: 587–589.

Kameo S, Aso M, Furukawa R, Matsumae R, Yokono M, Fujita T, Tanaka A, Tanaka R,

Takabayashi A. **2021**. Substitution of Deoxycholate with the Amphiphilic Polymer Amphipol A8-35

Improves the Stability of Large Protein Complexes during Native Electrophoresis. *Plant and Cell*

Physiology **62**: 348–355.

Katoh K, Standley DM. **2013**. MAFFT Multiple Sequence Alignment Software Version 7:

Improvements in Performance and Usability. *Molecular Biology and Evolution* **30**: 772–780.

Koziol AG, Borza T, Ishida KI, Keeling P, Lee RW, Durnford DG. **2007**. Tracing the evolution

of the light-harvesting antennae in chlorophyll *a*/*b*-containing organisms. *Plant Physiology* **143**: 1802–

1816.

Kumazawa M, Ifuku K. **2024**. Unraveling the evolutionary trajectory of LHCI in red-lineage

algae: Conservation, diversification, and neolocalization. *iScience* **27**: 110897.

Kumazawa M, Nishide H, Nagao R, Inoue-Kashino N, Shen J, Nakano T, Uchiyama I, Kashino

Y, Ifuku K. **2022**. Molecular phylogeny of fucoxanthin-chlorophyll *a* / *c* proteins from *Chaetoceros*

gracilis and Lhcq/Lhcf diversity. *Physiologia Plantarum* **174**: e13598.

Lavaud J, Kroth PG. **2006**. In diatoms, the transthylakoid proton gradient regulates the

photoprotective non-photochemical fluorescence quenching beyond its control on the xanthophyll

cycle. *Plant and Cell Physiology* **47**: 1010–1016.

Lavaud J, Lepetit B. **2013**. An explanation for the inter-species variability of the photoprotective

non-photochemical chlorophyll fluorescence quenching in diatoms. *Biochimica et Biophysica Acta -*

Bioenergetics **1827**: 294–302.

Leliaert F, Smith DR, Moreau H, Herron MD, Verbruggen H, Delwiche CF, De Clerck O. **2012**.

Phylogeny and Molecular Evolution of the Green Algae. *Critical Reviews in Plant Sciences* **31**: 1–46.

Lepetit B, Volke D, Gilbert M, Wilhelm C, Goss R. **2010**. Evidence for the existence of one

antenna-associated, lipid-dissolved and two protein-bound pools of diadinoxanthin cycle pigments in

diatoms. *Plant Physiology* **154**: 1905–1920.

Letunic I, Bork P. **2024**. Interactive Tree of Life (iTOL) v6: Recent updates to the phylogenetic

tree display and annotation tool. *Nucleic Acids Research* **52**: W78–W82.

Lohr M, Wilhelm C. **1999**. *Algae displaying the diadinoxanthin cycle also possess the violaxanthin*

cycle.

Malnoë A. **2018**. Photoinhibition or photoprotection of photosynthesis? Update on the (newly

termed) sustained quenching component qH. *Environmental and Experimental Botany* **154**: 123–133.

Manuel Zapata, Francisco Rodríguez, José L. Garrido. **2000**. Separation of chlorophylls and

carotenoids from marine phytoplankton: a new HPLC method using a reversed phase C8 column and

pyridine-containing mobile phases. *Marine Ecology Progress Series* **195**: 29–45.

Miloslavina Y, Grouneva I, Lambrev PH, Lepetit B, Goss R, Wilhelm C, Holzwarth AR. **2009**.

Ultrafast fluorescence study on the location and mechanism of non-photochemical quenching in

diatoms. *Biochimica et Biophysica Acta - Bioenergetics* **1787**: 1189–1197.

Minagawa J, Tokutsu R. **2015**. Dynamic regulation of photosynthesis in *Chlamydomonas*

reinhardtii. *The Plant Journal* **82**: 413–428.

Minh BQ, Dang CC, Vinh LS, Lanfear R. **2021**. QMaker: Fast and Accurate Method to Estimate

Empirical Models of Protein Evolution. *Systematic Biology* **70**: 1046–1060.

Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear

R. **2020**. IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic

Era (E Teeling, Ed.). *Molecular Biology and Evolution* **37**: 1530–1534.

Miyahara M, Aoi M, Inoue-Kashino N, Kashino Y, Ifuku K. **2013**. Highly efficient

transformation of the diatom *Phaeodactylum tricornutum* by multi-pulse electroporation. *Bioscience,*

Biotechnology and Biochemistry **77**: 874–876.

Nagao R, Ishii A, Tada O, Suzuki T, Dohmae N, Okumura A, Iwai M, Takahashi T, Kashino

Y, Enami I. 2007. Isolation and characterization of oxygen-evolving thylakoid membranes and

Photosystem II particles from a marine diatom *Chaetoceros gracilis*. *Biochimica et Biophysica Acta -*

Bioenergetics **1767**: 1353–1362.

Nagao R, Kato K, Ifuku K, Suzuki T, Kumazawa M, Uchiyama I, Kashino Y, Dohmae N,

Akimoto S, Shen J-R, *et al.* 2020a. Structural basis for assembly and function of a diatom

photosystem I-light-harvesting supercomplex. *Nature Communications* **11**: 2481.

Nagao R, Kato K, Kumazawa M, Ifuku K, Yokono M, Suzuki T, Dohmae N, Akita F, Akimoto

S, Miyazaki N, *et al.* 2022. Structural basis for different types of hetero-tetrameric light-harvesting

complexes in a diatom PSII-FCPII supercomplex. *Nature Communications* **13**: 1764.

Nagao R, Kato K, Suzuki T, Ifuku K, Uchiyama I, Kashino Y, Dohmae N, Akimoto S, Shen J-

R, Miyazaki N, *et al.* 2019. Structural basis for energy harvesting and dissipation in a diatom PSII–

FCPII supercomplex. *Nature Plants* **5**: 890–901.

Nagao R, Tomo T, Noguchi E, Nakajima S, Suzuki T, Okumura A, Kashino Y, Mimuro M,

Ikeuchi M, Enami I. 2010. Purification and characterization of a stable oxygen-evolving Photosystem

II complex from a marine centric diatom, *Chaetoceros gracilis*. *Biochimica et Biophysica Acta -*

Bioenergetics **1797**: 160–166.

Nagao R, Tomo T, Noguchi E, Suzuki T, Okumura A, Narikawa R, Enami I, Ikeuchi M. 2012.

Proteases are associated with a minor fucoxanthin chlorophyll *a*/*c*-binding protein from the diatom,

Chaetoceros gracilis. *Biochimica et Biophysica Acta - Bioenergetics* **1817**: 2110–2117.

Nagao R, Ueno Y, Akimoto S, Shen JR. 2020b. Effects of CO_2 and temperature on photosynthetic

performance in the diatom *Chaetoceros gracilis*. *Photosynthesis Research* **146**: 189–195.

Nagao R, Yokono M, Akimoto S, Tomo T. **2013**. High excitation energy quenching in fucoxanthin

chlorophyll *a*/*c*-binding protein complexes from the diatom *Chaetoceros gracilis*. *Journal of Physical*

Chemistry B **117**: 6888–6895.

Nagao R, Yokono M, Ueno Y, Shen JR, Akimoto S. **2020c**. Excitation-Energy Transfer and

Quenching in Diatom PSI-FCPI upon P700 Cation Formation. *Journal of Physical Chemistry B* **124**:

1481–1486.

Nakayasu M, Akimoto S, Yoneda K, Ikuta S, Shimakawa G, Matsuda Y. **2024**. Different

functions of Lhcx isoforms in photoprotective mechanism in the marine diatom *Thalassiosira*

pseudonana. *BioRxiv*.

Nedbal L, Trtflek M, Kaftan D. **1999**. *Flash fluorescence induction: a novel method to study*

regulation of Photosystem II.

Niyogi KK. **2000**. Safety valves for photosynthesis. *Current Opinion in Plant Biology* **3**: 455–460.

Park S, Steen CJ, Lyska D, Fischer AL, Endelman B, Iwai M, Niyogi KK, Fleming GR. **2019**.

Chlorophyll–carotenoid excitation energy transfer and charge transfer in *Nannochloropsis oceanica*

for the regulation of photosynthesis. *Proceedings of the National Academy of Sciences of the United*

States of America **116**: 3385–3390.

Peers G, Truong TB, Ostendorf E, Busch A, Elrad D, Grossman AR, Hippler M, Niyogi KK.

2009. An ancient light-harvesting protein is critical for the regulation of algal photosynthesis. *Nature*

462: 518–521.

Petroutsos D, Tokutsu R, Maruyama S, Flori S, Greiner A, Magneschi L, Cusant L, Kottke T,

Mittag M, Hegemann P, *et al.* 2016. A blue-light photoreceptor mediates the feedback regulation of

photosynthesis. *Nature* **537**: 563–566.

Pierella Karlusich JJ, Cosnier K, Zinger L, Henry N, Nef C, Bernard G, Scalco E, Dvorak E,

Wincker P, Sunagawa S, *et al.* 2025. Patterns and drivers of diatom diversity and abundance in the

global ocean. *Nature Communications* **16**.

Pierella Karlusich JJ, Ibarbalz FM, Bowler C. **2020**. Phytoplankton in the *Tara* Ocean. *Annual*

Review of Marine Science **12**: 233–265.

Steenwyk JL, Buida TJ, Li Y, Shen X-X, Rokas A. **2020**. ClipKIT: A multiple sequence alignment

trimming software for accurate phylogenomic inference (A Hejnol, Ed.). *PLOS Biology* **18**: e3001007.

Strassert JFH, Irisarri I, Williams TA, Burki F. **2021**. A molecular timescale for eukaryote

evolution with implications for the origin of red algal-derived plastids. *Nature Communications* **12**:

1879.

Taddei L, Chukhutsina VU, Lepetit B, Stella GR, Bassi R, van Amerongen H, Bouly JP,

Jaubert M, Finazzi G, Falciatore A. **2018**. Dynamic changes between two LHCX-related energy

quenching sites control diatom photoacclimation. *Plant Physiology* **177**: 953–965.

Tian L, Nawrocki WJ, Liu X, Polukhina I, Van Stokkum IHM, Croce R. **2019**. PH dependence,

kinetics and light-harvesting regulation of nonphotochemical quenching in Chlamydomonas.

Proceedings of the National Academy of Sciences of the United States of America **116**: 8320–8325.

Tokutsu R, Fujimura-Kamada K, Matsuo T, Yamasaki T, Minagawa J. **2019a**. The

CONSTANS flowering complex controls the protective response of photosynthesis in the green alga

Chlamydomonas. *Nature Communications* **10**.

Tokutsu R, Fujimura-Kamada K, Yamasaki T, Matsuo T, Minagawa J. **2019b**. Isolation of

photoprotective signal transduction mutants by systematic bioluminescence screening in

Chlamydomonas reinhardtii. *Scientific Reports* **9**.

Wang W, Yu L-J, Xu C, Tomizaki T, Zhao S, Umena Y, Chen X, Qin X, Xin Y, Suga M, *et al.*

2019. Structural basis for blue-green light harvesting and energy dissipation in diatoms. *Science* **363**.

Xu C, Pi X, Huang Y, Han G, Chen X, Qin X, Huang G, Zhao S, Yang Y, Kuang T, *et al.* **2020**.

Structural basis for energy transfer in a huge diatom PSI-FCPI supercomplex. *Nature Communications*

11: 5081.

Yamamoto Y, Leng J, Shen J-R. **2011**. Isolation of Photosystem II-Enriched Membranes and the

Oxygen-Evolving Complex Subunit Proteins from Higher Plants. In: Carpentier R, ed. Photosynthesis

Research Protocols. Totowa, NJ: Humana Press, 1–10.

Yang YN, Le TTL, Hwang JH, Zulfugarov IS, Kim EH, Kim HU, Jeon JS, Lee DH, Lee CH.

2022. High Light Acclimation Mechanisms Deficient in a PsbS-Knockout Arabidopsis Mutant.

International Journal of Molecular Sciences **23**.

Ye Z, Sawada M, Iwasa M, Moriyama R, Dey D, Furutani M, Kitao M, Hara T, Tanaka A,

Kishimoto J, *et al.* **2025**. Revisiting the early light-induced protein hypothesis in the sustained thermal

dissipation mechanism in yew leaves. *Journal of Experimental Botany* **76**: 513–531.

Yokono M, Akimoto S, Tanaka A. **2008**. Seasonal changes of excitation energy transfer and

thylakoid stacking in the evergreen tree *Taxus cuspidata*: How does it divert excess energy from

photosynthetic reaction center? *Biochimica et Biophysica Acta - Bioenergetics* **1777**: 379–387.

Zhang T, Gao Y, Wang R, Zhao Y. **2017**. Production of Guide RNAs in vitro and in vivo for

CRISPR Using Ribozymes and RNA Polymerase II Promoters. *Bio Protoc.* **7**.

Zhang H, Xiong X, Guo K, Zheng M, Cao T, Yang Y, Song J, Cen J, Zhang J, Jiang Y, *et al.*

2024. A rapid aureochrome opto-switch enables diatom acclimation to dynamic light. *Nature*

Communications **15**: 5578.

Zhao S, Shen L, Li X, Tao Q, Li Z, Xu C, Zhou C, Yang Y, Sang M, Han G, *et al.* **2023**. Structural

insights into photosystem II supercomplex and trimeric FCP antennae of a centric diatom *Cyclotella*

meneghiniana. *Nature Communications* **14**: 8164.

Zheng M, Pang X, Chen M, Tian L. **2024**. Ultrafast energy quenching mechanism of LHCSR3-

dependent photoprotection in Chlamydomonas. *Nature communications* **15**: 4437.

Zhou C, Feng Y, Li Z, Shen L, Li X, Wang Y, Han G, Kuang T, Liu C, Shen J-R, *et al.* **2024**.

Structural and spectroscopic insights into fucoxanthin chlorophyll a/c-binding proteins of diatoms in

diverse oligomeric states. *Plant Communications* **5**: 101041.

1000

1001

Figure 1. Molecular Phylogenetic Tree of Light-Harvesting Complexes (LHCs) from Red and

Green Evolutionary Lineages

LHC sequences were obtained from red algae, green algae *Mesostigma viride* NIES-296

(Mesvir1_1) and *Klebsormidium nitens* NIES-2285 (Klepit1_1), diatoms: *Chaetoceros gracilis* (Cg),

Thalassiosira pseudonana (Tp), *Cyclotella cryptica* (Cc), *Phaeodactylum tricornutum* (Pt), and

cryptophytes: *Guillardia theta* CCMP2712 and Cryptophyceae sp. (*Baffinella* sp.) CCMP2293. These

were combined with reference sequences for Lhcsr from *Chlamydomonas reinhardtii*: Lhcsr1

(UniProt: P93664) and Lhcsr3.1 (UniProt: P0DO19), which is identical to Lhcsr3.2 (UniProt:

P0DO18), and a molecular phylogenetic tree was constructed using IQ-TREE2. The phylogenetic tree

is displayed as an unrooted tree, and the inset tree is rerooted with the other LHC family members

except for Lhcx, Lhcf, and CgLhcr9 homologs. Detailed annotations are shown only for the

Lhcx/Lhcsr subfamily. Circles on nodes indicate ultra-fast bootstrap support values of 90% or higher.

Numbers in parentheses indicate SH-aLRT support values (%) / aBayes support values / ultra-fast

bootstrap support values (%). Numbers in the inset tree are ultra-fast bootstrap support values (%).

1016

1017

Figure 2. PAM Fluorescence Measurements of *C. gracilis* Wild-Type and *lhcx1* Strains Cultured

with Aeration for 7–8 Days Under Low Light (LL) and High Light (HL)

A) NPQ of LL and HL cultures. B) qP of LL and HL cultures. C) PSII effective quantum yield Y(II)

of LL and HL cultures. * indicates that adjusted $p < 0.05$ by Tukey's HSD test was satisfied between

the wild-type and all *lhcx1* strains. The points in each figure represent mean ± standard deviation with

biological replicates of $n = 3$.

1024

1025

Figure 3. Pigment Analysis of Low Light (LL) and High Light (HL) Cultures

A) Chlorophyll (Chl) *a* content per 10^9 cells. B) Fucoxanthin (Fx) content per Chl *a*. C) Diatoxanthin (Dtx) content per Chl *a*. D) Diadinoxanthin (Ddx) content per Chl *a*. E) De-epoxidation state (DEPS = [Dtx]/([Dtx]+[Ddx])). Data are presented as mean ± standard deviation (*n* = 3). Different alphabets indicate adjusted *p* < 0.05 by Tukey's HSD test.

1031

Figure 4. Low-Temperature Fluorescence Emission Spectra of Wild-Type (WT) and *lhcx1*

Mutants Cultured Under High Light

A) Low-temperature fluorescence emission spectra measured after 30 minutes of dark treatment

followed by rapid freezing in liquid nitrogen. B) Low-temperature fluorescence emission spectra

measured after 30 minutes of HL treatment followed by rapid freezing in liquid nitrogen. Each

spectrum was normalized at the peak corresponding to CP47 (693–695 nm). The wild type was

measured in two separately cultured samples (black solid line and black dashed line).

Figure 5. Fluorescence Decay-Associated Spectra of Time-Resolved Fluorescence

Black lines indicate WT, blue lines indicate *lhcx1-7*, solid lines represent measurements after 30 minutes of dark treatment, and dashed lines represent measurements after 30 minutes of treatment at 300 μmol photons $m^{-2}\, s^{-1}$.

1043

Figure 6. 2D-Amphipol-based-CN/SDS-PAGE of *C. gracilis* Thylakoid Membranes

A) Amphipol-based CN-PAGE of *C. gracilis* and spinach thylakoid membranes. Annotation of each

band for spinach follows previous reports (Järvi *et al.*, 2011; Kameo *et al.*, 2021). B) Protein

fluorescence staining image of two-dimensional electrophoresis of Amphipol-based CN-PAGE.

Annotation of each spot indicated by red arrowheads is based on previous reports (Nagao *et al.*, 2007).

FCP-A, FCP-B, and FCP-C correspond to CgLhcf1, CgLhcf12, and CgLhcf2, respectively (Nagao *et*

al., 2019, 2022; Zhou *et al.*, 2024). C) Identification of the Lhcx1 spot by immunoblotting using anti-

Lhcx1 antibody. The blue arrowhead indicates the detected spot.

1052

Figure 7. PAM Fluorescence Measurements of *C. gracilis* Wild-Type and *lhcx1* Strains

Cultured with 3% CO₂ Aeration for three Days Under Low Light (LL) and High Light (HL)

A) NPQ of LL and HL cultures. B) qP of LL and HL cultures. C) PSII effective quantum yield Y(II) of LL and HL cultures. The points in each figure represent mean ± standard deviation with biological replicates of *n* = 3.

A

Common HL-induced genes in Air condition in both WT and *lhcx1*

B

Common *lhcx1* enhanced genes in both Air and CO₂ conditions

1058

Figure 8. Gene Expression Analysis under CO₂ Enrichment

KEGG pathway enrichment analysis of differentially expressed genes in the *lhcx1* mutant under LL

and HL conditions. A**)** Pathways enriched in genes commonly induced by HL in both WT and the

lhcx1 mutant under ambient air conditions. B**)** Pathways enriched in genes showing enhanced HL

response in *lhcx1* mutant under both ambient air and CO₂-enriched conditions. Dot size represents

gene count, and color intensity indicates statistical significance (q-value shown as p.adjust). Pathways

related to typical HL response and photosynthesis are indicated with red letters.