# Investigating high pathogenicity avian influenza virus incursions to remote islands: Detection of H5N1 on Gough Island in the South Atlantic Ocean

Antje Steinfurth[1,@], Joshua G. Lynton-Jenkins[2, 3], Jaimie Cleeland[4], Benjamin C. Mollett[2], Holly A. Coombes[2], Andrea Moores[2], Robyn Neal[2], Ben Clifton[2], Marco Falchieri[2], Christopher W. Jones [1], Michelle M. Risi [1], Susannah Gold[1], Joe James[2, 3], Peter G. Ryan[5], Jacob González- Solís[6,7] and Ashley C. Banyard[2, 3,@]

[1] Royal Society for the Protection of Birds (RSPB), Centre for Conservation Science, Pembroke Street, Cambridge, CB2 3QZ, UK

[2] Influenza and Avian Virology workgroup, Department of Virology, Animal and Plant Health Agency (APHA-Weybridge), Woodham Lane, Addlestone, Surrey KT15 3NB, UK

[3] WOAH/FAO International Reference Laboratory for Avian Influenza, Animal and Plant Health Agency (APHA-Weybridge), Woodham Lane, Addlestone, Surrey KT15 3NB, UK

[4] British Antarctic Survey, High Cross, Madingley Road, Cambridge, CB3 0ET, UK

[5] FitzPatrick Institute of African Ornithology, University of Cape Town, Rondebosch 7700, South Africa

[6] Departament de Biologia Evolutiva, Ecologia i Ciències Ambientals (BEECA), Universitat de Barcelona (UB), Av. Diagonal 643, 08028 Barcelona, Spain

[7] Institut de Recerca de la Biodiversitat (IRBio), Universitat de Barcelona (UB), Barcelona, Spain

23    @ Corresponding author email: Antje.Steinfurth@rspb.org.uk/Ashley.Banyard@apha.gov.uk

## Abstract

Understanding the mechanisms underlying the emergence and spread of high pathogenicity avian

influenza virus (HPAIV) is critical for tracking its global dissemination, particularly via migratory

seabirds, given their role in transmission over long distances. Scavenging seabirds, such as skuas, may

act as both reservoirs and vectors, and have been linked to multiple outbreaks since 2021. Here, we

report the detection of HPAIV H5N1 clade 2.3.4.4b in three Tristan skua (*Stercorarius antarcticus*

*hamiltoni*) carcasses on Gough Island in the central South Atlantic Ocean. To investigate potential

incursion routes, we combined genomic analyses with year-round tracking data from global location

sensors. Although migratory movement patterns suggested southern Africa as the most obvious

pathway, the strain detected on Gough Island was more closely related to that identified in South

Georgia, indicating that infection may have occurred during the pre-laying exodus, when skuas

disperse into frontal waters south of the island. No further cases have been confirmed for Gough, but

more systematic monitoring is needed to understand the dynamics of virus infection. The detection of

HPAIV H5N1 in skuas on Gough Island highlights the importance of continued vigilance, proactive

and geographically inclusive surveillance strategies, and biosecurity measures globally, alongside

efforts to reduce other pressures on globally important seabird populations to help strengthen their

resilience.

**Keywords (5-8)** High pathogenicity avian influenza, Gough Island, Brown Skua, migration,

transmission, emerging

## Introduction

The ongoing expansion of high pathogenicity avian influenza virus (HPAIV) poses a major threat to

wildlife, livestock, and, through its zoonotic potential, human health. Since first emerging in poultry

in China in 1996, the A/goose/Guangdong/1/96 (GsGd)-lineage of HPAIV H5Nx has evolved to

spread efficiently among a broad range of bird species globally, causing mass mortality events in

unprecedented numbers [1-6]. Following its dissemination across Europe by migratory species, H5N1

reached North America via the East Atlantic flyway and, by October 2022, had expanded rapidly

throughout South America, with devastating impacts on both seabird and pinniped populations [7,8].

In the austral spring of 2023, HPAIV reached the sub-Antarctic islands and Antarctica, first detected

in brown skuas (*Stercorarius antarcticus*) on Bird Island, South Georgia [9]. Soon after, cases were

reported in southern fulmars (*Fulmarus glacialoides*) and black-browed albatrosses (*Thalassarche*

*melanophris*) in the Falkland Islands [9], and in multiple species across the Antarctic Peninsula [10].

Genetic analyses indicate that the detections in these regions were linked to strains originating in

South America, likely introduced via long-distance movements of scavenging seabirds such as skuas,

gulls, and giant petrels [9, 11]. In the following breeding season 2023/2024, the virus had spread

further east across the sub-Antarctic and into the southern Indian Ocean, causing significant die-offs

in wandering albatross (*Diomedea exulans*) chicks on Marion Island [12], before it was reported for

the Crozet and Kerguelen Archipelagos, where mass mortality events predominantly affected southern

elephant seals (*Mirounga leonina*) [11]. Genetic evidence suggests that the Crozet and Kerguelen

outbreaks resulted from independent introductions linked to South Georgia rather than from the nearer

southern African coast [11].

Gough Island in the central South Atlantic is situated approximately 3,600 km northeast of South

Georgia and 2,450 km northwest of Marion Island and considered one of the most important seabird

breeding sites globally [13]. Despite its small size (~65 km²), the island supports an estimated eight

million birds of at least 24 species, many of them endemic or near-endemic to the island or island

group of Tristan da Cunha, and several of global conservation concern, including the Critically

Endangered Tristan albatross (*Diomedea dabbenena*), MacGillivray's prion (*Pachyptila*

*macgillivrayi*), and Gough finch (*Rowettia goughensis*) [13-16]. Consequently, the potential for the

arrival of HPAIV into this sensitive environment has remained a significant conservation concern. A

risk assessment in 2022 considered the likelihood of the emergence of HPAIV on Gough Island to be

'low' [17]. However, the spread of the HPAIV epizootic into the southern hemisphere and its

continued expansion across the sub-Antarctic region have substantially increased the risk of incursion.

Recurrent HPAIV outbreaks have also been reported in coastal areas of southern Africa, following

viral spread from Europe via the Afro-Eurasian flyway [18-20]. Therefore, Gough Island's unique

geographical position in the South Atlantic along major migratory flyways and marine corridors,

linking the Atlantic and Southern Oceans, renders it vulnerable to HPAIV introduction from multiple

directions [21].

Antibodies against AIVs in Gough's breeding seabird population were first detected in 2009, when a

serosurvey on Gough Island revealed exposure in brown skuas (*S. a. hamiltoni;* hereafter referred to

as Tristan skua) and northern rockhopper penguins (*Eudyptes moseleyi*), demonstrating that influenza

A viruses have previously reached the island and circulated among these species [22, 23]. The Tristan

skua is endemic to the island group of Tristan da Cunha, with most of its population breeding on

Gough [24]. Given their scavenging behaviour, skuas represent both key reservoirs and vectors for

disease transmission, making the study of their movements critical for identifying introduction

pathways [23, 25].

Here, we describe the detection of HPAIV H5N1 clade 2.3.4.4b in three Tristan skuas from Gough

Island in September 2024 and investigate potential incursion routes by using a combination of genetic

analyses of the virus and year-round tracking data from global location sensors.

**Materials & Methods**

**Study site**

Gough Island (40°19′12″S 09°56′24″W) is part of the British Overseas Territory of St Helena,

Ascension, and Tristan da Cunha in the central South Atlantic Ocean (Fig.1). Gough Island lies

approximately 380 km south-southeast from the Tristan da Cunha archipelago, which consists of three

main islands: Tristan da Cunha (96 km²), Inaccessible (14 km²), and Nightingale (4 km²), along with

its satellite islets Middle (or Alex) and Stoltenhoff (both ~0.1 km²), all within 40 km of each other

(Fig. 1). Gough is uninhabited except for a small year-round team of 7-9 people operating the research

station. The station is resupplied once a year in September/October by the South African research

vessel *SA Agulhas II*. A field team of 2-3 biologists is on the island year-round as part of the island's

long-term term monitoring and science programme, trained to provide surveillance for unusual avian

mortality events.

**Tracking data**

Migrate Technology Intigeo□C330 geolocation tags (17 x 19 x 8 mm, 3.3 g) were deployed on 10

incubating adult Tristan skuas near the Gough Island research station in October-November 2017. The

tags were attached to standard stainless steel SAFRING rings. Eight tags were retrieved during the

following breeding season; the two other devices were not recovered.

Light-level data were processed in R version 4.4.1 using the *SGAT* package (v0.1.3) [26],

implementing a Bayesian state-space model to estimate probable locations based on the timing of

twilight transitions, while accounting for uncertainties in light detection and movement behaviour.

Raw light data were pre-processed using the *BAStag* package (v0.1.3) [27] to identify twilight events

(sunrise and sunset) based on light intensity thresholds. These twilight times were then manually

reviewed and edited to correct for anomalies caused by shading, cloud cover, or behavioural factors

e.g., incubation. The resultant data were used to infer two positions per day, corresponding to daily

twilight transitions.

To quantify space use, we calculated kernel utilisation distributions (KUDs) using the *adehabitatHR*

package (v0.4.22) [28]. *SGAT*-derived location estimates were projected onto an equal-area coordinate

reference system to allow for unbiased spatial analysis. For each bird, 50% and 95% KUDs were

calculated to represent core use areas and home ranges, respectively, across the pre-laying, breeding,

and non-breeding periods. Location data during incubation in 2017 and 2018 were pooled for KUD

analysis. To evaluate population-level space use, individual KUDs were averaged to assess spatial

habitat use. Migration pathways between breeding and non-breeding areas were excluded from the

KUD analysis and instead visualised using raw location data. For each individual, the onset of north-

east migration away from the breeding colony and that of south-west return migration was defined as

the point at which the individual's distance from the colony increased or decreased continuously,

without subsequent decreases/increases. The end point of each migration phase was defined as the

point at which the distance from the colony stabilised, indicating arrival at a new residency area. All

spatial operations were conducted using the *sp*, *sf*, and *raster* packages in R.

While we acknowledge that the GLS data used in this study were collected up to six years prior to the

HPAI detection, we expect movement patterns of this population to have remained largely consistent.

Long-term (multiyear) fidelity to specific wintering sites is common in migratory seabirds,

particularly when these sites are associated with predictable environments such as upwelling systems,

where repeated use can increase familiarity with local resources and prey dynamics [29-31]. Although

climate change may gradually alter the predictability of marine environments  [32,33], the persistence

of the wintering site used by Tristan skuas across multiple breeding seasons (2009/10 (5 tracks),

2010/11 (9), 2011/12 (6), and 2017/18 (8)) spanning nearly a decade [34, this study], suggests that this

fidelity continues to confer ecological benefits and thus is likely representative of the wider Gough

Island skua population today.

**Sample collection**

Swab samples from the brain, trachea, and cloaca of dead birds were collected, frozen at −20°C

within one hour of collection, and shipped to the World Organisation for Animal Health/Food and

Agriculture Organisation (WOAH/FAO) International Reference Laboratory for Avian Influenza at

the Animal and Plant Health Agency (APHA) in Weybridge, UK, for diagnostic evaluation.

Surveillance for suspicious wildlife mortality continued with special attention given to any spatial or

temporal clustering of cases or unusual behaviours in wildlife.

**Molecular methods**

Total nucleic acid was extracted from all swab samples as described previously [35] and viral RNA

was screened using four real-time reverse transcription polymerase chain reaction (rRT-PCR) assays

including: i) a Matrix (M)-gene assay for generic influenza A virus detection [36], ii) a HPAIV H5

clade 2.3.4.4b specific assay for HA subtype and pathotyping [35], iii) an N1-subtype specific rRT-

PCR to confirm the neuraminidase type [37], and iv) an avian paramyxovirus type-1 (APMV-1) large

polymerase (L)-gene assay [38]. All rRT-PCRs were undertaken on the AriaMx qPCR System

(Agilent, United Kingdom). Material from positive brain tissue swabs was used for virus isolation in

10-day-old specific pathogen-free (SPF) embryonated fowls' eggs according to internationally

recognised methods (Codes and Manuals - WOAH - World Organisation for Animal Health).

**Viral Sequencing**

Whole-genome sequencing (WGS) was undertaken on brain tissue swabs. Extracted vRNA was

converted to double-stranded cDNA and amplified using a one-step RT-PCR using SuperScript III

One-Step RT-PCR kit (Thermo Fisher Scientific) as previously described [39]. PCR products were

purified with Agencourt AMPure XP beads (Beckman Coultrer) prior to sequencing library

preparation using the Native Barcoding Kit (Oxford Nanopore Technologies) and sequenced using a

GridION Mk1 (Oxford Nanopore Technologies) according to manufacturer's instructions. Assembly

of the influenza A viral genomes was performed using a custom in-house pipeline as previously

described [9].

**Phylogenetic Analysis**

To identify the genetic relatedness of Gough Island viruses to other reported strains, HPAIV H5N1

clade 2.3.4.4b sequences from Europe, Africa, Antarctica and South America available in the EpiFlu

database between 1$^{st}$ January 2020 and 28$^{th}$ July 2025 were collated. To remove over-represented

groups, the sequences were subset to cover 0.5% sequence divergence using PARNAS [40]. The

remaining dataset was separated by segment and aligned using Mafft v7.525 [41] and manually

trimmed to the open reading frame using Aliview v.2021 [42]. The trimmed alignments were used to

infer maximum-likelihood (ML) phylogenetic trees using IQ-Tree v2.4.0 [43] with model finder, 1000

ultrafast bootstraps and SH-like approximate likelihood ratio test. Clade classification of Gough

Island sequences was confirmed with GenoFLU-multi https://github.com/moncla-lab/GenoFLU-

multi.

To produce a time scale phylogeny, all H5 HA sequences from South America and Antarctica were

used alongside a subset of North American sequences to improve temporal signal. The dataset was

aligned, trimmed, and an ML phylogenetic tree was inferred using the same approaches as described

above. The HA ML phylogeny was then used to infer a time scaled phylogeny using TreeTime v0.11.4

[44]. Discrete trait analysis of transitions between locations was carried out using the mugration

inference model in TreeTime with default settings.

**Results**

The first suspected HPAIV-related death of an adult Tristan skua was observed on Gough Island on 12

September 2024 at a club of 50-200 non-breeding skuas at the helipad near the research station. The

following day, another skua at the same location exhibited clinical signs consistent with HPAIV,

including drooping head and wings, lethargy, and an inability to walk or fly, and was found dead later

the same day [45]. Two additional carcasses were discovered on 15 September. Swab samples were

collected from three of the four carcasses on 20 September 2025, and the carcasses were buried

afterwards. The fourth carcass had disappeared by the time of sampling and was presumed to have

been scavenged by other skuas.

*Diagnostic evaluation of samples*

All three birds sampled on Gough Island tested positive for HPAIV H5N1 in oropharyngeal and brain

swabs but tested negative in cloacal swabs (Table 1). Brain samples indicated a high viral load with

HPH5 rRT-PCR values ranging from 19.5 - 23.8 Cts. Virus isolation was attempted on two of the

three brain samples and both yielded hemagglutinating virus.

Whole genome sequences (WGS) obtained from two samples confirmed the viruses belonged to HA

clade 2.3.4.4b, genotype B3.2, as previously detected in the Antarctic region (Fig. 2a, Supplementary

Fig. S1) [9]. The two WGS shared >99% identity across all genes. Phylogenetic analysis shows that

these viruses are part of a monophyletic clade within the Antarctic Peninsula and sub-Antarctic

region, including ancestral viruses from South Georgia and their descendants in Crozet and Kerguelen

(Fig. 2) which is consistent with an expansion of the introductory pathway described by Banyard et al.

[9], rather than a new incursion from Africa. Analysis of the asymmetric transition rate matrix reveals

notable patterns of viral movement among key sub-Antarctic and South Atlantic islands (Fig. 2b).

South Georgia appears to be a significant ancestral source, exhibiting outward transitions to Crozet,

Kerguelen, Gough Island and the Falkland Islands. Gough Island demonstrates moderate connectivity

across the region, particularly with Crozet and Kerguelen. However, transition rates from Gough

Island to South Georgia and back suggest less frequent bidirectional exchange. Overall, the data

supports a model in which South Georgia serves as a primary source, seeding transmission across the

sub-Antarctic and South Atlantic islands (Fig. 2b).

*Ringing and tracking data*

Tracking durations ranged from 343 to 353 days (mean ± SD: 345 ± 3.3 days), yielding 687 ± 7.8

locations per individual (Fig. 3 and Fig. 4). Kernel utilisation distributions revealed clear seasonal

shifts in space use by Tristan skuas (Fig. 4a). During the pre-laying exodus, birds concentrated their

foraging activity in waters surrounding the Subantarctic front, with a mean 95% utilisation

distribution (home-range) of 977,900 km² and a corresponding 50% core area of 151,100 km². During

the breeding period, space use became more restricted around Gough Island, with a reduced mean

95% UD of 567,000 km² and a 50% core area of 128,900 km², consistent with central-place foraging

constraints. In contrast, the non-breeding period was characterised by a marked expansion and

displacement of activity areas to the South African and Namibian continental shelves, where skuas

occupied a mean 95% UD of 1 711,300 km² and a 50% core area of 394,400 km². Across these

periods, core areas consistently represented a smaller, high-use subset of the broader home-range,

while migration corridors were excluded from KUD estimation and are presented separately as

movement tracks.

Following the breeding season, individuals departed the foraging grounds around Gough Island

between 12 January and 12 February 2018, *en route* to non-breeding foraging grounds off the coasts

of South Africa and Namibia. Arrival at these non-breeding areas occurred between 20 January and 25

February 2018, with the north-eastward migration taking 11.9 ± 4.9 days (range 4–17 days).

Individuals remained within the non-breeding wintering grounds for 141 ± 19.5 days (range 109–165

days). Departures from these areas occurred between 4 June and 15 July 2018, with arrival back at the

pre-laying foraging area near Gough Island occurring between 13 June and 1 August 2018. The south-

westward return migration took $13.8 \pm 6.1$ days (range 7–23 days).

Two of the three dead skuas sampled for HPAIV had been ringed on Gough Island in 2016 and 2017.

The individual ringed in 2017 (SAFRING 888666) was a known breeder on the island and had been

equipped with a GLS device (BH142) as part of the tracking study (Fig. 4). In contrast, the skua

ringed in 2016 (SAFRING 888054) was caught as a non-breeder at the skua club near the helipad.

**Discussion**

The arrival of HPAIV H5N1 on Gough Island represents one of the most geographically isolated

detections to date, highlighting the interconnectivity of pelagic seabird populations and their role in

assisting global transmission in the current HPAIV H5N1 epizootic into some of the world's most

remote (and sensitive) ecosystems. Confirmation of HPAIV in Tristan skuas on Gough prompted

suspicions that individuals had become infected by viruses originating from Africa due to the species'

extensive connectivity with southern African coastal habitats outside their breeding season [34, this

study]. The GLS data analysed in this study corroborated this migratory pattern, suggesting high

temporal and spatial consistency in the species' use of the Benguela upwelling system as the species'

wintering foraging ground. Multiple outbreaks of HPAIV H5 clade 2.3.4.4b among seabird colonies in

southern Africa [18-20, 46] could have provided opportunities for transmission to scavenging species

such as skua. However, the phylogenetic analysis of global H5 HA clade 2.3.4.4b sequences places

the Gough Island viruses within a cluster of South American strains, distinct from viral strains

detected in Africa, and closely related to those found in South Georgia. An expansion of the

transmission pathway via South America and the sub-Antarctic is also supported by considering the

temporal context of the outbreaks in relation to seasonal and migratory movements within and across

species. Based on tracking records for 2018, Tristan skua complete their south-westward migration to

return to breeding sites in one to three weeks, arriving near Gough Island between mid-June and

beginning of August. While the incubation period for HPAIV can vary considerably depending on

species susceptibility and viral genetic characteristics, given the first observation of HPAIV on Gough

in mid-September, it is unlikely symptoms would only present 2–4 months after infection (assuming

broadly similar migratory timings in 2024). Although we cannot rule out the possibility that the virus

arrived first on the African coast before spreading to Gough Island, the timing of infection makes it

improbable that the skuas picked up the virus during their winter migration. Therefore, based on both

the phylogenetic analysis and outbreak timing, it is more likely that the virus reached Gough via

Southern or Atlantic Ocean flyways, potentially during the species' pre-laying exodus period in

August and September, when Tristan skuas disperse into the frontal region south of the island.

While terrestrial aggregations such as breeding colonies represent clear and obvious hotspots for virus

transmission, at-sea connectivity may play an equally critical role in transmission to remote locations

such as Gough Island. Overlapping foraging grounds and shared marine habitats increase the potential

for interspecies contact and facilitating viral spread between geographically distant populations.

Although there are currently no known reports of HPAIV transmission occurring at sea, scavenging

species are more likely than other seabird species to interact with carcasses, spend extended periods of

time sitting on the water, and come into close contact with other foraging individuals [45, 47, 48].

Such behaviours increase opportunities for viral transmission in circumstances where dilution of

virions in the environment would otherwise likely preclude infection, thereby supporting the role of

skuas and giant petrels as the most likely vectors of AIV across the Southern Ocean [23].

The nutrient-rich boundary waters between the sub-tropical and Antarctic polar fronts are known to

support large congregations of pelagic seabirds and serve as important foraging grounds for marine

mammals [21, 29, 49]. The high frequency and likelihood of interspecies interactions at these shared

foraging grounds, whether through direct or indirect contact, make such zones high-risk areas for the

persistence and transmission of infectious diseases. Seabird species breeding on Gough Island use

both the Southern and Atlantic Ocean Flyways [21, 32, 50-52], which overlap considerably between

30-60°S, linking Antarctic seabird populations with those on islands in the South Atlantic [21]. In

addition, populations breeding on South Georgia have been found to utilise waters near Gough Island

including southern elephant seals [53, 54], brown skua [47], and black-browed albatross *Thalassarche*

*melanophris* [31]. Ultimately, however, the timing and location of transmission of HPAIV to the

Tristan skuas in this study remain unknown, along with the uncertainty about whether, in fact, skuas

introduced the virus to the island or other avian or marine mammal species, either vagrants or those

breeding on Gough [50, 55-57].

The detection of HPAIV H5N1 originating from the Antarctic peninsula, in a species regularly

foraging in African coastal waters, illustrates a potentially novel eastward trans-oceanic circulation

loop, whereby influenza viruses of South American origin could be introduced into wild bird

populations of Africa, representing a reverse transmission pathway into the Afro-Eurasian flyway. The

Benguela upwelling region, stretching from Cape Agulhas in South Africa north to southwest Angola,

is one of the most productive marine ecosystems globally [58] and plays a vital role as a migratory

corridor and seasonal feeding area for numerous migratory species breeding in the southern Indian

Ocean, Antarctic or sub-Antarctic (e.g. White-capped/Shy albatross (*T. cauta*) [34]; Indian yellow-

nosed albatrosses (*T. carteri*) and northern giant petrels (*Macronectes halli*) [59]; black-browed

albatrosses [31]) [21]. Therefore, understanding the spatial and temporal dynamics of at-sea dispersal

and interactions, especially in areas of high biodiversity and migratory fly- and swim-ways, is

essential for anticipating and mitigating future outbreaks.

Initial concerns that HPAIV detection could constitute the onset of a mass outbreak on Gough Island,

have not materialized; no further symptomatic birds or mortalities related to HPAIV have been

confirmed (aside from a suspicious death of a breeding female Tristan albatross in April and one skua

in October 2025). Although mass mortality has occurred in other skua populations as a result of

HPAIV, notably in the UK where great skua (*Stercorarius* skua) declined by 73% [4, 45], the apparent

failure of the virus to cause a large-scale outbreak on Gough may be due to several factors. Cloacal

swabs taken during the initial investigation returned negative results for AIV, suggesting minimal

environmental shedding. Additionally, while one carcass was presumed scavenged, the remaining

three were buried following sampling, preventing further scavenging. Furthermore, while Gough

hosts large populations of seabird species, many of these species nest at low densities (e.g., Tristan

albatross) or in burrows, such as petrels and prions [13], reducing opportunities for direct contact,

inter-colony mixing and thus, viral spread across the island. A serological study in 2009 demonstrated

some seropositivity against influenza in northern rockhopper penguins and Tristan skua on Gough

[22, 23]; however, how far pre-existing immunity against AIVs could have influenced transmission

dynamics in this outbreak remains unknown.

Detection of outbreaks of HPAIV in wildlife, to infer transmission routes, and understand population

impacts, is dependent on reporting of wildlife mortality events and subsequent sampling and testing of

carcasses for AIV, with levels of coverage varying substantially across regions. Gaps in surveillance

lead to unsampled ancestry in sequence data, limiting the ability to resolve transmission chains from

the phylogenetic analysis. However, although surveillance for HPAIV remains limited across many

African countries [60], governmental and non-governmental agencies in South Africa and Namibia

routinely conduct both active and passive monitoring [18-20, 46, 60, 61]. Phylogenetic analysis of

2023 outbreak sequences in poultry and wild birds shows these strains originated from Europe via

West Africa [19]. Viruses sequenced in 2025 are likewise most closely related to the dominant

European strain of 2024/2025 [60, L. Roberts pers. comm.], with no South American cluster strains

detected in Africa to date. Thus, the strains reported in southern Africa continue to represent a distinct

cluster, separate from strains that spread via South America into the Antarctic region and our detection

reported here.

Wildlife disease monitoring of populations, particularly in parts of the polar and sub-Antarctic region,

currently remain extremely limited. The sheer number of sites and their remote location make any

systematic surveillance inherently challenging, and early or subclinical infections in wild or migratory

birds or mammals may go unnoticed for extended periods. Avian influenza may have also been

present in the Indian Ocean or Pacific sectors of the Southern Ocean before 2024. The absence of

detections in these areas, however, may reflect under-sampling rather than true absence, which poses a

barrier to early awareness of emerging issues and understanding of patterns of spread.

Although there have been no further mortalities attributed to HPAIV on Gough Island or in the

neighbouring Tristan da Cunha Archipelago, the impact of HPAIV in the region could be substantially

underestimated. Large parts of this remote group of islands are inaccessible, limiting timely

monitoring to detect carcasses or individuals exhibiting clinical signs. While there has been consistent

presence of a field team of biologists on Gough year-round since 2008, there is less systematic

surveillance at the Tristan da Cunha Archipelago. In February 2024, the island's fishery observer

reported a dead skua floating at sea off Nightingale Island, but no further unusual mortality of skuas

(or other seabirds) was observed on Nightingale and Inaccessible islands when visited in March and

September 2024, respectively (PGR pers. obs.). Therefore, the absence of confirmed cases should not

be taken as evidence of the absence of the disease.

The detection of HPAI on Gough Island is a stark reminder of the vulnerability of remote ecosystems

to infectious diseases, and for the need to alleviate already persisting pressures on globally important

seabird populations including habitat degradation and disturbance, invasive species, and climate-

driven stressors to strengthen their resilience. Proactive and geographically inclusive surveillance

strategies and biosecurity measures are vital to understand and mitigate the impacts of this disease,

rather than reactive responses after visible outbreaks. Understanding the spatiotemporal dynamics of

seabird movements -in particular scavenging seabirds- disease ecology, and host-pathogen

interactions is also essential for mitigating the impact of HPAIV and other emerging disease threats,

particularly in high-risk and ecologically sensitive areas [62].

**Acknowledgements**

All samples for HPAI testing were collected with the permission of the Tristan da Cunha government.

We are extremely grateful to Dr Laura Roberts, Western Cape Government and Dr Gretna DeWit,

Directorate of Animal Health, South Africa for their invaluable help issuing the South African import

permit. A special thank you to Ashley Bennison and Jennifer Forster Davidson, British Antarctic

Survey, for their generous and insightful advice and guidance on the Gough HPAIV response plan. We

would also like to thank the crews of the *SA Agulhas II* for their kind support and collaboration to

receive the samples onboard the vessel. The engagement, shipment to the UK, testing and generation

of the viral sequences was funded by the Department for Environment, Food and Rural Affairs (Defra,

UK) and the Devolved Administrations of Scotland and Wales, through the following programmes:

SV3006, SV3032 and SE2227. This work was also supported by the Biotechnology and Biological

Sciences Research Council (BBSRC) and Department for Environment, Food and Rural Affairs

(Defra, UK) research initiative 'FluTrailMap' [grant number BB/Y007271/1]. Funded by the

European Union (EU) under grant agreement (101084171) - (Kappa-Flu). Views and opinions

expressed are however those of the author(s) only and do not necessarily reflect those of the EU or

REA. Neither the EU nor the granting authority can be held responsible for them. The Science Animal

Ethics Committee (SFAEC) of the University of Cape Town approved the research protocols for the

deployments of GLS devices on seabirds including skuas (UCT SFAEC 2014\V10\PR), with

permission from the Tristan government.

**Declaration of interest statement**

The authors declare that they do not have any competing interests to disclose.

**References**

[1] Caliendo V, Lewis NS, Pohlmann A, et al. Transatlantic spread of highly pathogenic avian

influenza H5N1 by wild birds from Europe to North America in 2021. Sci Rep.

2022;12(1):11729. doi:10.1038/s41598-022-13447-z

[2] Falchieri M, Reid SM, Ross CS et al. Shift in HPAI infection dynamics causes significant losses in

seabird populations across Great Britain. Veterinary Record. 2022;191(7):294-6.

[3] Lane JV, Jeglinski JW, Avery☐Gomm S, et al. High pathogenicity avian influenza (H5N1) in

Northern Gannets (*Morus bassanus*): Global spread, clinical signs and demographic consequences.

Ibis. 2024;166(2):633-50. doi:10.1111/ibi.13275

[4] Tremlett CJ, Cleasby IR, Bolton et al. Declines in UK breeding populations of seabird species of

conservation concern following the outbreak of high pathogenicity avian influenza (HPAI) in 2021–

2022. Bird Study. 2024;71(4):293-310. doi:10.1080/00063657.2024.2438641

[5] Kuiken T, Vanstreels RE, Banyard A, et al. Emergence, spread, and impact of high☐pathogenicity

avian influenza H5 in wild birds and mammals of South America and

Antarctica. Conserv Biol. 2025;e70052. doi:10.1111/cobi.70052

[6] Pardo-Roa C, Nelson MI, Ariyama N, et al. Cross-species and mammal-to-mammal transmission

of clade 2.3. 4.4 b highly pathogenic avian influenza A/H5N1 with PB2 adaptations. Nat Commun.

2025;16(1):2232. doi:10.1038/s41467-025-57338-zdoi:10.1038/s41467-025-57338-z

[7] Leguia M, Garcia-Glaessner A, Muñoz-Saavedra B, et al. Highly pathogenic avian influenza A

(H5N1) in marine mammals and seabirds in Peru. Nat Commun.

2023;14(1):5489. doi:10.1038/s41467-023-41182-0

[8] Gamarra-Toledo V, Plaza PI, Angulo F, et al. Highly pathogenic avian influenza (HPAI) strongly

impacts wild birds in Peru. Biol Conserv. 2023;286:110272. doi:10.1016/j.biocon.2023.110272

[9] Banyard AC, Bennison A, Byrne AM, et al. Detection and spread of high pathogenicity avian

influenza virus H5N1 in the Antarctic Region. Nat Commun. 2024;15(1):7433. doi:10.1038/s41467-

024-51490-8

[10] León F, Ulloa-Contreras C, Pizarro EJ, et al. Skuas mortalities linked to positives HPAIV A/H5

beyond Polar Antarctic Circle. bioRxiv. 2025;1. doi:10.1101/2025.03.02.640960

[11] Clessin A, Briand FX, Tornos J, et al. Mass mortality events in the sub-Antarctic Indian

Ocean caused by long-distance circumpolar spread of highly pathogenic avian influenza H5N1 clade

2.3. 4.4 b. 2025. BioRxiv;2025-02. doi:10.1101/2025.02.25.640068

[12] Ryan PG, Gill R, Roberts L, Stephen V. Avian flu reaches Marion Island. African Birdlife.

2025;13(5):14-15.

[13] Caravaggi A, Cuthbert RJ, Ryan PG, et al. The impacts of introduced House Mice on the

breeding success of nesting seabirds on Gough Island. Ibis. 2019;161(3):648-

61. doi:10.1111/ibi.12664

[14] Oppel S, Clark BL, Risi MM, et al. Cryptic population decrease due to invasive species predation

in a long‑lived seabird supports need for eradication. J Appl Ecol. 2022;59(8):2059-

70. doi:10.1111/1365-2664.14218

[15] Jones CW, Risi MM, Osborne AM, et al. Mouse eradication is required to prevent local

extinction of an endangered seabird on an oceanic island. Anim Conserv. 2021;24(4):637-

45. doi:10.1111/acv.12670

[16] BirdLife International. *Rowettia goughensis*. *The IUCN Red List of Threatened Species* 2024:

e.T22723149A252044473. 2024 [cited on 07 August 2025]. doi:10.2305/IUCN.UK.2024-

2.RLTS.T22723149A252044473.en

[17] Dewar ML, Vanstreels RET, Boulinier T, et al. Biological Risk Assessment of Highly Pathogenic

Avian Influenza in the Southern Ocean. Scientific Committee on Antarctic Research. Cambridge

UK. SCAR Antarctic Wildlife Health Network. 2023.

[18] Abolnik C, Phiri T, Peyrot B, et al. The molecular epidemiology of clade 2.3.4.4B H5N1 high

pathogenicity avian influenza in Southern Africa, 2021–2022. Viruses.

2023;15(6):1383. doi:10.3390/v15061383

[19] Abolnik C, Roberts LC, Strydom C, et al. Outbreaks of H5N1 High Pathogenicity Avian

Influenza in South Africa in 2023 Were Caused by Two Distinct Sub-Genotypes of Clade 2.3. 4.4 b

Viruses.⬜Viruses.⬜2024;16(6):896. doi:10.3390/v16060896

[20] Roberts LC, Abolnik C, Waller LJ, et al. Descriptive epidemiology of and response to the high

pathogenicity avian influenza (H5N8) epidemic in South African coastal seabirds,

2018. Transbound Emerg Dis. 2023;2023(1):2708458. doi:10.1155/2023/2708458

[21] Morten JM, Carneiro AP, Beal M, et al. Global Marine Flyways Identified for Long⬜Distance

Migrating Seabirds From Tracking Data. Glob Ecol Biogeogr.

2025;34(2):e70004. doi:10.1111/geb.70004

[22] Abad, F.X., N.Busquets, A.Sanchez, et al. Serological and virological surveys of the influenza A

viruses in Antarctic and sub-Antarctic penguins. Antarct Sci. 2013;25:339-

344. doi:10.1017/S0954102012001228

[23] Gittins O, Grau-Roma L, Valle R, et al. Serological and molecular surveys of influenza A viruses

in Antarctic and sub-Antarctic wild birds. Antarct Sci. 2020;32(1):15-

20. doi:10.1017/s0954102019000464

[24] Ramsar Convention Secretariat. Ramsar Information Sheet (RIS) for Gough Island. Ramsar Sites

Information Service; 2025. Site No. 1868. Available from: Ramsar Sites Information Service.

[25] Gamble A, Bazire R, Delord K, et al. Predator and scavenger movements among and

within endangered seabird colonies: Opportunities for pathogen spread. J Appl Ecol. 2020;57(2):367-

78. doi:10.1111/1365-2664.13531

[26] Sumner MD, Wotherspoon SJ, Hindell MA. Bayesian estimation of animal movement from

archival and satellite tags. PLoS One. 2009.13;4(10):e7324. doi:10.1371/journal.pone.0007324

[27] Colchero F, Jones OR, Rebke M. BaSTA: an R package for Bayesian estimation of age specific

survival from incomplete mark–recapture/recovery data with

covariates. Methods Ecol Evol. 2012;3(3):466-470. doi: 10.1111/j.2041-210x.2012.00186.x

[28] Calenge C. Home range estimation in R: the adehabitatHR package. Office national de

la classe et de la faune sauvage: Saint Benoist, Auffargis, France. 2011.

[29] Weimerskirch H. Are seabirds foraging for unpredictable resources? Deep Sea Res II:

Top Stud Oceanogr. 2007;1;54(3-4):211-23. https://doi.org/10.1016/j.dsr2.2006.11.013

[30] Orben RA, Kokubun N, Fleishman AB, et al. Persistent annual migration patterns of a specialist

seabird. Marine Ecology Progress Series. 2018 Apr 12;593:231-45.

https://doi.org/10.3354/meps12459

[31] Phillips RA, Silk JR, Croxall JP, et al. Summer distribution and migration of nonbreeding

albatrosses: individual consistencies and implications for conservation. Ecology. 2005;86(9):2386-

96. doi:10.1890/04-1885

[32] Green CP, Green DB, Ratcliffe N, et al. Potential for redistribution of post moult habitat

for *Eudyptes* penguins in the Southern Ocean under future climate conditions. Glob Chang Biol.

2023;29(3):648-67. doi:10.1111/gcb.16500

[33] Lewin PJ, Wynn J, Arcos JM, et al. Climate change drives migratory range shift via individual

plasticity in shearwaters. Proc Natl Academ Sci, 2024;121(6), e2312438121.

[34] Carneiro APB, Dias MP, Clark BL, et al. The BirdLife Seabird Tracking Database: 20 years of

collaboration for marine conservation. Biol Conserv.

2024;299:110813. doi:10.1016/j.biocon.2024.110813

[35] James J, Seekings AH, Skinner P, et al. Rapid and sensitive detection of high pathogenicity

Eurasian clade 2.3. 4.4 b avian influenza viruses in wild birds and poultry.

J Virol Methods. 2022;301:114454. doi: 10.1016/j.jviromet.2022.114454

[36] Nagy A, Černíková L, Kunteová K, et al. A universal RT-qPCR assay for "One Health" detection

of influenza A viruses. PloS one. 2021;16(1):e0244669. doi:10.1371/journal.pone.0244669

[37] Payungporn S, Chutinimitkul S, Chaisingh A, et al. Single step multiplex real-time RT-PCR for

H5N1 influenza A virus detection. J Virol Methods. 2006;131(2):143-7.

doi:10.1016/j.jviromet.2005.08.004

[38] Sutton DA, Allen DP, Fuller CM, et al. Development of an avian avulavirus 1 (AAvV-1) L-gene

real-time RT-PCR assay using minor groove binding probes for application as a routine diagnostic

tool. J Virol Methods. 2019;265:9-14. doi:10.1016/j.jviromet.2018.12.001

[39] Slomka MJ, Reid SM, Byrne AM, et al. Efficient and informative laboratory testing for rapid

confirmation of H5N1 (Clade 2.3. 4.4) high-pathogenicity avian influenza outbreaks in the United

Kingdom. Viruses. 2023;15(6):1344. doi:10.3390/v15061344

[40] Markin A, Wagle S, Grover S, et al. PARNAS: objectively selecting the most representative taxa

on a phylogeny. Syst Biol. 2023;72(5):1052-63. doi:10.1093/sysbio/syad028

[41] Katoh K, Standley DM. MAFFT multiple sequence alignment software version 7: improvements

in performance and usability. Mol Biol Evol. 2013;30(4): 772-780. doi:10.1093/molbev/mst010

[42] Larsson A. AliView: a fast and lightweight alignment viewer and editor for large

datasets. Bioinformatics 30.22. 2014:3276-3278. doi:10.1093/bioinformatics/btu531

[43] Minh BQ, Schmidt HA, Chernomor O, et al. IQ-TREE 2: new models and efficient methods for

phylogenetic inference in the genomic era. Mol Biol Evol 2020;37(5):1530-

1534. doi:10.1093/molbev/msaa015

[44] Sagulenko P, Puller V, Neher RA. TreeTime: Maximum-likelihood phylodynamic analysis.

Virus Evol. 2018;4(1):vex042. doi:10.1093/ve/vex042

[45] Banyard AC, Lean FZ, Robinson C, et al. Detection of highly pathogenic avian influenza virus

H5N1 clade 2.3. 4.4 b in great skuas: a species of conservation concern in Great Britain. Viruses.

2022;14(2):212. doi:10.3390/v14020212

[46] World Organisation for Animal Health (WOAH), 2025. WAHIS Event 5034 Influenza A viruses

of high pathogenicity (Inf. with) (non-poultry including wild birds) (2017-

). https://wahis.woah.org/#/in-event/5034/dashboard

[47] Carneiro AP, Manica A, Clay TA, et al. Consistency in migration strategies and habitat

preferences of brown skuas over two winters, a decade apart. Mar Ecol Prog Ser. 2016;553:267-

81. doi:10.3354/meps11781

[48] Sinclair JC. Subantarctic Skua *Catharacta antarctica* predation techniques on land and at

sea. Mar Ornithol. 1980;8:3-6. doi:10.5038/2074-1235.8.1.27

[49] Bost CA, Cotté C, Bailleul F, et al. The importance of oceanographic fronts to marine birds and

mammals of the southern oceans. J Mar Syst. 2009;78(3):363-

376. doi:10.1016/j.jmarsys.2008.11.022

[50] Dias MP, Oppel S, Bond AL, et al. Using globally threatened pelagic birds to identify priority

sites for marine conservation in the South Atlantic Ocean. Biol Conserv. 2017;211:76-

84. doi:10.1016/j.biocon.2017.05.009

[51] Schoombie S, Dilley BJ, Davies D, et al. The foraging range of Great

Shearwaters (*Ardenna gravis*) breeding on Gough Island. Polar Biol. 2018;41(12):2451-

8. doi:10.1007/s00300-018-2381-7

[52] Jones CW, Phillips RA, Grecian WJ, et al. Ecological segregation of two superabundant,

morphologically similar, sister seabird taxa breeding in sympatry. Mar Biol.

2020;167(4):45. https://doi.org/10.1007/s00227-020-3645-7

[53] Reisinger RR, Bester MN. Long distance breeding dispersal of a southern elephant seal. Polar

Biol. 2010;33(9):1289-91. doi:10.1007/s00300-010-0830-z

[54] Hindell MA, McMahon CR, Bester MN, et al. Circumpolar habitat use in the southern elephant

seal: implications for foraging success and population trajectories. Ecosphere.

2016;7(5):e01213. doi:10.1002/ecs2.1213

[55] Bester MN, Wege M, De Bryn, et al. Ranging and fiving behaviour of Subantarctic fur seals from

the Tristan da Cunha Islands, South Atlantic. Tristan da Cunha Islands Seal Research Report (2009 –

2019). 2019

[56] Ryan PG, Dilley BJ, Risi MM, et al. Three new seabird species recorded at Tristan da Cunha

archipelago. Seabird. 2019;32:122-5. doi:10.61350/sbj.32.122

[57] Bester MN, Somers MJ. Monitoring of vagrant seals on mid-oceanic islands of the South Atlantic

Ocean. Polar Biol. 2025;48(3):79. doi:10.1007/s00300-025-03397-3

[58] Hutchings L, Van der Lingen CD, Shannon LJ, et al. The Benguela Current: An ecosystem of

four components. Prog in Oceanogr. 2009;83(1-4):15-32. doi:10.1016/j.pocean.2009.07.046

[59] Reisinger RR, Raymond B, Hindell MA, et al. Habitat modelling of tracking data from multiple

marine predators identifies important areas in the Southern Indian Ocean. Divers Distrib.

2018;24(4):535-50. doi:10.1111/ddi.12702

[60] Abolnik C. Avian influenza situation report—Africa. Canadian Journal of Microbiology.

2025;26;71:1-4. https://doi.org/10.1139/cjm-2025-0199

[61] Molini U, Yabe J, Meki IK, et al. Highly pathogenic avian influenza H5N1 virus outbreak among

Cape cormorants (Phalacrocorax capensis) in Namibia, 2022. Emerg Microbes Infection.

2023;12(1):2167610. https://doi.org/10.1080/22221751.2023.2167610

[62] Sacristán C, Ewbank AC, Ibáñez Porras P, et al. Novel epidemiologic features of high

pathogenicity avian influenza virus A H5N1 2.3. 3.4 b panzootic: a review. Transbound Emerg Dis.

2024;2024(1):5322378. doi:10.1155/2024/5322378

**Tables**

Table 1: RT-PCR results, virus isolation and viral whole genome sequencing on samples collected from three Tristan skuas (*Stercorarius antarcticus*

*hamiltoni*) on Gough Island on the 20th September 2024.

| Bird number | Sample Type (Swab) | M Gene RT-PCR | H5HP RT-PCR | N1 RT-PCR | Interpretation | Strain | GISAID Accession number | Isolate |
|---|---|---|---|---|---|---|---|---|
| 1 | Tracheal | Positive | Positive | Positive | HPAIV H5N1 | A/Tristan_Skua/Gough_Island/047354/2024_\|H5N1\|_2024 -09-20 | EPI_ISL_201 51596 | Yes |
| | Cloacal | Negative | Negative | Negative | | | | |
| | Brain | Positive | Positive | Positive | | | | |
| 2 | Tracheal | Positive | Positive | Positive | HPAIV H5N1 | A/Tristan_Skua/Gough_Island/047355/2024_\|H5N1\|_2024 -09-20 | EPI_ISL_201 51597 | NA |
| | Cloacal | Negative | Negative | Negative | | | | |
| | Brain | Positive | Positive | Positive | | | | |
| 3 | Tracheal | Positive | Positive | Positive | HPAIV H5N1 | A/Tristan_Skua/Gough_Island/ 050656/2024_\|H5N1\|_2024-09-20_EP1 | NA | Yes |
| | Cloacal | Negative | Negative | Negative | | | | |
| | Brain | Positive | Positive | Positive | | | | |

572　**Figures**

573　Figure 1

585    Figure 2a

603    Figure 2b

610    Figure 3

612    Figure 4 a-c

615    Supplementary Figure S1:

**Figure Captions**

**Figure 1.** a) Map indicating the location for Gough Island in the context of locations with ongoing HPAIV H5N1 outbreaks reported to WAHIS (red dots), b) Gough Island with the research station located in the southeast of the island (black square) and the island helipad's (yellow diamond), c) view of research station and helipad, d) Tristan skua (*Stercorarius antarcticus hamiltoni*).

**Figure 2.** a) Maximum-likelihood tree of HA-gene including the two Gough Island sequences identified from this study (indicated by an arrow) and 944 HA-gene sequences from Antarctica, South and North America previously identified as 2.3.4.4b. Sequences from South America are coloured according to country, all sequences from North America in black, b) a source-sink heat map of transmission rate estimates between geographic regions.

**Figure 3.** Time series showing the distance from the colony of GLS-tracked Tristan skuas (8) during the 2017/18 and 2018/19 breeding seasons, across key stages of the breeding cycle.

**Figure 4.** Kernel utilisation distributions (KUDs) showing the combined home-range (95% UD; lighter shading) and core (50% UD; darker shading) foraging areas of tracked Tristan skuas during (a) the pre-laying exodus (green), breeding (purple), and non-breeding periods (orange) across the 2017/18 and 2018/19 seasons, overlaid on major oceanic frontal features. Panels (b) and (c) illustrate individual migration routes from the Gough Island region to non-breeding foraging grounds on the South African and Namibian continental shelves, and the corresponding return migrations, respectively. The solid bold red line denotes the movement of one individual (GLS tag BH142) that tested positive for highly pathogenic avian influenza virus (HPAIV) in 2024.

651     **Figure S1**: Maximum-likelihood phylogenies of all eight influenza A virus gene segments.

652     Phylogenetic trees were inferred using maximum-likelihood methods for each gene segment show

653     that the Gough Island viruses are closely related to Antarctic H5 viruses rather than South African H5

654     viruses. Tip shapes represent individual sequences and are coloured according to the continent of

655     origin.

677