1

Test-retest reliability of sensorimotor activity measured with spinal cord fMRI

Olivia S. Kowalczyk[1,2]*, Sonia Medina[3], Alessandra Venezia[1], Dimitra Tsivaka[4], Aminul I.
Ahmed[5,6], Steven C. R. Williams[1], Jonathan C. W. Brooks[7], David J. Lythgoe[1], Matthew A.
Howard[1]

[1]Department of Neuroimaging, Institute of Psychology, Psychiatry & Neuroscience, King's
College London, London, UK

[2]Functional Imaging Laboratory, Department of Imaging Neuroscience, UCL Queen
Square Institute of Neurology, University College London, London, UK

[3]Department of Clinical and Biomedical Sciences, University of Exeter, UK

[4]Medical Physics Department, Medical School, University of Thessaly, Greece

[5]Department of Neurosurgery, King's College Hospital NHS Foundation Trust, UK

[6]Wolfson Sensory, Pain and Regeneration Centre, Institute of Psychiatry, Psychology &
Neuroscience, King's College London, UK

[7]Department of Psychology, University of Liverpool, UK

*Corresponding author:

Dr Olivia S. Kowalczyk
Centre for Neuroimaging Sciences
Institute of Psychiatry, Psychology & Neuroscience
De Crespigny Park
London, SE5 8AF
olivia.kowalczyk@kcl.ac.uk

Abstract

Establishing the reliability of spinal cord functional magnetic resonance imaging (fMRI) is critical before employing it to assess experimental or clinical interventions. Previous studies have mapped human motor activity primarily to the ipsilateral ventral horn, aligning with myotomal and dermatomal projections. Despite these insights, the test-retest reliability of spinal fMRI remains under-investigated. Here we assessed spinal cord activation during a sensorimotor paradigm involving right-hand grasping and grip force estimation in 30 healthy volunteers. Participants completed two identical scanning visits, each time performing the same task twice, enabling the investigation of test-retest reliability both within a single experimental visit and between visits performed on different days. Aggregating all task runs, motor-evoked activation was observed in ipsilateral ventro-dorsal regions of spinal segmental levels C6-C8 and T2, as well as in medial regions of levels C1-C3. Despite highly reliable task performance (grip force) and fMRI signal quality (temporal signal-to-noise ratio), the reliability of motor activation was predominantly poor both within and between visits, with notable variability in spatial distribution observed across task runs. Increasing the number of task runs per individual improved the robustness of group-level activation, as indexed by higher activated voxel count, larger cluster spatial extent, and attenuated t-statistic distribution. Although we demonstrated that motor-evoked activation corresponds to the known neuroanatomical organisation of motor circuits, its low test-retest reliability presents a challenge for wider applications of spinal fMRI. Understanding the drivers of low reliability in functional imaging is warranted, but we suggest that looking beyond measurement error is required, including careful consideration of inherent within-individual variability underpinned by neurophysiological and psychological factors.

Keywords: Spinal Cord, Spinal fMRI, Motor function, Test-retest Reliability

1 Introduction

The spinal cord is an integral part of the motor system, playing a key role in movement execution and control. While the function of supraspinal elements of the motor system has been extensively characterised, less focus has been devoted to the study of the spinal cord contributions to human motor function (Lemon, 2008; Nielsen, 2016; Rowe & Siebner, 2012). Recent technological and analytical developments have optimised image acquisition (Barry et al., 2021; Chu et al., 2023; Islam et al., 2019; Tsivaka et al., 2023) and processing (Banerjee et al., 2025; Brooks et al., 2008; De Leener et al., 2017), allowing non-invasive study of the spinal cord using functional magnetic resonance imaging (fMRI). Since the early applications of spinal fMRI (Yoshizawa et al., 1996), a growing literature has assessed spinal cord motor function and somatosensation, as well as the presence of spatiotemporally organised intrinsic activity of the cord, i.e. resting states (for reviews see Kaptan et al., 2024; Kinany et al., 2022; Landelle et al., 2021).

In the study of motor function, the field has employed several paradigms, primarily focussed on the upper limbs, including hand grasping, finger tapping, and extension-adduction-abduction movements of isolated parts of the limb (Landelle et al., 2021). Across paradigms, unilateral motor activity in healthy adults has been shown to elicit activation predominantly in ipsilateral ventral (motor neuron-rich) regions of spinal segmental levels corresponding to the myotomal projections of the moving body part (Barry et al., 2021; Braaß et al., 2023; Hemmerling et al., 2023; Landelle et al., 2021), aligning with the known functional organisation of the cord (Lemon, 2008; Nielsen, 2016). However, motor-related activity has also been observed in contralateral and dorsal (somatosensory neuron-rich) regions of the same spinal segmental level, sometimes with rostrocaudal distribution spanning neighbouring or more distal segmental levels (Hemmerling et al., 2023; Landelle et al., 2021). Additionally, reports indicate that the magnitude of spinal cord activation is modulated by the force generated (Braaß et al., 2023; Madi et al., 2001; Oliva et al., 2025), movement complexity (Maieron et al., 2007; Oliva et al., 2025, 2025; Vahdat et al., 2015), and motor learning (Khatibi et al., 2022; Kinany et al., 2023; Vahdat et al., 2015). Together, this literature demonstrates that spinal fMRI is sensitive to detecting not only gross motor-related activity but also more intricate processes facilitating motor function, which may relate to early sensorimotor integration, proprioception, and motoneuron facilitation-inhibition (Landelle et al., 2021).

The findings from studies in healthy individuals suggest that spinal fMRI might be a promising clinical tool, potentially facilitating identification of disease biomarkers or predictors of treatment progression in conditions affecting sensorimotor pathways, including multiple sclerosis, spinal cord injury, and myelopathy, among others (Kinany et al., 2022; Landelle et al., 2021; Powers et al., 2018). To facilitate this clinical translation, spinal fMRI still needs to demonstrate reliable and reproducible outcomes. This is particularly important given that acquiring fMRI recordings from the spinal cord is

uniquely complicated by the cord's anatomy and location in the body. The small diameter and large rostrocaudal extent of the cord, its proximity to tissues of differing magnetic susceptibility and sources of physiological noise limits the quality of spinal fMRI data and can directly impact the reliability of techniques used to assess function (Kinany et al., 2022). Despite these important considerations, the test-retest reliability of the spinal fMRI remains under-investigated.

To date, test-retest reliability of spinal fMRI has been primarily assessed in resting-state paradigms (Barry et al., 2016; Hu et al., 2018; Kaptan et al., 2022; Kowalczyk et al., 2024; Liu et al., 2016), where reliability estimates were shown to be mixed but comparable to those of brain fMRI (Noble et al., 2019). Only two studies investigated test-retest reliability of motor-related spinal cord activation (Bouwman et al., 2008; Weber et al., 2016b), showing *good* reliability (intraclass correlation coefficient, ICC = 0.8) of average percentage signal change in neuroanatomically relevant section of the cord (Bouwman et al., 2008) but *poor* reliability (ICC range = 0.1-0.3) when assessed voxel-by-voxel (Weber et al., 2016b). Although these two studies offer early evidence of test-retest reliability of motor signalling recorded by spinal fMRI, reliability was only assessed within the same experimental session and in small samples (N = 3 and N = 11). Establishing the reliability of spinal fMRI measurements across distinct sessions and in well-powered samples is critical before employing it to assess experimental or clinical interventions.

The aims of this study were two-fold: (1) we investigated cervical spinal activation during a unilateral motor paradigm involving right hand grasping (squeezing) with grip force measurement in a well-powered sample of healthy adults (N = 30, tested four times), and (2) characterised test-retest reliability of the acquired signal both within the same testing visit and across visits conducted on separate days. To do so, we used our locally-developed blood oxygen level dependent (BOLD) sensitive echo planar imaging (EPI) sequence, optimised for cervical spinal imaging and previously shown to provide increases in signal-to-noise ratio (tSNR) and a marked reduction in signal dropout (Kowalczyk et al., 2024; Tsivaka et al., 2023).

Our preregistered hypotheses were (Kowalczyk et al., 2021):

(1) Task-related BOLD responses to the task will be observed in the ipsilateral dorsal and ventral spinal cord in the C6-C8 spinal segmental levels.

(2) Spinal cord responses will be reliable, with ICC test-retest reliability statistics greater than 0.4.

2 Methods

2.1 Participants

Data from 30 healthy right-handed (assessed using the Edinburgh Handedness Inventory; Oldfield, 1971) adult volunteers (17 females, 13 males; 19-38 years old, mean ± SD = 24 ± 5 years) survived data quality assurance and were retained for analysis. Two runs of the task were acquired on each of two identical study visits.

Full inclusion and exclusion criteria for this study are outlined in the study preregistration (Kowalczyk et al., 2021). Briefly, participants were excluded due to: (1) history of psychiatric, medical, or psychological conditions, (2) history of substance or alcohol abuse, (3) regular use of medications affecting the central nervous system (CNS), (4) irregular menstrual cycle for menstruating participants, (5) MRI-related contraindications. Additionally, participants were excluded if they were unwilling to adhere to the following lifestyle guidelines before each visit: (1) abstain from alcohol for 24 hours, (2) limit consumption of caffeinated drinks to one on each study day, (3) abstain from using non-steroidal anti-inflammatory drugs (NSAIDs) or paracetamol for 12 hours, (4) abstain from nicotine-containing products for 4 hours. Additional exclusion criteria based on data quality were used. Subjects with incomplete functional acquisition, poor image contrast, and geometric distortions were excluded. Full details of participant/data exclusions at each stage of the study are shown in Figure 1.

Written informed consent was obtained from all participants for being included in the study. This study was approved by the Psychiatry, Nursing, and Midwifery Research Ethics subcommittee at King's College London, UK (HR-16/17-4769) and conducted in accordance with the Declaration of Helsinki.

Figure 1. Selection of participants fulfilling the study eligibility criteria and data quality assurance. MR, magnetic resonance; MRI, magnetic resonance imaging.

2.2 Procedure

The study comprised three visits – a screening/familiarisation visit and two identical MRI visits (for test-retest assessment). Across all three study visits, the mean interval (± SD, range) was 21 (± 20, 1-84) days. The mean (± SD, range) interval between the two scanning visits was 27 (± 24, 1-84) days. Variability in between-scan intervals was driven by considerations regarding diurnal (Webb et al., 2024) and menstrual cycle (Schuster & Jansen, 2022) influences on test-retest estimates, as well as practical constraints relating to scanner/participant availability and COVID-19 isolation rules in place at the time of data collection. The wider study involved additional fMRI assessments of resting-state (Kowalczyk et al., 2024) and evoked pressure pain, which preceded the motor task described here. Details of those can be found in the study preregistration (Kowalczyk et al., 2021).

2.2.1 Screening and familiarisation visit

Compliance with study lifestyle guidelines (see Section 2.1) was assessed at the beginning of the visit. Participants underwent breath alcohol and urine drugs of abuse tests to check alcohol/substance use. Caffeine, nicotine, and NSAIDs/paracetamol intake were assessed by self-report. Participants were familiarised with the scanner environment by visiting a mock scanner.

2.2.2 MRI visits

Both MRI visits were performed at the same time of day and the same phase of the menstrual cycle for menstruating participants. The visits began with an assessment of compliance with the study lifestyle guidelines. Additionally, participants completed the state version of the State Trait Anxiety Inventory (STAI; Spielberger et al., 1971) to assess differences in anxiety levels between visits. No differences were observed ($t(29) = 1.80$, $p = 0.082$, Cohen's d = 0.33, 95% CI [-0.69; 0.04]; visit 1 mean ± SD = 28.43 ± 6.35; visit 2 mean ± SD = 30.30 ± 7.92). Subsequently, MRI data was collected in the following order: (1) optimisation of static 0^{th}, 1^{st}, and 2^{nd} order shims and linear slice-specific shims, (2) structural data acquisition, (3) 10 min 50 s resting-state scan (reported in Kowalczyk et al., 2024), (4) evoked pressure pain task (Kowalczyk et al., 2021), and (5) two identical runs of a hand grasping motor task. Respiratory and cardiac traces were recorded with respiratory bellows and a pulse oximeter respectively, along with scanner triggers (at the start of each repetition time; TR), throughout the scan.

The hand grasping motor task required participants to squeeze a rubberised air-filled ball with their right (dominant) hand in response to verbal cues displayed on a projector screen. The screen was positioned at the rear of the scanner bore and visible through a mirror mounted on the head coil. The task was an A/B (squeeze/rest) design comprising eight squeeze/rest block pairs. Each block contained either eight 'squeeze and release' or eight 'rest' instructions (trial duration = 2.5 s; block duration = 20 s). Participants completed two consecutive 5 min 20 s runs of the task per visit, resulting in a total of four task runs over the course of the study. Each run was preceded by individual hand-grip force thresholding, asking the participant to squeeze the ball as hard as they could three times. Grip force exerted was recorded throughout the task.

2.3 MRI acquisition

Data were acquired at the NIHR Wellcome King's Clinical Research Facility, King's College London using a 3T GE MR750 System (General Electric, Chicago, Illinois) equipped with a 12-channel head-neck-spine coil and a 4-channel neurovascular array. A sagittal 3D CUBE T2-weighted structural image was acquired at the beginning of the scanning session over 64 slices with a coverage of the whole brain and cervical spinal cord to vertebral level T1 (TR = 2.5 s, echo time (TE) = 120 ms, echo train length = 78, flip angle = 90°, field of view (FOV) = 300 mm, acquisition matrix = 320 × 320, slice

thickness = 0.8 mm). This acquisition was based on Cohen-Adad et al. (2021) with the FOV increased to 300 mm.

Static 0^{th}, 1^{st}, and 2^{nd} order shims were optimised prior to functional data acquisition. A spectral-spatial excitation pulse was used to excite only tissue water and slice-specific linear shims were implemented by adding 0.6 ms duration x, y, and z-gradient lobes after the excitation pulse. Second-order shimming and x, y, and z-shimming were optimised over elliptical regions of interest (ROIs) covering the cord drawn manually by the researcher present during scanning (OSK or SM). To maintain consistency and avoid potential systematic differences in ROI drawing affecting test-retest estimates, the same researcher drew ROIs for both MRI visits within participants. Full details of shimming optimisation for this acquisition sequence can be found in Tsivaka et al. (2023).

Functional data were acquired over 38 sequential slices in descending order (slice thickness = 4 mm, slice gap = 1 mm), with the inferior-most slices anchored at vertebral level T1 and covering the whole cervical spinal cord and the brainstem (TR = 2.5 s, TE = 30 ms, flip angle = 90°, ASSET factor = 2, FOV = 180 mm, acquisition matrix = 96 × 96, reconstruction matrix = 128 × 128, in-plane voxel size = 1.41 × 1.41 mm). Four dummy scans were acquired to enable the signal to reach steady-state, followed by 136 volumes.

For 13 participants, the manufacturer's EPI internal reference option was used. The internal reference acquires four non-phase-encoded echoes before the EPI echo train, which are used to apply a phase correction to the EPI data. Upon further inspection of the data this was shown to contribute to slice misalignment in the anterior-posterior direction in 10 of the 13 participants (20 out of 26 task runs) and thus the setting was disabled for subsequently recruited participants. To keep the two MRI visits identical, however, the internal reference was used on both MRI visits for the first 13 participants, even after the issue was discovered.

2.4 Data analysis

2.4.1 Behavioural data

Data were analysed using SPSS v29.0.2.0. The outcome measure was maximum hand-grip force sampled during each trial, expressed as a percentage of individual grip strength threshold. Average task performance was assessed across all squeeze blocks. A 2 × 2 repeated measures ANOVA with visit and run as factors compared grip force exerted.

2.4.2 Imaging data

2.4.2.1 Preprocessing

Data were processed using Spinal Cord Toolbox (SCT) version 6.5 (De Leener et al., 2017), AFNI's *3dWarpDrive* (Cox, 1996; Cox & Hyde, 1997), and FSL version 6.0.7.14 (Jenkinson et al., 2012; Smith et al., 2004). Visual quality assurance was performed on raw data and at each stage of processing. Twenty task runs acquired with an early version of the

functional sequence using the internal reference (see above) had several slices come out of alignment with the rostrocaudal axis of the spinal cord due to a shift in the anterior-posterior (EPI phase-encoding) direction. A custom in-house Matlab version 9.5.0 (Mathworks Inc.) script was used to move the slices back into alignment with the rest of the cord. Briefly, for each slice, a 1D projection along the anterior/posterior direction was calculated for each time-point by summing the voxels in the left/right direction across the spinal cord. The anterior/posterior shift was determined by calculating the maximum of the cross correlation of the projection at each time-point with the first time-point. The shift was then applied to the image data in a block circular manner. Only shifts by an integer number of voxels were applied to avoid the need for an extra interpolation step. This step was performed prior to any other preprocessing.

For all functional data, brainstem structures were separated from cervical volumes at the level of the odontoid process. Subsequently, spinal cord functional data were motion-corrected for x and y translations only, using an in-house implementation of AFNI's *3dWarpDrive* following the steps in the Neptune Toolbox (https://neptunetoolbox.com/). Motion-corrected data were smoothed with an in-plane 2D Gaussian kernel with full width at half maximum (FWHM) of 2 mm using a custom in-house script relying on tools from AFNI and FSL.

Warping parameters for spatial normalisation were determined by segmenting and registering the functional data to the PAM50 spinal cord template (De Leener et al., 2018), via an intermediary subject-specific T2-weighted 3D volume. Specifically, *sct_deepseg* (Gros et al., 2019), with EPI-specific segmentation algorithm (Banerjee et al., 2025), was used to segment the cord from the cerebrospinal fluid (CSF) on motion-corrected functional data and on the T2-weighted structural image. Manual adjustment was needed for accurate functional data segmentation, performed in FSLeyes (McCarthy, 2022). Warping parameters for registration of functional data to the PAM50 template were created by combining warp parameters from: (1) registering the structural T2-weighted image to functional data, using manually created disc labels on both images and (2) registering the segmented cord from the T2-weighted image to the PAM50 T2-weighted template via *sct_register_to_template* (De Leener et al., 2018). These warps were applied to contrast of parameter estimates (COPE) images obtained from subject-level modelling, via *sct_register_multimodal* (De Leener et al., 2018). Inverse warp parameters obtained from these steps were used to transform the PAM50 template CSF mask to each participant's functional space. Each mask was subsequently used in physiological denoising.

The Physiological Noise Modelling (PNM) toolbox (Brooks et al., 2008) was used to generate 33 slice-specific regressors accounting for physiological noise based on cardiac and respiratory traces, and CSF signal.

2.4.2.2 Subject-level modelling

Spinal cord activation was assessed using mass univariate general linear models (GLMs) in FSL FEAT, using FILM pre-whitening and highpass temporal filtering at 90 s. GLMs included explanatory regressors for squeeze trials, along with slice-wise nuisance regressors accounting for physiological noise (33 PNM regressors) and in-scan motion (x and y translations). Additionally, volume censoring regressors were added to exclude the first trial of each squeeze block due to large behavioural variability in the responses at the start of the squeeze blocks (see Supplementary Figure 1). Two contrasts identified: (1) motor activity greater than rest (COPE multiplier = 1), and (2) for additional quality assurance and inspection of potential task negative activity – rest greater than motor activity (COPE multiplier = -1).

We also conducted a parametric modulation analysis to account for variability in task performance. These GLMs included a parametric regressor representing the grip force exerted during each squeeze trial, in addition to the regressor modelling squeeze trials and slice-wise nuisance regressors described above. Three contrasts were estimated: (1) motor activity greater than rest, accounting for grip force variability (hereafter referred to as grip force-adjusted motor activity), (2) activity positively correlated with grip force, and (3) activity negatively correlated with grip force.

2.4.2.3 Group-level modelling

Task effects were assessed using a one-sample t-test comprising averages of subject-level COPEs across all four runs of the task (two runs from each visit). Additional group-level maps were computed to assess activation in each run and visit separately. These analyses were conducted for both standard and parametric modulation models.

Group-level analyses were conducted within a priori defined ROIs and across the whole cervical cord. ROIs included C5-T1 spinal segmental levels, derived from the PAM50 spinal levels atlas and corresponding to the dermatomal and myotomal projection of the hand (Sonoo, 2023; Weber et al., 2020).

2.4.2.4 Impact of number of task runs on group-level activation

To investigate the impact of data quantity on group-level motor activation maps, four group-level models were conducted. The group-level models included: only one run (visit 1, run 1), two runs (mean of runs 1 and 2 from visit 1), three runs (average of runs 1 and 2 from visit 1, and run 1 from visit 2), and all fours runs (average of runs 1 and 2 from visits 1 and 2). These analyses were conducted for standard and parametric modulation models.

2.4.2.5 Statistical inference

All group-level models used non-parametric permutation-based testing implemented in *randomise* (FSL) with threshold-free cluster enhancement (5000 permutations, $p < 0.05$) and family-wise error (FWE) rate correction.

2.4.3 Test-retest reliability

ICC was used to quantify test-retest reliability within and between visits. For within-visit reliability, we compared the mean of the first task run across visits $(\overline{run\ 1}=\frac{run\ 1_{visit\ 1}+run\ 1_{visit\ 2}}{2})$ with the mean of the second task run across visits $(\overline{run\ 2}=\frac{run\ 2_{visit\ 1}+run\ 2_{visit\ 2}}{2})$. For between-visit reliability, we compared the mean of both task runs from visit 1 $(\overline{visit\ 1}=\frac{visit\ 1_{run\ 1}+visit\ 1_{run\ 2}}{2})$ with the mean of both task runs from visit 2 $(\overline{visit\ 2}=\frac{visit\ 2_{run\ 1}+visit\ 2_{run\ 2}}{2})$.

Following previous recommendations, ICC values were categorised accordingly: <0.4 as *poor*, 0.4–0.59 as *fair*, 0.6–0.74 as *good*, and >0.75 as *excellent* (Cicchetti, 1994; Fleiss et al., 2013).

2.4.3.1 Behavioural data

Reliability of task performance was assessed using ICC(3, 2) and calculated in SPSS v29.0.2.0.

2.4.3.2 Imaging data

ICC values were calculated for each voxel (i.e. voxelwise) using the locally-developed ICC toolbox (Caceres et al., 2009) running in Matlab version 9.5.0 (Mathworks Inc.). Reliability was calculated for the whole cervical cord (PAM50 cord mask), the activation network, and within spinal segmental levels C5-T1 (PAM50 spinal segmental masks). The activation network was obtained using a one-sample *t*-test of the first measurement $(\overline{run\ 1}$ for within-visit comparison, $\overline{visit\ 1}$ for between-visit comparison) with a voxelwise *t*-statistic threshold of 3.5 (equivalent to p = 0.001) conducted in SPM8, as per the original manuscript describing this approach (Caceres et al., 2009). Median ICC values are reported, defined as the reliability measure obtained from the median of the ICC distributions within regions.

In addition to the pre-registered voxelwise approach, test-retest reliability of the average signal was computed. Mean signal was extracted from motor-related COPEs from first-level models from the whole cervical cord (PAM50), the activation network obtained from group-level *randomise* (p_{FWE} < 0.05) models ($\overline{run\ 1}$ for within-visit comparison, $\overline{visit\ 1}$ for between-visit comparison), and within spinal segmental levels C5-T1 (PAM50). Reliability of motor activation was assessed in both standard and parametric modulation analysis.

To supplement the assessments of reliability of motor activation and to test the reliability of overall fMRI signal, within- and between-visit ICC was calculated for temporal signal-to-noise ratio (tSNR). tSNR was calculated on motion-corrected data prior to any further preprocessing or modelling steps to avoid artificially inflating the measure. The mean motion-corrected functional image was divided by its standard deviation. Mean tSNR was extracted in native space for the whole cervical cord using subject-specific cord masks generated during preprocessing (see Section 2.4.2.1).

Reliability of average signal and tSNR was assessed using ICC(3, 2) in SPSS v29.0.2.0.

3   Results

3.1   Motor performance

Participants produced consistent hand-grip force during the squeeze blocks (mean across all task runs ± SD = 83.50% ± 12.40; visit 1 – run 1 = 84.64% ± 16.89; visit 1 – run 2 = 79.45% ± 13.62; visit 2 – run 1 = 88.43% ± 14.47; visit 2 – run 2 = 81.67% ± 12.88). No grip force was detected during rest blocks (Figure 2).

No significant differences were observed between visits ($F_{(1, 29)} = 3.14$, $p = 0.087$, $\eta^2p = 0.098$), however, lower grip force was observed in the second run of the task across both visits ($F_{(1, 29)} = 8.58$, $p = 0.007$, $\eta^2p = 0.228$). No interaction effects between visit and run were found ($F_{(1, 29)} = 0.23$, $p = 0.631$, $\eta^2p = 0.008$).

Figure 2. Average hand-grip force exerted by participants during each of the task runs on each visit. The values are expressed as percentage of individually thresholded grip force. Boxplots mark the median (thick black line), interquartile range (box), and minimum and maximum values (whiskers). Red dots and the associated labels mark the mean.

3.2   Motor activation

Group averaging across all four task runs (two runs per visit, two visits) showed ipsilateral right-lateralised motor-related activation in both ventral and dorsal regions of segmental levels C6 and C7, as well as bilateral ventro-dorsal activation at levels C1-C3 when assessed across the whole cervical spinal cord (Figure 3A). A similar extent of ipsilateral

ventro-dorsal activation was observed when using a priori defined ROIs as masks at spinal segmental levels C6 and C7 (Figure 3B). Analysis of grip force-adjusted motor activation yielded similar results, with additional right ventral activation at levels C8 and T2 in whole cord analysis, and right ventro-dorsal activation in ROI analysis of level C6.

When assessing group-level activation for each task run separately within the whole cord mask, motor-evoked activation was observed in ipsilateral ventro-dorsal regions spanning levels C6-C8 (runs 1 and 2, visit 2). Additionally, whole cord analyses revealed bilateral ventro-dorsal and dorsal activation clusters at levels C1-C3 (run 2, visit 2) and C2 (run 1, visit 1), respectively (Figure 4). ROI analyses of individual task runs, showed primarily ipsilateral ventro-dorsal activation at levels C6 (runs 1 and 2, visit 2), C7 (runs 1 and 2, visit 2; run 2, visit 1), C8 (runs 1, visit 1 and visit 2), and T1 (run 1, visit 2), with some clusters spanning bilaterally (Figure 5). Grip force-adjusted motor activation was observed in the same regions, with an additional right ventro-dorsal activation at level C8 in whole cord analysis (run 1, visit 1) and C6 in ROI analysis (run 1, visit 2).

No significant clusters were detected in the contrast exploring activation during rest periods, or when exploring the relationship between grip force and cord activation.

Figure 3. Group-level motor activation (yellow) and grip force-adjusted motor activation (red) averaged across all task runs. Statistically significant t-statistics are shown (p_{FWE} < 0.05). Labels denote spinal segmental levels. **A.** Activation obtained from whole cervical cord analysis shown on one representative coronal slice and representative axial slices from spinal segmental levels C1-C3 and C6-T2. **B.** Activation obtained from ROI analysis at spinal segmental levels C5-T1. D = dorsal, I = inferior, L = left, n.s. = non-significant, R = right, ROI = region of interest, S = superior, V = ventral.

Figure 4. Group-level motor activation (yellow) and grip force-adjusted motor activation (red) in each run of the task assessed within the whole cervical cord. One representative coronal slice per task run is presented. Statistically significant t-statistics are shown ($p_{FWE} < 0.05$). Left axis denotes spinal segmental levels. I = inferior, L = left, n.s. = non-significant, R = right, S = superior.

Figure 5. Group-level motor activation (yellow) and grip force-adjusted motor activation (red) in each run of the task assessed within each spinal segmental level ROI (C5-T1). Statistically significant t-statistics are shown ($p_{FWE} < 0.05$). D = dorsal, I = inferior, L = left, n.s. = non-significant, R = right, ROI = region of interest, S = superior, V = ventral.

3.3 Impact of data quantity on motor activation

To investigate the impact of data quantity on motor activation, we repeated group-level models, varying the number of task runs included from one to four. We conducted this comparison on both motor and grip force-adjusted motor activation. In both modelling approaches, increasing the number of task runs included in each analysis resulted in more active voxels ($p_{FWE} < 0.05$) across the cord (motor activation, r = 0.89, p = 0.114, 95% CI [-0.51; 1.00]; grip force-adjusted motor activation, r = 0.93, p = 0.074, 95% CI [-0.99; 0.87]; Figure 6B) and skewed the distribution of statistically significant t-values towards lower estimates (Figure 6C). Increasing the number of runs also led to a broader spatial extent of activated voxels across modelling approaches, including at expected segmental levels C6-C8 (Figure 6A).

Figure 6. Investigation of the impact of increasing the number of task runs included in analysis on group-level motor activation (yellow) and grip force-adjusted motor activation (red) of motor response across the cord (p_{FWE} < 0.05). **A.** Group-level motor activation maps. One representative coronal slice per group-level is presented. Left axis denotes spinal segmental levels. **B.** Number of active voxels across the cord. **C.** Density of t-statistics. I = inferior, L = left, N = number, R = right, S = superior.

3.4 Test-retest reliability

3.4.1 Behavioural data

Reliability of task performance was *good* within visit (ICC = 0.73) and *excellent* between visits (ICC = 0.84).

3.4.2 Imaging data

Voxelwise reliability estimates of motor activation were *poor* for both within- and between-visit comparisons (ICC range = 0-0.34; Figure 7). Within- and between-visit reliability of average regional activation was *poor* in most ROIs (ICC range = 0-0.38), with

only a few exceptions. *Fair* within-visit reliability of average signal was observed in the
motor activation network (ICC = 0.51) and at C8 (ICC = 0.56), as well as between-visit at
segments C5 (ICC = 0.57), T1 (ICC = 0.55), and across the whole cervical cord
(ICC = 0.50). *Excellent* within-visit reliability was observed at C5 (ICC = 0.76).

Accounting for grip force (grip force-adjusted motor activation) lowered ICC estimates by
an average of 0.05 (t(27) = 2.5, *p* = 0.020, Cohen's d = 0.47, 95% CI [0.07; 0.86];
Supplementary Figure 2).

In addition to motor activation, we also assessed the reliability of tSNR as a measure of
overall signal acquired during scanning. Average tSNR across all task runs was 27 ± 6.01
(visit 1 – run 1 = 26.43 ± 6.63; visit 1 – run 2 = 26.62 ± 6.51; visit 2 – run 1 = 27.46 ± 5.16;
visit 2 – run 2 = 27.5 ± 5.86). tSNR showed *excellent* reliability both within- (ICC = 0.96)
and between-visit (ICC = 0.79).

Figure 7. ICC estimates for within- and between-visit reliability of motor activation. **A.**
Forest plot showing median ICC for voxelwise assessments and ICC of mean COPE
across the cervical spinal cord, the motor activation network, and spinal segmental
levels C5-T1. **B.** Statistical maps showing ICC distribution for individual voxels within the
cervical cord for within- and between-visit reliability assessments.
COPE = contrast of parameter estimates, ICC = intraclass correlation coefficient,
SE = standard error.

4   Discussion

This study investigated cervical spinal cord activation during a unilateral hand grasping
motor task and its test-retest reliability. We observed the expected ipsilateral activation
in ventro-dorsal regions spanning spinal segmental levels C6-T1, with additional bilateral

ventro-dorsal and medial activation at segments C1-C3, and ipsilateral ventral activation at segment T2. Despite *good*-to-*excellent* reliability of task performance and overall fMRI signal (tSNR), reliability of motor activation was primarily within the *poor*-to-*fair* range for voxelwise and mean signal estimates. Herein, we discuss these findings and consider the challenges in establishing reliability of measurements, emphasising the pressing need to distinguish between true measurement error and biologically meaningful sources of variability within individuals.

4.1 Hand grasping evoked ipsilateral ventro-dorsal activation in spinal segmental level C6-C8

The first aim of this study was to characterise motor-evoked cervical spinal cord activation. The ventro-dorsal activation observed at spinal segmental levels C6-C8 was largely lateralised to the hand executing the movement, with some clusters extending medially. This ipsilateral predominance aligned with our predictions and the existing literature. Spinal cord anatomy is inherently lateralised, with afferent and efferent projections ipsilateral to the corresponding body part (Ganapathy et al., 2025). Previous spinal fMRI studies using unilateral motor tasks have reported mostly ipsilateral activation, which aligns with established spinal circuitry (Barry et al., 2021; Braaß et al., 2023; Hemmerling et al., 2023; Maieron et al., 2007; Weber et al., 2016b). However, additional bilateral and/or medial activation has also been observed during unilateral movements (Bouwman et al., 2008; Hemmerling et al., 2023; Oliva et al., 2025; Stroman & Ryner, 2001; Vahdat et al., 2015), similar to that observed in the medial parts of levels C6-C8 in our ROI analysis. This medial activation likely reflects neural processing in the intermediate zone, a gray matter region between the ventral and dorsal horns that connects the left and right hemicords and houses commissural interneurons and propriospinal neurons (Ganapathy et al., 2025). Electrophysiology recordings in non-human primates have shown that commissural interneurons are involved in hand grasping, facilitating motor coordination of the upper limb (Soteropoulos et al., 2013; Takei & Seki, 2010). Reports from human studies show that analogous processes in the lumbar cord may underpin motor coordination during lower limb movements (Hanna-Boutros et al., 2014; Mrachacz-Kersting et al., 2018). Nonetheless, it is important to note that methodological choices, such as in-plane voxel size and associated partial volume effects, along with spatial smoothing, limit the spatial specificity of spinal fMRI. Consequently, while medial activation may reflect commissural interneuron activity, finer spatial resolution is required to validate this interpretation.

In line with our hypothesis, motor-evoked activation observed in this study encompassed both ventral and dorsal regions of the cord. The ventral and dorsal gray matter horns primarily comprise motoneurons and sensory neurons, respectively (Ganapathy et al., 2025). Both ventral and dorsal activation are consistently reported in spinal fMRI studies probing motor function, with peaks often located more ventrally (Landelle et al., 2021). Recruitment of both ventral and dorsal regions of the cord is expected, given the nature

of the sensorimotor task used here, the complex circuitry of the cord, and the intricate connections among motoneurons, sensory neurons, and interneurons, which give rise to various reflex loops and facilitation-inhibition interdependencies (Pierrot-Deseilligny & Burke, 2012). Co-activation of ventral and dorsal horns might thus reflect sensory feedback accompanying movement or early sensorimotor integration (Enoka, 2008; Landelle et al., 2021; Yeganegi et al., 2018). Despite a somewhat larger voxel size than typically used in the field (1.41 × 1.41 mm versus 1 × 1 mm) and the use of spatial smoothing lowering the spatial specificity of our findings, studies using higher spatial resolution (Hemmerling et al., 2023; Oliva et al., 2025; Weber et al., 2016b) and not incorporating spatial smoothing (Hemmerling et al., 2023) reported similar activation patterns, supporting the interpretation that hand grasping recruits diffuse areas within spinal cord segments.

In agreement with established muscle innervation pathways (Schirmer et al., 2011), hand grasping led to activation at spinal segmental levels spanning C6-T1, with peaks at levels C6 and C7, consistent with our hypothesis. Hand grasping involves muscle groups innervated by median and ulnar nerves of the brachial plexus, which receive projections from C5-C8 and C8-T1 nerve roots, respectively (Dawson-Amoah & Varacallo, 2025; Schirmer et al., 2011). Previous spinal fMRI studies involving similar tasks have shown analogous activation spanning C5-T1 (Backes et al., 2001; Barry et al., 2021; Braaß et al., 2023; Giulietti et al., 2008; Hemmerling et al., 2023; Islam et al., 2019; Kinany et al., 2019; Ng et al., 2006; Oliva et al., 2025; Stroman et al., 1999; Stroman & Ryner, 2001; Weber et al., 2016b), with peaks for wrist movements occurring at C6-C7 and finger movements at C6-T1 (Landelle et al., 2021). The activation we observed at levels C6-C8 when combining all task runs, aligns with the existing literature and the known somatotopic organisation of spinal cord function.

4.2 Hand grasping involved additional activation in rostral cervical and thoracic cord segments

In addition to the expected recruitment of spinal levels C6-T1, we observed activation at spinal segmental levels C1-C3 and T2. Although previous studies have documented activation spreading to adjacent segments, few have reported distinct clusters in such rostral regions (Landelle et al., 2021). Engagement of C2-C4 spinal segmental levels has been observed in addition to the expected dermatomal/myotomal projections during finger tapping (Govers et al., 2007; Ng et al., 2008) and C3-C4 during sensory stimulation of the fingers (Stracke et al., 2005). Drawing from electrophysiology literature, activation of rostral segments of the cervical spine may be underpinned by the involvement of neuronal projections spanning multiple spinal segmental levels, which are involved in somatosensory integration, proprioception, and fine motor control (Flynn et al., 2011; Pierrot-Deseilligny & Burke, 2012). This interpretation is further supported by the location of the peaks of our C1-C3 activation mapping to the intermediate zone, which contains interneurons and propriospinal neurons (Ganapathy et al., 2025). Additionally, C2-C4

segments receive projections from the posterior head, as well as the neck and shoulders (Schirmer et al., 2011; Walji & Tsui, 2016). Similarly, T2 segment corresponds to myotomal and dermatomal projections to the upper chest and back (Harrow-Mortelliti et al., 2025). We speculate that the activation observed at these levels may reflect additional sensory feedback and/or contraction of muscles not directly involved in hand grasping, but that aid in maintaining appropriate posture during scanning.

Relating this finding to the spinal fMRI literature is complicated by the fact that the studies reporting rostral activation have been conducted with modest sample sizes (N = 9-12) and prior to the many developments in optimising spinal fMRI data acquisition and processing that have occurred in the last decade (Kinany et al., 2022; Powers et al., 2018). The consistency of these findings is difficult to establish given that the majority of current spinal fMRI literature focuses on spinal segmental levels C4-T1, corresponding to the afferents and efferents of the upper limb, either by using a smaller field of view (Barry et al., 2021; Hemmerling et al., 2023; Oliva et al., 2025; Weber et al., 2016b) or by restricting analyses to this area (Kinany et al., 2019; Vahdat et al., 2015). Despite these considerations, and given the specificity of the activation to motor periods of the task, the C1-C3 and T2 activation observed here likely reflects neurophysiological processes related to movement.

4.3 Poor test-retest reliability of motor-activation can stem from methodological choices

Our second objective was to assess the test-retest reliability of spinal motor-evoked activation. Despite highly reliable task performance and tSNR (ICC > 0.7), motor activation reliability was substantially lower than expected (average ICC = 0.3), falling primarily within the *poor*-to-*fair* range. Similarly, the spatial representation of activation maps, particularly their rostrocaudal extent, varied across task runs in both whole cord and ROI analyses. Furthermore, despite reliability estimates typically being higher for assessments conducted within the same visit, as compared to those conducted during different visits, this pattern was not consistent in our data, except for tSNR (Bennett & Miller, 2010, 2013).

Our reliability findings align with previous spinal fMRI studies reporting *poor* test-retest reliability in motor (Weber et al., 2016b), sensory (Dabbagh et al., 2024; Weber et al., 2016a), and resting-state paradigms (Kowalczyk et al., 2024). These findings parallel the broader neuroimaging literature, where brain fMRI studies similarly report modest reliability (average ICC = 0.5) (Bennett & Miller, 2010; Elliott et al., 2020; Kragel et al., 2021; Noble et al., 2017, 2019) and tSNR is notably higher (typically three times that found in the cord; Murphy et al., 2007). Measurement error is a large contributor to low reliability, and considering that spinal fMRI faces additional technical challenges leading to greater signal variability (Kinany et al., 2022; Powers et al., 2018), perhaps low ICC values should be expected. The small diameter of the cord, impact of physiological noise, close proximity of different tissue types, and additional in-scan motion related to CSF

pulsation can all reduce the signal-to-noise ratio of fMRI data obtained from the spinal
cord (Kinany et al., 2022; Landelle et al., 2021; Powers et al., 2018). Indeed, the impact
of noise on spinal fMRI reliability has been extensively characterised recently,
underscoring the importance of accounting for the various sources of noise present in
spinal fMRI recordings (Kaptan et al., 2022).

Nonetheless, low reliability does not equate to low validity. Past work has shown little
association between reliability and the ability to predict behaviour or reidentify
individuals from intrinsic brain activity, demonstrating that data with low reliability still
contains biologically meaningful information (Noble et al., 2017). Furthermore,
preprocessing approaches developed to account for artefactual signal and thus boost
validity, including in-scan motion or unrelated physiological processes, have been shown
to reduce test-retest estimates in both brain and spinal fMRI (Birn et al., 2014; Kaptan et
al., 2022; Lipp et al., 2014; Noble et al., 2017, 2019). This likely reflects more structured
properties of noise (e.g. cardiac cycle, respiration, CSF pulsation) compared to the more
dynamic nature of neural processing (Kinany et al., 2020). Consequently, while continued
efforts to improve image acquisition, preprocessing, and modelling approaches will
undoubtedly increase both the reliability and validity of spinal fMRI (Barry et al., 2016;
Kaptan et al., 2022), it is also important to consider that observations of low reliability
may be underpinned by biologically meaningful variability in the measured signal.

In simple terms, ICC is the ratio of between-participant to within-participant variance
(Fleiss et al., 2013). Consequently, a high ICC is obtained when the variance between
participants is large, while variance within participants is minimal. Traditionally, however,
fMRI tasks are designed to minimise between-participant variance and elicit a robust
group-level response (Fröhner et al., 2019; Hedge et al., 2018). Even highly repeatable
group-level activation maps are not always underpinned by reliable individual-level
activation maps (Fröhner et al., 2019). While the lower level of measurement error at
group level, compared to individual level, might play a part in this phenomenon,
simulation and longitudinal analyses of brain fMRI suggest an important role of trait- and
state-related individual differences (Fröhner et al., 2019; Hedge et al., 2018). The brain
and spinal cord form a dynamic system, constantly adapting to external and internal
demands (Finn & Rosenberg, 2021; Kaptan et al., 2024; Kinany et al., 2020; Uddin, 2020).
Low ICC values and variable activation patterns across task runs might be explained by
fluctuating psychological and neurophysiological states within individuals. To date,
validated mixed-effects approaches, which consider within-, as well as between-
individual variability, remain reliant on random field theory assumptions for statistical
inference (Worsley, 2003). Important assumptions of stationary smoothness across the
image volume are violated by the inherently anisometric shape of the cord, rendering
their application invalid for spinal fMRI. Nonetheless, biologically-meaningful variability
influencing our data is in line with *good*-to-*excellent* reliability of tSNR observed here,

indicating that while the spatial extent and magnitude of responses related to hand
grasping was variable, the overall signal remained stable over time.

4.4 Variability in motor activation has likely behavioural and neurophysiological bases

4.4.1 Behavioural variability

It is possible that the variability of activation patterns and magnitude reflects differences
in performance across task runs, such as the exerted force or recruited muscle groups.
Motor-related intensity coding has previously been observed in the spinal cord using
fMRI, showing that activation scales positively with the force exerted (Braaß et al., 2023;
Madi et al., 2001; Oliva et al., 2025). While all participants in our study achieved adequate
performance in all task runs (approximately 80% of individualised grip force), the grip
force they exerted in the second run of the task across both visits was approximately 6%
lower than that exerted in the first runs. This suggests a potential effect of fatigue on task
performance, inherent to conducting multiple identical assessments within the same
testing session and potentially leading to the lower test-retest reliability of performance
within- compared to between-visits. Nonetheless, the similar pattern and magnitude of
motor activation across task runs observed in the analysis accounting for grip force (grip
force-adjusted motor activation), suggests that trial-by-trial variability in evoked grip
force alone is an unlikely explanation for the differences in activation patterns across task
runs.

Given that hand grasping involves contractions of multiple muscle groups, including
those of the hand and forearm (Dawson-Amoah & Varacallo, 2025; Schirmer et al., 2011),
the variability in activation may represent differences in motor control strategies
employed by participants across task runs. Combining spinal fMRI with
electromyography has demonstrated distinct segmental localisation of motor-evoked
activity depending on the type of movement performed, such as wrist extension (C5-C7),
wrist adduction (C7-C8), and finger abduction (C7-T1) (Kinany et al., 2019). Since
multiple strategies can be employed to exert adequate grip-force in the task,
incorporating electromyography recordings would have allowed us to track the
recruitment of specific muscle groups during the task and identify any potential changes
in motor control strategies over time. We recommend the inclusion of orthogonal
readouts in imaging studies as an index of behavioural variability, facilitating the capture
of associated neural processing with improved precision (Nebe et al., 2023).

4.4.2 Dynamics of neural processing

The variability in activation maps across task runs may also be explained by the plastic
and dynamic nature of neural processing in the spinal cord. Motor skill learning has been
shown to modulate spinal cord activity (Khatibi et al., 2022; Kinany et al., 2023; Vahdat et
al., 2015), and among other things, to induce a caudal-to-rostral shift in activation over
time (Khatibi et al., 2022). While previous spinal fMRI literature has focussed on
functional plasticity induced by repetition of complex movements, repetition of even

simple movements, such as fist clenching, is known to lower activation in the brain over time (Dinstein et al., 2007; Hamilton & Grafton, 2008). Although not explicitly investigated here, we speculate that the differences in activation patterns observed across task runs may be partly attributed to similar repetition suppression taking place at the level of the cord. These insights may explain the lack of significant activation observed in the second task run of the first visit, as well as the inconsistency in the spatial extent of activation across runs. Furthermore, the caudal-to-rostral shift in significant clusters observed over the course of task runs completed in the second visit echoes that observed following complex motor skill learning, potentially signifying the development of more efficient motor control with practice (Khatibi et al., 2022). Consequently, changing activation patterns observed across task runs may reflect dynamically adapting neurophysiological processes taking place within the spinal cord to exert appropriate motor control. This interpretation challenges the historical view of the spinal cord as a stable information relay and highlights the importance of accounting for the dynamic nature of neural processing in study design.

4.4.3 Individual differences in neuroanatomy

There is known individual variability in spinal cord structure. Spinal cord segmental levels, which describe the rostrocaudal organisation of the spinal cord, are identified by nerve rootlet connection points. By contrast, vertebral levels characterise the rostrocaudal extent of the bony structure of the spine (De Leener et al., 2018; Landelle et al., 2021). Given the ease of vertebral identification and the availability of automated tools, the majority of currently used spatial normalisation pipelines for spinal fMRI rely on vertebral levels (Azad et al., 2021; Bozorgpour et al., 2023; Gros et al., 2019; Rouhier et al., 2020; Ullmann et al., 2014; Vania & Lee, 2021). However, there is a disparity between spinal and vertebral levels, with evidence from postmortem (Lang & Bartram, 1982) and imaging studies (Cadotte et al., 2015) reporting individual differences in nerve rootlet distribution. We suggest that variability in spinal cord architecture is another likely contributor to the inconsistency in the rostrocaudal extent of activation observed in this study and beyond. Recent efforts to automate nerve rootlet segmentation should offer improvements to spatial normalisation strategies (Valošek et al., 2024), however, further validation for ventral nerve rootlets is warranted before applying this strategy to motor spinal fMRI studies.

4.5 Further considerations about methodological choices

We have discussed several methodological and biological factors that can affect group-level activation maps and their test-retest reliability. These factors may act in isolation or interact, their effects likely magnified by the unique challenges associated with acquiring fMRI signals from the cord. The complexities of spinal fMRI have been adroitly and exhaustively reviewed elsewhere to which we refer the reader (Kaptan et al., 2024; Kinany et al., 2022; Landelle et al., 2021; Powers et al., 2018). Here, we consider select

701 methodological challenges affecting spinal fMRI research, including statistical power,
702 choice of fMRI task, and optimisation of imaging paradigms.

4.5.1 Number of observations per individual

An important secondary finding of this study concerns the importance of appropriate
statistical power in spinal fMRI studies (Murphy et al., 2007). While recent years have
seen more than doubling in sample sizes (Braaß et al., 2023; Dabbagh et al., 2024;
Hemmerling et al., 2023; Kaptan et al., 2022; Kowalczyk et al., 2024; Landelle et al., 2021;
Oliva et al., 2022), power depends not only on the number of participants but also on the
amount of observations per participant (Dabbagh et al., 2024; Hemmerling et al., 2023;
Weber et al., 2016b). Recent findings from a similar motor paradigm to that used here
demonstrated that increasing the amount of observations per participant in spinal fMRI
may be more effective at maximising detected activation than simply expanding sample
size (Hemmerling et al., 2023), aligning with early reports from motor (Weber et al.,
2016b) and sensory studies (Weber et al., 2016a). Such an approach also echoes
recommendations reported in brain fMRI, in which a greater number of observations per
individual increases both reliability and validity (Noble et al., 2017). Increasing the
number of observations can improve accuracy by reducing the prevalence of random
unstructured noise originating from transient factors, such as alertness, fatigue, head
and neck positioning, thereby distilling signal reflecting stable biological traits and
processes (Elliott et al., 2021). Our findings support this approach, showing more robust
activation maps when multiple task runs are combined. Furthermore, underpowered
imaging studies are prone to overestimating effect sizes (Cremers et al., 2017), a pattern
recently confirmed in spinal fMRI (Hemmerling et al., 2023). Extending this evidence, we
observed inflated t-statistics when fewer observations per participant were analysed,
highlighting that this bias can arise not only from limited sample sizes but also
insufficient quantity of data per participant. Consequently, we accord with previous
recommendations to acquire more data per individual within adequately powered
sample sizes to improve spinal fMRI precision (Hemmerling et al., 2023).

4.5.2 Optimisation of functional paradigms

A well optimised fMRI paradigm must balance technical efficiency in capturing and
isolating the BOLD signal and *human* factors, such as attention, habituation, fatigue.
Given the lower effect sizes and greater impact of extraneous noise in the cord compared
to the brain, one might approach optimising functional paradigms by maximising signal-
to-noise ratio (e.g. by increasing stimulus repetition and/or scan duration; Landelle et al.,
2021; Powers et al., 2018; Stroman et al., 2014). This focus on technical optimisation,
however, may inadvertently lead to dismissal of large portions of biologically-relevant,
within-individual variability as noise. fMRI task choice remains a critical design
consideration, with particular emphasis on the trade-offs likely in optimising larger
versus more specific cord activation. Here, we selected hand grasping due to the large-
scale engagement of the upper limb and the large effect sizes associated with the task

facilitating assessment of test-retest reliability (Dawson-Amoah & Varacallo, 2025; Landelle et al., 2021; Schirmer et al., 2011). Nonetheless, we reflect that the many strategies employable in executing hand grasping may also contribute to the variability of motor activation observed across task runs, including the spread in rostrocaudal extent. A task choice targeting particular myotomes may offer greater spatial specificity (Kinany et al., 2019) and more consistent cord activation over time.

4.5.3 Outlook

Despite some three decades of development, functional imaging remains in its relative infancy in characterising neurophysiological and psychological sources of within-individual variability. Studies searching for reliable trait-level indices of neural processing have predominated, inadvertently biasing against the largely accepted view of the CNS as a plastic and dynamic system (Finn & Rosenberg, 2021; Kaptan et al., 2024; Kinany et al., 2020; Ricchi et al., 2025; Uddin, 2020). We suggest that a thorough understanding of variability both *between* and *within* complex biological systems is required in spinal fMRI and, more broadly, functional imaging per se. Current paradigms, optimised for group-level inference, offer only limited clinical utility in understanding individual neural processing, for example, those relating to physical trauma, disease progression, or treatment response and recovery. Spinal fMRI still requires ongoing technical improvement and optimisation, but our opinion is that this should not be at the expense of understanding both group- and individual-level signal variability. The combination of traditional fMRI approaches with novel fingerprinting and dense sampling frameworks is a promising avenue for characterising fine-grained spatial and temporal properties of neural function (Finn & Rosenberg, 2021; Michon et al., 2022; Naselaris et al., 2021; Ricchi et al., 2025). We suggest that validating the utility of such approaches in preliminary healthy volunteer studies should catalyse the deployment of spinal fMRI in clinical research.

4.6 Conclusions

Our findings reveal motor-related activation in ipsilateral ventro-dorsal regions of the cervical spinal segments C6-T1, consistent with established dermatomal and myotomal organisation. We also detected activation in segments C1-C3 and T2, likely reflecting movement-related sensory feedback, propriospinal, or postural control mechanisms. Despite consistent task performance and stable overall fMRI signal, the reliability of motor-evoked activation was low. These findings highlight an ongoing challenge in disentangling the impact of measurement error from within-individual variability in understanding reliability in imaging research.

Data and code availability

The participants of this study did not give written consent for their data to be shared publicly, therefore raw study data are not available.

Spatial maps generated from group-level models and reliability assessments are available on NeuroVault: https://identifiers.org/neurovault.collection:21749. Please note that currently NeuroVault does not support a spinal cord (PAM50) template for online visualisation of spatial maps. Consequently, we recommend downloading the maps and overlaying them on a PAM50 spinal cord template in your software of choice.

All code used in the analysis of this data is openly accessible on the project's Open Science Framework (OSF) repository: https://osf.io/fjasd/ and GitHub: https://github.com/oliviakowalczyk/spain_squeeze. The data that support the findings of this study are available on request from the corresponding author. The data are not publicly available due to privacy or ethical restrictions.

Author contributions

Conceptualization: O.S.K., S.M., D.T., S.C.R.W., J.C.W.B., D.J.L., and M.A.H.

Data curation: O.S.K, A.V.

Formal analysis: O.S.K., A.V.

Funding acquisition: S.C.R.W., J.C.W.B., and M.A.H.

Investigation: O.S.K., S.M., and M.A.H.

Methodology: O.S.K., S.M., D.T., J.C.W.B., D.J.L., and M.A.H.

Project administration: O.S.K., S.M., and M.A.H.

Resources: O.S.K., S.M., and M.A.H.

Software: O.S.K. and D.J.L.

Supervision: M.A.H.

Validation: O.S.K., S.M., J.C.W.B., D.J.L., and M.A.H.

Visualization: O.S.K. and A.V.

Writing – original draft: O.S.K.

Writing – review and editing: O.S.K., S.M., A.V., D.T., A.I.A., S.C.R.W., J.C.W.B., D.J.L., and M.A.H.

Funding

This article represents independent research funded by the Medical Research Council (MRC) Experimental Medicine Challenge Grant (MR/N026969/1) and the National Institute for Health Research (NIHR) Maudsley Biomedical Research Centre at South London and Maudsley NHS Foundation Trust and King's College London. The views

expressed are those of the authors and not necessarily those of the NHS, the NIHR, the
MRC, or the Department of Health and Social Care. O.S.K. is supported by a King's Prize
Fellowship.

Declaration of competing interests

The authors declare no competing of interest.

Acknowledgements

We thank the radiographers for their help with MRI scanning and all volunteers for their
participation in the study.

References

Azad, R., Rouhier, L., & Cohen-Adad, J. (2021). Stacked Hourglass Network with a Multi-level Attention Mechanism: Where to Look for Intervertebral Disc Labeling. In C. Lian, X. Cao, I. Rekik, X. Xu, & P. Yan (Eds), *Machine Learning in Medical Imaging* (pp. 406–415). Springer International Publishing. https://doi.org/10.1007/978-3-030-87589-3_42

Backes, W. H., Mess, W. H., & Wilmink, J. T. (2001). Functional MR Imaging of the Cervical Spinal Cord by Use of Median Nerve Stimulation and Fist Clenching. *American Journal of Neuroradiology*, *22*(10), 1854–1859.

Banerjee, R., Kaptan, M., Tinnermann, A., Khatibi, A., Dabbagh, A., Büchel, C., Kündig, C. W., Law, C. S. W., Pfyffer, D., Lythgoe, D. J., Tsivaka, D., Van De Ville, D., Eippert, F., Muhammad, F., Glover, G. H., David, G., Haynes, G., Haaker, J., Brooks, J. C. W., … Cohen-Adad, J. (2025). EPISeg: Automated segmentation of the spinal cord on echo planar images using open-access multi-center data. *Imaging Neuroscience*. https://doi.org/10.1162/IMAG.a.98

Barry, R. L., Conrad, B. N., Maki, S., Watchmaker, J. M., McKeithan, L. J., Box, B. A., Weinberg, Q. R., Smith, S. A., & Gore, J. C. (2021). Multi-shot acquisitions for stimulus-evoked spinal cord BOLD fMRI. *Magnetic Resonance in Medicine*, *85*(4), 2016–2026. https://doi.org/10.1002/mrm.28570

Barry, R. L., Rogers, B. P., Conrad, B. N., Smith, S. A., & Gore, J. C. (2016). Reproducibility of resting state spinal cord networks in healthy volunteers at 7 Tesla. *NeuroImage*, *133*, 31–40. https://doi.org/10.1016/j.neuroimage.2016.02.058

Bennett, C. M., & Miller, M. B. (2010). How reliable are the results from functional Magnetic Resonance Imaging? *Annals of the New York Academy of Sciences*, *1191*, 133–155. https://doi.org/10.1111/j.1749-6632.2010.05446.x

Bennett, C. M., & Miller, M. B. (2013). fMRI reliability: Influences of task and experimental design. *Cognitive, Affective, & Behavioral Neuroscience*, *13*(4), 690–702. https://doi.org/10.3758/s13415-013-0195-1

Birn, R. M., Cornejo, M. D., Molloy, E. K., Patriat, R., Meier, T. B., Kirk, G. R., Nair, V. A., Meyerand, M. E., & Prabhakaran, V. (2014). The influence of physiological noise correction on test-retest reliability of resting-state functional connectivity. *Brain Connectivity*, *4*(7), 511–522. https://doi.org/10.1089/brain.2014.0284

Bouwman, C. J. C., Wilmink, J. T., Mess, W. H., & Backes, W. H. (2008). Spinal cord functional MRI at 3 T: Gradient echo echo-planar imaging versus turbo spin echo. *NeuroImage*, *43*(2), 288–296. https://doi.org/10.1016/j.neuroimage.2008.07.024

Bozorgpour, A., Azad, B., Azad, R., Velichko, Y., Bagci, U., & Merhof, D. (2023). *HCA-Net: Hierarchical Context Attention Network for Intervertebral Disc Semantic Labeling* (No. arXiv:2311.12486). arXiv. https://doi.org/10.48550/arXiv.2311.12486

Braaß, H., Feldheim, J., Chu, Y., Tinnermann, A., Finsterbusch, J., Büchel, C., Schulz, R., & Gerloff, C. (2023). Association between activity in the ventral premotor cortex and spinal cord activation during force generation—A combined cortico-spinal

fMRI study. *Human Brain Mapping*, *44*(18), 6471–6483. https://doi.org/10.1002/hbm.26523

Brooks, J. C. W., Beckmann, C. F., Miller, K. L., Wise, R. G., Porro, C. A., Tracey, I., & Jenkinson, M. (2008). Physiological noise modelling for spinal functional magnetic resonance imaging studies. *NeuroImage*, *39*(2), 680–692. https://doi.org/10.1016/j.neuroimage.2007.09.018

Caceres, A., Hall, D. L., Zelaya, F. O., Williams, S. C. R., & Mehta, M. A. (2009). Measuring fMRI reliability with the intra-class correlation coefficient. *NeuroImage*, *45*(3), 758–768. https://doi.org/10.1016/j.neuroimage.2008.12.035

Cadotte, D. W., Cadotte, A., Cohen-Adad, J., Fleet, D., Livne, M., Wilson, J. R., Mikulis, D., Nugaeva, N., & Fehlings, M. G. (2015). Characterizing the Location of Spinal and Vertebral Levels in the Human Cervical Spinal Cord. *American Journal of Neuroradiology*, *36*(4), 803–810. https://doi.org/10.3174/ajnr.A4192

Chu, Y., Fricke, B., & Finsterbusch, J. (2023). Improving T2*-weighted human cortico-spinal acquisitions with a dedicated algorithm for region-wise shimming. *NeuroImage*, *268*, 119868. https://doi.org/10.1016/j.neuroimage.2023.119868

Cicchetti, D. V. (1994). Guidelines, criteria, and rules of thumb for evaluating normed and standardized assessment instruments in psychology. *Psychological Assessment*, *6*(4), 284–290. https://doi.org/10.1037/1040-3590.6.4.284

Cohen-Adad, J., Alonso-Ortiz, E., Abramovic, M., Arneitz, C., Atcheson, N., Barlow, L., Barry, R. L., Barth, M., Battiston, M., Büchel, C., Budde, M., Callot, V., Combes, A. J. E., De Leener, B., Descoteaux, M., de Sousa, P. L., Dostál, M., Doyon, J., Dvorak, A., … Xu, J. (2021). Generic acquisition protocol for quantitative MRI of the spinal cord. *Nature Protocols*, *16*(10), 4611–4632. https://doi.org/10.1038/s41596-021-00588-0

Cox, R. W. (1996). AFNI: Software for analysis and visualization of functional magnetic resonance neuroimages. *Computers and Biomedical Research, an International Journal*, *29*(3), 162–173. https://doi.org/10.1006/cbmr.1996.0014

Cox, R. W., & Hyde, J. S. (1997). Software tools for analysis and visualization of fMRI data. *NMR in Biomedicine*, *10*(4–5), 171–178. https://doi.org/10.1002/(sici)1099-1492(199706/08)10:4/5%253C171::aid-nbm453%253E3.0.co;2-l

Cremers, H. R., Wager, T. D., & Yarkoni, T. (2017). The relation between statistical power and inference in fMRI. *PLOS ONE*, *12*(11), e0184923. https://doi.org/10.1371/journal.pone.0184923

Dabbagh, A., Horn, U., Kaptan, M., Mildner, T., Müller, R., Lepsien, J., Weiskopf, N., Brooks, J. C. W., Finsterbusch, J., & Eippert, F. (2024). Reliability of task-based fMRI in the dorsal horn of the human spinal cord. *Imaging Neuroscience*, *2*, 1–27. https://doi.org/10.1162/imag_a_00273

Dawson-Amoah, K., & Varacallo, M. A. (2025). Anatomy, Shoulder and Upper Limb, Hand Intrinsic Muscles. In *StatPearls*. StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK539810/

De Leener, B., Fonov, V. S., Collins, D. L., Callot, V., Stikov, N., & Cohen-Adad, J. (2018). PAM50: Unbiased multimodal template of the brainstem and spinal cord aligned with the ICBM152 space. *NeuroImage*, *165*, 170–179. https://doi.org/10.1016/j.neuroimage.2017.10.041

De Leener, B., Lévy, S., Dupont, S. M., Fonov, V. S., Stikov, N., Louis Collins, D., Callot, V., & Cohen-Adad, J. (2017). SCT: Spinal Cord Toolbox, an open-source software for processing spinal cord MRI data. *NeuroImage*, *145*, 24–43. https://doi.org/10.1016/j.neuroimage.2016.10.009

Dinstein, I., Hasson, U., Rubin, N., & Heeger, D. J. (2007). Brain areas selective for both observed and executed movements. *Journal of Neurophysiology*, *98*(3), 1415–1427. https://doi.org/10.1152/jn.00238.2007

Elliott, M. L., Knodt, A. R., & Hariri, A. R. (2021). Striving toward translation: Strategies for reliable fMRI measurement. *Trends in Cognitive Sciences*, *25*(9), 776–787. https://doi.org/10.1016/j.tics.2021.05.008

Elliott, M. L., Knodt, A. R., Ireland, D., Morris, M. L., Poulton, R., Ramrakha, S., Sison, M. L., Moffitt, T. E., Caspi, A., & Hariri, A. R. (2020). What Is the Test-Retest Reliability of Common Task-Functional MRI Measures? New Empirical Evidence and a Meta-Analysis. *Psychological Science*, *31*(7), 792–806. https://doi.org/10.1177/0956797620916786

Enoka, R. M. (2008). *Neuromechanics of Human Movement*. Human Kinetics.

Finn, E. S., & Rosenberg, M. D. (2021). Beyond fingerprinting: Choosing predictive connectomes over reliable connectomes. *NeuroImage*, *239*, 118254. https://doi.org/10.1016/j.neuroimage.2021.118254

Fleiss, J. L., Levin, B., & Paik, M. C. (2013). *Statistical methods for rates and proportions*. John Wiley & Sons.

Flynn, J. R., Graham, B. A., Galea, M. P., & Callister, R. J. (2011). The role of propriospinal interneurons in recovery from spinal cord injury. *Neuropharmacology*, *60*(5), 809–822. https://doi.org/10.1016/j.neuropharm.2011.01.016

Fröhner, J. H., Teckentrup, V., Smolka, M. N., & Kroemer, N. B. (2019). Addressing the reliability fallacy in fMRI: Similar group effects may arise from unreliable individual effects. *NeuroImage*, *195*, 174–189. https://doi.org/10.1016/j.neuroimage.2019.03.053

Ganapathy, M. K., Reddy, V., & Tadi, P. (2025). Neuroanatomy, Spinal Cord Morphology. In *StatPearls*. StatPearls Publishing. http://www.ncbi.nlm.nih.gov/books/NBK545206/

Giulietti, G., Giove, F., Garreffa, G., Colonnese, C., Mangia, S., & Maraviglia, B. (2008). Characterization of the functional response in the human spinal cord: Impulse-response function and linearity. *NeuroImage*, *42*(2), 626–634. https://doi.org/10.1016/j.neuroimage.2008.05.006

Govers, N., Béghin, J., Van Goethem, J. W. M., Michiels, J., van den Hauwe, L., Vandervliet, E., & Parizel, P. M. (2007). Functional MRI of the cervical spinal cord on 1.5 T with

fingertapping: To what extent is it feasible? *Neuroradiology*, *49*(1), 73–81.
https://doi.org/10.1007/s00234-006-0162-4

Gros, C., De Leener, B., Badji, A., Maranzano, J., Eden, D., Dupont, S. M., Talbott, J.,
Zhuoquiong, R., Liu, Y., Granberg, T., Ouellette, R., Tachibana, Y., Hori, M., Kamiya,
K., Chougar, L., Stawiarz, L., Hillert, J., Bannier, E., Kerbrat, A., … Cohen-Adad, J.
(2019). Automatic segmentation of the spinal cord and intramedullary multiple
sclerosis lesions with convolutional neural networks. *NeuroImage*, *184*, 901–915.
https://doi.org/10.1016/j.neuroimage.2018.09.081

Hamilton, A. F. D. C., & Grafton, S. T. (2008). Repetition suppression for performed hand
gestures revealed by fMRI. *Human Brain Mapping*, *30*(9), 2898–2906.
https://doi.org/10.1002/hbm.20717

Hanna-Boutros, B., Sangari, S., Karasu, A., Giboin, L.-S., & Marchand-Pauvert, V. (2014).
Task-related modulation of crossed spinal inhibition between human lower limbs.
Journal of Neurophysiology, *111*(9), 1865–1876.
https://doi.org/10.1152/jn.00838.2013

Harrow-Mortelliti, M., Reddy, V., & Jimsheleishvili, G. (2025). Physiology, Spinal Cord. In
StatPearls. StatPearls Publishing.
http://www.ncbi.nlm.nih.gov/books/NBK544267/

Hedge, C., Powell, G., & Sumner, P. (2018). The reliability paradox: Why robust cognitive
tasks do not produce reliable individual differences. *Behavior Research Methods*,
50(3), 1166–1186. https://doi.org/10.3758/s13428-017-0935-1

Hemmerling, K. J., Hoggarth, M. A., Sandhu, M. S., Parrish, T. B., & Bright, M. G. (2023).
Spatial distribution of hand-grasp motor task activity in spinal cord functional
magnetic resonance imaging. *Human Brain Mapping*, *44*(17), 5567–5581.
https://doi.org/10.1002/hbm.26458

Hu, Y., Jin, R., Li, G., Luk, K. D., & Wu, Ed. X. (2018). Robust spinal cord resting-state fMRI
using independent component analysis-based nuisance regression noise
reduction. *Journal of Magnetic Resonance Imaging*, *48*(5), 1421–1431.
https://doi.org/10.1002/jmri.26048

Islam, H., Law, C. S. W., Weber, K. A., Mackey, S. C., & Glover, G. H. (2019). Dynamic per
slice shimming for simultaneous brain and spinal cord fMRI. *Magnetic Resonance
in Medicine*, *81*(2), 825–838. https://doi.org/10.1002/mrm.27388

Jenkinson, M., Beckmann, C. F., Behrens, T. E. J., Woolrich, M. W., & Smith, S. M. (2012).
FSL. *NeuroImage*, *62*(2), 782–790.
https://doi.org/10.1016/j.neuroimage.2011.09.015

Kaptan, M., Horn, U., Vannesjo, S. J., Mildner, T., Weiskopf, N., Finsterbusch, J., Brooks, J.
C. W., & Eippert, F. (2022). *Reliability of resting-state functional connectivity in the
human spinal cord: Assessing the impact of distinct noise sources* (p.
2022.12.23.521768). bioRxiv. https://doi.org/10.1101/2022.12.23.521768

Kaptan, M., Pfyffer, D., Konstantopoulos, C. G., Law, C. S. W., Weber II, K. A., Glover, G.
H., & Mackey, S. (2024). Recent developments and future avenues for human
corticospinal neuroimaging. *Frontiers in Human Neuroscience*, *18*, 1339881.
https://doi.org/10.3389/fnhum.2024.1339881

Khatibi, A., Vahdat, S., Lungu, O., Finsterbusch, J., Büchel, C., Cohen-Adad, J., Marchand-Pauvert, V., & Doyon, J. (2022). Brain-spinal cord interaction in long-term motor sequence learning in human: An fMRI study. *NeuroImage*, *253*, 119111. https://doi.org/10.1016/j.neuroimage.2022.119111

Kinany, N., Khatibi, A., Lungu, O., Finsterbusch, J., Büchel, C., Marchand-Pauvert, V., Van De Ville, D., Vahdat, S., & Doyon, J. (2023). Decoding cerebro-spinal signatures of human behavior: Application to motor sequence learning. *NeuroImage*, *275*, 120174. https://doi.org/10.1016/j.neuroimage.2023.120174

Kinany, N., Pirondini, E., Martuzzi, R., Mattera, L., Micera, S., & Van de Ville, D. (2019). Functional imaging of rostrocaudal spinal activity during upper limb motor tasks. *NeuroImage*, *200*, 590–600. https://doi.org/10.1016/j.neuroimage.2019.05.036

Kinany, N., Pirondini, E., Micera, S., & Van De Ville, D. (2020). Dynamic Functional Connectivity of Resting-State Spinal Cord fMRI Reveals Fine-Grained Intrinsic Architecture. *Neuron*, *108*(3), 424-435.e4. https://doi.org/10.1016/j.neuron.2020.07.024

Kinany, N., Pirondini, E., Micera, S., & Van De Ville, D. (2022). Spinal Cord fMRI: A New Window into the Central Nervous System. *The Neuroscientist*, 10738584221101827. https://doi.org/10.1177/10738584221101827

Kowalczyk, O. S., Medina, S., Howard, M. A., Tsivaka, D., Lythgoe, D. J., & Brooks, J. C. W. (2021). *Examining test-retest reliability of evoked response and resting-state functional MRI endpoints in human brain and cervical spine*. https://doi.org/10.17605/osf.io/u8rbq

Kowalczyk, O. S., Medina, S., Tsivaka, D., McMahon, S. B., Williams, S. C. R., Brooks, J. C. W., Lythgoe, D. J., & Howard, M. A. (2024). Spinal fMRI demonstrates segmental organisation of functionally connected networks in the cervical spinal cord: A test–retest reliability study. *Human Brain Mapping*, *45*(2), e26600. https://doi.org/10.1002/hbm.26600

Kragel, P. A., Han, X., Kraynak, T. E., Gianaros, P. J., & Wager, T. D. (2021). Functional MRI Can Be Highly Reliable, but It Depends on What You Measure: A Commentary on Elliott et al. (2020). *Psychological Science*, *32*(4), 622–626. https://doi.org/10.1177/0956797621989730

Landelle, C., Lungu, O., Vahdat, S., Kavounoudias, A., Marchand-Pauvert, V., De Leener, B., & Doyon, J. (2021). Investigating the human spinal sensorimotor pathways through functional magnetic resonance imaging. *NeuroImage*, *245*, 118684. https://doi.org/10.1016/j.neuroimage.2021.118684

Lang, J., & Bartram, C. T. (1982). Fila radicularia of the ventral and dorsal radices of the human spinal cord. *Gegenbaurs morphologisches Jahrbuch*, *128*(4), 417–462.

Lemon, R. N. (2008). Descending Pathways in Motor Control. *Annual Review of Neuroscience*, *31*(Volume 31, 2008), 195–218. https://doi.org/10.1146/annurev.neuro.31.060407.125547

Lipp, I., Murphy, K., Wise, R. G., & Caseras, X. (2014). Understanding the contribution of neural and physiological signal variation to the low repeatability of emotion-

induced BOLD responses. *NeuroImage*, *86*, 335–342. https://doi.org/10.1016/j.neuroimage.2013.10.015

Liu, X., Zhou, F., Li, X., Qian, W., Cui, J., Zhou, I. Y., Luk, K. D. K., Wu, Ed. X., & Hu, Y. (2016). Organization of the intrinsic functional network in the cervical spinal cord: A resting state functional MRI study. *Neuroscience*, *336*, 30–38. https://doi.org/10.1016/j.neuroscience.2016.08.042

Madi, S., Flanders, A. E., Vinitski, S., Herbison, G. J., & Nissanov, J. (2001). Functional MR Imaging of the Human Cervical Spinal Cord. *American Journal of Neuroradiology*, *22*(9), 1768–1774.

Maieron, M., Iannetti, G. D., Bodurka, J., Tracey, I., Bandettini, P. A., & Porro, C. A. (2007). Functional Responses in the Human Spinal Cord during Willed Motor Actions: Evidence for Side- and Rate-Dependent Activity. *Journal of Neuroscience*, *27*(15), 4182–4190. https://doi.org/10.1523/JNEUROSCI.3910-06.2007

McCarthy, P. (2022). FSLeyes (1.5.0). *Zenodo*. https://doi.org/10.5281/zenodo.7038115

Michon, K. J., Khammash, D., Simmonite, M., Hamlin, A. M., & Polk, T. A. (2022). Person-specific and precision neuroimaging: Current methods and future directions. *NeuroImage*, *263*, 119589. https://doi.org/10.1016/j.neuroimage.2022.119589

Mrachacz-Kersting, N., Gervasio, S., & Marchand-Pauvert, V. (2018). Evidence for a Supraspinal Contribution to the Human Crossed Reflex Response During Human Walking. *Frontiers in Human Neuroscience*, *12*. https://doi.org/10.3389/fnhum.2018.00260

Murphy, K., Bodurka, J., & Bandettini, P. A. (2007). How long to scan? The relationship between fMRI temporal signal to noise and necessary scan duration. *NeuroImage*, *34*(2), 565–574. https://doi.org/10.1016/j.neuroimage.2006.09.032

Naselaris, T., Allen, E., & Kay, K. (2021). Extensive sampling for complete models of individual brains. *Current Opinion in Behavioral Sciences*, *40*, 45–51. https://doi.org/10.1016/j.cobeha.2020.12.008

Ng, M. C., Wong, K. K., Li, G., Lai, S., Yang, E. S., Hu, Y., & Luk, K. D. (2006). Proton-density-weighted spinal fMRI with sensorimotor stimulation at 0.2 T. *NeuroImage*, *29*(3), 995–999. https://doi.org/10.1016/j.neuroimage.2005.08.011

Ng, M. C., Wu, E. X., Lau, H.-F., Hu, Y., Lam, E. Y., & Luk, K. D. (2008). Cervical spinal cord BOLD fMRI study: Modulation of functional activation by dexterity of dominant and non-dominant hands. *NeuroImage*, *39*(2), 825–831. https://doi.org/10.1016/j.neuroimage.2007.09.026

Nielsen, J. B. (2016). Human Spinal Motor Control. *Annual Review of Neuroscience*, *39*(Volume 39, 2016), 81–101. https://doi.org/10.1146/annurev-neuro-070815-013913

Noble, S., Scheinost, D., & Constable, R. T. (2019). A decade of test-retest reliability of functional connectivity: A systematic review and meta-analysis. *NeuroImage*, *203*, 116157. https://doi.org/10.1016/j.neuroimage.2019.116157

Noble, S., Spann, M. N., Tokoglu, F., Shen, X., Constable, R. T., & Scheinost, D. (2017). Influences on the Test–Retest Reliability of Functional Connectivity MRI and its

Relationship with Behavioral Utility. *Cerebral Cortex*, *27*(11), 5415–5429. https://doi.org/10.1093/cercor/bhx230

Oldfield, R. C. (1971). The assessment and analysis of handedness: The Edinburgh inventory. *Neuropsychologia*, *9*(1), 97–113. https://doi.org/10.1016/0028-3932(71)90067-4

Oliva, V., Bédard, S., Kaptan, M., Pfyffer, D., Chy, B., Aufrichtig, S., Berhe, N., Chaudhari, A. S., Tharin, S., Hu, S. S., Ratliff, J., Smith, Z. A., Smith, A. C., Glover, G. H., Mackey, S., Law, C. S. W., & Weber, K. A. (2025). *Mapping Hand Function with Simultaneous Brain-Spinal Cord Functional MRI* (p. 2025.02.27.640504). bioRxiv. https://doi.org/10.1101/2025.02.27.640504

Oliva, V., Gregory, R., Brooks, J. C. W., & Pickering, A. E. (2022). Central pain modulatory mechanisms of attentional analgesia are preserved in fibromyalgia. *Pain*, *163*(1), 125–136. https://doi.org/10.1097/j.pain.0000000000002319

Pierrot-Deseilligny, E., & Burke, D. (2012). *The Circuitry of the Human Spinal Cord: Spinal and Corticospinal Mechanisms of Movement*. Cambridge University Press.

Powers, J. M., Ioachim, G., & Stroman, P. W. (2018). Ten Key Insights into the Use of Spinal Cord fMRI. *Brain Sciences*, *8*(9), E173. https://doi.org/10.3390/brainsci8090173

Ricchi, I., Santoro, A., Kinany, N., Landelle, C., Khatibi, A., Vahdat, S., Doyon, J., Barry, R. L., & Ville, D. V. D. (2025). *Spine-Prints: Transposing Brain Fingerprints to the Spinal Cord* (p. 2025.05.30.656545). bioRxiv. https://doi.org/10.1101/2025.05.30.656545

Rouhier, L., Romero, F. P., Cohen, J. P., & Cohen-Adad, J. (2020). *Spine intervertebral disc labeling using a fully convolutional redundant counting model* (No. arXiv:2003.04387). arXiv. https://doi.org/10.48550/arXiv.2003.04387

Rowe, J. B., & Siebner, H. R. (2012). The motor system and its disorders. *NeuroImage*, *61*(2), 464–477. https://doi.org/10.1016/j.neuroimage.2011.12.042

Schirmer, C. M., Shils, J. L., Arle, J. E., Cosgrove, G. R., Dempsey, P. K., Tarlov, E., Kim, S., Martin, C. J., Feltz, C., Moul, M., & Magge, S. (2011). *Heuristic map of myotomal innervation in humans using direct intraoperative nerve root stimulation*. https://doi.org/10.3171/2011.2.SPINE1068

Schuster, V., & Jansen, A. (2022). 'That Time of the Month' – Investigating the Influence of the Menstrual Cycle and Oral Contraceptives on the Brain Using Magnetic Resonance Imaging. *Experimental and Clinical Endocrinology & Diabetes*, *130*, 303–312. https://doi.org/10.1055/a-1816-8203

Smith, S. M., Jenkinson, M., Woolrich, M. W., Beckmann, C. F., Behrens, T. E. J., Johansen-Berg, H., Bannister, P. R., De Luca, M., Drobnjak, I., Flitney, D. E., Niazy, R. K., Saunders, J., Vickers, J., Zhang, Y., De Stefano, N., Brady, J. M., & Matthews, P. M. (2004). Advances in functional and structural MR image analysis and implementation as FSL. *NeuroImage*, *23 Suppl 1*, S208-219. https://doi.org/10.1016/j.neuroimage.2004.07.051

Sonoo, M. (2023). Recent advances in neuroanatomy: The myotome update. *Journal of Neurology, Neurosurgery & Psychiatry*, *94*(8), 643–648. https://doi.org/10.1136/jnnp-2022-329696

Soteropoulos, D. S., Edgley, S. A., & Baker, S. N. (2013). Spinal Commissural Connections to Motoneurons Controlling the Primate Hand and Wrist. *Journal of Neuroscience*, *33*(23), 9614–9625. https://doi.org/10.1523/JNEUROSCI.0269-13.2013

Spielberger, C. D., Gonzalez-Reigosa, F., Martinez-Urrutia, A., Natalicio, L. F., & Natalicio, D. S. (1971). The State-Trait Anxiety Inventory. *Revista Interamericana De Psicología/Interamerican Journal of Psychology*, *5*(3–4). https://journal.sipsych.org/index.php/IJP/article/view/620

Stracke, C. P., Pettersson, L. G., Schoth, F., Möller-Hartmann, W., & Krings, T. (2005). Interneuronal systems of the cervical spinal cord assessed with BOLD imaging at 1.5 T. *Neuroradiology*, *47*(2), 127–133. https://doi.org/10.1007/s00234-004-1318-8

Stroman, P. W., Nance, P. W., & Ryner, L. N. (1999). BOLD MRI of the human cervical spinal cord at 3 tesla. *Magnetic Resonance in Medicine*, *42*(3), 571–576. https://doi.org/10.1002/(SICI)1522-2594(199909)42:3%253C571::AID-MRM20%253E3.0.CO;2-N

Stroman, P. W., & Ryner, L. N. (2001). Functional MRI of motor and sensory activation in the human spinal cord. *Magnetic Resonance Imaging*, *19*(1), 27–32. https://doi.org/10.1016/S0730-725X(01)00226-0

Stroman, P. W., Wheeler-Kingshott, C., Bacon, M., Schwab, J. M., Bosma, R., Brooks, J., Cadotte, D., Carlstedt, T., Ciccarelli, O., Cohen-Adad, J., Curt, A., Evangelou, N., Fehlings, M. G., Filippi, M., Kelley, B. J., Kollias, S., Mackay, A., Porro, C. A., Smith, S., … Tracey, I. (2014). The current state-of-the-art of spinal cord imaging: Methods. *NeuroImage*, *84*, 1070–1081. https://doi.org/10.1016/j.neuroimage.2013.04.124

Takei, T., & Seki, K. (2010). Spinal Interneurons Facilitate Coactivation of Hand Muscles during a Precision Grip Task in Monkeys. *The Journal of Neuroscience*, *30*(50), 17041–17050. https://doi.org/10.1523/JNEUROSCI.4297-10.2010

Tsivaka, D., Williams, S. C. R., Medina, S., Kowalczyk, O. S., Brooks, J. C. W., Howard, M. A., Lythgoe, D. J., & Tsougos, I. (2023). A second-order and slice-specific linear shimming technique to improve spinal cord fMRI. *Magnetic Resonance Imaging*. https://doi.org/10.1016/j.mri.2023.06.012

Uddin, L. Q. (2020). Bring the Noise: Reconceptualizing Spontaneous Neural Activity. *Trends in Cognitive Sciences*, *24*(9), 734–746. https://doi.org/10.1016/j.tics.2020.06.003

Ullmann, E., Pelletier Paquette, J. F., Thong, W. E., & Cohen-Adad, J. (2014). Automatic Labeling of Vertebral Levels Using a Robust Template-Based Approach. *International Journal of Biomedical Imaging*, *2014*(1), 719520. https://doi.org/10.1155/2014/719520

Vahdat, S., Lungu, O., Cohen-Adad, J., Marchand-Pauvert, V., Benali, H., & Doyon, J. (2015). Simultaneous Brain–Cervical Cord fMRI Reveals Intrinsic Spinal Cord

Plasticity during Motor Sequence Learning. *PLOS Biology*, *13*(6), e1002186. https://doi.org/10.1371/journal.pbio.1002186

Valošek, J., Mathieu, T., Schlienger, R., Kowalczyk, O. S., & Cohen-Adad, J. (2024). Automatic segmentation of the spinal cord nerve rootlets. *Imaging Neuroscience*, *2*, 1–14. https://doi.org/10.1162/imag_a_00218

Vania, M., & Lee, D. (2021). Intervertebral disc instance segmentation using a multistage optimization mask-RCNN (MOM-RCNN). *Journal of Computational Design and Engineering*, *8*(4), 1023–1036. https://doi.org/10.1093/jcde/qwab030

Walji, A. H., & Tsui, B. C. H. (2016). Clinical Anatomy of the Dermatomes and Innervation of the Joints. In B. C. H. Tsui & S. Suresh (Eds), *Pediatric Atlas of Ultrasound- and Nerve Stimulation-Guided Regional Anesthesia* (pp. 205–222). Springer. https://doi.org/10.1007/978-0-387-79964-3_14

Webb, A. J. S., Klerman, E. B., & Mandeville, E. T. (2024). Circadian and Diurnal Regulation of Cerebral Blood Flow. *Circulation Research*, *134*(6), 695–710. https://doi.org/10.1161/CIRCRESAHA.123.323049

Weber, K. A., Chen, Y., Paliwal, M., Law, C. S., Hopkins, B. S., Mackey, S., Dhaher, Y., Parrish, T. B., & Smith, Z. A. (2020). Assessing the spatial distribution of cervical spinal cord activity during tactile stimulation of the upper extremity in humans with functional magnetic resonance imaging. *NeuroImage*, *217*, 116905. https://doi.org/10.1016/j.neuroimage.2020.116905

Weber, K. A., Chen, Y., Wang, X., Kahnt, T., & Parrish, T. B. (2016a). Functional magnetic resonance imaging of the cervical spinal cord during thermal stimulation across consecutive runs. *NeuroImage*, *143*, 267–279. https://doi.org/10.1016/j.neuroimage.2016.09.015

Weber, K. A., Chen, Y., Wang, X., Kahnt, T., & Parrish, T. B. (2016b). Lateralization of cervical spinal cord activity during an isometric upper extremity motor task with functional magnetic resonance imaging. *NeuroImage*, *125*, 233–243. https://doi.org/10.1016/j.neuroimage.2015.10.014

Worsley, K. J. (2003). *Developments in Random Field Theory*.

Yeganegi, H., Fathi, Y., & Erfanian, A. (2018). Decoding hind limb kinematics from neuronal activity of the dorsal horn neurons using multiple level learning algorithm. *Scientific Reports*, *8*(1), 577. https://doi.org/10.1038/s41598-017-18971-x

Yoshizawa, T., Nose, T., Moore, G. J., & Sillerud, L. O. (1996). Functional magnetic resonance imaging of motor activation in the human cervical spinal cord. *NeuroImage*, *4*(3 Pt 1), 174–182. https://doi.org/10.1006/nimg.1996.0068

A. Whole cervical cord

B. ROIs