Endogenous corazonin signaling modulates the post-mating switch in behavior and physiology in females of the brown planthopper and *Drosophila*

Ning Zhang (张宁)[1], Shao-Cong Su (苏少聪)[1], Ruo-Tong Bu (卜若彤)[1], Yi-Jie Zhang (张伊杰)[1], Lei Yang (杨磊)[1], Jie Chen (陈洁)[1], Dick R. Nässel[2], Cong-Fen Gao (高聪芬)[1], Shun-Fan Wu (吴顺凡)[1*]

Orcid ID numbers of Shun-Fan Wu: 0000-0003-0096-147X

Orcid ID numbers of Dick Nässel: 0000-0002-1147-7766

Affiliations:

[1]State Key Laboratory of Agricultural and Forestry Biosecurity, College of Plant Protection, Nanjing Agricultural University, Nanjing 210095, China

[2]Department of Zoology, Stockholm University, Stockholm, Sweden

[*]Correspondence: wusf@njau.edu.cn (S.F. Wu).

Abstract

Mating in insects typically triggers a post-mating response (PMR) in females, characterized by reduced receptivity to re-mating and increased oviposition, which ensures numerous and viable offspring and male paternity. This PMR is induced by male seminal factors, such as sex peptide in *Drosophila melanogaster*, as well as intrinsic female signaling components. The latter signaling remains poorly understood in most insects, including the devastating rice pest, the brown planthopper (BPH) *Nilaparvata lugens*. Here, we show that the neuropeptide corazonin (CRZ) and its receptor (CrzR) are critical for the PMR in female BPHs. Peptide injection, RNAi knockdown, and CRISPR/Cas9 mutagenesis confirm that intact CRZ signaling reduces re-mating frequency and increases ovulation in mated BPH females. The CrzR is highly expressed in the female reproductive tract, and CrzR knockdown phenocopies CRZ diminishment. Importantly, female CRZ/CrzR signaling is required for male seminal factors, such as the peptide maccessin, to induce the PMR; with disrupted *CrzR* signaling, injection of seminal fluid or maccessin fails to reduce female receptivity. Notably, CRZ is not produced in male accessory glands (MAGs) of BPHs and thus not transferred during copulation. We furthermore demonstrate that also in *D. melanogaster* disrupted CRZ signaling increases female re-mating and reduces oviposition, while CRZ injection suppresses virgin receptivity and increases oviposition. Finally, we detected no CRZ in the MAG of *D. melanogaster,* supporting its role as an endogenous signal in the female PMR also in this species. In summary, our findings reveal a conserved role of endogenous CRZ signaling in regulating the female PMR and demonstrate that female CRZ signaling and male-derived signals cooperate to induce post-mating transitions in BPHs and *D. melanogaster*. CRZ is a paralog of the peptide gonadotropin-releasing hormone, known to regulate reproduction in vertebrates, including humans, suggesting evolutionary conservation of an ancient function.

Keywords: Post mating response; *Nilaparvara lugens*; *Drosophila melanogaster*; corazonin; corazonin receptor

Introduction

Reproductive behavior in animals is influenced by a complex interplay between

external sensory inputs and internal states, and is regulated by neuronal and

hormonal signaling. In insects, males typically initiate mating through courtship, while

females determine acceptance by integrating external sensory cues with their internal

state. Consequently, female reproductive activity is modulated by multiple signaling

pathways [1-8]. Importantly, in insects, successful copulation commonly induces

profound physiological and behavioral changes in the females [5,9-18].

Following their first mating, insect females typically reject further courtship

attempts and initiate oviposition. This distinct shift in behavior and physiology is

referred to as a post-mating response (PMR) [5-7,19-23]. Induction of a PMR has been

documented across diverse insect taxa, for example in *D. melanogaster*, mosquitos

Anopheles gambiae, Aedes aegypti, crickets (*Gryllodes sigillatus*), and the brown

planthopper (BPH), *Nilaparvata lugens* [6,10,15,18,21,24-28].

Induction of a PMR commonly requires the transfer of specific seminal fluid

components during copulation. In *D. melanogaster*, seminal fluid peptides, including

sex peptide (SP) and DUP99B produced in male accessory glands (MAG), enter the

female reproductive tract, bind to specific receptors on sensory neurons, and activate

signaling pathways that drive post-mating behavior and physiology [6,13,14,18,29].

Additional factors such as Acp26Ab and ovulin (Acp26Aa) are also transferred with

the seminal fluid and modulate female reproductive physiology in *D. melanogaster* [30-33].

Notably, it has been shown that the PMR-inducing substances, produced in the MAG

of various insects, are highly species-specific and not even conserved across related

taxonomic groups [24,34].

While MAG-derived sex peptide (SP) and its receptor (SPR) are key regulators of

the female post-mating response (PMR) in *D. melanogaster* [5-7,16], additional signaling

pathways intrinsic to females also modulate this process. Interneurons within

abdominal neuromeres of the ventral nerve cord (VNC) promote receptivity to males

by signaling via myoinhibitory peptide (MIP) to brain circuits [35]. Furthermore, the

neuropeptide diuretic hormone 44 (DH44) modulates female sexual receptivity

through a sex-specific brain circuit; mating suppresses DH44 signaling, thereby

reducing receptivity [1]. During sexual maturation, leucokinin (LK)-expressing neurons

(ABLKs) in the abdominal ganglion, downstream of SP-activated circuitry, suppress

female receptivity [36]. In addition to changes in receptivity, the *D. melanogaster* PMR

includes a shift in diet preference toward yeast consumption. This altered feeding

behavior is mediated by specific peptidergic neurons (ALKs) in the brain through LK

signaling [12].

Although the mechanisms and neuronal pathways underlying the post-mating

response (PMR) in *D. melanogaster* are well characterized [3,4,6,7,37-40], this process

remains poorly understood in most non-model insects. A prominent example is the

brown planthopper (BPH), *N. lugens* (Hemiptera). As a monophagous rice pest, the

BPH exhibits robust reproductive capacity and remarkable environmental adaptability

[41]. It has also developed high resistance to diverse chemical pesticides [42-44].

Consequently, elucidating reproductive physiology and behavior in BPHs is crucial for

developing novel control strategies.

In BPHs, the recently identified MAG-derived peptide maccessin (macc) is

transferred during copulation and triggers a PMR in females, although its receptor

remains unknown [24]. Notably, the same study demonstrated that a second peptide,

ion transport-like peptide 1 (ITPL-1), also modulates the PMR. Whereas macc is

male-specific, ITPL-1 is produced both in the MAG and endogenously in females,

where it reduces female receptivity to courting males [24]. Given its endogenous female

expression, ITPL-1 likely functions as an intrinsic signaling component in the PMR of

BPHs, analogous to the role of myoinhibitory peptide (MIP) and DH44 in *D.*

melanogaster females [1,35].

Here, we explored the role of another neuropeptide, corazonin (CRZ), in the

reproductive behavior of BPHs. In *D. melanogaster* males, CRZ is released from

interneurons of the ventral nerve cord (VNC) and acts on serotonergic projection

neurons to regulate ejaculation and copulation duration [45], mitigates mating-induced

heart rate acceleration for physiological recovery [46] and influences post-mating dietary

preference [12]. Surprisingly, the role of CRZ signaling in female reproduction remains

unexplored. To address this gap, we investigated the role of CRZ in female

reproduction, including the PMR, in BPHs and *D. melanogaster.*

Here, we demonstrate that CRZ induces a post-mating switch in receptivity in

females of both species. CRZ injection in virgin females reduces receptivity to

courting males and increases oviposition, while *Crz* knockdown or knockout in mated

females restores receptivity to re-mating and decreases oviposition. Notably, CRZ is

not produced in the male accessory gland (MAG) of either species and thus the

peptide is not transferred during copulation, but acting as an endogenous female

peptide signal. In BPHs, the CRZ receptor (CrzR) is highly expressed in the female

reproductive tract and both CRISPR-Cas9-mediated *CrzR* knockout and

RNAi-mediated knockdown disrupt the PMR, phenocopying CRZ deficiency. Critically,

receptivity changes in virgin BPH female, whether induced by MAG extract

(containing seminal fluid proteins) or CRZ injection, depend on presence of the CrzR.

Collectively, our data demonstrate that endogenous female CRZ signaling and male

seminal cues are both required to coordinate post-mating changes in physiology and

behavior in females of both insects. CRZ is a paralog of the peptide

gonadotropin-releasing hormone, known to regulate reproduction in vertebrates,

including humans, suggesting evolutionary conservation of an ancient function.

Results

CRZ signaling reduces receptivity to courting males and stimulates oviposition in *N. lugens* females

While numerous studies have established roles of the neuropeptide CRZ in regulating aspects of male reproduction across diverse insect species [47-53], research addressing its potential functions in female reproduction remains conspicuously lacking.

This study investigates the role of CRZ in female reproduction, starting with the brown planthopper (BPH), *Nilaparvata lugens*, a major agricultural pest. Initially, the *Crz* gene (GenBank accession: AB817247.1) was cloned based on transcriptome data [54]. The deduced mature CRZ sequence in BPH, pQTFQYSRGWTNamide, is identical to that reported for most studied insect species (Figure 1-figure supplement 1A). Consistent with other known CRZ precursors, the BPH precursor encodes a single CRZ peptide starting at the first amino acid of the propeptide (Figure 1-figure supplement 1A and B). A comparison of the organization of selected CRZ precursor genes is shown in Figure 1-figure supplement 1B. Genomic analysis revealed that the CRZ precursor is encoded by two exons separated by a single intron (Figure 1-figure supplement 1B). The mature CRZ peptide and a scrambled control peptide (sCRZ) were synthesized for pharmacological experiments.

First, we investigated the role of CRZ signaling in the reproductive behavior and PMR of female BPHs. The PMR in BPHs, is characterized by reduced receptivity to courting males, including display of rejection behaviors, and increased oviposition [24].

We found that injection of CRZ significantly reduced receptivity of virgin female BPHs

to courting males (Figure 1A and B). Furthermore, virgin females injected with CRZ

actively rejected courting males, exhibiting a distinct rejection behavior characterized

by ovipositor extrusion and kicking or fleeing. This behavior was not observed in

control females injected with sCRZ (Figure 1 Supplementary Video 1). Notably, the

CRZ-induced mating refusal behavior in virgin females was no longer observed 24

hours after injection (Figure 1-figure supplement 2A). While injection of CRZ peptide

did not stimulate egg production in virgin BPHs (Figure 1-figure supplement 2B), it

significantly promoted oviposition in mated females (Figure 1C).

To further investigate CRZ action, we performed RNA interference (RNAi) by

injecting double-stranded *Crz* RNA (*dsNlCrz*) into female BPHs, which efficiently

reduced *Crz* expression (Figure 1-figure supplement 2). In *N. lugens*, female

receptivity is transiently suppressed after mating and does not begin to recover until at

least 72 h after the first mating[24]; therefore, remating assays were performed at 48 h

post-mating as shown in Figure 1D. Knockdown of *Crz* resulted in increased

receptivity with 10% of the mated females mating again, and significantly reduced

oviposition to approximately 60% of control levels (Figure 1E and F). In contrast, *Crz*

knockdown had no effect on the receptivity of virgin females (Figure 1E).

Since RNAi-mediated knockdown of *Crz* in female BPHs left a residual level of

gene expression (Figure 1-figure supplement 2), we generated two *Crz* deletion

alleles, *ΔCrz¹* and *ΔCrz²*, using CRISPR/Cas9 (Figure 1G-G') to achieve a complete

ablation. Both alleles harbor deletions eliminating the mature CRZ peptide coding

sequence (Figure 1G"). Sanger sequencing confirmed that the deletions

compromised essential triplet codons within the *Crz* open reading frame (Figure

1-figure supplement 4), thereby efficiently disrupting CRZ peptide expression (Figure

1G" and H). Immunohistochemical analysis further validated the mutants, revealing a

complete absence of anti-CRZ immunolabeling in homozygous *ΔCrz¹* and *ΔCrz²*

females (Figure 1H). Homozygous *ΔCrz¹* and *ΔCrz²* virgin females are fully viable and

fertile. They exhibit a courtship vigor similar to controls and their initial mating success

rate with wild-type males exceeds 60% (Figure 1I and J). However, 25% of the mated

mutant females were receptive to further mating (Figure 1J), and oviposition was

significantly reduced to approximately 50% of the wild-type levels (Figure 1K). To

further test the *Crz* mutants, we also analyzed mating and oviposition behaviors in *Crz*

heterozygous (*+/ΔCrz¹*) females. Notably, neither the remating rate nor egg

production in heterozygous females differed significantly from those of wild-type

controls (Figure 1J and K). These results indicate that the *Crz* loss-of-function

phenotype is recessive, and that a single functional copy of *Crz* is sufficient to sustain

a normal post-mating response (PMR). Collectively, these results demonstrate that

diminished CRZ signaling severely perturbs the PMR in female BPHs, seen as

increased post-mating receptivity and reduced egg-laying. Importantly, the *Crz*

deletion abolished the onset of typical post-mating rejection behavior. We, thus,

conclude that endogenous CRZ signaling is important for the induction of a PMR in

female BPHs.

Figure 1. CRZ signaling affects the post-mating response (PMR) and oviposition in female *N. lugens.*

Wild-type males were used in all behavioral assays.

(A) Experimental design for panels B-C. The white/black segments on the experimental time lines denote light/dark periods, respectively.

(B) Receptivity of virgin females 6 h post-injection, using different doses of CRZ (sCRZ as control), each virgin female BPH was injected with a calibrated glass capillary needle directly into the abdomen with 100 ng/10 ng/1 ng/0.1 ng of peptide dissolved in 50 nL of 1 × PBS

balanced salt solution. The graph shows percentage females copulating within 30 min. **$P <$ 0.01 vs. control; ns (non-significant): $P > 0.05$ (Mann-Whitney test). The small circles denote the number of replicates, $\geqslant$5 insects/replicate; the numbers above the curves denote total number of animals.

(C) Eggs laid per mated female 12 h post-copulation. We used the highest CRZ dose from panel B (sCRZ, control). Each mated female BPH was injected with a calibrated glass capillary needle directly into the abdomen with 100 ng of peptide dissolved in 50 nL of 1 × PBS balanced salt solution. *$P < 0.05$ (Student's t-test). Data: mean ± s.e.m ($\geqslant$4 biological replicates, $\geqslant$6 insects/replicate). The numbers below the bars denote total number of animals.

(D) Experimental design for panels E-F.

(E) Receptivity of virgin (1st mating) and mated females after *Crz*-RNAi (*dsgfp* as control). Graph displays the percentage of females copulating within 30 min. The small circles denote the number of replicates, $\geqslant$7 insects/replicate; the numbers below the bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, $P < 0.05$ was considered significant. (F) Total numbers of eggs laid per female over 72 h after *Crz*-RNAi (*dsgfp* as control). ****$P < 0.0001$ (Student's *t*-test). The small circles and the numbers below the bars denote total number of animals, ($\geqslant$4 biological replicates, $\geqslant$8 insects/replicate). Data are shown as mean $\pm$ s.e.m. ($\geqslant$3 biological replicates, $\geqslant$8 insects/replicate).

(G) Schematic diagram of the *Crz* gene and sgRNA (single-guide RNA) design for *Crz* mutant production. The *Crz* gene consists of 2 exons, and ATG and TAG are located on Exon 1. The target site of the 20 bp sgRNA on Exon 1 is highlighted in pink. The PAM sites are indicated in yellow. The two knockout strains *ΔCrz^1* and *ΔCrz^2* were unable to produce mature CRZ peptides. (G') The amino acid sequences obtained by translating the mutated base sequences.

(H) Absence of CRZ immunoreactivity in homozygous *ΔCrz^1* and *ΔCrz^2* mutants compared to wild type (WT) expression. Scale bars: 50 μm.

(I) Experimental design for panels J-K.

(J) Receptivity of virgin/mated females across genotypes. No effect was seen on virgin receptivity, only the re-mating was affected. The small circles denote the number of replicates; $\geqslant$6 insects/replicate, the numbers below the bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, $P < 0.05$ was considered significant.

(K) Eggs laid 72 h post-mating. The numbers below the bars denote total number of animals., ≥4 biological replicates, ≥8 insects/replicate. Data are shown as mean ± s.e.m. ****$P < 0.0001$; Student's *t* test.

Figure 1 - figure supplement 1. Evolutionary conservation of CRZ peptide sequences in

Nilaparvata lugens and other arthropods.

(A) Alignment of mature CRZ peptide sequences across species.

Abbreviations of species names are as follows: Bmori (*Bombyx mori*), Gmole (*Grapholita molesta*), Cpomo (*Cydia pomonella*), Sexig (*Spodoptera exigua*), Harmi (*Helicoverpa armigera*), Dplex (*Danaus plexippus*), Dmela (*Drosophila melanogaster*), Csupp (*Chilo suppressalis*), Dmagn (*Daphnia magan*), Dviri (*Drosophila virilis*), Cnodu *(Catagly nodus)*, Carcu (*Callinectes arcuatus*), Nnorv (*Nephrops norvegicus*), Cmaen (*Carcinus maenas*), Mrose (*Macrobrachium rosenbergii*), Mscal (*Megaselia scalaris*), Cglom (*Cotesia glomerata*), Rmicr (*Rhipicephalus microplus*), Derec (*Drosophila erecta*), Agamb (*Anopheles gambiae*), Pstal (*Plautia stali*), Nluge (*Nilaparvata lugens*), Cdraw (*Caerostris darwini*), Cextr (*Caerostris extrusa*), Dpule (*Daphnia pulex*), Hitam (*Heterotrigona itama*), Lmigr (*Locusta migratoria*), Cmoro (*Carausius morosus*), Sgreg (*Schistocerca gregaria*), Lsalm (*Lepeophtheirus salmonis*), Alabo (*Apis laboriosa*). Species within the blue area possess highly conserved mature peptide sequences, whereas those in the magenta zone exhibit relatively less conservation in their mature peptides.

(B) Schematic organization of CRZ precursors in representative species. Signal peptides (blue)

and mature CRZ peptides (red) are indicated.

Figure 1 - figure supplement 2. The *Crz* gene-silencing efficacy in female insects following

dsRNA injection assayed by qPCR. The small circles denote the number of replicates. Data

are shown as mean ± s.e.m. Mann–Whitney test. **$P<0.001$.

Figure 1 - figure supplement 3. Phenotypic effects of CRZ peptide injection.

(A) Receptivity of virgin females at indicated time points after CRZ injection (sCRZ as control).

Each virgin female BPH was injected with a calibrated glass capillary needle directly into the

abdomen with 100 ng of peptide dissolved in 50 nL of 1 × PBS balanced salt solution. Graph

shows percentage females copulating within 30 min. ** $P < 0.01$; ns: not significant Binomial,

GLM with logit link, followed by pairwise contrasts. The small circles denote the number of

replicates, ≥8 insects/replicate; the numbers associated with the curves denote total number of

animals.

(B) Eggs laid per virgin female after neuropeptide injection (sCRZ, control). Females were

mated with wild-type males for 12 h prior to assessment. Each virgin female BPH was injected

with a calibrated glass capillary needle directly into the abdomen with 100 ng of peptide

dissolved in 50 nL of 1 × PBS balanced salt solution. Data are shown as mean ± s.e.m. ns: $P >$

0.05 (Student's t-test). The numbers below the bars denote total number of animals, ≥4

biological replicates, ≥10 insects/replicate).

Figure 1 - figure supplement 4. Characterization of *Crz* mutants in *N. lugens*.

(A) Sequencing of *Crz* mutant (*ΔCrz1* and *ΔCrz2*) and wild type (WT) PCR products from BPHs.

The exact indel types of the G0 mutant individuals were confirmed by cloning and sequencing.

The wild-type sequences are shown at the top with the target site marked by in pink shading

and PAM in yellow shading. The change in the length of the mutant sequence is shown on the

right side of the sequence (+, insertion; -, deletion). (B) The Sanger sequence chromatograms

flanking the sgRNA target site for wild type (WT). (C and D) The Sanger sequence

chromatograms flanking the sgRNA target site for wild type (WT), heterozygous mutant

(*ΔCrz1/+* and *ΔCrz2/+*). (C' and D') The Sanger sequence chromatograms flanking the sgRNA

target site for wild type (WT), homozygous mutant (*ΔCrz1* and *ΔCrz2*).

(E) Developmental duration (egg to adult) across genotypes. ****P < 0.0001 (Student's *t*-test).

The numbers below the bars denote total number of animals. (F) Survival curves of adult *Crz*

mutants vs. WT. *P < 0.05 (Log-rank Mantel-Cox test; df = 1; n ≥ 50/group).

CRZ is produced by similar neurons in both sexes of BPHs and is

absent from the male accessory gland

To further understand how CRZ regulates the PMR in BPHs, we examined the expression of CRZ and *Crz* across different tissues in males and females. The *Crz* mRNA expression was examined in various BPH tissues using both RT-PCR and qPCR (Figure 2A). We detected *Crz* mRNA expression primarily in the central nervous system (CNS), regardless of the primer set or detection method used, while expression was nearly undetectable in other tissues, including reproductive organs of both sexes (Figure 2B and C).

A previous study reported CRZ in four male-specific interneurons within the abdominal ganglia of *D. melanogaster* [47]. To determine whether CRZ exhibits sexually dimorphic expression in BPHs, we performed immunolabeling using an anti-CRZ antiserum. This labeled four pairs of neurons in the protocerebrum of the brain (Figure 2D and E) and one pair of neurons in the subesophageal ganglion in both males and females (Figure 2D and E). Since CRZ immunolabeled neurons in brains of most, if not all, insects studied include three or more pairs of lateral neurosecretory cells [55,56], we suggest that CRZ signaling regulating the PMR may be hormonal (at least in part). Importantly, no CRZ-immunopositive neurons were detected in the abdominal ganglia of either sex (Figure 2D and E), and thus there is no evidence for sex-specific expression of CRZ in neurons in BPHs. This is in contrast to *D. melanogaster*, where we confirmed, as previously reported [45], the presence of CRZ-immunolabeled neurons specifically in the abdominal ganglia of males (Figure 2 - figure supplement

1A).

To exclude that CRZ acts as a seminal fluid peptide transferred to females during

copulation, similar to sex peptide (SP) in *D. melanogaster* and macc in BPHs [24], we

specifically examined CRZ expression in reproductive tissues of both sexes. However,

we observed no CRZ immunolabeling in the male reproductive organs (Figure 2F1 -

F3) or female reproductive organs (Figure 2G1 – G3). This aligns with our

transcriptional *Crz* data shown above (Figure 2B and C). Furthermore, *Crz* mRNA and

CRZ protein were absent from transcriptomic and proteomic datasets of the male

accessory glands (MAGs) in BPHs [24]. Collectively the available data suggest, that

CRZ is not produced by the MAG and is consequently unlikely to be transferred to

females as a seminal fluid component.

Figure 2. Expression of *Crz* and CRZ in the nervous system of the brown planthopper (BPH).

(A) Schematic of primer design for *Crz* detection (primer sequences provided in Table S1).

(B) RT-PCR analysis of *Crz* mRNA levels in various BPH tissues. Actin served as the loading

control. Tissues: Nervous system (NS), fat body (FB), Epidermis (EP), digestive system (DS),

female reproductive organs (FRO), male reproductive organs (MRO). Note: Tissues except

reproductive organs were pooled from both sexes.

(C) Relative *Crz* expression in female *N. lugens* tissues determined by RT-qPCR

(abbreviations as in B). Expression was normalized to *Actin* and *18SrRNA* and expressed

relative to the mean value of the highest-expressing tissue (set to 1). Data are shown as mean

± s.e.m. Groups that share at least one letter are statistically indistinguishable. One-way

ANOVA followed by Tukey's multiple comparisons test.

(D and E) Immunolocalization of CRZ peptide in the nervous system of male (D) and female (E)

BPHs. Four pairs of CRZ neurons in the brain and one pair in the subesophageal ganglion are

shown. Note that some of the CRZ neurons are likely to be neurosecretory cells with axon

terminations in the corpora cardiaca, others may be interneurons. The extensive branches

within the brain suggest CRZ signaling in brain circuits. VNC, ventral nerve cord. Scale bars:

50 µm. (E) CRZ immunolabeling in the nervous system of female BPHs. Scale bar: 50 µm.

(F and G) Lack of CRZ immunoreactivity in BPH reproductive organs. (F1-F3) Male

reproductive organs (MRO): testes (test). (G1-G3) Female reproductive organs (FRO): lateral

oviduct (Lat. ov), common oviduct (Com. ov). Scale bars: 100 µm.

Figure 2 - figure supplement 1. CRZ immunolabeling in abdominal ganglion and

reproductive organs of male and female BPHs and *D. melangaster*.

Anti-CRZ immunolabeling in the abdominal ganglia of the ventral nerve cord (VNC) in *N.*

lugens and *D. melanogaster* (*D. Melan.*). Scale bar: 50 µm. Note: Male-specific CRZ-positive

neurons are present in *D. melanogaster*, as previously reported.

The CRZ receptor is essential for mediating the post mating

response in female BPHs

Given our findings implicating CRZ in modulating the PMR in female BPHs, we

identified the CRZ receptor (*CrzR*) gene in this species; it encodes a structurally

conserved GPCR orthologous to other insect *CrzR*s (Figure 3-supplemental figure 1).

We next explored the effect of knocking down the *CrzR* on the PMR in BPHs.

RNAi-mediated knockdown of the receptor (*dsCrzR* injection) resulted in 15% of the

mated females re-mating, a phenotype not seen in *dsgfp*-injected controls (Figure 3A

and B) and in significantly reduced oviposition (Figure 3C). Importantly, CRZ injection

failed to diminish receptivity in *CrzR*-knockdown virgins (Figure 3D and E).

Furthermore, MAG extract (with seminal fluid proteins) injection into *dsCrzR*-treated

virgins did not affect receptivity (Figure 3F).

To further confirm the role of the *CrzR* in the PMR of BPHs, we applied

CRISPR/Cas9 genome editing to mutate the *CrzR* gene and constructed a

homozygous mutant strain of *CrzR* ($CrzR^M$) by genetic crosses (Figure 3G-3G'', Fig. 3

- figure supplement 2A). We found that the developmental time and life span of the

female *CrzR* mutants, like the *Crz* mutants, were slightly extended compared to wild

type animals (Figure 3 - supplement figure 2E and 2F). Homozygous $CrzR^M$ females

displayed increased remating rates (30%) (Figure 3I and J) and the number of eggs

laid by $CrzR^M$ females was also significantly decreased (Figure 3K). We also

examined mating and oviposition behaviors in *CrzR* heterozygous ($+/CrzR^M$) females.

Notably, both the remating rate and egg production in heterozygous females were

comparable to those observed in homozygous *CrzR* mutant (*CrzR^M*) females

(approximately 30%) (Figure 3I and J). Thus, heterozygosity for the *CrzR* was

sufficient to disrupt post-mating behaviors in female BPHs. These findings indicate

that the *CrzR* loss-of-function phenotype is dominant, and that a single functional

copy of *CrzR* is insufficient to maintain a normal post-mating response. Consistent

with the *CrzR* RNAi results, neither CRZ nor MAG extract injections reduced

receptivity in *CrzR^M* virgins (Figure 3K - M). Finally, macc peptide injection also failed

to decrease receptivity in *CrzR*-mutant virgins (70% acceptance rate), whereas it

significantly suppressed mating in wild-type controls (Figure 3N). These results

collectively demonstrate that intact CRZ/*CrzR* signaling in female BPHs is required for

transducing the PMR-inducing effects of male-derived seminal factors, including macc

[24].

Figure 3. The CRZ receptor (*CrzR*) is essential for the female PMR and is required for action

of MAG-derived factors.

(A) Experimental design for B and C.

(B) Receptivity after *CrzR*-RNAi (injection of ds*CrzR*; ds*GFP* as control). Percentage females

copulating within 30 min. The small circles denote the number of replicates, ≥9 insects/replicate; the numbers below the bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, $P <$ 0.05 was considered significant.

(C) Numbers of eggs laid within 72 h post-mating (*dsCrzR* females × WT males). Data are shown as mean ± s.e.m. ****$P < 0.0001$ (Student's *t*-test). The numbers below the bars denote total number of animals, ≥4 biological replicates, ≥7 insects/replicate.

(D) Experimental design for E-F.

(E) Receptivity after CRZ injection in *CrzR*-knockdown virgins (*dsGFP* + CRZ control). Graph shows percentage females copulating within 30 min. Each virgin female BPH was injected with a calibrated glass capillary needle directly into the abdomen with 10 ng of peptide dissolved in 50 nL of 1 × PBS balanced salt solution. The small circles denote the number of replicates, ≥7 insects/replicate; the numbers below the bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, $P <$ 0.05 was considered significant.

(F) Receptivity after MAG extract (with seminal fluid proteins) injection in *CrzR*-knockdown virgins (*dsGFP* + SFP control). Percentage copulating within 30 min. The small circles denote the number of replicates, ≥7 insects/replicate; the numbers below the bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, $P < 0.05$ was considered significant.

(G) *CrzR* targeting strategy for generation of mutants. *Top:* Genomic structure (ATG/TAG in Exons 2/7). sgRNA target (pink), PAM (yellow). *Bottom:* CRISPR alleles.

(G') Translated receptor mutant sequences.

(H) Experimental design for I-J.

(I) Receptivity of *CrzR* mutants compared to wild type animals (WT). Graph displays the percentage of females copulating within 30 min. The small circles denote the number of replicates, ≥9 insects/replicate; the numbers below the bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, $P < 0.05$ was considered significant.

(J) Number of eggs laid by mutants mated with WT males within 72 h. The numbers below the bars denote total number of animals, ≥4 biological replicates, ≥7 insects/replicate. Data are shown as mean ± s.e.m. ****$P < 0.0001$; Student's *t* test.

(K) Experimental design for L-M.

(L) Receptivity after CRZ injection in *CrzR* mutants compared to WT. Graph displays the percentage of females copulating within 30 min. Each virgin female BPH was injected with a calibrated glass capillary needle directly into the abdomen with 10 ng of peptide dissolved in 50 nL of 1 × PBS balanced salt solution. The small circles denote the number of replicates, ≥5 insects/replicate; the numbers below the bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, $P <$

0.05 was considered significant.

(M) Receptivity after SFP injection in *CrzR* mutants compared to WT. Graph displays the

percentage of females copulating within 30 min. The small circles denote the number of

replicates, ≥7 insects/replicate; the numbers below the bars denote total number of animals.

Statistical comparisons of proportions were performed using chi-square tests for contingency

tables, *P* < 0.05 was considered significant.

(N) Receptivity after macc injection in *CrzR* mutants compared to WT. Graph displays the

percentage of females copulating within 30 min. The small circles denote the number of

replicates, ≥10 insects/replicate; the numbers below the bars denote total number of animals.

Statistical comparisons of proportions were performed using chi-square tests for contingency

tables, *P* < 0.05 was considered significant.

Figure 3 - figure supplement 1. Structural conservation of the *CrzR* across insect species.

(A) Multiple sequence alignment highlighting conserved transmembrane domains (TM1-TM7,

shaded) in CRZ receptor orthologs. The species shown are: *N. lugens*: *Nilaparvata lugens; M.*

sexta: Manduca sexta; B. mori: *Bombyx mori*; *A. gamb*: *Anopheles gambiae*; *D. mel*: *D.*

melanogaster.

(B) Phylogenetic analysis of *CrzR* and *AkhR* based on the alignment of amino acid sequences

of insect species. Phylogenetic analysis was performed in MEGA 12 using the Maximum

Likelihood method with the LG amino-acid substitution model and Gamma-distributed rate

variation among sites (G, 5 discrete categories), with gaps treated by complete deletion.

Branch support was evaluated with 10,000 bootstrap replicates, and the numbers at the nodes

indicate bootstrap support values.

Figure 3 - figure supplement 2. Characterization of *CrzR* mutants in *N. lugens*.

(A) Sanger sequencing of *CrzR* mutants. WT sequence with target (pink), PAM (yellow). Indels:

+ (insertion), – (deletion).

(B) Sanger sequencing chromatograms of *CrzR* alleles: WT (wild-type).

(C) Sanger sequencing chromatograms of *CrzR* alleles: heterozygous (*CrzR^{M/+}*).

(C') Sanger sequencing chromatograms of *CrzR* alleles: homozygous (*CrzR ^M*) mutants.

(D) Developmental duration (egg to adult eclosion) in *CrzR* mutants compared to WT. Data:

mean ± SD. ****P < 0.0001 (Student's *t*-test; n ≥ 40/group).

(E) Survival curves of adult *CrzR* mutants compared to WT. ***P < 0.001 (Log-rank Mantel-Cox

test; df = 1; n ≥ 50/group).

The *CrzR* is highly expressed in the reproductive organs of female BPHs

To further elucidate the role of the CRZ signaling in regulation of the PMR, we

analyzed the *CrzR* expression in *N. lugens* tissues. We first monitored *CrzR*

expression by means of RT-PCR and qPCR. Both methods revealed predominant

CrzR transcript expression in the female reproductive tract (Figure 4A-C). qPCR

further demonstrated significantly higher *CrzR* expression in adult females than in

males (Figure 4 - figure supplement 1A), particularly localized to female abdomens

(Figure 4 - figure supplement 1B).

As we failed to generate a functional NlCrzR antibody, to determine the tissue

expression of the receptor in more detail, we used *in situ* hybridization and confirmed

strong *CrzR* expression within the female reproductive system (Figure 4D). No

hybridization signal was detected in the lateral or common oviducts (Figure 4D1).

However, specific labeling occurred in the sperm storage organs: the spermathecae

and pouched glands of the lower reproductive tract (Figure 4D2 and 4D2'). These

structures are the primary sites for long-term sperm storage. Specificity was validated

by absence of signal in sense-probe controls (Figure 4D3 - D4'). This enrichment in

sperm storage organs suggests that CRZ signaling may modulate post-mating

physiology in part by regulating the mobilization of stored sperm and seminal fluid.

Notably, while CrzR is highly expressed in the female reproductive tract, we

acknowledge that RT-PCR and qPCR analyses also detected low but detectable CrzR

expression in other tissues, including the CNS and fat body (Figure 4B-C). Thus, we

cannot exclude the possibility that *CrzR* in the brain (e.g., by modulating neural

circuits governing reproductive behavior) or fat body (e.g., by affecting post-mating

metabolism and nutrient/energy allocation) may also contribute to the PMR. These

non-reproductive tract sites of CrzR expression warrant further investigation to fully

delineate the systemic roles of CRZ/CrzR signaling in female PMR regulation.

Figure 4. Spatial expression profiling of *CrzR* in the female *N. lugens* reproductive tract.

(A) *CrzR* genomic structure with primer design for experiments in panels B-C.

(B) RT-PCR analysis of *CrzR* tissue distribution. Note weaker expression in central nervous system and fat body. Actin loading control is shown in top row. Tissue abbreviations: NS: central nervous system; FB: fat body; EP: Epidermis; DS: digestive system, FRO: female reproductive organs, MRO: male reproductive organs. Except for the reproductive system, the rest of the tissues come from a mixture of female and male.

(C) Relative *CrzR* expression in female *N. lugens* tissues determined by RT-qPCR (abbreviations as in B). Expression was normalized to *Actin* and *18SrRNA* and expressed relative to the mean value of the highest-expressing tissue (set to 1). Data are shown as mean ± s.e.m. Groups that share at least one letter are statistically indistinguishable. One-way

ANOVA followed by Tukey's multiple comparisons test.

(D1 - D2') *CrzR* mRNA localization via fluorescent *in situ* hybridization (antisense probe): D1:

There is no signal in lateral/common oviducts; D2/D2': Specific signal is detected in

spermatheca (SP) and pouched gland (PG).

(D3 - D4') Negative controls (sense probe) showing background-level signal. Scale bars: 100

µm.

Figure 4 - figure supplement 1. Developmental and tissue-specific expression profiling of

CrzR in *N. lugens* determined by qPCR.

(A) *CrzR* transcript levels across developmental stages. Relative expression levels were

quantified by RT-qPCR, normalized to the reference gene *Actin*, expressed relative to the

mean value of the highest-expressing tissue (set to 1). Data are shown as mean ± SEM shown.

Groups that share at least one letter are statistically indistinguishable. One-way ANOVA

followed by Tukey's multiple comparisons test.

(B) *CrzR* expression in distinct body regions. Relative expression levels were quantified by

RT-qPCR, normalized to the reference gene *Actin*, expressed relative to the mean value of the

highest-expressing tissue (set to 1). Data are shown as mean ± SEM shown. Groups that

share at least one letter are statistically indistinguishable. One-way ANOVA followed by

Tukey's multiple comparisons test.

Endogenous CRZ signaling modulates aspects of the PMR in female *Drosophila melanogaster*

Having established the critical role of CRZ signaling in regulating the PMR of the female BPH, we next investigated its potential function in female reproduction, including the PMR, of the genetically tractable fly *D. melanogaster*.

We first assessed re-mating frequency in *Crz* mutant female flies (*Crz^{attp}*). Approximately 40% of the *Crz^{attp}* females that had experienced a first successful copulation mated again, compared to only 3% of mated wild-type control females (Figure 5A and B). Importantly, the *Crz* mutation did not affect the first mating of females (Figure 5B). Furthermore, mated *Crz^{attp}* females laid approximately 45% fewer eggs than mated wild-type females (Figure 5C). These results demonstrate that CRZ signaling is necessary for aspects of the PMR in female *D. melanogaster*, specifically inhibiting re-mating and stimulating post-mating egg production. Immunohistochemical labeling confirmed the absence of CRZ protein in *Crz^{attp}* mutants (Figure 5D).

To determine whether release of CRZ from *Crz* neurons is responsible for inhibition of remating and stimulation of oviposition, we knocked down *Crz* expression specifically in CRZ-producing neurons using the GAL4-UAS system (*Crz-GAL4 > UAS-Crz-RNAi*). Consistent with the mutant phenotype, 22% of the mated *Crz-GAL4 > UAS-Crz-RNAi* females mated again after their first successful mating with wild-type males, whereas only 2% of the control females (*Crz-GAL4/+*) did so (Figure 5E). Virgin receptivity was again unaffected by *Crz* knockdown (Figure 5E). Additionally,

mated *Crz-GAL4 > UAS-Crz-RNAi* females laid approximately 20% fewer eggs than

mated controls (*Crz-GAL4/+*) (Figure 5F). To further confirm the role of CRZ

neuropeptide in modulation of the PMR in females, we injected this peptide in virgin

females. Consistent with the above results, CRZ injected virgins displayed a reduced

response to mating attempts (Figure 5G). To determine whether CRZ modulates the

PMR endogenously also in female *D. melanogaster*, we first examined whether the

male accessory gland (MAG) expresses *Crz*. RT-PCR analysis revealed no

detectable *Crz* transcripts in the *D. melanogaster* MAG (Figure 5H), indicating that the

MAG does not synthesize the CRZ peptide. To further exclude the possibility that CRZ

or any other *Crz*-precursor-derived peptide might enter the AG from other sources, we

used the GAL4–UAS system to visualize *Crz*/CRZ-producing neurons and performed

immunolabeling with anti-CRZ antibodies. Neither approach revealed any signal in the

MAG (Figure 5I). Some GFP signals were observed in cells of the testes (indicated by

white   arrows),   whereas   no   corresponding   signal   was   detected   by   CRZ

immunohistochemistry (Figure 5I).

To further evaluate whether the MAG could be a source of CRZ, we interrogated

publicly available single-cell transcriptomic datasets from the Fly Cell Atlas covering

the male reproductive glands (including MAG main and secondary cells, ejaculatory

duct/bulb epithelium, secretory and muscle cells) and the male testis (Figure 5 - figure

supplement 1). In the MAG atlas, *Crz* transcripts were undetectable across MAG main

and secondary cells as well as other gland-associated cell types (Figure 5 - figure

supplement 1A). In contrast, the MAG-derived sex peptide (SP) was robustly detected,

confirming correct tissue representation and cell-type annotation (Figure 5 - figure

supplement 1A). In the testis atlas, Crz expression was sparse and restricted to a

small subset of cells rather than enriched in any major spermatogenic lineage (Figure

5 - figure supplement 1B and C), whereas β2-tubulin prominently marked

spermatogenic populations (Figure 5 - figure supplement 1B). Hence, these

single-cell data support our RT-PCR and anti-CRZ immunolabeling findings. Together,

our data demonstrate that the male accessory gland does not produce or sequester

CRZ. In conclusion, like in BPHs, CRZ acts endogenously in females to modulate

post-mating behavior also in *D. melanogaster*.

.

Figure 5. CRZ regulates part of the PMR in *D. melanogaster*.

(A) Schematic of experimental design. White segments on the time axis indicate light periods; black segments indicate dark periods.

(B) Receptivity of virgin and mated *Crz* mutant (*Crz^attp^*) females. Graph displays the percentage of females copulating within 30 min. Small circles indicate the number of replicates, ≥6 insects/replicate; numbers below bars denote total number of animals. Statistical comparisons of proportions were performed using chi-square tests for contingency tables, *P* < 0.05 was considered significant.

(C) Graph shows eggs laid per female after mating. Numbers below bars denote total animals, ≥4 biological replicates, ≥8 insects/replicate. ****P < 0.0001 (Student's t-test).

(D) Anti-CRZ immunostaining in brains of *Crz* mutant (*Crz^attp^*), control (Cs) and after *Crz*-RNAi. Note CRZ-expressing cells at arrows in control. Scale bar: 100 μm.

(E) Receptivity of virgin and mated females after *Crz*-RNAi. Graph displays the percentage of females copulating within 30 min. Small circles indicate the number of replicates, ≥8

insects/replicate; numbers below bars denote total animals. Statistical comparisons of

proportions were performed using chi-square tests for contingency tables, $P < 0.05$ was

considered significant.

(F) Graph shows eggs laid per female after mating. Numbers below bars denote total animals,

≥4 biological replicates, ≥8 insects/replicate. ****$P < 0.0001$ (Student's *t*-test).

(G) Receptivity of virgin females following CRZ peptide injection (monitored 6 h after injection).

Graph displays the percentage of females copulating within 30 min. Small circles indicate the

number of replicates, ≥9 insects/replicate; numbers below bars denote total animals. Statistical

comparisons of proportions were performed using chi-square tests for contingency tables,$P <$

0.05 was considered significant.

(H) RT-PCR analysis of Crz tissue distribution. Note the lack of expression in the accessory

gland. Actin loading control is shown in the top row. Tissue abbreviations: MAG, male

accessory gland; Male–MAG, whole body without male accessory gland.

(I) Anti-GFP and anti-CRZ immunostaining in brains of *w⁻/y; Crz-Gal4/UAS-mCD8-GFP*. Note

no signal was detected in male accessory gland (MAG). Some GFP-stained signals were

observed in the testes (indicated by white arrows), whereas no corresponding signals were

detected via CRZ antibody staining. Scale bar: 50 μm.

A

B

C

Figure 5 - figure supplement 1. Crz expression is undetectable in the *Drosophila* male

accessory gland in Fly Cell Atlas single-cell datasets.

(A) Fly Cell Atlas male reproductive glands single-cell dataset. Left, schematic of male

reproductive glands. Middle, two-dimensional embedding colored by annotated cell types,

including male accessory gland (MAG) main and secondary cells. Right, feature plot showing

expression of Crz (green scale) together with the AG marker sex peptide (SP) (red scale). Crz

transcripts are not detected in MAG cell populations, whereas SP is robustly expressed.

(B) Fly Cell Atlas male testis single-cell dataset. Left, schematic of testis. Middle, embedding

colored by annotated testis cell types. Right, feature plot showing Crz (green), SP (red)   and

β2-tubulin (blue) expression, with β2-tubulin marking spermatogenic populations and Crz

detected only in a small subset of cells.

(C) Strip/dot plot summarizing normalized Crz expression across selected testis cell clusters,

illustrating sparse Crz expression without broad enrichment in a major lineage. The y-axis

signifies the expression level.

Discussion

Our study unveils a novel function of CRZ in an endogenous female signaling

pathway that modulates the post-mating reproductive behavior and physiology

(collectively PMR) in BPHs. It is, thus, the first report linking CRZ to a PMR in insects.

In contrast to SP and DUP99B in *D. melanogaster* [5,18], HP-1 in *Aedes aegypti* [15] and

maccessin (macc) recently identified in BPHs [24], CRZ is not produced in the MAG,

and thus not transferred from males during copulation. Instead, it is acting as an

endogenous peptide derived from neuroendocrine cells in the female brain. Moreover,

this CRZ-mediated endogenous modulation in females is seen also in *D.*

melanogaster. When *Crz* is knocked down or knocked out in flies, mated females

likewise exhibit an altered PMR.

Focusing on the endogenous CRZ signaling in female BPHs, our main findings

using mutants and RNAi of both *Crz* and *CrzR*, demonstrate that CRZ signaling

decreases the female responsiveness to courting males, especially in already mated

females. Also, oviposition in mated BPHs is increased by CRZ signaling. CRZ

immunolabeling revealed a small set of neurons in the brain and subesophageal

ganglion and no CRZ was detected in reproductive organs. Thus, the likely source of

CRZ affecting the PMR is the CRZ-producing neuroendocrine cells in the brain. In

insects studied so far (including other hemipterans), there are at least three pairs of

CRZ expressing lateral neurosecretory cells with axon terminations in the corpora

cardiaca [55,56], suggesting that in BPHs and *D. melanogaster* hormonal CRZ may

contribute to the modulation of the PMR. However local CRZ signaling in brain circuits

cannot be excluded since the CRZ neurons arborize extensively within the brain.

We found that the CRZ receptor, *CrzR*, is predominantly expressed in the female

reproductive tract in BPHs, more specifically in the lower tract including the

spermatheca and pouched gland, which are sites of sperm and seminal fluid storage.

Thus, in BPHs, CRZ may act to regulate the release of stored sperm and seminal fluid

in the female. This regulation would also gate the flow of male-derived factors that

induce and maintain the PMR, such as macc and ITPL-1. As mentioned, we cannot

exclude additional action of CRZ in neuronal circuits in the CNS that regulate female

sexual arousal and reproductive behavior, since our PCR data indicate expression of

the *CrzR* also in the CNS. Additionally, the fat body of BPHs, express *CrzR*, similar to

D. melanogaster [57], possibly suggesting a role of CRZ in post-mating metabolism.

The brain CRZ neurons of BPHs are likely to be activated by sensory signals from

the female reproductive tract upon copulation and hence release CRZ into the

circulation (and maybe within the brain). In *D. melanogaster,* sensory neurons in the

reproductive tract are activated by SP (via the SP receptor), and signal via a chain of

ascending axons of the VNC to sex-specific neurons in the brain to induce the PMR

switch in behavior and physiology [7,38,40,58]. Thus, in *D. melanogaster,* these

sex-specific circuits may also activate the CRZ neurons. Possibly similar sensory and

sex-specific neurons exist in BPHs and respond either to mechanical stimuli or factors

in the seminal fluid leading to activation of CRZ neurons and other brain neurons that

regulate the PMR.

The female PMR is complex in *D. melanogaster* and not only receptivity and

oviposition are affected, but also feeding, metabolism, sleep patterns and aggression

[5,16,18,19,59,60]. In BPHs only the receptivity (and associated rejection behavior) and

oviposition have been analyzed in any detail. We found that the MAG-derived

peptides transferred with the semen, macc and ITPL-1, only affected the receptivity [24],

whereas we show here that CRZ affects also oviposition. Thus, we have identified

three different peptides CRZ, macc and ITPL-1 that all influence the PMR in different

ways, suggesting complex regulatory mechanisms also in BPHs.

It is interesting to note that CRZ is an evolutionarily conserved neuropeptide that

has been found in most, but not all, insect species and it displays a wide array of

functions [56,57,61,62]. The CRZ function varies across different insect species, but a

common role seems to be in regulation of stress [62-64]. Other functions are for example:

in the American cockroach, CRZ accelerates the heart rate [65], in locusts, it triggers the

formation of pigments that darken the body color in gregarious insects [66], in the moth

Bombyx mori, CRZ affects developmental speed and reduces the rate of silk spinning

behavior [67]; in *D. melanogaster* larvae, CRZ regulates molting motor behavior [68] and

growth [69]; in adult *D. melanogaster* CRZ regulates stress responses, food intake and

metabolism [57,70], and protein food preference [12], as well as sperm ejection and mating

duration in males [45]; and in social ants, it regulates the social hierarchy and inhibits

egg-laying in the queen [48]. Finally, a recent study found that CRZ mediates a

diapause-induced suppression of reproduction in female bean bugs [71]. To all these

functions can now be added the role of CRZ in reproductive behavior and physiology

of female BPHs and *D. melanogaster*. Finally, building on the role of CRZ in

metabolism shown in *D. melanogaster* [57], it is possible that CRZ also regulates

post-mating metabolism in female BPHs and flies. Thus, there may be a link between

CRZ and the altered feeding and metabolism commonly seen in mated female insects

that ensures optimal egg development[5,9,16].

CRZ and the vertebrate peptide gonadotropin-releasing hormone (GnRH) are

paralogs that arose via an ancient gene duplication, and act on evolutionarily

conserved receptors[72,73]. *Bona fide* GnRHs have not been found in arthropods and the

CRZ peptide has been lost in the vertebrate lineage, but the cephalochordate

(lancelet) *Branchiostoma floridae* does have peptides of both types[73]. In vertebrates,

including humans, GnRH regulates female reproduction (specifically sexual

maturation)[74,75]. We have shown here that CRZ regulates reproductive behavior and

physiology in females of *D. melanogaster* and previous reports demonstrated a role in

female fecundity[76] and a reproductive function also in males of this species[45]. Thus,

CRZ and GnRH share a major function that may be ancient, although CRZ signaling

in insects has evolved to also regulate stress responses, metabolism and

development[57,62,77].

Some outstanding questions remain with respect to mechanisms of CRZ

signaling in the female PMR. The inputs triggering release of hormonal CRZ at mating

are not known in BPHs or flies. This can probably be best investigated in *Drosophila*

where brain neurons that are targets of the sex peptide receptor expressing neurons

in the oviduct have been identified[58], and synaptic inputs to CRZ producing brain

neurons have been described[78]. Once candidate interneurons interacting with the

CRZ neurons have been identified, genetic manipulations of these neurons may

unveil how mating triggers endogenous CRZ signaling and the PMR. Another aspect

of CRZ signaling that merits further study in *Drosophila* is the link between CRZ and

the post-mating change in feeding and metabolism. At present it is not known whether

also BPHs change their feeding and metabolism after mating, but would be valuable

to investigate to determine to what extent the PMR in the two insects are similar.

Finally, since BPHs are a major rice pest, it may be useful to further investigate the

regulation of the PMR to design novel strategies for control their reproduction.

Table legend:

733   **Table 1. Primers used in this study**

Primer name	Sequence (5'-3')	Purpose
NlCrz-F	ACCGCATCAGGACGACAATA	Gene clone
NlCrz-R	ACCATAGCTGTTGAGCTACAAAT	
NlCrz-RNAi-F	TAATACGACTCACTATAGGGGCTAG TGTTGCAATGCTGTTG	
NlCrz-RNAi-R	TAATACGACTCACTATAGGGGCTGTT GAGCGTTAGACTGT	*dsNlCrz* synthesis
QNlCrz-F	AGACAGTGACCACAGCTCAA	
QNlCrz-R	GAGTACTGGAACGTCTGTGC	qRT-PCR for NlCrz
NlCrz-sgRNA-F1	TAATACGACTCACTATAGGCACAGG AGTGATGCAGAC	
NlCrz-sgRNA-R1	TTCTAGCTCTAAAACGTCTGCATCAC TCCTGTGCC	sgRNA synthesis for NlCrz knock-out
NlCrz-sgRNA-F2	TAATACGACTCACTATAGCTACTAGT GCTGGGCTGCA	
NlCrz-sgRNA-R2	TTCTAGCTCTAAAACTGCAGCCCAG CACTAGTAGC	sgRNA synthesis for NlCrz knock-out
NlCrz Check F	ACCATAGCTGTTGAGCTACAAAT	
NlCrz Check R	TGCAATGCTGTTGTCGCGAC	To analyze NlCrz mutant
NlCrzR-F	AATCAGGCGATGAACCTCAGTT	
NlCrzR-R	ACGCAGTTCTATGACGTGAGG	Gene clone
NlCrzR-RNAi-F	TAATACGACTCACTATAGGGCGCCG TCTACACACTCATCT	
NlCrzR-RNAi-R	TAATACGACTCACTATAGGGCTACCA GCTTCGTACAGCGT	*dsNlCrzR* synthesis
QNlCrzR-F1	CTGGTGCTAATAGGGCTGGA	
QNlCrzR-R1	TCCGGTGTACAGTTGCTCTT	qRT-PCR for NlCrzR
QNlCrzR-F2	TGCTCACCATCGCCTCTATC	
QNlCrzR-R2	TGCCATGGTTCCGTGTAGAA	qRT-PCR for NlCrzR
NlCrzR-F1	CCCACTAGCCAGCAACCATT	
NlCrzR-R1	TCCTCCACAAATGGTCCACG	RT-PCR for NlCrzR
NlCrzR-sgRNA-F1	TAATACGACTCACTATAGGCACGCA CTCGAGCTGCCT	
NlCrzR-sgRNA-R1	TTCTAGCTCTAAAACAGGCAGCTCG AGTGCGTGCC	sgRNA synthesis for NlCrzR knock-out
NlCrzR-sgRNA-F2	TAATACGACTCACTATAGGCGATGG TGAGCAGGTTGC	
NlCrzR-sgRNA-R2	TTCTAGCTCTAAAACGCAACCTGCTC ACCATCGCC	sgRNA synthesis for NlCrzR knock-out
NlCrzR Check-F	GGCAGCAACCTGAGCCACCC	To analyze NlCrzR mutant

NlCrzR Check-R	GTCTTACCAGCGACCCAAGC	
QNlvg-F	GTGGCTCGTTCAAGGTTATGG	
QNlvg-R	GCAATCTCTGGGTGCTGTTG	qRT-PCR for Nlvitellogenin
QNlvgR-F	AGGCAGCCACACAGATAACCGC	
QNlvgR-R	AGCCGCTCGCTCCAGAACATT	qRT-PCR for Nlvitellogenin
QNl53903-R	GTCCACCGAACTTTTCCTGG	receptor
QNl Macc-F	TCTTTGGAAGTGGTGATTCCATCT	qRT-PCR for NlMacc
QNl macc-R	TGTCTACTTCACATACATGTGGCT	
QNl itpl-1-F	GATCAGATTCAATTCACATTCCTC	qRT-PCR for Nlitpl-1
QNl itpl-1-R	AGATTGTAGCAGTCCTCGCATA	

736 <div align="center">Key resources table</div>

REAGENT or RESOURCE	Source or reference	Identifiers
Antibodies		
Goat anti-Rabbit Alexa 488	Invitrogen	Cat#A11008; RRID: AB_143165
Goat anti-mouse Alexa 488	Invitrogen	Cat#A28175; RRID: AB_2536161
Goat anti-mouse Alexa 555	Invitrogen	Cat# A21428; RRID: AB_2535849
Anti-CRZ	Gift from Prof. Jan A. Veenstra	
Chemicals, Peptides, and Recombinant Proteins		
4% PFA	Leagene	Cat# DF0134
DAPI	Solarbio	Cat# C0060
PBS	Solarbio	Cat# P1022
ReadyProbes 2.5% Normal Goat Serum	Invitrogen	Cat# R37624
Experimental Models: Organisms/Strains		
Canton-S	Bloomington Drosophila Stock Center	64349
Crz^{attp}	Rao and Deng et al., 2019	BDSC: 84487
Crz-GAL4	Bloomington Drosophila Stock Center	51976
Crz-RNAi	Tsinghua Fly Center	THU2139
Software and Algorithms		
FIJI		https://doi.org/10.1038/nmeth.2019
Prism 9	GraphPad	RRID: SCR_002798
Photoshop	Adobe	RRID: SCR_014199
Illustrator	Adobe	RRID: SCR_010279
ZEN	Zeiss	N/A
Motic Images Plus3.0	Motic	N/A
Motic Images Devices	Motic	N/A
Other		
Egg-laying device	Gou et al., 2016	N/A
Anatomical tweezers	Dumont	Cat# 0209-5-PO

Method details

Experiment insects

All the brown planthoppers used in this research were collected from Hangzhou, Zhejiang Province in 1995. The insects were kept indoors on fresh rice seedlings (Taichung Native 1, TN1) in a growth chamber at 27 ± 1 °C and 70 ± 10% relative humidity, under a 16 h:8 h (Light: Dark) photoperiod [79].

All *D. melanogaster* stocks were raised in standard cornmeal–molasses–agar medium maintained at 25° C, 60% humidity, and under 12 hr:12 hr light:dark conditions. We used Canton s flies as the wild-type strain. The relevant information about fruit flies used in the experiments are shown in key resources table.

Male accessory gland protein extraction of *Nilaparvata lugens*

Protein extraction was performed as described previously[24]. Male accessory glands (MAGs) were dissected from 3–5-day-old virgin males in phosphate buffer (pH 7.2) under a stereomicroscope. For protein extraction, approximately 300 MAGs were collected, immediately frozen, and stored at −80 °C until use. Frozen MAGs were pre-ground in liquid nitrogen and then lysed in 500 μL protein cracking buffer (100 mM NH_4HCO_3, 8 M urea, 0.2% SDS, pH 8.0) with thorough homogenization. Lysates were centrifuged at 12,000 × g for 15 min at 4 °C, and the supernatants were collected as soluble protein extracts. Protein concentration was determined using a Bradford colorimetric assay (BSA as the standard; absorbance at 595 nm). Protein integrity and quality were assessed by SDS–PAGE. Protein extracts were stored at −80°C until next step experiment.

Behavior assays of *Nilaparvata lugens*

Mating and receptivity

All BPHs were raised in incubators with a 16 h:8 h dark:light cycle. Unmated female and male BPHs were collected within 12 hours of emergence. For RNAi or neuropeptide injection treatment, the protocol is shown in Figure 1A and Figure 1D. The mating behavior (or female receptivity behavior) test in this study usually conducted the first mating between 3 and 5 days after emergence. The mating experiment was completed in

a plastic tube (2.5cm in diameter and 10cm in height). Fresh rice seedlings of appropriate length were placed in each tube, and then a female and a male BPH was added. The mating conditions and other reproductive behavior parameters during this period were observed in a constant temperature room at 25°C for 30min, and the males were taken out after 30 min. In *N. lugens*, mating induces a transient suppression of female receptivity that is not permanent; females typically start to regain remating willingness 72 h after the first mating, as documented in our previous study[24]. This temporal window guided the design of our remating assays, in which females were paired with naive males at 48 h, post-initial mating to capture both the suppressed and recovered phases of receptivity. For the experiment requiring a second mating, females successfully mated for the first time were retained. After 48 hours, wild-type males were put in for the re-mating test. The males in all secondary mating experiments were wild type. The monitoring time for each mating behavior was 30 minutes; the receptivity rate refers to the proportion of successful matings of female individuals within the 30-minute measurement period.

Egg laying

After completing the first mating, the male is sucked out and a sufficient number of rice seedlings are added to the tube, and the female is retained to lay eggs in the tube. After 72 hours, the spawning female is transferred out of the tube. After the eggs have matured, about 4-5 days later, we used tweezers to gently manipulate the rice seedlings and counted the number of fertilized eggs produced by the females. Note that 4 to 5 days after the egg is produced, the successfully fertilized egg will display red eye-spots. Oviposition assay protocol: For quantitative assessment of post-mating responses, egg-laying activity was monitored over a 72-hour observation window following verified initial copulation. To eliminate confounding effects from sequential mating trials, experimental subjects undergoing oviposition measurement were permanently excluded from subsequent re-mating evaluations.

To assess oviposition in females injected with CRZ peptide, virgin or mated female BPHs (3–5 days post-emergence) were transferred into plastic tubes containing sufficient rice seedlings 6 h after recovery from peptide injection. Females were allowed to oviposit for

24 h and were then removed. Eggs were subsequently counted by gently dissecting the

rice seedlings with forceps.

Behavior assays of *D. melanogaster*

Mating and re-mating：During the receptivity assessment, all experimental fruit flies

were virgin flies that emerged within 12 hours in a 20°C incubator in a 12 h:12 h dark:light

cycle. After collection, they were kept at 25°C incubator in a 12 h:12 h dark:light cycle for

4 days before behavioral testing. Receptivity behavior measurement：a perforated plate

was placed on top of a non-perforated plate. Male fruit flies were briefly anesthetized on

ice and gently transferred into the holes of the perforated plate, with one male per hole.

Once the male transfer was completed, a plastic film was gently placed over the

perforated plate, followed by covering it with another perforated plate. Then, the virgin

female fruit fly, also anesthetized on ice, was placed in each hole. After the transfer was

completed, a non-perforated plate was placed on top. This setup was then transferred to a

25°C incubator to acclimate for 30 minutes. The plastic film separating the male and

female flies was quickly removed, and video recording began for 30 minutes. The

recorded videos were analyzed to calculate the acceptance rate of the females.

Successful mating females were collected and individually transferred into plastic tubes

containing sufficient artificial food for fruit flies to lay eggs. After 24 hours of laying eggs,

the female flies were transferred to a new finger-shaped tube with ample artificial feed for

another 24 hours of laying. After 48 hours, these egg-laying females were subjected to a

secondary mating assessment with wild-type, unmated males, using the same method as

described above, to evaluate the acceptance rate during the second mating process.

Additionally, the number of eggs laid by each female during the 48 hours was recorded.

Egg laying for mated females：First, collect virgin female fruit flies that eclosion within 12

hours at 20°C incubator in a 12 h:12 h dark:light cycle and then keep them in a 25°C

incubator in a 12 h:12 h dark:light cycle for 4 days. After that, unmated males of the wild

type and virgin females were mixed 1:1.5 times. 24 hours after full mating, female flies

were lightly anesthetized with ice and transferred to plastic vials containing enough

artificial fruit fly food for them to lay eggs for 24 hours. After 24 hours, the flies were

transferred again to a new bottle full of artificial food for another 24 hours of egg laying.

The number of eggs in each vial is calculated under a microscope, and the total number of

eggs in the two vials of each female fruit fly represents the egg-laying capacity of that

individual.

Developmental duration statistics

Nymphs of BPHs that hatched synchronously from distinct genotypes were collected and

reared in plastic cups containing fresh rice seedlings; the seedlings were replaced

regularly to maintain optimal conditions. From the day the nymphs reached the fifth instar,

cups were inspected daily for adult emergence, and the number of newly emerged adults

was recorded. After all surviving individuals had completed eclosion, the total

nymph-to-adult developmental duration was calculated.

BPHs survival rate statistics

For neuropeptide injection: After 4-day emergence, virgin BPHs were injected with CRZ

neuropeptide or scrambled CRZ, and after 6 hours of recovery, these insects were placed

into a plastic tube with sufficient quantity of rice seedlings. Survival was monitored every

24 hours until all specimens were dead.

For mutant: Wild-type and mutant BPHs adults that eclosed on the same day were

individually confined in plastic tubes, each supplied with ample fresh rice seedlings and

water renewed regularly to ensure optimal living and nutritional conditions. The number of

surviving individuals of each genotype was recorded every 24 h until all had died, and

survival curves were generated from these data.

Neuropeptide synthesis and injection.

Corazonin (CRZ) and scrambled CRZ (sCRZ) were synthesized by Genscript Co. Ltd

(Nanjing, China). Peptide masses were confirmed by Mass spectrometry and the amount

of peptide was quantified by amino acid analysis. The amino acid sequences of the

peptides used in this study are: *N. lugens* corazonin: pQTFQYSRGWNamide; scrambled

corazonin: pQSRYTTFGQWTNamide. On the fourth day after adult emergence, virgin

female BPHs were injected with different concentrations of synthetic CRZ peptide and

control scrambled peptide. The scrambled peptide control was used to monitor for

potential non-specific effects on behavior due to the process of injection. Each mg peptide

was dissolved in 100 μl DMSO solution (Acmec, #D54264), and then diluted to 1.5 mM,

0.15 mM, 0.015 mM and 0.0015 mM with 1×PBS as solvent. The BPHs were anesthetized

with CO_2 for about 30 seconds and placed abdomen up on 2% agarose (Solarbia, #A8201)

plate and 50 nl neuropeptide solution injected into each female. Six hours after

neuropeptide injection the insects were used for behavior assay.

Gene cloning and sequence analysis

We used the NCBI database with BLAST programs to carry out sequence alignment and

analysis. Then we predicted Open Reading Frames (ORFs) with EditSeq. The primers

were designed by Primer designing tool in NCBI. Total RNA Extraction was using the

TRIzol reagent (Invitrogen, Carlsbad, CA, USA) according to the manufacturer's

instructions. The cDNA template used for cloning was synthesized using the Biotech

M-MLV reverse transcription kit and the synthesized cDNA template was stored at -20˚C.

The transmembrane segments and topology of proteins were predicted by TMHMM v2.0

(http://www.cbs.dtu.dk/services/TMHMM-2.0/) [80]. Multiple alignments of the complete

amino acid sequences were performed with Clustal Omega

(http://www.ebi.ac.uk/Tools/msa/clustalo).

qRT-PCR for spatial expression pattern and RNAi efficiency

The nervous system (NS), fat body (Fb), digestive system (DS), epidermis (EP), male

reproductive organs (MRO) and female reproductive organs (FRO) from 3-5 days old

adults were dissected in 1×PBS, to determine the tissue-specific expression pattern. With

the exception of the reproductive system, the rest of the tissue comes from both sexes

equally. The adult BPHs were placed on ice under slight anesthesia and dissected under

a stereoscopic microscope (ZEISS). At least 50 individuals were dissected as one sample

for tissue-specific analysis with three replicates. All tissues were quickly transferred to

Trizol (Invitrogen, Carlsbad, CA, USA) solution at 0° C after dissection.

For RNAi efficiency testing, we collected the whole bodies of BPHs and transferred to a

1.5 mL EP tube containing Trizol reagent. Total RNA extraction was using Trizol reagent

(Invitrogen, Carlsbad, CA, USA) according to the manufacturer's instructions. The cDNA

template used for cloning was synthesized using the Biotech M-MLV reverse transcription

kit and the synthesized cDNA template was stored at -20°C.

Real-time qPCRs of the various samples used the UltraSYBR Mixture (with ROX) Kit

(CWBIO, Beijing, China). The PCR was performed in 20μl reaction including 2 μl of

10-fold diluted cDNA, 1 μl of each primer (10 μM), 10μl 2 × UltraSYBR Mixture, and 6 μl

RNase-free water. The PCR conditions used were as follows: initial incubation at 95°C for

10 min, followed by 40 cycles of 95°C for 10 s and 60°C for 45 s. *N. lugens* 18S rRNA and

actin were used as an internal control (primers listed in Table1). Relative quantification

was performed via the comparative $2^{-\Delta\Delta CT}$ method [81].

RNAi and RNAi efficiency determination

For lab-synthesized dsRNA, *gfp*, Nl*CrzR* and Nl*Crz* were amplified by PCR using specific

primers conjugated with the T7 RNA polymerase promoter (primers listed in Table1).

dsRNA was synthesized by the MEGAscript T7 transcription kit (Ambion, Austin, TX, USA)

according to the manufacturer's instructions. Finally, the quality and size of the dsRNA

products were verified by 1% agarose gel electrophoresis and the Nanodrop 1000

spectrophotometer and kept at -70°C until use.

3-5 days old virgin female BPHs were used for injection of 50nl of 5 μg/μl dsRNA per

insect. Injection of an equal volume of ds*gfp* was used as negative control. After 48 hours,

the insects were used behavior assays and gene relative expression analysis.

Neuropeptide injection after RNAi

48 hours after the ds*CrzR* injection, each insect was injected again with 10 ng of peptide

dissolved in 50 nL of 1 × PBS balanced salt solution. Recovery was for six hours after

injection. These insects were used for behavior assays. RNAi efficiency was examined by

qPCR using a pool of ten individuals 48 hours after dsRNA injections.

In vitro synthesis of sgRNA and Cas9 mRNA

The single guide RNA (sgRNA) was designed as previously reported [82]. Briefly, sgRNA was designed by manually searching genomic sequence around the region of the *Crz* and *CrzR* exon for the sequences corresponding to 5'-N_{17-20}NGG-3', where NGG is the protospacer-adjacent motif (PAM) of SpCas9 and N is any nucleotide. For in vitro transcription of sgRNA, were synthesized in vitro using the GeneArt™Precision gRNA Synthesis Kit (Thermo Fisher Scientific, Vilnius, Lithuania) according to the instructions of manufacturer. The Cas9 protein (TrueCut™Cas9 Protein v2, Cat. NO. A36497) was purchased from Thermo Fisher Scientific (Shanghai, China).

Embryo microinjection, crossing and genotyping

The embryonic injection was performed as previously reported [83]. The female BPHs, which had been mated for 3-6 days after emergence, were selected to lay eggs. After 30 minutes, the embryos were transferred onto double-sided tape attached to a microscope slide along the long edge by gently pressing the slide onto the dorsal surface of embryos. A few drops of halocarbon oil 700 was added to these eggs. About 0.5–1 nL mixture of Cas9 protein (200 ng/μL) and sgRNA (100 ng/μL) was injected into each egg using a FemtoJet and Inject Man NI 2 microinjection system (Eppendorf, Germany). After the injection, the eggs were placed in Petri dishes covered with moist filter paper, and the Petri dishes were placed in plant growth chamber at 27±1 °C, 70±10% RH for hatching. These eggs hatched into G0 generations.

Crossing and genotyping

To establish homozygous mutants of *Nilaparvata lugens* through CRISPR-mediated gene editing, newly emerged G0 adults were outcrossed with wild-type counterparts to generate G1 progeny. Following successful mating and oviposition, genomic DNA was extracted from G0 adults for mutation screening. The target region surrounding NlCrzR was amplified by PCR using specific primers NlCrzR-check-F and NlCrzR-check-R (primer sequences listed in Table 1), followed by Sanger sequencing to identify G0 individuals carrying mutations.

Progeny (G1) from mutation-positive G0 parents were maintained through sibling crosses

to establish G2 populations. Subsequent genotyping via Sanger sequencing was

performed on G1 individuals to select breeding pairs where both male and female parents

carried heterozygous mutations. The resulting G2 offspring from these validated pairs

were then subjected to intercrossing to ultimately generate homozygous mutant lines.

After homozygous mutant lines were established, virgin homozygous mutant females

were paired individually with virgin wild-type males. Following successful copulation,

inseminated females were transferred to fresh rice seedlings for oviposition, and fertilized

eggs were allowed to hatch. Hatched nymphs were continuously reared on fresh rice

seedlings until the fifth instar. Rice seedlings were replaced every 12 hours, and newly

emerged adults were collected and sex-separated for subsequent behavioral assays.

Quantitative RT-PCR

The first-strand cDNA was synthesized with HiScript II Q RT SuperMix for qPCR (+gDNA

wiper) kit (Vazyme, Nanjing, China) using an oligo (dT)18 primer and 500 ng total RNA

template in a 10 µl reaction, following the instructions. Real-time qPCRs in the various

samples used the UltraSYBR Mixture (with ROX) Kit (CWBIO, Beijing, China). The PCR

was performed in 20 µl reaction including 4 µl of 10-fold diluted cDNA, 1 µl of each primer

(10 µM), 10 µl 2 × UltraSYBR Mixture, and 6 µl RNase-free water. The PCR conditions

used were as follows: initial incubation at 95°C for 10 min, followed by 40 cycles of 95°C

for 10 s and 60°C for 45 s. *N. lugens* 18S rRNA or *D. melanogaster* rp49 were used as an

internal control (Table 1). Relative quantification was performed via the comparative

$2^{-\Delta\Delta CT}$ method (Livak, K. J,2001).

Immunohistochemistry

Adult BPHs, 3-5 days old, were dissected under 1x phosphate-buffered saline (PBS; pH

7.4) in Schneider's insect medium (S2). We dissected the brain, ventral nerve cord and

reproductive system of the BPHs. These tissues were fixed in 4% paraformaldehyde in

PBS for 30 min at room temperature. After extensive washing with PTX (0.5% Triton X100,

0.5% bovine serum albumin in PBS), blocked in 3% normal goat serum. Then, the tissues

were incubated in Anti - CRZ for 12 hours at 4℃ and in secondary antibody for 12 h at

4℃. Primary antibodies used were: rabbit anti - CRZ (1:1000, Anti - CRZ was gift from Jan

A. Veenstra), mouse anti-Bruchpilot (1:1000, Developmental Studies Hybridoma Bank

nc82). Secondary antibodies used: donkey anti-rabbit IgG conjugated to Alexa 488 (1:500,

R37118, Thermo Fisher Scientific) and donkey anti-mouse IgG conjugated to Alexa Alexa

555 (1:500, R37115, Thermo Fisher Scientific). The samples were mounted in

Vectorshield (Vector Laboratory). Images were acquired with Zeiss LSM 700 confocal

microscopes, and were processed with Image J software. All antibodies were diluted in

PTX solution. The primary antibody was diluted at a ratio of 1:1000, while the secondary

antibody was diluted at a ratio of 1:500.

Adult female *D. melanogaster*, 3-5 days old, were dissected under 1 x

phosphate-buffered saline (PBS; pH 7.4) in Schneider's insect medium (S2). We

dissected the brain, ventral nerve cord and reproductive system. These tissues were fixed

in 4% paraformaldehyde in PBS for 30 min at room temperature. After extensive washing

with PTX (0.5% Triton X100, 0.5% bovine serum albumin in PBS), blocked in 3% normal

goat serum. Then, the tissues were incubated in Anti - CRZ and Anti - nc82 for 12 hours at

4℃ and in secondary antibody for 12 h at 4℃. Primary antibodies used were: rabbit

anti-CRZ (1:1000, A11122, Provided by Jan A. Veenstra). Secondary antibodies used:

donkey anti-rabbit IgG conjugated to Alexa 488 and anti-mouse IgG conjugated to Alexa

555 (R37118, Thermo Fisher Scientific). The samples were mounted in Vectorshield

(Vector Laboratory). Images were acquired with Zeiss LSM 700 confocal microscopes,

and were processed with Image J software. All antibodies were diluted in PTX solution.

1001

Fluorscent in situ hybridization (FISH)

FISH was performed as previously reported by Yan et al. [84]. Briefly, the reproductive

system of the female BPH dissection was performed under 1x Phosphate Buffered Saline

(PBS) supplemented with a protease and phosphatase inhibitor cocktail (Shyuanye,

#R40012) on a clean, clear rubber pad using two fine forceps (Dumont, #0108-5-po).

Dissected tissues were fixed using 4% paraformaldehyde (PFA) (Shyuanye, #22039)

diluted in 1 x PBS with 0.1% Tween20 (0.1% PBST) at room temperature (RT) for 20 min

on a shaker. Fixed tissues were washed three times with 0.1% PBST for 15 min each on a

shaker. The tissues were dehydrated in 25%, 50%, 90% and 100% methanol, and then

rehydrated in reverse order, performed at room temperature for 10 minutes at a time. The

tissue was then washed twice with 0.1% PBST and shaken for 5 minutes. A post fixation

was performed by incubating the tissue in 4% PFA in 1 x PBS on a shaker for 20 minutes,

followed by two washes with PBST for 5 minutes each while shaking. The tissue was

permeabilized with proteinase K (10 µg/ml) at room temperature for 2 minutes. After

incubation, the tissue was washed with 0.1% PBST for 5 minutes while shaking. The

PBST was removed, and pre-hybridization solution (Boster, #AR0152) was applied at

55°C for approximately 2 hours. The DIG probes were denatured in a metal bath at 80°C

for 5 minutes and then immediately placed on ice. The denatured DIG-labeled probes

were added to the hybridization solution, and hybridization was carried out overnight at

55°C. Then, the hybridized tissue was washed twice at 55°C for 40 minutes with warm

pre-hybridization solution, followed by four washes at room temperature with 0.1% PBST

for 10 minutes each. The detection of the DIG-labeled probes was performed using an

anti-DIG conjugated fluorescent antibody (1:100 dilution, Jackson, #200-542-156),

incubated at room temperature in 0.1% PBST for 2 hours, followed by three washes with

0.1% PBST. Finally, the tissue was mounted in mounting medium and imaged using a

Zeiss confocal microscope.

1028

Quantification and statistical analysis

All graphs were generated using Prism 9 software (GraphPad Software, La Jolla, CA).

Data presented in this study were first verified for normal distribution by D'Agostino– Pearson

normality test. If normally distributed, Student's *t* test was used for pairwise comparisons, and

one-way ANOVA and chi-square was used for comparisons among multiple groups. If not

normally distributed, Mann–Whitney test was used for pairwise comparisons, and Kruskal–

Wallis test was used for comparisons among multiple groups, followed by Dunn's multiple

comparisons. All data are presented as mean ± s.e.m.

1037

1038   **Data availability statement:** All of the RNA sequence data in this article have been

1039   deposited in the China National Center for Bioinformation database and are accessible in

1040   PRJCA045433.

1041

1042   **ACKNOWLEDGMENTS:** We thank Dr. Jan A. Veenstra for providing CRZ antibody. This

1043   project was supported by National Key R&D Program of China (2022YFD1700200) to S.-F.W.,

1044   the National Natural Science Foundation of China (No. 32472542) to S.-F.W..

1045

References:

1. Do-Hyoung Kima , Y.-Hoon J., , Minsik Y., and , K.-Min L., and Young-Joon Kima,1 (2024). Long-term neuropeptide modulation of female sexual drive via the TRP channel in Drosophila melanogaster. Proceedings of the National Academy of Sciences. 10.1073/pnas.

2. Sun, M., Ma, M., Deng, B., Li, N., Peng, Q., and Pan, Y. (2023). A neural pathway underlying hunger modulation of sexual receptivity in Drosophila females. Cell Rep *42*. 10.1016/j.celrep.2023.113243.

3. Wang, F., Wang, K., Forknall, N., Parekh, R., and Dickson, B.J. (2020). Circuit and Behavioral Mechanisms of Sexual Rejection by Drosophila Females. Current biology : CB. 10.1016/j.cub.2020.07.083.

4. Wang, K., Wang, F., Forknall, N., Yang, T., Patrick, C., Parekh, R., and Dickson, B.J. (2021). Neural circuit mechanisms of sexual receptivity in Drosophila females. Nature *589*, 577-581. 10.1038/s41586-020-2972-7.

5. Kubli, E. (2003). Sex-peptides: seminal peptides of the Drosophila male. Cell Mol Life Sci *60*, 1689-1704. 10.1007/s00018-003-3052.

6. Yapici, N., Kim, Y.J., Ribeiro, C., and Dickson, B.J. (2008). A receptor that mediates the post-mating switch in Drosophila reproductive behaviour. Nature *451*, 33-37. 10.1038/nature06483.

7. Yang, C.H., Rumpf, S., Xiang, Y., Gordon, M.D., Song, W., Jan, L.Y., and Jan, Y.N. (2009). Control of the postmating behavioral switch in Drosophila females by internal sensory neurons. Neuron *61*, 519-526. 10.1016/j.neuron.2008.12.021.

8. Rezaval, C., Pavlou, H.J., Dornan, A.J., Chan, Y.B., Kravitz, E.A., and Goodwin, S.F. (2012). Neural circuitry underlying Drosophila female postmating behavioral responses. Current biology : CB *22*, 1155-1165. 10.1016/j.cub.2012.04.062.

9. Camus, M.F., Huang, C.C., Reuter, M., and Fowler, K. (2018). Dietary choices are influenced by genotype, mating status, and sex in Drosophila melanogaster. Ecol Evol *8*, 5385-5393. 10.1002/ece3.4055.

10. Craig, G.B., Jr. (1967). Mosquitoes: female monogamy induced by male accessory gland substance. Science *156*, 1499-1501. 10.1126/science.156.3781.1499.

11. Judson, C.L. (1967). Feeding and oviposition behavior in the mosquito Aedes aegypti (L.). I. Preliminary studies of physiological control mechanisms. Biol Bull *133*, 369-378. 10.2307/1539832.

12. Liu, C., Tian, N., Chang, P., and Zhang, W. (2024). Mating reconciles fitness and fecundity by switching diet preference in flies. Nature communications *15*, 9912. 10.1038/s41467-024-54369-w.

13. Liu, H., and Kubli, E. (2003). Sex-peptide is the molecular basis of the sperm effect in Drosophila melanogaster. Proceedings of the National Academy of Sciences of the United States of America *100*, 9929-9933. 10.1073/pnas.1631700100.

14. Chapman, T., Bangham, J., Vinti, G., Seifried, B., Lung, O., Wolfner, M.F., Smith, H.K., and Partridge, L. (2003). The sex peptide of Drosophila melanogaster: Female post-mating responses analyzed by using RNA interference. Proceedings of the National Academy of Sciences of the United States of America *100*, 9923-9928. 10.1073/pnas.1631635100.

15.   Duvall, L.B., Basrur, N.S., Molina, H., McMeniman, C.J., and Vosshall, L.B. (2017). A Peptide
Signaling System that Rapidly Enforces Paternity in the Aedes aegypti Mosquito. Current
Biology *27*, 3734-3742.e3735. 10.1016/j.cub.2017.10.074.

16.   Wolfner, M.F. (2002). The gifts that keep on giving: physiological functions and evolutionary
dynamics of male seminal proteins in Drosophila. Heredity *88*, 85-93.
10.1038/sj.hdy.6800017.

17.   Gorter, J.A., Jagadeesh, S., Gahr, C., Boonekamp, J.J., Levine, J.D., and Billeter, J.C. (2016). The
nutritional and hedonic value of food modulate sexual receptivity in Drosophila melanogaster
females. Sci Rep *6*, 19441. 10.1038/srep19441.

18.   Chen, P.S., Stumm-Zollinger, E., Aigaki, T., Balmer, J., Bienz, M., and Böhlen, P. (1988). A male
accessory gland peptide that regulates reproductive behavior of female D. melanogaster. Cell
54, 291-298. https://doi.org/10.1016/0092-8674(88)90192-4.

19.   Ellendersen, B.E., and von Philipsborn, A.C. (2017). Neuronal modulation of D. melanogaster
sexual behaviour. Curr Opin Insect Sci *24*, 21-28. 10.1016/j.cois.2017.08.005.

20.   Klowden, M.J. (1999). The check is in the male: male mosquitoes affect female physiology
and behavior. J Am Mosq Control Assoc *15*, 213-220.

21.   Manning, A. (1962). A Sperm Factor Affecting the Receptivity of Drosophila Melanogaster
Females. Nature *194*, 252-253. 10.1038/194252a0.

22.   McCall, P.J. (2000). The biology of mosquitoes: volume 2, sensory reception and behaviour. By
A.N. Clements (Wallingford, CABI Publishing, 1999). 740 pp. Hard cover £95.00 (US$175.00).
ISBN 0 85199 313 3. Bulletin of Entomological Research *90*, 99-99.
10.1017/S0007485300000171.

23.   Lung, O., Tram, U., Finnerty, C.M., Eipper-Mains, M.A., Kalb, J.M., and Wolfner, M.F. (2002).
The Drosophila melanogaster seminal fluid protein Acp62F is a protease inhibitor that is toxic
upon ectopic expression. Genetics *160*, 211-224.

24.   Zhang, Y.J., Zhang, N., Bu, R.T., Nässel, D.R., Gao, C.F., and Wu, S.F. (2025). A novel male
accessory gland peptide reduces female post-mating receptivity in the brown planthopper.
Plos Genet *21*, e1011699. 10.1371/journal.pgen.1011699.

25.   Rines, I.G., Harrod, A.E., Hunt, J., Sadd, B.M., and Sakaluk, S.K. (2023). Disentangling effects of
mating, nuptial gifts and accessory gland proteins on reproduction in female crickets. Anim
Behav *196*, 1-7. https://doi.org/10.1016/j.anbehav.2022.11.008.

26.   Tripet, F., Touré, Y.T., Dolo, G., and Lanzaro, G.C. (2003). Frequency of multiple inseminations
in field-collected Anopheles gambiae females revealed by DNA analysis of transferred sperm.
Am J Trop Med Hyg *68*, 1-5.

27.   Klowden, M.J., and Russell, R.C. (2004). Mating affects egg maturation in Anopheles gambiae
Giles (Diptera: Culicidae). J Vector Ecol *29*, 135-139.

28.   Naccarati, C., Audsley, N., Keen, J.N., Kim, J.H., Howell, G.J., Kim, Y.J., and Isaac, R.E. (2012).
The host-seeking inhibitory peptide, Aea-HP-1, is made in the male accessory gland and
transferred to the female during copulation. Peptides *34*, 150-157.
10.1016/j.peptides.2011.10.027.

29.   Saudan, P., Hauck, K., Soller, M., Choffat, Y., Ottiger, M., Sporri, M., Ding, Z., Hess, D., Gehrig,
P.M., Klauser, S., et al. (2002). Ductus ejaculatorius peptide 99B (DUP99B), a novel Drosophila
melanogaster sex-peptide pheromone. Eur J Biochem *269*, 989-997.
10.1046/j.0014-2956.2001.02733.x.

30. Herndon, L.A., and Wolfner, M.F. (1995). A Drosophila seminal fluid protein, Acp26Aa, stimulates egg laying in females for 1 day after mating. Proceedings of the National Academy of Sciences of the United States of America *92*, 10114-10118. 10.1073/pnas.92.22.10114.

31. Mackay, T.F.C., Ram, K.R., and Wolfner, M.F. (2007). Sustained Post-Mating Response in Drosophila melanogaster Requires Multiple Seminal Fluid Proteins. Plos Genet *3*. 10.1371/journal.pgen.0030238.

32. Heifetz, Y., Vandenberg, L.N., Cohn, H.I., and Wolfner, M.F. (2005). Two cleavage products of the Drosophila accessory gland protein ovulin can independently induce ovulation. Proceedings of the National Academy of Sciences of the United States of America *102*, 743-748. 10.1073/pnas.0407692102.

33. Desplan, C., Gligorov, D., Sitnik, J.L., Maeda, R.K., Wolfner, M.F., and Karch, F. (2013). A Novel Function for the Hox Gene Abd-B in the Male Accessory Gland Regulates the Long-Term Female Post-Mating Response in Drosophila. Plos Genet *9*. 10.1371/journal.pgen.1003395.

34. Hopkins, B.R., Angus-Henry, A., Kim, B.Y., Carlisle, J.A., Thompson, A., and Kopp, A. (2024). Decoupled evolution of the Sex Peptide gene family and Sex Peptide Receptor in Drosophilidae. Proceedings of the National Academy of Sciences of the United States of America *121*, e2312380120. 10.1073/pnas.2312380120.

35. Jang, Y.-H., Chae, H.-S., and Kim, Y.-J. (2017). Female-specific myoinhibitory peptide neurons regulate mating receptivity in Drosophila melanogaster. Nature communications *8*. 10.1038/s41467-017-01794-9.

36. Chen, J., Zhu, P., Jin, S., Zhang, Z., Jiang, S., Li, S., Liu, S., Peng, Q., and Pan, Y. (2025). A hormone-to-neuropeptide pathway inhibits sexual receptivity in immature Drosophila females. Proceedings of the National Academy of Sciences of the United States of America *122*, e2418481122. 10.1073/pnas.2418481122.

37. Avila, F.W., Sirot, L.K., LaFlamme, B.A., Rubinstein, C.D., and Wolfner, M.F. (2011). Insect seminal fluid proteins: identification and function. Annual review of entomology *56*, 21-40. 10.1146/annurev-ento-120709-144823.

38. Hasemeyer, M., Yapici, N., Heberlein, U., and Dickson, B.J. (2009). Sensory Neurons in the Drosophila Genital Tract Regulate Female Reproductive Behavior. Neuron *61*, 511-518. 10.1016/j.neuron.2009.01.009.

39. Rezával, C., Pavlou, H.J., Dornan, A.J., Chan, Y.B., Kravitz, E.A., and Goodwin, S.F. (2012). Neural circuitry underlying Drosophila female postmating behavioral responses. Current biology : CB *22*, 1155-1165. 10.1016/j.cub.2012.04.062.

40. Auer, T.O., and Benton, R. (2016). Sexual circuitry in Drosophila. Curr Opin Neurobiol *38*, 18-26. 10.1016/j.conb.2016.01.004.

41. Lin, P.X., Peng, Y.X., Xing, J.Y., Liu, Z.Y., Guo, F.R., Thia, J.A., Gao, C.F., and Wu, S.F. (2025). Cis-regulation of the CYP6CS1 gene and its role in mediating cross-resistance in a pymetrozine-resistant strain of Nilaparvata lugens. Insect biochemistry and molecular biology *177*, 104261. 10.1016/j.ibmb.2025.104261.

42. Wu, S.F., Zeng, B., Zheng, C., Mu, X.C., Zhang, Y., Hu, J., Zhang, S., Gao, C.F., and Shen, J.L. (2018). The evolution of insecticide resistance in the brown planthopper (Nilaparvata lugens Stål) of China in the period 2012-2016. Sci Rep *8*, 4586. 10.1038/s41598-018-22906-5.

43. Song, X.Y., Peng, Y.X., Wang, L.X., Ye, W.N., Pei, X.G., Zhang, Y.C., Zhang, S., Gao, C.F., and Wu, S.F. (2022). Monitoring, cross-resistance, inheritance, and fitness costs of brown

planthoppers, Nilaparvata lugens, resistance to pymetrozine in China. Pest management
science *78*, 3980-3987. 10.1002/ps.7017.

44. Ye, W.N., Li, Y., Zhang, Y.C., Liu, Z.Y., Song, X.Y., Pei, X.G., Wu, S.F., and Gao, C.F. (2024).
Area-wide survey and monitoring of insecticide resistance in the brown planthopper,
Nilaparvata lugens (Stål), from 2020 to 2023 in China. Pestic Biochem Physiol *205*, 106173.
10.1016/j.pestbp.2024.106173.

45. Tayler, T.D., Pacheco, D.A., Hergarden, A.C., Murthy, M., and Anderson, D.J. (2012). A
neuropeptide circuit that coordinates sperm transfer and copulation duration in
Drosophila. Proceedings of the National Academy of Sciences *109*, 20697-20702.
10.1073/pnas.1218246109.

46. Li, H., Cheng, S., Niu, Y., Diao, T., Zhang, Q., and Zhang, W. (2025). A heart releasing
neuropeptide that synchronizes brain-heart regulation during courtship behavior. bioRxiv,
2025.2003.2031.646145. 10.1101/2025.03.31.646145.

47. Tayler, T.D., Pacheco, D.A., Hergarden, A.C., Murthy, M., and Anderson, D.J. (2012). A
neuropeptide circuit that coordinates sperm transfer and copulation duration in Drosophila.
Proceedings of the National Academy of Sciences of the United States of America *109*,
20697-20702. 10.1073/pnas.1218246109.

48. Gospocic, J., Shields, E.J., Glastad, K.M., Lin, Y., Penick, C.A., Yan, H., Mikheyev, A.S., Linksvayer,
T.A., Garcia, B.A., Berger, S.L., et al. (2017). The Neuropeptide Corazonin Controls Social
Behavior and Caste Identity in Ants. Cell *170*, 748-759.e712. 10.1016/j.cell.2017.07.014.

49. Hou, Q.-L., Chen, E.-H., Jiang, H.-B., Yu, S.-F., Yang, P.-J., Liu, X.-Q., Park, Y., Wang, J.-J., and
Smagghe, G. (2018). Corazonin Signaling Is Required in the Male for Sperm Transfer in the
Oriental Fruit Fly Bactrocera dorsalis. Front Physiol *9*. 10.3389/fphys.2018.00660.

50. Shohat-Ophir, G., Kaun, K.R., Azanchi, R., Mohammed, H., and Heberlein, U. (2012). Sexual
deprivation increases ethanol intake in Drosophila. Science *335*, 1351-1355.
10.1126/science.1215932.

51. Zer-Krispil, S., Zak, H., Shao, L., Ben-Shaanan, S., Tordjman, L., Bentzur, A., Shmueli, A., and
Shohat-Ophir, G. (2018). Ejaculation Induced by the Activation of Crz Neurons Is Rewarding to
Drosophila Males. Current Biology *28*, 1445-1452.e1443. 10.1016/j.cub.2018.03.039.

52. Cheng, J., Zhao, P., Zhu, L., Zhu, F., Tian, Z., Shen, Z., Liu, X., and Liu, X. (2022). Corazonin
signaling modulates the synthetic activity of male accessory gland in Grapholita molesta. Int J
Biol Macromol *216*, 446-455. 10.1016/j.ijbiomac.2022.07.025.

53. Yang, C.h., Belawat, P., Hafen, E., Jan, L.Y., and Jan, Y.N. (2008). Drosophila Egg-Laying Site
Selection as a System to Study Simple Decision-Making Processes. Science *319*, 1679-1683.
10.1126/science.1151842.

54. Tanaka, Y., Suetsugu, Y., Yamamoto, K., Noda, H., and Shinoda, T. (2014). Transcriptome
analysis of neuropeptides and G-protein coupled receptors (GPCRs) for neuropeptides in the
brown planthopper Nilaparvata lugens. Peptides *53*, 125-133.
10.1016/j.peptides.2013.07.027.

55. Roller, L., Tanaka, Y., and Tanaka, S. (2003). Corazonin and corazonin-like substances in the
central nervous system of the Pterygote and Apterygote insects. Cell Tissue Res *312*, 393-406.
10.1007/s00441-003-0722-4.

56. Nässel, D.R., and Zandawala, M. (2019). Recent advances in neuropeptide signaling in
Drosophila, from genes to physiology and behavior. 10.7287/peerj.preprints.27515v2.

57. Kubrak, O.I., Lushchak, O.V., Zandawala, M., and Nässel, D.R. (2016). Systemic corazonin
signalling modulates stress responses and metabolism in Drosophila. Open Biol *6*.
10.1098/rsob.160152.

58. Nallasivan, M.P., Singh, D.N.D., Saleh, M.S.R.S., and Soller, M. (2024). Sex-peptide targets
distinct higher order processing neurons in the brain to induce the female post-mating
response. eLife Sciences Publications, Ltd.

59. Isaac, R.E., Li, C., Leedale, A.E., and Shirras, A.D. (2010). Drosophila male sex peptide inhibits
siesta sleep and promotes locomotor activity in the post-mated female. Proc Biol Sci *277*,
65-70. 10.1098/rspb.2009.1236.

60. Bath, E., Bowden, S., Peters, C., Reddy, A., Tobias, J.A., Easton-Calabria, E., Seddon, N.,
Goodwin, S.F., and Wigby, S. (2017). Sperm and sex peptide stimulate aggression in female
Drosophila. Nat Ecol Evol *1*, 0154. 10.1038/s41559-017-0154.

61. Predel, R., Neupert, S., Russell, W.K., Scheibner, O., and Nachman, R.J. (2007). Corazonin in
insects. Peptides *28*, 3-10. 10.1016/j.peptides.2006.10.011.

62. Boerjan, B., Verleyen, P., Huybrechts, J., Schoofs, L., and De Loof, A. (2010). In search for a
common denominator for the diverse functions of arthropod corazonin: a role in the
physiology of stress? Gen Comp Endocrinol *166*, 222-233. 10.1016/j.ygcen.2009.09.004.

63. Veenstra, J.A. (2009). Does corazonin signal nutritional stress in insects? Insect biochemistry
and molecular biology *39*, 755-762. 10.1016/j.ibmb.2009.09.008.

64. Zhao, Y., Bretz, C.A., Hawksworth, S.A., Hirsh, J., and Johnson, E.C. (2010). Corazonin neurons
function in sexually dimorphic circuitry that shape behavioral responses to stress in
Drosophila. Plos One *5*, e9141. 10.1371/journal.pone.0009141.

65. Veenstra, J.A. (1989). Isolation and structure of corazonin, a cardioactive peptide from the
American cockroach. Febs Lett *250*, 231-234. 10.1016/0014-5793(89)80727-6.

66. Tawfik, A.I., Tanaka, S., De Loof, A., Schoofs, L., Baggerman, G., Waelkens, E., Derua, R., Milner,
Y., Yerushalmi, Y., and Pener, M.P. (1999). Identification of the gregarization-associated
dark-pigmentotropin in locusts through an albino mutant. Proceedings of the National
Academy of Sciences of the United States of America *96*, 7083-7087.
10.1073/pnas.96.12.7083.

67. Tanaka, Y., Hua, Y., Roller, L., and Tanaka, S. (2002). Corazonin reduces the spinning rate in the
silkworm, Bombyx mori. J Insect Physiol *48*, 707-714. 10.1016/s0022-1910(02)00094-x.

68. Kim, Y.J., Spalovská-Valachová, I., Cho, K.H., Zitnanova, I., Park, Y., Adams, M.E., and Zitnan, D.
(2004). Corazonin receptor signaling in ecdysis initiation. Proceedings of the National
Academy of Sciences of the United States of America *101*, 6704-6709.
10.1073/pnas.0305291101.

69. Imura, E., Shimada-Niwa, Y., Nishimura, T., Hückesfeld, S., Schlegel, P., Ohhara, Y., Kondo, S.,
Tanimoto, H., Cardona, A., Pankratz, M.J., and Niwa, R. (2020). The Corazonin-PTTH Neuronal
Axis Controls Systemic Body Growth by Regulating Basal Ecdysteroid Biosynthesis in
Drosophila melanogaster. Current biology : CB *30*, 2156-2165.e2155.
10.1016/j.cub.2020.03.050.

70. Zandawala, M., Nguyen, T., Balanya Segura, M., Johard, H.A.D., Amcoff, M., Wegener, C.,
Paluzzi, J.P., and Nassel, D.R. (2021). A neuroendocrine pathway modulating osmotic stress in
Drosophila. Plos Genet *17*, e1009425. 10.1371/journal.pgen.1009425.

71. Xi, J., Hamanaka, Y., and Shiga, S. (2025). Corazonin mediates photoperiodically induced diapause in the bean bug Riptortus pedestris. J Exp Biol *228*. 10.1242/jeb.250528.

72. Tian, S., Zandawala, M., Beets, I., Baytemur, E., Slade, S.E., Scrivens, J.H., and Elphick, M.R. (2016). Urbilaterian origin of paralogous GnRH and corazonin neuropeptide signalling pathways. Sci Rep *6*, 28788. 10.1038/srep28788.

73. Yañez Guerra, L.A., and Zandawala, M. (2023). Discovery of Paralogous GnRH and Corazonin Signaling Systems in an Invertebrate Chordate. Genome Biol Evol *15*. 10.1093/gbe/evad108.

74. Knobil, E. (1988). The hypothalamic gonadotrophic hormone releasing hormone (GnRH) pulse generator in the rhesus monkey and its neuroendocrine control. Hum Reprod *3*, 29-31. 10.1093/oxfordjournals.humrep.a136647.

75. Millar, R.P. (2005). GnRHs and GnRH receptors. Anim Reprod Sci *88*, 5-28. 10.1016/j.anireprosci.2005.05.032.

76. Bergland, A.O., Chae, H.S., Kim, Y.J., and Tatar, M. (2012). Fine-scale mapping of natural variation in fly fecundity identifies neuronal domain of expression and function of an aquaporin. Plos Genet *8*, e1002631. 10.1371/journal.pgen.1002631.

77. Barredo, C.G., Gil-Marti, B., Deveci, D., Romero, N.M., and Martin, F.A. (2020). Timing the Juvenile-Adult Neurohormonal Transition: Functions and Evolution. Front Endocrinol (Lausanne) *11*, 602285. 10.3389/fendo.2020.602285.

78. McKim, T.H., Gera, J., Gayban, A.J., Reinhard, N., Manoli, G., Hilpert, S., Helfrich-Förster, C., and Zandawala, M. (2024). Synaptic connectome of a neurosecretory network in the Drosophila brain. bioRxiv. 10.1101/2024.08.28.609616.

79. Wu, S.-F., Jv, X.-M., Li, J., Xu, G.-J., Cai, X.-Y., and Gao, C.-F. (2017). Pharmacological characterisation and functional roles for egg-laying of a β-adrenergic-like octopamine receptor in the brown planthopper Nilaparvata lugens. Insect biochemistry and molecular biology *87*, 55-64. https://doi.org/10.1016/j.ibmb.2017.06.008.

80. Krogh, A., Larsson, B., von Heijne, G., and Sonnhammer, E.L. (2001). Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. J Mol Biol *305*, 567-580. 10.1006/jmbi.2000.4315.

81. Livak, K.J., and Schmittgen, T.D. (2001). Analysis of Relative Gene Expression Data Using Real-Time Quantitative PCR and the 2−ΔΔCT Method. Methods *25*, 402-408. https://doi.org/10.1006/meth.2001.1262.

82. Sun, H., Bu, L.-A., Su, S.-C., Guo, D., Gao, C.-F., and Wu, S.-F. (2023). Knockout of the odorant receptor co-receptor, orco, impairs feeding, mating and egg-laying behavior in the fall armyworm Spodoptera frugiperda. Insect biochemistry and molecular biology *152*. 10.1016/j.ibmb.2022.103889.

83. Zhang, Y.C., Gao, Y., Ye, W.N., Peng, Y.X., Zhu, K.Y., and Gao, C.F. (2023). CRISPR/Cas9-mediated knockout of NlCYP6CS1 gene reveals its role in detoxification of insecticides in Nilaparvata lugens (Hemiptera: Delphacidae). Pest management science *79*, 2239-2246. 10.1002/ps.7404.

84. Yan, H., Opachaloemphan, C., Carmona-Aldana, F., Mancini, G., Mlejnek, J., Descostes, N., Sieriebriennikov, B., Leibholz, A., Zhou, X., Ding, L., et al. (2022). Insulin signaling in the long-lived reproductive caste of ants. Science *377*, 1092-1099. 10.1126/science.abm8767.