Vitamin K-dependent carboxylation in osteoblasts regulates bone resorption through GAS6

Running title: Vitamin K-dependent carboxylation and bone resorption

Monica Pata[1], Diep Ngoc Thi Pham[1], Julie Lacombe[1], B. Ashok Reddy[1], Young Woong Kim[1], Abeer Gamal Ali Ahmed[3], Monzur Murshed[3,4], Mathieu Ferron[1,2,5]

[1] Molecular Physiology Research Unit, Institut de Recherches Cliniques de Montréal, Montréal, Québec, Canada, H2W 1R7.

[2] Department of Medicine, Université de Montréal, Montréal, Québec, Canada, H3T 1J4

[3] Faculty of Dental Medicine and Oral Health Sciences, McGill University, Montreal, Quebec, Canada, H3A 0C7

[4] Department of Medicine, McGill University, Montreal, Quebec, Canada, H4A 3J1; Shriners hospital for children, Montreal, Quebec, Canada, H4A 0A9.

[5] Division of Experimental Medicine, Faculty of Medicine and Health Sciences, McGill University, Montréal, Québec, Canada. H4A 3J1

Author's emails:

Monica.pata@ircm.qc.ca; phamtnd@hotmail.fr; Julie.lacombe@ircm.qc.ca; ashokbareddy@gmail.com; YoungWoong.Kim@hli.ubc.ca; abeer.ahmed@mail.mcgill.ca; monzur.murshed@mcgill.ca

* Correspondence to:
Mathieu Ferron, PhD
Institut de Recherches Cliniques de Montréal
110 Ave. des Pins O.
Montréal, QC,
H2W 1R7, Canada
Phone: 514-987-5754
Fax: 514-987-5649
Email: mathieu.ferron@ircm.qc.ca

Contributions
M.F. and M.P. assembled the figures. M.F. wrote the manuscript with suggestions from the other authors. M.P., M.F., D.N.T.P., J.L., B.A.R. and Y.W.K., collected and analyzed data. M.F. conceived and designed the project and provided supervision to M.P., D.N.T.P., J.L., B.A.R. and Y.W.K. A.G.A.A. and M.M. collected and analyzed the µCT data. M.F. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of data and the accuracy of data analysis.

Abstract

Studies in humans suggest that vitamin K is involved in the regulation of bone remodeling, but the precise mechanism at play remains unknown. In cells, vitamin K functions as a co-factor for the γ-glutamyl carboxylase (GGCX), an enzyme responsible for the conversion of glutamic acid residues (Glu) into γ-carboxyglutamic acid (Gla) residues in secreted proteins. We aim here at determining the role of γ-carboxylation in bone remodeling and at identifying the Gla protein(s) involved. We show that mice lacking γ-carboxylation specifically in osteoblasts ($Ggcx^{flox/flox}$;OCN-Cre) have increased bone mass at 6 months of age due to a reduced number of multinucleated bone resorbing osteoclasts. In co-culture experiments, $Ggcx$-deficient osteoblasts were less effective than control osteoblasts at supporting the generation of osteoclasts. Among all known Gla proteins, we identify GAS6 as an osteoblast-secreted γ-carboxylated factor which signals to differentiating osteoclasts. The GAS6 receptors MerTK and AXL are expressed in pre-osteoclasts and pharmacological inhibitors of AXL and MerTK block osteoclast generation in co-culture. Conversely, recombinant γ-carboxylated GAS6 dose-dependently increases the size of osteoclasts and the number of nuclei per osteoclast in culture. GAS6 marginally affected the induction of osteoclast-specific genes during osteoclast differentiation but significantly increased pre-osteoclast fusion. Finally, increasing bone marrow GAS6 level in transgenic mice was sufficient to increase the number and size of osteoclasts and to decrease bone mass. This work identifies GAS6 as a novel osteoblast derived vitamin K-dependent protein regulating osteoclast maturation.

Keywords: Osteoclast; osteoblast; vitamin K; gamma-carboxylation; GAS6; GGCX

Introduction

The acquisition and maintenance of an adequate skeleton depend on the process of bone turnover, which requires the activity of two cell types: the osteoclast, responsible for the resorption of the mineralized bone extracellular matrix (ECM) and the osteoblast, responsible for the synthesis, secretion and mineralization of the bone ECM. An imbalance between the activities of these two cell types can result in severe bone diseases such as osteoporosis. Like many chronic aging disorders, osteoporosis appears to be a multifactorial disease. Hence, although several genetic variants may contribute to the physiopathology of osteoporosis [1,2], epigenetic, environmental and behavioral factors have also been implicated [3,4]. For instance, several micronutrients including vitamins are known to play a role in the acquisition and the maintenance of a healthy skeleton and could therefore influence the etiology of osteoporosis [4,5]. This is evidenced by the profound impact that vitamins C, D or B12 deficiency has on bone density or quality in humans [6-9]. Cross-sectional and prospective studies suggest that vitamin K (VK) might be another micronutrient playing a key role in the regulation of bone mass.

Longitudinal prospective studies have reported that lower VK intake or serum levels were associated with decreased bone mineral density (BMD) [10,11] or increased fracture risk in postmenopausal women [12-14]. There are two naturally occurring forms of VK: VK_1 or phylloquinone, synthesized in plants, and VK_2 or menaquinones (e.g., MK-4 and MK-7), present in animals and fermented foods [15]. Bone was shown to contain both VK_1 and MK-4 [16,17]. A number of clinical trials have tested the effects of VK_1 or VK_2 supplementation on BMD or fracture risk, but have led to conflicting results [15]. While some studies did find that MK-4, MK-7 or VK_1 increased femoral and/or vertebral BMD and/or reduced fracture risk, others found no

significant impact [18-22]. A few meta-analyses of interventional studies suggest that VK

supplementation had a significant and positive effect on BMD at lumbar spine and forearm

[18,19,23], while another suggested that VK prevents fractures without impacting BMD [24]. Therefore,

whether VK is directly implicated in the regulation of bone mass or quality remains controversial.

Moreover, the molecular and cellular mechanisms through which VK influences bone density

and/or quality remain uncharacterized.

In vertebrates, the only well described cellular function of VK is to serve as a co-factor during

the γ-carboxylation reaction that converts glutamic acid (Glu) residues to γ-carboxyglutamic acid

(Gla) residues in specific proteins [25]. Hence, γ-carboxylated proteins (also called "Gla" proteins)

are either secreted or transmembrane polypeptides with the potential of endocrine, paracrine or

autocrine functions. Two enzymes are involved in γ-carboxylation: γ-glutamyl carboxylase (γ-

carboxylase or GGCX) and vitamin K oxidoreductase (VKORC1) [26]. γ-carboxylase requires

reduced VK (VKH_2) as an essential cofactor, which upon carboxylation is oxidized to VK

epoxide (VKO). VKO is next reconverted to VKH_2 by VKORC1.

Osteocalcin (Ocn) is an osteoblast-derived Gla protein presents at high concentration in the bone

ECM. Because γ-carboxylated Ocn has great affinity for hydroxyapatite, the mineral component

of bone ECM, it was originally believed to play a role in bone mineralization [27]. The analysis of

Ocn-deficient mice or rats (*$Ocn^{-/-}$*) suggested that the absence of osteocalcin may slightly increase

bone formation and bone density [28,29]. However, two other studies involving different strains of

$Ocn^{-/-}$ mice did not detected any impact on bone homeostasis [30,31]. Altogether, these genetic data

do not support a major role for Ocn in bone mass accrual or turnover, suggesting that the

mechanism and the Gla protein(s) through which VK may influence bone health remain to be

characterized.

We aim here at addressing specifically the role of γ-carboxylation in bone remodeling in vivo. We

show that an osteoblast-specific inactivation of *Ggcx* in mice is associated with an increased bone

density resulting from a reduced number of osteoclasts without significant impact on bone

formation. We establish that *Ggcx*-deficient osteoblasts are less effective at supporting osteoclast

formation ex vivo and identify GAS6 as an osteoblast-secreted γ-carboxylated factor which

signals to osteoclast precursors. Finally, using cell-based assays and a gain-of-function model in

mice, we shown that γ-carboxylated GAS6 promote pre-osteoclasts fusion. These data reveal an

unexpected role for γ-carboxylation in coupling bone resorption to bone formation and provides

new insights on the function of VK in bone homeostasis.

Results

The enzymes of the VK cycle are predominantly expressed in osteoblasts

To identify the bone cell type(s) mediating the potential impact of VK on bone, we examined the expression of GGCX and VKORC1 in osteoblasts and osteoclasts at the mRNA and protein levels. Quantitative PCR (qPCR) revealed that both *Ggcx* and *Vkorc1* expression levels are about 10-fold higher in mouse proliferating non-mineralized osteoblasts (pre-OB) compared to bone marrow derived monocytes (BMMC) or fully differentiated osteoclast (OCL) (**Fig.1A**). While *Vkorc1* mRNA level remains elevated in mineralized osteoblast culture (OB), the expression of *Ggcx* is reduced in the same cells compared to pre-OB cultures (**Fig.1A**). Western blotting confirmed that GGCX and VKORC1 are more abundant in primary osteoblast cultures compared to osteoclasts. Interestingly, the expression level of GGCX and VKORC1 in osteoblasts was comparable to the one observed in liver where these enzymes are known to play a critical role in the γ-carboxylation and activation of several coagulation factors (**Fig. 1B**).

Inactivation of GGCX in osteoblasts results in increased bone density in mice

Given the predominant expression of the VK-cycle enzymes in osteoblasts, we inactivated *Ggcx* specifically in this cell type and investigate the impact on bone remodeling. For this purpose, *Ggcx$^{flox/flox}$* mice were bred with *Osteocalcin-Cre* (OCN-Cre) mice expressing the Cre recombinase in mature osteoblasts only [32]. We previously reported that *Ggcx* was specifically and efficiently inactivated in osteoblasts in *Ggcx$^{fl/fl}$;OCN-Cre* mice [33]. The percentage of circulating carboxylated osteocalcin, an osteoblast-specific VK-dependent protein, was reduced by 80% to 90% between 2 and 6 months of age in these mice, confirming efficient inactivation of VK-

dependent carboxylation in osteoblasts in vivo (Supplementary Figure 1). In a previous study, we

reported that 3-month-old $Ggcx^{fl/fl}$;*OCN-Cre* male mice had normal trabecular bone density [33].

Since bone mass decreases with aging, we characterized the bone parameters of the

$Ggcx^{fl/fl}$;*OCN-Cre* at 6 months of age when bone mass has already started to decline in mice.

Histological analysis of lumbar vertebrae indicated that trabecular bone density was significantly

increased by about 30% in 6-month-old $Ggcx^{fl/fl}$;*OCN-Cre* mice (**Fig. 1C-D**). Trabecular bone

density was also increased by about 30% in the femur of $Ggcx^{fl/fl}$;*OCN-Cre* mice analyzed by

microcomputed tomography (μCT), although this parameter did not reach statistical significance

due to higher variation (**Fig. 1E-F**). However, trabecular bone surface density (BS/TV), was

significantly increased in the same animals (**Fig. 1F**). In addition, trabecular spacing (Tb.S) was

significantly reduced, while trabecular number (Tb.N) and connectivity density (Conn.D), a

computational measure of the inter-connectivity among trabeculae, were both significantly

increased (**Fig. 1G**). Trabecular thickness was unaffected in the $Ggcx^{fl/fl}$;*OCN-Cre* femur (**Fig.**

1G). Altogether, these data indicate that the absence of VK-dependent carboxylation in

osteoblasts is associated with an increased bone mass at 6 months of age.

Decreased bone resorption in $Ggcx^{fl/fl}$;*OCN-Cre* mice

To get insight into the cellular mechanism behind the increased bone mass phenotype of the

$Ggcx^{fl/fl}$;*OCN-Cre* mice, static and dynamic bone histomorphometry analyses were performed at

6 months of age. Mineral apposition rate (MAR), measured using double calcein labeling, was

slightly but significantly reduced in absence of GGCX in osteoblasts (**Fig.2A-B**). However, bone

formation rate (BFR/BS), osteoblast number and osteoblast surface were not significantly

affected (**Fig.2C-E**). In contrast, both osteoclast number and surface were significantly decreased

in $Ggcx^{fl/fl}$;*OCN-Cre* mice (**Fig. 2F-H**). Decrease circulating carboxy-terminal collagen crosslinks

(CTX) independently confirmed reduced osteoclast activity in the same animals (**Fig.2I**). Overall,

these analyses suggest that the increased bone mass observed in the absence of γ-carboxylation in

osteoblasts is mainly driven by a reduction in osteoclastic bone resorption.

Decreased osteoclastogenesis in absence of *Ggcx* in osteoblasts

Osteoclast differentiation is dependent on factors, such as RANKL and M-CSF, produced by

cells of the osteoblast lineage including pre-osteoblast and osteocytes. Because of the reduction

in osteoclasts number in the *Ggcx$^{fl/fl}$;OCN-Cre* mice, we hypothesized that *Ggcx*-deficient

osteoblasts may be less efficient at supporting osteoclast differentiation or maturation. This was

tested using co-cultures of wildtype bone marrow cells with either osteoblasts lacking GGCX

(i.e., *Ggcx$^{fl/fl}$* osteoblasts transduced with a CRE expressing adenovirus) or control osteoblasts

(i.e., *Ggcx$^{fl/fl}$* osteoblasts transduced with a GFP expressing adenovirus). We previously reported

very efficient inactivation of GGCX in osteoblasts at the mRNA and protein levels using this

approach [33] (see also **Fig. 1B**). The coculture was performed in the presence of 1,25 vitamin D_3

(VitD3) and prostaglandin E_2 (PGE2), which promote the expression of pro-osteoclastogenesis

factors by osteoblasts. These experiments indicated that the number of osteoclasts was

significantly reduced by approximately 3-fold in the co-cultures involving GGCX-deficient

osteoblasts (**Fig.3A-B**). This result suggests that γ-carboxylation in osteoblasts may regulate the

production or the activity of a pro-osteoclastogenic factor. We next tested if the absence of

osteocalcin (OCN), a γ-carboxylated protein specifically secreted by osteoblasts and osteocytes,

impacted osteoclasts formation in the same co-culture assay and we found that the absence of

Ocn in osteoblast did not affect the number of osteoclasts obtained from WT bone marrow cells

(**Fig. 3C-D**).

Cells from the osteoblast lineage produce factors that promote (i.e., RANKL and M-CSF) or

inhibit (i.e., osteoprotegerin; OPG) osteoclasts generation. We assessed the expression of these

factors in control and GGCX-deficient osteoblasts to determine if γ-carboxylation regulates the

pro-osteoclastogenic potential of osteoblasts. qPCR revealed that as expected, VitD3 and PGE2,

robustly increased the expression of *Rankl* and suppressed *Opg* expression (**Fig. 3E**).

Consequently, the *Rankl/Opg* ratio, which is positively associated with osteoclast formation, was

increased by more than 10-fold with VitD3 and PGE2. Importantly, the absence of *Ggcx* in

osteoblasts had no significant impact on *Rankl, Opg* and the *Rankl/Opg* ratio in the pro-

osteoclastogenic condition (**Fig. 3E**). The expression of *Csf1,* the gene encoding for M-CSF, was

also unaffected in the absence of GGCX in osteoblasts (**Fig. 3E**). These data suggest that γ-

carboxylation positively regulates osteoclast formation independently of osteocalcin and of the

expression of known pro-osteoclastogenic factors.

The γ-carboxylated protein GAS6 is expressed by osteoblasts and activates its receptors on

pre-osteoclasts

Based on the results obtained in vivo and in co-culture assays, we postulated the existence of a γ-

carboxylated protein, different from osteocalcin, produced by osteoblasts and promoting

osteoclast formation. Expression of all the known γ-carboxylated proteins encoding genes was

measured by qPCR in primary mouse osteoblasts. As expected, we detected expression of the γ-

carboxylated ECM proteins osteocalcin (*Ocn*), matrix Gla protein (*Mgp*) and periostin (*Postn*),

previously shown to be produced by the osteoblastic lineage (**Fig.4A**) [34-36]. Previous studies have

established that inactivation of *Mgp* or *Postn* in mice had no impact on osteoclast number and

surface [37,38], and our own data show that osteocalcin is no required for osteoclastogenesis in

coculture assays (**Fig.3A-D**). We therefore excluded periostin, osteocalcin and MGP, as potential

mediators of the positive effect of γ-carboxylation on osteoclasts formation. Of note, the γ-

carboxylated coagulation factors and the Proline Rich and Gla domain 1 to 4 proteins (*Prrg1-4*)

were undetectable in osteoblasts. However, the gene encoding Growth Arrest Specific 6 (*Gas6*)

was expressed at a level comparable or higher to *Mgp*, *Ocn* and *Postn* (**Fig. 4A**).

GAS6 is a secreted γ-carboxylated protein that functions as a ligand for the TAM family of

tyrosine kinase receptors that includes AXL, Tyro3 and MerTK. Given that GAS6 is a signaling

molecule and that TAM receptors were previously shown to be present on myeloid cells in other

tissues [39], we decided to assess the presence of functional GAS6 receptors on osteoclasts. The

expression of TAM receptors was evaluated at different stages of bone marrow-derived

monocytes (BMMC) differentiation into osteoclasts. Both *MerTK* and *Axl* were detected in

primary BMMC (d0) and in differentiating osteoclasts (d2, d4 and d6), while *Tyro3* expression

was below the detection limit in most samples (**Fig.4B**).

We previously established that purified recombinant GAS6 (recGAS6) produced in HEK293

cells in the presence of VK is fully γ-carboxylated and able to activate TAM receptors on muscle

cells (Supplemental Fig. 2) [40]. We thus tested if this ligand could also elicit TAM receptors

activation in primary pre-osteoclasts. For this purpose, BMMC were serum starved for 3 hours,

stimulated for various times with recGAS6 (200ng/mL) and the phosphorylation of AXL and

MerTK or their downstream target AKT [41] assessed by Western blotting. Following recGAS6

stimulation, we could detect an increase in the phosphorylation of MerTK (Y753) and AXL

(Y702) on a tyrosine residue previously shown to be a dominant autophosphorylation site in the

tyrosine kinase domain of TAM receptors that leads to the activation of their kinase activity [42]. In

the condition of this assay, recGAS6 also robustly induces the phosphorylation of AKT on serine

473, reflecting the activation of the phosphatidylinositol 3-kinase (PI3K) pathway. Together,

these results support the existence of functional GAS6-TAM signaling coupling osteoblasts to

osteoclasts.

GAS6 signaling promotes osteoclast formation ex vivo

We then tested if GAS6-TAM signaling influences osteoclast formation using two approaches. In

a first set of experiments, we assessed osteoclastogenesis in co-cultures of WT bone marrow cells

and WT osteoblasts in the presence of increasing concentrations of two pharmacological TAM

inhibitors, LDC1267 and R428 (bemcentinib) [43,44]. As shown in Figure 5A, LDC1267, a pan-

TAM inhibitor, dose dependently inhibits the formation of TRAP-positive multinucleated

osteoclasts in these co-cultures, with an IC50 of ~750nM. R428, which is >50-100-fold more

selective for AXL than MerTK and Tyro3, also inhibits osteoclast formation with a lower IC50 of

~250nM (**Fig.5B**). These data suggest that TAM receptor signaling promotes osteoclast

formation in co-culture assay.

In a second approach, we tested if exogenous recGAS6 (25-200 ng/mL) impacts osteoclast

formation in primary BMMC cultures in the presence of RANKL (20ng/mL) and M-CSF

(10ng/mL). In these experiments, recGAS6 significantly and dose-dependently increases

osteoclasts formation (**Fig.5C-E**). Quantification of the cultures following TRAP staining

indicated that recGAS6 increased both the surface covered by TRAP+ osteoclasts (**Fig.5D**) and

the number of multinucleated osteoclasts (**Fig. 5E)**, as early as 4 days of culture when small

multinucleated TRAP-positive osteoclasts begin to appear. In addition, the number of individual

nuclei per osteoclast was significantly increased in presence of recGAS6 (**Fig.5F**), indicating the

formation of larger osteoclasts. These results identify GAS6-TAM signaling as a modulator of

osteoclastogenesis ex vivo.

GAS6 increases the fusion of pre-osteoclasts in culture

Osteoclast formation involves first, the transcriptional activation of a differentiation program in

mononucleated pre-osteoclasts and second, the fusion of these pre-osteoclasts to form mature

multinucleated osteoclasts. We sought to determine if GAS6 signaling increases osteoclast

formation by promoting differentiation of BMMC into osteoclasts. The expression of genes

encoding several markers of osteoclast differentiation was measured by qPCR at 2, 4 and 6 days

of differentiation in the absence or presence of intermediate doses of recGAS6 (i.e., 50 and 100

ng/mL). The presence of recGAS6 in the media did not significantly impact the expression of

Acp5 (TRAP) and *Clcn7* which gradually increase between 2 and 6 days of differentiation as

expected (**Fig.6A-B**). The highest dose of GAS6 (100ng/mL) induces a small and significant

increase in the expression of *Ctsk*, the gene encoding cathepsin K, at day 4 and 6 of

differentiation, and of *Dcstamp* at day 6 of differentiation (**Fig.6C-D**). Overall, this marginal

impact of GAS6 on the expression of the osteoclast differentiation program cannot explain its

strong positive effect on the generation of multinucleated osteoclasts in the same culture

conditions. Indeed, 50 and 100 ng/mL of GAS6 increases by >50% the number of osteoclasts and

the area of TRAP+ osteoclast (**Fig.5C-E**).

Since recGAS6 increases the number of nuclei per OCL (**Fig.5F**), we next tested if GAS6

promotes pre-osteoclast fusion. For this purpose, we established a pre-osteoclast fusion assay in

which an equal number of $Ctsk^{Cre/+}$ and $Rosa26^{CAG-lox-stop-lox-tdTomato}$ BMMC were induced to

differentiate into osteoclasts. In this experimental setting, the fluorescent protein tdTomato (Tom)

is only expressed in cells resulting from the fusion of at least one $Ctsk^{Cre/+}$ and one $Rosa26^{CAG-lox-}$

$^{stop-lox-tdTomato}$ pre-osteoclasts (**Fig.6E**). We then monitored using time-lapse microscopy the

appearance of fluorescent Tom+ positive cells in the presence or absence of recGAS6. As shown

in Figure 6F, Tom+ cells begin to appear in osteoclast culture at day 4 and their number increases

considerably at day 5 and 6, validating our approach. Interestingly, the number of Tom+ cells

was significantly higher in the presence of recGAS6 in the culture media even with the lowest

tested dose of 50 ng/mL, suggesting that GAS6 signaling promotes osteoclast fusion.

GAS6 promotes osteoclast formation and bone resorption in vivo

In addition to GAS6, protein S (Pros1) was shown to activate TAM receptors [45]. Our data

indicate that osteoblasts also express Pros1 (**Fig. 4A**) and that both AXL and MerTK are present

in pre-osteoclasts (**Fig. 4B**). This redundancy in the TAM signaling system prevented us from

testing its importance in osteoclast formation through a genetic loss-of-function model. Instead, a

gain-of-function approach was employed to determine the impact of raising GAS6 on osteoclastic

bone resorption in vivo. We took advantage of a previously described transgenic mouse line

expressing GAS6 under the human *ApoE* promoter [40]. In these *ApoE-Gas6*^{Tg} mice, GAS6 is

produced by hepatocytes and therefore fully carboxylated and secreted in the circulation, leading

to a 4-fold increase in active GAS6 in the serum (**Fig.6A**). Since the skeleton, and particularly the

bone marrow cavity is highly vascularized, we reasoned that this rise in circulating GAS6 will

translate into an increased concentration of this protein in the bone marrow space, in which

osteoclast differentiation takes place. Accordingly, the concentration of GAS6 in the bone

marrow space was increased by 2-fold in *ApoE-Gas6^{Tg}* mice (**Fig.7B**). We next investigated bone

density and remodeling in this GAS6 gain-of-function mouse model.

Histology and µCT revealed a reduced trabecular bone volume (BV/TV) in vertebrae and femur

of 6-month-old *ApoE-Gas6^{Tg}* male mice compared to WT littermates (**Fig.7C-F**). The µCT

analyses showed that trabecular bone surface density (BS/TV), was also significantly reduced in

the same animals (**Fig. 7G**). Trabecular spacing (Tb.S) was significantly increased and trabecular

number (Tb.N) and connectivity density (Conn.D) significantly decreased, while trabecular

thickness was unaffected in *ApoE-Gas6^{Tg}* mice (**Fig. 7H**). Static and dynamic bone

histomorphometry indicated that *ApoE-Gas6^{Tg}* mice were characterized by a significant increase

in both the number and surface of osteoclasts in the trabecular bone (**Fig. 7I-J**). Importantly,

bone formation parameters were unaffected in the same animals (**Supplementary Figure 3**).

Altogether, these data indicate that increased concentration of GAS6 in bone is sufficient to

stimulate the formation of osteoclast in vivo, resulting in reduced bone density.

Discussion

Whether VK supplementation has any beneficial impact on skeletal health remains unclear since interventional studies in humans have led to conflicting results [22]. Our data suggest that VK-dependent carboxylation in osteoblasts is not required for normal bone formation and mineralization by this cell type. Instead, we uncover that γ-carboxylation, through GAS6, impacts the capacity of osteoblasts to regulate osteoclasts formation. Taken at face value, these findings suggest that VK should negatively regulate bone mass, potentially explaining the limited impact of VK supplementation on bone density in humans. However, osteoclastic bone resorption also plays a critical role in bone turnover and repair of microdamage to the bone structure [46]. Consequently, the production of active γ-carboxylated GAS6 by osteoblasts may be important for the fine tuning of osteoclast activity during bone repair processes, thereby promoting the maintenance of a healthy skeleton and preventing fractures.

Vitamin K-dependent carboxylation in osteoblast may influence the activity of several Gla proteins, including osteocalcin and MGP. The conclusion that γ-carboxylation in osteoblasts impact bone resorption through GAS6 is supported by several lines of evidence. First, the bone phenotype of the *ApoE-Gas6^{Tg}* mice, a GAS6 gain-of-function model, is the mirror image of the one observed in *Ggcx$^{fl/fl}$;OCN-Cre* mice. Second, in line with previous reports, we show that GAS6 is strongly expressed in osteoblasts [47,48]. Third, recombinant carboxylated GAS6 promotes the formation of larger osteoclasts in culture, while pharmacological inhibition of its receptors reduces osteoclastogenesis. Fourth, osteocalcin-deficient osteoblasts fully support osteoclast formation, while *Mgp*-deficient mice have normal bone resorption [38].

In line with our findings, a recent study reported that whole body inactivation of MerTK in mice

results in higher bone density with concomitant reduction in bone resorption [49]. TAM receptors

signaling may also play a role in osteoblasts function since it was reported that MerTK

inactivation specifically in osteoblasts using the *Col1a1-Cre* driver caused increased bone

formation and bone density, without affecting bone resorption [48]. Importantly, *Ggcx$^{fl/fl}$;OCN-Cre*

mice did not show this increase in bone formation, suggesting that osteoblasts-derived

carboxylated GAS6 does not suppress osteoblast activity through MerTK. Both protein S and

GAS6 can activate MerTK, while AXL is activated exclusively by GAS6 [45]. According to our

own data and previous reports [48,50], osteoblasts express protein S. We also found that pre-

osteoclast and differentiated osteoclasts express comparable level of AXL and MerTK. Due to

this redundancy in ligands and receptors, assessing the role of the Pros1/GAS6/TAM axis in bone

resorption using loss-of-function mouse models will likely require the generation of cell-specific

double knockout mice.

In our primary culture assays, low to intermediate doses of GAS6 (i.e., 25 to 100 ng/mL)

significantly increased osteoclasts number and surface at 5 and 6 days of culture. The same

concentration also increased the average number of nuclei per osteoclast. However, dose of 50 to

100 ng/mL had limited impact on the expression of the osteoclast differentiation program.

Together with the results of our fusion assay, these data suggest that GAS6 does not regulate

osteoclast differentiation but rather promotes the fusion of pre-osteoclasts.

To optimally activate TAM receptors, GAS6 requires its Gla domain, even though this domain is

not required for binding to the ectodomain of TAM [45]. In the context of its myeloid function,

GAS6 Gla domain binds to phosphatidylserines (PtdSer) exposed at the surface of a phagocytic

target (e.g., an apoptotic cell), and although the precise mechanism is not fully understood, this

PtdSer binding induces a conformational change in GAS6 that results in the activation of the

TAM receptor on the phagocytic cells [51]. Interestingly, recent studies have established that PtdSer

are exposed on the outer leaflet of the plasma membrane during pre-osteoclast differentiation and

that this exposure mediates the fusion of osteoclast precursors [52-54]. Therefore, GAS6 may be

contributing to the fusion process by binding PtdSer at the surface of one pre-osteoclast while

activating TAM receptors at the surface of another pre-osteoclast. Future studies will focus on

understanding how the signal downstream of TAM in pre-osteoclast contributes to the fusion.

In conclusion, this work uncovers a new vitamin K-dependent carboxylation pathway in

osteoblasts involved in the regulation of osteoclast formation.

Materials and Methods

Animals

All the mouse strains used in this study were backcrossed or generated on a pure C57BL6/J genetic background. The generation of $Ggcx^{fl/fl}$;OCN-Cre mice was previously described [33]. Briefly, $Ggcx^{flox/flox}$ mice were bred with $Osteocalcin$-Cre (OCN-Cre) mice, which express the Cre recombinase in mature osteoblasts only [32]. We previously established that $Ggcx$ was specifically and efficiently inactivated in osteoblasts in $Ggcx^{fl/fl}$;OCN-Cre mice [33]. The generation and characterization of $ApoE$-$Gas6^{Tg}$ mice expressing mouse $Gas6$ under the control of the liver specific human APOE promoter and hepatic enhancers was also previously reported [40]. The $tdTomato^{flox/flox}$ strain expressing the tdTomato reporter gene from the $Rosa26$ locus was obtained from the Jackson Laboratory ($B6.Cg$-$Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze/J}$; stock 007914). The $Ctsk$-Cre ($Ctsk^{tm1(cre)Ska}$) strain was previously described [55]. Osteocalcin-deficient mice (Ocn-/-) in which both osteocalcin genes ($Bglap1/Ocn1$ and $Bglap2/Ocn2$) are inactivated were previously reported [29]. C57BL6/J newborn or adult mice (Jackson Laboratory; strain 000664) were used as a source of primary wildtype (WT) osteoblasts and bone marrow cells. All strains were maintained in an IRCM specific pathogen-free animal facility under 12-hour dark/12-hour light cycles. Mice were fed ad libitum a normal chow diet (Teklad global 18% protein rodent diet; 2918; Envigo). Male mice were used in all experiments. All animal use complied with the guideline of the Canadian Committee for Animal Protection (CCAP) and was approved by IRCM institutional animal care committee.

Biochemical measurements

Serum carboxylated (Gla) and total osteocalcin were assessed using previously described ELISA assays [56]. Serum C-terminal telopeptide of collagen type I (CTX) was measured using a

commercially available assay (RatLaps; IDS, AC-06F1). Serum GAS6 was quantified using a

mouse GAS6 ELISA (DuoSet ELISA, R&D Systems, DY986) as we previously described [40]. To

measure GAS6 in bone marrow, the bone marrow of one femur per mouse was flushed using

1mL of PBS1X and the concentration of GAS6 quantified using the same ELISA assay. To

estimate the concentration of GAS6 in the BM, the amount of GAS6 per femur (i.e, GAS6

concentration $\times$ 1mL) was divided by 20µL, the estimated volume for a femur BM cavity of a 6-

month-old male mouse based on previous µCT data.

Bone histology and histomorphometry

Mice were injected intraperitoneally twice at a 3-day interval with 0.2 mL of a solution of calcein

(0.25% in a solution of 0.15M NaCl and 2% $NaHCO_3$). Following euthanasia, the skeletal tissue

was fixed in 4% formaldehyde in PBS 1X and processed for histomorphometry and micro-CT.

For non-decalcified histology, the vertebrae were embedded in methyl methacrylate resin (MMA)

as previously described [57] and sectioned (5 µm and 7µm thickness). Following deplastification,

the sections were stained with von Kossa and van Gieson (7µm), with Toluidine blue (5 µm) or

TRAP (5 µm). Bone histomorphometric analyses were completed using the OsteoMeasure

Analysis System (OsteoMetrics) connected to a light and fluorescent microscope (DM4000B

LED; Leica) with a 40$\times$ objective (HCX PL FLUOTAR 40$\times$; NA = 0.75).

Micro-CT

Ex vivo micro-computed tomography (µCT) scans were performed on harvested distal femurs

using the Skyscan 1272 system (Bruker, Kontich, Belgium) at 62 kV, 156 µA, and 6 µm isotropic

resolution with a 0.5 mm aluminum filter. Images were reconstructed with NRecon (beam

hardening correction: 50%; ring artifact reduction: 7) and analyzed using CTAn software (Bruker,

Skyscan). The trabecular bone region of interest (ROI) was defined 0.75 mm proximal to the

growth plate and extended 1 mm proximally. Morphometric parameters were calculated

following published guidelines for rodent bone analysis using μCT [58].

Production and Purification of Recombinant GAS6

Recombinant γ-carboxylated mouse GAS6 was produced as previously described [40]. Briefly,

HEK293 Flp-In T-REx cells (Thermo Fisher Scientific) stably transfected with a tetracycline-

inducible plasmid encoding mouse GAS6 with a C-terminal 6xHis tag were maintained in

DMEM (Wisent) supplemented with 10% fetal bovine serum (FBS; Sigma) and 200 μg/mL

hygromycin for selection. Cells were expanded and the expression of GAS6 induced with

1 μg/mL tetracycline in presence of 22 μM vitamin K_1 (Millipore Sigma, V3501) in serum-free

DMEM for 36 hours. The culture supernatant was collected, centrifuged, and filtered (0.22 μm)

to remove cells and debris. Recombinant GAS6 was purified by nickel affinity chromatography

using a HisTrap HP column (Cytiva) connected to an ÄKTAprime FPLC system (GE

Healthcare). Elution was performed using an imidazole-containing buffer, and the purified

protein was dialyzed against PBS. Protein purity and concentration were assessed by SDS-PAGE

followed by Coomassie staining (see **Supplemental Figure 2**).

Osteoblast cultures

Mouse calvaria pre-osteoblasts were isolated from 3-day-old pups and cultured as previously

described [33]. Briefly, dissected calvarias were digested at 37°C twice for 10 minutes and twice for

30 min with a solution of collagenase type 2 (0.1 mg/mL; Worthington Biochemical Corporation,

LS004176) containing 0.25% trypsin. Only the last two digestions were kept and cultured in

alpha modified minimum essential medium (α-MEM, Sigma, M0644) supplemented with 10%

fetal bovine serum (FBS; Sigma) and 1% penicillin-streptomycin (Wisent) to obtain proliferating

pre-osteoblasts (pre-OB). Differentiation into mineralizing osteoblasts (OB) was induced by

culturing these cells in the same media supplemented with 5mM β-glycerophosphate and 100

µg/mL L-ascorbic acid for 21 days. Media was changed every 2 days. To generate control and

Ggcx$^{-/-}$ pre-osteoblasts ex vivo, *Ggcx$^{flox/flox}$* cells were transduced with either GFP- or Cre-

expressing adenovirus (University of Iowa) at a MOI of 200 [33].

Bone marrow monocytes and osteoclasts cultures

Primary bone marrow cells were isolated from the femurs and tibias of 6- to 10-week-old

C57BL6/J mice. Bones were cleaned of soft tissue and flushed with α-MEM supplemented with

10% FBS and 1% penicillin-streptomycin using a 25G needle. Bone marrow-derived monocytes

(BMMC) were obtained by culturing the marrow cell suspension in non-adherent 10 cm dishes

in M-CSF–rich medium (20% L929 cell-conditioned medium; ATCC) for 5 days. Non-adherent

cells were removed by vigorous washing with PBS, and adherent cells were harvested by

incubation with 0.02% EDTA in PBS.

For osteoclast-like cell (OCL) differentiation, BMMCs were seeded at 2×10^4 cells per well in

24-well plates and cultured in medium supplemented with 20 ng/mL RANKL and 10 ng/mL M-

CSF, with or without recombinant γ-carboxylated GAS6 at various concentrations (25-200

ng/mL). Multinucleated mature osteoclasts were visualized by TRAP staining [59]. In selected

experiments, nuclei were stained with DAPI, and both brightfield and fluorescence images were

acquired (DM4000B LED; Leica) with a 5× objective (HCX PL FLUOTAR 5×; AP = 0.15). The

number of nuclei per osteoclast was quantified using ImageJ. In co-culture assays, 6×10^6 bone

marrow cells were added to confluent osteoblasts cultures in 12 well plates in αMEM

supplemented with 10% FBS, prostaglandin E_2 (PGE$_2$; 10^{-6} M) and 1,25 vitamin D_3 (VitD$_3$; 10^{-8}

M). In some experiments, TAM receptors inhibitors R428 or LDC1267 (Selleckchem, S2841 and

S7638) were added to the cultures at the indicated concentrations. After 8 days of coculture, the

cells were fixed and TRAP staining performed. Osteoclasts culture pictures were acquired using a

SteREO Discovery.V12 microscope (Zeiss) using Zen software. Osteoclasts number and surface

area were quantified from TRAP-stained cell cultures using ImageJ software.

Fusion assay

An equal number of *tdTomato*$^{flox/flox}$ and *Ctsk-Cre* BMMCs (1×10^4 cells each per well) were

seeded in 24-well plates and cultured in medium supplemented with 20 ng/mL RANKL and 10

ng/mL M-CSF, with or without recombinant γ-carboxylated GAS6 at various concentrations.

Plates were imaged using the Incucyte Live-Cell Analysis System (Sartorius). Fluorescence

(tdTomato) and phase contrast images were acquired every 24 h starting from day 3 of culture

using a 10× objective (16 fields per well). Images were analyzed using ImageJ and each

tdTomato-positive (red) cell with 2 or more nuclei was counted as a fusion event.

Gene expression

Total RNA was extracted from cells or tissues as previously [60]. mRNA was reverse transcribed

using M-MLV as previously reported [61]. Relative gene expression was measured by quantitative

PCR using PowerUp SYBR Green Master Mix (A25741; Applied Biosystems) with gene-

specific primers (Supplemental Table 1) and ViiA7 Real-Time PCR System (Applied

Biosystems).

Western blot

Proteins from cells or tissues were extracted with a lysis buffer containing 20mM Tris-HCl (pH

7.5), 150mM NaCl, 1mM EDTA (pH 8.0), 1mM EGTA, 2.5mM NaPyrophosphate, 1mM β-

glycerophosphate, 10mM NaF, 1% Triton, 1 mM vanadate, 1mM phenylmethylsulfonyl fluoride

(PMSF) and protease inhibitors (Roche Diagnostics, 4693132001). Proteins were detected by

Western blot with the indicated primary antibodies overnight at 4°C and quantified by

densitometry analyses using Image Lab software (Bio-Rad Laboratories). The antibodies used are

listed in Supplemental Table 2.

Statistical Analysis

Statistical analyses were performed with GraphPad Prism version 10.4.0. Results are given as

mean ± SEM. Unpaired two-tailed Student's t test was used to compare two groups. For

experiments involving multiple groups, one- or two-way ANOVA, followed by Bonferroni's

multiple comparisons test was used. In all figures, $* P < 0.05$; $** P < 0.01$; $*** P < 0.001$. All

experiments were repeated at least 3 times or performed on at least 3 independent animals.

Data and Resource Availability

The data sets generated and/or analyzed during the current study are available from the

corresponding author upon reasonable request. The $ApoE\text{-}Gas6^{Tg}$ transgenic and the $Ggcx^{flox/flox}$

mouse lines, the recombinant GAS6 expressing cell line, the phosphorylated AXL (Y702)

antibody and the mouse VKORC1 antibody used during the current study are available from the

corresponding author upon reasonable request.

Acknowledgements

We thank S. Kato and T. Clemens for providing *Ctsk-Cre* and *OCN*-Cre mice respectively. This work was supported by funding from the Fonds de Recherche du Québec–Santé (to M.F.), Canada Research Chairs program (to M.F.) and Canadian Institutes of Health Research grants PJT-159534 (to M.F.). B.AR. received a postdoctoral fellowship from the Fonds de Recherche du Québec–Santé.

Conflict of interests

No potential conflicts of interest relevant to this article were reported.

Contributions

M.F. and M.P. assembled the figures. M.F. wrote the manuscript with suggestions from the other authors. M.P., M.F., D.N.T.P., J.L., B.A.R. and Y.W.K., collected and analyzed data. M.F. conceived and designed the project and provided supervision to M.P., D.N.T.P., J.L., B.A.R. and Y.W.K. A.G.A.A. and M.M. collected and analyzed the μCT data. M.F. is the guarantor of this work and, as such, had full access to all the data in the study and takes responsibility for the integrity of data and the accuracy of data analysis.

References

Morris, J. A. *et al.* An atlas of genetic influences on osteoporosis in humans and mice. *Nat Genet* **51**, 258-266, doi:10.1038/s41588-018-0302-x (2019).

Zhu, X., Bai, W. & Zheng, H. Twelve years of GWAS discoveries for osteoporosis and related traits: advances, challenges and applications. *Bone Res* **9**, 23, doi:10.1038/s41413-021-00143-3 (2021).

Moller, A. M. J. *et al.* Aging and menopause reprogram osteoclast precursors for aggressive bone resorption. *Bone Res* **8**, 27, doi:10.1038/s41413-020-0102-7 (2020).

Wilson-Barnes, S. L., Lanham-New, S. A. & Lambert, H. Modifiable risk factors for bone health & fragility fractures. *Best Pract Res Clin Rheumatol* **36**, 101758, doi:10.1016/j.berh.2022.101758 (2022).

Nieves, J. W. Osteoporosis: the role of micronutrients. *Am J Clin Nutr* **81**, 1232S-1239S, doi:10.1093/ajcn/81.5.1232 (2005).

Veldurthy, V. *et al.* Vitamin D, calcium homeostasis and aging. *Bone Res* **4**, 16041, doi:10.1038/boneres.2016.41 (2016).

Brzezinska, O., Lukasik, Z., Makowska, J. & Walczak, K. Role of Vitamin C in Osteoporosis Development and Treatment-A Literature Review. *Nutrients* **12**, doi:10.3390/nu12082394 (2020).

Clements, M. *et al.* A 2-Year Randomized Controlled Trial With Low-Dose B-Vitamin Supplementation Shows Benefits on Bone Mineral Density in Adults With Lower B12 Status. *J Bone Miner Res* **37**, 2443-2455, doi:10.1002/jbmr.4709 (2022).

Tucker, K. L. *et al.* Low plasma vitamin B12 is associated with lower BMD: the Framingham Osteoporosis Study. *J Bone Miner Res* **20**, 152-158, doi:10.1359/JBMR.041018 (2005).

Booth, S. L. *et al.* Vitamin K intake and bone mineral density in women and men. *Am J Clin Nutr* **77**, 512-516 (2003).

Booth, S. L. *et al.* Associations between vitamin K biochemical measures and bone mineral density in men and women. *J Clin Endocrinol Metab* **89**, 4904-4909, doi:10.1210/jc.2003-031673 (2004).

Booth, S. L. *et al.* Dietary vitamin K intakes are associated with hip fracture but not with bone mineral density in elderly men and women. *Am J Clin Nutr* **71**, 1201-1208, doi:10.1093/ajcn/71.5.1201 (2000).

Feskanich, D. *et al.* Vitamin K intake and hip fractures in women: a prospective study. *Am J Clin Nutr* **69**, 74-79 (1999).

Hao, G. *et al.* Vitamin K intake and the risk of fractures: A meta-analysis. *Medicine (Baltimore)* **96**, e6725, doi:10.1097/MD.0000000000006725 (2017).

Hamidi, M. S., Gajic-Veljanoski, O. & Cheung, A. M. Vitamin K and bone health. *Journal of clinical densitometry : the official journal of the International Society for Clinical Densitometry* **16**, 409-413, doi:10.1016/j.jocd.2013.08.017 (2013).

16   Thijssen, H. H. & Drittij-Reijnders, M. J. Vitamin K distribution in rat tissues: dietary
phylloquinone is a source of tissue menaquinone-4. *The British journal of nutrition* **72**,
415-425 (1994).

17   Ronden, J. E., Thijssen, H. H. & Vermeer, C. Tissue distribution of K-vitamers under
different nutritional regimens in the rat. *Biochim Biophys Acta* **1379**, 16-22 (1998).

18   Fang, Y., Hu, C., Tao, X., Wan, Y. & Tao, F. Effect of vitamin K on bone mineral density:
a meta-analysis of randomized controlled trials. *J Bone Miner Metab* **30**, 60-68,
doi:10.1007/s00774-011-0287-3 (2012).

19   Cockayne, S. *et al.* Vitamin K and the prevention of fractures: systematic review and
meta-analysis of randomized controlled trials. *Archives of internal medicine* **166**, 1256-
1261, doi:10.1001/archinte.166.12.1256 (2006).

20   Inoue, T. *et al.* Randomized controlled study on the prevention of osteoporotic fractures
(OF study): a phase IV clinical study of 15-mg menatetrenone capsules. *J Bone Miner*
Metab **27**, 66-75, doi:10.1007/s00774-008-0008-8 (2009).

21   Cheung, A. M. *et al.* Vitamin K supplementation in postmenopausal women with
osteopenia (ECKO trial): a randomized controlled trial. *PLoS medicine* **5**, e196,
doi:10.1371/journal.pmed.0050196 (2008).

22   Dupuy, M. *et al.* Vitamin K: Metabolism, Genetic Influences, and Chronic Disease
Outcomes. *Food Sci Nutr* **13**, e70431, doi:10.1002/fsn3.70431 (2025).

23   Zhou, M., Han, S., Zhang, W. & Wu, D. Efficacy and safety of vitamin K2 for
postmenopausal women with osteoporosis at a long-term follow-up: meta-analysis and
systematic review. *J Bone Miner Metab* **40**, 763-772, doi:10.1007/s00774-022-01342-6
(2022).

24   Salma *et al.* Effect of Vitamin K on Bone Mineral Density and Fracture Risk in Adults:
Systematic Review and Meta-Analysis. *Biomedicines* **10**,
doi:10.3390/biomedicines10051048 (2022).

25   Lacombe, J. & Ferron, M. Vitamin K-dependent carboxylation in beta-cells and diabetes.
Trends Endocrinol Metab **35**, 661-673, doi:10.1016/j.tem.2024.02.006 (2024).

26   Berkner, K. L. & Runge, K. W. Vitamin K-Dependent Protein Activation: Normal
Gamma-Glutamyl Carboxylation and Disruption in Disease. *International journal of*
molecular sciences **23**, doi:10.3390/ijms23105759 (2022).

27   Hauschka, P. V., Lian, J. B. & Gallop, P. M. Direct identification of the calcium-binding
amino acid, gamma-carboxyglutamate, in mineralized tissue. *Proc Natl Acad Sci U S A*
72, 3925-3929 (1975).

28   Lambert, L. J. *et al.* Increased trabecular bone and improved biomechanics in an
osteocalcin-null rat model created by CRISPR/Cas9 technology. *Disease models &*
mechanisms **9**, 1169-1179, doi:10.1242/dmm.025247 (2016).

29   Ducy, P. *et al.* Increased bone formation in osteocalcin-deficient mice. *Nature* **382**, 448-
452 (1996).

30   Diegel, C. R. *et al.* An osteocalcin-deficient mouse strain without endocrine
abnormalities. *PLoS genetics* **16**, e1008361, doi:10.1371/journal.pgen.1008361 (2020).

31 Moriishi, T. *et al.* Osteocalcin is necessary for the alignment of apatite crystallites, but not
glucose metabolism, testosterone synthesis, or muscle mass. *PLoS genetics* **16**, e1008586,
doi:10.1371/journal.pgen.1008586 (2020).

32 Zhang, M. *et al.* Osteoblast-specific knockout of the insulin-like growth factor (IGF)
receptor gene reveals an essential role of IGF signaling in bone matrix mineralization. *J*
Biol Chem **277**, 44005-44012 (2002).

33 Ferron, M., Lacombe, J., Germain, A., Oury, F. & Karsenty, G. GGCX and VKORC1
inhibit osteocalcin endocrine functions. *J Cell Biol* **208**, 761-776,
doi:10.1083/jcb.201409111 (2015).

34 Coutu, D. L. *et al.* Periostin, a member of a novel family of vitamin K-dependent
proteins, is expressed by mesenchymal stromal cells. *J Biol Chem* **283**, 17991-18001
(2008).

35 Barone, L. M. *et al.* Developmental expression and hormonal regulation of the rat matrix
Gla protein (MGP) gene in chondrogenesis and osteogenesis. *J Cell Biochem* **46**, 351-
365, doi:10.1002/jcb.240460410 (1991).

36 Ducy, P. & Karsenty, G. Two distinct osteoblast-specific cis-acting elements control
expression of a mouse osteocalcin gene. *Mol Cell Biol* **15**, 1858-1869 (1995).

37 Gerbaix, M., Vico, L., Ferrari, S. L. & Bonnet, N. Periostin expression contributes to
cortical bone loss during unloading. *Bone* **71**, 94-100, doi:10.1016/j.bone.2014.10.011
(2015).

38 Marulanda, J., Gao, C., Roman, H., Henderson, J. E. & Murshed, M. Prevention of
arterial calcification corrects the low bone mass phenotype in MGP-deficient mice. *Bone*
57, 499-508, doi:10.1016/j.bone.2013.08.021 (2013).

39 Lemke, G. Biology of the TAM receptors. *Cold Spring Harb Perspect Biol* **5**, a009076,
doi:10.1101/cshperspect.a009076 (2013).

40 Schott, C. *et al.* GAS6 and AXL Promote Insulin Resistance by Rewiring Insulin
Signaling and Increasing Insulin Receptor Trafficking to Endosomes. *Diabetes* **73**, 1648-
1661, doi:10.2337/db23-0802 (2024).

41 Goruppi, S., Ruaro, E., Varnum, B. & Schneider, C. Requirement of phosphatidylinositol
3-kinase-dependent pathway and Src for Gas6-Axl mitogenic and survival activities in
NIH 3T3 fibroblasts. *Mol Cell Biol* **17**, 4442-4453 (1997).

42 Pao-Chun, L., Chan, P. M., Chan, W. & Manser, E. Cytoplasmic ACK1 interaction with
multiple receptor tyrosine kinases is mediated by Grb2: an analysis of ACK1 effects on
Axl signaling. *J Biol Chem* **284**, 34954-34963, doi:10.1074/jbc.M109.072660 (2009).

43 Paolino, M. *et al.* The E3 ligase Cbl-b and TAM receptors regulate cancer metastasis via
natural killer cells. *Nature* **507**, 508-512, doi:10.1038/nature12998 (2014).

44 Holland, S. J. *et al.* R428, a selective small molecule inhibitor of Axl kinase, blocks tumor
spread and prolongs survival in models of metastatic breast cancer. *Cancer research* **70**,
1544-1554, doi:10.1158/0008-5472.CAN-09-2997 (2010).

45 Lew, E. D. *et al.* Differential TAM receptor-ligand-phospholipid interactions delimit
differential TAM bioactivities. *eLife* **3**, doi:10.7554/eLife.03385 (2014).

46   Schindeler, A., McDonald, M. M., Bokko, P. & Little, D. G. Bone remodeling during
fracture repair: The cellular picture. *Semin Cell Dev Biol* **19**, 459-466,
doi:10.1016/j.semcdb.2008.07.004 (2008).

47   Shiozawa, Y. *et al.* GAS6/AXL axis regulates prostate cancer invasion, proliferation, and
survival in the bone marrow niche. *Neoplasia* **12**, 116-127, doi:10.1593/neo.91384
(2010).

48   Engelmann, J. *et al.* Regulation of bone homeostasis by MERTK and TYRO3. *Nat*
Commun **13**, 7689, doi:10.1038/s41467-022-33938-x (2022).

49   Ryu, K. Y. *et al.* Mer tyrosine kinase regulates bone metabolism, and its deficiency
partially ameliorates periodontitis- and ovariectomy-induced bone loss in mice. *JBMR*
Plus **8**, ziad014, doi:10.1093/jbmrpl/ziad014 (2024).

50   Maillard, C., Berruyer, M., Serre, C. M., Dechavanne, M. & Delmas, P. D. Protein-S, a
vitamin K-dependent protein, is a bone matrix component synthesized and secreted by
osteoblasts. *Endocrinology* **130**, 1599-1604, doi:10.1210/endo.130.3.1531628 (1992).

51   Lemke, G. Phosphatidylserine Is the Signal for TAM Receptors and Their Ligands. *Trends*
in biochemical sciences **42**, 738-748, doi:10.1016/j.tibs.2017.06.004 (2017).

52   Krishnacoumar, B. *et al.* Caspase-8 promotes scramblase-mediated phosphatidylserine
exposure and fusion of osteoclast precursors. *Bone Res* **12**, 40, doi:10.1038/s41413-024-
00338-4 (2024).

53   Verma, S. K. *et al.* Cell-surface phosphatidylserine regulates osteoclast precursor fusion. *J*
Biol Chem **293**, 254-270, doi:10.1074/jbc.M117.809681 (2018).

54   Whitlock, J. M. *et al.* Cell surface-bound La protein regulates the cell fusion stage of
osteoclastogenesis. *Nat Commun* **14**, 616, doi:10.1038/s41467-023-36168-x (2023).

55   Nakamura, T. *et al.* Estrogen prevents bone loss via estrogen receptor alpha and induction
of Fas ligand in osteoclasts. *Cell* **130**, 811-823 (2007).

56   Ferron, M., Wei, J., Yoshizawa, T., Ducy, P. & Karsenty, G. An ELISA-based method to
quantify osteocalcin carboxylation in mice. *Biochem Biophys Res Commun* **397**, 691-696
(2010).

57   Chappard, D., Palle, S., Alexandre, C., Vico, L. & Riffat, G. Bone embedding in pure
methyl methacrylate at low temperature preserves enzyme activities. *Acta Histochem* **81**,
183-190 (1987).

58   Bouxsein, M. L. *et al.* Guidelines for assessment of bone microstructure in rodents using
micro-computed tomography. *J Bone Miner Res* **25**, 1468-1486, doi:10.1002/jbmr.141
(2010).

59   Ferron, M. *et al.* Inositol polyphosphate 4-phosphatase B as a regulator of bone mass in
mice and humans. *Cell Metab* **14**, 466-477 (2011).

60   Chomczynski, P. & Sacchi, N. The single-step method of RNA isolation by acid
guanidinium thiocyanate-phenol-chloroform extraction: twenty-something years on.
Nature protocols **1**, 581-585, doi:10.1038/nprot.2006.83 (2006).

61   Lacombe, J. *et al.* Vitamin K-dependent carboxylation regulates Ca(2+) flux and
adaptation to metabolic stress in beta cells. *Cell Rep* **42**, 112500,
doi:10.1016/j.celrep.2023.112500 (2023).

 FIGURES LEGENDS

Figure 1. Increased bone mass in mice lacking γ-carboxylation in osteoblasts. (A) Gene

expression analysis by qPCR of *Ggcx* and *Vkorc1* in bone marrow derived monocytes (BMMC),

osteoclasts (OCL), proliferating pre-osteoblasts (pre-OB) and mineralized osteoblast (OB)

cultures (n=2-3). **(B)** Protein expression in liver (Liv) and bone cells by Western blot. GGCX was

analyzed on a 7.5% SDS Tris Glycine gel using 20μg of extracts, while VKORC1 was resolved

on a 10% SDS Tris Tricine gel using 10μg of extracts. **(C-I)** Six-month-old *Ggcx^{ff}* and

Ggcx^{ff};OCN-Cre male mice were analyzed (n=7-10). **(C)** Representative pictures of sections from

lumbar vertebrae stained with von Kossa and van Gieson. **(D)** Quantification of trabecular bone

volume over tissue volume (BV/TV) from the L4 and L5 lumbar vertebrae sections. **(E)**

Representative μCT images of the distal femur trabecular bones. **(F)** Quantification of trabecular

bone volume (BV/TV) and trabecular bone surface density (BS/TV) from the μCT data. **(G)**

Trabecular bone μCT derived data. Tb.Sp., Tb.N., and Tb.Th., trabecular spacing, number, and

thickness respectively; Conn.Dn., connectivity density. Unpaired, 2-tailed Student's t test was

used in (D), (F), and (G). ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns : non-significant.

Figure 2. Reduced osteoclasts number and surface in *Ggcx^{ff};OCN-Cre* mice. (A-H) Bone

histomorphometry analysis of lumbar vertebrae in six-month-old *Ggcx^{ff}* and *Ggcx^{ff};OCN-Cre*

male mice (n=6-10). **(A)** Representative pictures of calcein double labeling and toluidine blue

staining. **(B)** Mineral apposition rate (MAR). **(C)** Bone formation rate over bone surface

(BFR/BS). **(D)** Number of osteoblasts per bone perimeter (N.Ob/B.Pm). **(E)** Osteoblasts surface

over bone surface (Ob.S/BS). **(F)** Representative pictures of TRAP staining. **(G)** Number of

osteoclasts per bone perimeter (N.Oc/BPm). **(H)** Osteoclast surface over bone surface (Oc.S/BS).

(I) Fasting serum CTx levels (n=12-17). Results represent the mean ± SEM. Unpaired, 2-tailed

Student's t test was used in (B-E) and (G-I). **$p < 0.01$, *$p < 0.05$, ns : non-significant.

Figure 3. *Ggcx* inactivation impairs the ability of osteoblasts to support osteoclastogenesis

ex vivo. (A) Representative TRAP staining of osteoblasts (OB) and bone marrow cells (BM) co-

cultures at day 8 in the presence of prostaglandin E_2 (PGE$_2$; 10^{-6} M) and 1,25 vitamin D_3 (VitD$_3$;

10^{-8} M). *Ggcx^{ff}* osteoblasts were transduced with either Ad-GFP (control) or Ad-Cre (knockout)

before the addition of the WT bone marrow cells. **(B)** Quantification of the number of TRAP+

osteoclasts per well (Nb of OCL/well) (n=3). **(C)** Representative TRAP staining of osteoblasts

and bone marrow cells co-cultures at day 8. Control (*Ocn+/+*) or osteocalcin-deficient (*Ocn-/-*)

osteoblasts were cultured with WT bone marrow cells. **(D)** Quantification of the number of

TRAP+ osteoclasts per well (Nb of OCL/well) (n=3). **(E)** Gene expression analysis by qPCR in

Ggcx^{ff} + Ad-GFP and *Ggcx^{ff}* + Ad-Cre osteoblasts cultured in presence (+) or absence (-) of PGE_2

and $VitD_3$. Results represent the mean ± SEM. Unpaired, 2-tailed Student's t test was used in (B)

and (D). Two-way ANOVA with Bonferroni's post tests was used in (E). ***$p < 0.001$, **$p <$

0.01, *$p < 0.05$, ns : non-significant.

Figure 4. The γ-carboxylated protein GAS6 is expressed by osteoblasts and activates its

receptors on pre-osteoclasts. (A) Expression analysis by qPCR of genes encoding the known γ-

carboxylated proteins. *Gas6* is highlighted in red. **(B)** Gene expression analysis by qPCR of the

TAM receptors *Axl*, *Mertk* and *Tyro3* in bone marrow derived monocytes (d0: day 0) and in

differentiating osteoclasts cultures in the presence of RANKL and M-CSF (d2-d6: day 2 to 6).

(C) Western blot analysis of the phosphorylation (P) of AKT (S473) and TAM (Y702 in AXL

and Y753 in MerTK) in bone marrow derived monocytes (BMMC) serum starved for 3h and

treated with GAS6 (200ng/mL) for the indicated times. Total AKT, MerTK and AXL were used

as loading controls. **(D)** Quantification of P-AKT (S473), P-MerTK (Y753) and P-AXL (Y702)

normalized to the amount of total protein (n=2-3). Results represent the mean ± SEM.

Figure 5. Carboxylated GAS6 signaling promotes osteoclasts formation in culture. (A-B)

Representative TRAP staining (left) and quantification of the TRAP+ osteoclast area (right) in

WT osteoblasts (OB) and bone marrow cells (BM) co-cultures at day 8 in the presence of PGE_2

and $VitD_3$, with or without the TAM inhibitors LDC1267 (A) or R428 (B), at the indicated

concentrations. **(C-F)** Bone marrow derived monocytes (BMMC) were cultured in the presence

of RANKL (20 ng/mL) and M-CSF (10 ng/mL) with or without recombinant carboxylated GAS6

at the indicated concentrations for up to 6 days. **(C)** Representative TRAP staining at day 5 and 6

of differentiation. **(D)** Quantification of the TRAP+ osteoclast area at day 4, 5 and 6 of

differentiation. **(E)** Quantification of the number of TRAP+ multinucleated osteoclasts at day 4, 5

and 6 of differentiation. **(F)** Quantification of the number of nuclei per osteoclast at day 5 of

differentiation. Results represent the mean ± SEM. One-way ANOVA with Bonferroni's post tests

was used in (A-B) and (D-F). ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns : non-significant.

Figure 6. Carboxylated GAS6 impact on osteoclasts differentiation and fusion. (A-D) Gene

expression analysis by qPCR of osteoclasts differentiation markers *Acp5* (TRAP), *Clcn7*, *Ctsk*

and *Dcstamp*. Bone marrow derived monocytes (BMMC) were cultured in the presence of

RANKL (20 ng/mL) and M-CSF (10 ng/mL) with or without recombinant carboxylated GAS6 at

the indicated concentrations for 2, 4 and 6 days (n=4 per condition). **(E)** Schematic representation

of the assay used to assess the impact of GAS6 on pre-osteoclasts fusion in culture using a

conditionally activated tdTomato (Tom) reporter. **(F)** Representative pictures of live osteoclasts

cultures at the indicated time and concentration of recombinant GAS6. The stars indicate the

presence of fusion events (Tom+ cells) in the GAS6 conditions at Day 4. **(G)** Quantification of

the number of fusion events per 10 mm^2 at the indicated time of osteoclasts differentiation (n=16

fields per condition). Results represent the mean ± SEM. One-way ANOVA with Bonferroni's

post tests was used in (A-D) and (G). ***$p < 0.001$, **$p < 0.01$.

Figure 7. GAS6 promotes osteoclasts formation and bone resorption in vivo. (A-J) Six-

month-old *WT* (non-transgenic littermates) and *ApoE-Gas6^{Tg}* male mice were analyzed. **(A-B)**

GAS6 concentration in the serum (A) and bone marrow cavity (B) (n=4). **(C)** Representative

pictures of sections from lumbar vertebrae stained with von Kossa and van Gieson. **(D)**

Quantification of trabecular bone volume over tissue volume (BV/TV) from the L4 and L5

lumbar vertebrae sections (n=15-17). **(E-H)** μCT analysis of the distal femur trabecular bone

(n=12). **(E)** Representative μCT images. **(F)** Quantification of trabecular bone volume (BV/TV).

(G) Quantification of trabecular bone surface density (BS/TV). **(H)** Trabecular bone μCT derived

data. Tb.Sp., Tb.N., and Tb.Th., trabecular spacing, number, and thickness, respectively;

Conn.Dn., connectivity density. **(I-J)** Bone histomorphometry analysis of lumbar vertebrae in six-

month-old *WT* and *ApoE-Gas6^{Tg}* male (n=19-20). **(I)** Representative pictures of TRAP staining.

(J) Number of osteoclasts per bone perimeter (N.Oc/BPm) and osteoclasts surface over bone

surface (Oc.S/BS). Results represent the mean ± SEM. Unpaired, 2-tailed Student's t test was

used in (A,B, D, F, G and J). ***$p < 0.001$, **$p < 0.01$, *$p < 0.05$, ns : non-significant.

Figure 1

Figure 2

Figure 4

Figure 5

scale bar = 100μm

Figure 6

Figure 7