BeWo-derived extracellular vesicles downregulate IL-6Rα

expression via miRNAs on CD4+ T lymphocytes

Gábor Seregélyes[1], Bence Nagy[1], Árpád Ferenc Kovács[2,3], Nóra Fekete[1], Edit I Buzás[1,4,5],

Éva Pállinger[1]

[1] Institute of Genetics, Cell- and Immunobiology, Semmelweis University, Budapest,

Hungary

[2] Heart and Vascular Center, Semmelweis University, Budapest, Hungary

[3]Department of Pathology and Experimental Cancer Research, Semmelweis University,

Budapest, Hungary

[4]MTA-SE Immune-Proteogenomics Extracellular Vesicle Research Group, Budapest,

Hungary

[5] HCEMM-SE Extracellular Vesicle Research Group

Correspondence:

Éva Pállinger

1089 Budapest Üllői út 26, Hungary

Institute of Genetics, Cell and Immunobiology, Semmelweis University

pallinger.eva@semmelweis.hu

0036309008343

Abstract

Regulatory T lymphocytes are essential for maternal immunotolerance. Their de novo

differentiation in the placenta is regulated by local intercellular interactions involving primed

uterine immune cells, fetal syncytiotrophoblasts, and the cytokine environment. Trophoblast-

derived, HLA-G-positive extracellular vesicles (EVs) can bind to T lymphocytes, thereby

influencing their differentiation and cytokine production. Thus, these EVs play a role in

establishing and maintaining a tolerogenic environment. In our study, we used the BeWo

choriocarcinoma cell line to model the effects of trophoblast-derived EVs. Large EVs derived

from BeWo cells (BeWo-12.5K lEVs) reduce IL-6Rα expression on CD4+ T cells. This

modifies the IL-6 pathway by downregulating the transcription factors STAT3 and NFKB1,

and PIAS3, while upregulating STAT1. This may be caused by specific microRNAs

(miRNAs), such as hsa-mir-92a-3p, hsa-mir-520f-3p, and hsa-mir-25-3p. These microRNAs

are present in BeWo-12.5K lEVs and target the IL-6 pathway. BeWo-12.5K lEVs induce

phenotypic and functional changes in T cells, enhancing the ratio of CD4+/CD25+ T cells

that produce IL-10. Pregnancy-associated IL-6Rα downregulation has been demonstrated in

clinical samples. Significantly lower levels of IL-6Rα were detected on circulating

CD4+CD25+ T cells in healthy pregnant women than in healthy non-pregnant individuals.

This finding reflects the in vivo significance of our in vitro studies. Our studies suggest that

communication between maternal and fetal cells significantly influences the development and

maintenance of local T cell polarity. The microRNA content of HLA-G+ 12.5K lEVs appears

to be key to this process, as it alters the IL-6 pathway.

Keywords: human pregnancy, immune tolerance, CD4+ T cells, regulatory T cells,

extracellular vesicles, interleukin-6, IL-6Rα, IL-6 pathway, miRNA

Introduction

The adaptation of the maternal immune system to pregnancy is strictly regulated and can be characterized by immunological stages, or the "immune clock" [1]. Recognizing the semi-allogeneic fetus and developing maternal immune tolerance are prerequisites for a successful pregnancy. Although an inflammatory environment is necessary for embryo implantation, the immune system transitions to an anti-inflammatory state by the end of the first trimester [2]. The presence of regulatory T cells (Treg) and Th2 cells in the placenta is essential for fetal growth [3-6]. Alterations to this immune milieu can result in recurrent spontaneous abortion [7, 8]. The local cytokine environment determines the polarization of naïve CD4+ T helper (Th) cells into specific subsets, such as Th1, Th2, Th17, and Treg cells. Transforming growth factor beta (TGF-β) and interleukin 2 (IL-2) are critical for Treg differentiation. The final effects of TGF-β signaling are also highly influenced by surrounding cytokines. Proinflammatory cytokines, such as IL-1β, IL-6, and TNF-α, can neutralize the effects of TGF-β and favor Th17 development over Treg polarization [9, 10]. IL-6 is a Janus faced player during pregnancy [11]. Although IL-6 is present in the amniotic fluid and the placenta, already from the earliest period of pregnancy [12, 13], the complete role of it has not been clarified yet. According to our knowledge IL-6, as a pleiotropic cytokine contributes to the main physiological processes during pregnancy, including implantation, embryogenesis and parturition [14]. On the other hand, IL-6 directly acts on CD4+ Th cells and plays a role in the orchestration of Th cell differentiation and function [15, 16].

The tolerogenic immune environment at the feto-maternal interface is maintained through direct cell-to-cell interactions, soluble mediators, and extracellular vesicles (EVs) [17, 18]. The presence of   human leukocyte antigen-G (HLA-G) -positive trophoblast-derived EVs at the feto-maternal interface and in the maternal bloodstream is a classic feature of pregnancy [19 -21]. EVs are membrane-covered intercellular messengers that are

constitutively secreted by eukaryotic and prokaryotic cells [22]. Two types of EVs can be

distinguished according to their production: ectosomes and exosomes. However, available

isolation methods separate them based on size [23]. In this context, the following definitions

apply to EV populations: small EVs (sEVs), which have an average diameter under 200 nm

and are enriched in exosomes, and large EVs, which have an average diameter higher than

200 nm. The large EV fraction contains the "12.5K lEVs" (formerly known as iEVs or

microvesicles), which have been isolated and purified by centrifugation at 12,500 g. [24]

These EV subpopulations may differ in their membrane components, including proteins and

lipids, as well as in their cargo, such as nucleic acids (e.g., microRNAs, intact mRNAs, long

non-coding RNAs, and double-stranded DNA) [25]. Placental EVs are critical modulators of

the maternal immune system [26]. Because EVs are involved in many physiological and

pathological processes, they can serve as diagnostic and prognostic biomarkers for various

pathological conditions [27]. In our previous study, we found that both platelet- and

trophoblast-derived EVs target immune cells [28] and trophoblastic-derived 12.5K lEVs

induced Treg expansion is associated with the HSPE1 expression and the specific miRNA

cargo of EVs. [29]

In the present study, we demonstrate that trophoblast-derived EVs modulate the IL-6

sensitivity of Th cells by regulating their interleukin-6 receptor alpha (IL6Ra) expression via

miRNA, thereby playing an important role in local Treg cell polarization.

Material and methods

Patients and sample collection

Peripheral venous blood samples were collected from the median cubital vein of 37 healthy

pregnant women in their 18th week of pregnancy and 15 healthy non-pregnant women using

Vacutainer® Brand Plus Sodium Heparin Tubes or Vacutainer® EDTA Tubes from Becton

Dickinson (BD, San Jose, California, USA). The blood samples arrived at the laboratory

within two hours and were immediately processed. The pregnant women underwent routine

obstetrical control at the Department of Obstetrics and Gynecology at Semmelweis

University's Faculty of Medicine. An obstetrical history of previous spontaneous abortions,

the course of previous pregnancies, hypertension, gestational diabetes, and preeclampsia were

exclusion criteria. Samples from the healthy non-pregnant women were obtained from the

Hungarian National Blood Transfusion Service. The clinical data of the studied groups are

presented in Supplementary Table 1. The study was approved by the Ethics Committee of

Scientific Research of Hungary (ETT-TUKEB: 10147-4/2015/EKU (93/2014)).

Lymphocyte separation

Peripheral blood mononuclear cells (PBMCs) were isolated using density gradient

centrifugation with Histopaque® 1077 (Sigma-Aldrich, St. Louis, Missouri), and the

monocytes were removed by adherence.

Cell culture (BeWo cells and human lymphocyte co-culturing)

BeWo choriocarcinoma cell line (ATCC® CCL-98™) was obtained from the American Type

Culture Collection. The BeWo cells were cultured as described earlier [29].

To address the interactions between BeWo cells and lymphocytes we designed a co-culture

system: 2 x 10^5 BeWo cells and 1 x 10^6 lymphocytes were plated into the wells of a BD

Falcon™ 12-well Cell Culture Insert Companion Plates (BD San Jose, California, USA) in 2

mL culture media, for 24 hours. BD Falcon™ Cell Culture Inserts (with 1 µm pores) have

been used to study the effects of BeWo-derived EVs and soluble mediators on lymphocytes

(Supplementary Figure 1).

BeWo cells were tested regularly for mycoplasma infection with a PCR-based method

(Supplementary Figure 2A). We examined the viability of the co-cultured cells regularly

using a microscope. A flow cytometric analysis was performed at the end of the culturing

period. (Supplementary Figure 2B).

Flow cytometry was used to demonstrate the transport of EVs through 1 µm inserts.

(Supplementary Figure 2C).

Isolation of BeWo-derived large vesicles (12.5K lEVs), small-sized vesicles (sEVs) and

EV free supernatant by differential centrifugation

2×10^6 BeWo cells were seeded in 75 cm^2 tissue culture treated flasks in 10 mL of culture

media. Once the cells reached 80-90% confluence, we discarded the culture medium and

replaced it with 10 mL of FBS-free medium. Then, we cultured the cells for an additional 24

hours. We applied the differential centrifugation method for EV isolation, which yields

intermediate recovery and specificity. Next, BeWo cell-culture supernatant was centrifuged at

300 *g* for 5 minutes at room temperature (RT) to remove cell debris. The obtained

supernatant was centrifuged at 2 000 *g* for 15 minutes to remove the apoptotic body fraction.

The supernatant of the 2 000 *g* speed centrifugation was sedimented at 12 500 *g* for 20

minutes at 16°C (Z216 MK Microlite centrifuge, fixed angle 200.88 rotor, Hermle

Labortechnik GmbH, Wehingen, Germany) and subsequently washed with 0.20 µm filtered

phosphate-buffered saline (PBS) at 12 500 *g* for 15 minutes at 16°C. The supernatant of

12.5K centrifugation was further processed at 100 000 *g* for 70 minutes at 4°C (Optima

MAX-XP, fixed angle MLA-55 rotor, Beckman Coulter Inc, Brea, USA). A subsequent

washing step with filtered (0.20 µm pore size, Millipore) PBS at 100 000 *g* for 70 minutes at

4°C was done. The supernatant of the 100 000 *g* washing step was further centrifuged for

another 24 hours to obtain the EV free supernatant fraction. The obtained pellets enriched in

12.5K lEV fraction (12.5K pellet) or sEV fraction (100K pellet) were resuspended in 20-100

µL PBS or serum-free Ham's F12/K media depending on the downstream analysis

(Supplementary Figure 3).

EV characterization

Following the recommendations of the International Society for Extracellular Vesicles'

(ISEV) latest position paper [24], the isolated EV preparations were characterized by several

methods. (Supplementary Figure 4)

miRNA sequencing

Three biological replicates of BeWo-12.5K lEVs were isolated and resuspended in 12 µL of

DNase, RNase free water for miRNA sequencing. The RNA integrity was evaluated with

Agilent Tapestation and samples with RIN values ≥7 were further processed. For library

preparation multiplex Small RNA Libr. Prep Kit was used according to the manufacturer's

instructions. The sequencing was performed on an Illumina MiSeq instrument and the paired-

end read value was > 20 million/sample. Next, we aligned the transcriptome sequence reads

to the reference genome (Ensembl GRCh38 release) with STAR version 2.5.3a using 2-pass

alignment mode. The mirBase annotation was used in both mapping and read counting. After

alignment, the reads were associated with known miRNAs and the number of reads assigned

to each miRNA was counted using featureCounts from R package Rsubread. The data were

normalised using the TMM normalisation method of the edgeR R/Bioconductor package (R

version 3.4.1, Bioconductor version 3.5). After pre-processing and quality control we

analysed the target genes of the miRNAs (miRDB target score ≥80) and the IL-6R pathway

targeting miRNAs with Funrich 3.1.3 [32], g:Profiler [30] and NetworkAnalyst 3.0 [33]. The

miRNA-Seq data has been deposited to the ExoCarta database [34].

Investigation of target cells of BeWo-12.5K lEVs and characterization of molecular

interactions between 12.5K lEVs and lymphocytes

One hundred µL of the separated 12.5K lEVs were labeled with a freshly prepared PKH26

fluorescent dye solution at a final concentration of 5 µM according to the manufacturer's

instructions (Merck, USA). After staining, the labeled vesicles were washed by adding 1 mL

of RPMI medium and centrifuging at 12,500 g for 15 minutes at 16 °C. We evaluated the

effectiveness of staining by flow cytometry. Validation was performed using a mock control

(the same staining procedure without adding EV samples) and differential detergent lysis

(Supplementary Figure 5). The number of EVs was determined by adding internal counting

beads (PKH Reference Beads, Merck, USA). A constant amount of BeWo-derived PKH

labelled 12.5K lEVs (1×10^6) were added to immunophenotyped lymphocytes (5×10^5) for

30 min at 4 °C to investigate EV binding. T lymphocyte subsets were defined by

CD3+/CD4+ (Th cells), CD3+/CD4+/CD25+ (activated Th and regulatory Th cells) and

CD3+/CD8+ (Tc cells) positivity. B cells were characterized by their CD3-/CD19+

expression, and NK cells by their CD3-/CD56+ immunophenotype. Antibodies were titrated

for the optimal concentration. At the end of the incubation period, cells were washed with

PBS and analyzed by flow cytometry.

To characterize the molecular interactions between EVs and lymphocytes, we measured the

binding of PKH-labeled EVs after masking surface PS, CD95L, or HLA-G molecules using

annexin V, anti-CD95L, or anti-HLA-G antibodies. On the other hand, the surface

phosphatidylserine receptor (PSR) and CD95 molecules on lymphocytes were also blocked

with anti-PSR polyclonal antibody or anti-CD95 antibodies. PKH-labeled BeWo-derived

12.5K lEVs and immunophenotyped lymphocytes PKH-labeled BeWo-derived 12.5K lEVs

and immunophenotyped lymphocytes were incubated together in the presence or in absence

of monoclonal antibodies in PBS containing 4% bovine serum albumin (BSA) for 30 minutes

at 4 C.

Investigation of IL-4, IL-6, IL-17 and IFNγ production by *Cytometric Bead Array (CBA)*

IL-4, IL-6, IL-17 and IFN-γ production was detected from cell culture supernatants, by the

BD™ CBA (BD San Jose, California, USA) system according to the manufacturer's

instructions (BD Cytometric Bead Array (CBA) Human Th1/Th2/Th17 Cytokine Kit).

Intracellular IL-10 staining of CD4+ T lymphocytes

5×10^5 human lymphocytes were stimulated with BeWo-12.5K lEVs for 24 hours in the

presence of the protein transport inhibitor Brefeldin A (Thermo Fisher Scientific, USA).

Stimulated cells were washed and incubated in 100 µL of staining buffer (PBS containing 1%

BSA) with a pre-titrated optimal concentration of PerCP-conjugated anti-human CD4

monoclonal antibody (15 minutes at RT on ice). The immunophenotyped cells were fixed in a

4% paraformaldehyde solution (Sigma-Aldrich Co., St. Louis, MO) for 10 minutes at RT ,

permeabilized with a 0.1% saponin solution, and incubated in 50 µL of staining buffer (1%

FBS containing PBS) containing a pre-titrated optimal concentration of phycoerythrin-

conjugated anti-human IL-10 monoclonal antibody for 20 minutes at RT on ice. The cells

were washed in 2 mL of PBS and centrifuged at 300 g for five minutes. The pellet was

resuspended in 300 µL of 2% PFA and analyzed by flow cytometry.

Study of IL-6Rα expression on lymphocytes and BeWo cells

We examined the expression of IL-6Rα in lymphocytes obtained from co-culture systems

through immunophenotyping. Staining was performed according to the manufacturer's

instructions. A total of 5,000 cells were analyzed per tube.

Adherent BeWo cells were scraped off by cell scraper after a 20 minutes incubation period on

ice and were stained using the same staining method described for lymphocyte

immunophenotyping.

Detection of soluble IL-6Rα

Human Instant ELISA™ (Thermo Fisher, USA) was used for detection of soluble IL-6Rα

concentrations according to the manufacturer's instructions. IL-6Rα levels were detected in

pg/mL at 450 nm and 620 wavelengths by Multiskan MS Microplate Reader (Thermo

Fisher, USA). The sensitivity of the kit was 0.01 ng/mL.

IL-6 pathway signalling

For gene expression analysis, first total RNA isolated from untreated and BeWo-12.5K lEVs-

treated lymphocytes was purified with RNeasy Mini Kit (Qiagen, CA, USA). The RNA

concentration of each sample was determined using the Qubit RNA HR Assay Kit (Qubit

Fluorometer, Life Technologies). Samples were stored at -80°C until cDNA synthesis. From

each sample 600 ng total RNA was reverse transcribed with Sensifast cDNA Synthesis Kit

(Bioline, London, UK). Quantitative PCR reactions using SensiFAST SYBR Hi-ROX Kit

with SYBR Green primers (Supplementary table 2) were carried out on an ABI 7900 real-

time PCR instrument (Thermo Fisher Scientific, USA) according to the manufacturer's

instructions. Real-time PCR results were calculated according to the following protocol:

relative expression level = $2^{-\Delta Ct}$, where ΔCt = Ct (of gene of interest) – Ct (of houseskeeping

gene) with *HPRT* as an endogenous control.

To establish the optimal timepoint for the gene expression analysis we quantified the mRNA

levels of key IL-6 pathway factors (*IL6, IL6R, IL6ST, AKT1, STAT1, STAT3* and *PIAS3*) in

lymphocytes at 2, 4, 6, 8 and 24 hours after recombinant IL-6 (Sigma-Aldrich Co., St. Louis,

USA) (0.02 ng/ mL = 5.0×10^7 units/ mg) treatment. The concentration of IL-6 was chosen

accordingly to physiological circulating IL-6 concentration in healthy pregnancy [35, 36].

Based on the results of this experiment, we chose the four-hour treatment as the optional time

point for IL-6 pathway analysis.

Flow cytometry

Antibodies and dies used in flow cytometry are listed in Supplementary Table 3.

Measurements were carried out using a FACSCalibur flow cytometer (BD, San Jose, CA,

USA) on the day of the staining. In the case of cellular measurements, $5x10^4$ cells/ tube were

collected. Data were analysed with CellQuestPro (BD, San Jose, California, USA) and

Flowjo v10.3 (Tree Star Inc., OR, USA) software.

Statistics

GraphPad Prism version 8.0 (GraphPad Software, La Jolla California, USA) was used for

statistical analysis. For normally distributed data Two-sided Student's unpaired t-test,

respectively for multiple parameter analysis ANOVA test was used, followed by Bonferroni

correction. For data not showing normal distribution Mann-Whitney U or Wilcoxon test was

used, for multiple parameter analysis ANOVA test was used, followed by Kruskal-Wallis

test. The level of significance was set at $p < 0.05$.

The datasets generated during and/or analysed during the current study are available from the

corresponding author on reasonable request.

Results

Target cells of BeWo-EVs

The 12.5k lEVs primarily bound to CD3+ T lymphocytes; however, the binding ability of T

cell subsets differed. BeWo-lEVs preferred Th cells (5.04 ± 0.98 %) compared to Tc

lymphocytes (2.01 ± 0.14 %) ($p < 0.05$). On the other hand, significantly fewer CD19+ B

cells (0.92 ± 0.45 %, $p < 0.01$) and CD56+ NK cells bound BeWo-lEVs (0.14 ± 0.05 %) ($p <$

0.001) like T lymphocytes (Figure 1A-B). There was no difference in the binding ability to

the lymphocytes of pregnant and non-pregnant donors (non-pregnants: 3.55 ± 0.48 % vs.

pregnant: 5.04 ± 0.98 %) ($p > 0.05$). We used monoclonal antibodies or AnnexinV

pretreatments for blocking the BeWo-lEVs - lymphocyte interactions (Figure 1C-D). Anti-

phosphatidylserine receptor (PSR) mAb pre-treatment of lymphocytes partially inhibited the

binding of 12.5K lEVs to CD3+/CD4+ (untreated vs. anti-PSR Mab: 215 ± 106 mean

fluorescence intensity (MFI) vs. 112.5 ± 24.06 MFI) and to CD3+/CD8+ T cells (untreated

vs. anti-PSR Mab: 33.5 ± 9.3 MFI vs. 21.43 ± 7.17 MFI). Similarly, Annexin V pre-treatment

of BeWo-lEVs resulted in minimal binding inhibition (untreated vs. Annexin V treated: 13.2

± 0.46 MFI vs. 7.06 ± 0.4 MFI). Masking of HLA-G on BeWo-lEVs (untreated vs. anti-

HLA-G mAb: 13.05 ± 0.55 MFI vs. 7.83 ± 0.06 MFI) could also prevent the binding of

12.5K lEVs to lymphocytes. Neutralization of CD95 molecules on lymphocytes (untreated

vs. anti-CD95L mAb: 43.53 ±1.88 MFI vs. 39.5 ± 4.8 MFI) or CD95L (untreated vs. anti-

CD95 mAb: 13.05 ± 0.55 MFI vs. 16.35 ± 0.35 MFI) on 12.5K lEVs did not have a

significant effect on the 12.5K lEV-lymphocyte interaction. Based on these results we

concluded, that interactions via PS and HLA-G could coordinate, at least partially the binding

of 12.5K lEVs to CD4+ lymphocytes (Figure 1C-D).

The effects of BeWo-lymphocyte interactions on the IL-6Rα expression of T cells

IL-6Rα expressing cells were detected in all of the tested T cell subsets, though the

proportion of IL-6Rα+ cells varied among them (CD4+ T cells: 61.8 ± 6.3%; CD8+ T cells:

11.4 ± 1.9%). Since IL6 sensitivity depends on the number of IL-6Rα, we compared their

expression levels in T cell subsets. Exofacial IL-6Rα was significantly higher on CD4+ T

cells (13.72 ± 0.68 MFI) than on CD8+ T cells (9.31 ± 0.16 MFI). (p < 0.0001)

(Supplementary Figure 6A-D).

We examined the impact of BeWo-12.5K lEVs on the IL-6Rα expression of T lymphocytes

in BeWo-lymphocyte co-culture systems. BeWo-derived 12.5K lEVs significantly

downregulated the exofacial IL-6Rα on lymphocytes (Ly + 12.5K lEV: 9.7 ± 0.06 MFI) (p <

0.05) (Figure 2A-B). Since T lymphocytes were identified as targets of BeWo-12.5K lEVs,

the IL-6Rα levels of CD4+ T cell subsets were further analysed. The Th cells (untreated:

13.71 ± 0.68 MFI; co-cultured: 9.2 ± 0.04 MFI) and the CD25+ Th subset (untreated: 16.52 ±

1.18 MFI; co-cultured: 9.49 ± 0.27 MFI) exhibited significantly lower IL6Ra expression

levels when the co-cultured cells were separated by insert membranes (p<0.0001)

(Supplementary Figure 6 E-F). Direct cellular interaction (Ly + BeWo) did not have any

effect (data not shown). Soluble IL-6Rα can be formed by ectodomain shedding or

alternative splicing, but cells can also lose their exofacial IL-6Rα via vesicle-associated

shedding [37]. To determine the pathways of IL-6Rα shedding, the concentration of soluble

IL-6Rα and also the EV-bound IL-6Rα was determined by ELISA or flow cytometry,

respectively. Treatment of lymphocytes with BeWo-derived 12.5K lEVs did not change

either the soluble IL-6Rα concentration in cell culture supernatant (untreated vs. 12.5K lEV

treated: 114 ± 45.8 pg/ mL vs. 178.3 ± 24.1 pg/ mL p>0.05) or the number of IL-6Rα positive

12.5K lEVs (untreated vs. 12.5K lEV treated: 741 ± 326 events/ μL vs 1071 ± 351 events/

μL, data not shown), which may suggest that the IL-6Rα downregulation could happen at the

gene expression level.

To investigate the mechanism of IL-6 downregulation we performed miRNA-Seq of BeWo-

derived 12.5K lEVs. Altogether we identified 517 ± 70 miRNAs in the 12.5K lEV samples,

out of which 366 were common in all samples. The biological pathway analysis revealed that

the interleukin-6 receptor binding (GO:0005138) to be significantly enriched ($p_{adj} = 0.02$)

(Figure 2C) and 21 miRNAs from the 366 were targeting IL-6R. Overall, in the

NetworkAnalyst database there are 103 known miRNAs targeting IL-6R, through gene

silencing of *IL6ST* and *ERAP*. The top 5 highly expressed miRNAs targeting IL-6R found in

BeWo-derived 12.5K lEVs were the hsa-miR-92a-3p (18170 ± 5221 RPM), hsa-miR-520f-3p

(5274 ± 1002 RPM), hsa-miR-25-3p (5065 ± 838.8 RPM), hsa-miR-519d-3p (3703 ± 1261

RPM) and hsa-miR-26b-5p (1348 ± 363.3 RPM) (Figure 2D-E).

Further analysis of the miRNA pattern revealed that there is a strong miRNA signature

pattern among the biological replicates (Figure 3A and C). Most of the target genes of the

miRNAs found in BeWo-12.5K lEVs are expressed in the placental tissue and one of the

main target cell types is the CD4+ T cell (Figure 3B). Beside the IL-6 signalling pathway,

other significantly enriched targeted biological pathways were FAS and Arf6 downstream

pathways, the plasma membrane estrogen receptor and VEGF signalling, followed by the

β1integrin cell surface interaction biological pathway (Figure 3D). Additionally, BeWo-

12.5K lEVs induced molecular function network analysis reveals cell differentiation as one of

the main significant molecular function (p=0.001, Figure 3E). The in-silico analysis of cell

differentiation depicts that the process is initiated by signal transduction (IL6R

downregulation) followed by regulation of cell proliferation and cell cycle coupled with anti-

apoptotic signalling events. Furthermore, cell growth and immune response balancing

accompany cell differentiation (Figure 3F).

IL-6 pathway analysis

Lymphocytes pre-treated with 12.5K lEVs and further treated with recombinant IL-6 showed

a decreased *IL6R* expression. *STAT1* showed a 3.5-fold change increase in mRNA expression

upon 12.5K lEVs treatment (relative gene expression of untreated: 265.8 ± 125.8, 12.5K lEVs

treated: 1014 ± 102.4, Ly + BeWo: 308.4 ± 148.3, Ly insert BeWo: 990.8 ± 75.2). *PIAS3*

showed a 10 fold decrease in 12.5K lEVs treated lymphocytes (relative gene expression of

untreated: 44.7 ± 0.07, 12.5K lEVs treated: 4.48 ± 0.017, Ly + BeWo: 1.09 ± 0.37, Ly insert

Bewo: not detected), *AKT1* showed an 1.4 fold increase upon 12.5K lEVs stimulation

(relative gene expression of untreated: 16.8 ± 0.68, 12.5K lEV treated: 23.3 ± 0.4, Ly +

BeWo: 17.39 ± 3.15, Ly insert BeWo: 21.15 ± 4.0), *RELA*, *NFKBIA* showed an increased

mRNA expression (Figure 4). In BeWo-lymphocyte co-culture system we detected a decrease

in *CXCL12*, *IL1R1*, *IL6ST*, *OSMR*, *TNFRSF10B* and *STAT3* expression. Separation of BeWo

cells and lymphocytes with 1 μm pore sized membrane led to increased *IL10*, *IL2RA*, *IL1R1*,

CD80, *JAK2*, *JAK3*, *MAPK1*, *MAPK14*, *RAC1* and decreased *STAT3* expression detectable in

lymphocytes (Supplementary Table 4).

IL-6 production induced by BeWo-lymphocyte interaction or BeWo-derived 12.5K lEVs

Neither the BeWo cells nor the lymphocytes secreted IL-6 without stimulation; however,

both populations began producing it after co-culturing. Higher levels of IL-6 could be

measured after direct cell-to-cell interaction than with the transwell system (direct contact:

1,833 ± 30.02 pg/mL and 1,296 ± 15.61 pg/mL, p < 0.001). IL-6 production in the co-culture

system was also validated at the mRNA level by qPCR. (Figure 5.).

IFN-γ, IL-4, IL-10 and IL-17 production induced by BeWo-lymphocyte interactions

Neither the direct cellular interaction nor the membrane separated co-culture system of

lymphocytes and BeWo cells induced detectable levels of IFN-γ and IL-4. In the case of IL-

17, there was no significant difference compared to control lymphocyte secretion levels (Ly:

58 ± 0.01, Ly insert BeWo: 61.44 ± 0.27; Ly + BeWo: 61.77 ± 0.02, data not shown).

However, BeWo-derived 12.5K lEVs induced a significant IL-10 production in CD4+ T

cells. To elicit the specific potential of 12.5K lEVs induced IL-10 production, lymphocytes

were treated with BeWo-derived 12.5K lEVs, sEVs or the EV-free supernatant. Only the

12.5K lEV induced increased IL-10 expression in CD4+ T cells (12.5K lEVs treatment 305.9

± 29.1 MFI; sEV treatment: 252 ± 9 MFI; EV-free supernatant: 258.5 ± 45.5 MFI; non-

treated CD4+ T cells: 255.3 ± 46.2 MFI, p < 0.01 Wilcoxon test), as determined by

multicolour flow cytometry. (Supplementary figure 7.).

IL-6Rα expression of circulating lymphocytes of pregnant and non-pregnant women

As our in vitro studies showed a strong IL-6Rα downregulation in CD4+ lymphocytes, we

aimed to assess the *in vivo* significance of IL-6Rα downregulation during pregnancy.

Peripheral blood samples from first trimester healthy pregnant and non-pregnant control

women were assessed for IL-6Rα expression on circulating lymphocytes. We detected the

significantly lower percentage of IL-6Rα expressing CD3+/CD4+/CD25+ lymphocytes in the

blood of pregnant women compared to the healthy non-pregnant group (81.53 ± 4.2 % vs.

92.78 ± 1.95 %; p < 0.05) (Figure 6 A-D).

Discussion

Local immune tolerance is essential for a successful pregnancy. This antigen-specific

immune suppression is mediated by cellular and humoral factors, as well as the recently

recognized 12.5K lEV-related network. Through their complex molecular cargo, EVs may

play a role in maintaining local immune tolerance. The existence of suppressor regulatory T

lymphocytes was revealed in the early 1980s. CD4+/CD25+/FoxP3+ natural T_{reg} cells are

characterized by their low proliferating ability, anergy and the production of TGF-β and IL-

10 [38, 39]. Although the exact process of T_{reg} cells development is still unknown, peripheral

differentiation seems to be an existing pathway beside thymic differentiation [40].

The presence of T_{reg} cells can be shown in the feto-maternal interface during the entire

pregnancy and increased percentage of circulating CD4+/CD25+ Treg cells has been found in

the blood of healthy pregnant women; however, their level is dependent on gestational age:

the highest T_{reg} number can be measured during the second trimester; and their number

significantly decreases in 6-8 weeks after delivery [41]. An elevated level of T_{reg} cells can

also be detected in local lymph nodes and in vaginal secretions even in the earliest gestational

weeks. The question arises whether this early presence of regulatory T cells in pregnant

tissues would be a consequence of local differentiation from naïve T cells. One of the theories

suggests that fetal antigens presented by local professional antigen presenting cells induce the

differentiation of CD4+/CD25- Th lymphocytes into CD4+/CD25+ T_{reg} cells [42, 43].

It is well known that trophoblast cells present fetal alloantigens, playing a role in the

induction and maintenance of local immune tolerance at the feto-maternal interface. The role

of HLA-G in inducing Treg cells has also been described. In our earlier studies, we

demonstrated that HLA-G-expressing, trophoblast-derived 12.5K lEVs, which were isolated

from maternal plasma, bind to T lymphocytes. This binding inhibits the IL-2-induced signal

transduction pathway of T cells [28]. In the present work, we examined the role of

trophoblastic cells and their 12.5K lEVs in the putative local development of Treg cells by

using the BeWo choriocarcinoma in vitro model system. Co-culturing of BeWo cells and

lymphocytes resulted in IL-6 production. In the present work, we investigated the role of

trophoblastic cells and their 12.5K lEVs in the putative local development of Treg cells by

using the BeWo choriocarcinoma in vitro model system. Co-culturing of BeWo cells and

lymphocytes resulted in IL-6 production. IL-6 has been shown to be a critical cytokine that

controls the transition between the induction of regulatory T lymphocytes and Th17 cells

[44]. This shift is inauspicious for the pregnancy outcome, so it is cardinal that the immune

system counterbalances this effect. The modulation of cytokine-induced cellular effects via

regulation of receptor expression is a highly sensitive control mechanism. We determined the

IL-6Rα expression of the lymphocytes in order to check their IL-6 sensitivity. BeWo-derived

12.5K lEVs downregulated IL-6Rα expression of CD4+ T lymphocytes, indicating the

decreased IL-6 sensitivity. Decrease of IL-6Ra resulted in profound functional changes of

lymphocytes including disrupted IL-6 signalization and altered cytokine production.

Regulation of cell surface receptor expression can occur through multiple pathways including

ectodomain shedding [45], vesicle-mediated removal [46] or transcriptional and

posttranscriptional modification [47]. Ectodomain shedding or vesicle-mediated receptor

removal could be excluded based on our experiments, therefore, we focused on

transcriptional regulation. We premised that the regulation happens by the alteration of the

IL-6 pathway, and consequently, the mRNA level of *IL6R* would be downregulated, which

we have confirmed by qPCR analysis. Additionally, we observed the downregulation of key

players of the IL-6 pathway, including the central component *STAT3* transcription factor. To

elicit the downregulation mechanism of *IL6R*, we analysed whether miRNAs are mediating

the observed process. miRNAs represent a class of small, noncoding RNA molecules which

have major role in the posttranscriptional regulation of protein expression [48]. We employed

next-generation sequencing to analyse miRNA profiles in BeWo-12.5K lEVs and identified

21 miRNAs targeting downregulation of *IL6R* through gene silencing.

Our findings indicate that HLA-G expressing trophoblastic 12.5K lEVs have a central role in

the regulation of local Th cell polarization via regulation of their IL-6 sensitivity. The

appearance of placental EVs in maternal circulation raises the possibility that they play a role

in the maintenance of maternal immune tolerance at the systemic level as well. The lower

circulating level of IL-6Rα bearing CD4+/CD25+ T cell subset in healthy pregnant women

also supports this hypothesis.

It should be noted that in the placental microenvironment, EV-mediated interactions between

trophoblast cells and lymphocytes orchestrate the local immune response, which is critical for

the maintenance of maternal immune tolerance during pregnancy.

Acknowledgement:

This work was funded by STIA_2024 (19279/PMKP/2025) to ÉP,   Hungarian National

Research, Development and Innovation Office—NKFIH, OTKA PD_21 138521 grant to

ÁFK. We are grateful for the electronmicroscopic measurements of Tamás Visnovitz.

Author contribution

BN, GS, NF and ÁFK carried out the experiments; ÁFK, BN, GS and ÉP participated in data

analysis; ÁFK, EIB, and ÉP wrote the manuscript; ÉP designed the experiments.

Conflicts of interest

The authors declare no conflict of interests.

1. Aghaeepour N, Ganio EA, Mcilwain D, Tsai AS, Tingle M, Van Gassen S, Gaudilliere

DK, Baca Q, McNeil L, Okada R, Ghaemi MS, Furman D, Wong RJ, Winn VD,

Druzin ML, El-Sayed YY, Quaintance C, Gibbs R, Darmstadt GL, Shaw GM,

Stevenson DK, Tibshirani R, Nolan GP, Lewis DB, Angst MS, Gaudilliere B. An

immune clock of human pregnancy. Sci Immunol. 2017 Sep 1;2(15):eaan2946. doi:

10.1126/sciimmunol.aan2946.

2. Griffith OW, Chavan AR, Protopapas S, Maziarz J, Romero R, Wagner GP. Embryo

implantation evolved from an ancestral inflammatory attachment reaction. Proc Natl

Acad Sci U S A. 2017 Aug 8;114(32):E6566-E6575. doi: 10.1073/pnas.1701129114.

3. Ahn SH, Nguyen SL, Petroff MG. Exploring the Origin and Antigenic Specificity of

Maternal Regulatory T Cells in Pregnancy. Front Immunol. 2020 Jun 25;11:1302. doi:

10.3389/fimmu.2020.01302.

4. Saito S. Reconsideration of the Role of Regulatory T Cells during Pregnancy: Differential

Characteristics of Regulatory T Cells between the Maternal-Fetal Interface and

Peripheral Sites and between Early and Late Pregnancy. Med Princ Pract.

2022;31(5):403-414. doi: 10.1159/000527336.

5. Morelli S, Mandal M, Goldsmith LT, Kashani BN, Ponzio NM. The maternal immune

system during pregnancy and its influence on fetal development. Research and Reports

in Biology. 2015;6:171-189 https://doi.org/10.2147/RRB.S80652

6. Esparvarinha M, Madadi S, Aslanian-Kalkhoran L, Nickho H, Dolati S, Pia H, Danaii S,

Taghavi S, Yousefi M. Dominant immune cells in pregnancy and pregnancy

complications: T helper cells (TH1/TH2, TH17/Treg cells), NK cells, MDSCs, and the

immune checkpoints. Cell Biol Int. 2023 Mar;47(3):507-519. doi: 10.1002/cbin.11955.

7. Li QH, Zhao QY, Yang WJ, Jiang AF, Ren CE, Meng YH. Beyond Immune Balance: The

Pivotal Role of Decidual Regulatory T Cells in Unexplained Recurrent Spontaneous

Abortion. J Inflamm Res. 2024 May 1;17:2697-2710. doi: 10.2147/JIR.S459263.

8. Tang C, Hu W. The role of Th17 and Treg cells in normal pregnancy and unexplained

recurrent spontaneous abortion (URSA): New insights into immune mechanisms.

Placenta. 2023 Oct;142:18-26. doi: 10.1016/j.placenta.2023.08.065.

9. Sakaguchi S, Vignali DAA, Rudensky AY, Niec RE, Waldmann H. The plasticity and

stability of regulatory T cells. Nat Rev Immunol (2013) 13:461–7.

doi:10.1038/nri3464

10. Malviya V, Yshii L, Junius S, Garg AD, Humblet-Baron S, Schlenner SM. Regulatory T-

cell stability and functional plasticity in health and disease. Immunol Cell Biol. 2023

Feb;101(2):112-129. doi: 10.1111/imcb.12613.

11. Omere C, Richardson L, Saade GR, Bonney EA, Kechichian T, Menon R. Interleukin

(IL)-6: A Friend or Foe of Pregnancy and Parturition? Evidence From Functional

Studies in Fetal Membrane Cells. Front Physiol. 2020 Jul 24;11:891. doi:

10.3389/fphys.2020.00891.

12. Del Barco E, Franco-Jarava C, Vargas M, Maíz N, Arevalo S, Sánchez MÁ, Avilés MT,

Rodó C, Mendoza M, Goya M, Hernández-González M, Carreras E. Reference values

for interleukin-6 in the amniotic fluid of asymptomatic pregnant women. Acta Obstet

Gynecol Scand. 2023 Apr;102(4):480-485. doi: 10.1111/aogs.14524.

13. Chang Y, Li W, Shen Y, Li S, Chen X. Association between interleukin-6 and preterm

birth: a meta-analysis. Ann Med. 2023;55(2):2284384. doi:

10.1080/07853890.2023.2284384.

14. Prins JR, Gomez-Lopez N, Robertson SA. Interleukin-6 in pregnancy and gestational

disorders. J Reprod Immunol. 2012 Sep;95(1-2):1-14. doi: 10.1016/j.jri.2012.05.004.

15. Dienz O, Rincon M. The effects of IL-6 on CD4 T cell responses. Clin Immunol. 2009

Jan;130(1):27-33. doi: 10.1016/j.clim.2008.08.018.

488 16. Korn T, Hiltensperger M. Role of IL-6 in the commitment of T cell subsets. Cytokine.

489   2021 Oct;146:155654. doi: 10.1016/j.cyto.2021.155654.

490 17. Huang N, Chi H, Qiao J. Role of Regulatory T Cells in Regulating Fetal-Maternal

491   Immune Tolerance in Healthy Pregnancies and Reproductive Diseases. Front

492   Immunol. 2020 Jun 26;11:1023. doi: 10.3389/fimmu.2020.01023.

493 18. Muñoz EL, Fuentes FB, Felmer RN, Yeste M, Arias ME. Extracellular vesicles in

494   mammalian reproduction: a review. Zygote. 2022;30(4):440-463.

495   doi:10.1017/S0967199422000090

496 19. Tantengco OAG, Richardson L, Lee A, Kammala A, Silva MC, Shahin H, Sheller-Miller

497   S, Menon R. Histocompatibility Antigen, Class I, G (HLA-G)'s Role during Pregnancy

498   and Parturition: A Systematic Review of the Literature. Life (Basel). 2021 Oct

499   9;11(10):1061. doi: 10.3390/life11101061.

500 20 Cho K, Kook H, Kang S, Lee J. Study of immune-tolerized cell lines and extracellular

501   vesicles inductive environment promoting continuous expression and secretion of

502   HLA-G from semiallograft immune tolerance during pregnancy. J Extracell Vesicles.

503   2020 Jul 20;9(1):1795364. doi: 10.1080/20013078.2020.1795364.

504 21. Nakahara A, Nair S, Ormazabal V, Elfeky O, Garvey CE, Longo S, Salomon C.

505   Circulating Placental Extracellular Vesicles and Their Potential Roles During

506   Pregnancy. Ochsner J. 2020 Winter;20(4):439-445. doi: 10.31486/toj.20.0049. PMID:

507   33408584; PMCID: PMC7755551.an Niel G, D'Angelo G, Raposo G. Shedding light

508   on the cell biology of extracellular vesicles. Nat Rev Mol Cell Biol (2018) Available

509   at: http://dx.doi.org/10.1038/nrm.2017.125

510 22. Liu, YJ., Wang, C. A review of the regulatory mechanisms of extracellular vesicles-

511   mediated intercellular communication. Cell Commun Signal 21, 77 (2023).

512   https://doi.org/10.1186/s12964-023-01103-6

23. Buzas EI. The roles of extracellular vesicles in the immune system. Nat Rev Immunol.

2023 Apr;23(4):236-250. doi: 10.1038/s41577-022-00763-8.

24. Welsh JA, Goberdhan DCI, O'Driscoll L, Buzas EI, Blenkiron C, Bussolati B, Cai H, Di

Vizio D, Driedonks TAP, Erdbrügger U, Falcon-Perez JM, Fu QL, Hill AF, Lenassi

M, Lim SK, Mahoney MG, Mohanty S, Möller A, Nieuwland R, Ochiya T, Sahoo S,

Torrecilhas AC, Zheng L, Zijlstra A, Abuelreich S, Bagabas R, Bergese P, Bridges

EM, Brucale M, Burger D, Carney RP, Cocucci E, Crescitelli R, Hanser E, Harris AL,

Haughey NJ, Hendrix A, Ivanov AR, Jovanovic-Talisman T, Kruh-Garcia NA,

Ku'ulei-Lyn Faustino V, Kyburz D, Lässer C, Lennon KM, Lötvall J, Maddox AL,

Martens-Uzunova ES, Mizenko RR, Newman LA, Ridolfi A, Rohde E, Rojalin T,

Rowland A, Saftics A, Sandau US, Saugstad JA, Shekari F, Swift S, Ter-Ovanesyan D,

Tosar JP, Useckaite Z, Valle F, Varga Z, van der Pol E, van Herwijnen MJC, Wauben

MHM, Wehman AM, Williams S, Zendrini A, Zimmerman AJ; MISEV Consortium;

Théry C, Witwer KW. Minimal information for studies of extracellular vesicles

(MISEV2023): From basic to advanced approaches. J Extracell Vesicles. 2024

Feb;13(2):e12404. doi: 10.1002/jev2.12404. Erratum in: J Extracell Vesicles. 2024

May;13(5):e12451.

25. Liu T, Zhang Q, Zhang J, Li C, Miao YR, Lei Q, Li Q, Guo AY. EVmiRNA: A database

of miRNA profiling in extracellular vesicles. Nucleic Acids Res (2019)

doi:10.1093/nar/gky985

26. Paul N, Sultana Z, Fisher JJ, Maiti K, Smith R. Extracellular vesicles- crucial players in

human pregnancy. Placenta. 2023 Sep 7;140:30-38. doi:

10.1016/j.placenta.2023.07.006.

27. Ciferri MC, Quarto R, Tasso R. Extracellular Vesicles as Biomarkers and Therapeutic

Tools: From Pre-Clinical to Clinical Applications. Biology (Basel). 2021 Apr

23;10(5):359. doi: 10.3390/biology10050359.

28. Pap E, Pállinger E, Falus A, Kiss AA, Kittel A, Kovács P, Buzás EI. T lymphocytes are

targets for platelet- and trophoblast-derived microvesicles during pregnancy. Placenta.

2008; 29: 826-32. doi: 10.1016/j.placenta.2008.06.006.

29. Kovács ÁF, Fekete N, Turiák L, Ács A, Kőhidai L, Buzás EI, Pállinger É. Unravelling

the Role of Trophoblastic-Derived Extracellular Vesicles in Regulatory T Cell

Differentiation. Int J Mol Sci. 2019 Jul 14;20(14):3457. doi: 10.3390/ijms20143457.

30. Kovács ÁF, Láng O, Turiák L, Ács A, Kőhidai L, Fekete N, Alasztics B, Mészáros T,

Buzás EI, Rigó J, et al. The impact of circulating preeclampsia-associated extracellular

vesicles on the migratory activity and phenotype of THP-1 monocytic cells. *Sci Rep*

(2018) **8**:5426. doi:10.1038/s41598-018-23706-7

31. György B, Módos K, Pállinger É, Pálóczi K, Pásztói M, Misják P, Deli MA, Sipos Á,

Szalai A, Voszka I, et al. Detection and isolation of cell-derived microparticles are

compromised by protein complexes resulting from shared biophysical parameters.

Blood 2011Jan 27;117(4):e39-48. doi: 10.1182/blood-2010-09-307595.32. Pathan M,

Keerthikumar S, Ang CS, Gangoda L, Quek CYJ, Williamson NA, Mouradov D,

Sieber OM, Simpson RJ, Salim A, et al. FunRich: An open access standalone

functional enrichment and interaction network analysis tool. *Proteomics* (2015)

15:2597–2601. doi:10.1002/pmic.201400515

33. Raudvere U, Kolberg L, Kuzmin I, Arak T, Adler P, Peterson H, Vilo J. g:Profiler: a web

server for functional enrichment analysis and conversions of gene lists (2019 update).

Nucleic Acids Res (2019) doi:10.1093/nar/gkz369

34. Zhou G, Soufan O, Ewald J, Hancock REW, Basu N, Xia J. NetworkAnalyst 3.0: a visual

analytics platform for comprehensive gene expression profiling and meta-analysis.

Nucleic Acids Res (2019) doi:10.1093/nar/gkz240

35.   Soriano D, Hugol D, Quang NT, Darai E. Serum concentrations of interleukin-2R (IL-

2R), IL-6, IL-8, and tumor necrosis factor alpha in patients with ectopic pregnancy.

Fertil Steril (2003) **79**:975–980. https://doi.org/10.1016/S0015-0282(02)04853-7

36.   Li Y, Wang Y, Ding X, Duan B, Li L, Wang X. Serum Levels of TNF-α and IL-6 Are

Associated with Pregnancy-Induced Hypertension. *Reprod Sci* 2016 Oct;23(10):1402-

8.doi:10.1177/1933719116641760

37.   Schumacher N, Meyer D, Mauermann A, Von Der Heyde J, Wolf J, Schwarz J,

Knittler K, Murphy G, Michalek M, Garbers C, et al. Shedding of endogenous

interleukin-6 receptor (IL-6R) is governed by a disintegrin and metalloproteinase

(ADAM) proteases while a full-length IL-6R isoform localizes to circulating

microvesicles. *J Biol Chem* 2015 Oct 23;290(43):26059-71.

doi:10.1074/jbc.M115.649509

38.   Papiernik M, De Moraes ML, Pontoux C, Vasseur F, Pénit C. Regulatory CD4 T cells:

Expression of IL-2Rα chain, resistance to clonal deletion and IL-2 dependency. *Int*

Immunol 1998 Apr;10(4):371-8. doi:10.1093/intimm/10.4.371

39.   Dieckmann D, Plottner H, Berchtold S, Berger T, Schuler G. Ex Vivo Isolation and

Characterization of Cd4 [+] Cd25 [+] T Cells with Regulatory Properties from Human

Blood. *J Exp Med* 2001 Jun 4;193(11):1303-10. doi:10.1084/jem.193.11.1303

40.   Kim B-S, Park Y-J, Kang C-Y. Peripheral Generation of CD4+ CD25+ Foxp3+

Regulatory T Cells. *Immune Netw* (2007) **7**:1–9. https://doi.org/10.4110/in.2007.7.1.1

41.   Somerset DA, Zheng Y, Kilby MD, Sansom DM, Drayson MT. Normal human

pregnancy is associated with an elevation in the immune suppressive CD25 + CD4 +

regulatory T-cell subset. *Immunology* 2004 May;112(1):38-43.doi:10.1111/j.1365-

2567.2004.01869.x

42.   Zenclussen AC, Gerlof K, Zenclussen ML, Sollwedel A, Bertoja AZ, Ritter T, Kotsch

588   K, Leber J, Volk HD. Abnormal T-cell reactivity against paternal antigens in

589   spontaneous abortion: Adoptive transfer of pregnancy-induced CD4+CD25+T

590   regulatory cells prevents fetal rejection in a murine abortion model. *Am J Pathol* 2005

591   Mar;166(3):811–822.doi:10.1016/S0002-9440(10)62302-4

592 43. Robertson SA. Seminal plasma and male factor signalling in the female reproductive

593   tract. *Cell Tissue Res* 2005 Oct;322(1):43-52. doi:10.1007/s00441-005-1127-3

594 44. Harbour SN, DiToro DF, Witte SJ, Zindl CL, Gao M, Schoeb TR, Jones GW, Jones

595   SA, Hatton RD, Weaver CT. TH17 cells require ongoing classic IL-6 receptor

596   signaling to retain transcriptional and functional identity. Sci Immunol. 2020 Jul

597   17;5(49):eaaw2262. doi: 10.1126/sciimmunol.aaw2262.

598 45. Croucher PI, Wang F, Hargreaves PG. Interleukin-6 receptor shedding: a possible role

599   for members of the ADAM family. *Biochem Soc Trans* (1999) **27**:224 LP-228.

600   doi:10.1042/bst0270224

601 46. Park, EJ., Lee, CW. Soluble receptors in cancer: mechanisms, clinical significance,

602   and therapeutic strategies. *Exp Mol Med* **56**, 100–109 (2024).

603   https://doi.org/10.1038/s12276-023-01150-647.  Luo Y, Zheng SG. Hall of fame

604   among pro-inflammatory cytokines: Interleukin-6 gene and its transcriptional

605   regulation mechanisms. *Front Immunol* (19.dec. 2016) doi:10.3389/fimmu.2016.00604

606 48. Wahid F, Shehzad A, Khan T, Kim YY. MicroRNAs: Synthesis, mechanism, function,

607   and recent clinical trials. *Biochim Biophys Acta - Mol Cell Res* 2010

608   Nov;1803(11):1231-43.doi:10.1016/j.bbamcr.2010.06.013

Abbreviations

EV – extracellular vesicles

12.5K lEV – intermediate-sized extracellular vesicles

FSC – Forward Scatter

HLA-G – human leukocyte antigen G (major histocompatibility complex, class I, G)

IL-6Rα – interleukin 6 receptor alpha

miRNA – micro ribonucleic acid

PS – phosphatidylserine

PSR – phosphatidylserine receptor

sEV – small-sized extracellular vesicles

SSC – Side Scatter

Figure 1. Target cells of BeWo-derived 12.5K lEVs

A. Binding of BeWo-12.5K lEVs to immunophenotyped lymphocytes isolated from pregnant

women BeWo-12.5K lEVs bound significantly more CD3+/CD4+ T cells (n= 6 pregnant

donors, one-way ANOVA followed by Bonferroni correction). B. Representative dot plots

show the cell-bound 12.5K lEVs to the different lymphocyte subpopulations. C. Analysis of

selected blocking between 12.5K lEVs and their target CD4+ T cells. Anti-PSR, annexin V

and anti-HLA-G blocking resulted in significant inhibition (n=6 pregnant donors, one-way

ANOVA followed by Bonferroni correction).   D. Representative histograms show the

changes of expression levels after selective blocking of a specific interaction between 12.5K

lEVs and target lymphocytes (red histogram: PKH2+ surface bound 12.5K lEVs, blue

histogram: PKH+ surface bound 12.5K lEVs after targeted blocking, n= 6 different pregnant

donors; measurements were repeated 2 times from each donor).

Figure 2. IL-6Rα downregulation mediated by miRNAs

A. Analysis of IL-6 downregulation upon BeWo-12.5K lEV treatment (n=10 donors). B.

Representative FACS histogram showing the IL-6Rα downregulation upon 12.5K lEV treatment

marked with red (untreated control - blue) C. Molecular function analysis of BeWo-derived

miRNAs target genes revealing the interleukin-6 receptor binding as one of the main targets. D.

Overall, we identified 366 common miRNA in 12.5K lEVs. 21 miRNAs are targeting the

downregulation of IL-6R, together with the expression values (RPM). E. IL-6R downregulating

miRNA network with highlighted (dark blue) miRNA found in the BeWo-12.5K lEVs. Values

represent mean ± SEM (n=10 ** p < 0.01, *** p < 0.001) Ly - lymphocytes

Figure 3. BeWo-derived 12.5K lEV miRNA content

A. Chord diagram analysis showing the overlap between the biological replicates. B. Bar chart

showing the site expression of genes targeted by BeWo-12.5K lEVs. Based on the analysis the

main target tissue is the placenta. One of the prominent cell types are the leukocytes, most of the

targeted cells are the CD4+ T cells (blue bars represent the percentage of target genes, red bars

are representing the p-values, and the yellow bars represent the p=0.05 reference). C. Venn

diagram showing the 366-common miRNA in 12.5K lEVs. D. Biological pathway analysis

showing the miRNA targeted pathways, among the IL-6 signalling, the following pathways are

significantly targeted: FAS signalling and Arf6 downstream pathways, the plasma membrane

estrogen receptor and VEGF signalling. E. BeWo-derived 12.5K lEV induced molecular function

network analysis reveals as the main molecular function the cell differentiation. F. The in-silico

detailed analysis of cell differentiation function reveals the molecular function is mainly induced

through alteration of signal transduction (IL-6R downregulation), followed by regulation of cell

proliferation and cell cycle coupled with an anti-apoptotic signalling event. Furthermore, cell

growth and immune response balancing accompany the cell differentiation.

Figure 4. IL-6 pathway analysis

A. Heatmap analysis of IL-6 pathway of non-pregnants' lymphocytes treated with BeWo-12.5K

lEVs, lymphocytes co-cultured with BeWo cells with or without a 1µm pore sized insert

membrane (n=3, each sample was run in triplicates). Heatmap showing the differential expressed

genes of 12.5K lEV treated or BeWo co-cultured lymphocytes normalized to the untreated

control lymphocytes. B. Schematic figure of the analysed IL-6 pathway molecules. Green arrows

show significant downregulation at mRNA level upon BeWo-derived 12.5K lEV treatment. Red

arrows mark the upregulated molecules upon 12.5K lEV treatment. C. Figures showing the

relative gene expression: down- or upregulation of *IL6R*, *STAT1, STAT3* and *NFKBIA* mRNAs

upon different treatments of lymphocytes (one-way ANOVA followed by Bonferroni correction,

* $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, ****$p < 0.0001$).

Figure 5. Interleukin 6 secretion

674 A. IL-6 could be detected both in the cell culture supernatant of insert separated BeWo co-culture

675 system and also in the media of BeWo-12.5K lEV treated non-pregnants' lymphocytes.

676 Unstimulated lymphocytes did not secrete IL-6 (n=11, **** $p < 0.0001$, One-Way ANOVA,

677 followed by Bonferroni's correction). B. Validation of IL-6 secretion at the mRNA level.

679 Figure 6. IL6-Rα expression on pregnant and non-pregnant lymphocytes

680 A. Analysis of IL-6Rα expressing circulating CD4+ T cells in healthy non-pregnant and pregnant

681 donors (n=12). B. Representative overlay histograms show the IL-6Rα expression on CD4+

682 lymphocytes of a pregnant (blue histogram) and a non-pregnant donor (red histogram). C.

683 Percentage of IL-6Rα expressing CD4+/CD25+ T_{reg} cells in healthy non-pregnant and pregnant

684 donors (n=12). D. Representative overlay histograms show the IL-6Rα expression on

685 CD4+/CD25+ lymphocytes of a pregnant (blue histogram) and a non-pregnant donor (red

686 histogram) (unpaired t-test, * $p < 0.05$).

A.

B.

A.

B.

C.

D.