The macroecology of viral coinfection

Cecilia A. Sánchez[1], Colin J. Carlson[1], Amy R. Sweeny[2*]

[1]Department of Epidemiology of Microbial Diseases, Yale School of Public Health, New Haven,
CT, USA
[2]Institute of Ecology and Evolution, University of Edinburgh, Edinburgh, UK
[*]Corresponding author: Amy R. Sweeny. Email: amyrsweeny@gmail.com

ORCID iDs:
CAS: https://orcid.org/0000-0002-1141-6816
CJC: https://orcid.org/0000-0001-6960-8434
ARS: https://orcid.org/0000-0003-4230-171X

Abstract

Coinfection is common in wild animals, and can have profound consequences for disease
outcomes and transmission. However, most research on coinfection is based on laboratory
experiments or well-studied wildlife systems. Here, we use the testing data from the PREDICT
project – the largest standardized wildlife disease surveillance project ever conducted – to
describe patterns of viral coinfection, and evaluate their association with host and virus factors.
We find that coinfection is rare because viral infections are rare, but coinfection is more
common than expected by random chance – especially among coronaviruses,
paramyxoviruses, and influenza A virus. We further find that coinfection risk changes over the
host lifespan, but bats and rodents exhibit divergent responses to aging. Finally, we find that
cave-roosting bats do not have a significantly higher rate of coinfection, but wild animals in
captivity do – highlighting potential risks to wildlife and human health from the wildlife trade. Our
results suggest that macroecological patterns of coinfection are driven by a mix of
macroevolutionary and host-level variation in immunity, virus-virus interactions, and human
influence, and point to coinfection as a potentially important (and frequently, undetected) driver
of viral dynamics in nature.

Introduction

Parasites exist in complex communities, where coinfection with multiple parasites at any time is the norm for wild animals [1,2]. These coinfections provide an opportunity for direct interactions, such as competition for resources [3] or space [4], or indirect interactions, such as those mediated by host immune responses [5]. These parasite interactions can have a dramatic effect on outcomes at the host level, and transmission dynamics at the population level [6–9].

Several theoretical frameworks in disease ecology have incorporated the dynamics and consequences of coinfection [8–11], but it can be difficult to test their predictions against empirical evidence. Within-host processes are easiest to study in laboratory settings [12], but the controlled settings that make experimental approaches useful (e.g., laboratory organisms from a single host species, with known ancestry and no unknown parasites) also limit their similarity to real-world dynamics in natural communities. Well-developed (and often long-term) field systems have also led to a number of insights about interactions within diverse host and parasite communities [5], and some systems allow for natural experiments such as parasite removal or nutritional supplementation [4,13,14]. However, neither field nor experimental case studies can provide insight into the full network of interactions occurring within parasite communities (or their consequences) at the ecosystem scale or above.

Whereas large-scale biodiversity data have reached volumes that allow for unprecedented research and monitoring, wildlife disease data collection has historically been fragmented between emerging infectious disease surveillance, wildlife management efforts, and ecological research. Across all three fields, there are significant global gaps in surveillance [15,16], and – despite broader progress towards open science in ecology and epidemiology – the sharing of spatially-explicit, host-level infection data remains rare [17,18]. Some studies have used data synthesis to develop global datasets [16,19–21], but in most cases, they have focused on relationships between hosts and their parasites, rather than among the parasites themselves.

Here, we explore coinfection dynamics at the global scale for the first time, using the single largest standardized study of wildlife disease ever conducted. The PREDICT project was a decade-long, multi-country initiative funded by the U.S. Agency for International Development (USAID) that conducted viral surveillance primarily in wild animals [22]. The project produced one of the largest standardized multi-virus datasets ever assembled and resulted in the discovery of new viruses [23,24], an improved understanding of drivers of virus infection and

diversity [25,26], and insight into human and virus-related risk factors for zoonotic spillover [27].

However, minimal work has been done using the PREDICT data to test macroecological

hypotheses. Using these data, we set out to answer three questions: (1) Is viral coinfection

common in nature? (2) Is coinfection more common in certain host groups, life stages, and

settings? And finally: (3) Are there non-random patterns of coinfection driven by specific virus

taxa, or by interactions between specific pairs of virus taxa?

Methods

Analyses were performed using the R statistical environment v4.4.2 [28]. Prior to analysis, we

cleaned the PREDICT data by excluding human test results and fixing minor errors in animal

host taxonomy. We combined an animal sampling dataset that contained details such as

species, sex, age class with another dataset of virus test results for individual specimens, so

that each test result was paired with the relevant animal data.

Data overview

Data from the PREDICT-2 project (the second half of the decade-long initiative) was openly

shared on the USAID Development Data Library prior to the takedown of the USAID website.

Specimen collection (n = 126,952 unique specimens from 65,662 unique animals) was

concentrated in West and Central Africa as well as South and Southeast Asia, with > 10,000

specimens collected in Sierra Leone, Guinea, and Bangladesh each (Fig. S1). Most specimens

were collected from bats (n = 77,334, 60.9%) and rodents (n = 25,683, 20.2%). The dataset

includes some farm and companion animals, including poultry, dogs, and swine. Virus screening

was conducted using consensus polymerase chain reaction (PCR) assays (714,954 total tests),

mostly targeting viral families of significance to human health (e.g., coronaviruses, filoviruses,

and paramyxoviruses), but in some cases testing for specific viruses of interest (e.g., influenza

viruses and ebolaviruses).

Analyses

For each animal associated with one or more virus detections, we quantified the number of

viruses and virus families detected in that animal's specimens, and created binary

classifications noting whether virus or virus family coinfection was detected. For each taxonomic

group, we calculated the overall infection and coinfection prevalence. For bats, rodents, and

birds (the three largest taxonomic groups), we compared the observed coinfection prevalence to

the expected coinfection prevalence using a permutation testing approach. Specifically, we

randomized infection across individual animals while maintaining the observed prevalence of
each virus within the taxonomic group, then calculated the number of simulated coinfected
animals. We repeated this procedure 1000 times for each taxonomic group, then plotted the
simulated number of coinfected animals compared to the true number.
104
We created a coinfection matrix at the virus family level for use in network analyses, where each
count within the matrix represents a coinfection of two virus families occurring in the same
animal. For the small number of animals that were infected with three viruses, we counted each
coinfection separately (e.g., an animal infected with coronavirus X, coronavirus Y, and
paramyxovirus Z would count for two coronavirus-paramyxovirus coinfections and one
coronavirus-coronavirus coinfection). We used the *ggraph* package [29] to visualize the
frequency of virus family coinfection. To test if certain virus families co-occurred more often than
expected by chance, we first calculated the expected coinfection prevalence of two virus
families as the product of each virus family's overall prevalence in the dataset. We then
performed a binomial test for each virus family–virus family combination to determine if the
observed coinfection prevalence was significantly greater than expected. To account for multiple
comparisons, we adjusted *P* values using a Bonferroni correction. Finally, we repeated the
above analyses for two subsets of the full data: bats and rodents.
118
We used generalized linear models to explore host correlates of virus coinfection. **1)** For an all-
animals model (n = 2,462 animals), we excluded taxonomic groups with ≤ 10 animals, animals
that were missing sex or age class data, and a small number of neonates and fetuses. Using
the 'glm' function in the *stats* package, we modeled animal coinfection status (binary: either
single infection or coinfection) as a function of geographic region (Africa, Asia), taxonomic group
(bats, birds, rodents, shrews), sex (female, male), age class (adult, subadult, juvenile), and
captivity status (wild free-ranging, owned domesticated, wild in captivity). Given inconsistency in
host identification (i.e., not all animals could be identified to species level), we did not include
any species-level traits as model predictors. **2)** For a bats-only model (n = 1,519), we classified
whether each bat species roosts in caves using data from Willoughby *et al.* [30] and the IUCN
Red List. We excluded any bats for which cave roosting behavior or taxonomic family was not
available and included the same restrictions on sex and age class as above. Using the
'glmmTMB' function in the *glmmTMB* package [31], we modeled bat coinfection status as a
function of four fixed effects variables (geographic region, sex, age class, and cave roosting
status), and one random effects variable (family). **3)** For a rodents-only model (n = 488), we

included the same restrictions on sex and age class as above and modeled rodent coinfection
status as a function of those two predictors (collapsing subadults and juveniles into one
category due to sample size restrictions). Family and geographic region were not included as
variables because nearly all rodents were murids and were collected in Asia.
138

Results

140   We found that both infection and coinfection were rare overall. Out of 65,662 unique animals,
141   3,271 (4.98%) were infected with one or more viruses (Fig. 1A); viral prevalence in birds
142   (487/5,617; 8.67%) and bats (2,059/36,181; 5.69%) was somewhat higher than average, while
143   prevalence in rodents (549/12,876; 4.26%) was slightly lower than average. Animals with at
144   least one virus detection were most often captured in South and Southeast Asia (Fig. 1B).
145

A single virus was detected in most infected animals (n = 3,048, 93.18%); coinfections of two (n
= 211, 6.45%) or three (n = 12, 0.37%) viruses in the same animal were infrequent. Of the 223
animals in which virus coinfection was detected, over half were bats (n = 128, 57.4%) with the
remainder being 50 rodents, 39 birds, 5 swine, and 1 shrew. Within infected animals, virus
coinfection rates were highest in rodents (50/549, 9.11%), birds (39/487, 8.01%), and bats
(128/2,059, 6.22%). Roughly half of the triple coinfections (7/12; Table S1) were domesticated
ducks and geese, which were all infected with duck coronavirus, influenza A virus, and either
Newcastle virus or avian paramyxovirus 6; the remainder (5/12) were bats infected with a mix of
coronaviruses, paramyxoviruses, influenza A virus, and a novel rhabdovirus.
155

Coinfection is non-random across geography, phylogeny, and life stage
Bats, rodents, and birds exhibited unique patterns of coinfection, even accounting for their over-
representation in the dataset. Coinfections were more common than expected based on the
observed prevalence of all viruses: for all three taxonomic groups, we applied a permutation
testing approach, and found that true coinfection rates exceeded the maximum random
frequency observed across 1,000 simulations (Fig. S2). These results suggest that coinfection is
more common than expected, indicating that animals either share risk factors for exposure, or
viral infections tend to increase susceptibility to secondary infections. However, after accounting
for other host factors, we found that birds were significantly less likely to be coinfected than bats
(odds ratio (OR) = 0.30, 95% confidence interval (CI): 0.11-0.67, P = 0.007; Fig. 2A, Table S2),
and rodents and shrews were marginally less likely than bats to be coinfected (rodents: OR =
0.76, 95% CI: 0.47-0.1.19, P = 0.239; shrews: OR = 0.43, 95% CI: 0.02-2.03, P = 0.404). These

findings could support the widespread hypothesis (and growing evidence) that bats are able to

tolerate viral infection, and potentially coinfection, through a unique set of immune adaptations

[32–34]. However, we note that virus coinfection rates within the all-animals model dataset (i.e.,

animals with available sex and age data) differed compared to the coinfection rates in the entire

dataset: while coinfection was still high in rodents (46/488, 9.43%) and bats (112/1,628, 6.88%),

it was noticeably lower in birds (12/316, 3.80%).

Animals sampled in Asia were more likely to be coinfected than those sampled in Africa in both

the all-animals model (OR = 2.07, 95% CI: 1.40-3.08, P < 0.001; Fig. 2A, Table S2) and the

bats-only model (OR = 2.61, 95% CI: 1.65-4.13, P < 0.001; Fig. 2C, Table S3). Rather than

reflecting any characteristic of one region versus another, we believe that this result was instead

driven by the geographic mosaic of specific coevolutionary associations between host and virus

groups [35]. In particular, flying foxes (*Pteropus* spp.; n = 3,084 animals) and rats (*Rattus* spp.;

n = 5,109) were heavily sampled but infected animals were primarily found in Asia, and both of

these genera had high coinfection rates (*Pteropus*: 37/245; 15.1%; *Rattus*: 44/403; 10.9%).

Coinfection was associated with life history, but those relationships varied between groups.

Based on previous studies [25,36], we hypothesized that infection prevalence – and therefore

coinfection rates – would be higher in younger animals, due to waning of maternal antibodies

and less-developed immune systems [37]. We found that younger rodents (juveniles and

subadults) were more likely to be coinfected than adults (OR = 2.14, 95% CI: 1.12-4.00, P =

0.018; Fig. 2E, Table S4). However, juvenile bats were less likely to be coinfected than adults

(OR = 0.14, 95% CI: 0.03-0.57, P = 0.006), and this effect was strong enough that it dominated

the all-animals model (OR = 0.34, 95% CI: 0.10-0.84, P = 0.040). This pattern is aligned with the

widespread hypothesis that bats' unique immune adaptations allow them to tolerate persistent

viral infections [38,39], which might accumulate over the course of an individual's lifespan.

Alternatively, depending on the timing of sampling by the PREDICT study, juvenile bats may

have been captured at times when levels of maternal antibodies were still high. Though

logistically challenging, longitudinal sampling within populations can help elucidate patterns of

virus infection and coinfection in relation to host maturation [36,40,41].

Finally, we found that host social structure could drive coinfection patterns, but found divergent

effects between natural and un-natural settings. For bats, cave-roosting behavior was not a

significant predictor of coinfection status. Cave-roosting bats can exist in large groups in close

proximity, potentially increasing infection rates, but other roosting strategies can also create
large, close aggregations of bats (e.g., tree-roosting *Pteropus* species that cluster together).
Finer-scale contact patterns that we could not examine here (e.g., segregation by age or sex, or
centrality in social networks) might lead to more correlated infection risk across viruses at the
individual level. Sickness behaviors (e.g., reduced social interactions) after an initial infection
could also reduce an individual's subsequent likelihood of infection with other viruses [42,43].
208
In contrast, across all animals, we found that wild animals in captivity were significantly more
likely to be coinfected than free-ranging wild animals (OR = 2.16, 95% CI: 1.19-3.88, *P* = 0.010;
Fig. 2A, Table S2). These results support the hypothesis that captivity can increase parasite
transmission through both crowding, stress, and unusual patterns of interspecific contact. This
hypothesis has been strongly supported by previous analyses of coronavirus testing conducted
by the PREDICT project [23,44], and is a particular point of concern, given the potential for
coinfection to lead to recombinant coronaviruses [45], and the high risk of spillover in wildlife
farms and markets [46]. However, we note that our model was limited to an unrepresentative
sample of the data for which age and sex were available, and so of the 163 animals classified
as "wild animals in captivity," 159 (98%) were either rats (*Rattus* spp.) or mallards (*Anas*
platyrhynchos).
220

Evidence of virus-virus interactions
Coinfection networks revealed that certain viruses–namely, coronaviruses (CoVs),
paramyxoviruses (PMVs), and orthomyxoviruses (specifically, influenza A virus)–were most
frequently detected together (Fig. 2B, 2D, 2F). Notably, there were 108 total coinfections with
multiple CoVs (57 in bats, 48 in rodents), 69 CoV-PMV coinfections (44 in bats, 2 in rodents), 45
CoV-influenza coinfections (23 in bats), and 10 PMV-influenza coinfections (1 in bats). Binomial
tests indicated that CoV-PMV, CoV-influenza, and PMV-rhabdovirus coinfections occurred
significantly more than expected in all animals and in bats, while coinfections with multiple
coronaviruses occurred significantly more than expected in rodents (Tables S5-S7). All other
virus family coinfections occurred < 10 times, and no coinfections involving arenaviruses,
filoviruses, or hantaviruses were detected.
232

233   Examining coinfection patterns at the level of individual viruses, we found that viruses that are
234   transmitted by poultry and other livestock (influenza A virus, Newcastle disease virus, and
235   infectious bronchitis virus) had the highest centrality in the network. However, a handful of

236 commonly occurring betacoronaviruses, including murine coronavirus, PREDICT CoV-16 and

237 PREDICT CoV-17, and Bat coronavirus HKU9, were also well connected within the network.

Discussion

Here, we conducted the first comprehensive analysis of viral coinfection patterns across multiple

continents and an entire host community. We found that coinfection is rare, because viral

infections are rare, but coinfection occurs more frequently than expected, particularly among

specific virus groups. Our findings suggest that widely-occurring viruses like influenza A virus

could have important effects on host susceptibility to secondary infections, and might be

important to account for in research on other pathogens. Future work could explore how specific

viral coinfections influence factors such as the magnitude and duration of shedding, particularly

as it could relate to zoonotic spillover risk [47]. We also found that coinfection dynamics may be

modulated by both broad differences in host immunology and fine-scale host contact processes,

including in anthropogenic settings like farms and the wildlife trade. These findings are also of

potential importance for public health, as they highlight the need for both basic research on

host-parasite interactions, and surveillance of environments where coinfection can produce new

reassortant and recombinant viruses with epidemic or pandemic potential [25].

Although our study is the largest macroecological study of coinfection to date, our findings

should be interpreted in light of the limitations of our data. Most importantly, the PREDICT

project was not designed with ecological research as the primary objective, and so the testing

data have uneven coverage both across and within taxonomic groups. For instance, the

PREDICT project was almost single-handedly responsible for bats becoming the best-studied

wildlife hosts of viruses [48]. Given that over 60% of specimens were collected from bats, these

data probably drove many of our broad-scale findings. However, only half of all bat families (10

of 21) were sampled, and an emerging body of work highlights that while bats are unique viral

reservoirs, there is still considerable important (and interesting) variation within the group [34].

Future work could build on initial steps here by designing surveillance to more explicitly search

for and report coinfection. A recent systematic review found that coinfections were common in

bats, but were incidentally reported within study findings [49]. More deliberate focus on sampling

to capture coinfection dynamics may also require specific sampling strategies: for example,

reliably detecting coinfection in an individual host requires capture of animals in-hand (versus,

e.g., pooled or environmental sampling). Similarly, different parasites are associated with

different organs and sites in the body, and so prioritizing particular samples or assays may be

necessary, depending on study aims. Although broad sample and parasite coverage presents

substantial cost considerations, several recent developments offer possible solutions. Multiplex,

metagenomic, and metatranscriptomic methods all provide a fuller view of the host virome, and

are especially useful when sample sizes are small due to biological limitations (e.g., collecting

blood from small animals) [50–52]. Longitudinal surveillance also improves the odds of viral

detection, and allows researchers to test hypotheses about asynchronous interactions between

parasites, host immunity, and external environmental factors [53]. Finally, host-level data on

wildlife disease are also shared infrequently [17], preventing the kinds of re-analyses we

conducted here, or approaches that synthesize findings through systematic reviews and meta-

analyses [49]. Developing a better global view of wildlife disease dynamics is an essential step

to conducting more macroecological research on the consequences of coinfection.

Acknowledgements

284 This work was supported by NSF DBI 2515340. We thank our colleagues on the Verena team

285 for helpful feedback and brainstorming, and the countless members of the PREDICT consortium

286 who contributed to one of the largest and most unique ecological datasets ever collected.

Data Accessibility

289 PREDICT data on animal sampling and viral testing will be made available to the reviewers and

290 editors at the time of submission as .csv files. A public link to the data will be provided upon

291 manuscript publication. All code to reproduce analyses is available at

292 https://github.com/viralemergence/macroecology-coinfection.

Figures

Figure 1. A. Number of unique animals infected by one or more viruses, separated by taxonomic group. Groups with only single infections are not displayed (dogs: n = 10; goats/sheep: n = 4; cattle/buffalo: n = 2; carnivores: n = 1; other: n = 1). **B.** Distribution of unique animals associated with one or more virus detections. Points are partially transparent and jittered for improved visibility.

Figure 2. Predictors of virus coinfection (**A**, **C**, **E**) and virus family coinfection networks (**B**, **D**, **E**). In the left column, generalized linear model outputs (odds ratios and 95% confidence intervals (CIs)) for predictors of coinfection are shown for all animals (**A**), bats only (**C**), and rodents only (**E**). CIs that do not cross the vertical line at 1 (i.e., statistically significant) are purple, while CIs that do cross are gray. Reference levels are Africa (geographic region), bats (taxonomic group), female (sex), adult (age class), wild free-ranging (captivity status), and non-cave roosting (roosting status). In the right column, networks of virus family coinfection are shown for all animals (**B**), bats only (**D**), and rodents only (**F**). Nodes represent virus families, while edges represent the frequency of virus family coinfection. Asterisks indicate coinfections that occurred significantly more than expected.

317   **Figure 3.** Virus coinfection network. Nodes represent individual viruses, while edges represent
318   the frequency of virus coinfection. Only nodes with degree centrality ≥ 4 are labeled.
319

320
321

References

1. Cox FEG. 2001 Concomitant infections, parasites and immune responses. Parasitology 122, S23–S38.

2. Petney TN, Andrews RH. 1998 Multiparasite communities in animals and humans: frequency, structure and pathogenic significance. Int. J. Parasitol. 28, 377–393.

3. Budischak SA, Hansen CB, Caudron Q, Garnier R, Kartzinel TR, Pelczer I, Cressler CE, van Leeuwen A, Graham AL. 2017 Feeding immunity: Physiological and behavioral responses to infection and resource limitation. Front. Immunol. 8, 1914.

4. Knowles SCL, Fenton A, Petchey OL, Jones TR, Barber R, Pedersen AB. 2013 Stability of within-host - parasite communities in a wild mammal system. Proceedings of the Royal Society of London B: Biological Sciences 280.

5. Ezenwa VO, Budischak SA, Buss P, Seguel M, Luikart G, Jolles AE, Sakamoto K. 2021 Natural resistance to worms exacerbates bovine tuberculosis severity independently of worm coinfection. Proc. Natl. Acad. Sci. U. S. A. 118. (doi:10.1073/pnas.2015080118)

6. Telfer S, Lambin X, Birtles R, Beldomenico P, Burthe S, Paterson S, Begon M. 2010 Species interactions in a parasite community drive infection risk in a wildlife population. Science 330, 243–246.

7. Gorsich EE, Etienne RS, Medlock J, Beechler BR, Spaan JM, Spaan RS, Ezenwa VO, Jolles AE. 2018 Opposite outcomes of coinfection at individual and population scales. Proceedings of the National Academy of Sciences 115, 7545–7550.

8. Pedersen AB, Fenton A. 2007 Emphasizing the ecology in parasite community ecology. Trends in Ecology & Evolution 22, 133–139.

9. Graham AL. 2008 Ecological rules governing helminth–microparasite coinfection. Proceedings of the National Academy of Sciences 105, 566–570.

10. Fenton A, Pedersen AB. 2005 Community Epidemiology Framework for Classifying Disease Threats. Emerg. Infect. Dis. 11, 1815–1821.

11. Sweeny AR, Albery GF, Becker DJ, Eskew EA, Carlson CJ. 2021 Synzootics. J. Anim. Ecol. 90, 2744–2754.

12. Fenton A, Knowles SCL, Petchey OL, Pedersen AB. 2014 The reliability of observational approaches for detecting interspecific parasite interactions: comparison with experimental results. Int. J. Parasitol. 44, 437–445.

13. Sweeny AR, Thomason CA, Carbajal EA, Hansen CB, Graham AL, Pedersen AB. 2020 Experimental parasite community perturbation reveals associations between Sin Nombre virus and gastrointestinal nematodes in a rodent reservoir host. Biol. Lett. 16, 20200604.

14. Ezenwa VO, Jolles AE. 2015 Epidemiology. Opposite effects of anthelmintic treatment on
microbial infection at individual versus population scales. Science 347, 175–177.
15. Carlson CJ et al. 2025 Pathogens and planetary change. Nat. Rev. Biodivers. 1, 32–49.
16. Cohen LE, Fagre AC, Chen B, Carlson CJ, Becker DJ. 2023 Coronavirus sampling and
surveillance in bats from 1996–2019: a systematic review and meta-analysis. Nature
Microbiology 8, 1176–1186.
17. Schwantes CJ et al. 2025 A minimum data standard for wildlife disease research and
surveillance. Sci. Data 12, 1054.
18. Albery GF, Sweeny AR, Becker DJ, Bansal S. 2022 Fine-scale spatial patterns of wildlife
disease are common and understudied. Funct. Ecol. 36, 214–225.
19. Cohen JM, Sauer EL, Santiago O, Spencer S, Rohr JR. 2020 Divergent impacts of warming
weather on wildlife disease risk across climates. Science 370.
(doi:10.1126/science.abb1702)
20. Heckley A, Becker D. 2024 Parasite traits shape the association between forest loss and
infection: A global meta-analysis. EcoEvoRxiv. (doi:10.32942/x2pd2q)
21. Warmuth VM, Metzler D, Zamora-Gutierrez V. 2023 Human disturbance increases
coronavirus prevalence in bats. Sci Adv 9, eadd0688.
22. PREDICT Consortium. 2020 Advancing Global Health Security at the Frontiers of Disease
Emergence. University of California, Davis: One Health Institute.
23. Anthony SJ et al. 2017 Further evidence for bats as the evolutionary source of Middle East
respiratory syndrome Coronavirus. MBio 8. (doi:10.1128/mBio.00373-17)
24. Goldstein T et al. 2018 The discovery of Bombali virus adds further support for bats as
hosts of ebolaviruses. Nat. Microbiol. 3, 1084–1089.
25. Anthony SJ et al. 2017 Global patterns in coronavirus diversity. Virus Evol 3, vex012.
26. Eskew EA, Olival KJ, Mazet JAK, Daszak P. 2025 A global-scale dataset of bat viral
detection suggests that pregnancy reduces viral shedding. Proc. Biol. Sci. 292, 20242381.
27. Saylors K et al. 2021 Socializing One Health: an innovative strategy to investigate social
and behavioral risks of emerging viral threats. One Health Outlook 3, 11.
28. R Core Team. 2016 R: A language and environment for statistical computing. R Foundation
for Statistical Computing, Vienna, Austria. See https://www.R-project.org/.
29. Pedersen TL. 2024 ggraph: An implementation of grammar of graphics for graphs and
networks. R package version 2.2.1. See https://github.com/thomasp85/ggraph,
https://ggraph.data-imaginist.com.
30. Willoughby A, Phelps K, Olival K, PREDICT Consortium. 2017 A comparative analysis of

390   viral richness and viral sharing in cave-roosting bats. Diversity (Basel) 9, 35.

391 31. Brooks M, Kristensen K, Benthem K van, Magnusson A, Berg C, Nielsen A, Skaug H,

392   Mächler M, Bolker B. 2017 GlmmTMB balances speed and flexibility among packages for

393   zero-inflated generalized linear mixed modeling. R J. 9, 378.

394 32. Letko M, Seifert SN, Olival KJ, Plowright RK, Munster VJ. 2020 Bat-borne virus diversity,

395   spillover and emergence. Nat. Rev. Microbiol. 18, 461–471.

396 33. Irving AT, Ahn M, Goh G, Anderson DE, Wang L-F. 2021 Lessons from the host defences

397   of bats, a unique viral reservoir. Nature 589, 363–370.

398 34. Becker DJ et al. 2025 Diverse hosts, diverse immune systems: Evolutionary variation in bat

399   immunology. Ann. N. Y. Acad. Sci. (doi:10.1111/nyas.15395)

400 35. Forero-Muñoz NR, Muylaert RL, Seifert SN, Albery GF, Becker DJ, Carlson CJ, Poisot T.

401   2024 The coevolutionary mosaic of bat betacoronavirus emergence risk. Virus Evol 10,

402   vead079.

403 36. Peel AJ et al. 2025 Synchronized seasonal excretion of multiple coronaviruses coincides

404   with high rates of coinfection in immature bats. Nat. Commun. 16, 6579.

405 37. Goenka A, Kollmann TR. 2015 Development of immunity in early life. J. Infect. 71 Suppl 1,

406   S112–20.

407 38. Calisher CH, Childs JE, Field HE, Holmes KV, Schountz T. 2006 Bats: Important reservoir

408   hosts of emerging viruses. Clin. Microbiol. Rev. 19, 531–545.

409 39. Plowright RK, Peel AJ, Streicker DG, Gilbert AT, McCallum H, Wood J, Baker ML, Restif O.

410   2016 Transmission or within-host dynamics driving pulses of zoonotic viruses in reservoir-

411   host populations. PLoS Negl. Trop. Dis. 10, e0004796.

412 40. Wacharapluesadee S et al. 2018 Longitudinal study of age-specific pattern of coronavirus

413   infection in Lyle's flying fox (Pteropus lylei) in Thailand. Virol. J. 15, 38.

414 41. Brook CE et al. 2019 Disentangling serology to elucidate henipa- and filovirus transmission

415   in Madagascar fruit bats. J. Anim. Ecol. 88, 1001–1016.

416 42. Ripperger SP, Stockmaier S, Carter GG. 2020 Tracking sickness effects on social

417   encounters via continuous proximity sensing in wild vampire bats. Behav. Ecol. 31, 1296–

418   1302.

419 43. Moreno KR, Weinberg M, Harten L, Salinas Ramos VB, Herrera M LG, Czirják GÁ, Yovel

420   Y. 2021 Sick bats stay home alone: fruit bats practice social distancing when faced with an

421   immunological challenge. Ann. N. Y. Acad. Sci. 1505, 178–190.

422 44. Huong NQ et al. 2020 Coronavirus testing indicates transmission risk increases along

423   wildlife supply chains for human consumption in Viet Nam, 2013-2014. PLoS One 15,

e0237129.

45. Wells HL, Bonavita CM, Navarrete-Macias I, Vilchez B, Rasmussen AL, Anthony SJ. 2023
The coronavirus recombination pathway. Cell Host Microbe 31, 874–889.

46. Sánchez CA, Li H, Phelps KL, Zambrana-Torrelio C, Wang L-F, Olival KJ, Daszak P. 2021
A strategy to assess spillover risk of bat SARS-related coronaviruses in Southeast Asia.
medRxiv (doi:10.1101/2021.09.09.21263359)

47. Schuh AJ, Amman BR, Guito JC, Graziano JC, Sealy TK, Towner JS. 2025 Modeling
natural coinfection in a bat reservoir shows modulation of Marburg virus shedding and
spillover potential. PLoS Pathog. 21, e1012901.

48. Gibb R, Albery GF, Mollentze N, Eskew EA, Brierley L, Ryan SJ, Seifert SN, Carlson CJ.
2022 Mammal virus diversity estimates are unstable due to accelerating discovery effort.
Biol. Lett. 18, 20210427.

49. Jones BD, Kaufman EJ, Peel AJ. 2023 Viral co-infection in bats: A systematic review.
Viruses 15, 1860.

50. Boyd V, Smith I, Crameri G, Burroughs AL, Durr PA, White J, Cowled C, Marsh GA, Wang
L-F. 2015 Development of multiplexed bead arrays for the simultaneous detection of
nucleic acid from multiple viruses in bat samples. J. Virol. Methods 223, 5–12.

51. Raghwani J et al. 2023 Seasonal dynamics of the wild rodent faecal virome. Mol. Ecol. 32,
4763–4776.

52. Chang W-S, Eden J-S, Hall J, Shi M, Rose K, Holmes EC. 2020 Metatranscriptomic
analysis of virus diversity in urban wild birds with paretic disease. J. Virol. 94.
(doi:10.1128/JVI.00606-20)

53. Plowright RK, Becker DJ, McCallum H, Manlove KR. 2019 Sampling to elucidate the
dynamics of infections in reservoir hosts. Philos. Trans. R. Soc. Lond. B Biol. Sci. 374,
20180336.

Supplementary Information

Figure S1. Map of sampling effort in the PREDICT dataset. Only one point is plotted for each country, with the size corresponding to the number of unique specimens collected in that country.

458 **Figure S2.** Density plots of 1000 coinfection simulations each for bats (n = 36,181), rodents (n =
459 12,876), and birds (n = 5,617). The true number of coinfected animals for each taxonomic group
460 is plotted with a dashed vertical line.
461

462
463

464   **Table S1.** Characteristics of the 12 animals that were infected with three viruses. Note that host
465   and virus names are provided as originally described in the PREDICT dataset, and may not
466   conform to the latest taxonomies.
467

PREDICT sample ID	Country	Taxonomic group	Species	Virus family	Virus
56344	Egypt	Bats	Rousettus aegyptiacus	Coronaviridae	Bat coronavirus HKU9
				Orthomyxoviridae	Influenza A
				Paramyxoviridae	PREDICT_PMV-116
108315	Thailand	Bats	Pteropus lylei	Coronaviridae	PREDICT_CoV-17
				Paramyxoviridae	PREDICT_PMV-6
				Paramyxoviridae	PREDICT_PMV-128
108328	Thailand	Bats	Pteropus giganteus	Coronaviridae	PREDICT_CoV-16
				Coronaviridae	PREDICT_CoV-68
				Paramyxoviridae	PREDICT_PMV-125
133237	Cameroon	Bats	Rhinolophus cf. alcyone	Coronaviridae	Kenya bat coronavirus/BtKY83/59/58
				Coronaviridae	PREDICT_CoV-70
				Coronaviridae	PREDICT_CoV-109
145203	Cambodia	Bats	Scotophilus kuhlii	Coronaviridae	Bat coronavirus 512/2005
				Paramyxoviridae	PREDICT_PMV-66
				Rhabdoviridae	PREDICT_RbdV-21
54944	Bangladesh	Birds	Anas	Coronaviridae	Duck Coronavirus

				Orthomyxoviridae	Influenza A
			platyrhynchos domesticus	Paramyxoviridae	Newcastle Disease Virus
108865	Bangladesh	Birds	Anas platyrhynchos domesticus	Coronaviridae	Duck Coronavirus
				Orthomyxoviridae	Influenza A
				Paramyxoviridae	Newcastle Disease Virus
126366	Bangladesh	Birds	Cairina moschata	Coronaviridae	Infectious bronchitis virus (IBV)
				Coronaviridae	Duck Coronavirus
				Orthomyxoviridae	Influenza A
126464	Bangladesh	Birds	Anas platyrhynchos domesticus	Coronaviridae	Duck Coronavirus
				Orthomyxoviridae	Influenza A
				Paramyxoviridae	Newcastle Disease Virus
126519	Bangladesh	Birds	Anas platyrhynchos domesticus	Coronaviridae	Duck Coronavirus
				Orthomyxoviridae	Influenza A
				Paramyxoviridae	Avian Paramyxovirus 6
126536	Bangladesh	Birds	Anser cygnoides domesticus	Coronaviridae	Duck Coronavirus
				Orthomyxoviridae	Influenza A
				Paramyxoviridae	Avian Paramyxovirus 6
126561	Bangladesh	Birds	Anas platyrhynchos domesticus	Coronaviridae	Duck Coronavirus
				Orthomyxoviridae	Influenza A
				Paramyxoviridae	Avian Paramyxovirus 6

469   **Table S2.** Model coefficients for a generalized linear model to examine effects of taxonomy,
470   sex, age, and captivity status on virus coinfection status in bats, birds, rodents, and shrews. *P*
471   values < 0.05 are in bold.
472

Variable		Estimate	SE	z	*P*	Odds ratio (95% CI)
Intercept		-2.92	0.18	-16.52	**< 0.001**	0.05 (0.04-0.08)
Geographic region	Africa	–	–	–	–	–
	Asia	0.73	0.20	3.63	**< 0.001**	2.07 (1.40-3.08)
Taxonomic group	bats	–	–	–	–	–
	birds	-1.20	0.45	-2.69	**0.007**	0.30 (0.11-0.67)
	rodents	-0.28	0.24	-1.18	0.239	0.76 (0.47-1.19)
	shrews	-0.86	1.02	-0.84	0.404	0.43 (0.02-2.03)
Sex	female	–	–	–	–	–
	male	-0.05	0.16	-0.28	0.776	0.96 (0.70-1.31)
Age class	adult	–	–	–	–	–
	subadult	-0.004	0.20	-0.02	0.986	1.00 (0.66-1.47)
	juvenile	-1.07	0.52	-2.05	**0.040**	0.34 (0.10-0.84)
Captivity status	free-ranging wild animal	–	–	–	–	–
	owned domesticated animal	0.18	0.60	0.31	0.760	1.20 (0.36-4.01)
	wild animal in captivity	0.77	0.30	2.57	**0.010**	2.16 (1.19-3.88)

473
474

475 **Table S3.** Model coefficients (fixed effects only) for a generalized linear mixed model to
476 examine effects of sex, age, and cave roosting behavior on virus coinfection status in bats. *P*
477 values < 0.05 are in bold.
478

Variable		Estimate	SE	z	*P*	Odds ratio (95% CI)
Intercept		-3.13	0.47	-6.72	**< 0.001**	0.04 (0.02-0.11)
Geographic region	Africa	–	–	–	–	–
	Asia	0.96	0.23	4.10	**< 0.001**	2.61 (1.65-4.13)
Sex	female	–	–	–	–	–
	male	-0.09	0.21	-0.42	0.675	0.92 (0.61-1.37)
Age class	adult	–	–	–	–	–
	subadult	-0.59	0.32	-1.86	0.063	0.55 (0.30-1.03)
	juvenile	-2.00	0.73	-2.73	**0.006**	0.14 (0.03-0.57)
Cave roosting	no	–	–	–	–	–
	yes	-0.12	0.28	-0.42	0.673	0.89 (0.51-1.54)

480 **Table S4.** Model coefficients for a generalized linear model to examine effects of sex and age
481 on virus coinfection status in rodents. *P* values < 0.05 are in bold.
482

Variable		Estimate	SE	z	*P*	Odds ratio (95% CI)
Intercept		-2.48	0.26	-9.44	**< 0.001**	0.08 (0.05-0.14)
Sex	female	–	–	–		–
	male	-0.03	0.31	-0.11	0.913	0.97 (0.52-1.80)
Age class	adult	–	–	–		–
	subadult/juvenile	0.76	0.32	2.36	**0.018**	2.14 (1.12-4.00)

484   **Table S5.** Expected versus observed coinfection prevalence for virus families, for all animals. *P*
485   values were calculated from binomial tests and adjusted using a Bonferroni correction. P_{adj}
486   values < 0.05 are in bold.
487

Virus family combination	Expected coinfection prevalence	Observed coinfection prevalence	Observed > expected	P_{adj}
Arenaviridae-Arenaviridae	5.80e-9	0	No	1
Arenaviridae-Coronaviridae	2.76e-6	0	No	1
Arenaviridae-Filoviridae	1.39e-8	0	No	1
Arenaviridae-Flaviviridae	5.80e-9	0	No	1
Arenaviridae-Hantaviridae	4.64e-9	0	No	1
Arenaviridae-Orthomyxoviridae	5.49e-7	0	No	1
Arenaviridae-Paramyxoviridae	5.42e-7	0	No	1
Arenaviridae-Rhabdoviridae	5.68e-8	0	No	1
Coronaviridae-Coronaviridae	1.31e-3	1.64-3	Yes	0.841
Coronaviridae-Filoviridae	6.62e-6	0	No	1
Coronaviridae-Flaviviridae	2.76e-6	0	No	1
Coronaviridae-Hantaviridae	2.21e-6	0	No	1
Coronaviridae-Orthomyxoviridae	2.61e-4	6.85e-4	Yes	**< 0.001**
Coronaviridae-Paramyxoviridae	2.58e-4	1.05e-3	Yes	**< 0.001**
Coronaviridae-Rhabdoviridae	2.70e-5	4.57e-5	Yes	1
Filoviridae-Filoviridae	3.34-8	0	No	1
Filoviridae-Flaviviridae	1.39e-8	0	No	1
Filoviridae-Hantaviridae	1.11e-8	0	No	1
Filoviridae-Orthomyxoviridae	1.32e-6	0	No	1
Filoviridae-Paramyxoviridae	1.30e-6	0	No	1
Filoviridae-Rhabdoviridae	1.36e-7	0	No	1
Flaviviridae-Flaviviridae	5.80e-9	0	No	1

Flaviviridae-Hantaviridae	4.64e-9	0	No	1
Flaviviridae-Orthomyxoviridae	5.49e-7	1.52e-5	Yes	1
Flaviviridae-Paramyxoviridae	5.42e-7	0	No	1
Flaviviridae-Rhabdoviridae	5.68e-8	0	No	1
Hantaviridae-Hantaviridae	3.71e-9	0	No	1
Hantaviridae-Orthomyxoviridae	4.39e-7	0	No	1
Hantaviridae-Paramyxoviridae	4.33e-7	0	No	1
Hantaviridae-Rhabdoviridae	4.55e-8	0	No	1
Orthomyxoviridae-Orthomyxoviridae	5.19e-5	0	No	1
Orthomyxoviridae-Paramyxoviridae	5.12e-5	1.52e-4	Yes	0.090
Orthomyxoviridae-Rhabdoviridae	5.38e-6	0	No	1
Paramyxoviridae-Paramyxoviridae	5.06e-5	7.61e-5	Yes	1
Paramyxoviridae-Rhabdoviridae	5.31e-6	9.14e-5	Yes	**< 0.001**
Rhabdoviridae-Rhabdoviridae	5.57e-7	0	No	1

489 **Table S6.** Expected versus observed coinfection prevalence for virus families, for bats only. *P*
490 values were calculated from binomial tests and adjusted using a Bonferroni correction. P_{adj}
491 values < 0.05 are in bold.
492

Virus family combination	Expected coinfection prevalence	Observed coinfection prevalence	Observed > expected	P_{adj}
Coronaviridae-Coronaviridae	1.80e-3	1.58e-3	No	1
Coronaviridae-Filoviridae	1.41e-5	0	No	1
Coronaviridae-Flaviviridae	1.17e-6	0	No	1
Coronaviridae-Orthomyxoviridae	2.37e-4	6.36e-4	Yes	**< 0.001**
Coronaviridae-Paramyxoviridae	3.89e-4	1.22e-3	Yes	**< 0.001**
Coronaviridae-Rhabdoviridae	5.75e-5	8.29e-5	Yes	1
Filoviridae-Filoviridae	1.10e-7	0	No	1
Filoviridae-Flaviviridae	9.17e-9	0	No	1
Filoviridae-Orthomyxoviridae	1.85e-6	0	No	1
Filoviridae-Paramyxoviridae	3.03e-6	0	No	1
Filoviridae-Rhabdoviridae	4.49e-7	0	No	1
Flaviviridae-Flaviviridae	7.64e-10	0	No	1
Flaviviridae-Orthomyxoviridae	1.54e-7	0	No	1
Flaviviridae-Paramyxoviridae	2.53e-7	0	No	1
Flaviviridae-Rhabdoviridae	3.74e-8	0	No	1
Orthomyxoviridae-Orthomyxoviridae	3.12e-5	0	No	1
Orthomyxoviridae-Paramyxoviridae	5.11e-5	2.76e-5	No	1
Orthomyxoviridae-Rhabdoviridae	7.56e-6	0	No	1
Paramyxoviridae-Paramyxoviridae	8.37e-5	1.11e-4	Yes	1
Paramyxoviridae-Rhabdoviridae	1.24e-5	1.66e-4	Yes	**< 0.001**
Rhabdoviridae-Rhabdoviridae	1.83e-6	0	No	1

494   **Table S7.** Expected versus observed coinfection prevalence for virus families, for rodents only.
495   *P* values were calculated from binomial tests and adjusted using a Bonferroni correction. P_{adj}
496   values < 0.05 are in bold.
497
498

Virus family combination	Expected coinfection prevalence	Observed coinfection prevalence	Observed > expected	P_{adj}
Arenaviridae-Arenaviridae	1.51e-7	0	No	1
Arenaviridae-Coronaviridae	1.47e-5	0	No	1
Arenaviridae-Hantaviridae	6.03e-8	0	No	1
Arenaviridae-Orthomyxoviridae	6.03e-8	0	No	1
Arenaviridae-Paramyxoviridae	1.60e-6	0	No	1
Coronaviridae-Coronaviridae	1.44e-3	3.73e-3	Yes	**< 0.001**
Coronaviridae-Hantaviridae	5.90e-6	0	No	1
Coronaviridae-Orthomyxoviridae	5.90e-6	0	No	1
Coronaviridae-Paramyxoviridae	1.56e-4	1.55e-4	No	1
Hantaviridae-Hantaviridae	2.41e-8	0	No	1
Hantaviridae-Orthomyxoviridae	2.41e-8	0	No	1
Hantaviridae-Paramyxoviridae	6.39e-7	0	No	1
Orthomyxoviridae-Orthomyxoviridae	2.41e-8	0	No	1
Orthomyxoviridae-Paramyxoviridae	6.39e-7	0	No	1
Paramyxoviridae-Paramyxoviridae	1.69e-5	0	No	1

499