A phylogenetic protein-coding genome-phenome map of complex traits across 224 primate species

Alejandro Valenzuela[1],[2]**, Fabio Barteri[1],[4],[9]**, Claudia Vasallo[4], Lukas Kuderna[10], Joseph Orkin[11], Jean Boubli[12], Amanda Melin[13], Hafid Laayouni[1], Kyle Farh[10], Jeffrey Rogers[14], Tomàs Marquès-Bonet[1], Gerard Muntané[1],[5],[6]*, Arcadi Navarro[1],[2],[3],[4]*, David Juan[15]*

**These authors contributed equally to this work.

*****To whom correspondence should be addressed: gerard.muntane@upf.edu; arcadi.navarro@upf.edu; david.juan@cnb.csic.es

---

[1] Institute of Evolutionary Biology (IBE, UPF–CSIC), Department of Medicine and Life Sciences, Universitat Pompeu Fabra, Parc de Recerca Biomèdica de Barcelona (PRBB), Barcelona, Spain; [2] Institució Catalana de Recerca i Estudis Avançats (ICREA) and Universitat Pompeu Fabra, Barcelona, Spain; [3] Center for Genomic Regulation (CRG), The Barcelona Institute of Science and Technology, Barcelona, Spain; [4] BarcelonaBeta Brain Research Center, Pasqual Maragall Foundation, Barcelona, Spain; [5] Hospital Universitari Institut Pere Mata, Institut de Recerca Biomèdica Catalunya Sud, Universitat Rovira i Virgili, Reus, Spain; [6] Centro de Investigación Biomédica en Red en Salud Mental (CIBERSAM), Spain; [9] Museu de Ciències Naturals de Barcelona, Barcelona, Spain; [10] Illumina, San Diego, CA, USA; [11] Département d'Anthropologie, Université de Montréal, Montréal, Québec, Canada; [12] University of Salford, Salford, UK; [13] University of Calgary, Calgary, Canada; [14] Baylor College of Medicine, Houston, TX, USA; [15] Systems Biology Department, Spanish National Center for Biotechnology (CNB-CSIC), Madrid, Spain

Abstract.

Complex traits arise from networks of coding and regulatory loci, making their genetic basis difficult to resolve. Macroevolutionary studies leverage millions of years of divergence across species to uncover fixed genomic changes invisible to within-species approaches like GWAS. Studying variation in mammals and primates offers insights into the evolution of complex traits and provides a complementary framework for generating hypotheses in biomedical research. Here, we present the first phylogenetic protein-coding, primate-wide genome–phenome map (P3GMap), spanning 263 traits across 224 primate species, which we release through the Primate Genome-Phenome Archive (PGA, https://pgarchive.github.io). Using two complementary approaches, convergent amino acid substitutions and relative evolutionary rates, we link protein-coding variation to complex phenotypes and identify thousands of gene-trait associations, including lineage-specific adaptations in diet, immunity, and lifespan.

One sentence summary: Cross-species genome-phenome mapping in primates reveals thousands of protein-coding variants linked to complex trait evolution.

Authors Contributions

Author contributions were defined according to the CRediT taxonomy (https://credit.niso.org/). Alejandro Valenzuela (affiliations 1,2) led data curation, formal analysis, investigation, methodology, visualization, validation, and contributed to the original draft as first author. Fabio Barteri (1,4,9) contributed to investigation, methodology, visualization, and writing as co–first author. Claudia Vasallo (4) carried out investigation and validation. Lukas Kuderna (10) and Joseph Orkin (11) contributed to data curation. Jean Boubli (12), Amanda Melin (13), Hafid Laayouni (1), Kyle Farh (10), and Jeffrey Rogers (14) provided supervision. Tomàs Marquès-Bonet (1) contributed to funding acquisition and supervision. Gerard Muntané (5) contributed to conceptualization and methodology and served as co–corresponding author. Arcadi Navarro (1,2,3,5,6,7,8,9) contributed to conceptualization, funding acquisition, methodology, project administration, and resources, serving as co–corresponding author. David Juan (15) contributed to conceptualization and methodology and served as corresponding author. All authors participated in writing – review and editing.

1. Introduction.

Complex traits arise from the interplay of genetic and non-genetic factors. Genetic contributions are typically distributed across interacting coding and regulatory elements. This multilayered organization makes the genetic basis of phenotypic variation difficult to resolve. Much of our knowledge on the genetic architecture of complex traits originates from studies of model organisms (1–3) and human genome-wide association studies (GWAS) (4–7), which link genomic and phenotypic variation within populations. Although powerful, GWAS capture only segregating variation within contemporary populations and miss fixed substitutions along evolutionary lineages, changes that often underlie major phenotypic shifts and whose study may unveil loci of strong biological interest (8,9).

Macroevolutionary analyses address this gap by testing whether genetic elements are associated with traits across species. Phylogenetic Comparative Methods (PCMs) (9–12) exploit millions of years of divergence to reveal associations involving gene gain and loss, protein evolution rates, structural variants, and point mutations (13–16). PCMs have revealed the genetic architecture of key traits in human evolution, such as brain size (17,23,24) and maximum lifespan (25–28).

The growing availability of high-quality genomes now enables scalable, cross-species genome–phenome mapping. Leveraging high-quality assemblies from 224 primate species (29), we created the first phylogenetic protein-coding genome–phenome map (P3GMap), publicly available through the Primate Genome–Phenome Archive (PGA, https://pgarchive.github.io). This resource integrates 263 complex traits across primates and reveals thousands of coding variants associated with phenotypic evolution.

To illustrate the scope of the resource, we examined three case studies: insectivorous diet (30,118–120), white blood cell count (140,141), and maximum lifespan (25–28). For each of these traits, we defined phenotypic *Contrasts* as pairs of species from different lineages that show marked differences in trait values, and we used these to identify genes and individual substitutions consistently associated with the trait. These case studies exemplify how researchers can use the PGA to move from a trait of interest to candidate coding changes and pathways that may underlie its evolution.

2. Results

2.1 The Primate Phenomic Space

We compiled phenotypic data for 224 primate species by integrating the average species values of 263 quantitative and qualitative traits derived from over 800 measurements obtained from public sources (30–33). These traits were classified into five primary domains (behavior, 29 traits; ecology, 7; life history, 29; morphology, 93; and physiology, 67) and 15 secondary domains (Fig. 1, Methods, Table S1), thereby defining the primate phenomic space for comparative analyses. As data density varies across clades, comparisons should be constrained to sets of species with similar information. To do so, we applied supervised clustering to group traits into 13 categories based on phylogenetic sampling (Fig. 1, SM). Body mass-related traits were among the most consistently represented, explaining among 15–23% of the variance in the first component for clusters VIII–X (see Fig. 1A). To control for allometry (34–38), we corrected 74 morphological, 14 life history, and two physiology traits for *Adult Body Mass* (Data S1). The curated dataset is publicly available from the Primate Genome–Phenome Archive (PGA, https://github.com/pgarchive/data.2.2).

We conducted a genome-phenome analysis of protein-coding sequences to create the first P3GMap (also available on the PGA web). This resource catalogues associations between protein-coding genes and diverse phenotypes across primates. The map comprises alignments for 16,133 one-to-one orthologous genes, over 75% of which include more than 200 species (Fig. S1, SM). To detect genetic factors that correlate to phenotypic evolution, we applied two conceptually distinct approaches: Convergent Amino Acid Substitutions (CAAS) using CAAStools (40; SM CAAStools), and relative evolutionary rates (RERs) using RERconverge (39; SM RERconverge). These methods capture genome-phenome associations across a broad range of evolutionary scenarios, beyond classical positive selection signatures (9,11,13). To ensure consistency and scalability, we implemented a standardized strategy for cross-trait comparisons, providing a unified framework for mapping genome-phenome relationships in primates.

2.2.1 Convergent Amino Acid Substitutions Associated with Phenotypic Diversity

For each trait, we identified CAASs between species with diverging trait values. For continuous traits, we selected species with higher trait values (*Top*) and species with lower trait values (*Bottom*) (see Fig. 1B–C, Figs. S24–S27, Data S1, and for details SM A.1–A.4). For categorical traits, we compared species annotated with different labels (*e.g.*, *Diet*: gummivore vs insectivore). To ensure independence, the standard strategy emphasized *Contrasts* within families. Of the analyzed traits, 83 (37%) showed at least one intra-family

Contrast, and 63 showed at least one *Contrast* in two or more families. Traits in widely distributed clusters (WD) more frequently exhibited multiple *Contrasts (~23%)*. In contrast, intermediately distributed (ID; ~60%) and Strepsirrhine-centered (STR; ~54%) clusters showed a high fraction of traits without *Contrasts*, reflecting both coverage limitations and the predominance of inter-family over intra-family variation in quantitative traits.

To apply CAAS and RER methods, we constructed a dataset of curated multiple sequence alignments of one-to-one orthologs extracted from a recently sequenced set of 233 primate genomes (29). Using CAAStools for these alignments with phylogenetic correction via permulations (41) and per-trait multiple testing (SM A.4), we detected 29,155 significant CAASs in 8,780 genes across 76 traits ($p_{permul-FDR} < 0.05$). This yielded in a median of 49 genes and 51 positions per trait (Fig. 2A, Figs. S28–S31, Data S2), providing a large-scale resource for studying candidate molecular convergence. These associations reflect fixed amino acid substitutions accumulated along evolutionary lineages rather than segregating variation within species.

We assessed the consistency of these discoveries using the set of primate species not involved in the discovery process, employing *PGLS* for quantitative traits and *phyloglm* for qualitative traits (see SM Internal phylogenetic test in primates; Fig. 2C, Fig. S39). This approach is conservative because it excludes extreme species from the consistency-checking step. As a quality control step, we focused on the ten traits with fewer than 100 validated associations, reducing the impact of non-specific associations ($p_{PGLS-FDR} < 0.05$, $p_{phyloglm-FDR} < 0.05$), recovering both known signals and novel candidate associations (Suppl. Table S3). Representative examples include associations between *Body Mass* and mitochondrial genes *NDUFS7* (substitutions Y223L, R230H, K231G, I232A; $p_{PGLS} = 1.37 \times 10^{-5}$–$3.57 \times 10^{-5}$) and *NOXO1* T/V223A ($p_{PGLS} = 2.06 \times 10^{-7}$); the immunoglobulin gene *ICAM1* R/V418M ($p_{PGLS} = 2.39 \times 10^{-5}$) (52); also the association of *Lactation Period Length* with *MEFV* K/R546Q ($p_{PGLS} = 1.66 \times 10^{-6}$), a regulator of innate immunity and familial Mediterranean fever (53–55). Additionally, we tested ten equivalent traits in non-primate mammals to determine whether their 4,968 CAASs exhibited consistent effects in the mammal phylogeny. Despite being highly conservative, this test yielded far more consistency than expected by chance ($p_{binomial}$: 4.07×10e−6, see SM, Suppl. Table S4). Representative cross-mammalian examples include *NELL1* Q176RN for *Lactation Period Length* ($p_{PGLS} = 1.09 \times 10^{-4}$), which has been previously associated with delayed milk production in goats (56) and to cranial bone development (57–59), and *MPDZ* H/P956Q for *Gestation Period Length* ($p_{PGLS} = 1.65 \times 10^{-5}$), a gene whose mutations cause congenital non-communicating hydrocephalus (60,61).

2.2.2 Relative Evolutionary Rates and Trait Associations

While CAASs detect genome–phenome associations independently of the underlying mechanism, relative evolutionary rates (RERs) identify genes whose evolutionary rates covary with trait changes, which is consistent with positive or purifying selection. We applied RERconverge to 16,109 the multiple sequence alignments of orthologous coding genes and 200 quantitative traits (see SM 3.1.1, RERconverge), assessing significance using permulations (41; SM 3.1.1.1). This analysis yielded 3,911 genes significantly associated with 194 different traits ($p_{permul-FDR} < 0.05$), with a median of 19 genes per trait, summarized across domains (Fig. 2A, Figs. S28–S31, Data S2). Because RERconverge applies less conservative corrections than CAAStools (SM 3.1.1), we also required FDR-corrected parametric p-values, which were broadly consistent with permulation statistics (SM 3.1.1.1).

RERconverge revealed biologically relevant signals (SM 3.1.1, Table S2), including *NKX6-2* for *Percentage of Time Moving* (Rho = 0.62; $p_{permul-FDR} < 10^{-3}$) which has been linked to spastic ataxia (42,43); *DEFA5* for *Percentage of Fauna Ingest* (Rho = –0.5; $p_{permul-FDR} < 10^{-3}$) which is a defensin gene associated with gastritis (44); and *DESI2* for *Mesencephalon Relative Size* (Rho = 0.69; $p_{permul-FDR} < 10^{-3}$), which is implicated in amyotrophic lateral sclerosis type 5 (47).

2.3 Functional analysis of the P3GMap

We performed over-representation analyses to evaluate biological functions associated with the discovered P3GMap genes. Genes with significant CAAS were associated with 199 unique enriched terms across 27 traits, each exhibiting at least one enrichment (Suppl. Data S3). Body mass–related traits were enriched for DNA repair categories, consistent with previous findings in *Carnivora* (87) and with the identification of DNA repair genes as anti-cancer targets in long-lived species (28,48,88–90) (Suppl. Table S5). The trait *Percentage of Time with Social Activity* showed enrichment for chromosome breakage (ER = 7.21; $p = 4.53 \times 10^{-7}$) and Fanconi anemia (ER = 4.73; $p = 3.03 \times 10^{-5}$), which are linked to developmental disorders (91,92) (Fig. 2D-E).

Across 48 traits, 94 enriched terms were found for significant RERconverge genes. These included *Long Philtrum* (distance >2SD between nasal base and upper lip) which is associated with *Pallidum* (ER = 3.35; $p = 2.97 \times 10^{-6}$) and *Subthalamus Size* (ER = 3.34; $p = 3.12 \times 10^{-6}$). Nose morphology is a key distinguishing feature between Strepsirrhines and Haplorhines (62,63) and has been linked to human syndromes involving cognitive, sensory, and craniofacial abnormalities (64–68). Traits under sexual selection also showed enrichment: *Hair Dimorphism* with the MHC complex ($p = 7.76 \times 10^{-6}$), which has long been proposed to affect mate choice (73–77), although this remains debated in some taxa (78–82); and *Body Mass Dimorphism* with mammary

neoplasms (ER = 2.16; p = 4.65 $\times 10^{-6}$), which is consistent with the effects of sex-dimorphism on cancer incidence (83–85).

Examining trait relationships further refined functional interpretation. Gene set overlaps across domains (Fig. 2F) revealed 16 significant pairs for CAAS (p < 1 $\times 10^{-50}$), concentrated in *Trichromatic Color Vision* and *Brain/Body Mass*, suggesting pleiotropic effects across sensory and morphological systems. Five overlaps involved life history traits, such as *Age of Female at First Reproduction* with *Trichromatic Color Vision* and with *Maximum Young Adult Weight*; and *Gestation Period Length* with *Maximum Young Adult Weight* (including sex-specific values). These reproductive overlaps were prominent in Atelidae, where Ateles and Lagothrix show polymorphic color vision and Alouatta displays routine trichromacy (117), pointing to coordinated evolution of vision, reproduction, and somatic traits under ecological and social pressures.

RERconverge overlaps (11 pairs, p < 10^{-30}) emphasized longevity-related traits. Genes for *Percentage of Infant Mortality* overlapped with *Maximum Male Longevity* and *Gamma-Glutamyl Transferase Levels* (GGT) (p = 4.46 $\times 10^{-74}$, p = 7.13 $\times 10^{-39}$), an antioxidant enzyme linked to cardiovascular and neurological disease (100–106), lung inflammation (107), cancers (108–110), and mortality prediction (111). Genes linked to *Cholesterol Levels* overlapped with *Median Male Longevity* (p = 6.74 $\times 10^{-48}$), although cholesterol–mortality links remain debated (112). Genes linked to *Serum Oxaloacetic Transaminase Levels* (AST) overlapped with *Median Female Longevity* (p = 2.18 $\times 10^{-38}$); elevated AST is associated with mortality in liver, neurological, and cardiovascular diseases (113–116).

Family-level analyses revealed lineage-specific patterns. *Cathemeral Activity* (sporadic intense activity periods during day and night) shared gene sets with *Dichromatic Color Vision* (p = 7.72 $\times 10^{-49}$) and *Seasonal Breeding* (p = 2.34 $\times 10^{-200}$), mainly in Lemuridae. The trait *Solitary Social System* overlapped with *Monochromatic Color Vision* (p = 2.98 $\times 10^{-41}$) in Lorisidae and Galagidae, and with *Dichromatic Color Vision* (p = 9.85 $\times 10^{-48}$) in Lepilemuridae, Microcebus, and Nycticebus. These results suggest co-evolution of vision, reproduction, and social behavior in response to nocturnality, seasonality, and mating strategies.

Finally, we asked whether P3GMap-associated genes overlap with human GWAS loci. No significant enrichment was found, consistent with limited overlap between cross-species and within-population variation (SM 6.1–6.2). Moreover, 86.16% of CAAS positions are fixed in humans. This suggests a decoupling between macroevolutionary divergence and microevolutionary variation, reinforcing the complementarity of cross- and within-species approaches to disentangle the architecture of complex traits.

2.4 Targeted Case Studies of Complex Primate Traits

While the general analysis in the P3GMap framework enables consistent cross-trait comparisons across hundreds of phenotypes with RERconverge, CAAS, and other methods, tailored lineage-specific CAAS analyses can reveal mechanisms obscured in large-scale scans. To demonstrate this, we examined three unrelated traits from different domains: *Insectivorous Diet*, *White Blood Cell Count*, and *Maximum Lifespan*. First, *Diet* (a widely distributed trait) displayed one *Contrast* for insectivory, reflecting the propensity of broadly distributed, often qualitative traits to capture intra-family extremes. Secondly, *White Blood Cell Count* (Strepsirrhine-focused trait) presented two *Contrasts*, providing an example within a cluster that generally yields fewer contrasting cases. By comparison, *Maximum Lifespan* showed no contrasts, providing an example of a quantitative trait whose variation tends to manifest between families rather than within them. By tailoring species *Contrasts* to the particular phylogenetic distribution of each one of these traits, we identified additional gene sets that refined the functional interpretation and highlighted recurrent adaptive pathways. Thus, the case studies (Fig. 3A–C) demonstrate the potential of the PGA for trait-focused exploration, which complements the global view of the general analysis presented above. Additionally, to leverage the biological relevance of our approach, we also performed a natural selection analysis on these positions, revealing directional selection toward the adaptative amino acids (SM 3.2)

2.4.1 *Insectivorous Diet*

Diet is closely associated with traits such as *Brain Size* (118–120) and *Dentition* (121,122) in primates, including Strepsirrhines (120). For insectivory, the standard *Contrasts* involved Lorisidae (slender *vs.* slow lorises) and *Daubentonia madagascariensis* (Daubentiidae) as insectivorous, compared with non-insectivorous species from other families. In the tailored design, we included Tarsiidae as an additional insectivorous lineage alongside alternative non-insectivorous selections (Fig. 3A). This approach identified 619 positions across 537 genes, all of which were significant by permulations (Fig. 3D). These were enriched in four categories (pFDR < 0.05): apical part of the cell (GO:0045177, 19 genes), cargo receptor activity (GO:0038024, 8 genes), lipase activity (GO:0016298, 7 genes), and endopeptidase activity (GO:0004175, 14 genes). These categories were not detected in alternative species selections separately. These enrichments suggest that the epithelial surface is a hub for nutrient absorption, which is consistent with structural changes implicated in nutritional and inflammatory disorders (123–128). Among candidate genes, *Sucrase–Isomaltase* (*SI*) carried CAASs A/R/T917K across all species sets and I627M in those including Tarsiidae, both under directional selection. *SI* is essential for carbohydrate digestion, and loss-of-function variants in Arctic human populations cause sucrose intolerance in childhood (129–131). Lipases also featured prominently, including *ASPG* (p.G374D), *HPSE* (A/F/L44V), *PLA2G3* (L432P), and *PNLIP* (K394R), all of them subject to directional selection (132,133). Of

the 27 genes and 14 positions shared across selections, DEFB113 V52I was the only site under selection in all comparisons. DEFB113, a β-defensin gene central to host defense, has undergone repeated gain and loss in mammals (134–136), with diet suggested as a driver of diversification (137–139).

2.4.2 *White Blood Cells Count*

White Blood Cell Count (WBC) metrics have rarely been explored in macroevolutionary analyses, though they correlate with mating systems and sexually dimorphic traits such as *Testis Size* (140,141). We contrasted species with extreme WBC counts from Hominidae and Lorisidae (standard strategy) and added Callitrichidae, which present outlier values. This identified 1,901 positions in 1,191 genes with significant permulation results. Enrichments included olfactory receptor activity (GO:0004984), also recovered in the general P3GMap analysis above, and two novel extracellular matrix (ECM) categories—ECM constituent (GO:0005201) and ECM–receptor interaction (hsa04512). These ECM categories partially overlapped with olfactory receptor genes (9 and 6 genes, respectively) but were only detected in the analysis tailored for the case study. Olfactory receptors (ORs), while canonical sensory genes, are also expressed in immune cells such as macrophages (142) and leukemia cell lines (143–145). Four ORs were particularly noteworthy: *OR4D5* (I48T) under directional selection in Hominidae; *OR10G9* with CAASs R89K and F/Y114L in Callitrichidae and Lorisidae (pPGLS < 1 × 10^{-3}) plus four additional sites under selection; and OR52B6 and OR10AD1, the latter with L/M102I under directional selection. These findings suggest repeated co-option of ORs for immune as well as sensory roles. We also detected a xenobiotic metabolism cluster. *SULT1C3* carried four CAASs (K45R, H/R196Q, E227D, Q243Y) under directional selection in Callitrichidae and Hominidae, plus 19 further associations (pPGLS < 1 × 10^{-3}). CES2 showed a CAAS (L599F) with directional selection and consistent phylogeny-wide association (pPGLS = 5.46 × 10^{-4}). Both genes are central to detoxification and belong to neuro-immune pathways involved in xenobiotic avoidance, allergy, and immune dysregulation (146–148). Immune recognition genes also showed lineage-specific signals: LILRB3 under directional selection in Hominidae, LILRA6 under selection despite lacking CAASs, HLA-DQA1 with directional selection in Hominidae, and HLA-DPB1 with a CAAS (Q75R) in Callitrichidae and Lorisidae (pPGLS = 1.67 × 10^{-4}). Collectively, these results underscore the interplay of immune function, environmental exposure, and lineage-specific adaptation in shaping WBC variation.

2.4.3 *Maximum Lifespan*

Longevity is a central focus of comparative genomics, with humans living markedly longer than their closest relatives (48,149–151). After correcting for *Body Mass*, the *Contrasts* used in the general analysis we performed for the P3GMap included humans (Hominidae), long-lived Cebidae, and several short-lived lineages. For the tailored use case, we refined these comparisons by adding Hylobatidae as another long-lived

clade and including additional short-lived species (Fig. 3C). This analysis identified 30,886 positions in 8,565 genes (Fig. 3D). However, because lifespan *Contrasts* often span entire families, detection can be noisy and inflated. To refine this, we assessed consistency. In primates, from 62.72% of informative positions (containing *Top* and *Bottom* states across more than five additional species), the analysis yielded 5,225 positions in 3,338 genes with consistent associations. In mammals, this figure was 30.71% of informative positions, yielding 1,061 positions in 921 genes with consistent associations across primates and non-primate mammals. These curated sets provided a robust basis for functional interpretation. First, among primate-consistent genes, enrichment analysis revealed four categories, including chromosome segregation (GO:0007059) and complement/coagulation cascades (hsa04610). Eight categories emerged for mammal-consistent genes, including interleukin-10 production (GO:0032613) and the regulation of multi-organism processes (GO:0043900). Across both clades, 139 amino acid positions in 136 genes were consistently associated and were enriched in cranial nerve paralysis (HP:0006824) and abnormal cranial nerve physiology (HP:0031910), conditions linked to diabetes (152,153), ischemic stroke (154), or aneurysm (155). From the gene sets linked to these terms in this case study, seven overlapped genes in the general strategy, highlighting the added value of examining overlaps across different selection configurations.

Studying the overlap between all strategies with the most phylogenetically restrictive set of consistent genes, six genes presented CAASs in all species selections. Notable examples include IL1B with CAAS L152M, a key regulator of immune and inflammatory responses (96,156,157), complemented with the detection of two other interleukines, IL12B and IL4I1. Additionally, we also recovered IL23R and IL5 in the directional selection analysis. A second example to highlight is *KLK4* Q227P (Kallikrein-related peptidase 4), known for exerting tumorigenic effects through activating growth factors like insulin-like growth factor (IGF), detected in comparative studies searching for genomic correlates of long-lived species (28,158). We also detected CAASs and directional selection in KLK1 P57R in Hylobatidae and Hominidae, though this site was not consistently associated across primates or mammals. Together, these results implicate immunity, inflammation, and growth factor pathways as recurrent contributors to lifespan evolution.

3. Discussion

This study presents the first P3GMap, which integrates 263 complex traits across 224 primate species. By combining CAAS (40) and RERs (39), we uncovered thousands of candidate associations linking coding variation to phenotypic evolution. The PGA and the associated P3GMap offer a unified resource for studying the genetic basis of traits relating to ecology, morphology, life history, and physiology. This kind of approach complements within-species studies such as GWAS (4–7) by capturing evolutionary signals that are fixed

along lineages (8,9), as illustrated by the observation that ≈86% of CAAS positions are fixed in humans. This reinforces the decoupling between macroevolutionary divergence, which establishes species-specific trait differences, and microevolutionary variation, which can be readily observed in within-species studies (28,48,88–90).

Our framework, has limitations. It focuses on single-copy protein-coding genes (29), excluding regulatory variation, gene duplications, and structural variants, all of which are also important for trait evolution (164). CAAS and RERconverge rely on distinct statistical and conceptual foundations, which can result in partially different gene–trait associations (39,40). Although permulation tests are rigorous (41), they are based on Brownian motion models and may not fully capture all evolutionary dynamics (11,159,160). These caveats highlight the need for future expansions that incorporate regulatory, structural, and expression data (164), as well as refined evolutionary models (11,159,160).

Despite these constraints, our results demonstrate the power of macroevolutionary analyses to illuminate the genetic architecture of complex traits (13–16). The P3GMap reveals recurrent pathways ranging from immunity, inflammation, and DNA repair, to growth factor signaling, linked to phenotypic change, echoing prior evidence from long-lived mammals (28,48,88–90) and other comparative frameworks (25–28,87). Our case studies connect genomic signatures to insectivory (118–122), immunity and WBC variation (140–145), and lifespan (25–28,149–151), illustrating the value of targeted *C*ontrasts for dissecting lineage-specific adaptations. As an open and scalable platform, the PGA provides a foundation for extending genome–phenome inference beyond protein-coding variation, strengthening the link between comparative genomics and human biomedical research.

4. Figures

Figure 1. Overview of Primate Phenomic Data.

(**A**) Hierarchical clustering of phenomic data, mapped onto the primate phylogeny and grouped by the primate family. Based on data availability, the traits were classified into 13 data clusters [see Supplementary Material, Trait clustering by phylogenetic sampling]. The area within the square highlights traits from one of the clusters, Cluster VIII, which includes a broad range of ecological, life history, and morphological traits. (**B**) Total number of traits (x-axis) showing extreme phenotypic differences between species within the same family. Top and Bottom values refer to species with contrasting trait values (see text for definitions). (**C**) Examples

illustrating the distribution of values for three traits: Body Mass (left), Maximum Lifespan (center), and Gummivore, sap-eating, Diet (right). Each point represents a species. The vertical bars indicate median family values. Dashed lines correspond to the median global trait value. Species selected as Top and Bottom extremes (based on criteria explained in the text) are indicated by triangles and squares, respectively.

Figure 2. Exploring the Phylogenetic Protein-coding Genome–Phenome Map (P3GMap).

(**A**) Gene–trait associations recovered across secondary biological domains using RERs (left) and CAAS (right), showing the median number of significant genes per trait (log10 scale) and the number of traits with associations (center). (**B**, **C**) Manhattan plots highlighting top gene–trait associations detected by

RERconverge and CAAS, with significance thresholds indicated (green, FDR; red, Bonferroni). Genes surpassing strict thresholds are labeled. (**D & E**) Overlap of significant gene sets across secondary domains, with strip-lined boxes marking domain pairs without shared associations. . (**F**) Heatmap showing the overlap between functional enrichments identified by CAAS and RER across traits, highlighting pathways recurrently implicated by both approaches.

Figure 3. Case studies illustrating the CAAS selection approach for Insectivorous Diet, Maximum Lifespan, and White Blood Cell Count.

(**A–C**) Selection of Top species (triangles) and Bottom species (colored lines) using the standard genome–phenome strategy (*S-*) and two alternative approaches (*A-* and *B-*). Pie charts show ancestral states for qualitative traits, while internal nodes with strong quantitative shifts are depicted as posterior distributions (histograms). (**D**) Venn diagrams of gene overlap (p-perm < 0.05) between alternative family selections for

each trait. (**E**) Counts of sites under directional selection per trait; stray lines indicate CAAS positions also showing positive directional selection.

5. Bibliography

1.   Ogienko AA, Omelina ES, Bylino OV, Batin MA, Georgiev PG, Pindyurin AV. Drosophila as a Model Organism to Study Basic Mechanisms of Longevity. Int J Mol Sci. 2022 Sep 24;23(19):11244.

2.   Zhang S, Li F, Zhou T, Wang G, Li Z. Caenorhabditis elegans as a Useful Model for Studying Aging Mutations. Front Endocrinol [Internet]. 2020 [cited 2024 Jan 17];11. Available from: https://www.frontiersin.org/articles/10.3389/fendo.2020.554994

3.   Irion U, Nüsslein-Volhard C. Developmental genetics with model organisms. Proc Natl Acad Sci. 2022 Jul 26;119(30):e2122148119.

4.   Sakaue S, Kanai M, Tanigawa Y, Karjalainen J, Kurki M, Koshiba S, et al. A cross-population atlas of genetic associations for 220 human phenotypes. Nat Genet. 2021 Oct;53(10):1415–24.

5.   Watanabe K, Stringer S, Frei O, Umićević Mirkov M, de Leeuw C, Polderman TJC, et al. A global overview of pleiotropy and genetic architecture in complex traits. Nat Genet. 2019 Sep;51(9):1339–48.

6.   Liu X, Tian D, Li C, Tang B, Wang Z, Zhang R, et al. GWAS Atlas: an updated knowledgebase integrating more curated associations in plants and animals. Nucleic Acids Res. 2023 Jan 6;51(D1):D969–76.

7.   Canela-Xandri O, Rawlik K, Tenesa A. An atlas of genetic associations in UK Biobank. Nat Genet. 2018 Nov;50(11):1593–9.

8.   de Magalhães JP, Wang J. The fog of genetics: what is known, unknown and unknowable in the genetics of complex traits and diseases. EMBO Rep. 2019 Nov 5;20(11):e48054.

9.   Smith SD, Pennell MW, Dunn CW, Edwards SV. Phylogenetics is the New Genetics (for Most of Biodiversity). Trends Ecol Evol. 2020 May 1;35(5):415–25.

10.   Garamszegi LZ, editor. Modern Phylogenetic Comparative Methods and Their Application in Evolutionary Biology: Concepts and Practice [Internet]. Berlin, Heidelberg: Springer; 2014 [cited 2023 Sep 25]. Available from: http://link.springer.com/10.1007/978-3-662-43550-2

11.   Nagy LG, Merényi Z, Hegedüs B, Bálint B. Novel phylogenetic methods are needed for understanding gene function in the era of mega-scale genome sequencing. Nucleic Acids Res. 2020 Mar 18;48(5):2209–19.

12.   Harmon L. Phylogenetic Comparative Methods: Learning From Trees. 2019 May 20 [cited 2023 Sep 27]; Available from: https://ecoevorxiv.org/repository/view/4486/

13.   Hiller M, Schaar BT, Indjeian VB, Kingsley DM, Hagey LR, Bejerano G. A "forward genomics" approach links genotype to phenotype using independent phenotypic losses among related species. Cell Rep. 2012 Oct 25;2(4):817–23.

14.   De Bie T, Cristianini N, Demuth JP, Hahn MW. CAFE: a computational tool for the study of gene family evolution. Bioinformatics. 2006 May 15;22(10):1269–71.

15.   Yan H, Hui Z, Thomas GWC, Edwards SV, Sackton TB, Liu JS. PhyloAcc-GT: A Bayesian method for inferring patterns of substitution rate shifts on targeted lineages accounting for gene tree discordance. Mol Biol Evol. 2023 Sep 4;msad195.

16.   Rey C, Guéguen L, Sémon M, Boussau B. Accurate Detection of Convergent Amino-Acid Evolution with PCOC. Mol Biol Evol. 2018 Sep 1;35(9):2296–306.

17.   DeCasien AR, Trujillo AE, Janiak MC, Harshaw EP, Caes ZN, Galindo GA, et al. Equivocal evidence for a link between megalencephaly-related genes and primate brain size evolution. Sci Rep. 2022 Jun 28;12(1):10902.

18.   Sun Y, Liu Y, Sun X, Lin Y, Yin D, Xu S, et al. Insights into body size variation in cetaceans from the evolution of body-size-related genes. BMC Evol Biol. 2019 Jul 27;19(1):157.

19.   Chen L, Qiu Q, Jiang Y, Wang K, Lin Z, Li Z, et al. Large-scale ruminant genome sequencing provides insights into their evolution and distinct traits. Science. 2019 Jun 21;364(6446):eaav6202.

20.   Silva FA, Souza ÉMS, Ramos E, Freitas L, Nery MF. The molecular evolution of genes previously associated with large sizes reveals possible pathways to cetacean gigantism. Sci Rep. 2023 Jan 19;13(1):67.

21.   Xu S, Sun X, Niu X, Zhang Z, Tian R, Ren W, et al. Genetic basis of brain size evolution in cetaceans: insights from adaptive evolution of seven primary microcephaly (MCPH) genes. BMC Evol Biol. 2017 Aug 29;17(1):206.

22.   Shao Y, Zhou L, Li F, Zhao L, Zhang BL, Shao F, et al. Phylogenomic analyses provide insights into primate evolution. Science. 2023 Jun 2;380(6648):913–24.

23.   Montgomery SH, Capellini I, Venditti C, Barton RA, Mundy NI. Adaptive Evolution of Four Microcephaly Genes and the Evolution of Brain Size in Anthropoid Primates. Mol Biol Evol. 2011 Jan 1;28(1):625–38.

24.   Montgomery SH, Mundy NI. Microcephaly genes evolved adaptively throughout the evolution of eutherian mammals. BMC Evol Biol. 2014 Jun 5;14(1):120.

25.   Farré X, Molina R, Barteri F, Timmers PRHJ, Joshi PK, Oliva B, et al. Comparative Analysis of Mammal Genomes Unveils Key Genomic Variability for Human Life Span. Mol Biol Evol. 2021 Oct 27;38(11):4948–61.

26.   Muntané G, Farré X, Rodríguez JA, Pegueroles C, Hughes DA, de Magalhães JP, et al. Biological Processes Modulating Longevity across Primates: A Phylogenetic Genome-Phenome Analysis. Mol Biol Evol. 2018 Aug 1;35(8):1990–2004.

27.   Kowalczyk A, Partha R, Clark NL, Chikina M. Pan-mammalian analysis of molecular constraints underlying extended lifespan. Kaeberlein M, Tautz D, editors. eLife. 2020 Feb 11;9:e51089.

28.   Yu Z, Seim I, Yin M, Tian R, Sun D, Ren W, et al. Comparative analyses of aging-related genes in long-lived mammals provide insights into natural longevity. The Innovation. 2021 May 28;2(2):100108.

29.   Kuderna LFK, Gao H, Janiak MC, Kuhlwilm M, Orkin JD, Bataillon T, et al. A global catalog of whole-genome diversity from 233 primate species. Science. 2023 Jun 2;380(6648):906–13.

30.   DeCasien AR, Higham JP. Primate mosaic brain evolution reflects selection on sensory and cognitive specialization. Nat Ecol Evol. 2019 Oct;3(10):1483–93.

31.   Kemnitz JW. DATABASE FOR INDICES OF AGING IN NONHUMAN PRIMATES. Innov Aging. 2019 Nov 8;3(Suppl 1):S957.

32.   Galán-Acedo C, Arroyo-Rodríguez V, Andresen E, Arasa-Gisbert R. Ecological traits of the world's primates. Sci Data. 2019 May 13;6(1):55.

33.   Zehr SM, Roach RG, Haring D, Taylor J, Cameron FH, Yoder AD. Life history profiles for 27 strepsirrhine primate taxa generated using captive data from the Duke Lemur Center. Sci Data. 2014;1:140019.

34.   Simmen B, Morino L, Blanc S, Garcia C. The energy allocation trade-offs underlying life history traits in hypometabolic strepsirhines and other primates. Sci Rep. 2021 Jul 9;11(1):14196.

35.   Isler K, van Schaik CP. Allomaternal care, life history and brain size evolution in mammals. J Hum Evol. 2012 Jul 1;63(1):52–63.

36.   Antoł A, Kozłowski J. Scaling of organ masses in mammals and birds: phylogenetic signal and implications for metabolic rate scaling. ZooKeys. 2020 Nov 2;982:149–59.

37.   Burger JR, George MA Jr, Leadbetter C, Shaikh F. The allometry of brain size in mammals. J Mammal. 2019 Apr 24;100(2):276–83.

38.   Sansalone G, Allen K, Ledogar JA, Ledogar S, Mitchell DR, Profico A, et al. Variation in the strength of allometry drives rates of evolution in primate brain shape. Proc R Soc B Biol Sci. 2020 Jul 8;287(1930):20200807.

39.   Kowalczyk A, Meyer WK, Partha R, Mao W, Clark NL, Chikina M. RERconverge: an R package for associating evolutionary rates with convergent traits. Bioinformatics. 2019 Nov 15;35(22):4815–7.

40.   Barteri F, Valenzuela A, Farré X, Juan D de, Muntané G, Esteve-Altava B, et al. CAAStools: a toolbox to identify and test Convergent Amino Acid Substitutions [Internet]. bioRxiv; 2023 [cited 2023 Sep 25]. p. 2022.12.14.520422. Available from: https://www.biorxiv.org/content/10.1101/2022.12.14.520422v3

41.   Saputra E, Kowalczyk A, Cusick L, Clark N, Chikina M. Phylogenetic Permulations: A Statistically Rigorous Approach to Measure Confidence in Associations in a Phylogenetic Context. Mol Biol Evol. 2021 Jul 1;38(7):3004–21.

42.   Chelban V, Patel N, Vandrovcova J, Zanetti MN, Lynch DS, Ryten M, et al. Mutations in NKX6-2 Cause Progressive Spastic Ataxia and Hypomyelination. Am J Hum Genet. 2017 Jun 1;100(6):969–77.

43.   Toch M, Harris A, Schakman O, Kondratskaya E, Boulland JL, Dauguet N, et al. Onecut-dependent Nkx6.2 transcription factor expression is required for proper formation and activity of spinal locomotor circuits. Sci Rep. 2020 Jan 22;10(1):996.

44.   Wu E, Zhu J, Ma Z, Tuo B, Terai S, Mizuno K, et al. Gastric alarmin release: A warning signal in the development of gastric mucosal diseases. Front Immunol [Internet]. 2022 [cited 2023 Oct 17];13. Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2022.1008047

45.   Trieschmann G, Wilhelm C, Berweck S, Zech M. De novo retinoic acid receptor beta (RARB) variant associated with microphthalmia and dystonia. Eur J Med Genet. 2023 Aug 1;66(8):104802.

46.   Srour M, Chitayat D, Caron V, Chassaing N, Bitoun P, Patry L, et al. Recessive and Dominant Mutations in Retinoic Acid Receptor Beta in Cases with Microphthalmia and Diaphragmatic Hernia. Am J Hum Genet. 2013 Oct 3;93(4):765–72.

47.   Foran E, Rosenblum L, Bogush AI, Trotti D. Sumoylation of critical proteins in amyotrophic lateral sclerosis: emerging pathways of pathogenesis. Neuromolecular Med. 2013 Dec;15(4):10.1007/s12017-013-8262–x.

48.   Tejada-Martinez D, Avelar RA, Lopes I, Zhang B, Novoa G, de Magalhães JP, et al. Positive Selection and Enhancer Evolution Shaped Lifespan and Body Mass in Great Apes. Mol Biol Evol. 2022 Feb 3;39(2):msab369.

49.   Orkin JD, Montague MJ, Tejada-Martinez D, de Manuel M, Del Campo J, Cheves Hernandez S, et al. The genomics of ecological flexibility, large brains, and long lives in capuchin monkeys revealed with fecalFACS. Proc Natl Acad Sci U S A. 2021 Feb 16;118(7):e2010632118.

50.   Benssouina FZ, Parat F, Villard C, Leloup L, Garrouste F, Sabatier J marc, et al. Overexpression of a Novel Noxo1 Mutant Increases Ros Production and Noxo1 Relocalisation. Int J Mol Sci. 2023 Feb 28;24(5):4663.

51.   Burch TC, Rhim JS, Nyalwidhe JO. Mitochondria Biogenesis and Bioenergetics Gene Profiles in Isogenic Prostate Cells with Different Malignant Phenotypes. BioMed Res Int. 2016;2016:1785201.

52.   Bui TM, Wiesolek HL, Sumagin R. ICAM-1: A master regulator of cellular responses in inflammation, injury resolution, and tumorigenesis. J Leukoc Biol. 2020 Sep 1;108(3):787–99.

53.   Makay B, Unsal E. Does breast-feeding affect severity of familial Mediterranean fever? Clin Rheumatol. 2009 Dec;28(12):1389–93.

54.   Özen S, Batu ED, Demir S. Familial Mediterranean Fever: Recent Developments in Pathogenesis and New Recommendations for Management. Front Immunol [Internet]. 2017 [cited 2023 Sep 26];8. Available from: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5362626/

55.   Yanmaz MN, Özcan AJ, Savan K. The impact of familial Mediterranean fever on reproductive system. Clin Rheumatol. 2014 Oct;33(10):1385–8.

56.   Massender E, Oliveira HR, Brito LF, Maignel L, Jafarikia M, Baes CF, et al. Genome-wide association study for milk production and conformation traits in Canadian Alpine and Saanen dairy goats. J Dairy Sci. 2023 Feb 1;106(2):1168–89.

57.   Zhang X, Zara J, Siu RK, Ting K, Soo C. The Role of NELL-1, a Growth Factor Associated with Craniosynostosis, in Promoting Bone Regeneration. J Dent Res. 2010 Sep;89(9):865–78.

58.   Lai K, Xi Y, Du X, Jiang Z, Li Y, Huang T, et al. Activation of Nell-1 in BMSC Sheet Promotes Implant Osseointegration Through Regulating Runx2/Osterix Axis. Front Cell Dev Biol [Internet]. 2020 [cited 2023 Sep 26];8. Available from: https://www.frontiersin.org/articles/10.3389/fcell.2020.00868

59.   Chen W, Zhang X, Siu RK, Chen F, Shen J, Zara JN, et al. Nfatc2 Is a Primary Response Gene of Nell-1 Regulating Chondrogenesis in ATDC5 Cells. J Bone Miner Res. 2011 Jun;26(6):1230–41.

60.   Abebe MS, Seyoum G, Emamu B, Teshome D. Congenital Hydrocephalus and Associated Risk Factors: An Institution-Based Case–Control Study, Dessie Town, North East Ethiopia. Pediatr Health Med Ther. 2022 May 11;13:175–82.

61.   Tully HM, Dobyns WB. Infantile hydrocephalus: a review of epidemiology, classification and causes. Eur J Med Genet. 2014 Aug;57(8):359–68.

62.   Rossie JB, Smith TD. Ontogeny of the nasolacrimal duct in primates: functional and phylogenetic implications. J Anat. 2007 Feb;210(2):195–208.

63.   Rossie JB, Smith TD, Beard KC, Godinot M, Rowe TB. Nasolacrimal anatomy and haplorhine origins. J Hum Evol. 2018 Jan 1;114:176–83.

64.   Low KJ, Ansari M, Abou Jamra R, Clarke A, El Chehadeh S, FitzPatrick DR, et al. PUF60 variants cause a syndrome of ID, short stature, microcephaly, coloboma, craniofacial, cardiac, renal and spinal features. Eur J Hum Genet EJHG. 2017 May;25(5):552–9.

65.   Popova S, Charness ME, Burd L, Crawford A, Hoyme HE, Mukherjee RAS, et al. Fetal alcohol spectrum disorders. Nat Rev Dis Primer. 2023 Feb 23;9(1):1–21.

66.   Pehlivan D, Karaca E, Aydin H, Beck CR, Gambin T, Muzny DM, et al. Whole-exome sequencing links TMCO1 defect syndrome with cerebro-facio-thoracic dysplasia. Eur J Hum Genet EJHG. 2014 Sep;22(9):1145–8.

67.   Verloes A, Di Donato N, Masliah-Planchon J, Jongmans M, Abdul-Raman OA, Albrecht B, et al. Baraitser–Winter cerebrofrontofacial syndrome: delineation of the spectrum in 42 cases. Eur J Hum Genet. 2015 Mar;23(3):292–301.

68.   Kline AD, Moss JF, Selicorni A, Bisgaard AM, Deardorff MA, Gillett PM, et al. Diagnosis and management of Cornelia de Lange syndrome: first international consensus statement. Nat Rev Genet. 2018 Oct;19(10):649–66.

69.   Ockeloen CW, Willemsen MH, de Munnik S, van Bon BWM, de Leeuw N, Verrips A, et al. Further delineation of the KBG syndrome phenotype caused by ANKRD11 aberrations. Eur J Hum Genet EJHG. 2015 Sep;23(9):1176–85.

70.   Guo L, Elcioglu NH, Wang Z, Demirkol YK, Isguven P, Matsumoto N, et al. Novel and recurrent COL11A1 and COL2A1 mutations in the Marshall-Stickler syndrome spectrum. Hum Genome Var. 2017;4:17040.

71.   Panarotto M, Davidson IF, Litos G, Schleiffer A, Peters JM. Cornelia de Lange syndrome mutations in NIPBL can impair cohesin-mediated DNA loop extrusion. Proc Natl Acad Sci U S A. 2022 May 3;119(18):e2201029119.

72.   Fusco C, Micale L, Augello B, Teresa Pellico M, Menghini D, Alfieri P, et al. Smaller and larger deletions of the Williams Beuren syndrome region implicate genes involved in mild facial phenotype, epilepsy and autistic traits. Eur J Hum Genet EJHG. 2014 Jan;22(1):64–70.

73.   E M, Song X, Wang L, Yang Y, Wei X, Yu J, et al. Mate choice for major histocompatibility complex (MHC) complementarity in the Yellow-rumped Flycatcher (Ficedula zanthopygia). Avian Res. 2021 May 29;12(1):27.

74.   Havlíček J, Winternitz J, Roberts SC. Major histocompatibility complex-associated odour preferences and human mate choice: near and far horizons. Philos Trans R Soc Lond B Biol Sci. 2020 Jun 8;375(1800):20190260.

75.   Roved J, Hansson B, Tarka M, Hasselquist D, Westerdahl H. Evidence for sexual conflict over major histocompatibility complex diversity in a wild songbird. Proc R Soc B Biol Sci. 2018 Aug;285(1884):20180841.

76.   Han QH, Sun RN, Yang HQ, Wang ZW, Wan QH, Fang SG. MHC class I diversity predicts non-random mating in Chinese alligators (Alligator sinensis). Heredity. 2019 Jun;122(6):809–18.

77.   Sommer S. The importance of immune gene variability (MHC) in evolutionary ecology and conservation. Front Zool. 2005 Oct 20;2:16.

78.   Auer KE, Primus J, Istel S, Dahlhoff M, Rülicke T. Mate genetic similarity affects mating behaviour but not maternal investment in mice. Sci Rep. 2023 Jun 29;13(1):10536.

79.   Achorn AM, Rosenthal GG. It's Not about Him: Mismeasuring 'Good Genes' in Sexual Selection. Trends Ecol Evol. 2020 Mar 1;35(3):206–19.

80.   Thonhauser KE, Raveh S, Penn DJ. Multiple paternity does not depend on male genetic diversity. Anim Behav. 2014 Jul;93(100):135–41.

81.   Amo L, Paz GA de, Kabbert J, Machordom A. House sparrows do not exhibit a preference for the scent of potential partners with different MHC-I diversity and genetic distances. PLOS ONE. 2022 Dec 21;17(12):e0278892.

82.   Westerdahl H. No evidence of an MHC-based female mating preference in great reed warblers. Mol Ecol. 2004 Aug;13(8):2465–70.

83.   He F, Furones AR, Landegren N, Fuxe J, Sarhan D. Sex dimorphism in the tumor microenvironment - From bench to bedside and back. Semin Cancer Biol. 2022 Nov;86(Pt 3):166–79.

84.   Zheng D, Trynda J, Williams C, Vold JA, Nguyen JH, Harnois DM, et al. Sexual dimorphism in the incidence of human cancers. BMC Cancer. 2019 Jul 12;19(1):684.

85.   Rubin JB. The spectrum of sex differences in cancer. Trends Cancer. 2022 Apr;8(4):303–15.

86.   Ly D, Forman D, Ferlay J, Brinton LA, Cook MB. An International Comparison of Male and Female Breast Cancer Incidence Rates. Int J Cancer J Int Cancer. 2013 Apr 15;132(8):1918–26.

87.   Huang X, Sun D, Wu T, Liu X, Xu S, Yang G. Genomic insights into body size evolution in Carnivora support Peto's paradox. BMC Genomics. 2021 Jun 9;22(1):429.

88.   Ma S, Upneja A, Galecki A, Tsai YM, Burant CF, Raskind S, et al. Cell culture-based profiling across mammals reveals DNA repair and metabolism as determinants of species longevity. Wagers AJ, editor. eLife. 2016 Nov 22;5:e19130.

89.   Zhang G, Cowled C, Shi Z, Huang Z, Bishop-Lilly KA, Fang X, et al. Comparative analysis of bat genomes provides insight into the evolution of flight and immunity. Science. 2013 Jan 25;339(6118):456–60.

90.   Keane M, Semeiks J, Webb AE, Li YI, Quesada V, Craig T, et al. Insights into the evolution of longevity from the bowhead whale genome. Cell Rep. 2015 Jan 6;10(1):112–22.

91.   Bogliolo M, Surrallés J. Fanconi anemia: a model disease for studies on human genetics and advanced therapeutics. Curr Opin Genet Dev. 2015 Aug 1;33:32–40.

92.   Helbling-Leclerc A, Garcin C, Rosselli F. Beyond DNA repair and chromosome instability-Fanconi anaemia as a cellular senescence-associated syndrome. Cell Death Differ. 2021 Apr;28(4):1159–73.

93.   Geremia A, Biancheri P, Allan P, Corazza GR, Di Sabatino A. Innate and adaptive immunity in inflammatory bowel disease. Autoimmun Rev. 2014 Jan;13(1):3–10.

94.   Nie S, Yuan Y. The Role of Gastric Mucosal Immunity in Gastric Diseases. J Immunol Res. 2020 Jul 24;2020:7927054.

95.   Kelley N, Jeltema D, Duan Y, He Y. The NLRP3 Inflammasome: An Overview of Mechanisms of Activation and Regulation. Int J Mol Sci. 2019 Jul 6;20(13):3328.

96.   Boraschi D, Italiani P, Weil S, Martin MU. The family of the interleukin-1 receptors. Immunol Rev. 2018 Jan;281(1):197–232.

97.   Huang J, Huffman JE, Huang Y, Do Valle Í, Assimes TL, Raghavan S, et al. Genomics and phenomics of body mass index reveals a complex disease network. Nat Commun. 2022 Dec 29;13(1):7973.

98.   Kuningas M, Altmäe S, Uiterlinden AG, Hofman A, van Duijn CM, Tiemeier H. The relationship between fertility and lifespan in humans. Age. 2011 Dec;33(4):615–22.

99.   Gorbunova V, Seluanov A, Zhang Z, Gladyshev VN, Vijg J. Comparative genetics of longevity and cancer: insights from long-lived rodents. Nat Rev Genet. 2014 Aug;15(8):531–40.

100.   Cho EJ, Jeong SM, Chung GE, Yoo JJ, Cho Y, Lee KN, et al. Gamma-glutamyl transferase and risk of all-cause and disease-specific mortality: a nationwide cohort study. Sci Rep. 2023 Jan 31;13(1):1751.

101.   Koenig G, Seneff S. Gamma-Glutamyltransferase: A Predictive Biomarker of Cellular Antioxidant Inadequacy and Disease Risk. Dis Markers. 2015;2015:818570.

102.   Kalirawna TR, Rohilla J, Bairwa SS, Gothwal SK, Tak P, Jain R. Increased concentration of serum gamma-glutamyl transferase in ischemic stroke patients. Brain Circ. 2021;7(2):71–6.

103.   Ndrepepa G, Kastrati A. Gamma-glutamyl transferase and cardiovascular disease. Ann Transl Med. 2016 Dec;4(24):481.

104.   Zhou Y, Zhao S, Chen K, Hua W, Zhang S. Predictive value of gamma-glutamyltransferase for ventricular arrhythmias and cardiovascular mortality in implantable cardioverter-defibrillator patients. BMC Cardiovasc Disord. 2019 May 30;19(1):129.

105.   Li S, Liao X, Pan Y, Xiang X, Zhang Y. Gamma-glutamyl transferase levels are associated with the occurrence of post-stroke cognitive impairment: a multicenter cohort study. BMC Neurol. 2022 Feb 23;22(1):65.

106.   Ruttmann E, Brant LJ, Concin H, Diem G, Rapp K, Ulmer H, et al. Gamma-glutamyltransferase as a risk factor for cardiovascular disease mortality: an epidemiological investigation in a cohort of 163,944 Austrian adults. Circulation. 2005 Oct 4;112(14):2130–7.

107.   Sun D, Liu H, Ouyang Y, Liu X, Xu Y. Serum Levels of Gamma-Glutamyltransferase During Stable and Acute Exacerbations of Chronic Obstructive Pulmonary Disease. Med Sci Monit Int Med J Exp Clin Res. 2020 Oct 22;26:e927771-1-e927771-7.

108.   Akaydın SY, Salihoğlu EM, Güngör DG, Karanlık H, Demokan S. Correlation Between Gamma-Glutamyl Transferase Activity and Glutathione Levels in Molecular Subgroups of Breast Cancer. Eur J Breast Health. 2019 Dec 5;16(1):72–6.

109.   Xiao Y, Yang H, Lu J, Li D, Xu C, Risch HA. Serum gamma-glutamyltransferase and the overall survival of metastatic pancreatic cancer. BMC Cancer. 2019 Oct 29;19(1):1020.

110.   Hong SW, Lee HJ, Han K, Moon JM, Park S, Soh H, et al. Risk of gastrointestinal cancer in patients with an elevated level of gamma-glutamyltransferase: A nationwide population-based study. PloS One. 2021;16(2):e0245052.

111.   Du G, Song Z, Zhang Q. Gamma-glutamyltransferase is associated with cardiovascular and all-cause mortality: A meta-analysis of prospective cohort studies. Prev Med. 2013 Jul 1;57(1):31–7.

112.   Ridker PM. LDL cholesterol: controversies and future therapeutic directions. Lancet Lond Engl. 2014 Aug 16;384(9943):607–17.

113.   Ndrepepa G. Aspartate aminotransferase and cardiovascular disease—a narrative review. J Lab Precis Med [Internet]. 2021 Jan 30 [cited 2024 Jan 18];6(0). Available from: https://jlpm.amegroups.org/article/view/5898

114.   Ke P, Zhong L, Peng W, Xu M, Feng J, Tian Q, et al. Association of the serum transaminase with mortality among the US elderly population. J Gastroenterol Hepatol. 2022 May;37(5):946–53.

115.   Xie K, Chen CH, Tsai SP, Lu PJ, Wu H, Zeng Y, et al. Loss of Life Expectancy by 10 Years or More From Elevated Aspartate Aminotransferase: Finding Aspartate Aminotransferase a Better Mortality Predictor for All-Cause and Liver-Related than Alanine Aminotransferase. Off J Am Coll Gastroenterol ACG. 2019 Sep;114(9):1478.

116.   Ndrepepa G, Holdenrieder S, Cassese S, Xhepa E, Fusaro M, Laugwitz KL, et al. Aspartate aminotransferase and mortality in patients with ischemic heart disease. Nutr Metab Cardiovasc Dis. 2020 Nov 27;30(12):2335–42.

117.   Henriques LD, Hauzman E, Bonci DMO, Chang BSW, Muniz JAPC, da Silva Souza G, et al. Uniform trichromacy in Alouatta caraya and Alouatta seniculus: behavioural and genetic colour vision evaluation. Front Zool. 2021 Jul 8;18(1):36.

118.   DeCasien AR, Williams SA, Higham JP. Primate brain size is predicted by diet but not sociality. Nat Ecol Evol. 2017 Mar 27;1(5):1–7.

119.   Grabowski M, Kopperud BT, Tsuboi M, Hansen TF. Both Diet and Sociality Affect Primate Brain-Size Evolution. Syst Biol. 2023 Mar 1;72(2):404–18.

120.   López-Aguirre C, Lang MM, Silcox MT. Diet drove brain and dental morphological coevolution in strepsirrhine primates. PLOS ONE. 2022 Jun 6;17(6):e0269041.

121.   Marcé-Nogué J, Püschel TA, Kaiser TM. A biomechanical approach to understand the ecomorphological relationship between primate mandibles and diet. Sci Rep. 2017 Aug 21;7(1):8364.

122.   Berthaume MA, Lazzari V, Guy F. The landscape of tooth shape: Over 20 years of dental topography in primates. Evol Anthropol. 2020;29(5):245–62.

123.   Sallee MD, Pickett MA, Feldman JL. Apical PAR complex proteins protect against programmed epithelial assaults to create a continuous and functional intestinal lumen. eLife. 10:e64437.

124.   Qi M, Tan B, Wang J, Li J, Liao S, Yan J, et al. Small intestinal transcriptome analysis revealed changes of genes involved in nutrition metabolism and immune responses in growth retardation piglets. J Anim Sci. 2019 Sep;97(9):3795–808.

125.   Stojanović O, Altirriba J, Rigo D, Spiljar M, Evrard E, Roska B, et al. Dietary excess regulates absorption and surface of gut epithelium through intestinal PPARα. Nat Commun. 2021 Dec 2;12(1):7031.

126.   Hossain MS, Begum SMKN, Rahman MM, Mazumder RN, Parvez M, Gazi MA, et al. Alterations in the histological features of the intestinal mucosa in malnourished adults of Bangladesh. Sci Rep. 2021 Jan 27;11(1):2355.

127.   Farràs M, Chandwe K, Mayneris-Perxachs J, Amadi B, Louis-Auguste J, Besa E, et al. Characterizing the metabolic phenotype of intestinal villus blunting in Zambian children with severe acute malnutrition and persistent diarrhea. PloS One. 2018;13(3):e0192092.

128.   Bein A, Fadel CW, Swenor B, Cao W, Powers RK, Camacho DM, et al. Nutritional deficiency in an intestine-on-a-chip recapitulates injury hallmarks associated with environmental enteric dysfunction. Nat Biomed Eng. 2022 Nov;6(11):1236–47.

129.   Andersen MK, Skotte L, Jørsboe E, Polito R, Stæger FF, Aldiss P, et al. Loss of Sucrase-Isomaltase Function Increases Acetate Levels and Improves Metabolic Health in Greenlandic Cohorts. Gastroenterology. 2022 Apr 1;162(4):1171-1182.e3.

130.   Senftleber NK, Ramne S, Moltke I, Jørgensen ME, Albrechtsen A, Hansen T, et al. Genetic Loss of Sucrase-Isomaltase Function: Mechanisms, Implications, and Future Perspectives. Appl Clin Genet. 2023 Mar 23;16:31–9.

131.   Senftleber NK, Skøtt Pedersen K, Schnoor Jørgensen C, Pedersen H, Bjerg Christensen MM, Kabel Madsen E, et al. The effect of sucrase-isomaltase deficiency on metabolism, food intake and preferences: protocol for a dietary intervention study. Int J Circumpolar Health. 82(1):2178067.

132.   Pirahanchi Y, Sharma S. Biochemistry, Lipase. In: StatPearls [Internet]. Treasure Island (FL): StatPearls Publishing; 2024 [cited 2024 Feb 9]. Available from: http://www.ncbi.nlm.nih.gov/books/NBK537346/

133.   Zhu G, Fang Q, Zhu F, Huang D, Yang C. Structure and Function of Pancreatic Lipase-Related Protein 2 and Its Relationship With Pathological States. Front Genet [Internet]. 2021 [cited 2024 Feb 9];12. Available from: https://www.frontiersin.org/journals/genetics/articles/10.3389/fgene.2021.693538

134.   Fu J, Zong X, Jin M, Min J, Wang F, Wang Y. Mechanisms and regulation of defensins in host defense. Signal Transduct Target Ther. 2023 Aug 14;8(1):1–30.

135.   Islam S, Akhand MRN, Hasan M. Evolutionary trend of bovine β-defensin proteins toward functionality prediction: A domain-based bioinformatics study. Heliyon. 2023 Mar 4;9(3):e14158.

136.   Zhang ZY, Zhang HM, Li DS, Xiong TY, Fang SG. Characterization of the β-defensin genes in giant panda. Sci Rep. 2018 Aug 17;8(1):12308.

137.   Robinson K, Ma X, Liu Y, Qiao S, Hou Y, Zhang G. Dietary modulation of endogenous host defense peptide synthesis as an alternative approach to in-feed antibiotics. Anim Nutr. 2018 Jun;4(2):160–9.

138.   Chen J, Zhai Z, Long H, Yang G, Deng B, Deng J. Inducible expression of defensins and cathelicidins by nutrients and associated regulatory mechanisms. Peptides. 2020 Jan 1;123:170177.

139.   Castellanos FX, Moreno-Santillán D, Hughes GM, Paulat NS, Sipperly N, Brown AM, et al. The evolution of antimicrobial peptides in Chiroptera. Front Immunol [Internet]. 2023 [cited 2024 Jan 18];14. Available from: https://www.frontiersin.org/articles/10.3389/fimmu.2023.1250229

140.   Nunn CL, Gittleman JL, Antonovics J. Promiscuity and the primate immune system. Science. 2000 Nov 10;290(5494):1168–70.

141.   Anderson MJ, Hessel JK, Dixson AF. Primate mating systems and the evolution of immune response. J Reprod Immunol. 2004 Feb 1;61(1):31–8.

142.   Orecchioni M, Matsunami H, Ley K. Olfactory receptors in macrophages and inflammation. Front Immunol. 2022 Oct 13;13:1029244.

143.   Manteniotis S, Wojcik S, Göthert JR, Dürig J, Dührsen U, Gisselmann G, et al. Deorphanization and characterization of the ectopically expressed olfactory receptor OR51B5 in myelogenous leukemia cells. Cell Death Discov. 2016 May 9;2(1):1–9.

144.   Guardia GDA, Naressi RG, Buzzato VC, da Costa JB, Zalcberg I, Ramires J, et al. Acute Myeloid Leukemia Expresses a Specific Group of Olfactory Receptors. Cancers. 2023 Jun 6;15(12):3073.

145.   Manteniotis S, Wojcik S, Brauhoff P, Möllmann M, Petersen L, Göthert JR, et al. Functional characterization of the ectopically expressed olfactory receptor 2AT4 in human myelogenous leukemia. Cell Death Discov. 2016 Jan 25;2(1):1–10.

146.   Florsheim EB, Bachtel ND, Cullen JL, Lima BGC, Godazgar M, Carvalho F, et al. Immune sensing of food allergens promotes avoidance behaviour. Nature. 2023 Aug;620(7974):643–50.

147.   Pollard KM, Christy JM, Cauvi DM, Kono DH. Environmental Xenobiotic Exposure and Autoimmunity. Curr Opin Toxicol. 2018 Aug;10:15–22.

148.   Talbot S, Foster SL, Woolf CJ. Neuroimmunity: Physiology and Pathology. Annu Rev Immunol. 2016 May 20;34:421–47.

149.   Colchero F, Aburto JM, Archie EA, Boesch C, Breuer T, Campos FA, et al. The long lives of primates and the 'invariant rate of ageing' hypothesis. Nat Commun. 2021 Jun 16;12(1):3666.

150.   Byrne H, Webster TH, Brosnan SF, Izar P, Lynch JW. Signatures of adaptive evolution in platyrrhine primate genomes. Proc Natl Acad Sci U S A. 2022 Aug 30;119(35):e2116681119.

151.   Finch CE. Evolution in health and medicine Sackler colloquium: Evolution of the human lifespan and diseases of aging: roles of infection, inflammation, and nutrition. Proc Natl Acad Sci U S A. 2010 Jan 26;107 Suppl 1(Suppl 1):1718–24.

152.   Billerot E, Nguyen TH, Sedira N, Espinoza S, Vende B, Heron E, et al. Ocular motor nerve palsy in patients with diabetes: High-resolution MR imaging of nerve enhancement. J Fr Ophtalmol. 2023 Sep 1;46(7):726–36.

153.   Choi DD, Han K, Oh SY, Park KA. Association between metabolic syndrome and incidence of ocular motor nerve palsy. Sci Rep. 2021 Nov 29;11(1):23033.

154.   Kang DW, Ha SY, Sung JJ, Nam H. Cerebral small vessel disease as imaging biomarker predicting ocular cranial nerve palsy of presumed ischemic origin at admission. Sci Rep. 2022 Jul 18;12(1):12251.

155.   Suda T, Matsushita E, Minamide H. Oculomotor nerve palsy owing to cerebral aneurysm. QJM Int J Med. 2022 Jun 1;115(6):395–6.

156.   Boraschi D. What Is IL-1 for? The Functions of Interleukin-1 Across Evolution. Front Immunol. 2022 Apr 6;13:872155.

157.   Kaneko N, Kurata M, Yamamoto T, Morikawa S, Masumoto J. The role of interleukin-1 in general pathology. Inflamm Regen. 2019 Jun 6;39(1):12.

158.   Brown-Borg HM. Growth Hormone, Not IGF-1 Is the Key Longevity Regulator in Mammals. J Gerontol Ser A. 2022 Sep 1;77(9):1719–23.

159.   Scossa F, Fernie AR. Ancestral sequence reconstruction - An underused approach to understand the evolution of gene function in plants? Comput Struct Biotechnol J. 2021;19:1579–94.

160.   Uyeda JC, Bone N, McHugh S, Rolland J, Pennell MW. How should functional relationships be evaluated using phylogenetic comparative methods? A case study using metabolic rate and body temperature. Evolution. 2021;75(5):1097–105.

161.   Sander PM, Griebeler EM, Klein N, Juarbe JV, Wintrich T, Revell LJ, et al. Early giant reveals faster evolution of large body size in ichthyosaurs than in cetaceans. Science. 2021 Dec 24;374(6575):eabf5787.

162.   Benson RBJ, Hunt G, Carrano MT, Campione N. Cope's rule and the adaptive landscape of dinosaur body size evolution. Palaeontology. 2018;61(1):13–48.

163.   Ksepka DT, Balanoff AM, Smith NA, Bever GS, Bhullar BAS, Bourdon E, et al. Tempo and Pattern of Avian Brain Size Evolution. Curr Biol. 2020 Jun 8;30(11):2026-2036.e3.

164.   Fukushima K, Pollock DD. Detecting macroevolutionary genotype–phenotype associations using error-corrected rates of protein convergence. Nat Ecol Evol. 2023 Jan;7(1):155–70.

165.   Altschul SF, Gish W, Miller W, Myers EW, Lipman DJ. Basic local alignment search tool. J Mol Biol. 1990 Oct 5;215(3):403–10.

166.   Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. Nucleic Acids Res. 2004;32(5):1792–7.

167.   Capella-Gutiérrez S, Silla-Martínez JM, Gabaldón T. trimAl: a tool for automated alignment trimming in large-scale phylogenetic analyses. Bioinformatics. 2009 Aug 1;25(15):1972–3.

168.   Ranwez V, Douzery EJP, Cambon C, Chantret N, Delsuc F. MACSE v2: Toolkit for the Alignment of Coding Sequences Accounting for Frameshifts and Stop Codons. Mol Biol Evol. 2018 Oct 1;35(10):2582–4.

169.   Ranwez V, Harispe S, Delsuc F, Douzery EJP. MACSE: Multiple Alignment of Coding SEquences Accounting for Frameshifts and Stop Codons. PLOS ONE. 2011 Sep 16;6(9):e22594.

170.   Criscuolo A, Gribaldo S. BMGE (Block Mapping and Gathering with Entropy): a new software for selection of phylogenetic informative regions from multiple sequence alignments. BMC Evol Biol. 2010 Jul 13;10(1):210.

171.   Revell LJ. phytools: an R package for phylogenetic comparative biology (and other things). Methods Ecol Evol. 2012;3(2):217–23.

172.   Revell LJ, Chamberlain SA. Rphylip: an R interface for PHYLIP. Methods Ecol Evol. 2014;5(9):976–81.

173.   Clavel J, Escarguel G, Merceron G. mvmorph: an r package for fitting multivariate evolutionary models to morphometric data. Methods Ecol Evol. 2015;6(11):1311–9.

174.   Stefen C, Wagner F, Asztalos M, Giere P, Grobe P, Hiller M, et al. Phenotyping in the era of genomics: MaTrics—a digital character matrix to document mammalian phenotypic traits. Mamm Biol. 2022 Feb 1;102(1):235–49.

175.   Soria CD, Pacifici M, Di Marco M, Stephen SM, Rondinini C. COMBINE: a coalesced mammal database of intrinsic and extrinsic traits. Ecology. 2021;102(6):e03344.

176.   Devine J, Kurki HK, Epp JR, Gonzalez PN, Claes P, Hallgrímsson B. Classifying high-dimensional phenotypes with ensemble learning. BioRxiv Prepr Serv Biol. 2023 May 29;2023.05.29.542750.

177.   Alghamdi SM, Hoehndorf R. Improving the classification of cardinality phenotypes using collections. J Biomed Semant. 2023 Aug 7;14(1):9.

178.   Kwak HS, Im HG, Shim EB. A model for allometric scaling of mammalian metabolism with ambient heat loss. Integr Med Res. 2016 Mar 1;5(1):30–6.

179.   Jones JH. Primates and the Evolution of Long-Slow Life Histories. Curr Biol CB. 2011 Sep 27;21(18):R708–17.

180.   Freckleton RP, Harvey PH, Pagel M. Phylogenetic analysis and comparative data: a test and review of evidence. Am Nat. 2002 Dec;160(6):712–26.

181.   Junier T, Zdobnov EM. The Newick utilities: high-throughput phylogenetic tree processing in the Unix shell. Bioinformatics. 2010 Jul 1;26(13):1669–70.

182.   Yang Z. PAML: a program package for phylogenetic analysis by maximum likelihood. Comput Appl Biosci CABIOS. 1997 Oct;13(5):555–6.

183.   Collyer ML, Adams DC. RRPP: An r package for fitting linear models to high-dimensional data using residual randomization. Methods Ecol Evol. 2018;9(7):1772–9.

184.   Ives AR, Garland T Jr. Phylogenetic Logistic Regression for Binary Dependent Variables. Syst Biol. 2010 Jan 1;59(1):9–26.

185.   Hussain T, Asghar S. EVALUATION OF SIMILARITY MEASURES FOR CATEGORICAL DATA. The Nucleus. 2013 Nov 27;50(4):387–94.

186.   Liao Y, Wang J, Jaehnig EJ, Shi Z, Zhang B. WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. Nucleic Acids Res. 2019 Jul 2;47(W1):W199–205.

187.   Herrendörfer G. JOHNSON, N. L., KOTZ, S.; KEMP, A. W.: Univariate discrete distributions. Second Edition — John Wiley & Sons 1992, 565 pp., £58.00. Biom J. 1994;36(2):164–164.

188.   Pond SLK, Frost SDW, Muse SV. HyPhy: hypothesis testing using phylogenies. Bioinformatics. 2005 Mar 1;21(5):676–9.

189.   Murrell B, Moola S, Mabona A, Weighill T, Sheward D, Kosakovsky Pond SL, et al. FUBAR: a fast, unconstrained bayesian approximation for inferring selection. Mol Biol Evol. 2013 May;30(5):1196–205.

190.   Tong C, Avilés L, Rayor LS, Mikheyev AS, Linksvayer TA. Genomic signatures of recent convergent transitions to social life in spiders. Nat Commun. 2022 Nov 22;13(1):6967.

191.   Shank SD, Weaver S, Kosakovsky Pond SL. phylotree.js - a JavaScript library for application development and interactive data visualization in phylogenetics. BMC Bioinformatics. 2018 Jul 25;19(1):276.

192.   Zhou W, Chen T, Chong Z, Rohrdanz MA, Melott JM, Wakefield C, et al. TransVar: a multilevel variant annotator for precision genomics. Nat Methods. 2015 Nov;12(11):1002–3.

193.   Christmas MJ, Kaplow IM, Genereux DP, Dong MX, Hughes GM, Li X, et al. Evolutionary constraint and innovation across hundreds of placental mammals. Science. 2023 Apr 28;380(6643):eabn3943.

194.   Kirilenko BM, Munegowda C, Osipova E, Jebb D, Sharma V, Blumer M, et al. Integrating gene annotation with orthology inference at scale. Science. 2023 Apr 28;380(6643):eabn3107.

195.   Nature [Internet]. 2020 [cited 2023 Sep 28]. gnomAD. Available from: https://www.nature.com/collections/afbgiddede

196.   Jones KE, Bielby J, Cardillo M, Fritz SA, O'Dell J, Orme CDL, et al. PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. Ecology. 2009;90(9):2648–2648.

197.   Leeuw CA de, Mooij JM, Heskes T, Posthuma D. MAGMA: Generalized Gene-Set Analysis of GWAS Data. PLOS Comput Biol. 2015 Apr 17;11(4):e1004219.

198.   Finucane HK, Bulik-Sullivan B, Gusev A, Trynka G, Reshef Y, Loh PR, et al. Partitioning heritability by functional annotation using genome-wide association summary statistics. Nat Genet. 2015 Nov;47(11):1228–35.

199.   Lee BT, Barber GP, Benet-Pagès A, Casper J, Clawson H, Diekhans M, et al. The UCSC Genome Browser database: 2022 update. Nucleic Acids Res. 2022 Jan 7;50(D1):D1115–22.

200.   Hamosh A, Scott AF, Amberger JS, Bocchini CA, McKusick VA. Online Mendelian Inheritance in Man (OMIM), a knowledgebase of human genes and genetic disorders. Nucleic Acids Res. 2005 Jan 1;33(suppl_1):D514–7.