1 **Using gas exchange measurements to monitor growth, energy expenditure, and**
2 **body composition of black soldier fly larvae (*Hermetia illucens*) on diets varying**
3 **in protein-to-carbohydrate ratio**
4

M.L. Schøn[1,2], K. Jensen[1,*], M. Schow-Madsen[1], M. Gold[2], A. Mathys[2], T.M. Schou[3], I.E. Berggreen[1],
J.V. Nørgaard[1], and J. Overgaard[4,*]

[1] Department of Animal and Veterinary Sciences, Aarhus University, Blichers Allé 20, 8830 Tjele, Denmark
[2] Sustainable Food Processing Laboratory, Department for Health Sciences and Technology, ETH Zurich,
Schmelzbergstrasse 9, 8092 Zurich, Switzerland
[3] ENORM Biofactory A/S, Hedelundvej 15, 8762 Flemming, Denmark
[4] Zoophysiology, Department of Biology, Aarhus University, C.F. Møllers Allé 3, 8000 Aarhus C, Denmark
*Corresponding authors: Johannes.Overgaard@bio.au.dk and kj@anivet.au.dk

13
14

Running header: Metabolism and growth of BSFL on diets varying in protein-to-carbohydrate ratio

Abstract

Optimizing black soldier fly larvae (BSFL) production requires a better understanding of how
diet composition shapes growth, nutrient allocation, and metabolic efficiency. This study in-
vestigated whether respiratory measurements could serve as predictive, non-invasive tools for
BSFL performance on diets differing in protein-to-carbohydrate (P:C) ratio. Larvae were
reared for seven days on four experimental diets, with growth, survival, gas exchange (O_2
consumption, CO_2 production, respiratory exchange ratio (RER)), and body composition
(crude protein and lipid contents) measured across seven replicates per treatment. O_2 con-
sumption correlated strongly with final larval biomass, while RER was closely associated
with protein and lipid deposition. Larvae on low-protein, high-carbohydrate diets accumulated
proportionally more lipid, whereas protein-rich diets increased crude protein content but re-
quired higher energetic investment. This demonstrates that lipid-rich larvae grew more effi-
cient in energetic terms per unit accumulated biomass than protein-rich larvae. Across diets,
gas exchange variables reliably reflected growth dynamics, metabolic activity, and nutrient
assimilation. Integrating respirometry into rearing systems could enable real-time monitoring
of insect biomass yield and nutrient composition.

Keywords: carbon dioxide, dietary macronutrient composition, energetic cost, growth monitoring,
metabolism, oxygen, respiratory exchange ratio

ORCID:
Maja Lermark Schøn, https://orcid.org/0009-0007-6251-9034
Kim Jensen, https://orcid.org/0000-0003-0261-3831
Mattias Schow-Madsen https://orcid.org/0009-0004-1289-3579
Moritz Gold https://orcid.org/0000-0001-7285-6164
Alexander Mathys https://orcid.org/0000-0003-1633-848X
Ida Elisabeth Berggreen, https://orcid.org/0000-0002-8546-6517
Toke Munk Schou, https://orcid.org/0000-0001-7545-7572
Jan Værum Nørgaard, https://orcid.org/0000-0002-0580-0184
Johannes Overgaard, https://orcid.org/0000-0002-2851-4551

Introduction

The world faces pressing environmental challenges that impact global food production, including climate change, soil degradation, habitat loss, and the depletion of natural resources (Foley *et al*., 2011). More sustainable food systems are essential to meet the increasing food and feed demand driven by population growth while simultaneously mitigating the impacts associated with greenhouse gas emissions, land-use, water scarcity, and food wastage (FAO, 2009; Foley *et al*., 2011). One component of this more sustainable food and feed production is the use of black soldier fly larvae (BSFL; *Hermetia illucens* (L.)) for converting low quality organic material into nutrient-rich biomass (Broeckx *et al*., 2021; Gold *et al*., 2020). This biomass can supplement traditional unsustainable protein sources (e.g. soybean and fishmeal) in livestock feeds (van der Heide *et al*., 2021) and pet foods (Bosch & Swanson, 2021), and as an added benefit, the remaining frass after harvest of larvae represents a high-quality fertilizer (Fuhrmann *et al*., 2022; Lomonaco *et al*., 2024).

The ability of BSFL to feed on a wide range of organic materials is a major reason why this species has become increasingly important commercially (Purkayastha & Sarkar, 2022). Nevertheless, BSFL growth rates, energy conversion efficiencies, and body compositions can vary considerably with diet quality and macronutrient balance (Barragan-Fonseca *et al*., 2021; Eggink *et al*., 2023), and with other conditions such as moisture content (Bekker *et al*., 2021) or temperature (Schøn *et al*., 2024). Diet composition, in particular the protein-to-carbohydrate (P:C) ratio and not just the total diet mass, has been shown to strongly affect larval growth and body composition. Studies have demonstrated that nutrient-deficient diets that deprive larvae of adequate protein or carbohydrate supply, as well as diets with unbalanced P:C ratios, reduce larval growth and alter nutrient deposition patterns (Barragan‐Fonseca *et al*., 2019; Berggreen *et al*., 2025; Eggink *et al*., 2023). As a result, differences in diet can lead to substantial variation in BSFL body composition, with larvae ranging from 32–58% crude protein and 15–38% crude lipids on a dry matter (DM) basis (see review by Gold *et al.*, 2018). Because variations in macronutrient composition (encompassing both the levels and specific forms of protein, fat, and carbohydrate) affect larval development, size, and composition, it is important to understand in more detail how these dietary factors influence metabolism and energy conversion during growth. Similarly, there is a growing interest in developing methods to monitor the progression of growth in actively rearing larvae.

Measurements of gas exchange (O_2 consumption and CO_2 production) as well as the respiratory exchange ratio (RER, VCO_2/VO_2) are powerful indicators of metabolism, energy expenditure, and nutrient assimilation under different dietary conditions (Parodi et al., 2020;

Schøn *et al*., 2024; Talal *et al*., 2021). BSFL utilizes aerobic energy both for general mainte-

nance of basal physiological processes, but also in association with the direct cost of growth

and nutrient assimilation (Ferrannini, 1988; Talal *et al*., 2021) and studies have previously

found a close association between O_2 consumption or CO_2 production and total larval mass

gain (Eriksen, 2024; Schøn *et al*., 2024). However, this association may be affected by the

rearing conditions, including the dietary conditions, as the energetics of growth is expected to

differ depending on the ratio of lipid and protein assimilation and because different nutritional

quality may affect the relative magnitude of maintenance metabolism versus growth metabo-

lism in ectotherms (Goodrich *et al*., 2024; Laganaro *et al*., 2021). This may be further compli-

cated by the fact that in the BSFL rearing system the diet and digestive tract microbiome also

contribute to overall gas exchange rates (Bekker *et al.*, 2021; Parodi *et al.*, 2020). Neverthe-

less, continuous monitoring of gas exchange rates holds the potential to provide important

information on growth and body composition in production systems. In this context, RER

may be particularly relevant to examine, as this metric reflects the combined input from a

range of metabolic processes related to both anabolism and catabolism. RER reflects the ca-

tabolism of macronutrients where oxidation of lipid, protein and carbohydrate results in RER

of 0.7, 0.83 or 1.0 (Schmidt-Nielsen, 1997). In addition, RER will increase as a consequence

of lipogenesis from carbohydrates or amino acids (Ferrannini, 1988; Talal *et al*., 2021), and

RER measurements therefore hold potential to indirectly monitor which macronutrients are

metabolized and assimilated.

The main objectives of this study were: (*i*) to assess how altering diet macronutrient compo-

sition affects the growth, metabolism, and body composition of BSFL, and (*ii*) to determine

whether gas exchange measurements during rearing are predictive of BSFL production output

in terms of growth rate, quantity, and quality. To address these objectives, larval growth, gas

exchange (O_2 and CO_2) and body composition (crude protein and lipid content) were meas-

ured on BSFL reared on diets with varying P:C ratio.

Materials and methods

Black soldier flies

BSFL of 5-6 days upon hatching (4[th] or 5[th] instar) (Gligorescu *et al*., 2019)　were supplied

from a commercial BSFL facility (ENORM Biofactory A/S, Flemming, Denmark). After col-

lection at the factory, the larvae were transported (1 h) to Aarhus University, Denmark, where

all experiments were performed. For each experimental replicate, 600 counted larvae with

average individual mass of 5.0 ± 0.9 mg were placed in 1 L respirometer chambers (15 cm

diameter, 6 cm height) each containing approximately 340 g of fresh diet. This resulted in a

density of ~1.76 larvae per g wet diet or ~0.133 g dry diet per larva.

Experimental protocol

The study investigated how BSFL metabolism is affected by dietary macronutrient composi-

tion by measuring gas exchange, growth, and body composition of BSFL during 7 days of

rearing on four experimental diets. Experiments were performed over four consecutive weeks

using a blocked design where one or two replicates of each dietary treatment was included

every week. This resulted in 7 replicates of each dietary treatment. The experimental system

allowed for hourly recordings of O_2 consumption rate ($\dot{V}O_2$) and CO_2 production rate

($\dot{V}CO_2$). Larval growth was estimated daily by sub-sampling and weighing 15 randomly

selected larvae from each replicate. At termination of the 7-day experiment, larvae from each

replicate were separated from the frass using wet sieving, and larvae were subsequently

quickly dried with paper towels. Larvae were then weighed and photographed to count the

number of survivors using CountThings (Dynamic Ventures Inc., California, US). All experi-

ments were performed in a climate cabinet at 27 °C (Termaks KB8182, Bergen, Norway) and

constant darkness. Larvae sampled at the termination of the experiment were stored at -20 °C

before analyses of DM, crude lipid, and crude protein contents. Samples of the initial seed

larvae were also collected and stored for analysis.

Experimental diets

The ingredients and nutritional composition of the four experimental diets are shown in Table

1. All four diets contained commercial chicken feed (KylleKræs 1, Danish Agro, Galten,

Denmark). The reference (REF) diet consisted of 100% chicken feed and was used as a high-

protein diet control to monitor if the system supported growth and metabolism to a similar

degree as previous studies (e.g., Berggreen *et al*., 2025; Eggink *et al*., 2023; Nguyen *et al*.,

2015). The semi-artificial diets where mixed to represent different ratios of protein to non-

fiber carbohydrate (NFC) (P:C). Specifically, P:C 1:3, P:C 1:5, and P:C 1:9 diets contained

35% chicken feed to ensure supplementation of all required micronutrients, and the remaining

65% was included to create as semi-synthetic diets of desired macronutrient contents follow-

ing Berggreen *et al*. (2025) (Table 1). These treatment diets contained different proportions of

a protein mix (casein and peptone, 1:1) and a carbohydrate mix (sucrose and dextrin, 1:1) to

achieve desired P:C ratios. A lipid mix (rapeseed oil, sunflower oil, and lard, 1:1:1) and a mix

of additives including vitamins (Vanderzant modification vitamin mixture), minerals (Wes-

sons salt mixture), fatty acids (linoleic acid and oleic acid, 4:1), cholesterol, and nipagin

(1:6:1:1:1), as well as cellulose and agar were added at similar concentrations in the three

mixed diets (Table 1). Cellulose and agar were added as structural components, and nipagin

was added as a preservative. All experimental diets were prepared by mixing 320 g of agar

solution (agar dissolved in boiling water and then cooled to 40 °C) with 80 g of pre-mixed dry

diet ingredients. The diets were then left to solidify during which water evaporated, leaving

341.3 ± 6.3 g wet diet (~77% water) in the chambers at the onset of the experiment.

Table 1

Gas exchange measurements

The open-flow respirometry setup and protocol for gas exchange measurements were similar

to that described with greater detail in Schøn *et al.* (2024). Briefly, 8 custom built, airtight

respiration chambers (15 cm diameter, 6 cm height, 1 L volume) were placed in the climate

cabinet. Each chamber was provided with a controlled and constant flow of room air (~600

mL min^{-1}) that continuously provided oxygen and removed water vapor and CO_2. The air

flow for the 8 sequential chambers was regulated and recorded using a Flowbar-8 Mass Flow

Meter System (Sable Systems).

To measure $\dot{V}O_2$ and $\dot{V}CO_2$, air was sequentially sub-sampled at a rate of ~200 mL

min^{-1} using a SS-4 Sub-Sampler Pump (Sable Systems). Sequential shifting between cham-

bers was performed using an eight-channel RM-8 Respirometry Flow Multiplexer (Sable Sys-

tems). The sampled air passed through a calcium chloride column for removal of water vapor,

and syringe filters (Acrodisc syringe filter, 25 mm, 0.2 µm Supor membrane, Pall Corpora-

tion) filtered the air before O_2 and CO_2 concentrations were analysed using an O_2 analyser

(FC-2 Differential Oxygen Analyser (Oxzilla), Sable Systems) and a CO_2 analyser (LI-850

Gas Analyzer, LI-COR Biosciences), respectively. The multiplexer shifted between each of

the eight respiration chambers (7 experimental chambers and one empty control chamber)

every 7.5 minutes, providing hourly measurements from each chamber over the entire 7-day

period. Unstable recordings of O_2 or CO_2 as well as outliers were removed and replaced by

interpolation from preceding and subsequent samples of high quality using the data quality

assessment protocol described in Schøn *et al.* (2024).

$\dot{V}O_2$ and $\dot{V}CO_2$ were calculated hourly from the product of the airflow rate and the

difference in gas concentration between the experimental chambers and the empty control

chamber. The total volumes of O_2 consumed (VO_2) and CO_2 produced (VCO_2) were cal-

culated by summation of hourly measurements of VO_2 and VCO_2 over the entire 7-day pe-

riod. The respiratory exchange ratio (RER) was calculated hourly from $V̇CO_2/V̇O_2$ or

cumulatively from total VCO_2/ total VO_2.

Growth analysis

Respiration chambers with growing larvae were daily removed briefly from the temperature

cabinet to track changes in total mass (frass and larvae) between days. On average the repli-

cates lost 27.3 ± 3.4 g day^{-1}, mainly due to removal of water vapor and CO_2. During this pro-

cedure, 15 randomly selected larvae were quickly collected and weighed to assess average

larval mass within each replicate. Three of the 15 larvae were sampled and frozen, and the

other 12 larvae were returned. Thus, during the experiment 18 larvae were removed from each

replicate. At the termination of the experiment all remaining larvae were counted and weighed

to enable calculation of survival and total mass gain.

The total daily larval mass was estimated by multiplying the average daily larval mass with

the estimated number of live larvae on each day. The estimate of live larvae accounted for the

daily removal of 3 larvae and assumed that mortality was evenly distributed across the 7-day

rearing period. Specific growth rates (SGR) were calculated as the difference in the natural

logarithm of average body mass between two time points divided by the elapsed time:

$$SGR = \frac{\ln Final\ wet\ mass - \ln Initial\ wet\ mass}{Time}$$

Water, lipid, and protein content

At the termination of the experiment, all harvested larvae from each replicate were oven-dried

at 60 °C for three to four days until they had reached a stable mass. Water content was calcu-

lated from the difference between wet mass and DM, and sub-samples of dried larvae were

taken to estimate lipid and protein contents. To obtain crude lipid content, samples of $1.76 \pm$

0.01 g dried larvae were weighed before they were washed 5 times in 25 mL petroleum ether

to remove the lipid fraction (Jensen *et al*., 2010). Each round of washing lasted 48 h and after

the final round the defatted samples were again oven-dried at 60°C until stable mass to allow

for estimation of the crude lipid fraction (calculated as the difference between initial DM and

lean DM). Nitrogen analysis of defatted, finely ground sub-samples (4.2–7.2 mg per sample)

was performed using a combustion analyser (Na 2000, Italy). The proportional crude protein

content was then estimated by multiplying the measured nitrogen content by a factor of 4.76,

a standardized conversion factor for the nitrogen-to-protein ratio in BSFL (Janssen *et al*.,

2017). Total protein was estimated from lean DM and expressed as a fraction of the total DM

(before lipid removal).

Analysis of gross cost of growth and assimilation

To examine the relationship between energy consumption and energy assimilation, mass gain

was converted into energy equivalents based on mass gain of lipid and protein. For these cal-

culations an assumption of 37 kJ per g lipid and 17 kJ per g protein as energy conversion fac-

tors was used (FAO, 2002), respectively. Further, to assess the energy used for aerobic me-

tabolism VO_2 was converted to energy expenditure using a standard caloric conversion factor

of 20.5 kJ per L O_2 (Schmidt-Nielsen, 1997).

Data analysis

Separate one-way ANOVA were performed to analyse BSFL growth, gas exchange, and body

composition between treatment groups. Assumptions of normality and homogeneity of vari-

ances were assessed using Shapiro-Wilk tests, visual inspection of Q-Q plots, and Levene's

test. Tukey's Honest Significant Difference (HSD) test was used for post hoc comparisons,

with statistical significance set at $p < 0.05$.

Linear regressions were used to evaluate the relationships between final larval DM and

VO_2, as well as between assimilated energy from lipid and protein and the estimated aerobic

energy expenditure. Pearson correlation coefficients were calculated to assess the strength and

direction of these relationships. In selected analyses, linear models were constrained to pass

through the origin (0,0) to test the assumption of a direct proportional relationship between

the dependent and independent variables.

Data processing and visualization were performed in RStudio version 2024.04.2 with R ver-

sion 4.3.2 (R Core Team, 2023; RStudio Team, 2023).

Results

Effects of diet on growth and body composition of BSFL

Larvae survived well and developed successfully on all diets, with average survival rates

ranging from 89–97% and with no significant difference among diets (Table 2). While all di-

ets supported substantial larval growth, the final mass of larvae was significantly different

across diets after the 7-day growth period (Figure 1A). Total larval yield was highest with the

REF diet, followed by the mixed diets with highest protein fraction, P:C 1:3, P:C 1:5 and P:C

1:9 diets, respectively (Figure 1A, Table 2). The difference in growth between diets was also

evident from our analysis of specific growth rates (SGR) (Figure 1B, Table 2). From the SGR

(Figure 1B), it is possible to see the change in estimated total biomass between two consecu-

tive days, where the biggest increase for P:C 1:3, P:C 1:5, and P:C 1:9 was found to be from

day 3 to 4, and thereafter a more moderate increase (and even slight decrease for the REF

diet) in SGR per day.

The larvae reared on different diets had significantly different body compositions at harvest

(Table 2). Larvae reared on the P:C 1:9 diet had significantly lower water content compared to

those from the other three diets. While the proportions of crude protein and crude lipid in the

DM did not differ significantly between REF and P:C 1:3, they were significantly higher than

found in diets with lower protein content (P:C 1:5 and P:C 1:9) (Table 2). As the P:C ratio

increased in the experimental diets, larval crude protein levels rose and lipid levels dropped,

resulting in larvae from the low protein diets exhibiting significantly elevated proportional

lipid fraction and reduced protein fraction (Table 2).

Effects of diet on gas exchange and metabolism

The dynamic changes in $\dot{V}O_2$ and $\dot{V}CO_2$ over the 7-day growth period followed a

broadly similar pattern (see Figure 2A and Figure S1). Here, gas exchange data is presented

with a focus on $\dot{V}O_2$ and RER, showing that gas exchange rates generally increased over the

7-day growth period but with different temporal progressions and distinct patterns depending

on dietary treatment (Figure 2). For the commercial high-protein diet, REF, the $\dot{V}O_2$ in-

creased slowly during the first two days, after which the rate accelerated. $\dot{V}O_2$ spiked at

three distinct time points: on day 1 (0.7 mL O_2 min^{-1}), day 3.5 (2.2 mL O_2 min^{-1}), and day

5.5 (4.6 mL O_2 min^{-1}), before $\dot{V}O_2$ declined considerably during the last 1.5 days of the

experiment. A somewhat similar pattern was found with P:C 1:3 and P:C 1:5, but here the

spikes occurred later and were more moderate. P:C 1:9 displayed an even more delayed and

reduced response with a small spike around day 1.5-2, another by day 5.5, with indications of

a third increase in $\dot{V}O_2$ at the end of the experiment. Total $\dot{V}O_2$ for REF was significantly

higher than for all other diets, and the two mixed diets with highest protein content (P:C 1:3

and P:C 1:5) were associated with a significantly higher total VO_2 compared to P:C 1:9 (Fig-

ure 2A, Table 2).

RER values were >1 across diets when estimated from the total gas exchange of the cham-

bers over the 7-day growth period (Figure 2B, Table 2). RERs differ significantly among all

four diets with lowest RER for the REF and increasing RER for the P:C 1:3, P:C 1:5, and P:C

1:9 diet, respectively.

Respiratory variables as predictors of growth and body composition

Combining the data of growth and body composition with the data from gas exchange meas-

ures revealed several strong statistical correlations. First, there was a strong positive linear

correlation between VO_2 and final total larval DM when data was analysed across all diets

($R^2 = 0.93$, Figure 3A, black solid line). This correlation persists to be strong also when the

linear regression model is forced through origo (0,0) ($R^2 = 0.99$, Figure 3A, red dashed line).

As seen in Figure 3B, the correlation of energy consumption per energy assimilation results in

a significant, albeit weaker, correlation ($R^2 = 0.42$, Figure 3B, black solid line) suggesting that

across diets 0.91 kJ are expended for each kJ assimilated. Importantly, the gross cost of

growth (amount of metabolic energy used in relation to assimilation of 1 unit of energy as

biomass) varies significantly and considerably among diets (Figure 3B). The cost of growth

was significantly higher for the REF and lowest for the low protein diet (P:C 1:9). To examine

this further, the correlation between overall energy consumption and protein energy assimila-

tion was analysed. This analysis also found a strong positive correlation among all treatments

rendering an overall energetic cost of 1.88 kJ per kJ of protein assimilated ($R^2 = 0.87$, Figure

3C, black solid line). Importantly when analysed within each dietary treatment the cost of

protein assimilation varied significantly between diets (Figure 3C) such that protein energy

assimilation was cheapest on the high protein diets and more costly in the low protein diets.

Because RER is influenced by both catabolic and anabolic processes (i.e. lipogenesis) we

hypothesized that this metric could reveal information about the body composition of growing

larvae. Thus, lipid rich larvae should be characterized by higher proportional CO_2 contribution

from lipogenesis than more lean larvae. As seen in Figure 4, there was a strong correlation

between RER and lipid and protein contents when analysed across larvae from all treatments

($R^2 = 0.65$ and $R^2 = 0.64$, respectively, Figure 4, black lines). Interestingly, when this correla-

tion was investigated within each diet, the correlations were weaker and, in some cases, oppo-

site to the expected pattern (see Supplementary material for statistical analysis).

Discussion

The purpose of this study was to evaluate BSFL growth performance, metabolic activity, and

energy conversion efficiency (energetic cost of growth) using respirometry as a monitoring

tool in larvae reared on diets with varying protein-to-carbohydrate (P:C) ratios. Four experi-

mental diets were used; three semi-artificial diets with identical formulations except for vary-

ing P and C content to different ratios, and a reference diet of 100% chicken feed. Over a 7-

day period, measurements for larval biomass gain, oxygen consumption, carbon dioxide pro-

duction, and final crude protein and lipid content in the larvae was obtained. As expected,

diets with a higher P:C ratio accelerated larval growth and development by increasing protein

deposition. We could then link diet composition to metabolic activity and nutrient allocation

and evaluate whether gas exchange patterns can provide useful information on larval output in

terms of both larval biomass and nutritional body composition. We find that gas exchange

measurements are powerful indicators of both biomass yield and body composition in BSFL,

capturing not only the intensity but also the pulsatile nature of metabolic activity over time.

Effects of diet on growth and body composition of BSFL

The present study confirms previous findings that higher dietary protein levels support greater

larval biomass and protein content, whereas carbohydrate-rich diets increase lipid deposition

and produce smaller larvae (Barragan-Fonseca *et al*., 2021; Eggink *et al*., 2023; Gold *et al*.,

2020). In our experiments, the protein-rich diet with a P:C ratio of 1:3 (P:C 1:3) resulted in

69.0% higher larval DM and 59.1% higher crude protein content compared to larvae reared on

the carbohydrate-rich diet (P:C 1:9) (Table 1). P:C ratios around 1:2–1:3 appear optimal for

biomass yield (consistent with Eggink *et al*., 2023), while diets richer in carbohydrate pro-

moted lipid deposition, reaching 52.7% in this study and >44% in Eggink *et al*. (2023). These

results reflect how BSFL (and other insects) metabolically can channel surplus carbohydrates

in an unbalanced diet into lipid storage by lipogenesis (Schow-Madsen *et al*., 2025; Talal *et*

al., 2021) (Further addressed in the last section of the discussion).

The highest larval yield and protein accumulation in the present study was observed for the

reference diet (REF). While the REF diet (100% commercial chicken feed) is a valuable refer-

ence, it differs from the semi-synthetic diets (35% commercial chicken feed) in nutritional

aspects other than the P:C ratio (Table 1). The macronutrient content alone does not capture

the full nutritional complexity of a diet, which is also shaped by the types of protein, carbohy-

drates, and micronutrients it contains (Schow-Madsen *et al*., 2025). Therefore, the three semi-

synthetic diets, identical apart from their P:C content, provide the best basis for directly com-

paring P:C effects.

Oxygen consumption as predictor of growth and development in BSFL

Metabolic activity is known to increase with growth in BSFL, as in other insects (Eriksen,
2022, 2024). In our study, higher oxygen consumption was positively correlated with higher
larval end-biomass (Figure 3A), reflecting the energetic expenditure of increased nutrient as-
similation, tissue synthesis, and maintenance. In this experiment, total oxygen consumption
served as a reliable predictor of final biomass following larval growth in BSFL (Figure 3A).
Although some of the measured gas exchange may arise from microbial metabolism (Fuhr-
mann *et al.*, 2025), the data still reveals a direct relationship between respiration and BSFL
biomass yield.

Examining cumulative daily oxygen consumption in relation to total daily biomass provides
insight into the cost of growing and maintaining biomass. Across all diets, larvae on average
used less oxygen per g biomass as they grew, declining from ~0.07–0.08 L O_2 per g biomass
early in rearing to ~0.04–0.05 L O_2 per g biomass by day 7 (Figure S2). This cost reduction
likely reflects a shift in energy allocation from high-cost rapid structural growth in early de-
velopment to slower growth in later stages, where more metabolism is directed to mainte-
nance and energy storage, reducing the proportion of metabolic energy from respiration
needed per unit biomass as larvae approach peak size.

Monitoring temporal respiratory gas exchange showed that diet composition significantly
influenced oxygen consumption, corresponding closely with observed biomass growth curves
(Figures 1A and 2A). During the 7-day growth period, larvae grew progressively on all diets
(Figure 1A) and the signal of metabolic intensity ($V O_2$ and $V CO_2$; Figure 2A and S1)
showed increases and decreases according to growth. These pulsatile growth patterns likely
reflect instar transitions, with molting phases temporarily halting feeding and lowering meta-
bolic rates before the next growth phase (Gligorescu *et al.*, 2019). Temporal patterns differed
among diets: higher protein content accelerated development and shortened instar duration,
whereas carbohydrate-rich diets slowed progression and extended development periods (Fig-
ure 2A and S1). Although larval instar was not measured directly, the respiration profiles seem
to capture the cohort's ontogeny through larval stages, which demonstrate the value of respi-
rometry as a non-invasive tool for monitoring developmental dynamics in BSFL production
systems.

Continuous gas measurements identified the last growth phase, and thereby the maximum
biomass, for REF diet larvae by day 6, with respiration on the final day likely dominated by
non-feeding 7th instar individuals (Figure 1A and 2A) (also observed in Schow-Madsen *et al.*,
2025). For larvae reared on diet P:C 1:3 and P:C 1:5 biomass likely peaked between day 6 and
7, and for those on P:C 1:9 it would have been even later. This aligns with Eggink *et al.*

(2023), where the most carbohydrate-rich diet (P:C 1:9) prolonged development time by two

days compared to diets with P:C ratios of 1:3 or less.

Shifts in development time may influence end results, as BSFL body composition (e.g., pro-

portions of water, protein, and lipid) changes throughout development (Eggink & Dalsgaard,

2023). If larvae reared on different diets are at different developmental stages at harvest on

day 7, this can directly affect measured body composition as well as observed growth and

metabolic profiles. Harvesting larvae at the same developmental stage can help control for this

but may also introduce other limitations (see discussion on timing of harvest in Deruytter *et*

al. (2025). Notably, Eggink *et al*. (2023) found that even when carbohydrate-rich diets (P:C

1:5 and P:C 1:9) were harvested later to match developmental stage, they still produced larvae

with significantly different body compositions compared to protein-rich diets, aligning with

results from the present study.

Energy expenditure and energy assimilation in BSFL

To evaluate how efficiently the BSFL converted consumed energy into body mass, the gross

cost of growth was analysed; the amount of metabolic energy expended per unit of energy

assimilated into biomass. Energy expenditure was calculated from VO_2 measurements using

a caloric conversion factor (20.5 kJ L^{-1} O_2) (Schmidt-Nielsen, 1997), while biomass gain

was converted into energy equivalents based on lipid and protein content (37 kJ g^{-1} lipid and

17 kJ g^{-1} protein) (FAO, 2002). This approach assumes predominantly aerobic metabolism

and integrates both growth and maintenance related costs unlike more advanced studies that

can partition these costs in individual larvae (Eriksen, 2022, 2024).

Substantial variation in the gross cost of growth was observed among diets. BSFL reared on

the high-protein diet (P:C 1:3) spent 0.6 kJ per kJ lipid and protein assimilated, which is 33%

more energy than the high-carbohydrate diet (P:C 1:9) (Figure 3B). When considering protein

assimilation alone, larvae on P:C 1:3 required ~47% less energy to store one unit of protein

than larvae reared on P:C 1:9, who required 4.5 kJ of energy per kJ of protein stored (Figure

3C). However, these differences should be interpreted with great caution, as it is somewhat

naive to isolate assimilation of only one component in the BSFL biomass and then correlate

this to the overall energy use of the whole bioconversion system. Furthermore, a substantial

part of energetic costs is not directly associated with assimilation, but rather with general

maintenance (Eriksen, 2022, 2024) and it is possible that protein-rich larvae have higher

maintenance costs per mass unit than do lipid rich larvae. Deviations between the uncon-

strained (black) and origin-forced (red) linear models in Figure 3C further suggest that factors

beyond a simple proportional relationship between metabolic cost and nutrient gain influence

these patterns. While the biological interpretation of these energetic balances should be inter-

preted with caution it remains interesting for producers to evaluate at what cost either protein

or energy is assimilated in different diets. Our data show that the energetic cost per unit of

assimilated energy varies with diet composition (Figure 3). While both lipid and protein depo-

sition are closely linked to metabolic activity, protein synthesis is inherently more energeti-

cally demanding than lipid accumulation. Storing lipid appears to be metabolically less costly

per energy unit gained than synthesizing protein (Goodrich *et al.*, 2024; Laganaro *et al.*, 2021;

Talal *et al.*, 2021). As a result, larvae allocating more of their nutrient accumulation to lipid

biomass effectively grow "cheaper" in energetic terms per unit accumulated biomass than

those accumulating more protein.

Respiratory exchange ratio as predictor of BSFL body composition

Oxygen consumption provides a strong measure of total respiratory energy expenditure, while

carbon dioxide production reflects both respiratory metabolism and the biochemical conver-

sion of excess dietary carbohydrates (or amino acids) into lipids via lipogenesis (Laganaro *et*

al., 2021; Talal *et al.*, 2021). The respiratory exchange ratio (RER = VCO_2/VO_2) therefore

reflects the combined inputs of multiple metabolic processes related to aerobic and anaerobic

anabolism and catabolism of a system. RER values typically range from ~0.7 for lipid oxida-

tion to ~1.0 for carbohydrate oxidation, reflecting the macronutrient used for energy produc-

tion (Schmidt-Nielsen, 1997). However, in fast-growing systems, such as BSFL, lipogenesis

can drive values above 1.0 by producing CO_2 without consuming O_2, making RER a useful

indicator relative to anabolic activity associated with lipid synthesis (Ferrannini, 1988; Talal *et*

al., 2021; Parodi *et al.*, 2020; Schøn *et al.*, 2024). In accordance, higher RER values were

observed for the high-carbohydrate diet (P:C 1:9) that produced larvae with higher lipid and

lower protein content (Figure 4, Table 2). This pattern is consistent with carbohydrate-driven

lipogenesis, where surplus dietary energy from carbohydrates is stored as lipid. The correla-

tion between RER values and the larval protein and lipid content for different treatment diets

(Figure 4) highlights the potential of RER to indicate final body composition of BSFL.

Since RER is calculated from two separately measured values (CO_2 and O_2), it is sensitive

to measurement variability, so averaging RER over several days or the full 7-day growth pe-

riod may provide a more robust estimate. However, if the data is sampled with sufficient qual-

ity RER holds the potential to indicate to what degree larvae are allocating energy primarily

toward lipogenesis or other metabolic processes. It should be noted that VCO_2 and VO_2

measurements, and thereby the estimation of RER, can be confounded by CO_2 production
and O_2 consumption from diet-associated microbes and the microbial communities within
the larval digestive tract. For instance, other studies, such as Fuhrmann *et al.* (2025) have
demonstrated substantial diet microbial CO_2 production, which may complicate or reduce the
precision of nutrient allocation using RER. In contrast, the CO_2 production curves observed
in this present study are typical for bioconversion systems with low microbial respiration from
the diet, suggesting a more accurate reflection of larval metabolism. Under these conditions,
RER can serve as a reliable indicator of larval metabolism, allowing both cumulative and real-
time tracking of energy use and nutrient allocation, and acting as a non-invasive tool to pre-
dict nutrient assimilation in BSFL.
459
Conclusion and perspectives
This study focused on evaluating BSFL diet effects through the lens of respirometry, demon-
strating how respiratory measurements can be applied as predictive, non-invasive tools for
growth performance, nutrient allocation, and energetic efficiency. Across diets, the volume of
O_2 consumed correlated strongly with final larval biomass, while RER reflected differences
in protein and lipid deposition, confirming that dietary P:C ratio shapes both body composi-
tion and the energetic cost of growth. Lipid-rich larvae on a high-carbohydrate diet grew more
efficient in energetic terms than protein-rich larvae on a high-protein diet, highlighting practi-
cal implications for feed formulation and economic optimization.
Three key conclusions emerge: (i) dietary P:C ratio influences both growth and body com-
position, (ii) nutrient balance shapes the cost of growth and metabolic activity, and (iii) respi-
ratory variables (VO_2, VCO_2, RER) are robust predictors of larval performance, energetics,
and nutrient allocation. In a BSFL production facility, integrating O_2 with existing CO_2
monitoring could link gas exchange to biomass yield and nutrient profiles, enabling real-time
production monitoring and potentially improving decision-making, resource use, and valorisa-
tion of organic side streams. Combined with growth data, respirometry may serve as an indi-
cator of larval energy investment, offering prospects for more precise management and the
development of efficient, sustainable insect production systems.
478
479 **Data availability**
480 Data supporting this study will be made available upon publication.
481
482 **Declaration of competing interest**

The authors declare that they have no known competing financial interests or personal relationships that could have influenced the work reported in this paper. This includes Moritz Gold who was employed at REPLOID Group AG in parallel to ETH Zurich during the writing phase of the presented research.

References

Barragan☐Fonseca, K. B., Gort, G., Dicke, M., & van Loon, J. J. A. (2019). Effects of dietary protein and carbohydrate on life☐history traits and body protein and fat contents of the black soldier fly *Hermetia illucens*. *Physiological Entomology*, *44*(2), 148–159. https://doi.org/10.1111/phen.12285

Barragan-Fonseca, K. B., Gort, G., Dicke, M., & van Loon, J. J. A. (2021). Nutritional plasticity of the black soldier fly (*Hermetia illucens*) in response to artificial diets varying in protein and carbohydrate concentrations. *Journal of Insects as Food and Feed*, *7*, 51–61. https://doi.org/10.3920/JIFF2020.0034

Bekker, N. S., Heidelbach, S., Vestergaard, S. Z., Nielsen, M. E., Riisgaard-Jensen, M., Zeuner, E. J., Bahrndorff, S., & Eriksen, N. T. (2021). Impact of substrate moisture content on growth and metabolic performance of black soldier fly larvae. *Waste Management*, *127*, 73–79. https://doi.org/10.1016/j.wasman.2021.04.028

Berggreen, I. E., Schøn, M. L., Nørgaard, J. V., & Jensen, K. (2025). Optimal dietary protein content and essential amino acid limitation in larvae of the black soldier fly (Hermetia illucens). *Journal of Insects as Food and Feed*, 1–12. https://doi.org/10.1163/23524588-bja10221

Bosch, G., & Swanson, K. S. (2021). Effect of using insects as feed on animals: Pet dogs and cats. *Journal of Insects as Food and Feed*, *7*(5), 795–805. https://doi.org/10.3920/JIFF2020.0084

Broeckx, L., Frooninckx, L., Slegers, L., Berrens, S., Noyens, I., Goossens, S., Verheyen, G., Wuyts, A., & Van Miert, S. (2021). Growth of Black Soldier Fly Larvae Reared on Organic Side-Streams. *Sustainability*, *13*(23), 12953. https://doi.org/10.3390/su132312953

Deruytter, D., Coudron, C., Wiklicky, V., El Deen, S. N., Van Peer, M., Rumbos, C., Peguero, D. A., Adamaki-Sotiraki, C., Athanassiou, C., Veldkamp, T., Sandrock, C., & Yakti, W. (2025). *BugBook: Considerations for designing and performing insect larvae production experiments.* https://doi.org/10.1163/23524588-bja10252

Eggink, K. M., & Dalsgaard, J. (2023). Chitin contents in different black soldier fly (*Hermetia illucens*) life stages. *Journal of Insects as Food and Feed*, *9*(7), 855–864. https://doi.org/10.3920/JIFF2022.0142

Eggink, K. M., Donoso, I. G., & Dalsgaard, J. (2023). Optimal dietary protein to carbohydrate ratio for black soldier fly (*Hermetia illucens*) larvae. *Journal of Insects as Food and Feed*, *9*(6), 789–798. https://doi.org/10.3920/JIFF2022.0102

Eriksen, N. T. (2022). Dynamic modelling of feed assimilation, growth, lipid accumulation, and CO2 production in black soldier fly larvae. *PLOS ONE*, *17*(10), e0276605. https://doi.org/10.1371/journal.pone.0276605

Eriksen, N. T. (2024). Metabolic performance and feed efficiency of black soldier fly larvae. *Frontiers in Bioengineering and Biotechnology*, *12*. https://doi.org/10.3389/fbioe.2024.1397108

FAO. (2002). *Food energy: Methods of analysis and conversion factors□; report of a technical workshop, Rome, 3 - 6 December 2002*. Food and Agriculture Organization of the United Nations.

FAO. (2009). *How to Feed the World in 2050*. Food and Agriculture Organization of the United Nations. https://www.fao.org/fileadmin/templates/wsfs/docs/expert_paper/How_to_Feed_the_World_in_2050.pdf

Ferrannini, E. (1988). The theoretical bases of indirect calorimetry: A review. *Metabolism*, *37*(3), 287–301. https://doi.org/10.1016/0026-0495(88)90110-2

Foley, J. A., Ramankutty, N., Brauman, K. A., Cassidy, E. S., Gerber, J. S., Johnston, M., Mueller, N. D., O'Connell, C., Ray, D. K., West, P. C., Balzer, C., Bennett, E. M., Carpenter, S. R., Hill, J., Monfreda, C., Polasky, S., Rockström, J., Sheehan, J., Siebert, S., … Zaks, D. P. M. (2011). Solutions for a cultivated planet. *Nature*, *478*(7369), 337–342. https://doi.org/10.1038/nature10452

Fuhrmann, A., Wilde, B., Conz, R. F., Kantengwa, S., Konlambigue, M., Masengesho, B., Kintche, K., Kassa, K., Musazura, W., Späth, L., Gold, M., Mathys, A., Six, J., & Hartmann, M. (2022). Residues from black soldier fly (*Hermetia illucens*) larvae rearing influence the plant-associated soil microbiome in the short term. *Frontiers in Microbiology*, *13*, 994091. https://doi.org/10.3389/fmicb.2022.994091

Fuhrmann, A., Gold, M., Loh, R. K., Chu, C. X., Haberkorn, I., Puniamoorthy, N., & Mathys, A. (2025). Physical properties of food waste influence the efficiency of black soldier fly larvae bioconversion via microbial activity. Journal of Environmental Management, 387, 125777. https://doi.org/10.1016/j.jenvman.2025.125777

Gligorescu, A., Toft, S., Hauggaard□Nielsen, H., Axelsen, J. A., & Nielsen, S. A. (2019). Development, growth and metabolic rate of *Hermetia illucens* larvae. *Journal of Applied Entomology*, *143*(8), 875–881. https://doi.org/10.1111/jen.12653

Gold, M., Cassar, C. M., Zurbrügg, C., Kreuzer, M., Boulos, S., Diener, S., & Mathys, A. (2020). Biowaste treatment with black soldier fly larvae: Increasing performance through the formulation of biowastes based on protein and carbohydrates. *Waste Management*, *102*, 319–329. https://doi.org/10.1016/j.wasman.2019.10.036

Gold, M., Tomberlin, J. K., Diener, S., Zurbrügg, C., & Mathys, A. (2018). Decomposition of biowaste macronutrients, microbes, and chemicals in black soldier fly larval treatment: A review. *Waste Management*, *82*, 302–318. https://doi.org/10.1016/j.wasman.2018.10.022

556   Goodrich, H. R., Wood, C. M., Wilson, R. W., Clark, T. D., Last, K. B., & Wang, T. (2024). Specific
557   dynamic action: The energy cost of digestion or growth? *Journal of Experimental Biology*,
558   *227*(7). https://doi.org/10.1242/jeb.246722

559   Janssen, R. H., Vincken, J.-P., an den Broek, L. A. M., Fogliano, V., & Lakemond, C. M. M. (2017).
560   Nitrogen-to-Protein Conversion Factors for Three Edible Insects: *Tenebrio molitor*,
561   *Alphitobius diaperinus*, and *Hermetia illucens*. *Journal of Agricultural and Food Chemistry*,
562   *65*(11), 2275. https://doi.org/10.1021/acs.jafc.7b00471

563   Jensen, K., Mayntz, D., Wang, T., Simpson, S. J., & Overgaard, J. (2010). Metabolic consequences of
564   feeding and fasting on nutritionally different diets in the wolf spider *Pardosa prativaga*. *Jour-*
565   *nal of Insect Physiology*, *56*(9), 1095–1100. https://doi.org/10.1016/j.jinsphys.2010.03.001

566   Laganaro, M., Bahrndorff, S., & Eriksen, N. T. (2021). Growth and metabolic performance of black
567   soldier fly larvae grown on low and high-quality substrates. *Waste Management*, *121*, 198–
568   205. https://doi.org/10.1016/j.wasman.2020.12.009

569   Lomonaco, G., Franco, A., De Smet, J., Scieuzo, C., Salvia, R., & Falabella, P. (2024). Larval Frass of
570   Hermetia illucens as Organic Fertilizer: Composition and Beneficial Effects on Different
571   Crops. *Insects*, *15*(4), Article 4. https://doi.org/10.3390/insects15040293

572   Nguyen, T. T. X., Tomberlin, J. K., & Vanlaerhoven, S. (2015). Ability of Black Soldier Fly (Diptera:
573   Stratiomyidae) Larvae to Recycle Food Waste. *Environmental Entomology*, *44*(2), 406–410.
574   https://doi.org/10.1093/ee/nvv002

575   Parodi, A., de Boer, I. J. M., Gerrits, W. J. J., van Loon, J. J. A., Heetkamp, M. J. W., van Schelt, J.,
576   Bolhuis, J. E., & van Zanten, H. H. E. (2020). Bioconversion efficiencies, greenhouse gas and
577   ammonia emissions during black soldier fly rearing – A mass balance approach. *Journal of*
578   *Cleaner Production*, *271*. Scopus. https://doi.org/10.1016/j.jclepro.2020.122488

579   Purkayastha, D., & Sarkar, S. (2022). Sustainable waste management using black soldier fly larva: A
580   review. *International Journal of Environmental Science and Technology*, *19*(12), 12701–
581   12726. https://doi.org/10.1007/s13762-021-03524-7

582   R Core Team. (2023). *R: A language and environment for statistical computing. R Foundation for*
583   *Statistical Computing* [Computer software].

584   RStudio Team. (2023). *RStudio: Integrated Development for R. RStudio, PBC.* [Computer software].

585   Schmidt-Nielsen, K. (1997). *Animal Physiology: Adaptation and Environment* (5th ed.). Cambridge
586   University Press.

587   Schøn, M. L., Mikkelsen, M. V. N., Jensen, K., Poulsen, J. M., Berggreen, I. E., Schou, T. M.,
588   Nørgaard, J. V., & Overgaard, J. (2024). Effect of temperature on growth, metabolism, and gas
589   exchange in *Hermetia illucens* larvae reared under commercial and laboratory conditions.
590   *Journal of Insects as Food and Feed*, 1–16. https://doi.org/10.1163/23524588-00001268

Schow-Madsen, M., Jensen, K., Toribio, M., Schøn, M. L., Schou, T. M., Nørgaard, J. V., & Over-
gaard, J. (2025). Consequences of dietary amino acid imbalance on growth and gas exchange
in black soldier fly (*Hermetia illucens*) larvae. *In Preparation*.

Talal, S., Cease, A., Farington, R., Medina, H. E., Rojas, J., & Harrison, J. (2021). High carbohydrate
diet ingestion increases post-meal lipid synthesis and drives respiratory exchange ratios above
1. *Journal of Experimental Biology*, *224*(4), jeb240010. https://doi.org/10.1242/jeb.240010

van der Heide, M. E., Stødkilde, L., Værum Nørgaard, J., & Studnitz, M. (2021). The Potential of Lo-
cally-Sourced European Protein Sources for Organic Monogastric Production: A Review of
Forage Crop Extracts, Seaweed, Starfish, Mussel, and Insects. *Sustainability*, *13*(4), 2303.
https://doi.org/10.3390/su13042303

Table 1: Macronutrient contents of the four experimental diets. The reference diet (REF) is an
entirely commercial chicken feed used as a reference diet supporting high growth. The three other
diets are formulated in house and designed to represent different protein-to-carbohydrate (NFC) ratios.

Diet treatment	REF	P:C 1:3	P:C 1:5	P:C 1:9
Protein-to-carbohydrate ratio (P:C)	1 : 3.2	1 : 3.2	1 : 5.0	1 : 9.4
Ingredients **(% of dry mass)**				
Chicken feed (CF)	97.6	34.1	34.1	34.1
Protein mix	0	10.4	5.2	0
Carbohydrate mix	0	33.3	38.6	43.7
Lipid mix	0	2.7	2.7	2.7
Additive mix	0	3.9	3.9	3.9
Cellulose	0	13.1	13.1	13.1
Agar	2.4	2.4	2.4	2.4
Diet composition **(% of dry mass)**				
Protein	19.9	17.3	12.1	7.0
Carbohydrate (NFC)	63.3	55.5	60.7	65.8
Carbohydrate (fibre, agar)	5.6	16.7	16.7	16.7
Lipid	5	4.5	4.5	4.5
Ash	6.2	6.1	6.1	6.1

4 The diet composition is based on the ingredient list and the mass used. The non-fibre carbohydrate (NFC) fraction from
5 chicken feed is estimated by subtracting protein, fat, fibre, and ash from the total dry mass.

Table 2: Survival, growth, body composition and gas exchange of black soldier fly larvae after 7 days of rearing on diets with different protein and carbohydrate contents.

Growth performance	5–6 day old larvae* At onset	REF After 7 days	P:C 1:3 After 7 days	P:C 1:5 After 7 days	P:C 1:9 After 7 days
Survival rate (%)		$97.0 \pm 4.4\%$ [a]	$94.9 \pm 3.6\%$ [a]	$92.9 \pm 5.5\%$ [a]	$88.8 \pm 8.1\%$ [a]
Larvae end biomass (g DM per replicate)	0.78 ± 0.15 g	25.9 ± 0.5 g [a]	22.6 ± 0.7 g [b]	20.1 ± 1.1 g [c]	12.6 ± 3.0 g [d]
Specific growth rate (per day)		0.50 ± 0.03 [a]	0.47 ± 0.03 [ab]	0.45 ± 0.02 [b]	0.36 ± 0.03 [c]
Larvae percent dry matter (% DM)	26.3%	$27.7 \pm 1.5\%$ [a]	$27.7 \pm 0.8\%$ [a]	$29.0 \pm 2.2\%$ [a]	$32.7 \pm 1.5\%$ [b]
Larvae crude protein (% of DM)	40.9%	$31.5 \pm 0.8\%$ [a]	$32.5 \pm 0.6\%$ [a]	$25.5 \pm 2.3\%$ [b]	$19.8 \pm 0.8\%$ [c]
Larvae crude lipid (% of DM)	10.0%	24.3 ± 1.9 % [a]	27.5 ± 3.5 % [a]	$39.2 \pm 6.7\%$ [b]	$52.7 \pm 2.7\%$ [c]
Gas exchange measurements					
VCO_2 (L, total for 7 days)		19.5 ± 1.2 L [a]	16.9 ± 1.1 L [b]	16.2 ± 1.6 L [b]	10.8 ± 2.2 L [c]
VO_2 (L, total for 7 days)		17.7 ± 1.3 L [a]	14.2 ± 1.2 L [b]	12.9 ± 1.5 L [b]	8.0 ± 1.7 L [c]
RER		1.10 ± 0.02 [a]	1.20 ± 0.03 [b]	1.26 ± 0.04 [c]	1.35 ± 0.06 [d]

* Samples of 5-6-day old seed larvae were sampled for analysis at the onset of the experiment.
DM = dry matter; V = volume; RER = respiratory exchange ratio, calculated from total VCO_2/VO_2 over the 7-day experiment. All values are mean ± standard deviation and dissimilar letters indicate significant differences (one-way ANOVA followed by Tukey HSD test, $p < 0.05$, n = 7)

13
Figure 1: Effect of diets with different protein-to-carbohydrate ratios on total growth and estimated daily
specific growth rate (SGR). A) Average individual larvae mass (mg wet mass) over time, where the insert
shows the final total dry larvae biomass at the termination of the experiment (approximately 600 larvae). B)
Daily estimates of SGR, calculated from the increase in average body mass estimated between two consecutive
days, where the insert shows SGR calculated for the entire growth period. Bold lines represent treatment means,
while semi-transparent lines show the individual replicates (n=7). Inserted bar plots show treatment means ±
standard deviations, and dissimilar letters indicate significant differences (one-way ANOVA followed by Tukey
HSD test, $p < 0.05$, n = 7).

22
23

26
Figure 2: Effect of diet with different protein-to-carbohydrate ratios on gas exchange measurements. A)
Change in O☐ consumption rate ($\dot{V}O$☐, mL/min/chamber) over time, where the insert shows the total VO☐
consumed for the entire period (approximately 600 larvae). B) Change in respiratory exchange ratio (RER) over
time $\dot{V}CO$☐/$\dot{V}O$☐ ($\dot{V}$ is the rate, mL/min), where insert show total RER VO☐/VCO☐ (V is the volume, L) for
each treatment. Bold lines represent treatment means, while semi-transparent lines show the individual replicates
(n=7). Inserted bar plots show treatment means ± standard deviations, and dissimilar letters indicate significant
differences (one-way ANOVA followed by Tukey HSD test, $p < 0.05$, n = 7).

34
Figure 3: Analysis of the metabolic cost of growth and assimilation in larvae reared on diets with different
protein-to-carbohydrate (P:C) ratio. A) Total oxygen consumption over 7 days as a function of final biomass
across replicates reared on 4 different diets with varying P:C ratio. Correlation shows a linear relationship
between the total volume of O☐ consumed (L) and the final dry biomass (g). B) and C) show the energy
expenditure of the system (using the energy equivalent of VO☐) in relation to the assimilated energy in the
biomass (kJ) represented by gain in lipid and protein, or protein only. Bold points represent treatment means
with error bars showing standard deviations, and semi-transparent points represent treatment replicates (n=7).
Black lines represent a linear model for the replicates (n=28), and red dashed lines represent the linear model
when forced through origo (0,0). Linear formulas and R^2 values are displayed for each regression model.
44
45
46
47

48
Figure 4: Effect of diets with different protein-to-carbohydrate ratios on larval body composition.
Relationship between the measured RER values and A) crude lipid and B) crude protein content (% of dry mass).
Bold points represent treatment means with error bars showing standard deviations, and semi-transparent points
represent treatment replicates (n=7). The black lines represent a linear model for all the replicate data (n=28).
Linear models for each treatment group are shown in their respective colours (n=7). Linear formulas and R^2
values are displayed for each regression model.

Supplementary material
56

57
58
Figure S1: Effect of diet with different protein-to-carbohydrate ratios on CO₂ production. Change in
CO₂ production rate (V̇CO₂, mL/min/chamber) over time, where the insert shows the total VCO₂ produced for
the entire period (approximately 600 larvae). Bold lines represent treatment means, while semi-transparent lines
show the individual replicates (n=7). Inserted bar plots show treatment means ± standard deviations, and
dissimilar letters indicate significant differences (one-way ANOVA followed by Tukey HSD test, p < 0.05, n =
7).
65
66

67
68   **Figure S2: Effect of diet with different protein-to-carbohydrate ratios on the cost of biomass.** Daily
69   estimates of the metabolic costs of biomass, when calculated as consumed VO₂ in relation to biomass gain. Bold
70   lines represent treatment means, while semi-transparent lines show the individual replicates (n=7). Inserted bar
71   plot show treatment means ± standard deviations, and dissimilar letters indicate significant differences (one-way
72   ANOVA followed by Tukey HSD test, p < 0.05, n = 7).

Model illustration	Model formula	Intercept	Slope	Slope SE	R^2	Adj. R^2	Residual SE	DF	F-statistic	p-value	Pearson correlation coefficient
Fig. 3A black	Total VO_2 (L) ~ Final larvae biomass (g DM)	-1.04	0.70	0.04	0.933	0.931	1.00	26	363	**<2.2 · 10⁻¹⁶**	0.966
Fig. 3A red	Total VO_2 (L) ~ Final larvae biomass (g DM) + 0	0.00	0.65	0.01	0.995	0.995	1.02	27	5087	**<2.2 · 10⁻¹⁶**	
Fig. 3B black	Energy expenditure (kJ) ~ Protein and lipid energy gain (kJ)	-37.71	0.90	0.21	0.425	0.403	60.18	26	19	**1.7 · 10⁻⁴**	0.652
Fig. 3B red	Energy expenditure (kJ) ~ Protein and lipid energy gain (kJ) + 0	0.00	0.80	0.03	0.957	0.955	59.37	27	601	**<2.2 · 10⁻¹⁶**	
Fig. 3C black	Energy expenditure (kJ) ~ Protein energy gain (kJ)	96.41	1.88	0.14	0.875	0.870	28.07	26	181	**3.1 · 10⁻¹³**	0.935
Fig. 3C red	Energy expenditure (kJ) ~ Protein energy gain (kJ) + 0	0.00	2.77	0.09	0.974	0.973	46.34	27	1005	**<2.2 · 10⁻¹⁶**	
Fig. 4A black	Larvae crude lipid (% DM) ~ RER	-82.59	96.53	14.00	0.647	0.633	7.28	26	48	**2.5 · 10⁻⁷**	0.804
Fig. 4A treatments	Larvae crude lipid (% DM) ~ RER · Treatment	See Fig. 4A	See Fig. 4A		0.914	0.883	4.10	20	30	**2.7 · 10⁻⁹**	
Fig. 4B black	Larvae crude protein (% DM) ~ RER	79.86	-42.79	6.34	0.636	0.622	3.30	26	46	**3.7 · 10⁻⁷**	-0.798
Fig. 4B treatments	Larvae crude protein (% DM) ~ RER · Treatment	See Fig. 4B	See Fig. 4B		0.954	0.938	1.34	20	59	**5.8 · 10⁻¹²**	

76   DM = dry matter; V = volume; RER = respiratory exchange ratio, calculated from total VCO_2/VO_2 over the 7-day experiment; Treatments = REF, P:C 1:3, P:C 1:5, and P:C 1:9; SE = standard
77   error.