Phasing single-molecule nano-NOMe-seq reveals chromatin state heterogeneity in the context of transcription and long-range interactions

Stephanie Bellini [1], Catherine Do[1*] & Jane A Skok[1,2,*]

1.   Department of Pathology, NYU Grossman School of Medicine, New York, NY, USA
2.   Perlmutter Cancer Center, NYU Langone Health, New York, NY, USA
*   These authors contributed equally

Corresponding author - Catherine Do

ABSTRACT

A central challenge in molecular biology is determining how 3D chromatin architecture, particularly enhancer-promoter looping and insulating CTCF-mediated interactions, influences gene transcription in individual cells, which has significant implications for healthy and diseased states. To overcome current limitations in imaging and genomic technologies, we developed a cluster-based phasing strategy using long read nano-NOMe-seq to link distinct CTCF binding states—captured at the single molecule level—to the transcriptional status of genes. By stitching partially overlapping long reads and clustering them by shared GpC-accessibility patterns, we stratify CTCF into graded binding states on individual molecules, classify RNA polymerase states at promoters/gene bodies, and infer when spatially separated loci are coordinately activated and occupy loop-competent configurations on the same molecules. When applied to *Sox2*, *Hoxa*, and *Klf1* regions, cluster-based nano-NOMe-seq phasing reveals how specific topologies bias polymerase behavior and multi-locus activity in ways that bulk assays or locus-engineered imaging cannot fully capture.

INTRODUCTION

Three-dimensional (3D) organization of chromatin is central to gene regulation. Bulk chromosome conformation capture approaches such as Hi-C and Micro-C generate genome-wide contact maps that reveal two nested axes of organization. First, chromatin segregates into higher-order active (A) and inactive (B) compartments [3]. Second, within and between these compartments, CTCF and cohesin extrude and stabilize loops that partition the genome into highly self-interactive topologically associating domains (TADs) of ~1 Mb or less [4-6]. TADs act as functional units that insulate enhancer–promoter activity, thereby constraining gene regulation within defined boundaries, while this hierarchical architecture brings distant elements into proximity to enable long-range enhancer–promoter interactions that control transcription [7].

Beyond these domains, finer-scale interactions within microcompartments were uncovered by Region Capture Micro-C (RCMC) [8], that constitute highly focal contacts linking active enhancers and promoters. These interactions persist after acute cohesin depletion or short-term transcriptional blockade by triptolide, indicating that microcompartments do not require loop extrusion or ongoing transcription on short timescales [8]. However, with extended RNAPII depletion or chronic inhibition, enhancer–promoter contacts weaken globally, consistent with transcription contributing indirectly to the long-term maintenance of these interactions [9].

More recently, single-cell Hi-C and related technologies have extended these observations to the level of individual nuclei [10]. By capturing contact maps in single cells, these approaches can quantify the probability of specific loops or compartmental interactions, though they cannot yet determine whether loop formation and gene expression are coordinated within the same cell. This highlights a key limitation in current technologies: resolving chromatin into its component, heterogeneous states, which reflect the dynamics of interactions in the context of gene activation remains a largely unresolved issue. Indeed, conformation and expression profiling approaches, whether in bulk or single-cell, are uncoupled such that although they can reveal which chromatin contacts are possible or prevalent, and which genes are expressed in a population, they are unable to determine whether active transcription and looping co-occur, or are independent of one another in the same cell for a given locus, and what proportion of alleles/cells are both looped and actively transcribing within a sample.

Live-cell imaging approaches that simultaneously label chromatin and nascent RNA can address this issue. CRISPR/dCas9-based genomic tagging of non-repetitive loci [11] has been enhanced by

signal-amplifying scaffold systems such as Casilio and CRISPR-Sirius [12], which make it possible

to visualize weakly fluorescent, low-copy regions in living cells. These chromatin labels can be

combined with RNA reporters, such as MS2 or PP7 stem-loop systems that tag nascent

transcripts [13,14] or more recently dCas13-based RNA imaging strategies. Using this integrated

approach, Alexander *et al.,* [15] showed that transcriptional activation at the *Sox2* locus can occur

without stable enhancer–promoter juxtaposition. More recently, Kim *et al.* [16] introduced SiCLAT, a

dual-color system that simultaneously labels loop anchors and nascent RNA in primary cells,

extending live imaging beyond immortalized cell lines. Despite these advances, such techniques

remain restricted to a handful of engineered loci.

Complementary fixed-cell imaging methods have provided a powerful means to interrogate

chromatin architecture at high resolution in single cells. Optical Reconstruction of Chromatin

Architecture (ORCA [17]) uses sequential oligo-FISH to trace chromatin folding with ~2 kb resolution

across hundreds of kilobases. Applied to the *Sox2* locus, ORCA revealed fine-scale

heterogeneous rosette conformations in embryonic stem and neural progenitor cells [1]. Hi-M [18,19]

in contrast, was specifically designed to integrate sequential DNA FISH with RNA FISH in intact

Drosophila embryos, enabling simultaneous visualization of chromatin architecture and

transcriptional output at the same loci. Together, these approaches highlight how fixed-cell

imaging can achieve either high-resolution structural reconstruction (ORCA) or joint analysis of

structure and transcription (Hi-M), though both remain restricted to targeted loci rather than

genome-wide views.

Emerging sequencing-based multi-omic single-cell platforms now enable genome-wide coupling

of chromatin structure with transcription. Simultaneous single-cell Hi-C and RNA-seq [20] directly

link genome-wide contact maps with full transcriptomes from the same cells, revealing context-

dependent coupling between enhancer–promoter interactions and gene expression. Other

strategies, including HiRES [20], LiMCA [20], and GAGE-seq [21], employ variations in library

preparation and indexing to balance sensitivity and throughput. These technologies overcome the

locus restriction of imaging but remain limited by high sequencing depth requirements and cost.

Together, these imaging and sequencing approaches highlight both the progress and current

limitations in resolving the temporal and functional coupling of chromatin looping and transcription

at single-cell resolution.

To overcome these constraints, we made use of long read nano-NOMe-seq to link distinct CTCF

binding states—captured at the single molecule level—to the transcriptional status of genes.

Using a cluster-based approach analogous to pseudobulk analyses in single-cell datasets, we

could grade CTCF binding at a near single cell level, classify Pol II states at promoters and across

gene bodies, and infer when dispersed loci are co-activated and in loop-ready configurations on

the same molecules. Applied to the *Sox2*, *Hoxa*, and *Klf1* regions, this cluster-phased view reveals

how specific topologies bias polymerase behavior and multi-locus activity—features not fully

captured by ensemble assays or engineered-locus imaging.

RESULTS

Clustering of nucleosome occupancy reveals graded CTCF binding and Pol II–dependent

transcription states.

To identify distinct within-sample binding states for CTCF and Pol II, we performed long-read

Nano-NOMe-seq in mESCs. This enabled us to assess genome-wide nucleosome positioning

using exogenous 5mC at HGC (GC excluding CGC) at the single molecule (SM) level[22] (**Figure**

1A). Unlike MNA-seq analyses, the single molecule (SM) data can be clustered to reveal different

CTCF and RNA Pol II binding states based on nucleosome occupancy patterns. For CTCF, we

analyzed a 2 kb region centered on CTCF binding sites (CBSs) validated by ChIP-seq. As shown

in the heatmap, this uncovered a continuum of binding states previously described by us [22]

(**Figure 1B**). Since nano-NOMe-seq libraries are PCR-free, for a given locus, the proportion of

each SM cluster is likely to reflect the percentage of cells with similar CTCF binding states.

To classify CTCF binding states, we aggregated clusters as unbound (light blue – no CTCF

footprint), weak/transitioning (purple – CTCF footprint adjacent or overlapping a nucleosome) or

strong (red – CTCF footprint within a nucleosome-free region -NFR). A similar analysis was

performed to identify within-sample Pol II transcription states, centering on ChIP-seq validated,

Pol II–bound transcription start sites (TSSs). SM clustering in the heatmap was ordered and

annotated based on a Pol II pausing index (PI) and enrichment for Pol II peaks (**Figure 1C**). Using

this approach, we classified sites as elongating, active with +1 nucleosome, paused and inactive.

Clustered-based phasing of Nano-NOMe-seq single molecules.

The goal of these studies was to link distinct CTCF binding states—captured at the single

molecule level—to the chromatin and transcriptional status of genes and their regulatory

elements, however the median read length of the SM nano-NOMe-seq was not sufficient for this

purpose (**Figure 2A** left) and stitching reads together was necessary to be able to incorporate

CTCF loop bases with intervening regulatory elements and genes across a region of interest.

Figure 2A (right) shows an example of SM methylation sequence at the GpC level across a locus

spanning 25 kb.

Figure 1. Clustering of nucleosome occupancy patterns from Nano-NOMe-seq reveals graded CTCF binding and Pol II–dependent transcription states. A. Cartoon illustrating the Nano-NOMe-seq strategy to identify single-molecule (SM) nucleosome positioning and transcription factor (TF) footprinting, based on exogenous 5mC levels in a GpC context. **B**. Heatmap and average profiles of 2 kb regions centered on CTCF binding sites, showing clustering of SMs in mESCs to define within-sample CTCF binding states. **C**. Heatmap and average profiles of 2 kb regions centered on Pol II–bound transcription start sites (TSSs), showing clustering of SMs in mESCs to define within-sample Pol II transcription states. Hyper-accessible clusters, C1-C3, were color-coded in blue and annotated as nascent chromatin [2]. Clusters are ordered and annotated based on Pol II pausing index (PI) and enrichment for Pol II peaks. ε=0.5.

Our approach for stitching is described in the scheme in **Figure 2B.** Nano-NOMe-seq reads from

a given long-range locus were binned into 50 bps, to obtain sufficient resolution to detect both

transcription factor and nucleosome footprints. The locus was then divided into overlapping

windows to facilitate stitching. SMs were clustered within each window using agglomerative

hierarchical clustering (**Figure S1A**), and clusters matched across consecutive overlapping

windows using the Hungarian algorithm (**Figure S1B**). After generating aggregated methylation

Figure 2. Clustered-based phasing of the Nano-NOMe-seq single molecules. A: Left, distribution of Nano-NOMe seq read lengths. The read dashed line indicates the median. Right, example of SM methylation sequence at the GpC level in a locus spanning 25 kb. **B**: For a given long-range locus, single molecules (Nano-NOMe-seq reads) are binned into 50 bp. The region is divided into overlapping windows to enable stitching: SMs are first clustered within each window using agglomerative hierarchical clustering, and clusters are then matched across consecutive overlapping windows using the Hungarian algorithm. After generating aggregated methylation profiles per cluster, each SM is reassigned to its best-matched cluster based on Hamming distance. This step corrects inconsistent assignments across windows and yields long-range phased SM clusters, which can be analyzed at either the bin or GpC level.

profiles for each cluster, SMs were reassigned to their best-matched neighboring clusters based

on the Hamming distance (**Figure S1C**). This step corrects inconsistent assignments across

windows and yields long-range phased SM clusters, which can be analyzed at either the bin or

GpC level.

An example of cluster-based phasing for a locus spanning 50 kb is shown in **Figure 3A**, with a

cartoon illustrating the approach used to estimate the accuracy of the algorithm across 90

extended loci, each mapped by at least one long (≥50 kb) read that is kept out of the analysis and

used for validation (**Figure 3B** and **Table S1**). For each locus, the methylation sequence of the

long read is compared to the mean methylation sequences of all clusters (c_j), and the closest

cluster is identified using the Hamming distance (d_H). $A(r_i)$ denotes the read accuracy, reflecting

how well read r_i matches its best-aligned cluster (higher values = better match). The overall

clustering accuracy is defined as the average of $A(r_i)$ across all long reads. The distribution of

read accuracy across all long reads is shown in **Figure 3C**. The global clustering accuracy was

0.78. Examples of long-read methylation sequences and their closest-matching clusters for three

loci spanning 50–60 kb are shown in **Figure 3D**.

Cluster-based phasing of nano-NOMe-seq data confirms CTCF-independent Sox2

transcription.

As proof of principle, cluster-based stitching/phasing of nano-NOMe-seq data was implemented

at the well-studied *Sox2* locus to determine whether there is regulatory heterogeneity in mESCs.

Sox2 expression is regulated by the *Sox2* control. The SCR contains two distal enhancers,

SRR107 and SRR111, which flank a CTCF binding site known as SRR109. Previous studies have

shown that, while deletion of the SCR in mice significantly reduces Sox2 expression [23], deletion

of the CTCF motif at SRR109 results in only a modest, non-significant decrease in Sox2 levels [24]

[25], suggesting that transcription is primarily mediated by CTCF-independent enhancer–promoter

interactions [26].

To determine whether we could confirm these results, we performed cluster-based phasing of

SMs using our previously generated nano-NOMe-seq data [22]. The average coverage and median

read lengths mapping the *Sox2* locus is reported in **Table S2**. Region Capture Micro-C (RCMC) [8]

(at 200 bp resolution) of the 150 kb *Sox2* locus in mESCs revealed a previously described CTCF-

mediated loop between the *Sox2* control region (SCR) and a CTCF binding site (window 1)

located 2 kb upstream of the *Sox2* transcription start site (TSS, window 2) (**Figure 4A**). The two

Figure 3. **Examples and accuracy of Nano-NOMe-seq single molecules clustered-based phasing. A**: Example of the clustered-based phasing algorithm for one locus spanning 50 kb. Missing data are color-coded in blue. **B**: Cartoon illustrating the approach used to estimate the accuracy of cluster-based phasing across 90 long-range reference loci, each mapped by at least one long (≥50 kb). For each locus, the methylation sequence of a long read is compared to the mean methylation sequences of all clusters (c_j), and the closest cluster is identified using the Hamming distance (d_H). $A(r_i)$ denotes the *read accuracy*, reflecting how well read r_i matches its best-aligned cluster (higher values = better match). The overall clustering accuracy is defined as the average of $A(r_i)$ across all long reads. **C**: Distribution of read accuracies across all long reads. The global clustering accuracy is 0.78. **D**: Examples of long-read methylation sequences and their closest-matching clusters for three loci spanning 50–60 kb.

179   confirmed binding of CTCF and active Sox2 transcription / elongation respectively in mESCs.

Figure 4: Cluster-based phasing of nano-NOMe-seq data confirms CTCF-independent Sox2 transcription.
A: Capture Micro-C (300 bp resolution) of the 150 kb Sox2 locus in mESCs (GSE207225) reveals a CTCF-mediated loop between the Sox2 control region (SCR) and a CTCF binding site (window 1) located 2 kb upstream of the Sox2 transcription start site (TSS, window 2). The SCR includes two enhancers, SRR107 (window 4) and SRR111 (window 6), flanking the downstream CTCF anchor (SRR109, window 5). CTCF motif orientations are shown below the CTCF ChIP-seq track. Pol II ChIP-seq (GSE207225) confirms active Sox2 transcription and elongation in mESCs. **B:** Single-molecule nucleosome occupancy clustering identifies distinct transcriptional states within the sample, reflecting dynamic Pol II pausing, release, and elongation. This window is centered on Pol II with the TSS located 30 bp downstream. **C:** Nano-NOMe-seq at CTCF anchors shows that only a subset of molecules exhibits a clear CTCF footprint and associated nucleosome-free region (NFR) at both anchors. **D:** Cluster-based phasing resolves the coupling between loop extrusion, enhancer–promoter interactions, and Sox2 transcription. Clusters 1, 2 and 5 exhibit CTCF-independent Sox2 transcription, whereas cluster 0 links Sox2 Pol II paused transcription to CTCF-mediated loop extrusion. SRR107 is accessible (active) across all clusters with active Sox2 transcription, while SRR111 is inactive in Cluster 2, where elongation is reduced. In cluster 0 and cluster 7, both SRR107 and SRR111 are inaccessible (inactive). CTCF motifs are colored according to orientation: blue for positive and green for negative.

In this locus, our cluster-based phasing approach allowed us to resolve regulatory elements within

9 clusters of single molecules (**Figure S2**). Clusters 0, 1, 2, 5, 7 displayed nucleosome patterns

indicative of distinct transcriptional states within the sample, reflecting dynamic Pol II activation

with +1 nucleosome, elongation and pausing, (**Figure 4B**). Nano-NOMe-seq at CTCF anchors

shows that only a subset of molecules exhibited a clear CTCF footprint and associated

nucleosome-free region (NFR) at both anchors (**Figure 4C**). We hypothesized that clusters
exhibiting CTCF footprint at chromatin loop anchors identified by chromatin capture assays and
overlapping CTCF motifs arranged in a convergent orientation (blue – forward, green – reverse),
likely represent the subset of cells harboring a CTCF-mediated loop. At the *Sox2* locus, only
~10% of the SMs/cells harbored a CTCF-mediated loop between the SCR and the CTCF binding

Figure 5: Cluster-based phasing of nano-NOMe-seq data identifies CTCF footprints at the bases of long-range multiway loops surrounding the *Sox2* locus. A: Capture Micro-C (1 kp resolution), CTCF, SMC3 and Pol II ChIP-seq across the extended 1 Mb *Sox2* locus in mESCs. Arcs depict CTCF-dependent loops predicted by nucleosome and CTCF footprinting in clusters with at least one predicted loop. The black rectangles highlight CTCF anchors of the *Sox2* rosette previously validated by ORCA [1]. **B**: Nano-NOMe-seq profiles of the 2 kb windows centered on the CTCF motifs involved in looping and on *Sox2* TSS. Arcs represent predicted CTCF-dependent loops with dashed line corresponding to loops connecting at least one anchor with a weaker CTCF footprint.

192   site located directly upstream of *Sox2* TSS, confirming the high within-sample heterogeneity of
193   chromatin architecture and CTCF binding states.
194
Figure 4D shows the relationship between *Sox2* transcription, enhancer activity at SRR107 and
SRR111, and CTCF-mediated looping, resolved at the SM cluster level. Clusters 1, 2 and 5 exhibit
CTCF-independent *Sox2* transcription (no CTCF footprints present), whereas cluster 0 links *Sox2*
Pol II paused transcription to the presence of CTCF footprints that could mediate the formation of
loops detected by RCMC. SRR107 is accessible (active) across all clusters with active *Sox2*
transcription, while SRR111 is inactive in cluster 2, where elongation is reduced. In cluster 0 and
cluster 7, both SRR107 and SRR111 are inaccessible (inactive). These results validate previous
findings [15,24-26], and explain why deletion of the CTCF motif in this region does not impact *Sox2*
expression. Furthermore, they highlight the connections between SRR107 and SRR111 activity
and elongation strength: elongation is more pronounced when both enhancers are accessible and
active, and reduced when only SRR107 is active.
206

Cluster-based phasing identifies CTCF footprints at the bases of long-range multiway loops including the *Sox2* locus.

We next extended our analyses downstream of the *Sox2* locus and SCR, to examine CTCF
footprints at bases of CTCF-mediated rosettes identified by ORCA [1] and loops also shown to be
present in RCMC [8] (**Figure 5A** and **Table S2**). Because of the reduced coverage within
overlapping windows over the extended region, we identified fewer clusters (7 in total, **Figure
S3**). We detected CTCF footprints and binding sites in the appropriate orientation to form loops
in clusters 1, 2 and 5 (arcs in **Figure 5A and 5B**). Our cluster-based analysis confirmed the
presence of CTCF footprints at CTCF anchors of the *Sox2* rosette (windows 1, 5 and 6) previously
validated by ORCA [1], but revealed a relatively weak CTCF footprint in window 6. Based on the
strength of the CTCF footprint and the finding that strong footprints highlight strong CTCF signals
that are better able to stop cohesin [27], we expect that stronger loops form between CBS sites with
stronger CTCF footprints (solid lines) and weaker loops form between CBS sites with weaker
CTCF footprints (dashed lines). Interestingly, we identified two CTCF footprints that could
mediate long-range loops connecting window 1 with windows 9 and 10 in cluster 5, and window
1 with window 10 in cluster 6 that can be visualized by RCMC, but that were not detected with
ORCA. These findings suggest that loops detected by bulk interaction data in the *Sox2* locus arise
from a mixture of two-way and higher-order chromatin interactions consistent with distinct rosette
formations (**Figure 5B**).

Figure 6: Cluster-based phasing of nano-NOMe-seq identifies long range high-order multiway CTCF dependent loops across the *Hoxa* region. A: Hi-C (5 kb resolution), CTCF, SMC3 and Pol II ChIP-seq across the 2.3 Mb *Hoxa1* locus in mESCs. Arcs show CTCF-dependent loops predicted by nucleosome and CTCF footprinting in each cluster. The black rectangles show CTCF anchors of the *Hoxa1* rosette previously validated by ORCA[1]. **B:** Nano-NOMe-seq profiles of the 2 kb windows centered on the CTCF motifs overlapping with the anchors of loops identified by Hi-C and ORCA. Sites with CTCF footprints and their potential for loop formation are highlighted by the orientation of the CTCF motif (blue – forward, green – reverse). **C:** Nano-NOMe-seq profiles of the 2 kb windows centered on Pol II summits at the TSS of genes within CTCF loops, colored according to Pol II states defined in **Figures 1** and **4**. The cartons on the right represent the chromatin conformation predicted by CTCF footprint (shown in **B**) for each cluster.

We did not observe an association between predicted *Sox2* transcription and higher-order CTCF-
dependent interactions. Consistent with our findings in **Figure 3**, both cluster 5 and cluster 6
chromatin conformations were permissive for *Sox2* expression in the absence of a potential
contact between the CTCF site at the TSS and the SCR (SRR109). In contrast, co-occurring
multiway loops between the downstream and both upstream CTCF sites (cluster 5) were linked
to weaker Pol II elongation. Since deletion of CTCF at the downstream sites has no effect on
Sox2 expression [26], we conclude that the multiway loops we detect do not contribute to SCR-
mediated effects on transcription in ESCs. However, we cannot rule out that in other contexts
these may be important for Sox2 gene regulation. Together, our cluster-based nano-NOMe-seq
CTCF and Pol II footprint analysis highlight the potential for identifying rosette structures across
extended regions, and the connections between loop formation and Pol II status.

Cluster-based phasing of nano-NOMe-seq identifies long-range high-order multiway CTCF
dependent loops across the *Hoxa* region.

CTCF-mediated rosette structures were also previously identified by ORCA across the *Hoxa*
region [1]. Using our cluster-based nano-NOMe-seq we asked if we could identify nucleosome and
CTCF footprinting at the bases of these loops, together with the concomitant Pol II occupancy
state (**Table S2**). Our previously generated Hi-C data [27] revealed pairwise chromatin interactions
between ORCA-identified loop anchors and additional CTCF/SMC3 sites across the 2.3 Mb
domain (**Figure 6A)**. ORCA loop anchors corresponded exclusively to bound CTCF motifs in the
negative orientation, suggesting that additional upstream CTCF binding sites in the positive
orientation contribute to the rosette structure identified by imaging analysis. Among the nano-
NOMe-seq SM clusters, only cluster 5 exhibited co-occurring CTCF footprints at all three ORCA
loop anchors (**Figure 6B**). In this cluster, we also identified two weak upstream CTCF footprints
(window 1 and 2) mapping to motifs in the positive orientation that could enable the rosette
formation by providing the required convergent CTCF binding. These were not included in the
ORCA analyses of three-way interactions. Cluster 5 additionally showed footprints at multiple
CTCF binding sites in both orientations between window 3 and 9, consistent with higher-order
multiway interactions beyond those originally described. The other clusters displayed a spectrum
of two-way and multiway interactions involving distinct CTCF binding sites within the domain
(**Figure 6B**), again underscoring chromatin conformation heterogeneity not captured by bulk
assays and a complexity of high-order interactions that might not be fully anticipated in SM
targeted designs.

Figure 7: **Cluster-based phasing of nano-NOMe-seq data reveals long range high-order multiway CTCF independent micro-compartments loops across the *Klf1* region. A:** RCMC (300 bp resolution), CTCF, SMC3, Pol II ChIP-seq and RNA-seq across the 2.3 Mb *Klf1* region in mESCs. The micro-compartment score (PC2 values from Hi-C eigenvector decomposition) highlights A-micro-compartments (red), which overlap Pol II peaks. Arcs depict CTCF-independent loops predicted by nucleosome occupancy at Pol II sites in each cluster. Red rectangles indicate anchors with predicted interactions in ≥ 5 out of 8 clusters. Clusters supporting the highest-order multiway interaction are also highlighted in red. **B**: Nano-NOMe-seq profiles of the 2 kb windows centered on Pol II summits overlapping A-micro-compartments involved in looping. **C**: ClinVar single-nucleotide variant (SNV) track in the syntenic human region, showing peaks of pathogenic SNVs overlapping chromatin interaction hubs that connect the center and boundaries of the region. (L)P (likely) Pathogenic; (L)B, (likely) begnin; VUS, variant of unknown significance; OTH, other.

While ORCA has identified the presence of rosettes within extended regions, it does not determine

whether these support or inhibit gene expression. In contrast, our nano-NOMe-seq cluster-based

approach has the potential to predict within-sample looping associated with Pol II activity. The Pol

II activation status across the *Hoxa* region in each rosette-forming cluster, demonstrates that

some rosette structures are compatible with activity across the entire region, while others are

linked to Pol II pausing and inactivity at particular sites (**Figure 6C**). As a general rule, we observe

that higher connectivity across the *Hoxa* region is associated with higher Pol II activity.

Cluster-based phasing of nano-NOMe-seq data reveals long-range high-order multiway

CTCF independent micro-compartments loops across the *Klf1* region

We next turned our attention to the *Klf1* region, which RCMC identified as being part of a

microcompartment with multiway interactions (**Figure 7A** and **Table S2**). Although

microcompartments are not (at least in the short term) reliant on either loop extrusion or

transcription, contacts largely occurred between active enhancers and promoters that are

enriched for Pol II, as highlighted by RCMC (**Figure 7A**) and Pol II peaks were overlapping A-

micro-compartments (red) as shown by the micro-compartment score (PC2 values from Hi-C

eigenvector decomposition).

Using our cluster-based nano-NOMe-seq we analyzed the nucleosome and Pol II status across

the region, centering our analysis on Pol II peaks. We assumed that TSSs exhibiting co-occurring

active Pol II-associated nucleosome patterns likely represent co-activated genes engaged in

multiway interactions across A micro-compartments identified by RCMC [8]. Across 8 windows, we

identified clusters with distinct combinations of elongating, paused or inactive Pol II sites (**Figure**

7B and **S4**). Clusters containing multiple simultaneously elongating Pol II sites, predicted by

nucleosome occupancy, are shown in **Figure 7B**. Arcs indicate the connections between

simultaneously active sites in each cluster, which are highlighted as CTCF-independent contacts

in the RCMC map. For visualization, only consecutive predicted interactions are displayed.

Although we cannot exclude the existence of additional pairwise combinations within a given

cluster, no cluster exhibited a single loop spanning the whole 220 kb region. Cluster 1,

representing only 10 % of the SM population, was the only case with a single loop spanning 115

kb, consistent with the reported maximum loop length achievable without CTCF-mediated loop

extrusion [28,29].   Red rectangles in **Figure 7B** indicate windows with coordinated activity in ≥ 5 out

of 8 clusters. The coalescence of active sites in these regions suggests coordinated multiway

interactions forming activity hubs at three locations: a central region (window 5) bridging two edge

regions located beyond the distance where chromatin interactions typically form without CTCF.
These hubs encompass genes including *Klf1, RNaseH, Jun2, Tnp2* and *Dhps,* suggesting
coordinated regulation of these genes.
298
299 Since many of the genes in these activity hubs are important for function and survival, we turned
300 to the ClinVar single-nucleotide variant (SNV) track in the syntenic human region. Intriguingly,
301 peaks of pathogenic SNVs overlap all three major activity hubs, connecting the center and
302 boundaries of the region, highlighting the relevance of the hubs.
303

DISCUSSION:
Despite advances in long-read sequencing, current technologies such as nanopore remain
insufficient to generate ultra-long reads that span entire TADs at the depth required for
comprehensive analysis. We generated a single-molecule, cluster-based phasing strategy for
nano-NOMe-seq (analogous to pseudobulk in single-cell data) that resolves heterogeneous
chromatin states and their coordination with transcription and long-range CTCF/cohesin
architecture. Stitched single-molecule clustering of nano-NOMe-seq delivered near single-cell
resolution to: (i) map how discrete Pol II states (elongating, active, paused, inactive) align with
CTCF footprints at defined loop anchors—explicitly revealing when dual-anchor occupancy is
compatible or incompatible with elongation; (ii) detect simultaneous multi-anchor CTCF
occupancy consistent with cohesin-mediated rosette topologies, while on the same molecules,
identifying which conformations are permissive for Pol II activation; and (iii) identify co-elongating
loci within microcompartments on single molecules, exposing recurrent, coordinated activity hubs.
PCR-free nano-NOMe preserves molecule counts, so cluster proportions approximate within-
sample allele/cell fractions. This framework not only reproduces prior findings but also adds
mechanistic resolution into how topology partitions transcriptional states across complex
regulatory domains.
321

By integrating cluster-level CTCF footprint and nucleosome patterns with ChIP-seq and chromatin
capture data, we could predict cluster-specific chromatin interactions and transcriptional activity
that were validated at the bulk level, thereby deconvolving distinct chromatin conformations that
enforce distinct transcriptional outcomes. The graded spectrum of CTCF footprints (unbound →
transitioning → strong NFR-embedded) likely reflects differences in residence time and cohesin-
stopping efficiency, providing a mechanistic framework for interpreting how anchor occupancy
shapes transcriptional compatibility. At the *Sox2* locus, for example, strong simultaneous CTCF

footprints at the promoter-proximal anchor and the SCR-embedded SRR109 anchor (window 5)

were selectively enriched on paused molecules but absent in elongating clusters, indicating that

the 1↔5 configuration is not compatible with elongation. In contrast, elongation was observed

when SRR109 (window 5) lacked a CTCF footprint, or when the promoter-proximal anchor paired

with the further distal downstream anchors (windows 9/10). These observations reconcile why

deleting the SRR109 motif does not affect steady-state Sox2 expression as enhancer activity

drives transcription when the 1↔5 loop cannot form. In mESCs, deletion of CTCF at the promoter-

proximal anchor has been shown to have no effect on Sox2 expression [26], indicating that the

longer distance multiway loops that contribute to rosette formation are dispensable for SCR-

mediated transcription in ESCs. However, they may be important for Sox2 gene regulation in other

contexts. Consistent with this, the enrichment of paused Pol II in the 1↔5 configuration suggests

a dynamic poised state in which Pol II remains primed for rapid activation or repression. Such

poised states are frequently involved the regulation of developmental genes [30]. In this view, the

1↔5 loop may not simply be an architectural feature, but a regulatory checkpoint that modulates

transcriptional responsiveness—stabilizing a paused Pol II complex until appropriate contextual

signals trigger either productive elongation or silencing. Importantly, even when bulk contact

frequencies appear stable, transcriptional changes may reflect a redistribution of single-molecule

state ensembles (e.g., looped-paused vs unlooped-elongating) rather than loss of specific

contacts. This highlights how specific CTCF-mediated topologies can gate transcriptional

plasticity, linking three-dimensional genome organization with dynamic gene expression.

Cluster-based phasing of SMs across hundreds of kilobases enabled us to deconvolve bulk

pairwise chromatin interactions into a mixture of two-way and multiway interactions consistent

with 'rosette' architectures, previously described at the *Sox2* and *Hoxa* domains using SM imaging

[17]. However, the non-targeted design of NOMe-seq data uncovered an unexpectedly rich

spectrum of cluster-specific interaction patterns that extended beyond the originally queried

regions, allowing unbiased discovery of higher-order conformations and revealing a level of

complexity not fully appreciated in earlier assays. Crucially, this molecule-resolved view separates

structural compatibility from transcriptional activity. Unlike long-read chromatin capture methods

such as Pore-C, which identify multiway interactions at SM resolution [31], but cannot resolve their

impact on transcription, our approach couples 3D architecture with Pol II and nucleosome status,

thereby linking specific conformations to their functional output. For example, we showed that

Hoxa clusters with higher predicted connectivity were more often permissive for polymerase

engagement across the region, but distinct rosette ensembles were compatible with the activation

of different gene combinations. Together, these results refine the rosette concept: rather than a

single hub, rosettes constitute ensembles of loop-competent configurations, distinguished by

which anchors are simultaneously involved. These distinct arrangements bias promoter states

toward pausing or activation, depending on local regulatory grammar, anchor strength, and

enhancer wiring.

Our nucleosome occupancy-based approach is not limited to predicting CTCF-mediated multiway

loops. At the *Klf1* locus, for example, microcompartment organization has previously been

identified using bulk RCMC. Although such microcompartment contacts persist after acute

inhibition of transcription, our single-molecule data show that their most active instances

correspond to concurrent elongation constellations across promoters and enhancers—a

coordination uniquely captured by nano-NOMe. Similar to the rosette structures, stitched clusters

revealed synchronous elongation across multiple Pol II peaks on the same molecules, generating

an even more diverse repertoire of multiway combinations. Despite this complexity, three

recurrent activity hubs emerged, coinciding with A-compartment foci. The central hub bridged

distal regions that, in the absence of this configuration, would otherwise require CTCF-mediated

looping to interact. These findings suggest that such hubs encompass genes requiring

coordinated regulation. Supporting this hypothesis, these activity regions overlap with peaks of

pathogenic human SNVs in the syntenic interval, suggesting that coordinated microcompartment

organization may mark regulatory hotspots particularly vulnerable to disease-associated

variation.

As with any predictive framework, our conclusions are restricted by the indirect nature of

accessibility footprints, inference of Pol II states, coverage depth, and gaps in GpC density, which

together constrain resolution and sensitivity; nevertheless, our approach already overcomes key

limitations of existing methods by linking chromatin architecture to transcriptional outcomes at

near single-molecule resolution, and future methodological advances will only expand its power.

In conclusion, cluster-based single-molecule phasing from nano-NOMe-seq data reveals that

regulatory loci exist as ensembles of loop-competent and transcriptionally distinct states whose

proportions—rather than population averages—govern output. Our framework bridges the gap

between contact maps and functional engagement and provides a roadmap for perturbations that

shift state distributions to modulate gene expression.

MATERIALS AND METHODS

Datasets and Data Processing

Nano-NOMe-seq

Nano-NOMe-seq data (GSE280711) were previously generated in mESCs and processed as described in Do *et al*.,[22]. Briefly, base modification calling and alignment against the mm10 reference genome were performed using Dorado. GpC methylation of single molecules (SMs) mapping to the regions of interest was extracted using modkit extract.

Hi-C

Hi-C data (GSE270669) were previously generated using the Arima Hi-C kit (A510008), aligned to mm10, and processed with HiC-Pro and HiCExplorer, as described in Do *et al*., [27]. Hi-C matrices were generated at 10 kb resolution.

Capture Micro-C

Capture Micro-C interaction matrices (50 bp resolution) aligned to mm39 were downloaded from Goel *et al*., [8] (GSE207225). The 2 replicates were merged, zoomified and balanced using cooler.

ChIP-seq, RNA-seq, and ATAC-seq

CTCF and SMC3 ChIP-seq, RNA-seq, and ATAC-seq data (GSE270669) were previously generated in mESCs, aligned to mm10, and processed as described in Do *et al*., [27], using the Seq-N-Slide pipeline. Pol II ChIP-seq data aligned to mm39 were downloaded from Goel *et al*., [8] (GSE207225), and subsequently realigned to mm10 using the Seq-N-Slide pipeline.

Visualization

Genomic maps were generated using HiGlass and IGV. For the *Sox2* and *Klf1* regions, ChIP-seq and RNA-seq data were lifted over to mm39 to ensure consistency with the Capture Micro-C data. For the *Klf1* locus, the micro-compartment score was defined as the second component (PC2) eigenvector values derived from eigenvector decomposition of the Hi-C correlation matrix and was generated using hicPCA (HiCExplorer). For the *Sox2* and *Hoxa1* loci, coordinates of anchors of multiway interactions identified by ORCA were obtained from *Hafner et al*., [1].

Nano-NOMe-Seq clustered-based phasing

Formatting and Filtering

Preprocessed files were imported and formatted to group GpC methylation calls by read identifier.

For each GpC site, the mean coverage was calculated. Reads were then filtered to exclude low-

quality or uninformative molecules based on length (<10 bp) and the number of methylated sites

retained (<10 GpCs).

Binning

Methylation sequences were partitioned into 50 bp bins spanning from the first read start to the

last read end. Smoothed methylation matrices were generated across fixed bins. *These matrices*

enabled visualization of methylation density profiles and identification of fully methylated versus

fully unmethylated reads. Because those reads were less informative to assess footprinting and

can be ambiguous for stitching, they were pre-assigned to a fully methylated and unmethylated

cluster, respectively and ignored for downstream analyses.

Window Separation

To enable localized clustering, sliding windows were defined across the genomic bins, and valid

methylation sequences were extracted for each window. Windows were defined as 60 valid bins

(valid bin were defined as bin with at least one read with methylation data) with at least 20

overlapping bins. Reads with less than 30 valid bins were excluded.

Window-Based Clustering

Within each window, pairwise similarity between methylation sequences was computed using a

masked Hamming distance metric that ignores missing values. Agglomerative hierarchical

Clustering was then performed within windows, ensuring rare cell states could be captured even

if not present across the full region. This algorithm separates the reads within each window into *t*

clusters based on the similarity of their methylation patterns: each data point is first treated as an

individual cluster. At each iteration, the two closest clusters are merged based on the average

linkage **(**average of all pairwise distances between points in the two clusters). This process

continues until the desired number of clusters, *t*, is reached. The agglomerative hierarchical

clustering procedure is described in detail below:

Let W *d*enote a window of the dataset containing n reads, with corresponding methylation

sequence vectors $\{x_1, x_2, ..., x_n\}$.

The clustering algorithm input is a distance matrix D, where each element $d_H(u, v)$ represents the

masked Hamming distance between two methylation sequence vectors u and v. The distance is

defined as:

465
$$d_H(u,v) = \begin{cases} 0 & if\ \sum m_i = 0 \\ \dfrac{(\sum m_i \cdot |u_i - v_i|)}{\sum m_i} & otherwise \end{cases}$$

466   where $m_i = 1$ if both u_i and v_i are observed (non-missing) and $m_i = 0$ otherwise.

467   Step 1. Initialization: Start with the n reads in the window, each considered as its own cluster.

468   Step 2. Define cluster distances (average linkage): The distance between two clusters $\boldsymbol{C_p}$ and $\boldsymbol{C_q}$

469   is defined as the average of all pairwise distances between points in $\boldsymbol{C_p}$ and points in $\boldsymbol{C_q}$:

470
$$d(C_p, C_q) = \frac{1}{|C_p| \cdot |C_q|} \sum_{u \in C_p} \sum_{v \in C_q} d_H(u, v)$$

471   where $|C_p|$ denotes the number of reads in cluster $\boldsymbol{C_p}$.

472   Step 3. Compute all inter-cluster distances: At this stage, compute the complete distance matrix

473   between every pair of clusters.

474   Step 4. Merge closest clusters: Identify the pair of clusters (C_r, C_s) such that

475
$$(C_r, C_s) = \arg \min_{p \neq q} d\left(C_p, C_q\right)$$

476   Merge these two clusters into a new cluster $C_{rs} = C_r \cup C_s$

477   Step 5. Update the distance matrix: Remove the rows and columns corresponding to C_r and C_s.

478   Add a row/column for the new cluster C_{rs}. Compute distances between C_{rs} and each remaining

479   cluster C_k using the average linkage formula.

480
$$d(C_{rs}, C_k) = \frac{|C_r| \cdot d(C_r, C_k) + |C_s| \cdot d(C_s, C_k)}{|C_r| + |C_s|}$$

Step 6. Repeat until target cluster number is reached: Continue iteratively merging clusters until

the number of clusters equals the predefined parameter t.

Cluster phasing/stitching

After per-window clustering, each of the M windows yields t clusters. To generate t consistent

clusters across the entire region, clusters must be stitched between successive windows. This

task is an assignment problem: matching clusters in window N to the most similar clusters in

window $N + 1$. The problem can be expressed as a mathematical optimization:

Let the windows W_N and W_{N+1}, two successive windows each containing t clusters

$$c_1^N, \ c_2^N, \ \dots, \ c_t^N \text{ and } c_1^{N+1}, \ c_2^{N+1}, \ \dots, \ c_t^{N+1}.$$

Given a distance matrix $D = [d_{ij}]$ with $i, j = 1, \dots, n$ where d_{ij} is the distance between cluster

c_i^N and c_j^{N+1} , the assignment problem is to find a binary matrix $\mathbf{X} = [x_{ij}]$ where each entry x_{ij}

indicates whether cluster c_i^N is matched to cluster c_j^{N+1} ($x_{ij} = 1$) or not ($x_{ij} = 0$) and which

minimizes the total cost:

496
$$\min_{X \in \{0,1\}^{t \times t}} \sum_{i=1}^{n} \sum_{j=1}^{n} d_{ij} x_{ij}$$

497　Subject to:

498
$$\sum_{j=1}^{n} x_{ij} = 1 \ \forall i, \ \sum_{i=1}^{n} x_{ij} = 1 \ \forall j,$$

499　so that every cluster in window N is matched uniquely to one cluster in window $N + 1$

500　To solve this problem, we used the Hungarian algorithm, a classical method for optimal

501　assignment in combinatorial optimization, with as input the distance matrix

503
$$\mathbf{D} = \{d_H(\mu_i^{(N)}, \mu_j^{(N+1)})\}_{i,j \in [\![1,n]\!]^2} \, ,$$

where $d_H(u, v)$ is the Masked Hamming distance between two vectors u and v and $\mu_i^{(N)}$ is the

mean methylation sequence vector of cluster c_i^N

$$\mu_i^{(N)} = \frac{1}{|C_i^{(N)}|} \sum_{r \in C_i^{(N)}} \mathbf{m_r},$$ where $\mathbf{m_r}$ is the methylation sequence vector of read r

and the following procedure:

Step 1. Row reduction: For each row in the cost matrix **D**, subtract the minimum value of that row

from all elements in the row.

Step 2. Column reduction: For each column, subtract the minimum value of that column from all

elements in the column.

Step 3. Cover zeros: Cover all zeros with the minimum number of horizontal/vertical lines. If the

number of lines = t, proceed to Step 5, otherwise, continue to Step 4.

Step 4. Adjust uncovered elements: Find the smallest uncovered value δ (i.e., not crossed by any

line), subtract δ from all uncovered elements, add δ to elements covered twice (i.e., by both a row

and column) then repeat Step 3.

Step 5. Find an optimal assignment: Find a zero in the matrix, assign it, and eliminate its row and

column from further assignment. Continue until every row has a unique assignment. This gives

you the optimal assignment minimizing total cost.

Repeating the procedure for each pair of adjacent windows yields an optimal one-to-one matching

of clusters across all M windows. This ensures that each of the t stitched clusters is continuous

and consistent throughout the region.

Global Clustering

Each read was assigned to its most similar stitched cluster series based on masked Hamming

distance. Final clusters thus represented groups of single-molecule methylation profiles spanning

the entire locus. Cluster assignments were assessed by plotting individual reads alongside their

corresponding series profiles at the bin and GpC level.

Long read accuracy

To evaluate the accuracy of our clustering approach, we selected 100 loci each covered by at

least one long read (>50 kb). After filtering for informative reads, 90 loci with sufficient coverage

were retained (**Table S1**). For each locus, the cluster-based stitching procedure was performed

while leaving out the long read. After clustering, the long read was assigned to its best-matched

global cluster. For each locus, the methylation sequence of the long read was compared to the

mean methylation sequence of each cluster (c_i). The closest cluster was identified using the

masked Hamming distance d_H.

The read accuracy for a long read r_i was defined as:

542
$$A(r_i) = 1 - \min_{c_i \in Clusters} d_H(r_i, c_i)$$

544 The overall clustering accuracy was defined as the average accuracy across all long reads:

546
$$Accuracy = \frac{1}{N} \sum_{r_i \in Long\ reads} A(r_i)$$

Data availability

No data were generated in this study. Accession numbers of the datasets used are reported in

the datasets section.

Code availability

The codes for the clustered-based phasing strategy will be made publicly available on zenodo

upon acceptance (https://doi.org/10.5281/zenodo.17064564).

Acknowledgements

This work was supported by 1R35GM122515 (J.S) and NIH P01CA229086 (J.S).

The authors thank Skok lab members for helpful scientific discussions, New York University

School of Medicine High Performance Computing Facility (HPCF) for computing technical

support, Adriana Heguy and the Genome Technology Center (GTC) core for sequencing efforts.

GTC is a shared resource partially supported by the Cancer Center Support Grant P30CA016087

at the Laura and Isaac Perlmutter Cancer Center.

Author contributions

These studies were designed by Jane Skok, Catherine Do and Stephanie Bellini. The analysis

was performed by Stephanie Bellini and Catherine Do. The paper was written by Jane Skok and

Catherine Do.

Declaration of interests

The authors declare no competing interests.

SUPPLEMENTARY INFORMATION

Supplemental Tables

Table S1. Cluster-based phasing accuracy. Table of long reads used for the accuracy
estimation. This table relates to **Figure 3**.

Table S2. Coverage and read length median for Sox2, Hoxa and Klf1 loci. This table relates
to **Figures 4-7**.

Supplemental Figures

Figure S1. Clustered-based phasing of the Nano-NOMe-seq single molecules. Cartoon
depicting the agglomerative hierarchical clustering (**A**), the Hungarian algorithm (**B**) and (**C**) the
Hamming distance used in the clustered based phasing pipeline. This figure relates to **Figure 2**.

Figure S2. Cluster-based phasing of nano-NOMe-seq data reveals CTCF-independent Sox2
transcription. Profiles of mean methylation across all 9 clusters (excluding the fully methylated
and unmethylated clusters, C1 and C9) and studied windows shown in **Figure 4A**. This figure
relates to **Figure 4**.

Figure S3. Cluster-based phasing of nano-NOMe-seq data reveals long range multiway
CTCF dependent loops at the *Sox2* locus. A. Profiles of mean methylation across all 7 clusters
(excluding the fully methylated and unmethylated clusters, C1 and C7) and studied windows
shown in **Figure 5A**. This figure relates to **Figure 5**.

Figure S4. Cluster-based phasing of nano-NOMe-seq data reveals long range high-order
multiway CTCF independent loops between micro-compartments at the *Klf1* locus. A.
Profiles of mean methylation across all 8 clusters (excluding the fully methylated and
unmethylated clusters, C8 and C9) and studied windows shown in **Figure 7A**. This figure relates
to **Figure 7**.

Figure S1. Clustered-based phasing of the Nano-NOMe-seq single molecules. Cartoon depicting the agglomerative hierarchical clustering (**A**), the Hungarian algorithm (**B**) and (**C**) the Hamming distance used in the clustered based phasing pipeline. This figure relates to **Figure 2**.

604
605

Figure S2. Cluster-based phasing of nano-NOMe-seq data reveals CTCF-independent Sox2 transcription. Profiles of mean methylation across all 9 clusters (excluding the fully methylated and unmethylated clusters, C1 and C9) and studied windows shown in **Figure 4A**. This figure relates to **Figure 4**.

606
607

Figure S3. Cluster-based phasing of nano-NOMe-seq data reveals long range multiway CTCF dependent loops at the *Sox2* locus. A. Profiles of mean methylation across all 7 clusters (excluding the fully methylated and unmethylated clusters, C1 and C7) and studied windows shown in **Figure 5A**. This figure relates to **Figure 5**.

609
610

Figure S4. Cluster-based phasing of nano-NOMe-seq data reveals long range high-order multiway CTCF independent loops between micro-compartments at the *Klf1* locus. A. Profiles of mean methylation across all 8 clusters (excluding the fully methylated and unmethylated clusters, C8 and C9) and studied windows shown in **Figure 7A**. This figure relates to **Figure 7**.

611
612

REFERENCES:

1. Hafner, A., Park, M., Berger, S.E., Murphy, S.E., Nora, E.P., and Boettiger, A.N. (2023). Loop stacking organizes genome folding from TADs to chromosomes. Mol Cell *83*, 1377-1392 e1376. 10.1016/j.molcel.2023.04.008.

2. Ostrowski, M.S., Yang, M.G., McNally, C.P., Abdulhay, N.J., Wang, S., Renduchintala, K., Irkliyenko, I., Biran, A., Chew, B.T.L., Midha, A.D., et al. (2025). The single-molecule accessibility landscape of newly replicated mammalian chromatin. Cell *188*, 237-252 e219. 10.1016/j.cell.2024.10.039.

3. Fudenberg, G., Imakaev, M., Lu, C., Goloborodko, A., Abdennur, N., and Mirny, L.A. (2016). Formation of Chromosomal Domains by Loop Extrusion. Cell Rep *15*, 2038-2049. 10.1016/j.celrep.2016.04.085.

4. Dekker, J., Rippe, K., Dekker, M., and Kleckner, N. (2002). Capturing chromosome conformation. Science *295*, 1306-1311.

5. Lieberman-Aiden, E., van Berkum, N.L., Williams, L., Imakaev, M., Ragoczy, T., Telling, A., Amit, I., Lajoie, B.R., Sabo, P.J., Dorschner, M.O., et al. (2009). Comprehensive mapping of long-range interactions reveals folding principles of the human genome. Science *326*, 289-293. 326/5950/289 [pii]. 10.1126/science.1181369.

6. Krietenstein, N., Abraham, S., Venev, S.V., Abdennur, N., Gibcus, J., Hsieh, T.S., Parsi, K.M., Yang, L., Maehr, R., Mirny, L.A., et al. (2020). Ultrastructural Details of Mammalian Chromosome Architecture. Mol Cell *78*, 554-565 e557. 10.1016/j.molcel.2020.03.003.

7. Bonev, B., and Cavalli, G. (2016). Organization and function of the 3D genome. Nat Rev Genet *17*, 661-678. 10.1038/nrg.2016.112.

8. Goel, V.Y., Huseyin, M.K., and Hansen, A.S. (2023). Region Capture Micro-C reveals coalescence of enhancers and promoters into nested microcompartments. Nat Genet *55*, 1048-1056. 10.1038/s41588-023-01391-1.

9. Zhang, S., Ubelmesser, N., Barbieri, M., and Papantonis, A. (2023). Enhancer-promoter contact formation requires RNAPII and antagonizes loop extrusion. Nat Genet *55*, 832-840. 10.1038/s41588-023-01364-4.

10. Chang, L., Xie, Y., Taylor, B., Wang, Z., Sun, J., Armand, E.J., Mishra, S., Xu, J., Tastemel, M., Lie, A., et al. (2024). Droplet Hi-C enables scalable, single-cell profiling of chromatin architecture in heterogeneous tissues. Nat Biotechnol. 10.1038/s41587-024-02447-1.

11.   Fu, Y., Rocha, P.P., Luo, V.M., Raviram, R., Deng, Y., Mazzoni, E.O., and Skok, J.A.
(2016). CRISPR-dCas9 and sgRNA scaffolds enable dual-colour live imaging of satellite
sequences and repeat-enriched individual loci. Nat Commun *7*. 10.1038/ncomms11707.

12.   Ma, H., Tu, L.C., Naseri, A., Huisman, M., Zhang, S., Grunwald, D., and Pederson, T.
(2016). Multiplexed labeling of genomic loci with dCas9 and engineered sgRNAs using
CRISPRainbow. Nat Biotechnol *34*, 528-530. 10.1038/nbt.3526.

13.   Bertrand, E., Chartrand, P., Schaefer, M., Shenoy, S.M., Singer, R.H., and Long, R.M.
(1998). Localization of ASH1 mRNA particles in living yeast. Mol Cell *2*, 437-445.
10.1016/s1097-2765(00)80143-4.

14.   Darzacq, X., Shav-Tal, Y., de Turris, V., Brody, Y., Shenoy, S.M., Phair, R.D., and Singer,
R.H. (2007). In vivo dynamics of RNA polymerase II transcription. Nat Struct Mol Biol *14*,
796-806. 10.1038/nsmb1280.

15.   Alexander, J.M., Guan, J., Li, B., Maliskova, L., Song, M., Shen, Y., Huang, B., Lomvardas,
S., and Weiner, O.D. (2019). Live-cell imaging reveals enhancer-dependent Sox2
transcription in the absence of enhancer proximity. Elife *8*. 10.7554/eLife.41769.

16.   Wan, X., Kong, J., Hu, X., Liu, L., Yang, Y., Li, H., Liu, G., Niu, X., Chen, F., Zhang, D., et
al. (2025). SiCLAT: simultaneous imaging of chromatin loops and active transcription in
living cells. Genome Biol *26*, 1. 10.1186/s13059-024-03463-9.

17.   Mateo, L.J., Murphy, S.E., Hafner, A., Cinquini, I.S., Walker, C.A., and Boettiger, A.N.
(2019). Visualizing DNA folding and RNA in embryos at single-cell resolution. Nature *568*,
49-54. 10.1038/s41586-019-1035-4.

18.   Espinola, S.M., Gotz, M., Bellec, M., Messina, O., Fiche, J.B., Houbron, C., Dejean, M.,
Reim, I., Cardozo Gizzi, A.M., Lagha, M., and Nollmann, M. (2021). Cis-regulatory
chromatin loops arise before TADs and gene activation, and are independent of cell fate
during early Drosophila development. Nat Genet *53*, 477-486. 10.1038/s41588-021-
00816-z.

19.   Cardozo Gizzi, A.M., Cattoni, D.I., Fiche, J.B., Espinola, S.M., Gurgo, J., Messina, O.,
Houbron, C., Ogiyama, Y., Papadopoulos, G.L., Cavalli, G., et al. (2019). Microscopy-
Based Chromosome Conformation Capture Enables Simultaneous Visualization of
Genome Organization and Transcription in Intact Organisms. Mol Cell *74*, 212-222 e215.
10.1016/j.molcel.2019.01.011.

20.   Liu, Z., Chen, Y., Xia, Q., Liu, M., Xu, H., Chi, Y., Deng, Y., and Xing, D. (2023). Linking
genome structures to functions by simultaneous single-cell Hi-C and RNA-seq. Science
380, 1070-1076. 10.1126/science.adg3797.

21.   Zhou, T., Zhang, R., Jia, D., Doty, R.T., Munday, A.D., Gao, D., Xin, L., Abkowitz, J.L.,
680   Duan, Z., and Ma, J. (2024). GAGE-seq concurrently profiles multiscale 3D genome
681   organization and gene expression in single cells. Nat Genet *56*, 1701-1711.
682   10.1038/s41588-024-01745-3.

683   22.   Catherine Do, G.J., Paul Zappile, Adriana Heguy, Jane A. Skok (2025). The coordination
684   between CTCF, cohesin and TFs impacts nucleosome repositioning and chromatin
685   insulation to define state specific 3D chromatin folding. Biorxiv.

686   23.   Zhou, H.Y., Katsman, Y., Dhaliwal, N.K., Davidson, S., Macpherson, N.N., Sakthidevi, M.,
687   Collura, F., and Mitchell, J.A. (2014). A Sox2 distal enhancer cluster regulates embryonic
688   stem cell differentiation potential. Genes Dev *28*, 2699-2711. 10.1101/gad.248526.114.

689   24.   de Wit, E., Vos, E.S., Holwerda, S.J., Valdes-Quezada, C., Verstegen, M.J., Teunissen,
690   H., Splinter, E., Wijchers, P.J., Krijger, P.H., and de Laat, W. (2015). CTCF Binding Polarity
691   Determines Chromatin Looping. Mol Cell *60*, 676-684. 10.1016/j.molcel.2015.09.023.

692   25.   Taylor, T., Sikorska, N., Shchuka, V.M., Chahar, S., Ji, C., Macpherson, N.N., Moorthy,
693   S.D., de Kort, M.A.C., Mullany, S., Khader, N., et al. (2022). Transcriptional regulation and
694   chromatin architecture maintenance are decoupled functions at the Sox2 locus. Genes
695   Dev *36*, 699-717. 10.1101/gad.349489.122.

696   26.   Chakraborty, S., Kopitchinski, N., Zuo, Z., Eraso, A., Awasthi, P., Chari, R., Mitra, A.,
697   Tobias, I.C., Moorthy, S.D., Dale, R.K., et al. (2023). Enhancer-promoter interactions can
698   bypass CTCF-mediated boundaries and contribute to phenotypic robustness. Nat Genet
699   *55*, 280-290. 10.1038/s41588-022-01295-6.

700   27.   Do, C., Jiang, G., Cova, G., Katsifis, C.C., Narducci, D.N., Sakellaropoulos, T., Vidal, R.,
701   Lhoumaud, P., Tsirigos, A., Regis, F.F.D., et al. (2025). Binding domain mutations provide
702   insight into CTCF's relationship with chromatin and its contribution to gene regulation. Cell
703   Genom *5*, 100813. 10.1016/j.xgen.2025.100813.

704   28.   Uyehara, C.M., and Apostolou, E. (2023). 3D enhancer-promoter interactions and multi-
705   connected hubs: Organizational principles and functional roles. Cell Rep, 112068.
706   10.1016/j.celrep.2023.112068.

707   29.   Hsieh, T.S., Cattoglio, C., Slobodyanyuk, E., Hansen, A.S., Darzacq, X., and Tjian, R.
708   (2022). Enhancer-promoter interactions and transcription are largely maintained upon
709   acute loss of CTCF, cohesin, WAPL or YY1. Nat Genet *54*, 1919-1932. 10.1038/s41588-
710   022-01223-8.

711   30.   Core, L., and Adelman, K. (2019). Promoter-proximal pausing of RNA polymerase II: a
712   nexus of gene regulation. Genes Dev *33*, 960-982. 10.1101/gad.325142.119.

713   31.   Deshpande, A.S., Ulahannan, N., Pendleton, M., Dai, X., Ly, L., Behr, J.M., Schwenk, S.,
714   Liao, W., Augello, M.A., Tyer, C., et al. (2022). Identifying synergistic high-order 3D
715   chromatin conformations from genome-scale nanopore concatemer sequencing. Nat
716   Biotechnol *40*, 1488-1499. 10.1038/s41587-022-01289-z.

717
718