Mitochondria-Lysosome Crosstalk Shapes Metabolic Transition in Neonatal Enterocytes

Gonzalo Herranz,[1,2] Diego Alonso-Larre,[1,2,10] Tamara González,[1,2,10] Laura Akintche,[1,2] Alejandra
Ramos-Manzano,[1,2,7] Marta Iborra-Pernichi,[1,2,8], María Velasco de la Esperanza[1,2], Covadonga
Díaz-Díaz,[1,2,9] Ian G Ganley,[3] Patricia Boya,[4,5] Sara Cogliati,[6] Nuria Martínez-Martín,[1,2]* and
Fernando Martín-Belmonte[1,2,11]**
[1]Program of Tissue and Organ Homeostasis, Centro de Biología Molecular Severo Ochoa, Consejo Superior de
Investigaciones Científicas-Universidad Autónoma de Madrid, Madrid, Spain
[2]Intestinal Morphogenesis and Homeostasis Group, Área 3-Cancer, Instituto Ramón y Cajal de Investigación Sanitaria
(IRYCIS), Madrid, Spain
[3]MRC Protein Phosphorylation and Ubiquitylation Unit, University of Dundee, Dundee, UK
[4]Dept of Cellular and Molecular Biology, Centro de Investigaciones Biológicas Margarita Salas, CSIC, Madrid, Spain
[5]Department of Neuroscience and Movement Science, Section of Medicine, University of Fribourg, Fribourg,
Switzerland
[6]Program of Physiological and Pathological Processes, Centro de Biología Molecular Severo Ochoa, Consejo Superior
de Investigaciones Científicas-Universidad Autónoma de Madrid, Madrid, Spain
[7]Present addresses: Immunology Department, Hospital Universitario de La Princesa, Instituto de Investigación Sanitaria (IIS)-
Princesa, Madrid, Spain; Medicine Department, School of Medicine, Universidad Autónoma of Madrid, Madrid, Spain
[8]Present address: B Lymphocyte Biology Lab, Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid, Spain
[9]Present addresses: Cardiovascular Regeneration Program, Centro Nacional de Investigaciones Cardiovasculares (CNIC), Madrid,
Spain; Centro de Investigación Biomédica en Red de Enfermedades Cardiovasculares (CIBERCV), Madrid, Spain
[10]These authors contributed equally
[11]Senior author and Lead contact
*Correspondence: nmartinez@cbm.csic.es
**Correspondence: fmartin@cbm.csic.es
24

SUMMARY

The neonatal gastrointestinal tract mediates nutrient absorption and the establishment of immune tolerance

to commensal microbiota. In early life, lysosome-rich enterocytes (LREs) in the ileum are necessary for the

intracellular digestion of maternal milk proteins. However, the molecular mechanisms sustaining their

function remain incompletely characterized. Here, we demonstrate that LRE mitochondrial homeostasis

and autophagic capacity are critical for efficient nutrient uptake and maintenance of their specialized

identity, as disruption of either process leads to premature differentiation into post-weaning enterocytes

(PECs) with diminished endolysosomal and metabolic activity. Transcriptomic profiling further revealed that

neonatal LREs exhibit a distinctive antioxidant signature, which preserves redox balance and safeguards

the expression of the transcriptional regulators MAFB and BLIMP1, both central repressors of the neonatal-

to-adult metabolic transition. These findings establish the mitochondria–lysosome axis as a key determinant

of LRE function and neonatal metabolic programming. They also provide a mechanistic framework for

understanding how organelle dysfunction and redox imbalance may contribute to early-life malnutrition

syndromes, such as Kwashiorkor, and suggest therapeutic strategies aimed at preserving mitochondrial

and lysosomal integrity.

KEYWORDS

Intestine, Enterocyte, Autophagy, Mitochondria, Lysosome, Neonatal, Metabolic Switch, ROS, Nutrient

Absorption, Mouse

INTRODUCTION

The small intestine is a specialized section of the gastrointestinal tract (GIT) engaged in nutrient digestion and absorption, while simultaneously acting as a barrier against harmful substances. It consists of a variety of cell types, with epithelial and immune cells playing key roles in maintaining intestinal function. Through their interactions with one another and the gut microbiota, these cells help create a finely balanced microenvironment that supports both nutritional and immunological functions (Beumer and Clevers, 2021; Choo et al., 2022). The main absorptive epithelial cells in the intestine are called enterocytes. They are the most numerous cells in the intestinal epithelium and, along with secretory epithelial cells (including Paneth, goblet, and enteroendocrine cells), originate from stem cells of the intestinal crypts (Beumer and Clevers, 2021).

Dietary protein is a macronutrient essential for growth and overall organismal homeostasis. In weaned mammals, protein digestion takes place in the lumen of GIT, where gastric pepsin and intestinal proteases sequentially break down proteins into oligopeptides and free amino acids, which are then absorbed by enterocytes via specific transporters (Henning, 1981, 1985; Rombout et al., 1985).

However, in the intestinal lumen of suckling mammals and stomachless fishes, protease activity remains low and protein digestion is incomplete (Henning, 1981, 1985; Rombout et al., 1985). For this reason, their posterior-mid small intestine contains a population of lysosome-rich enterocytes (LREs), formerly referred to as vacuolated enterocytes (Kleinman and Walker, 1984; Park et al., 2019; Rodríguez-Fraticelli et al., 2015). These cells possess a highly developed endolysosomal system, including a prominent lysosomal vacuole (LV), which is capable of internalising and metabolising intact proteins via non-specific fluid-phase processes (pinocytosis) and receptor-mediated endocytosis and membrane adsorption (Fujita et al., 1990; Gonnella and Neutra, 1984; Park et al., 2019; Rombout et al., 1985). Moreover, LREs facilitate the transcytosis of antigens, maternal immunoglobulins, and antibodies across the epithelial barrier while preserving their biological activity, contributing to the immune system development (Kulkarni and Newberry, 2019; Rombout et al., 1985). Recent studies have highlighted the nutritional importance of LREs during the neonatal period. Key endolysosomal proteins, whose absence leads to developmental defects, have been identified, including Mucolipins 1 and 3 (Remis et al., 2014), Pllp (Rodríguez-Fraticelli et al., 2015), Mamdc4 (also known as endotubin) (Cox et al., 2018) and Dab2 (Park et al., 2019). Despite these findings, the molecular mechanisms that regulate endocytic and lysosomal pathways in LREs are still not fully understood.

Autophagy is a conserved lysosome-dependent catabolic pathway that recycles intracellular components and nutrients. This process involves the sequestration of cytoplasmic material into double-membraned autophagosomes, which subsequently fuse with lysosomes for degradation and breakdown. The lipidation of LC3, a key step in autophagosome formation, is mediated by the ATG12–ATG5–ATG16L1 complex (He, 2022; Kaur and Debnath, 2015). Mutations in components of this complex have been linked to intestinal pathologies (Foerster et al., 2022; Larabi et al., 2020), and *Atg5*-deficient neonates exhibit increased

neonatal mortality under fasting conditions, underscoring the essential role of autophagy during early
postnatal life (Kuma et al., 2004). Despite this, the homeostatic role of autophagy throughout the entire
neonatal period, particularly in LRE biology, remains unclear.

Autophagy plays a crucial role in cellular metabolism and differentiation, primarily through the selective
elimination of mitochondria (mitophagy), thereby contributing to the control of reactive oxygen species
(ROS) levels (Wang et al., 2023; Zhong et al., 2024). Beyond their role in energy production, mitochondria
and ROS also function as key regulators of cell signalling and differentiation (Lennicke and Cochemé, 2021;
Morris and Jasper, 2021; Popov, 2023). In the intestinal epithelium, a gradient of increasing mitochondrial
mass, activity, and ROS levels has been described along the crypt–villus axis, correlating with higher
degrees of cellular differentiation (Rath et al., 2018). However, how mitochondrial homeostasis and ROS
dynamics influence the developmental remodelling of enterocytes remains poorly understood. In particular,
their role in regulating the metabolic and functional transition of neonatal LREs into post-weaning
enterocytes (PECs) has not yet been explored.

This study reveals that the mitochondria–lysosome axis is essential for the function and maturation of LREs.
We show that LREs rely on high autophagic and mitophagic activity to sustain nutrient absorption and
lysosomal integrity, and that disruption of either autophagy (*Atg16*cKO) or mitochondrial homeostasis
(*Tfam*cKO) leads to impaired endolysosomal function, oxidative stress, and premature weaning transition
through the repression of the transcription factors BLIMP1 and MAFB. These findings establish organelle
homeostasis as a critical determinant of neonatal enterocyte identity and provide mechanistic insights into
how its disruption contributes to early-life malnutrition.

RESULTS

Lysosome-rich enterocytes (LREs) display heightened basal autophagy levels essential for
absorptive function

Unlike PECs, LREs rely on intracellular digestion within lysosomes to process macromolecular nutrients.
This process demands a highly active endolysosomal system that breaks down complex proteins into
absorbable units, which confers a particular architecture to LREs (**Figures S1A and S1B**) (Fujita et al.,
1990). Autophagy, the cellular mechanism responsible for degrading and recycling cellular components,
plays a central role in maintaining lysosomal function and turnover (Yim and Mizushima, 2020). In LREs,
elevated basal autophagy may support endolysosomal activity. Exploring this hypothesis could clarify the
role of autophagy in the absorption and processing of nutrients in neonates, as well as its contribution to
early development.

Utilising the GFP-LC3 transgenic mouse model (*GFP-Map1lc3a*$^{TG/+}$) (Mizushima et al., 2004), we visualised
autophagosomes as GFP-LC3 positive puncta and monitored autophagy flux via western blot (WB) by
assessing the levels of free GFP, a byproduct of lysosomal degradation of GFP-LC3 II (**Figures 1A, 1B and**

S1C) (Mizushima et al., 2010; Ni et al., 2011). The presence and activity of LREs was determined by probing

with mCherry/ fluorescent dextrans that are internalised as a macromolecule by LREs, but not by PECs

(**Figures 1A, 1B, S1C, andS1D**) (Park et al., 2019). Following gavage, fluorescent-dextran was rapidly

internalised from the lumen into early and late endosomes (**Figure S1D**, arrowheads), accumulating

progressively in lysosomal vacuoles (LVs) (**Figure S1D**, arrows). WB analysis was performed on lysates

from epithelial cells in both proximal and distal sections of small intestines from GFP-LC3 mice at neonatal

stages (P1, P7 and P14), and post-weaning (P21–30). This analysis revealed a proximal-distal gradient of

protein absorption and autophagic degradation, mirroring the anterior-posterior intestinal maturation (**Figure**

1B) (Spence et al., 2011).

In the distal intestine, protein absorption preceded the induction of autophagy, with a subtle onset already

detectable at P1, although no free GFP signal was observed. By P7, both mCherry uptake and free GFP

increased markedly, with a strong accumulation of free GFP (**Figure 1B**, pink rectangle), indicating peak

autophagic activity in LREs—coinciding with the establishment of neonatal intestinal features (Hansson et

al., 2011). At P14, mCherry absorption persisted in the distal region, while free GFP levels declined, and

by weaning (P21) both signals were absent (**Figure 1B**). Quantification of mCherry and free GFP in ileal

epithelial lysates at P7 and P21 confirmed these developmental differences (**Figures 1C and S1C**).

In contrast, the proximal intestine displayed a different temporal profile. At P1, both mCherry and free GFP

were already present, but their levels progressively declined, and at P14 only low residual signals were

detectable. Note that undegraded GFP-LC3 forms (I and II) and intermediate degradation products

accumulated at P21, in both distal and proximal samples (**Figure 1B**).

Given these observations, we conclude that LREs maintain elevated autophagy levels relative to PECs,

most probably as a result of their enhanced lysosomal function. To explore the consequences of autophagy

disruption in these cells during early development, we utilised a genetic mouse model. ATG16L1 is crucial

for both autophagosome formation and LC3 lipidation, two processes essential for autolysosomal

degradation (Fujita et al., 2008; Mizushima et al., 2003). This gene has been extensively studied for its

association with inflammatory bowel diseases (Foerster et al., 2022; Larabi et al., 2020); nonetheless, its

specific role in LRE functionality during the neonatal period remains largely unexplored. For this, we crossed

with the *Atg16l1*$^{fl/fl}$ mice (Hwang et al., 2012), with mice expressing the *Villin-creER*^{T2} transgene (el Marjou

et al., 2004). This breeding strategy generated an *Atg16l1* knockout offspring in which the *Atg16l1* gene is

conditionally deleted in the intestinal epithelium upon administration of tamoxifen (*Atg16*cKO) (**Figure 1D**).

Initially, we validated our model via WB analysis by measuring the levels of ATG16L1 and LC3 isoforms

(cytoplasmic LC3-I and lipidated LC3-II) in the ileum of both WT and *Atg16*cKO mice at P14 (**Figures 1D**

and 1E). We observed a significant reduction in ATG16L1 protein levels across all *Atg16*cKO samples,

which led to a corresponding decrease in LC3-II/LC3-I ratio (**Figure 1D**). Specifically, our data showed an

84.5% reduction in ATG16L1 protein levels and an approximately 8-fold decrease in LC3-II/LC3-I ratio

(**Figure 1E**). Furthermore, we observed elevated levels of P62, indicating a reduction in its autophagy-

dependent degradation (**Figures 1D and 1E**). These outcomes are consistent with previous results (Cadwell
et al., 2008).

Subsequently, we examined the developmental trajectory of *Atg16*cKO mice, with a primary focus on their
intestinal morphology and function. We observed that the mice reached the post-weaning stage (P30)
without apparent issues, showing no weight defects throughout early postnatal development (**Figure S2A**).
No significant changes were noted in the length or histological state of the small and large intestines
(**Figures S2B and S2C**). However, a reduction in villus length and a decrease in the number of Goblet cells
(GCs) were detected in the ileum at P14 (**Figures 1F and 1G**), along with mild oedema at P30, an indicator
of inflammation (**Figure S2D**, arrowhead). Inspired by these observations, we quantified specific intestinal
epithelial (IEC) and immune cell subpopulations in the ileum of *Atg16*cKO mice at both neonatal and post-
weaning stages using flow cytometry (**Figures S2E and S2F**). For this, we applied specific markers to
identify various IEC subtypes, including enterocytes and their progenitors (EC), Goblet cells (GC), Paneth
cells (PC), and enteroendocrine cells (EEC) (Asano et al., 2017) (**Figure S3**, middle row). We also focused
on cells related to innate immunity (IIC), such as CD11b+ cells (including NK cells, granulocytes, and
macrophages/monocytes), as well as adaptive immunity, encompassing B lymphocytes (B cells) and CD4+
T helper cells (Th cells). We observed an increase in IIC and IEC populations during the neonatal period,
accompanied by a concurrent decrease in GC and EEC in adult IECs (**Figure S2E**). In contrast, post-
weaning samples showed a rise in B cells, stabilization in IIC and IEC levels, an increase in GC, and a
decrease in PC and EEC (**Figure S2F**). These results corroborate our histological findings regarding GCs
and inflammation.

Next, we investigated the effect of autophagy inhibition on the absorption of macromolecules in the ileum.
We gaveged rhodamine B–dextran (Rd-dextran) as a probe in *Atg16*cKO and WT mice as controls (**Figure
1H**). Our analysis differentiated between the apical and basal regions of enterocytes to assess the
sequential entry of Rd-dextran, according to the internalization dynamics observed in these cells (**Figure
S1D**). Confocal microscopy demonstrated that at P7, *Atg16*cKO enterocytes internalized Rd-dextran
similarly to WT counterparts. Nevertheless, it is worth noting that in *Atg16*cKO enterocytes, Rd-dextran
levels were slightly elevated in the apical region but reduced in the basal region compared to WT (**Figure
1H**). This observation suggests a deceleration in the internalisation of macromolecules during this
developmental period. Next, at P14, *Atg16*cKO enterocytes exhibited a complete abrogation of
macromolecule absorption, resembling post-weaning enterocytes and indicating an advancement in
intestinal maturation (**Figure 1H**). To further validate the observed decrease in absorption at P14, we probed
neonates from both WT and *Atg16*KO groups with mCherry and quantified the internalised mCherry in the
ileum via WB. The results confirmed consistently lower levels of nutrient absorption in *Atg16*KO mice
(**Figures 1I and 1J**). Altogether, we conclude that blocking autophagy severely disrupts macromolecule
absorption in the neonatal ileum.

To further investigate this role autophagy in macromolecule absorption, we examined the expression of the
endocytic genes *Dab2*, *Amn*, and *Mamdc4*, as well as the lysosomal genes *Mcoln3*, *Lamp2*, and *Ctsl1*.
Previous studies have demostrated the specific role of these genes on the endolysosomal function of LREs
(Cox et al., 2018; Park et al., 2019; Remis et al., 2014). Indeed, we validated their expression during early
postnatal intestinal development (**Figure 2A**). As expected, an upregulation of these genes was observed
in distal samples at P7, coinciding with the presence of LREs (**Figure 2A**). Moreover, a proximodistal
expression pattern was observed, consistent with the antero-posterior maturation of the intestine (Spence
et al., 2011) (**Figure 2A**).

Then, we studied the mRNA expression of *Dab2*, *Mcoln3*, and *Lamp2* genes in both the proximal and distal
regions of the small intestines of WT and *Atg16*cKO mice at P7 and P14 (**Figures 2B and S2G**). Since LREs
predominantly localise in the distal region, the proximal segment was also analysed to assess potential
compensatory mechanisms. In distal *Atg16*cKO samples at P7, mRNA levels of these LRE markers were
significantly reduced (**Figure 2B**), while the proximal region showed no significant changes compared to
WT mice (**Figure S2G**). By P14, a substantial reduction in gene expression was evident in both proximal
and distal *Atg16*cKO mice samples (**Figures 2B and S2G**). Additionally, we analysed *Amn*, *Mamdc4,* and
Ctsl1 mRNA expression in the ileum at P14, finding lower expression levels in the *Atg16*cKO samples
(**Figure S2H**).

To delve deeper into the protein levels of Dab2, we performed WB and immunofluorescence (IF) analysis
in the ileum of *Atg16*cKO mice at P7 and P14. While P7 samples exhibited reduced mRNA expression
(**Figure 2B**), Dab2 protein levels remained comparable to WT; however, by P14, a marked reduction in
Dab2 was observed in *Atg16*cKO samples (**Figures 2C–2F**).

Furthermore, we assess the lysosomal occupancy and content in the apical and basal regions of WT and
*Atg16*cKO ileal enterocytes using confocal microscopy with an anti-LAMP1 antibody as a lysosomal marker,
as the *Lamp1* and *Lamp2* genes exhibit similar expression patterns (data not shown). Post-weaning (P21–
30) WT mice were used as controls (**Figures 2G and 2H**). At P7, both WT and *Atg16*cKO enterocytes
exhibited comparable lysosomal occupancy and content, with peak intensity observed in both conditions.
However, by P14, while lysosomal occupancy remained similar, *Atg16*cKO enterocytes demonstrated
reduced intensity in both the apical and basal regions (**Figures 2G and 2H**). Magnified images also revealed
that lysosomes in *Atg16*cKO enterocytes were smaller in P14 neonates, resembling those of PECs (**Figure
2G**, arrowheads). This reduction in lysosomal content in *Atg16*cKO P14 ileum was further confirmed by WB
analysis (**Figures 2I and 2J**).

In summary, our results indicate that autophagy plays a crucial role in preserving endolysosomal
homeostasis of LREs. Disruption of autophagy in LREs using the *Atg16*cKO mice results in significant
defects in macromolecule uptake and lysosomal function, accompanied by the downregulation of key
endocytic and lysosomal genes, particularly in the distal ileum of late neonatal mice. These outcomes
highlight the importance of autophagy in maintaining LRE function and, consequently, ensuring proper

postnatal development. However, the precise mechanisms by which autophagy exerts its effects remain
unclear.

LREs show a significant induction of mitophagy to control mitochondrial homeostasis

To elucidate the primary type of autophagy in neonatal enterocytes, we defined the gene expression
signature of LREs through bulk RNA sequencing. Leveraging the high levels of macromolecular absorption
and autophagy in LREs, we probed neonatal GFP-LC3 mice (P7) with Rd-dextran and isolated LREs from
the cellular suspension of the ileum by fluorescence-activated cell sorting (FACS), selecting them as the
epithelial cells with high levels of autophagy and Rd-dextran (Dextran$^+$GFP^{high}). Moreover, to highlight their
genetic neonatal signature, LREs were compared to post-weaning enterocytes (PECs) from mice (P42).
Following RNA extraction, we conducted next-generation sequencing (NGS) analysis (**Figures 3A and**
S4A).

We detected a total of 1,537 differentially expressed genes (DEGs), of which 1,036 were upregulated and
501 were downregulated in LREs (**Figure 3B**). Differential expression analysis and principal component
analysis demonstrated that each population exhibited unique enrichment signatures (**Figures S4B and**
S4C). The computational method of over-representation analysis (ORA) identified which biological
processes were overrepresented in the differentially expressed genes (DEGs) (Boyle et al., 2004). The
majority of differentially regulated pathways were associated with lysosomal function, nutrient absorption,
processing, and transport (**Figures 3C, 3D, and S4D**). These findings are consistent with earlier RNA-Seq
data from LREs in zebrafish (Park et al., 2019). Nonetheless, the regulation of lipid metabolism pathways
differed (**Figure 3C and S4D**), suggesting that mammalian LREs significantly modulate these pathways to
process lipids from maternal milk.

Among the most upregulated genes were those associated with endocytosis, such as *Mamdc4*, *Dab2*, and
Amn, as well as lysosome-associated genes like *Mcoln3* and *Ctsl1* (**Figure 3B**). The upregulation of *Dab2*,
Amn, and *Lamp2* genes corresponded with previously published ileum transcriptome data from mice and
zebrafish (Park et al., 2019). Additionally, the overexpression of *Mamdc4* and *Mcoln3* genes also aligned
with earlier results in LREs from rodents (Remis et al., 2014; Wilson et al., 1991). We confirmed elevated
levels of *Ctsl1* mRNA expression and lysotracker staining in LREs compared to PECs in WT mice (**Figures**
S4E and S4F), which validated the RNA-Seq data and ensured that the lysosome-associated pathway was
not positively regulated due to the selection of neonatal enterocytes with high autophagy levels.

Interestingly, the 'positive regulation of autophagy of mitochondrion ' and 'mitochondrial outer membrane'
pathways appeared significantly enriched in LREs (**Figures 3C and 3D**). These outcomes, along with the
upregulation of several genes involved in mitophagy, such as *Bnip3*, *Atg10*, and *Map1lc3a* (**Figure 3B**) (Li
et al., 2022; Yang et al., 2024), indicated that mitophagy is a significant component of the elevated
autophagic activity observed in LREs. To explore this hypothesis, we examined the mRNA expression of

the canonical mitophagy genes *Prkn* (Parkin) and *Pink1* (Picca et al., 2023), together with *Bnip3* as a
control, in ileal samples from WT mice across developmental stages (E18.5-P42). While *Bnip3* exhibited a
progressive decrease in expression throughout development, *Prkn* and *Pink1* showed upregulation in
neonatal samples, especially *Prkn* (**Figure 3E**). Moreover, we observed that LREs displayed increased
Parkin protein levels compared to PECs through IF (**Figure 3F**).

Next, we assessed mitophagy in LREs and PECs using the fixable *mito*-QC reporter C57BL/6J mouse
(McWilliams et al., 2016), which ubiquitously expresses a chimeric protein consisting of pH-insensitive
mCherry, a pH-sensitive GFP, and the mitochondria-targeting sequence (MTS) from the outer mitochondrial
membrane protein FIS1 (amino acids 101–152). This allowed us to differentiate between cytoplasmic
mitochondria (mCherry$^+$GFP$^+$) and mitochondria degraded by mitophagy in lysosomes (mCherry$^+$GFP$^-$)
(**Figure 3G**). Confocal images of LREs revealed a high concentration of red-fluorescent mitochondria inside
their endolysosomal system (**Figures 3H and 3I**). In fact, when we inhibited autophagic flux by treating the
neonates with the protease inhibitor Leupeptin (40 mg/kg for 16 hours) (Jiménez-Loygorri et al., 2024), we
observed a significant expansion of LVs and a considerable accumulation of red-only mitochondria within
them (**Figure 3H**, arrows; **and 3I**). In contrast, PECs exhibited fewer red mitochondria in lysosomes and a
uniform distribution of yellow mitochondria throughout the cytoplasm (**Figures 3H and 3I**).

Thus, we confirmed the high mitophagic activity of LREs, suggesting severe impairment of mitophagy and
mitochondrial integrity in *Atg16*cKO enterocytes. Indeed, we found that mRNA expression levels of both
Bnip3 and *Prkn* (**Figure 3J**), along with Parkin protein levels (**Figure 3K**), were significantly lowered in
neonatal *Atg16*cKO enterocytes. To further investigate the robust mitophagic activity of LREs, we generated
an autophagy-deficient *mito*-QC mouse line (*mQ-Atg16*cKO) to evaluate their status. We observed that in
*Atg16*cKO LREs exhibited reduced mitochondrial mass within the lysosomes (**Figures 3L and 3M**),
consistent with the low mitophagy observed in PECs (**Figures 3L**, lower panels). Therefore, the suppression
of autophagy in LREs results in the downregulation of mitophagy-related genes and a reduced uptake of
mitochondria into lysosomes.

Motivated by these results, we addressed the mitochondrial status of *Atg16*cKO mice. Using transmission
electron microscopy (TEM), we analysed mitochondrial damage, quantity, and morphology in ileal
enterocytes from neonatal *Atg16*cKO mice (**Figure 4A–4C**). Mitochondria were categorized into three
classes based on electron density, cristae appearance, matrix loss, and outer membrane integrity: class I
(normal), class II (partially affected), and class III (damaged) (**Figure 4B**) (Moschandrea et al., 2023). For
each class, the number, percentage, area, and dimensional ratio (aspect ratio, AR: major axis/minor axis)
were quantified (**Figure 4C**). The data showed that number and percentage of mitochondria within each
class were similar across both WT and *Atg16*cKO enterocytes. However, while class I mitochondria in
*Atg16*cKO enterocytes exhibited a smaller area and a higher AR, class III mitochondria showed a larger
area, suggesting an accumulation of damaged mitochondrial mass due to impaired autophagy (**Figure 4C**).

To further validate the mitochondrial status, we performed flow cytometry on neonatal *Atg16*cKO
enterocytes using mitochondrial probes *ex vivo*: Mitotracker Green for mitochondrial mass, Mitotracker Red
CMXROS for membrane potential, and CM-H$_2$DCFDA for ROS (Asano et al., 2017; Iborra-Pernichi et al.,
2024). This analysis confirmed an increase in mitochondrial mass, membrane potential, and ROS in
*Atg16*cKO enterocytes (**Figure 4D**). In conclusion, the inhibition of mitophagy in LREs leads to an increased
in the mass of dysfunctional mitochondrial.

Collectively, mitophagy represents a significant component of the heightened autophagic activity observed
in LREs. Its inhibition not only contributes to the mitochondrial defects seen in neonatal *Atg16*cKO
enterocytes but may also account for the impairment of their endolysosomal system. These findings
highlight a compelling link between the mitochondria-lysosomes crosstalk and neonatal development, as
well as the associated metabolic adaptations.

Dysfunction of the mitochondria-lysosome axis in the neonatal epithelium causes an early
metabolic switch dependent on BLIMP1 and MAFB

An interesting phenotype observed in neonatal *Atg16*cKO enterocytes was their resemblance to PECs,
particularly in terms of the abrogation of macromolecule absorption and the reduction in endolysosomal
system functionality. This phenotype could suggest an acceleration of their metabolic transition. At the end
of lactation, the mammalian intestine becomes capable of digesting nutrients extracellularly, replaces LREs
with PECs (without LVs), and regulates the expression of enzymes to adapt to the new solid diet (Henning,
1981; Skrzypek et al., 2007). Previous studies have demonstrated that the transcriptional factors BLIMP1,
MAFB, and c-MAF (two members of the Maf family of proteins) mediate the metabolic shift in enterocytes
during neonatal development (Bara et al., 2022; Harper et al., 2011; Muncan et al., 2011). At weaning,
Blimp1 expression decreases, leading to a downregulation of neonatal enzymes such as β-galactosidase
(Glb1) and argininosuccinate synthase 1 (Ass1), and an increase in expression of adult enzymes, including
sucrase-isomaltase (Sis), trehalase (Treh), and arginase 2 (Arg2) (Harper et al., 2011; Muncan et al., 2011).
Similarly, early inhibition of MAFB and c-MAF in neonatal intestine induces a comparable metabolic switch
(Bara et al., 2022).

Our RNA-Seq analysis comparing LREs and PECs confirmed these dynamics: LREs expressed higher
levels of *Mafb*, *Blimp1*, *Glb1*, and *Ass1*, along with a decrease of *Sis*, *Treh*, and *Arg2* (**Figure S5A**). *c-maf*
showed no significant differences (**Figure S5A**), though transient regulation at specific neonatal stages
cannot be excluded. Consistently, RT-qPCR analysis of proximal and distal small intestine samples from
WT mice across development (E18.5–P42) revealed neonatal enrichment of transcription factors and
neonatal enzymes, whereas post-weaning samples displayed the opposite profile, with strong induction of
adult enzymes (**Figures S5B and S5C**). *Arg2* expression levels closely paralleled those of Sis and Treh
(data not shown).

To investigate whether autophagy mediated by ATG16 contributes to this transition, we analyzed the
temporal expression patterns of these genes in our model. Gene expression analysis revealed a moderate
downregulation of the transcription factor genes *Blimp1* and *Mafb*, except for *c-maf*, as well as a more
pronounced downregulation of neonatal enzyme genes *Ass1* and *Glb1*. In contrast, there was an
upregulation of adult enzyme genes *Treh*, *Sis*, and *Arg2* (**Figure 5A**). Similarly, confocal microscopy
analysis revealed reduced levels of MAFB, BLIMP1 (**Figures 5B and 5C**), and ASS1, along with increased
levels of TREH in neonatal *Atg16*cKO enterocytes (**Figures 5D and 5E**). This later result was also confirmed
by WB analysis in neonates (**Figure 5F and 5G**). In summary, *Atg16*cKO mice advance their metabolic
transition in enzymes regulated by MAFB/BLIMP1 in their neonatal epithelium.

Next, we studied the metabolic profiles of neonatal enterocytes from *Atg16*cKO mice, in comparison with
neonatal and post-weaning WT mice. In order to evaluate deeply the metabolic transition during the
neonatal development and the impact of autophagy, we used a pioneer technique called Met-flow. The Met-
Flow technique is a flow cytometry-based method that examines the metabolic state of cells by evaluating
the levels of rate-limiting enzymes across key metabolic pathways. Specifically, we studied ASS1 and GLS
as markers of amino acid (AA) metabolism, HK1 as a marker of glycolysis, CPT1A and ACAC as markers
of fatty acid (FA) metabolism, CS and IDH2 for the tricarboxylic acid (TCA) cycle, and ATP5 for oxidative
phosphorylation (OXPHOS) (Ahl et al., 2020; Iborra-Pernichi et al., 2024) (**Figure S5D**). Met-Flow analysis
revealed higher expression of most metabolic pathways in LREs compared to PECs (**Figure S5E**), reflecting
a their hyperconsumptive signature aligned with the high energy demands of this developmental period
(Hulzebos and Sauer, 2007). Notably, neonatal *Atg16*cKO enterocytes exhibited a moderate reduction in
enzyme protein levels, positioning them between those of WT neonates and post-weaning mice (**Figure
5H**). This intermediate profile is indicative of a globally accelerated metabolic transition toward a more
mature state.

In summary, our results reveal that autophagy is essential for coordinating mitochondrial and metabolic
remodeling during neonatal ileum development and the weaning transition. Suppression of autophagy
perturbs endolysosomal and mitochondrial integrity, thereby altering the transcriptional control of LREs.
Consequently, LREs undergo a premature metabolic switch, marked by MAFB and BLIMP1 dysregulation
and reduced activity of key metabolic pathways.

Mitochondrial homeostasis is critical for maintaining LREs function and their metabolic transition

After identifying mitochondrial homeostasis defects as a downstream consequence of impaired
endolysosomal function in the *Atg16*cKO model, we next investigated whether mitochondria themselves
play an intrinsic role in sustaining LRE integrity. For this purpose, we used the *Tfam*cKO mouse model
(*Tfam^{fl/fl}/Villin-creER^{T2}*), which enables inducible deletion of Tfam in the neonatal intestinal epithelium. This
approach allowed us to directly test whether mitochondrial dysfunction is sufficient to compromise
lysosomal homeostasis in LREs.

TFAM is a transcription factor that regulates the transcription and replication of mitochondrial DNA (mtDNA)
(Campbell et al., 2012). While the role of mitochondria in maintaining intestinal homeostasis has been
extensively explored (Rath et al., 2018), and the consequences of Tfam deletion have been reported in
both embryonic and adult intestinal epithelium (Srivillibhuthur et al., 2018), its specific contribution during
the neonatal stage, and particularly in nutrient absorption and processing by LREs, remains unexplored.

We first characterized the general phenotype of *Tfam*cKO pups during the neonatal period. These mice
exhibited developmental defects, a generally poor appearance, and significantly reduced body weight,
starting around P8, which ultimately compromised their survival and prevented successful weaning (**Figures**
6A–6C). Given the severity of the phenotype and in adherence with animal welfare guidelines, euthanasia
was performed after two consecutive days without weight gain or upon observing weight loss of around
10% (**Figure 6C**). Based on the timing of euthanasia, we defined two experimental groups: mice euthanised
between P9–P11 (hereafter referred to as P10), and those euthanised between P13–P15 (hereafter
referred to as P14). In the ileum of both groups, we confirmed a marked reduction in *Tfam* mRNA expression
(**Figure 6D**), accompanied by decreased levels of mtDNA, as assessed through the quantification of
electron transport chain-encoded genes including *mt-Nd1*, *mt-Co*1, *mt-Co2*, and *mt-Atp6* (**Figure S6A**). In
addition, *Tfam* deletion resulted in a selective loss of mitochondrial gene expression at P10 (*mt-Nd1*, *mt-*
Co2), while nuclear *Sdha* expression was unaffected (**Figure S6B**). We also observed a significant reduction
in MTCO1 (complex IV subunit) protein levels in ileal sections of *Tfam*cKO at P14 (**Figures S6C and S6D**).
These results validated our neonatal *Tfam*cKO model and were consistent with findings from other *Tfam*KO
models (Baixauli et al., 2015; Chung et al., 2019; Iborra-Pernichi et al., 2024).

Further histological examination of the distal small intestine of *Tfam*cKO mice at P10 and P14 revealed
substantial structural alterations. The epithelium exhibited clear signs of disorganization and inflammation,
characterised by notably shorter, malformed villi, oedema, reduced GCs, and a thicker submucosa and
lower mucosa (**Figures S6E–S6H**). Enterocytes also exhibited aberrant vacuoles, suggesting dysfunctions
in nutrient processing (**Figure S6E**, arrows).

Moreover, flow cytometry analysis of the ileum at P10 and P14 revealed increased percentages of IIC (B
cells, and Th cells) in *Tfam*cKO samples (**Figures S6I–S6K**). Given the elevated OXPHOS activity
characteristic of intestinal stem cells (ISCs) (Rodríguez-Colman et al., 2017), we assessed ISC populations
in ileal sections from P10 and P14 *Tfam*cKO mice via Olfm4 immunofluorescence (**Figure S6L**, arrows),
which revealed a marked reduction in Olfm4$^+$ crypts (**Figures S6M**). Additionally, lysozyme staining showed
a striking increase in mislocalised PCs (**Figures S6N and S6O**). These PCs exhibited mislocalization,
aberrant morphologies and enlarged granules, indicative of disrupted crypt architecture and compromised
ISC niche homeostasis (**Figure S6N**, arrowheads).

With the systemic and intestinal phenotypes of *Tfam*cKO mice defined, we next investigated how loss of
TFAM affects mitochondrial homeostasis in LREs. Firstly, mitochondria from *Tfam*cKO enterocytes at P10
and P14 were analyzed by TEM, following the same approach previously used in *Atg16*cKO mice. At P10,

*Tfam*cKO enterocytes displayed more class I and II mitochondria, with reduced area and AR (**Figures 6E,**
6F, and S6P). By P14, damaged class III mitochondria increased while class I mitochondria decreased in
number, size, shape and AR (**Figures 6E, 6F, and S6P**). This progressive mitochondrial damage resembles
that observed in a previously described *Tfam*KO model in renal epithelial cells (Chung et al., 2019). These
data were corroborated by subsequent flow cytometry analysis, which revealed increased mitochondrial
mass, membrane potential, and ROS levels in neonatal *Tfam*cKO enterocytes (**Figure 6G**), features
reminiscent of those observed in TFAM-deficient lymphocytes (Baixauli et al., 2015; Iborra-Pernichi et al.,
2024).

In summary, deletion of *Tfam* in the neonatal intestinal epithelium severely disrupted mucosal structure and
function, leading to developmental defects that culminated in early mortality. Specifically, in LREs, TFAM
loss caused the accumulation of damaged mitochondria and increased oxidative stress.

We next investigated whether mitochondrial dysfunction could similarly drive an accelerated metabolic
transition in neonatal enterocytes, as observed in the *Atg16*cKO model. The microscopy analysis indicated
a reduction in the expression levels of BLIMP1 and MAFB, with a more pronounced decrease in BLIMP1
(**Figures 6H and 6I**). As a result, we also observed lower levels of ASS1 and elevated expression of TREH
in the *Tfam*cKO enterocytes (**Figures 6J and 6K**). These results were further supported by gene expression
analyses of metabolic transition genes that are regulated by BLIMP1, MAFB, and c-MAF. We obtained a
progressive downregulation of *Mafb*, *c-maf*, *Blimp1*, *Ass1,* and *Glb1*, accompanied by an upregulation of
Treh, *Sis*, and *Arg2* at P14 (**Figure S7A**). Finally, the activity of key metabolic pathways was assessed in
neonatal *Tfam*cKO enterocytes using Met-Flow (Ahl et al., 2020; Iborra-Pernichi et al., 2024), revealing a
marked reduction in nearly all metabolic proteins, except for ATP5A (**Figure 6L**), which was increased,
possibly due to mitochondrial accumulation or as a compensatory response to mitochondrial dysfunction.
Taken together, these findings indicate that neonatal *Tfam*cKO enterocytes undergo a premature metabolic
transition, resembling the metabolic profile of PECs.

Following the analysis of mitochondrial defects in TfamcKO neonatal enterocytes, we next investigated
whether impaired mitochondrial function also perturbs the endolysosomal system of LREs. Although
lysosome–mitochondria crosstalk has been predominantly characterized in the context of mitophagy, our
findings suggest a reciprocal layer of regulation, whereby mitochondria control lysosomal biogenesis and
function. This bidirectional communication implies that mitochondrial dysfunction not only disrupts energy
metabolism but also secondarily compromises lysosomal integrity, thereby exacerbating the metabolic and
absorptive defects observed in neonatal *Atg16*cKO enterocytes. Indeed, neonatal *Tfam*cKO enterocytes
exhibited a complete loss of Rd-dextran uptake (**Figures 6M and 6N**) and a progressive downregulation of
the endocytic genes *Dab2*, *Amn*, and *Mamdc4* (**Figures S7B–S7D**). Subsequently, we characterised a
gradual reduction in the expression of lysosomal genes *Lamp2*, *Mcoln3*, and *Ctsl1* (**Figure S7E**),
accompanied by a decrease in lysosomal content and size (**Figures 6M, 6O, S7F and S7G**). Thus, neonatal

*Tfam*cKO enterocytes recapitulated the phenotype observed in the *Atg16*cKO model, validating a reciprocal
interaction between mitochondria and lysosomes in LREs.

In conclusion, the *Tfam*cKO model enabled us to establish a bidirectional relationship between mitochondria
and lysosomes by recapitulating the endolysosomal defects observed in autophagy-deficient neonatal
enterocytes. Furthermore, we confirmed that mitochondrial and lysosomal dysfunction drive an early
metabolic switch in LREs. These alterations, alongside epithelial inflammation, developmental delay, and
premature death, position the *Tfam*cKO model as a valuable tool for investigating the pathophysiology of
childhood malnutrition.

Antioxidant activity in LREs modulates the onset of metabolic switch

After defining the metabolic switch underlying neonatal development, and the early metabolic switch in both
*Atg16*cKO and *Tfam*cKO models, we focused on uncovering the mechanisms responsible for this
phenotype. The observation that both *Atg16*cKO and *Tfam*cKO models presented heightened ROS levels
in enterocytes (**Figures 4D and 6G**) suggested that redox dysregulation may underlie their accelerated
metabolic maturation. Indeed, the high mitophagic activity observed in LREs may contribute to the control
of ROS (Wang et al., 2023; Zhong et al., 2024). Previous studies have shown that oxidative stress has
been reported to repress *c-maf* expression (Chen et al., 2015) and that antioxidant treatment can induce
Blimp1 expression (Qi et al., 2020). Therefore, we hypothesise ROS regulation could represent a critical
mechanism for controlling the timing of the metabolic transition in LREs.

We found that LREs exhibited significantly higher levels of mitochondrial ROS (mROS) compared to PECs
(**Figure 7A**), possibly reflecting their elevated OXPHOS activity (Murphy, 2008). In contrast, total ROS levels
were lower in LREs than in PECs (**Figure 7A**), indicating the presence of mechanisms that tightly regulate
oxidative stress.

Then, we further explored our LREs vs PECs RNA-Seq dataset analysis to identify additional pathways
involved in ROS regulation. Numerous pathways related to ROS, ROS response, and antioxidant activity
stood out (**Figure 7B and 7C**). Further, there was an overexpression of genes for the antioxidant enzymes
Glutathione Peroxidase-3 (Gpx3) and Catalase (Cat) (not shown), suggesting that these antioxidants could
be reducing ROS levels in LREs. Indeed, we observed that neonatal ileal samples showed induction of
Gpx3 and *Cat* expression compared to the embryo and post-weaning ileal samples, with the most drastic
changes in *Gpx3* (**Figure 7D**). Moreover, in both *Atg16*cKO and *Tfam*cKO neonates, *Gpx3* and *Cat*
expression were significantly decreased in the ileum (**Figure 7E and 7F**), and GPX3 protein levels were
likewise reduced, as shown by IF (**Figure 7G-J**). Thus, oxidative stress may act as a trigger for the metabolic
transition in LREs. To validate this hypothesis, neonatal mice were treated with hydrogen peroxide (H_2O_2)
to induce oxidative stress (Liang et al., 2024). mRNA expression analysis revealed a downregulation of the
transcription factors *Blimp1*, *Mafb*, and *c-maf*, alongside an upregulation of adult enzymes such as *Treh*

and *Arg2*. However, the expression of neonatal enzyme genes *Ass1* and *Glb1* remained unchanged (**Figure**
7K). Subsequently, we characterised a reduction of BLIMP1 and ASS1 levels by both IF and flow cytometry
(**Figure 7L–7N**), while TREH expression was found to be increased (**Figure 7L and 7M**).

The data collectively suggest that elevated antioxidant activity is a key characteristic of LREs. When this
antioxidant function is compromised in neonatal *Atg16*cKO and *Tfam*cKO enterocytes, alongside
mitochondrial dysfunction, the oxyidative stress rise. This increase in ROS contributes to the repression of
genes such as *Blimp1*, *Mafb*, and *c-maf*, ultimately triggering an early metabolic transition (**Figure 7O**).
Therefore, this study emphasiszes the critical role of mitochondrial-lysosomal coordination in mitigating
oxidative stress and regulating the transcriptional programs essential for neonatal intestinal development.
Additionally, it paves the way for future research into therapeutic strategies aimed at restoring mitochondria-
lysosome homeostasis and redox balance in early gut disorders.

DISCUSSION

In this work, we have elucidated new molecular mechanisms fundamental to the function and differentiation
of mammalian LREs, highlighting the role of the mitochondria-lysosome axis. Disruption of autophagy in
the neonatal gut, either through *Atg16*cKO or mitochondrial dysfunction via *Tfam*cKO, resulted in defective
nutrient absorption and a reduction in lysosomal content. Most importantly, *Tfam*cKO neonates exhibited
weight loss and premature death, resembling hallmarks of infant malnutrition. Notably, the early enzymatic
maturation of *Atg16*cKO and *Tfam*cKO enterocytes indicates a direct connection between autophagic and
mitochondrial processes in enterocytic maturation. Additionally, the associated increase in ROS observed
in both models may contribute to this metabolic switch by suppressing the expression of transcription factors
BLIMP1, MAFB, and c-MAF. These findings are consistent with our transcriptomic data, which reveal
upregulation of mitophagy- and antioxidant-related pathways in LREs compared to PECs, supporting the
concept that mitochondrial quality control and redox homeostasis are defining features of LRE identity and
function. Taken together, our outcomes offer interesting mechanistic insights into the mitochondria-
lysosome crosstalk and its role in postnatal intestinal physiology, with implications for understanding the
pathophysiology of neonatal malnutrition and related disorders.

Autophagy and endocytosis are vesicle-mediated pathways that converge at the lysosome to degrade and
recycle intra- and extracellular components. These processes intersect at multiple stages (vesicle
formation, trafficking, and fusion) and share several key regulators. Mounting evidence indicates reciprocal
dependence between these processes (Birgisdottir and Johansen, 2020; Yim and Mizushima, 2020). Using
GFP-LC3 mice, we demonstrated that neonatal ileal samples exhibit increased autophagosome
degradation compared to post-weaning counterparts (**Figures 1A–1C and S1C**). Consistently, *Atg16*cKO
neonates displayed impaired absorption (**Figures 1H–1J**) and showed reduced expression of crucial
endocytic genes characteristic of LREs (**Figures 2B–2F, S2G and S2H**). Moreover, *Atg16*cKO neonatal

enterocytes displayed reduced lysosomal size and content (**Figures 2G–2J**), along with downregulation of
key lysosomal genes (**Figures S2G and S2H**). These outcomes reveal a functional interplay between
autophagy and the endolysosomal system in LREs.

Despite these endolysosomal defects, *Atg16*cKO mice reached the post-weaning stage without apparent
physiological abnormalities, aside from mild signs of intestinal inflammation (**Figures S2A–S2D**). This
observation aligns with findings in mice lacking the lysosomal cation channels Mucolipin-1 and -3 (Mcoln1
and Mcoln3), which also exhibit pronounced endolysosomal abnormalities in neonatal enterocytes but
develop into phenotypically normal adults (Remis et al., 2014). Together, these results suggest that
endolysosomal and autophagic processes primarily affect LREs. Accordingly, once the neonatal period
concludes and LREs are replaced by PECs, the phenotype becomes attenuated or fully resolves. However,
further investigations will be needed to assess intestinal physiology and metabolism in adult *Atg16*cKO
mice comprehensively.

We characterized a significant increase in autophagic activity in LREs as primarily due to the lysosomal
degradation of mitochondria, a process known as mitophagy. Indeed, mitophagy-related genes exhibit a
higher expression in neonatal ileum compared to post-weaning samples (**Figures 3B–3F**). Correspondingly,
neonatal *mito*-QC mice (McWilliams et al., 2016) demonstrated enhanced mitochondrial degradation within
the LVs of LREs compared to PECs (**Figures 3H and 3I**). Furthermore, this process was inhibited either by
leupeptin treatment (Jiménez-Loygorri et al., 2024) (**Figures 3H and 3I**) or through genetic ablation of
autophagy (*Atg16*cKO) (**Figures 3L and 3M**), confirming the dependence of mitochondrial clearance on
lysosomal-autophagic activity. Consistent with this, neonatal *Atg16*cKO enterocytes accumulated
dysfunctional mitochondrial mass (**Figure 4**). Specifically, these cells showed an increase in the area of
damaged mitochondria (class III), whereas healthy mitochondria (class I) exhibited a smaller area but an
increased aspect ratio (**Figures 4A–4C**). These differences possibly reflect not only mitophagy blockade
but also alterations in mitochondrial dynamics, such as enhanced fusion or reduced fission, which are aimed
at preserving oxidative phosphorylation (OXPHOS) capacity (Zemirli et al., 2018). Future studies should
investigate the expression of proteins involved in mitochondrial fusion and fission in the *Atg16*cKO model
to define better the molecular landscape that integrates mitochondrial dynamics with mitophagy in LREs.

After identifying mitochondrial homeostasis defects secondary to impaired endolysosomal function in the
*Atg16*cKO model, we next investigated the reverse relationship by directly disrupting mitochondrial function
using the *Tfam*cKO model. *Tfam*cKO neonates exhibited severe growth impairment and premature death
(**Figures 6A–6C**), and their enterocytes accumulated markedly damaged mitochondria (**Figures 6E–6G and
S6P**). These mice also reproduced key features of the *Atg16*cKO phenotype, including loss of uptake
capacity, reduced lysosomal content, and downregulation of endolysosomal genes (**Figures 6M-6O and
S7B–S7G**). Interestingly, these lysosomal defects differ from what has been observed in *Tfam*-deficient T
and B lymphocytes, where dysfunctional acidic lysosomal compartments are expanded (Baixauli et al.,
2015; Iborra-Pernichi et al., 2024). In neonatal enterocytes, however, Tfam deletion results in diminished

lysosomal size and content, highlighting that mitochondria–lysosome interactions are highly cell-type

specific. Beyond this bidirectional crosstalk, our data raise the possibility of a retrograde signalling

mechanism. Mitochondria are increasingly recognized as signalling hubs that influence nuclear

transcriptional programs, as shown in immune cells. In LREs, two interconnected processes may be at play:

first, endolysosomal-mediated nutrient internalization, which is defective in both *Atg16*cKO and *Tfam*cKO

models; and second, mitophagy-driven mitochondrial degradation, which provides retrograde signals to the

nucleus. This latter pathway could be essential for controlling the transcriptional networks that govern the

metabolic switch (via BLIMP1 and MAFB) and for sustaining lysosomal functionality itself. In this view,

lysosomes must degrade mitochondria not only to preserve organelle quality control but also to control the

transcriptional reprogramming required for the neonatal-to-adult metabolic transition.

At weaning, the mammalian intestine undergoes a shift to extracellular nutrient digestion and adjusts its

enzymatic profile to meet the metabolic demands of a carbohydrate-rich solid diet (Henning, 1981). At the

same time, LREs are replaced by PECs lacking large lysosomal vacuoles (Skrzypek et al., 2007). The

transcription factors BLIMP1 (PRDM1), along with MAFB and c-MAF (both members of the Maf protein

family) have been positioned as central repressors of this switch. Neonatal mice lacking these transcription

factors in the intestinal epithelium exhibited impaired nutrient absorption, premature downregulation of

neonatal enzymes, and upregulation of adult counterparts, ultimately leading to developmental defects

(Bara et al., 2022; Harper et al., 2011; Muncan et al., 2011). Furthermore, our research demonstrated that

dysfunctions in autophagic, lysosomal, and mitochondrial processes in LREs —induced by the deletion of

Atg16l1 and *Tfam*— led to a repression of these transcription factors, resulting in similar phenotypes

(**Figures 1H, 5, 6H–6N and S7A**).

We further analysed the activity of core metabolic pathways in WT, *Atg16*cKO, and *Tfam*cKO neonatal

enterocytes, comparing them to PECs (**Figures 5H, 6L, and S5E**). This analysis revealed a moderate

reduction in metabolic activity in *Atg16*cKO cells and a more noticeable decline in *Tfam*cKO enterocytes,

suggesting a shift toward an adult-like metabolic state. These findings contrast with previous transcriptomic

data from neonatal mice lacking MAFB and c-MAF, which reported an upregulation of OXPHOS and fatty

acid degradation pathways (Bara et al., 2022). However, that study did not include a direct comparison to

PECs, leaving open the possibility of compensatory mechanisms. In this context, our results may provide

a more accurate reflection of the metabolic trajectory during neonatal-to-adult transition and its dependence

on the mitochondria-lysosome axis.

We have defined the metabolic evolution of enterocytes during neonatal development beyond the genes

classically associated with the transcription factors BLIMP1, MAFB, and c-MAF, and established a

functional link between this transition and the maintenance of lysosomal and mitochondrial homeostasis.

At the same time, our data reinforce the idea that, in enterocytes, genetic programs prevail over the sensing

of external cues: despite the neonatal context (milk-based diet, reduced microbiota, and limited immune

activation) (Walker, 2013), the repression of these transcription factors leads to inevitable epithelial

maturation. Yet, the cellular mechanisms that regulate these genetic pathways still need to be clarified.

Reactive oxygen species (ROS) are oxygen-derived molecules that play a crucial role in various cellular

processes, including energy production, mitochondrial signalling, and cell differentiation (Sena and

Chandel, 2012; Sies et al., 2017). Their intracellular levels are tightly regulated through multiple

mechanisms: the mitochondrial electron transport chain constitutes a major source of ROS, while

antioxidant enzymes serve as key systems for neutralizing their excess (Murphy, 2008; Sena and Chandel,

2012; Sies et al., 2017). In the intestine, ROS have been primarily associated with more differentiated

epithelial cells exhibiting higher OXPHOS activity (Rath et al., 2018). However, their role in epithelial

maturation during neonatal intestinal development continues to be unexplored.

Our results indicate that LREs display a transcriptional signature consistent with elevated antioxidant

activity, possibly as a compensatory mechanism to counteract high levels of mitochondrial ROS resulting

from hyperactivated OXPHOS (**Figures 7A–7D**). Beyond this homeostatic role, recent studies suggest that

this antioxidant signature may also be linked to the repression of transcription factor genes *Blimp1*, *Mafb*,

and *c-maf*, and consequently, to the maintenance of LRE identity and function. The antioxidant vitamin

ascorbic acid has been shown to increase the expression of Blimp1 in B lymphocytes through the activity

of DNA demethylases TET2 and TET3 (Qi et al., 2020). Conversely, treatment with H_2O_2 decreases the

expression of *c-maf*, whereas the ROS scavenger N-acetyl cysteine (NAC) can restore *its* levels in human

adipose tissue-derived mesenchymal stem cells (Chen et al., 2015).

This hypothesis is further supported by the elevated ROS levels and the loss of the antioxidant signature

observed in our knockout models (**Figures 4D, 6G, 7E–7J**), as well as by our experiments in which H_2O_2

treatment of neonatal WT mice accelerated the metabolic transition of enterocytes (**Figures 7K–7N**).

Together, these outcomes suggest that both ROS levels and ROS-regulated genetic programs actively

shape epithelial differentiation and metabolic programming in the neonatal intestine (**Figure 7O**). This may

represent a physiological mechanism to prevent premature transitions until the tissue is developmentally

prepared. In this context, therapeutic strategies aimed at fine-tuning redox homeostasis could offer a

powerful approach to supporting neonatal intestinal health.

In addition to their role in nutrient absorption, LREs may act as immunometabolic sentinels during the

neonatal period, integrating microbial cues and intracellular metabolic status to modulate immune

responses. Due to their high endolysosomal activity, LREs could internalise and process luminal antigens,

including microbial components, thereby contributing to the development of the intestinal immune system

(Massaquoi and Guillemin, 2019). In line with this, our RNA-Seq analysis revealed multiple differentially

regulated immune-related pathways (data not shown).

Disruption of lysosomal and mitochondrial homeostasis in *Atg16*cKO and *Tfam*cKO mice likely impairs

these immune functions, as suggested by the inflammatory status observed in both models (**Figures S2D–**

S2F, S6E, and S6H–S6K). In fact, mice with autophagy defects also exhibit gut microbiota alterations and

immune dysregulation (Larabi et al., 2020). Furthermore, cytosolic mtDNA resulting from mitochondrial
damage due to TFAM deficiency has been shown to activate the cGAS–STING inflammatory pathway
(Chung et al., 2019), while mitophagy induction, along with improved mitochondrial status, attenuates this
response (Jiménez-Loygorri et al., 2024). Additionally, *c-maf* KO mice exhibit changes in intraepithelial
lymphocytes and microbiota composition (Cosovanu et al., 2022), suggesting that genetic programs
involved in epithelial maturation, affected in our KO models, may be tightly linked to intestinal immune and
microbial homeostasis.

Given the immunometabolic imbalance observed in *Atg16*cKO and *Tfam*cKO neonates, it is plausible that
these models experience alterations in their gut microbiota. These changes may not only reflect enterocytic
dysfunction but could also actively contribute to the phenotype by disrupting microbial-derived cues
essential for LRE maintenance and programming. Future studies involving microbiota profiling and
transplantation will help determine whether restoring a balanced microbial ecosystem can rescue LRE
function.

Finally, the potential microbiota alterations described above, together with the inflammation and oxidative
stress observed in our KO models, suggest parallels with known hallmarks of children's malnutrition
disorders. Particularly, *Tfam*cKO mice display a more pronounced phenotype and marked developmental
defects, resembling features of Kwashiorkor, a severe form of protein deficiency prevalent in low-income
settings (Bhutta et al., 2017; Trehan and Manary, 2015). Notably, the redox imbalance observed in our
models resembles clinical features reported in malnourished children, who typically exhibit reduced levels
of antioxidants (Bhutta et al., 2017). Accordingly, pilot studies have demonstrated that antioxidant
supplementation may provide therapeutic benefit in conditions such as Kwashiorkor (Becker et al., 2005).
Moreover, our H_2O_2 treatment experiments, which replicated the epithelial maturation phenotype
characterized in our gene-deficient mice, might similarly induce nutritional, microbial, and inflammatory
disturbances. Indeed, intestinal tissues from H_2O_2-treated neonates displayed signs of inflammation (data
not shown).

Collectively, these considerations suggest that malnourished children may also experience premature
intestinal maturation and highlight the need for deeper investigation into antioxidant therapies as potential
interventions. Furthermore, our KO models may serve as valuable experimental platforms to investigate
the mechanisms underlying neonatal malnutrition and to explore interesting avenues for its treatment.
Altogether, future studies aimed at restoring LRE mitophagy and metabolism, as well as modulating the
redox balance, could yield promising strategies to counteract the devastating consequences of malnutrition-
related disorders in early life.

In conclusion, our study uncovers novel mechanisms underlying the function and metabolic transition of
neonatal enterocytes. We demonstrate that lysosomal and mitochondrial homeostasis is indispensable for
efficient nutrient absorption and for sustaining the high metabolic activity required during this energy-
demanding stage of life. Further investigation of the mitochondria-lysosome axis in LREs will not only

advance our understanding of neonatal intestinal physiology but also contribute significantly to the
development of targeted interventions against childhood malnutrition.

Limitations of the study

This work demonstrates that neonatal intestinal deficits in autophagy and mitochondrial homeostasis lead
to endolysosomal dysfunction in enterocytes, impairing development and ultimately triggering inflammation
and early lethality. However, since we used *Villin-cre*-based models, where gene deletion occurs in all
intestinal epithelial cells, we cannot exclude their contribution to the observed phenotypes. Indeed, we
detected altered abundance and morphology of these cells in both *Atg16*cKO (**Figures 1F, 1G, and S2E–
S2F**) and *Tfam*cKO mice (**Figures S6G, S6L–S6O**). Previous studies support this interpretation: defects in
GCs and PCs are linked to intestinal inflammatory diseases (Hodzic et al., 2017; Wallaeys et al., 2023),
and intestinal autophagy loss has been associated with increased oxidative stress in ISCs (Asano et al.,
2017), which could partially explain the elevated ROS levels in *Atg16*cKO enterocytes (**Figure 4D**).
Similarly, *Tfam* deletion in the adult intestinal epithelium causes ISC loss and lethality (Srivillibhuthur et al.,
2018), suggesting that depletion of ISCs may contribute to the early death of neonatal *Tfam*cKO mice,
limiting our ability to study mitochondrial function in enterocytes at later developmental stages.

To address these limitations, we plan to develop an LRE-specific tamoxifen-induced conditional knockout
(cKO) model to target critical genes specifically in neonatal enterocytes. Although selectively targeting
neonatal enterocytes would yield a more precise model, this is hindered by the lack of fully specific
promoters and the technical difficulty of isolating and culturing LREs, which are highly sensitive to
dissociation-induced cell death. To overcome these challenges, we are also studying *in vitro* systems that
recapitulate the properties of LREs and may serve as a platform for future functional validation.

RESOURCE AVAILABILITY

Lead contact
Further information and requests for resources and reagents should be directed to and will be fulfilled by
the lead contact, Fernando Martin-Belmonte (fmartin@cbm.csic.es).

Materials availability
This study did not generate new unique reagents.

Data and code availability
RNA-seq Illumina paired-end reads (FASTQ) generated for this study are available at the European
Nucleotide Archive (ENA; http://www.ebi.ac.uk/ena/) under the study accession number PRJEB46511.

ACKNOWLEDGMENTS

We thank Carmen M. Ruiz-Jarabo for her comments on the manuscript and members of the Martin-
Belmonte and Martinez-Martin labs for their support and helpful discussions. This work was supported by
MICINN (BFU2017-83243-P; PID2020-120367GB-I00; PID2023-151844OB-I00; RED2022-134927-T),
Fondos FEDER investigación (Horizon-MSCA-2022-DN-101119504) and Fundación Ramon Areces
(CIVP18A3904). HR22-00447 from ¨La Caixa¨, RTI2018-101586-A-I00, PID2021-126298OB-100 to N.M.-
M. G.H. and M.I.-P. were supported by FPI fellowships BES-2018-077789 and PRE2019-089076,
respectively, funded by the Spanish Ministry of Science and Innovation and the State Research Agency
(MCIN/AEI /10.13039/501100011033). D.A.L was supported by FPU and JAE fellowships FPU23/03569
and JAEINT_22_01549. L.A was supported by Horizon-MSCA-2022-DN -101119504. I.G.G. was funded
by the Medical Research Council, UK (MC_UU_00038/2). We thank Victor Lazaro Morato (currently at IMN-
CNM, CSIC, Madrid, Spain) for his contribution to the initial analyses of the general phenotype and
mitochondrial status by TEM in the *Tfam*cKO model. We acknowledge Michel Bagnat (Duke University,
USA) for the generous donation of mCherry plasmid, and Juan M. Serrador, Catalina Grabowski and
Dionisio Ureña (CBM), for their essential contribution to the protocol for the growth and purification of
mCherry plasmid. We are thankful to the Animal, Flow Cytometry, Advanced Optical Microscopy, and
Electron microscopy facilities at the CBM, as well as the Histology Service at the Centro Nacional de
Biotecnologia (CNB, CSIC, Madrid, Spain), for their excellent technical support. We acknowledge the
Centro de Análisis Genómico (CNAG-CRG, Barcelona, Spain) for the RNA sequencing, and to
Biocomputational Analysis Core Facility at the CBM, specially to Sandra González de la Fuente, for the
NGS data analysis and plotting. We thank the computational resources and support provided by the Centro
de Computación Científica at Universidad Autónoma de Madrid (CCC-UAM).
Certain Figures were inspired by or adapted from previously published sources, some of which are licensed
under Creative Commons (CC BY 4.0) (https://creativecommons.org/licenses/by/4.0/). While every effort
has been made to credit original sources, any omissions are unintentional and will be corrected in future
revisions if identified. Some figures were created with BioRender.com.

DECLARATION OF INTERESTS

The authors declare no competing interests.

FIGURE TITLES AND LEGENDS

Figure 1. High Autophagic Activity of LREs Is Required for Nutrient Absorption

(A and B) Assessment of autophagic flux and protein absorption along the small intestine of GFP-LC3 mice
during postnatal development. GFP-LC3 mice at postnatal days (P) 1, 7, 14, and 21 were orally gavaged
with mCherry and sacrificed 3 hours later (n = 3) (A). Proximal and distal intestinal segments were harvested
(A), and epithelial cell lysates analysed by immunoblotting for mCherry and GFP-LC3 isoforms (B). The

representative western blot in (B) distinguishes cytosolic LC3-I, membrane-bound LC3-II, and cleaved "free GFP," a marker of autolysosomal degradation. The pink rectangle highlights mCherry and free GFP in distal-P7, indicating maximal protein uptake and autophagic degradation. USP7, loading control. Figure 1A includes images adapted from Birchenough et al. (2023) (CC BY 4.0) and Zhao et al. (2021), and the GFP-LC3 isoforms scheme is based on Ni et al. (2011).

(C) Quantification of mCherry and free GFP levels in lysates of ileal epithelial cells from P7 (n = 4) and P21 (n = 5) GFP-LC3 mice. Data derived from Figure S1C.

(D and E) Generation and validation of *Atg16l1* intestinal knockout (*Atg16*cKO) mice. Ileal epithelial cell lysates from WT and *Atg16*cKO P14 mice were immunoblotted for ATG16L1, LC3, and P62 (D). (E) Quantification of ATG16L1 (n = 5,6) and P62 (n = 6,7), and LC3-II/LC3-I ratio (n = 17). In ATG16L1 and P62 graphs, data relative to WT.

(F and G) Ileal architecture in *Atg16*cKO neonates. Periodic acid-Schiff (PAS)-haematoxylin staining of P14 ileal sections from WT and *Atg16*cKO mice (F) shows villus shortening (double red arrow) and a decrease in goblet cells (PAS+, arrowheads) in *Atg16*cKO sections (n = 3). L, lumen; scale bar, 100 μm. (G) Quantification of villus length and goblet cells per villus. Dots represent individual villi (n = 30 villi from 3 mice; 10 villi per mouse).

(H) Confocal analysis of Rd-dextran uptake in ileal *Atg16*cKO enterocytes. WT and *Atg16*cKO neonates at P7 (early) and P14 (late), and post-weaning (P21–30) WT mice were gavaged with rhodamine B–dextran (Rd-dext, red) 16 hours prior to sacrifice (n = 3). Ileal sections labelled with β-catenin (βcat, white) and DAPI for DNA (blue). Dashed-line squares indicate magnified areas. Arrows point to lysosomal vacuoles; arrowheads, apical endosomes. L, lumen. Scale bars: 10 μm (main); 5 μm (insets). Dot plot shows the quantification of Rd-dextran intensity (integrated density) in the apical and basal regions of enterocytes. Dots represent individual enterocytes (n = 25–40).

(I and J) *Atg16*cKO mice show impaired mCherry absorption. P14 mice were gavaged with mCherry and ileal epithelial cell lysates collected 16 hours later for immunoblotting (I). Actin, loading control. (J) Quantification of mCherry signal, normalised to WT (n = 9,11).

In (C, E, J), dots represent individual mice and charts depict mean + SEM. In (G, H), dot plots show mean ± SEM (black lines).

(C, E, G): Student's t-test (mCherry, ATG16L1, P62, villus length) and Mann–Whitney test (free GFP, LC3 ratio, goblet cells/villus). (J): Mann–Whitney test. (H): Kruskal–Wallis test; only statistically significant comparisons between WT and *Atg16*cKO are shown within each region (apical or basal). *p < 0.05; ****p < 0.0001.

See also Figure S1.

Figure 2. Loss of Autophagy Compromises the Endolysosomal System in LREs

(A) Reverse transcription quantitative polymerase chain reaction (RT-qPCR) analysis of *Dab2*, *Mcoln3*, *Lamp2* and *Ctsl1* genes in epithelial cells from proximal and distal samples of small intestines from WT mice at the indicated time points (n = 3). Data relative to values of proximal region of small gut from embryonic day (E) 18.5 mice embryos, and presented as $\log_{10}$ fold changes ($\log_{10}$FC).

(B) RT-qPCR of *Dab2*, *Mcoln3*, and *Lamp2* in ileal epithelial cells from WT and *Atg16*cKO mice at P7 and P14. Sample sizes: *Dab2* and *Lamp2* (n = 10,6,6,6), *Mcoln3* (n = 9,8,6,6).

(C and D) Immunoblot analysis of Dab2 protein in ileal epithelial cell lysates from WT and *Atg16*cKO mice at P7 and P14 (C). Actin, loading control. (D) Quantification of Dab2 protein levels (n = 6,8,9,14).

(E and F) Confocal images of ileal sections from WT and *Atg16*cKO mice at P7 and P14 (n = 3) labelled with anti-Dab2 (green), β-catenin (magenta), and DAPI (blue) (E). Arrowheads mark apical Dab2 localization. (F) Violin plot quantifying apical Dab2 intensity (integrated density) of P14 sections, normalised to epithelial area (n = 10,8; 2–4 images per mouse).

(G and H) Lysosomal occupation and content in WT and *Atg16*cKO mice at P7 and P14, and in post-weaning (P21–30) WT mice. Confocal sections labelled with anti-LAMP1 (green), β-catenin (white), and DAPI (blue) (G). Arrows: endolysosomal system in WT LREs; arrowheads: reduced LAMP1[+] vesicles in *Atg16*cKO and post-weaning enterocytes. (H) Dot plots quantify the percentage of LAMP1[+] area and LAMP1 intensity (integrated density) within apical and basal regions of interest (ROIs) (n = 26–36).

(I and J) Immunoblot analysis of LAMP1 protein in ileal epithelial cell lysates from WT and *Atg16*cKO mice at P14 (I). Actin, loading control. (J) Quantification of LAMP1 protein levels (n = 10,13).

In (E, F), dashed-line squares indicate magnified areas; L, lumen. Scale bars: 10 µm (main); 5 µm (insets).

Dots represent individual mice (B, D, J) or images (F) or cells (H). In (B, D, F, J), data relative to WT values. Graphs of (A) represent mean + SD; graphs of (B, D, J), mean + SEM; violin plot of (F), median (thick line) and interquartile range (IQR, thin lines); dot plots of (H), mean ± SEM (black lines).

Statistical analyses: Kruskal–Wallis test (A [*Dab2*, *Mcoln3*], H), two-way ANOVA (A: *Lamp2*, *Ctsl1*), Student's *t*-test (B, D, J), Mann–Whitney test (F). In (A), only the most relevant statistically significant comparisons are shown. In (H), only statistically significant comparisons are displayed within each group (apical or basal). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

See also Figure S2.

Figure 3. LREs Present High Levels of Mitophagy

(A) Workflow of RNA sequencing (RNA-seq) analysis comparing LREs from P7 GFP-LC3 mice gavaged with rhodamine B–dextran (16 hours prior) to PECs from post-weaning (P42) GFP-LC3 mice. Enterocytes

were isolated from ileal samples by fluorescence-activated cell sorting (FACS) and used for RNA-seq. See
Figure S4A for gating strategy. Scheme includes images adapted from Treuting et al. (2018) and Park et
al. (2019).

(B) Volcano plot of differentially expressed genes (DEGs) between LREs and PECs. Each point represents
a gene, plotted by $\log_2$ fold change (x-axis) and $-\log_{10}$ adjusted p-value (padj) (y-axis). DEGs (n = 1537;
shown in steel blue) were defined as genes with padj < 0.05 and $|\log_2 FC| > 1$. Upregulated genes associated
with endocytosis (amber), lysosome (purple), and mitophagy (red) are highlighted. Dashed lines indicate
significance (padj < 0.05) and fold-change thresholds (±1). Genes with $-\log_{10}(padj) > 60$ are not shown.

(C) Bubble plot of selected Gene Ontology (GO) terms from over-representation analysis (ORA) of the
differential gene expression data from the LREs vs PECs. Note the presence of regulation of autophagy,
positive regulation of autophagy of mitochondrion and mitochondrial outer membrane pathways. The x-axis
represents the z-score, bubble size reflects the number of associated genes, and bubble colour indicates
statistical significance (adjusted p-value), with a red-to-blue gradient from most to least significant. GO
terms are grouped into three categories: BP, biological process; CC, cellular component; MF, molecular
function.

(D) Enrichment plots of lysosome organization and mitochondrial outer membrane gene sets from gene set
enrichment analysis (GSEA) comparing LREs and PECs. The top panel shows the Enrichment Score (ES)
across ranked genes, with the peak indicating maximum enrichment. The middle panel marks gene
positions in the ranked list. The bottom colour bar reflects gene correlation with phenotype (red: positive,
blue: negative).

(E) RT-qPCR analysis of *Bnip3*, *Prkn*, and *Pink1* genes in ileal epithelial cells from WT mice at indicated
time points (n = 3). Values are expressed as $\log_{10}$ fold change relative to E18.5 embryonic levels.

(F) Confocal images of ileal sections from WT mice at P7 (LREs) and P21–30 (PECs), labelled with anti-
Parkin (green) and DAPI (blue). Left: violin plot of Parkin intensity (integrated density) normalised to
epithelial area.

(G) Schematic of the *mito*-QC reporter mouse model. Mitochondria are tagged with the chimeric protein
mCherry-GFP-FIS1[101–152]. Upon mitophagy, mitochondria are delivered to the lysosome where acidic pH
quenches GFP but not mCherry, allowing identification of mitolysosomes as mCherry-only puncta.

(H) Confocal images of ileal sections from *mito*-QC mice at P7 (LREs) (± leupeptin, LP) and P21–30 (PECs).
Samples stained with anti-LAMP1 (white) and DAPI (blue). Arrows indicate enlarged, blocked lysosomes
with red-only mitochondrial aggregates in LREs+LP; arrowheads point to LAMP1[+]/mCherry[+] puncta lacking
GFP, validating the *mito*-QC model.

(I) Violin plot of lysosomal mCherry intensity normalised to cellular mCherry intensity from (H), based on
integrated density measurements.

(J) RT-qPCR of *Bnip3* and *Prkn* in ileal epithelial cells from WT and *Atg16*cKO mice at P14. Dots represent
individual mice. Sample sizes: *Bnip3* (n = 6, 6), *Prkn* (n = 5, 6).

(K) Confocal images of ileal sections from WT and *Atg16*cKO mice at P14 stained with anti-Parkin (green)
and DAPI. Left: violin plot of Parkin intensity (Integrated Density) normalised to epithelial area.

(L) Confocal images of ileal sections from *mQ*-WT and *mQ-Atg16*cKO mice at P14 (n = 3), labelled with
anti-LAMP1 (white) and DAPI (blue).

(M) Violin plot of lysosomal mCherry intensity normalised to cellular mCherry intensity from (L), based on
integrated density measurements (n = 12–11; 3–4 images per mouse).

In images of (F, H, K, L), L, lumen. Scale bars: 5 μm. Images of (H, L) deconvolved with Huygens software.

In (A–I) and (K–M), n = 3; in (F, I, K), n = 9 images from 3 mice (3 images per mouse). In graphs of (F, I),
data relative to LREs values; in (J, K, M), relative to WT values. In (E), graph represents mean + SD; in (J),
mean + SEM. In violin plots of (F, I, K, M), the median (thick line) and IQR (thin lines) are shown and dots
represent individual images.

Statistical analyses: Kruskal–Wallis test (E: *Bnip3*), one-way ANOVA (E [*Prkn*, *Pink1*], I), Student's *t*-test
(F, J, K, M). *p< 0.05, **p< 0.01, ***p < 0.001.

See also Figure S4.

Figure 4. *Atg16*cKO Neonatal Mice Exhibit Increased Mitochondrial Damage and Oxidative Stress in
Enterocytes

(A) Representative transmission electron microscopy (TEM) micrographs of ileal sections from WT and
*Atg16*cKO mice at P14 (n = 3). Note the reduction of the endolysosomal system and the absence of
lysosomal vacuole in the *Atg16*cKO enterocytes. Dashed boxes indicate magnified areas. N, nucleus; LV,
lysosomal vacuole; AEC, apical endosomal complex; L, lumen. Asterisks indicate damaged mitochondria.
Scale bars: 2 μm (main); 500 nm (insets).

(B) TEM-based classification of mitochondria into three categories based on electron density and cristae
integrity: Class I (normal), Class II (partially affected), and Class III (damaged). Classification based on
Moschandrea et al. (2023). See Electron Microscopy Section in Methods. Scale bar: 500 nm.

(C) Distribution of mitochondrial integrity classes in *Atg16*cKO neonatal enterocytes. Quantification of the
number, percentage, area and aspect ratio (long axis / short axis) of each mitochondria class in (A), as
defined in (B). In count and percentage graphs, dots represent individual enterocytes; in area and aspect
ratio graphs, dots represent individual mitochondria. Sample sizes are indicated in Table S1.

(D) Flow cytometry analysis of mitochondrial status of *Atg16*cKO neonatal enterocytes. Graphs show
relative mean fluorescence intensity (MFI) for Mitotracker Green (mitochondrial mass, MM), Mitotracker

Red CMXROS (mitochondrial membrane potential, MMP) and CM-H$_2$DCFDA (ROS) in ileal EpCAM$^+$CD24$^-$

cKit$^-$ cells from WT and *Atg16*cKO mice at P14. See Figure S3 for gating strategy. Dots represent individual

mice (MM, n = 23,17; MMP, n = 10,9; ROS, n = 19,13).

In (C), dot plots represent mean ± SEM (black lines). In (D), bar chart show mean + SEM.

Statistics analyses: Mann–Whitney test (C, D [MM]), Student's *t*-test (D: MMP, ROS). *$p < 0.05$, **$p < 0.01$,

***$p < 0.001$, ****$p < 0.0001$.

See also Figure S3 and Table S1.

Figure 5. Autophagy Deficiency Causes Premature Metabolic Transition in LREs

(A) RT-qPCR analysis of transcription factor (*Mafb*, *c-maf*, *Blimp1*), neonatal (*Ass1*, *Glb1*) and adult (*Treh*,

Sis, *Arg2*) enzyme genes in ileal epithelial cells from WT and *Atg16*cKO mice at late neonatal stage (P14)

and from WT mice at post-weaning stage (P21–30). Sample sizes: *Mafb* (n = 8, 9, 3), *c-maf* (n = 8, 11, 3),

Blimp1 (n = 11, 9, 4), neonatal enzymes (n = 6, 5, 4), *Treh* (n = 5, 3, 4), *Sis* (n = 5, 5, 4), *Arg2* (n = 5, 6, 4).

(B) Confocal images of ileal sections from WT and *Atg16*cKO mice at P14 (n = 5), and from post-weaning

(P21–30) WT mice (n = 2–3), labelled with anti-MAFB (green, top) or anti-BLIMP1 (green, bottom), and with

phalloidin for F-actin (magenta) and DAPI (blue). Arrowheads point to nuclear localization of MAFB and

BLIMP1 in WT LREs.

(C) Violin plot of nuclear MAFB (n = 20, 21, 11) and BLIMP1 (n = 21, 16, 9) intensity (integrated density) in

(B), normalised to epithelial nuclear area.

(D) Confocal images of ileal sections from WT and *Atg16*cKO mice at P14, labelled with anti-ASS1 (green,

top) (n = 3) or anti-TREH (green, bottom) (n = 5), anti-E-cadherin (Ecad, magenta), and DAPI (blue).

(E) Violin plots of ASS1 (top; n = 12, 10) and TREH (bottom; n = 15,16) intensity (integrated density) in (D),

normalised to epithelial area.

(F) Western blot of TREH protein from ileal epithelial cells of WT and *Atg16*cKO mice at P14. USP7, loading

control.

(G) Bar chart quantifying TREH protein levels from (F) (n = 5, 4).

(H) Met-Flow profiling of ileal EpCAM$^+$CD24$^-$ cells from WT (n = 10–13) and *Atg16*cKO (n = 7–9) mice at

P14, and from post-weaning (P21–30) WT mice (n = 8). Bubble colour and size reflect group means for

each marker. See Figure S3 for the gating strategy and Table S2 for statistical significance.

In (B, D), dashed boxes indicate magnified areas; L, lumen. Scale bars: 10 μm (main); 5 μm (insets).

In (A, C, E, G, H), data relative to WT P14 values. In (A, G), graphs show mean + SEM and dots represent individual mice. In (C, E), violin plots depict the median (thick line) and IQR (thin lines) and dots are individual images (2–5 images per mouse, 2–5 mice per group).

Statistical analyses: one-way ANOVA (A, C [MAFB], H), Kruskal–Wallis test (C: BLIMP1), Mann–Whitney test (E), Student's t test (G). $*p < 0.05$, $**p < 0.01$, $***p < 0.001$, $****p < 0.0001$.

See also Figures S3 and S5, and Table S2.

Figure 6. *Tfam*cKO Neonates Recapitulate Endolysosomal and Metabolic Defects Observed in *Atg16*cKO Enterocytes, and Exhibit Weight Loss and Early Lethality

(A) Representative images of WT and *Tfam*cKO mice at postnatal day 14 (P14).

(B) Kaplan–Meier survival curves of WT and *Tfam*cKO mice. Animals were euthanized if they failed to gain weight for two consecutive days or exhibited marked weight loss beginning at P9.

(C) Relative body weight of WT and *Tfam*cKO mice at P8, P10, P12, and P14, expressed as percentage. Sample sizes are indicated at the bottom of the graph.

(D) RT-qPCR analysis of *Tfam* mRNA in ileal epithelial cells from WT and *Tfam*cKO mice at P10 and P14 (n = 12,12,12,10).

(E) *Tfam*cKO neonatal enterocytes show damaged mitochondria. TEM micrographs of ileal sections from WT (n = 3) and *Tfam*cKO (n = 4) mice at P10 and P14. Asterisks indicate damaged mitochondria. Scale bar, 1 µm.

(F) Quantification of mitochondrial classes in (E), as defined in Figure 4B. Dots represent individual enterocytes. Sample sizes are indicated in Table S1.

(G) Flow cytometry analysis of mitochondrial status of *Tfam*cKO neonatal enterocytes. Bar charts show relative MFI for Mitotracker Green (mitochondrial mass, MM), Mitotracker Red CMXROS (mitochondrial membrane potential, MMP) and CM-H_2DCFDA (ROS) in ileal EpCAM$^+$CD24$^-$cKit$^-$ cells from WT and *Tfam*cKO mice at P10 and P14. Sample sizes: MM (n = 5, 5, 8, 14), MMP (n = 3, 4, 10, 13), ROS (n = 9, 6, 10, 9). See Figure S3 for gating strategy.

(H) Confocal images of ileal sections from WT and *Tfam*cKO mice at P14 (n = 3–4), stained with anti-MAFB (green, top) or anti-BLIMP1 (green, bottom), phalloidin (magenta), and DAPI (blue). Arrowheads mark nuclear localization of MAFB and BLIMP1 in WT LREs.

(I) Violin plots quantifying nuclear MAFB (top; n = 15, 10) and BLIMP1 (bottom; n = 16, 16) intensity in (H), normalised to epithelial nuclear area.

(J) Confocal images of ileal sections from WT and *Tfamc*KO mice at P14 (n = 3–4). Top: labelled with anti-ASS1 (green), anti-E-cadherin (Ecad, magenta), and DAPI (blue). Bottom: labelled with anti-TREH (green) and anti-Ecad (magenta).

(K) Violin plots quantifying ASS1 (top; n = 14, 16) and apical TREH (bottom; n = 9, 10) intensity in (J), normalised to epithelial area.

(L) Met-Flow profiling of ileal EpCAM$^+$CD24$^-$ cells from WT (n = 6) and *Tfamc*KO (n = 5) mice at P14. Bubble colour and size reflect group means for each marker. See Figure S3 for the gating strategy and Table S2 for statistical significance.

(M) Confocal images of ileal sections from WT (n = 4) and *Tfamc*KO (n = 3) mice at P14 gavaged with rhodamine B–dextran (Rd-dext, red) 3 hours before sacrifice. Sections were stained with anti-LAMP1 (green), anti-β-catenin (white), and DAPI (blue). Arrow: endolysosomal system in WT LREs; arrowhead: reduced LAMP1$^+$ vesicles in *Tfamc*KO enterocytes. Images deconvolved with Huygens software.

(N) Violin plot quantifying Rd-dextran uptake (integrated density) in (M), normalised to epithelial area (n = 11, 7).

(O) Quantification of lysosomal occupation and content in (M). Dot plots show the LAMP1$^+$ area as a percentage of cell area (n = 167, 145) and LAMP1 intensity (integrated density) per enterocyte (n = 148, 139).

In (D–O), mice aged P9–11 and P13–15 were grouped as "P10" and "P14," respectively.

In (H, J, M), dashed boxes indicate magnified areas; L, lumen. Scale bars: 10 µm (main); 5 µm (insets).

In (C, D, G, I, K, L, N), data relative to WT values. In (C, D, G), graphs depict mean + SEM and dots represent individual mice. In (F, O), dot plots show mean ± SEM (red lines) and dots represent individual enterocytes. In (I, K, N), violin plots show median (thick line) and IQR (thin lines) and dots represent individual images (2–4 per mouse).

Statistical analyses: Gehan–Breslow–Wilcoxon test (B), Student's *t*-test (C, D, F [P14 Class III only], G, K, L, N), Mann–Whitney test (F [except P14 Class III], I, O). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

See also Figures S3, S6 and S7, and Tables S1 and S2.

Figure 7. High Antioxidant Activity in LREs Regulates the Timing of Metabolic Transition

A) Bar chart showing the relative mean fluorescence intensity (MFI) of MitoSOX (mitochondrial ROS) and CM-H$_2$DCFDA (total ROS) in ileal EpCAM$^+$CD24$^-$ cells from WT mice at P7 (LREs) and P21–30 (PECs) (n = 4). Data relative to LREs values. See Figure S3 for gating strategy.

(B) Bubble plot and list of ROS-related GO terms from over-representation analysis (ORA) of DEGs between LREs and PECs. The x-axis represents the z-score, the y-axis shows the significance as –

log$_{10}$(adjusted p-value), bubble size reflects the number of associated genes, and colors indicate GO

categories (BP, biological process; CC, cellular component; MF, molecular function).

(C) GSEA enrichment plot of antioxidant activity gene set (LREs vs PECs).

(D) RT-qPCR analysis of *Gpx3* and *Cat* genes in ileal epithelial cells from WT mice at the indicated time

points (n = 3). Values are expressed as log$_{10}$ fold change relative to E18.5 embryonic levels.

(E) RT-qPCR analysis of *Gpx3* and *Cat* genes in ileal epithelial cells from WT and *Atg16*cKO mice at P14

(n = 6).

(F) RT-qPCR analysis of *Gpx3* (n = 7, 5) and *Cat* (n = 5, 7) in ileal epithelial cells from WT and *Tfam*cKO

mice at P14.

(G) Confocal images of ileal sections from WT and *Atg16*cKO mice at P14 (n = 3), stained with anti-GPX3

(green), anti-E-cadherin (Ecad, magenta), and DAPI (blue).

(H) Violin plot of GPX3 intensity from (G), normalised to epithelial area (n = 12, 9).

(I) Confocal images of ileal sections from WT and *Tfam*cKO mice at P14 (n = 4), labelled as in (F).

(J) Violin plot of GPX3 intensity from (H), normalised to epithelial area (n = 16).

(K) RT-qPCR analysis of transcription factor (*Mafb*, *c-maf*, *Blimp1*), neonatal (*Ass1*, *Glb1*) and adult (*Treh*,

Arg2) enzyme genes in ileal epithelial cells from control and H$_2$O$_2$ treated mice at P12. Sample sizes: *Mafb*

(n = 7, 6), *c-maf* (n = 7, 6), *Blimp1* (n = 7, 7), neonatal enzymes (n = 7, 7), *Treh* (n = 5, 6), *Arg2* (n = 6, 6).

(L) Confocal images of ileal sections from control (n = 4) and H$_2$O$_2$-treated (n = 4–5) mice at P12, stained

with anti-BLIMP1 (green, left) or anti-ASS1 (green, centre) or anti-TREH (green, right), and DAPI (blue). L,

lumen. Scale bars, 10 µm.

(M) Violin plots of BLIMP1, ASS1 and TREH intensity (integrated density) from (L). BLIMP1 data were

normalised to epithelial nuclear area; ASS1 and TREH data, to epithelial area. Sample sizes: BLIMP1 (n =

12, 15), ASS1 (n = 12, 16), TREH (n = 13, 12).

(N) Flow cytometry analysis of BLIMP1 (n = 7, 7) and ASS1 (n = 6, 7) in control and H$_2$O$_2$-treated mice at

P12. Bar charts show relative median fluorescent intensity of BLIMP1 and ASS1 in ileal EpCAM$^+$CD24$^-$cKit$^-$

cells. See Figure S3 for gating strategy.

(O) Working model: In *Atg16*cKO and *Tfam*cKO neonatal enterocytes, mitochondrial dysfunction and

reduced antioxidant responses elevate ROS levels, which repress *Blimp1* and *Mafb* expression and drive

premature metabolic transition. Enterocyte's design inspired by Fujita et al. (1990). Figure created with

BioRender.com.

In (G, I), dashed boxes indicate magnified areas; L, lumen. Scale bars: 10 µm (main); 5 µm (insets).

In (E, F, H, J), data relative to WT values; in (K, M, N), relative to control values. In (A, E, F, K, N), graphs

represent mean + SEM and dots represent individual mice; in (C), mean + SD. In (H, J, M), violin plots show

median (thick line) and IQR (thin lines) and dots represent individual images (2–4 per mouse, 3–5 mice per

group).

Statistical analyses: Student's t test (A, E, F, H, K, M [ASS1], N), one-way ANOVA (D), Mann–Whitney test

(J, M [BLIMP1, TREH]). $^*p < 0.05$, $^{**}p < 0.01$, $^{***}p < 0.001$, $^{****}p < 0.0001$.

See also Figure S3.

SUPPLEMENTAL INFORMATION

Figure S1. Characterization of LREs in the Neonatal Mouse Ileum

Related to Figures 1 and 2

(**A**) Representative TEM image of LREs from the distal small intestine of a P7 mice (n = 3). The profile of

an LRE is outlined by a dashed line. Dashed boxes indicate magnified areas. N, nucleus; LV, lysosomal

vacuole; AEC, apical endosomal complex; e, endosome; mvb, multivesicular body; L, lumen. Scale bars:

5 µm (main); 2 µm (insets).

(**B**) Confocal microscopy images of ileal sections from P7 mice (n = 3), stained with anti-LAMP1 (green),

anti-Dab2 (magenta), anti-β-catenin (white), and DAPI (cyan). Dashed boxes highlight magnified areas.

Arrows indicate the endolysosomal system; arrowheads point to apical Dab2 localization. Scale bars: 20 µm

(main); 10 µm (first magnification); 5 µm (second magnification).

(**C**) Western blot of mCherry and GFP-LC3 isoforms in ileal epithelial cells from GFP-LC3 mice at P7 and

P21, gavaged with mCherry 3 hours before sacrifice. USP7, loading control.

(**D**) Sequential fluorescent dextran uptake assay. P7 mice were gavaged with Alexa 647-dextran (FR-dext,

green), followed by rhodamine B–dextran (Rd-dext, red) 3 hours later. Mice were sacrificed at 4 and 8 hours

post-initial gavage to assess uptake via confocal microscopy (n = 3). Arrows point to lysosomal vacuoles

containing both dextrans at 8 hours. Arrowheads indicate endosomes. Scale bars: 5 µm (main); 2 µm

(insets).

1000

1001   **Figure S2. Phenotype of *Atg16*cKO Mice**

1002   *Related to Figures 1 and 2*

1003   (**A**) Relative body weight of WT and *Atg16*cKO mice at postnatal days P7, P14, and P30. Sample sizes are

1004   indicated in the graph.

1005   (**B**) Representative images of intestines from WT and *Atg16*cKO mice at the post-weaning stage (P30).

(C) Quantification of intestinal lengths shown in (B). Bar graphs show the relative lengths of the small intestine (SI) and the colon–rectum (CR) segments (n = 7, 9).

(D) Representative histological images of ileum and colon sections from WT and *Atg16*cKO post-weaning (P30) mice (n = 3). Ileal sections are stained with periodic acid-Schiff (PAS) and haematoxylin; colon sections are stained with haematoxylin and eosin (H&E). Asterisks denote oedema in *Atg16*cKO ileum; arrowheads indicate goblet cells. Scale bars: 200 µm (ileum), 400 µm (colon).

(E and F) Flow cytometry analysis of immune and epithelial cell populations in the ileum of WT and *Atg16*cKO mice at P14 (E) and P30 (F). Left: B lymphocytes (LB) and T-helper lymphocytes (LTh) percentages. Centre: Innate immune cells (IIC) and intestinal epithelial cells (IEC) percentages. Right: Enterocyte and enterocytic progenitors (EC), goblet cell (GC), Paneth cell (PC), and enteroendocrine cell (EEC) percentages. See Figure S3 for gating strategy. (E): Left, n = 17,8–11; Centre, n = 22,16–17; Right, n = 15,12–13. (F): n = 6–7, 9–10.

(G) RT-qPCR analysis of *Dab2* (n = 10,6,6,5), *Mcoln3* (n = 9,8,6,5), and *Lamp2* (n = 10,6,6,6) in epithelial cells from the proximal small intestine of WT and *Atg16*cKO mice at P7 and P14.

(H) RT-qPCR analysis of *Amn* (n = 6,5), *Mamdc4* (n = 9,10), and *Ctsl1* (n = 9,11) in ileal epithelial cells from WT and *Atg16*cKO mice at P14.

In (A, C, G, H), graphs represent mean + SEM. In (E, F), violin plots depict median (thick line) and interquartile range (IQR, thin lines). In all graphs, data relative to WT values, dots represent individual mice and Student's t test was conducted. *p < 0.05, **p < 0.01, ***p < 0.001, ****p < 0.0001.

Figure S3. Gating Strategies Used for Cell Population Characterization by Flow Cytometry
Related to Figures 4–7, S2 and S4–S6

Biopool is the combination of immune cell markers anti-CD11c, anti-CD3e, anti-CD11b, anti-CD19, and anti-CD45.2. The strategy for selecting IEC subtypes is based on Asano et al. (2017). GD, ghost dye; LTh, T-helper lymphocytes; LB, B lymphocytes; IIC, innate immune cells; IEC, intestinal epithelial cells; EEC, enteroendocrine cells; PC, Paneth cells; EC, enterocytes and enterocytic progenitors; GC, goblet cells.

Figure S4. RNA-seq Analysis Comparing Gene Expression of LREs and PECs
Related to Figure 3

(A) FACS gating strategy used to isolate LREs and PECs from GFP-LC3 mice at P7 (gavaged with Rd-dextran, 16 hours) and P42, respectively, for RNA-seq analysis. Single-living EpCAM⁺CD24⁻Biopool⁻ cells were gated. For LREs, Dextran⁺GFP^high cells were selected.

(B) Heatmap with hierarchical clustering of differentially expressed genes (DEGs) with q-value < 0.05 reveals distinct expression patterns between LREs and PECs. Each row represents a single gene and each

column represents a condition sample. Each sub-column represents an individual sample (3 samples per
group). Gene expression intensities were $\log_2$-transformed and visualized as a colour gradient from yellow
to light blue, representing the shift from downregulation to upregulation, as indicated in the key.

(C) Principal component analysis (PCA) plotting for 2 groups based on normalised overall transcriptomic
reads. The PCA figure represents a two-dimensional scatterplot of the first two principal components of the
RNA-Seq data. Analyses in panels (B) and (C) show that replicates of the same cell population are more
similar to each other than to replicates from a different population.

(**D**) Bubble plot of selected Kyoto Encyclopaedia of Genes and Genomes (KEGG) pathways from over-
representation analysis (ORA) of the differential gene expression data from the LREs vs PECs. The x-axis
represents the z-score, the y-axis shows the significance as –log10(adjusted p-value), bubble size reflects
the number of associated genes, and colours indicate KEGG pathway categories.

(E) RT-qPCR validation of *Ctsl1* expression in LREs and PECs from WT mice at P7 (gavaged with Rd-
dextran, 16 hours) and P42, respectively. FACS was used for cell sorting, selecting single-living
EpCAM$^+$CD24$^-$Biopool$^-$ cells, with Dextran$^+$ cells further gated for LREs. In LREs, each dot represents one
of three pooled samples (n = 3), each composed of multiple mice; in PECs, each dot corresponds to an
individual mouse (n = 3). Data relative to LREs values.

(F) Analysis of lysosomal content in LREs and PECs by measuring Lysotracker MFI by flow cytometry. Bar
chart depicts Lysotracker MFI of LREs and PECs from WT mice at P7 (gavaged with mCherry, 16 hours)
and P42 respectively. Single-living EpCAM$^+$CD24$^-$Biopool$^-$ cells were gated. For LREs, mCherry$^+$ cells were
selected. See Figure S3 for gating strategy. Dots represent individual mice (n = 3).

Graphs in (E, F) represent mean + SEM. In (E), Student's t test; in (F), Mann–Whitney test. *p < 0.05, **p
< 0.01.

1062

1063

Figure S5. Metabolic Transition in the Small Gut During Postnatal Development
Related to Figure 5
(**A**) Expression and statistical data of selected genes encoding transcription factors (TFs: *Mafb, c-maf,*
Blimp1), neonatal enzymes (*Ass1, Glb1*), and adult enzymes (*Treh, Sis, Arg2*) from the LREs vs PECs
RNA-seq analysis.

(B and C) RT-qPCR analysis of *Mafb* and *Blimp1* (B), and of *Ass1, Glb1, Treh,* and *Sis* (C) genes in
epithelial cells isolated from proximal and distal small intestinal segments of WT mice at the indicated
postnatal time points (n = 3). Data are normalised to the expression levels in the proximal small intestine of
E18.5 embryos and expressed as $\log_{10}$fold change ($\log_{10}$FC).

(D) Schematic representation of the cellular metabolic pathways. Enzymes used in the Met-flow panel are
shown in red. Scheme adapted from Iborra et al. (2024) (CC BY 4.0).

(E) Met-Flow analysis of ileal EpCAM$^+$CD24$^-$ cells from WT mice at P7 (LREs) and P42 (PECs) (n = 2–4).
Data relative to LREs values. Bubble colour and size reflect group means for each marker. See Figure S3
for the gating strategy and Table S2 for statistical significance.

In (B, C), graphs represent mean + SD, two-way ANOVA was conducted and only the most relevant
statistically significant comparisons are shown. In (E), Student's *t*-test. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$,
****$p < 0.0001$.

1081

Figure S6. Validation and Intestinal Phenotype of *Tfamc*KO Neonatal Mice
Related to Figure 6

(A) Relative mitochondrial DNA (mtDNA) levels of *Nd1*, *Co1*, *Co2*, and *Atp6* in ileal epithelial cells from WT
and *Tfamc*KO mice at P10 (n = 2) and P14 (n = 4), measured by qPCR.

(B) RT-qPCR analysis of mitochondrial genes *mt-Nd1*, *mt-Co2*, and *Sdha* in ileal epithelial cells from WT
and *Tfamc*KO mice at P10 (n = 5–6).

(C) Confocal images of ileal sections from WT (n = 3) and *Tfamc*KO (n = 2) mice at P14, labelled with anti-
MTCO1 (green) and anti-β-catenin (magenta). Arrows point to mitochondria in WT LREs.

(D) Violin plot shows MTCO1 intensity (integrated density) from (C), normalised to epithelial area. Dots
represent individual images (n = 9,7; 3–4 images per mouse).

(E) PAS and haematoxylin staining of ileal sections from WT and *Tfamc*KO mice at P10 and P14 (n = 4–6)
shows inflamed and disorganized epithelium in *Tfamc*KO mice. Phenotypes include shortened villi (double
red arrows), thickened submucosa and lower mucosa (double black arrows), oedema (asterisks), reduced
PAS+ goblet cells (arrowheads), and abnormal vacuoles in enterocytes (arrows).
Scale bars: 200 μm (main); 100 μm (insets).

(F–H) Quantification of villi length (F), percentage of villi with goblet cells (G), and the maximum thickness
of submucosa and lower mucosa (H) from (E). In (F), each dot represents a villus (n = 50,60,50,50; 10 villi
per mouse). For (G–H), n = 4–6 mice.

(I–K) Flow cytometry analysis of innate immune cells (IIC, I), B lymphocytes (LB, J), and T-helper cells (LT$_h$,
K) percentages in the ileal of WT and *Tfamc*KO mice at P10 and P14 (n = 3–5). See Figure S3 for gating
strategy.

(L–O) Confocal imaging and quantification of stem and Paneth cells in WT and *Tfamc*KO ileal at P10 and
P14:

• (**L**) Olfm4 (green) labels crypt stem cells; arrows point to Olfm4$^+$ crypts.

- (**M**) Quantification of Olfm4$^+$ crypts (30-100 crypts/mouse).

- (**N**) Lysozyme (green) marks Paneth cells; arrows show normal crypt localization, arrowheads indicate mislocalised Paneth cells in villi.

- (**O**) Quantification Paneth cells/mm² of epithelial area.

For (L, N), sections are also stained with anti-Ecad (white) and DAPI (blue). Scale bars: 200 μm (main); 50 μm (insets). For all panels, n = 3 mice.

(**P**) Distribution of mitochondrial integrity classes in *Tfam*cKO neonatal enterocytes. Quantification of the percentage, area and aspect ratio (long axis / short axis) of each mitochondria class in Figure 6E, as defined in Figure 4B. In percentage graph, dots represent individual enterocytes; in area and aspect ratio graphs, dots are individual mitochondria. Sample sizes are indicated in Table S1.

In (C, E, L, N), dashed boxes indicate magnified areas; L, lumen.

In (A, B, D, I–K), data relative to WT values. In (A, G–K, M, O), graphs represent mean + SEM and dots represent individual mice. In (B, D), violin plots show the median (thick line) and IQR (thin lines). In (F, P), dot plots show mean ± SEM (red lines).

Statistical analyses: Student's t test (A, B, I–K, M, O), Mann–Whitney test (D, P), two-way ANOVA (F, H), Kruskal–Wallis test (G). In (F–H), only statistically significant comparisons between WT and *Tfam*cKO are indicated within each time group. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Figure S7. *Tfam*cKO Neonatal Enterocytes Exhibit a Decreased Endolysosomal System and Undergo an Early Metabolic Switch
Related to Figure 6

(**A**) RT-qPCR analysis of transcription factor (*Mafb*, *c-maf*, *Blimp1*), neonatal (*Ass1*, *Glb1*) and adult (*Treh*, *Sis*, *Arg2*) enzyme genes in ileal epithelial cells from WT and *Tfam*cKO mice at P10 and P14.

(**B**) RT-qPCR analysis of *Dab2*, *Amn*, and *Mamdc4* genes in ileal epithelial cells from WT and *Tfam*cKO mice at P10 and P14.

(**C**) Confocal images of ileal sections from WT and *Tfam*cKO mice at P10 (n = 3) and P14 (n = 2–3), labelled with anti-Dab2 (green), anti-β-catenin (white), and DAPI (blue). Arrowheads indicate apical Dab2 localisation.

(**D**) Violin plot shows apical Dab2 intensity (integrated density) from (C), normalised to epithelial area and relative to WT values. Dots represent individual images (n = 6, 6, 6, 4; 2 images per mouse).

(**E**) RT-qPCR analysis of *Lamp2*, *Mcoln3*, and *Ctsl1* in ileal epithelial cells from WT and *Tfam*cKO mice at P10 and P14.

(F) Confocal images of ileal sections from WT and *Tfam*cKO mice at P10 (n = 3), labelled with anti-LAMP1 (green), anti-β-catenin (white), and DAPI (blue). Arrows indicate endolysosomal system in WT cells; arrowheads highlight the reduced lysosomal vesicles in *Tfam*cKO enterocytes.

(G) Quantification of lysosomal occupation and content in (E). Dot plots show the LAMP1$^+$ area as a percentage of cell area (n = 157,99) and LAMP1 intensity (integrated density) per enterocyte (n = 117,74). Dots represent individual enterocytes.

In (C, F), dashed boxes indicate magnified areas; L, lumen. Scale bars: 10 µm (main); 5 µm (insets).

In (A, B, E), data relative to WT-P10 values, dots represent individual mice (n = 4–7), and graphs represent mean + SEM. In (D), violin plot shows the median (thick line) and IQR (thin lines). In (G), dot plots depict mean ± SEM (red lines).

Statistical analyses: Kruskal–Wallis test (A [*Treh*], B [*Dab2*], E [*Lamp2*]), two-way ANOVA (A, B, E; except *Treh*, *Dab2*, *Lamp2*), Student's t test (D), Mann–Whitney test (G). In (A, B, E), only statistically significant comparisons between WT and *Tfam*cKO are indicated within each time group. *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$.

Table S1. Sample sizes in TEM graphs
Related to Figures 4, 6 and S6
Values in the number (No.) and percentage columns correspond to enterocytes; in the area and aspect ratio columns, values correspond to individual mitochondria. For more detailed information on sample size of the analyses, see the TEM section in the Methods.

Table S2. Statistical Significance of Met-Flow Experiments
Related to Figures 5, 6 and S5
For Figure 5H, statistical results are shown only for comparisons between WT and the two other groups. No statistical test was applied for HK1 of Figure S5E (n = 2). *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$. NS, not significant.

REFERENCES

Ahl, P.J., Hopkins, R.A., Xiang, W.W., Au, B., Kaliaperumal, N., Fairhurst, A.-M., Connolly, J.E., 2020. Met-Flow, a strategy for single-cell metabolic analysis highlights dynamic changes in immune subpopulations. Commun. Biol. 3, 305. https://doi.org/10.1038/s42003-020-1027-9

Asano, J., Sato, T., Ichinose, S., Kajita, M., Onai, N., Shimizu, S., Ohteki, T., 2017. Intrinsic Autophagy Is Required for the Maintenance of Intestinal Stem Cells and for Irradiation-Induced Intestinal Regeneration. Cell Rep. 20, 1050–1060. https://doi.org/10.1016/j.celrep.2017.07.019

Baixauli, F., Acín-Pérez, R., Villarroya-Beltrí, C., Mazzeo, C., Nuñez-Andrade, N., Gabandé-Rodriguez, E.,
Ledesma, M.D., Blázquez, A., Martin, M.A., Falcón-Pérez, J.M., Redondo, J.M., Enríquez, J.A.,
Mittelbrunn, M., 2015. Mitochondrial Respiration Controls Lysosomal Function during Inflammatory
T Cell Responses. Cell Metab. 22, 485–498. https://doi.org/10.1016/j.cmet.2015.07.020

Bara, A.M., Chen, L., Ma, C., Underwood, J., Moreci, R.S., Sumigray, K., Sun, T., Diao, Y., Verzi, M.,
Lechler, T., 2022. Maf family transcription factors are required for nutrient uptake in the mouse
neonatal gut. Dev. Camb. Engl. 149, dev201251. https://doi.org/10.1242/dev.201251

Becker, K., Pons-Kühnemann, J., Fechner, A., Funk, M., Gromer, S., Gross, H.-J., Grünert, A., Schirmer,
R.H., 2005. Effects of antioxidants on glutathione levels and clinical recovery from the malnutrition
syndrome kwashiorkor--a pilot study. Redox Rep. Commun. Free Radic. Res. 10, 215–226.
https://doi.org/10.1179/135100005X70161

Beumer, J., Clevers, H., 2021. Cell fate specification and differentiation in the adult mammalian intestine.
Nat. Rev. Mol. Cell Biol. 22, 39–53. https://doi.org/10.1038/s41580-020-0278-0

Bhutta, Z.A., Berkley, J.A., Bandsma, R.H.J., Kerac, M., Trehan, I., Briend, A., 2017. Severe childhood
malnutrition. Nat. Rev. Dis. Primer 3, 17067. https://doi.org/10.1038/nrdp.2017.67

Birchenough, G.M.H., Schroeder, B.O., Sharba, S., Arike, L., Recktenwald, C.V., Puértolas-Balint, F.,
Subramani, M.V., Hansson, K.T., Yilmaz, B., Lindén, S.K., Bäckhed, F., Hansson, G.C., 2023.
Muc2-dependent microbial colonization of the jejunal mucus layer is diet sensitive and confers local
resistance to enteric pathogen infection. Cell Rep. 42, 112084.
https://doi.org/10.1016/j.celrep.2023.112084

Birgisdottir, Å.B., Johansen, T., 2020. Autophagy and endocytosis – interconnections and
interdependencies. J. Cell Sci. 133, jcs228114. https://doi.org/10.1242/jcs.228114

Boyle, E.I., Weng, S., Gollub, J., Jin, H., Botstein, D., Cherry, J.M., Sherlock, G., 2004. GO::TermFinder—
open source software for accessing Gene Ontology information and finding significantly enriched
Gene Ontology terms associated with a list of genes. Bioinformatics 20, 3710–3715.
https://doi.org/10.1093/bioinformatics/bth456

Cadwell, K., Liu, J.Y., Brown, S.L., Miyoshi, H., Loh, J., Lennerz, J.K., Kishi, C., Kc, W., Carrero, J.A., Hunt,
S., Stone, C.D., Brunt, E.M., Xavier, R.J., Sleckman, B.P., Li, E., Mizushima, N., Stappenbeck,
T.S., Virgin IV, H.W., 2008. A key role for autophagy and the autophagy gene Atg16l1 in mouse
and human intestinal Paneth cells. Nature 456, 259–263. https://doi.org/10.1038/nature07416

Campbell, C.T., Kolesar, J.E., Kaufman, B.A., 2012. Mitochondrial transcription factor A regulates
mitochondrial transcription initiation, DNA packaging, and genome copy number. Biochim. Biophys.
Acta 1819, 921–929. https://doi.org/10.1016/j.bbagrm.2012.03.002

Chen, P.-M., Lin, C.-H., Li, N.-T., Wu, Y.-M., Lin, M.-T., Hung, S.-C., Yen, M.-L., 2015. c-Maf regulates
pluripotency genes, proliferation/self-renewal, and lineage commitment in ROS-mediated
senescence of human mesenchymal stem cells. Oncotarget 6, 35404–35418.
https://doi.org/10.18632/oncotarget.6178

Choo, J., Glisovic, N., Matic Vignjevic, D., 2022. Gut homeostasis at a glance. J. Cell Sci. 135.
https://doi.org/10.1242/jcs.260248

Chung, K.W., Dhillon, P., Huang, S., Sheng, X., Shrestha, R., Qiu, C., Kaufman, B.A., Park, J., Pei, L.,
Baur, J., Palmer, M., Susztak, K., 2019. Mitochondrial Damage and Activation of the STING
Pathway Lead to Renal Inflammation and Fibrosis. Cell Metab. 30, 784-799.e5.
https://doi.org/10.1016/j.cmet.2019.08.003

Cosovanu, C., Resch, P., Jordan, S., Lehmann, A., Ralser, M., Farztdinov, V., Spranger, J., Mülleder, M.,
Brachs, S., Neumann, C., 2022. Intestinal epithelial c-Maf expression determines enterocyte
differentiation and nutrient uptake in mice. J. Exp. Med. 219, e20220233.
https://doi.org/10.1084/jem.20220233

Cox, C.M., Lu, R., Salcin, K., Wilson, J.M., 2018. The Endosomal Protein Endotubin Is Required for Enterocyte Differentiation. Cell. Mol. Gastroenterol. Hepatol. 5, 145–156. https://doi.org/10.1016/j.jcmgh.2017.11.001

el Marjou, F., Janssen, K.-P., Chang, B.H.-J., Li, M., Hindie, V., Chan, L., Louvard, D., Chambon, P., Metzger, D., Robine, S., 2004. Tissue-specific and inducible Cre-mediated recombination in the gut epithelium. Genes. N. Y. N 2000 39, 186–193. https://doi.org/10.1002/gene.20042

Foerster, E.G., Mukherjee, T., Cabral-Fernandes, L., Rocha, J.D.B., Girardin, S.E., Philpott, D.J., 2022. How autophagy controls the intestinal epithelial barrier. Autophagy 18, 86–103. https://doi.org/10.1080/15548627.2021.1909406

Fujita, M., Reinhart, F., Neutra, M., 1990. Convergence of apical and basolateral endocytic pathways at apical late endosomes in absorptive cells of suckling rat ileum in vivo. J. Cell Sci. 97, 385–394. https://doi.org/10.1242/jcs.97.2.385

Fujita, N., Itoh, T., Omori, H., Fukuda, M., Noda, T., Yoshimori, T., 2008. The Atg16L complex specifies the site of LC3 lipidation for membrane biogenesis in autophagy. Mol. Biol. Cell 19, 2092–2100. https://doi.org/10.1091/mbc.e07-12-1257

Gong, G., Song, M., Csordas, G., Kelly, D.P., Matkovich, S.J., Dorn, G.W., 2015. Parkin-mediated mitophagy directs perinatal cardiac metabolic maturation in mice. Science 350, aad2459. https://doi.org/10.1126/science.aad2459

Gonnella, P.A., Neutra, M.R., 1984. Membrane-bound and fluid-phase macromolecules enter separate prelysosomal compartments in absorptive cells of suckling rat ileum. J. Cell Biol. 99, 909–917. https://doi.org/10.1083/jcb.99.3.909

Hansson, J., Panchaud, A., Favre, L., Bosco, N., Mansourian, R., Benyacoub, J., Blum, S., Jensen, O.N., Kussmann, M., 2011. Time-resolved Quantitative Proteome Analysis of *In Vivo* Intestinal Development. Mol. Cell. Proteomics 10, M110.005231. https://doi.org/10.1074/mcp.M110.005231

Harper, J., Mould, A., Andrews, R.M., Bikoff, E.K., Robertson, E.J., 2011. The transcriptional repressor Blimp1/Prdm1 regulates postnatal reprogramming of intestinal enterocytes. Proc. Natl. Acad. Sci. U. S. A. https://doi.org/10.1073/pnas.1105852108

He, C., 2022. Balancing nutrient and energy demand and supply via autophagy. Curr. Biol. 32, R684–R696. https://doi.org/10.1016/j.cub.2022.04.071

Henning, S.J., 1985. Ontogeny of Enzymes in the Small Intestine. Annu. Rev. Physiol. 47, 231–245. https://doi.org/10.1146/annurev.ph.47.030185.001311

Henning, S.J., 1981. Postnatal development: coordination of feeding, digestion, and metabolism. Am. J. Physiol.-Gastrointest. Liver Physiol. https://doi.org/10.1152/ajpgi.1981.241.3.G199

Hodzic, Z., Bolock, A.M., Good, M., 2017. The Role of Mucosal Immunity in the Pathogenesis of Necrotizing Enterocolitis. Front. Pediatr. 5. https://doi.org/10.3389/fped.2017.00040

Hulzebos, C.V., Sauer, P.J.J., 2007. Energy requirements. Semin. Fetal. Neonatal Med., Nutrition 12, 2–10. https://doi.org/10.1016/j.siny.2006.10.008

Hwang, S., Maloney, N.S., Bruinsma, M.W., Goel, G., Duan, E., Zhang, L., Shrestha, B., Diamond, M.S., Dani, A., Sosnovtsev, S.V., Green, K.Y., Lopez-Otin, C., Xavier, R.J., Thackray, L.B., Virgin, H.W., 2012. Nondegradative role of Atg5-Atg12/ Atg16L1 autophagy protein complex in antiviral activity of interferon gamma. Cell Host Microbe 11, 397–409. https://doi.org/10.1016/j.chom.2012.03.002

Iborra-Pernichi, M., Ruiz García, J., Velasco de la Esperanza, M., Estrada, B.S., Bovolenta, E.R., Cifuentes, C., Prieto Carro, C., González Martínez, T., García-Consuegra, J., Rey-Stolle, M.F., Rupérez, F.J., Guerra Rodriguez, M., Argüello, R.J., Cogliati, S., Martín-Belmonte, F., Martínez-Martín, N., 2024. Defective mitochondria remodelling in B cells leads to an aged immune response. Nat. Commun. 15, 2569. https://doi.org/10.1038/s41467-024-46763-1

Jiménez-Loygorri, J.I., Villarejo-Zori, B., Viedma-Poyatos, Á., Zapata-Muñoz, J., Benítez-Fernández, R., Frutos-Lisón, M.D., Tomás-Barberán, F.A., Espín, J.C., Area-Gómez, E., Gomez-Duran, A., Boya,

P., 2024. Mitophagy curtails cytosolic mtDNA-dependent activation of cGAS/STING inflammation
during aging. Nat. Commun. 15, 830. https://doi.org/10.1038/s41467-024-45044-1

Kaur, J., Debnath, J., 2015. Autophagy at the crossroads of catabolism and anabolism. Nat. Rev. Mol. Cell
Biol. 16, 461–472. https://doi.org/10.1038/nrm4024

Kawamura, N., Sun-Wada, G.-H., Aoyama, M., Harada, A., Takasuga, S., Sasaki, T., Wada, Y., 2012.
Delivery of endosomes to lysosomes via microautophagy in the visceral endoderm of mouse
embryos. Nat. Commun. 3, 1071. https://doi.org/10.1038/ncomms2069

Kleinman, R.E., Walker, W.A., 1984. Antigen processing and uptake from the intestinal tract. Clin. Rev.
Allergy 2, 25–37. https://doi.org/10.1007/BF02991209

Kulkarni, D.H., Newberry, R.D., 2019. Intestinal Macromolecular Transport Supporting Adaptive Immunity.
Cell. Mol. Gastroenterol. Hepatol. 7, 729–737. https://doi.org/10.1016/j.jcmgh.2019.01.003

Kuma, A., Hatano, M., Matsui, M., Yamamoto, A., Nakaya, H., Yoshimori, T., Ohsumi, Y., Tokuhisa, T.,
Mizushima, N., 2004. The role of autophagy during the early neonatal starvation period. Nature
432, 1032–1036. https://doi.org/10.1038/nature03029

Larabi, A., Barnich, N., Nguyen, H.T.T., 2020. New insights into the interplay between autophagy, gut
microbiota and inflammatory responses in IBD. Autophagy 16, 38–51.
https://doi.org/10.1080/15548627.2019.1635384

Larsson, N.-G., Wang, J., Wilhelmsson, H., Oldfors, A., Rustin, P., Lewandoski, M., Barsh, G.S., Clayton,
D.A., 1998. Mitochondrial transcription factor A is necessary for mtDNA maintance and
embryogenesis in mice. Nat. Genet. 18, 231–236. https://doi.org/10.1038/ng0398-231

Lennicke, C., Cochemé, H.M., 2021. Redox metabolism: ROS as specific molecular regulators of cell
signaling and function. Mol. Cell 81, 3691–3707. https://doi.org/10.1016/j.molcel.2021.08.018

Li, A., Gao, M., Liu, B., Qin, Y., Chen, L., Liu, H., Wu, H., Gong, G., 2022. Mitochondrial autophagy:
molecular mechanisms and implications for cardiovascular disease. Cell Death Dis. 13, 444.
https://doi.org/10.1038/s41419-022-04906-6

Liang, H., Wu, S., Zhang, Z., Khan, M.Z., Zhan, Y., Zhu, M., Wang, S., Liu, W., Wang, C., Cao, G., Han,
Y., 2024. Establishment of a mouse model of ovarian oxidative stress induced by hydrogen
peroxide. Front. Vet. Sci. 11. https://doi.org/10.3389/fvets.2024.1484388

Massaquoi, M.S., Guillemin, K., 2019. Packing in the Proteins. Dev. Cell 51, 1–2.
https://doi.org/10.1016/j.devcel.2019.09.013

McWilliams, T.G., Prescott, A.R., Allen, G.F.G., Tamjar, J., Munson, M.J., Thomson, C., Muqit, M.M.K.,
Ganley, I.G., 2016. mito-QC illuminates mitophagy and mitochondrial architecture in vivo. J. Cell
Biol. 214, 333–345. https://doi.org/10.1083/jcb.201603039

Mizushima, N., Kuma, A., Kobayashi, Y., Yamamoto, A., Matsubae, M., Takao, T., Natsume, T., Ohsumi,
Y., Yoshimori, T., 2003. Mouse Apg16L, a novel WD-repeat protein, targets to the autophagic
isolation membrane with the Apg12-Apg5 conjugate. J. Cell Sci. 116, 1679–1688.
https://doi.org/10.1242/jcs.00381

Mizushima, N., Yamamoto, A., Matsui, M., Yoshimori, T., Ohsumi, Y., 2004. In vivo analysis of autophagy
in response to nutrient starvation using transgenic mice expressing a fluorescent autophagosome
marker. Mol. Biol. Cell 15, 1101–1111. https://doi.org/10.1091/mbc.e03-09-0704

Mizushima, N., Yoshimorim, T., Levine, B., 2010. Methods in Mammalian Autophagy Research. Cell 140,
313–326. https://doi.org/10.1016/j.cell.2010.01.028

Morris, O., Jasper, H., 2021. Reactive Oxygen Species in intestinal stem cell metabolism, fate and function.
Free Radic. Biol. Med. 166, 140–146. https://doi.org/10.1016/j.freeradbiomed.2021.02.015

Moschandrea, C., Kondylis, V., Evangelakos, I., Herholz, M., Schneider, F., Schmidt, C., Yang, M., Ehret,
S., Heine, M., Jaeckstein, M.Y., Szczepanowska, K., Schwarzer, R., Baumann, L., Bock, T.,
Nikitopoulou, E., Brodesser, S., Krüger, M., Frezza, C., Heeren, J., Trifunovic, A., Pasparakis, M.,

2023. Mitochondrial dysfunction abrogates dietary lipid processing in enterocytes. Nature 1–8. https://doi.org/10.1038/s41586-023-06857-0

Muncan, V., Heijmans, J., Krasinski, S.D., Büller, N.V., Wildenberg, M.E., Meisner, S., Radonjic, M., Stapleton, K.A., Lamers, W.H., Biemond, I., Van Den Bergh Weerman, M.A., O'Carroll, D., Hardwick, J.C., Hommes, D.W., Van Den Brink, G.R., 2011. Blimp1 regulates the transition of neonatal to adult intestinal epithelium. Nat. Commun. https://doi.org/10.1038/ncomms1463

Murphy, M.P., 2008. How mitochondria produce reactive oxygen species. Biochem. J. 417, 1–13. https://doi.org/10.1042/BJ20081386

Ni, H.-M., Bockus, A., Wozniak, A.L., Jones, K., Weinman, S., Yin, X.-M., Ding, W.-X., 2011. Dissecting the dynamic turnover of GFP-LC3 in the autolysosome. Autophagy 7, 188–204. https://doi.org/10.4161/auto.7.2.14181

Park, J., Levic, D.S., Sumigray, K.D., Bagwell, J., Eroglu, O., Block, C.L., Eroglu, C., Barry, R., Lickwar, C.R., Rawls, J.F., Watts, S.A., Lechler, T., Bagnat, M., 2019. Lysosome-Rich Enterocytes Mediate Protein Absorption in the Vertebrate Gut. Dev. Cell. https://doi.org/10.1016/j.devcel.2019.08.001

Picca, A., Faitg, J., Auwerx, J., Ferrucci, L., D'Amico, D., 2023. Mitophagy in human health, ageing and disease. Nat. Metab. 5, 2047–2061. https://doi.org/10.1038/s42255-023-00930-8

Popov, L.-D., 2023. Mitochondria as intracellular signalling organelles. An update. Cell. Signal. 109, 110794. https://doi.org/10.1016/j.cellsig.2023.110794

Qi, T., Sun, M., Zhang, C., Chen, P., Xiao, C., Chang, X., 2020. Ascorbic Acid Promotes Plasma Cell Differentiation through Enhancing TET2/3-Mediated DNA Demethylation. Cell Rep. 33. https://doi.org/10.1016/j.celrep.2020.108452

Rath, E., Moschetta, A., Haller, D., 2018. Mitochondrial function — gatekeeper of intestinal epithelial cell homeostasis. Nat. Rev. Gastroenterol. Hepatol. 15, 497–516. https://doi.org/10.1038/s41575-018-0021-x

Remis, N.N., Wiwatpanit, T., Castiglioni, A.J., Flores, E.N., Cantú, J.A., García-Añoveros, J., 2014. Mucolipin Co-deficiency Causes Accelerated Endolysosomal Vacuolation of Enterocytes and Failure-to-Thrive from Birth to Weaning. PLoS Genet. 10. https://doi.org/10.1371/journal.pgen.1004833

Rodríguez-Colman, M.J., Schewe, M., Meerlo, M., Stigter, E., Gerrits, J., Pras-Raves, M., Sacchetti, A., Hornsveld, M., Oost, K.C., Snippert, H.J., Verhoeven-Duif, N., Fodde, R., Burgering, B.M.T., 2017. Interplay between metabolic identities in the intestinal crypt supports stem cell function. Nature 543, 424–427. https://doi.org/10.1038/nature21673

Rodríguez-Fraticelli, A.E., Bagwell, J., Bosch-Fortea, M., Boncompain, G., Reglero-Real, N., García-León, M.J., Andrés, G., Toribio, M.L., Alonso, M.A., Millán, J., Perez, F., Bagnat, M., Martín-Belmonte, F., 2015. Developmental regulation of apical endocytosis controls epithelial patterning in vertebrate tubular organs. Nat. Cell Biol. 17, 241–250. https://doi.org/10.1038/ncb3106

Rombout, J.H., Lamers, C.H., Helfrich, M.H., Dekker, A., Taverne-Thiele, J.J., 1985. Uptake and transport of intact macromolecules in the intestinal epithelium of carp (Cyprinus carpio L.) and the possible immunological implications. Cell Tissue Res. 239, 519–530. https://doi.org/10.1007/BF00219230

Schneider, C.A., Rasband, W.S., Eliceiri, K.W., 2012. NIH Image to ImageJ: 25 years of image analysis. Nat. Methods 9, 671–675. https://doi.org/10.1038/nmeth.2089

Sena, L.A., Chandel, N.S., 2012. Physiological roles of mitochondrial reactive oxygen species. Mol. Cell 48, 158–167. https://doi.org/10.1016/j.molcel.2012.09.025

Sies, H., Berndt, C., Jones, D.P., 2017. Oxidative Stress. Annu. Rev. Biochem. 86, 715–748. https://doi.org/10.1146/annurev-biochem-061516-045037

Skrzypek, T., Valverde Piedra, J.L., Skrzypek, H., Kazimierczak, W., Biernat, M., Zabielski, R., 2007. Gradual disappearance of vacuolated enterocytes in the small intestine of neonatal piglets. J. Physiol. Pharmacol. Off. J. Pol. Physiol. Soc. 58 Suppl 3, 87–95.

Spence, J.R., Lauf, R., Shroyer, N.F., 2011. Vertebrate intestinal endoderm development. Dev. Dyn. https://doi.org/10.1002/dvdy.22540

Srivillibhuthur, M., Warder, B.N., Toke, N.H., Shah, P.P., Feng, Q., Gao, N., Bonder, E.M., Verzi, M.P., 2018. TFAM is required for maturation of the fetal and adult intestinal epithelium. Dev. Biol. 439, 92–101. https://doi.org/10.1016/j.ydbio.2018.04.015

Stirling, D.R., Swain-Bowden, M.J., Lucas, A.M., Carpenter, A.E., Cimini, B.A., Goodman, A., 2021. CellProfiler 4: improvements in speed, utility and usability. BMC Bioinformatics 22, 433. https://doi.org/10.1186/s12859-021-04344-9

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette, M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., Mesirov, J.P., 2005. Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide expression profiles. Proc. Natl. Acad. Sci. 102, 15545–15550. https://doi.org/10.1073/pnas.0506580102

Trehan, I., Manary, M.J., 2015. Management of severe acute malnutrition in low-income and middle-income countries. https://doi.org/10.1136/archdischild-2014-306026

Treuting, P.M., Arends, M.J., Dintzis, S.M., 2018. 11 - Upper Gastrointestinal Tract, in: Treuting, P.M., Dintzis, S.M., Montine, K.S. (Eds.), Comparative Anatomy and Histology (Second Edition). Academic Press, San Diego, pp. 191–211. https://doi.org/10.1016/B978-0-12-802900-8.00011-7

Walker, W.A., 2013. Initial Intestinal Colonization in the Human Infant and Immune Homeostasis. Ann. Nutr. Metab. 63, 8–15. https://doi.org/10.1159/000354907

Wallaeys, C., Garcia-Gonzalez, N., Libert, C., 2023. Paneth cells as the cornerstones of intestinal and organismal health: a primer. EMBO Mol. Med. 15, e16427. https://doi.org/10.15252/emmm.202216427

Walter, W., Sánchez-Cabo, F., Ricote, M., 2015. GOplot: an R package for visually combining expression data with functional analysis. Bioinformatics 31, 2912–2914. https://doi.org/10.1093/bioinformatics/btv300

Wang, S., Long, H., Hou, L., Feng, B., Ma, Z., Wu, Y., Zeng, Y., Cai, J., Zhang, D., Zhao, G., 2023. The mitophagy pathway and its implications in human diseases. Signal Transduct. Target. Ther. 8, 304. https://doi.org/10.1038/s41392-023-01503-7

Wilson, J.M., Whitney, J.A., Neutra, M.R., 1991. Biogenesis of the apical endosome–lysosome complex during differentiation of absorptive epithelial cells in rat ileum. J. Cell Sci. 100, 133–143. https://doi.org/10.1242/jcs.100.1.133

Yang, Y., Chen, X., Liao, X., Jiang, W., Zhou, Y., Sun, Y., Zheng, B., 2024. Identification of MAP1LC3A as a promising mitophagy-related gene in polycystic ovary syndrome. Sci. Rep. 14, 16982. https://doi.org/10.1038/s41598-024-67969-9

Yim, W.W.-Y., Mizushima, N., 2020. Lysosome biology in autophagy. Cell Discov. 6, 6. https://doi.org/10.1038/s41421-020-0141-7

Yu, G., Wang, L.-G., Han, Y., He, Q.-Y., 2012. clusterProfiler: an R Package for Comparing Biological Themes Among Gene Clusters. OMICS J. Integr. Biol. 16, 284–287. https://doi.org/10.1089/omi.2011.0118

Zemirli, N., Morel, E., Molino, D., 2018. Mitochondrial Dynamics in Basal and Stressful Conditions. Int. J. Mol. Sci. 19, 564. https://doi.org/10.3390/ijms19020564

Zhao, Y.G., Codogno, P., Zhang, H., 2021. Machinery, regulation and pathophysiological implications of autophagosome maturation. Nat. Rev. Mol. Cell Biol. 22, 733–750. https://doi.org/10.1038/s41580-021-00392-4

Zhong, Y., Xia, S., Wang, G., Liu, Q., Ma, F., Yu, Y., Zhang, Y., Qian, L., Hu, L., Xie, J., 2024. The interplay between mitophagy and mitochondrial ROS in acute lung injury. Mitochondrion 78, 101920. https://doi.org/10.1016/j.mito.2024.101920

1410

METHODS

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Mouse monoclonal anti-GFP, HRP-conjugated (clone GG4-2C2.12.10)	Miltenyi Biotec	130-091-833
Rat monoclonal anti-RFP (clone 5F8)	Chromotek	5F8
Rabbit polyclonal anti-USP7	Bethyl	A300-033A
Rabbit polyclonal anti-ATG16L1	MBL	PM040
Mouse monoclonal anti-LC3C (clone E5Q2K)	CST	83506
Rabbit polyclonal anti-Actin	Sigma	A2103
Mouse monoclonal anti-SQSTM1 (clone 2C11)	Abnova	H00008878-M01
Mouse monoclonal anti-Dab2 (clone 52/p96)	BD Biosciences	610464
Rat monoclonal anti-LAMP1 (clone 1D4B)	DSHB	1D4B
Rabbit polyclonal anti-Trehalase	GeneTex	GTX109979
Rabbit polyclonal anti-β-catenin	Santa Cruz Biotechnology	sc-7199
Rabbit polyclonal anti-Parkin	Abcam	ab15494
Rabbit polyclonal anti-MafB	Novus Biologicals	NBP1-81342
Rabbit polyclonal anti-Blimp1	Invitrogen	PA5-20310
Mouse monoclonal anti-E-Cadherin (clone 36/E)	BD Biosciences	610181
Goat polyclonal anti-GPX3	R&D Systems	AF4199
Mouse monoclonal anti-MTCO1 (clone 1D6E1A8)	Thermo Fisher Scientific	459600
Rabbit monoclonal anti-Olfm4 (clone D6Y5A)	CST	39141
Rabbit polyclonal anti-Lysozyme	Dako	A0099
Alexa Fluor 647 rabbit monoclonal anti-ASS1 (clone EPR12398)	Abcam	ab311083
Rat monoclonal anti-CD16/32 (clone 93)	Biolegend	101330
PerCP-eFluor710 rat monoclonal anti-EpCAM (clone G8.8)	Invitrogen	46-5791-82
APC rat monoclonal anti-EpCAM (clone G8.8)	Biolegend	118214
BV605 rat monoclonal anti-EpCAM (clone G8.8)	Biolegend	118227
Pacific Blue rat monoclonal anti-CD24 (Clone M1/69)	Biolegend	101820
PE rat monoclonal anti-c-Kit (clone 2B8)	Biolegend	105807
BV785 rat monoclonal anti-c-Kit (clone 2B8)	Biolegend	105841
Alexa Fluor 700 rat monoclonal anti-CD11b (clone M1/70)	Biolegend	101222
Biotin rat monoclonal anti-CD11c (clone N418)	Biolegend	105841
APC-Fire810 rat monoclonal anti-B220 (clone RA3-6B2)	Biolegend	103278
APC-Fire810 rat monoclonal anti-CD4 (clone GK1.5)	Biolegend	100480
RY586 rat monoclonal anti-CD4 (clone GK1.5)	BD Biosciences	568162
Biotin rat monoclonal anti-CD19 (clone 6D5)	Biolegend	115504
Biotin rat monoclonal anti-CD3e (clone 145-2C11)	Biolegend	79751
Biotin rat monoclonal anti-CD45.2 (clone 104)	Biolegend	109804
Rabbit polyclonal anti-GLS	Thermo Fisher Scientific	PA5-35365
Alexa Fluor 647 rabbit monoclonal anti-HK1 (clone EPR10134(B))	Abcam	ab197864

Alexa Fluor 488 mouse monoclonal anti-CPT1A (clone 8F6AE9)	Abcam	ab171449
Rabbit monoclonal anti-ACAC (clone EPR23235-147)	Abcam	ab269273
Alexa Fluor 488 rabbit monoclonal anti-CS (clone EPR8067)	Abcam	ab197488
PE rabbit monoclonal anti-IDH2 (clone EPR7577)	Abcam	ab212122
Alexa Fluor 594 rabbit monoclonal anti-ATP5A (clone EPR23235- 147)	Abcam	ab216385
BB700 rat monoclonal anti-BLIMP1 (clone 5E7)	BD Biosciences	567830
Alexa Fluor 488 donkey polyclonal anti-mouse IgGs	Thermo Fisher Scientific	A-21202
Alexa Fluor 488 donkey polyclonal anti-rabbit IgGs	Thermo Fisher Scientific	A-21206
Alexa Fluor 488 donkey polyclonal anti-rat IgGs	Thermo Fisher Scientific	A-21208
Alexa Fluor 555 donkey polyclonal anti-mouse IgGs	Thermo Fisher Scientific	A-31570
Alexa Fluor 647 donkey polyclonal anti-rabbit IgGs	Thermo Fisher Scientific	A-31573
Alexa Fluor 647 donkey polyclonal anti-mouse IgGs	Thermo Fisher Scientific	A-31571
Peroxidase AffiniPure donkey polyclonal Anti-Mouse IgG	Jackson Immunoresearch	715-035-151
HRP donkey anti-rabbit IgG	Cytiva	NA934
Peroxidase AffiniPure goat polyclonal Anti-Rat IgG	Jackson Immunoresearch	112-035-003
Chemicals, peptides, and recombinant proteins		
Tamoxifen	Merck	T5648
Leupeptin	Sigma	L2884
Hydrogen Peroxide 30%	Merck	108597
Dextran rhodamine B, 10000MW, neutral	Invitrogen	D1824
Dextran Alexa Fluor 647, 10000MW, anionic, fixable	Invitrogen	D22914
Paraformaldehyde, powder, TECHNICAL	VWR Chemicals	28794.295
Tissue-Tek O.C.T. Compound	Sakura	4583
Sucrose	Merck	1076511000
DAPI	Merck	268298
Phalloidin Alexa Fluor™ 488	ThermoFisher	A12379
Phalloidin Alexa Fluor™ 647	ThermoFisher	A22287
ProLong™ Glass Antifade Mountant	Thermo Fisher	P36984
Ghost Dye Violet 540	TONBO Biosciences	13-0879-T100
Ghost Dye Red 780	TONBO Biosciences	13-0865-T100
Streptavidin-BV421	Biolegend	405225
CM-H2DCFDA	Invitrogen	C6827
MitoTracker Green FM	Invitrogen	M7514
MitoTracker Red CMXRos	Invitrogen	M7512
MitoSOX	Invitrogen	M36008
LysoTracker Green DND-26	Invitrogen	L7526
eBioscience FOXP3/Transcription Factor Staining Buffer Set	Invitrogen	00-5523-00
ECL Western Blotting Reagents	Cytiva	RPN2106
DTT	Thermo Scientific	R0861
Oligonucleotides		
Genotyping primers for $Atg16l1^{fl/fl}$: forward: 5' CTTTCTCAACAGAACCAGCAGTAC 3'	IDT	N/A
Genotyping primers for $Atg16l1^{fl/fl}$: reverse: 5' GTTCAGGTCAATCACCAACTGAGC 3'	IDT	N/A

Genotyping primers for *Tfam*^{fl/fl}: forward: 5' CTGCCTTCCTCTAGCCCGGG 3'	IDT	N/A
Genotyping primers for *Tfam*^{fl/fl}: reverse: 5' GTAACAGCAGACAACTTGTG 3'	IDT	N/A
Genotyping primers for *Tfam*^{fl/fl}: reverse-2: 5' CTCTGAAGCACATGGTCAAT- 3'	IDT	N/A
Genotyping primers for *Villin-creER*^{T2}: forward: 5'- CAAGCCTGGCTCGACGGCC -3'	Sigma	N/A
Genotyping primers for *Villin-creER*^{T2}: reverse: 5'-CGCGAACATCTTCAGGTTCT-3'	Sigma	N/A
Genotyping primers for *mito-QC*: forward: 5'- CAAAGACCCCAACGAGAAGC -3'	IDT	N/A
Genotyping primers for *mito-QC*: forward-2: 5'- CCCAAGGCACACAAAAAACC -3'	IDT	N/A
Genotyping primers for *mito-QC*: reverse: 5'-CTCTTCCCTCGTGACCTGCAACTCC-3'	IDT	N/A
Genotyping primers for *mito-QC*: reverse-2: 5'- CATGTCTTTAATCTACCTCGATGG -3'	IDT	N/A
Genotyping primers for *GFP-LC3*: forward: 5'-ATAACTTGCTGGCCTTTCCACT -3'	IDT	N/A
Genotyping primers for *GFP-LC3*: reverse: 5'- CGGGCCATTTACCGTAAGTTAT-3'	IDT	N/A
Genotyping primers for *GFP-LC3*: reverse-2: 5'-GCAGCTCATTGCTGTTCCTCAA-3'	IDT	N/A
RT-qPCR primers for *Atg16l1*: forward: 5' ACATGATGGTGCGTGGAATG 3'	Sigma	N/A
RT-qPCR primers for *Atg16l1*: reverse: 5' TCCTTGTCCTTCTGCTGCAT 3'	Sigma	N/A
RT-qPCR primers for *Tfam*: forward: 5' CAGGAGGCAAAGGATGATTC 3'	Sigma	N/A
RT-qPCR primers for *Tfam*: reverse: 5' CCAAGACTTCATTTCATTGTCG 3'	Sigma	N/A
RT-qPCR primers for *Dab2*: forward: 5' CGAAAGCGCCATCAAAGAAG 3'	IDT	N/A
RT-qPCR primers for *Dab2*: reverse: 5' CCTTTGAACCTGGCCAACAA 3'	IDT	N/A
RT-qPCR primers for *Mcoln3*: forward: 5' ATGGAGTTCATCAACGGGTG 3'	IDT	N/A
RT-qPCR primers for *Mcoln3*: reverse: 5' ATAGTTGACGTCCCGAGAAG 3'	IDT	N/A
RT-qPCR primers for *Lamp2*: forward: 5' TGCAGAATGGGAGATGAATTT 3'	IDT	N/A
RT-qPCR primers for *Lamp2*: reverse: 5' GGCACTATTCCGGTCATCC 3'	IDT	N/A
RT-qPCR primers for *Ctsl1*: forward: 5' CCAGCTACACAACGGGGAAT 3'	IDT	N/A
RT-qPCR primers for *Ctsl1*: reverse: 5' AGCATCAGCGGTTCCTGAAA 3'	IDT	N/A
RT-qPCR primers for *Bnip3*: forward: 5' TGTTGGATGCCCAGCATGAA 3'	IDT	N/A
RT-qPCR primers for *Bnip3*: reverse: 5' AATCTTCCTCAGACAGAGTGCT 3'	IDT	N/A
RT-qPCR primers for *Prkn*: forward: 5' AAACCGGATGAGTGGTGAGT 3'	IDT	N/A

RT-qPCR primers for *Prkn*: reverse: 5' AGCTACCGACGTGTCCTTGT 3'	IDT	N/A
RT-qPCR primers for *Pink1*: forward: 5' GCTGATCGAGGAGAAGCAG 3'	IDT	N/A
RT-qPCR primers for *Pink1*: reverse: 5' GATAATCCTCCAGACGGAAGC 3'	IDT	N/A
RT-qPCR primers for *Mafb*: forward: 5' AGAGGCGGACCCTGAAAAAC 3'	IDT	N/A
RT-qPCR primers for *Mafb*: reverse: 5' GTTTTCTCGGAAGCCGTTGC 3'	IDT	N/A
RT-qPCR primers for *c-maf*: forward: 5' GCATGCTGGACATGTATGGT 3'	IDT	N/A
RT-qPCR primers for *c-maf*: reverse: 5' ATGTACAACGGGAGGCTGAA 3'	IDT	N/A
RT-qPCR primers for *Blimp1*: forward: 5' TTCTTGTGTGGTATTGTCGGGACTT 3'	Sigma	N/A
RT-qPCR primers for *Blimp1*: reverse: 5' TTGGGGACACTCTTTGGGTAGAGTT 3'	Sigma	N/A
RT-qPCR primers for *Ass1*: forward: 5' CATTGGAATGAAGTCCCGAG 3'	IDT	N/A
RT-qPCR primers for *Ass1*: reverse: 5' GATTTTGCGTACTTCCCGAT 3'	IDT	N/A
RT-qPCR primers for *Glb1*: forward: 5' ACTGCTGCAACTGCTGGG 3'	Sigma	N/A
RT-qPCR primers for *Glb1*: reverse: 5' AATGGCTGTCCATCCTTGAG 3'	Sigma	N/A
RT-qPCR primers for *Treh*: forward: 5' GCACCTGCTGCTTCTGCT 3'	Sigma	N/A
RT-qPCR primers for *Treh*: reverse: 5' CATCTGAACTTGGTGCAGGA 3'	Sigma	N/A
RT-qPCR primers for *Sis*: forward: 5' GCCCACGCGTTTGGTGAGAC 3'	Sigma	N/A
RT-qPCR primers for *Sis*: reverse: 5' ATCAGCAGTGCAGGGCCCCA 3'	Sigma	N/A
RT-qPCR primers for *Arg2*: forward: 5' TGATTGGCAAAAGGCAGAGG 3'	Sigma	N/A
RT-qPCR primers for *Arg2*: reverse: 5' CTAGGAGTAGGAAGGTGGTC 3'	Sigma	N/A
RT-qPCR primers for *Gpx3*: forward: 5' AATTAGGCACAAAGCCCCCA 3'	IDT	N/A
RT-qPCR primers for *Gpx3*: reverse: 5' CATTCTGGGCTTCCCTTCCA 3'	IDT	N/A
RT-qPCR primers for *Cat*: forward: 5' GAACGGCAATAGGGGTCCTC 3'	IDT	N/A
RT-qPCR primers for *Cat*: reverse: 5' TGGTACACGCAAAAGGAGCA 3'	IDT	N/A
RT-qPCR primers for *Amn*: forward: 5' ATCTGTGAGAGCCTCGGCG 3'	IDT	N/A
RT-qPCR primers for *Amn*: reverse: 5' GTTGGAGGCAGTATCGAAGGA 3'	IDT	N/A
RT-qPCR primers for *Mamdc4*: forward: 5' TCACCACTGCCAGAACAAGG 3'	IDT	N/A
RT-qPCR primers for *Mamdc4*: reverse: 5' CTAACTGGTTCCACGGTCCC 3'	IDT	N/A

RT-qPCR primers for *mt-Nd1*: forward: 5' TCCCCTACCAATACCACACC 3'	Sigma	N/A
RT-qPCR primers for *mt-Nd1*: reverse: 5' TGAAATTGTTTGGGCTACGG 3'	Sigma	N/A
RT-qPCR primers for *mt-Co2*: forward: 5' ACGAAATCAACAACCCCGTA 3'	Sigma	N/A
RT-qPCR primers for *mt-Co2*: reverse: 5' GGCAGAACGACTCGGTTATC 3'	Sigma	N/A
RT-qPCR primers for *Sdha*: forward: 5' GCGGCTTTCACTTCTCTGTT 3'	IDT	N/A
RT-qPCR primers for *Sdha*: reverse: 5' CTCAGAAAGGCCAAATGCAG 3'	IDT	N/A
qPCR primers for *mt-Co1*: forward: 5' ATCCCTTGACATCGTGCTTC 3'	Sigma	N/A
qPCR primers for *mt-Co1*: reverse: 5' AAGTGGGCTTTTGCTCATGT 3'	Sigma	N/A
qPCR primers for *mt-Atp6*: forward: 5' ACAGGCTTCCGACACAAACT 3'	Sigma	N/A
qPCR primers for *mt-Atp6*: reverse: 5' GGTAGCTGTTGGTGGGCTAA 3'	Sigma	N/A
qPCR primers for *Hk2*: forward: 5' GCCAGCCTCTCCTGATTTTAGTGT 3'	Sigma	N/A
qPCR primers for *Hk2*: reverse: 5' GGGAACACAAAAGACCTCTTCTGG 3'	Sigma	N/A
RT-qPCR primers for *Ppib*: forward: 5' CACCAATGGCTCACAGTTCTT 3'	IDT	N/A
RT-qPCR primers for *Ppib*: reverse: 5' ATGACATCCTTCAGTGGCTTG 3'	IDT	N/A

1413

1414

1415

Mice

1416

Atg16l1^{fl/fl} (Hwang et al., 2012), *Tfam*^{fl/fl} (Larsson et al., 1998), and GFP-LC3 (Mizushima et al., 2004) mice

were kindly provided by N. Martínez-Martín (Centro de Biología Molecular "Severo Ochoa", CBM, CSIC-

UAM, Madrid, Spain). The *mito*-QC mouse line (McWilliams et al., 2016) was obtained from P. Boya (Centro

de Investigaciones Biológicas Margarita Salas, CSIC, Madrid, Spain; University of Fribourg, Switzerland).

Villin-creER^{T2} mice (el Marjou et al., 2004) were provided by N. Djouder (Spanish National Cancer Research

Centre, CNIO, Madrid, Spain). *Atg16l1* ^{fl/fl}, *Tfam* ^{fl/fl} mice were crossed at the CBM animal facility with *Villin-*

creER^{T2} mice to generate intestinal epithelial-specific conditional knockout (cKO) mice: *Atg16*cKO and

*Tfam*cKO, respectively. In turn, *Atg16*cKO mice were crossed with *mito*-QC mice to establish *the mQ-*

*Atg16*cKO mouse line. Tamoxifen (0.12–0.5 mg in corn oil containing 10% ethanol) was administered

subcutaneously at postnatal days 1 and 2 to pups from *Atg16*cKO, *mQ-Atg16*cKO, and *Tfam*cKO litters,

following a protocol similar to that described by Gong et al. (2015). Cre-positive animals were classified as

KO, and Cre-negative mice as WT. Neonatal mice were humanely sacrificed by decapitation; adult mice

were euthanized via CO_2 inhalation. *Tfam*cKO neonates were sacrificed after two consecutive days without

weight gain or upon observing approximately a 10% weight loss, in accordance with humane endpoint

criteria. No test was performed to determine the influence of sex, as no apparent differences were found.

Mouse colonies were maintained at CBM under specific pathogen-free conditions and on a C57BL/6 genetic

background. Mice were housed four/five per cage with food and water available ad libitum and maintained

in a temperature-controlled environment on a 12 h–12 h light–dark cycle with light onset at 08:00. Animal

experiments were performed under protocols approved by the CBM and Comunidad Autónoma de Madrid

ethical committees under the following protocol approval number (PROEX 307.2/21). All animal procedures

conformed to EU Directive 86/609/EEC and Recommendation 2007/526/EC regarding the protection of

animals used for experimental and other scientific purposes, enforced in Spanish law under Real Decreto

1201/2005.

1440

mCherry / fluorescent-dextran uptake assay

Mice from GFP-LC3, *Atg16*cKO, and *Tfam*cKO lines were fasted for 1 hour and orally gavaged with either

mCherry or 10 kDa fluorescent-dextran (0.3–1 mg/ml in PBS; 10 µl/g body weight). Following gavage, mice

were sacrificed either 3 or 16 hours later, adapted from the protocol reported by Remis et al. (2014). To

test the endocytic function in neonatal enterocytes, WT mice at postnatal day 7 (P7) were sequentially

gavaged with two distinct 10 kDa fluorescent-dextrans (0.7 mg/ml in PBS; 10 µl/g body weight), following a

modified version of the protocol established in mouse embryos by Kawamura et al. (2012). After a 1-hour

fasting period, pups were first gavaged with Alexa Fluor 647-dextran and then returned to their mother.

Three hours later, they were given rhodamine B–dextran again. Pups were subsequently sacrificed either

1 or 5 hours after the second gavage, and ileal enterocytes were analysed by confocal microscopy to assess

dextran uptake and intracellular localization.

1452

Leupeptin and hydrogen peroxide treatment

1454   To inhibit lysosomal proteases and block autophagy, mice were injected subcutaneously with 40 mg/Kg

1455   leupeptin (Sigma) dissolved in PBS and sacrificed after 16h (Jiménez-Loygorri et al., 2024). To assess the

1456   effect of ROS on intestinal epithelial maturation, mice were injected intraperitoneally with hydrogen peroxide

1457   (5.1 µmol/g body weight; Merck) once daily from postnatal day (P)5 to P12 (day of sacrifice), following a

1458   protocol adapted from Liang et al. (2024).

1459

Histology and Immunofluorescence

Portions of the distal small intestine from mice were fixed in 4% PFA for 16 hours at 4 °C, embedded in

paraffin resin, and cut transversely into 5-µm sections. Some samples were stained with hematoxylin-eosin

(HE&E; Sigma and Merck, respectively), while others underwent Periodic Acid-Schiff (PAS) staining.

Additionally, some samples were used for immunofluorescence. For immunofluorescence, sections were

deparaffinized using xylene and then rehydrated through a series of decreasing ethanol concentrations

(100%, 96%, 70%, 50%). For antigen retrieval, tissue was boiled at 100 °C on a heat block in sodium citrate

buffer (10 mM Sodium citrate, 0.05% Tween 20, pH 6.0) for 20 min. Next, the slides were washed with

water and PBS containing calcium and magnesium, and then incubated for 1 hour at room temperature

with 3% bovine serum albumin (BSA) as a blocking solution. For O.C.T. embedded sections, portions of

the distal small intestine were previously fixed in 4% PFA for 16 hours at 4 °C, incubated in 30% sucrose

in PBS for 48-72 hours at 4 °C, and then embedded in O.C.T. compound and stored at -80 °C until use.

O.C.T. blocks were cut transversely into 15 µm sections using Superfrost Plus Adhesion Microscope Slides

(Epredia) and stored at -80 °C. For immunofluorescence, sections were dried at room temperature for 30

minutes, washed with 0.1% Tween 20, and blocked for 1 hour at room temperature with BGT buffer (0.3%

BSA, 0.75% Glycine, 0.3% Triton X-100 in PBS). For both paraffin-embedded and O.C.T.-embedded

samples, the primary antibody was added and incubated overnight at 4 °C in the respective blocking buffer.

After washing with PBS, the secondary antibody was incubated with DAPI for 1h at RT. Lastly, slides were

assembled with coverslips with Prolong Glass Antifade (Thermo Fisher). Antibodies used are specified in

the key resources table.

1480

Image Acquisition and Analysis

HE&E and PAS staining images were acquired using an Axio Imager M1 vertical wide-field microscope,

coupled with a Leica DMC6200 CMOS Colour camera. The objective used was 10X/0.3 Air EC Plan-Neo.

LASXv4.13 software was used to stitch the tiles into an image. The analysis of goblet cells, villi length, and

submucosa–lower mucosa thickness was performed using Fiji (ImageJ 1.54f; https://fiji.sc/) (Wayne

Rasband, NIH, USA) (Schneider et al., 2012). Lower mucosa was defined as the sum of the *muscularis*

mucosae, *lamina propria* and epithelial compartment of the crypts. For villi length analysis, the 10 longest

villi per section were chosen. One section per mouse was analysed from 3–6 mice per genotype. For

immunofluorescence, images were acquired using a Zeiss LSM 900 confocal microscope mounted on an

Axio Imager Z2 upright stand, with sequential acquisition and bidirectional scanning. For individual fields, a

63×/1.4 NA oil-immersion Plan-Apochromat objective was used, with a voxel size of 50 × 50 × 520 nm. For

tile scans, images were acquired using a 20×/0.8 NA air Plan-Apochromat objective, with a voxel size of

0.624 × 0.624 × 1.9 µm, and stitched using ZEN Blue software version 3.4 (Zeiss;

https://www.zeiss.com/zen). For the analysis of Paneth cells and Olfm4[+] crypts in the ileum of *Tfam*cKO

mice, all cells and all crypts were quantified per section (one section per mouse; 3 mice per genotype). For

intensity analysis, all images were acquired with 16-bit resolution and under the same conditions. An

average intensity projection was made, and the endothelial cells (lacteal vessel) were cropped. The same

threshold was applied to all images to select positive areas. Images were analysed using CellProfiler 4.2.8

(https://cellprofiler.org/) (Stirling et al., 2021) and Fiji (ImageJ 1.54f) (Schneider et al., 2012). Intensity

values correspond to the integrated density, normalized to the epithelial nuclear area (defined by DAPI in

epithelial cells) for BLIMP1 and MAFB, or to the epithelial area (delimited by β-catenin, E-cadherin, or F-

actin) for all other markers, except for LAMP1 (in both KO models) and Rd-dextran (only in *Atg16*cKO mice),

which were not normalized. For LAMP1 intensity and area analysis in *Tfam*cKO mice, images were

deconvolved using Huygens Professional software (version 24.10; Scientific Volume Imaging,

https://svi.nl/Huygens-Professional). For the analysis of Rd-dextran intensity and LAMP1 area and intensity

in *Atg16*cKO mice, 25–40 and 26–36 enterocytes per condition were analysed, respectively. For LAMP1

area and intensity analysis in *Tfamc*KO mice, a minimum of 30 enterocytes per mouse were quantified,

across 3 mice per group (99–167 enterocytes per group). For all other intensity analyses, 2–5 random

images per mouse were acquired from comparable villi, each containing approximately 50 cells, with 2–5

mice per group.

1511

Transmission Electron Microscopy

Samples were processed by the Electron Microscopy Unit at CBM. Distal small intestine portions were fixed

using 4% PFA (paraformaldehyde) and 2% glutaraldehyde in phosphate buffer 0.1M, pH 7.4, for 2 hours at

room temperature. Then they were washed 3 times for 10 minutes with phosphate buffer. After fixation, the

material was post-fixed in 1% osmium tetroxide and potassium ferricyanide for 1 hour at 4 °C, then

embedded in tannic acid for 1 minute and block-stained with 2% uranyl acetate for 1 hour in the dark. Next,

they were dehydrated in crescent concentrations of ethanol (30%, 50%, 70%, 95%, 3x100%; 10 minutes

each) and infiltrated in propylene oxide for 45 minutes and Epon 100% for 1 hour. Finally, the samples were

introduced into BEEM capsules (ESBE Scientific). Samples were observed using a JEM1400 Flash

Transmission Electron Microscope (JEOL) equipped with a CMOS OneView camera (Gatan). Fiji (ImageJ

1.54f) (Schneider et al., 2012) was used for image analysis. Mitochondria were classified into three

categories, as in Moschandrea et al. (2023): class I (normal), characterized by mitochondria with narrow

and well-defined cristae; class II (partly affected), with mitochondria exhibiting undefined cristae; and class

III (damaged), including mitochondria with wide, short, and less electron-dense cristae, or those with

ruptured membranes or internal inclusions. For the analysis of mitochondrial status in *Atg16*cKO mice, 291–

340 mitochondria were analysed from 10–11 enterocytes (3 mice per genotype). For *Tfamc*KO mice, in the

analysis of number and percentage of mitochondrial classes, 444–1040 mitochondria from 13–24

enterocytes (3–4 mice per group) were used; for area and aspect ratio analysis, 261–520 mitochondria

from 11–17 enterocytes (2–4 mice per group) were analysed. The graphs show data after outlier exclusion.

1531

Isolation of IEC

Mice were euthanized, and the abdominal cavity was opened to expose the intestine. The region of interest

(proximal or distal intestine) was dissected, removed, and rinsed in PBS. The tissue was cut transversely

into ~5 mm fragments, which were then opened longitudinally and incubated in epithelial dissociation buffer

(1 mM DTT, 1 mM EDTA, 2% FBS in PBS) at 37°C with agitation for 20 minutes. After incubation, the

samples were centrifuged at 1200 rpm for 5 minutes. The supernatant was discarded, and the pellet was

resuspended in 10 mL of DMEM supplemented with 10% FBS at 4 °C. The suspension was passed through

70 µm cell strainers (Falcon) and then centrifuged again at 1,200 rpm for 5 minutes. The resulting pellet

was resuspended in FACS buffer (1 mM EDTA, 2% FBS in PBS) at 4 °C and centrifuged again under the

same conditions. The final pellet containing epithelial cells was used for western blotting, flow cytometry,

and RT-qPCR analysis.

1543

Immunoblotting

1544
IECs pellets were lysed with lysis buffer (50 mM Tris-HCl, 150 mM NaCl, 1% NP-40, 50 mM NaF, 0.1%
SDS, 1 mM Na_3VO_4, 30 mM β-glycerophosphate, 5 mM EDTA, protease inhibitors) and centrifuged for 10
minutes at 4 °C, 15,000 rpm. Supernatants were collected and quantified using the Bradford protein assay
(Biorad Protein Assay, Biorad). 25 µg of protein was loaded in 10 or 12% acrylamide gels for SDS-PAGE.
After electrophoresis, proteins were transferred to an Immobilon-P transfer membrane (Millipore), blocked
for 1 hour, and incubated overnight with primary antibodies at 4 °C. After washing with 0.05% Tween 20,
they were incubated with HRP-conjugated secondary antibodies for 1 hour at room temperature. ECL
western blotting reagents (Cytiva) were used for the chemiluminescence reaction. Bands were analysed
using Fiji (ImageJ 1.54f) (Schneider et al., 2012). Antibodies used are specified in the key resources table.
1554

Flow Cytometry

1555
Disaggregated epithelial cells were plated into a 96-well plate and incubated with FACS buffer with anti-
CD16/CD32 Fc block and viability markers (Ghost Dye) for 10 min at room temperature. After centrifugation,
cells were incubated for 20 min at 4ºC with the appropriate combination of the antibodies in FACS buffer to
discern between the different epithelial and immune populations. Following protocols adapted from
immunometabolic (Baixauli et al., 2015; Iborra-Pernichi et al., 2024) and intestinal studies (Asano et al.,
2017), we assessed mitochondrial mass and membrane potential, cellular and mitochondrial ROS levels,
and lysosomal content labelling cells with MitoTracker Green FM (50 nM), MitoTracker Red CMXRos (100
nM), CM-H2DCFDA (2.5 µM), MitoSOX (1 µM), and Lysotracker Green DND-26 (100 nM), respectively.
Probes were incubated for 15 min at room temperature and 15 min at 37 °C in DMEM with 5% FBS
(Lysotracker), without FBS (MitoTracker and CM-H2DCFDA), or in HBSS (MitoSOX). Antibodies used are
specified in the key resources table. Data were acquired using a Cytek Aurora full-spectrum cytometer at
the CBM, and analysed with FlowJo software (versions 10 and 10.8.1; BD Biosciences). See Figure S3 for
details on the gating strategies.
1569

Met-flow cytometry staining

1570
Based on recent works on immune cell metabolism (Ahl et al., 2020; Iborra-Pernichi et al., 2024), we
selected eight key metabolic enzymes to analyse their expression by flow cytometry in our epithelial cells:
ASS1, GLS, HK1, CPT1A, ACAC, CS, IDH2, and ATP5A. After the disaggregated epithelial cells were
incubated with surface antibodies according to the flow cytometry protocol, they were fixed and
permeabilized using the eBioscience Foxp3/Transcription Factor Staining Buffer Set (Invitrogen). Next, they
were washed in the provided wash buffer and blocked with 20% FBS in wash buffer and incubated for 2h
at room temperature with the appropriate antibodies to detect the rate-limiting enzymes from different
metabolic pathways. Lastly, they were rewashed with wash buffer. Antibodies used are specified in the key

resources table. Data were acquired using a Cytek Aurora full-spectrum cytometer at the CBM, and
analysed with FlowJo software (versions 10 and 10.8.1; BD Biosciences).

1581

RNA Sequencing and Bioinformatic Analyses

The FACS gating strategy used to isolate LREs and PECs for RNA sequencing (RNA-seq) is shown in
Figure S4A. Cell sorting was performed using a FACSAria Fusion cytometer (BD Biosciences). Each
population was collected in biological triplicate samples of 300,000–500,000 cells. For LREs, each replicate
consisted of pooled cells from multiple neonatal litters; for PECs, each replicate corresponded to a single
mouse at postnatal day 42 (P42). Cells were collected in RLT buffer (Qiagen), and total RNA was extracted
using the RNeasy Plus Micro Kit (Qiagen). Samples with an RNA integrity number (RIN) >7.0 were selected
for RNA-seq. Ultra-low input poly(A)-selected libraries (single-end) were prepared in triplicate and
sequenced using the Illumina HiSeq 2500 platform (50 bp, single-end reads) at the Centro de Análisis
Genómico (CNAG-CRG, Barcelona, Spain). Reads were aligned to the Mus musculus reference genome
(mm39) using HISAT2 (v2.1.0), and gene counts were obtained using HTSeq-count (v0.11.2). Differential
gene expression analysis was performed with DESeq2. Gene Set Enrichment Analysis (GSEA)
(Subramanian et al., 2005) was conducted using the clusterProfiler (Yu et al., 2012) and enrichplot
packages within the R environment. Predefined gene sets from the Gene Ontology (GO) database were
assessed for statistically significant enrichment based on a ranked list of gene expression values. The full
gene dataset, annotated with ENSEMBL identifiers and their associated statistical metrics (stat), was
utilized as input for the GSEA. Enrichment was quantified by normalized enrichment scores (NES), with
significance determined at an adjusted p-value threshold of < 0.05 following Benjamini-Hochberg
correction. Over-Representation Analysis (ORA) (Boyle et al., 2004) was performed in R using the
clusterProfiler package. Differentially expressed genes (DEGs) with an adjusted p-value < 0.05 and
mapped using ENTREZID identifiers were used. Enriched GO terms and KEGG pathways were also filtered
at an adjusted p-value < 0.05 (Benjamini-Hochberg correction). Selected GO and KEGG terms were
visualized using bubble plots generated in R with the GOplot package (GOBubble function) (Walter et al.,
2015). The biocomputational data analysis was carried out by the Biocomputational Analysis Service
(SABio) at the Centro de Biologia Molecular "Severo Ochoa" (CSIC-UAM).

1607

RNA Extraction, cDNA Synthesis, and RT-qPCR

Epithelial cells were lysed in 500 µL of TRIzol (Invitrogen) and stored at -80 °C until RNA extraction. RNA
was isolated using the RNeasy Micro Kit (Qiagen) according to the manufacturer's instructions. For cDNA
synthesis, 500 ng of RNA was reverse transcribed using the High-Capacity cDNA Reverse Transcription
Kit (Applied Biosystems). Gene expression was assessed by quantitative PCR (qPCR) using gene-specific
primer pairs (listed in Kay Resources Table). Reactions were performed in 384-well plates using either the
CFX384 Real-Time System (Bio-Rad) or the ABI 7900HT Fast Real-Time PCR System (Applied
Biosystems), with SYBR Green Master Mix (Agilent, Santa Clara, CA). Each well contained 10 ng of cDNA

and 2.5 µM of each primer in a final volume of 10 µL. All reactions were run in technical triplicates.
Cyclophilin B (*Ppib*) was used as the reference gene for normalisation.
1618

mtDNA Quantification
Genomic DNA (gDNA) was extracted using the NZY Tissue gDNA Isolation Kit (NZYTech). Relative
mitochondrial DNA (mtDNA) content was determined by qPCR using primers for the following mitochondrial
genes: NADH dehydrogenase 1 (*mt-Nd1*), cytochrome c oxidase subunits 1 and 2 (*mt-Co1* and *mt-Co2*),
and ATP synthase subunit 6 (*mt-Atp6*). Reactions were performed in 384-well plates using the CFX384
Real-Time System (Bio-Rad) with SYBR Green Master Mix (Agilent, Santa Clara, CA). Each well contained
10 ng of gDNA and 2.5 µM of each primer in a final volume of 10 µL. All reactions were run in technical
triplicates. Hexokinase 2 (*HK2*) was used as the reference gene for normalization, with primers listed in the
Kay Resources Table.
1628

QUANTIFICATION AND STATISTICAL ANALYSIS
GraphPad software (versions 8.0.1 and 9.5.1) and Microsoft Excel were used for data analysis and plotting.
Results shown are either representative or cumulative from at least two independent experiments. Except
for gene expression analyses during development, RT-qPCR and WB data were normalized to the average
of at least two WT or control mice of their corresponding litter. Sample size, statistical tests, and the meaning
of the dots in each graph are specified in figure legends, except for TEM graphs (Figures 4C, 6F and S6O;
samples sizes indicated in Table S1). In violin plots, median and interquartile range (IQR) are shown; in dot
plots, mean ± SEM; in bar plots with dots, the mean + SEM; and in standard bar plots, the mean + SD.
Outliers were identified using the ROUT method (Q = 5%) or Grubbs' test (α = 0.05) for datasets with n ≥
6. Normality was assessed using the Shapiro–Wilk test (n < 50) or the Kolmogorov–Smirnov test (n > 50).
Statistical significance between two groups was determined using Student's t-test for parametric data or
the Mann–Whitney test for nonparametric data. Welch's correction was applied when variances between
groups were unequal. For comparisons among more than two groups, one-way ANOVA or two-way ANOVA
with Tukey's post hoc test was used for parametric data, and the Kruskal–Wallis test, followed by Dunn's
multiple comparison test, was used for nonparametric data. Survival curves were compared using the
Gehan–Breslow–Wilcoxon test. A critical value for significance of $p < 0.05$ was used throughout the study
(*$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, ****$p < 0.0001$). In Figures 1H, 2A, 2H, S5B–C, S6F–H, S7A, S7B, S7E,
and Table S2, only the most relevant statistically significant comparisons are shown; non-significant results
are not displayed. In all other figures, statistical significance is indicated only when differences are
significant.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure S1

A P7 Distal small gut

LREs | Magnification

N | LV | AEC | L | LV | mvb | mvb | e | e

B P7 ileum | Magnification

LAMP1 | Dab2 | βcat | DNA

L | L

LAMP1 | Dab2

C GFP-LC3 mice

P7 mCherry
P21

KDa	P7	P21	
32			mCherry
45			GFP-LC3 I / GFP-LC3 II
25			Free GFP
130			USP7

D

0hrs FR-Dextran | 3hrs Rd-Dextran | 4hrs analysis | 8hrs analysis

4hrs | Magnification
FR-dext
Rd-dext
FR/Rd-dext DNA

8hrs | Magnification
FR-dext
Rd-dext
FR/Rd-dext DNA

Figure S2

Figure S3

Gate Strategy Fig S4F

Gate Strategy Figs 4D, 6G, 7N, S2E, S2F, S6I–K

Gate Strategy
Figs 5H, 6L, 7A, S5E

Figure S4

Figure S5

Figure S6

Figure S7

Table S1

	Class	No. mitochondria / μm²	% of mitochondria	Mitochondrial area	Mitochondrial aspect ratio
Figure 4C	I	11, 10	11, 10	239, 271	239, 276
	II	11, 10	11, 10	13, 32	12, 28
	III	11, 9	11, 10	14, 12	18, 12
Figures 6F and S6P	I	23, 24, 13, 17	23, 24, 13, 17	184, 321, 259, 128	169, 329, 251, 130
	II	23, 24, 13, 17	23, 24, 12, 17	19, 73, 14, 45	21, 74, 15, 44
	III	23, 24, 13, 17	23, 24, 13, 17	45, 78, 17, 134	47, 78, 16, 149

Table S2

Protein	Figure 5H		Figure 6L	Figure S5E
	WT vs *Atg16*cKO	WT-P14 vs WT-Preadult		
ASS1	*	****	***	***
GLS	*	****	****	**
HK1	***	**	**	—
CPT1A	*	****	***	****
ACAC	NS	****	***	**
CS	****	****	***	****
IDH2	****	****	****	*
ATP5A	NS	****	*	NS