The effect of hunger and state preferences on the neural processing of food images

Denise Moerel[a,b], Cecilia Chenh[a], Sophie A. Bowman[a], Thomas A. Carlson[a]

a) School of Psychology, University of Sydney, Sydney, Australia
b) The MARCS Institute for Brain, Behaviour and Development, Western Sydney University, Sydney, Australia

ABSTRACT

Visual information plays a key role in guiding food-related decisions. While previous studies have shown that features such as calories and naturalness are encoded by the visual brain, it remains unclear how this encoding is shaped by the observer's current state. In this study, we explore the effect of 1) hunger state, 2) current individual preference, and 3) task relevance on the processing of visual food information. Participants underwent two EEG sessions: one after fasting overnight and the another after eating normally. We used multivariate decoding methods to assess the impact of hunger on the decoding of edibility and food identity, and Representational Similarity Analysis to determine the time-course of information related to food flavour, personal appeal, and arousal, both when participants attended to the stimuli and when their attention was distracted away. Results showed that the visual brain quickly encodes information about edibility and food identity, and this was not influenced by whether the participants were in a fasted or sated state. Information about the flavour profile was found to be represented regardless of attentive state. Personal appeal and arousal information emerged later and were only observed when the food was task relevant. Using RSA, we found that food appeal and arousal encoding was more closely aligned with behavioural ratings within rather than between sessions, suggesting the nature of the encoding was driven by current state. In conclusion, the study provides insights into how personal preferences and physiological states influence the representation of food information in the brain.

1. INTRODUCTION

Humans rely on smell, taste, and vision to perceive, categorise, and choose foods. Among the senses, vision is unique in that it allows us to gather information about foods at a distance. Whether the food is sitting on a plate in front of us or 20m away on a billboard, the brain's visual system enables us to rapidly gather information to inform our decisions. Evidence from fMRI suggests that the brain uses similar mechanisms to encode food cues through vision and taste (Avery et al., 2021; Simmons et al., 2005). In addition, recent studies have provided evidence that food stimuli may drive the coarse scale organisation of the ventral visual stream (Jain et al., 2022; Khosla et al., 2022). Recent studies have also provided insight into how different features of food are encoded in the brain. For example, multiple studies have shown the visual brain distinguishes between edible and inedible (i.e., food and non-food) objects (Avery et al., 2021; Chen et al., 2016; Killgore et al., 2003; Moerel, Psihoyos, et al., 2024; Rothemund et al., 2007; Simmons et al., 2005; Stingl et al., 2010; Tsourides et al., 2016). Avery and colleagues (2021) further showed that the visual cortex carried information about the taste of the food (sweet, salty, or sour) when viewing food images. In addition, studies have shown that the neural response to food stimuli is modulated by the energetic/caloric content of the food (Chae et al., 2025; Killgore et al., 2003; Mengotti et al., 2019; Moerel, Psihoyos, et al., 2024; Toepel et al., 2009), and by whether food is natural or prepared (Coricelli et al., 2019; Moerel, Psihoyos, et al., 2024; Pergola et al., 2017; Vignando et al., 2019). However, it is still unclear how the encoding of visually presented food by the brain is mediated by the state of the observer. In this study, we investigate the effect of 1) hunger state, 2) current personal appeal, which could be influenced by both temporary cravings and long-term preferences, and 3) task relevance on the neural coding of visual food information.

Several studies have found an effect of hunger on the neural response to food stimuli. For example, fMRI studies have shown a greater response to pictures of foods when participants had fasted (Charbonnier et al., 2018; Cheng et al., 2007; Führer et al., 2008; Holsen et al., 2005; LaBar et al., 2001; Mohanty et al., 2008). In a meta-analysis, van der Laan and colleagues (2011) found an influence of hunger on visual food stimulus activation in the right parahippocampal gyrus and amygdala, areas suggested to process reward, and left lateral OFC, suggested to process the expected pleasantness of the food. EEG studies also have found effects of hunger on the activation in response to food stimuli (Ilse et al., 2020; Nijs et al., 2010; Stockburger et al., 2008, 2009; Sultson et al., 2019) and non-food stimuli (Sultson et al., 2019). Other work has suggested that appeal and calorie content might mediate hunger effects. For example, an fMRI study showed that medial and lateral OFC responses were enhanced for high compared to low appeal foods, but only when the participant was

hungry (Piech et al., 2009). In addition, several fMRI studies have found effects of hunger are mediated by the calorie content of the food (Frank et al., 2010; Goldstone et al., 2009; Siep et al., 2009). It is, however, possible that this interaction is mediated by an attentional bias. That is, hunger could increase the salience of food items, especially if they are highly appealing. Finally, it is not yet clear whether hunger can affect the representation of information about the food, as previous work used univariate analysis methods to show differences in activation driven by hunger (Charbonnier et al., 2018; Cheng et al., 2007; Führer et al., 2008; Holsen et al., 2005; Ilse et al., 2020; LaBar et al., 2001; Mohanty et al., 2008; Nijs et al., 2010; Stockburger et al., 2008, 2009; Sultson et al., 2019). There is a distinction to be made about whether hunger diffusely increases brain activity, possibly mediated by arousal or inhibitory control, or whether hunger functionally enhances the representation of food stimuli.

The findings about the whether the visual brain encodes personal food appeal information have been mixed (Chae et al., 2025; Moerel, Psihoyos, et al., 2024; Schubert et al., 2021). One study used multivariate pattern analysis on EEG data to show there was information about the subjective 'tastiness' of the food from approximately 530 ms after stimulus onset (Schubert et al., 2021). Notably, the same result was found when participants did not have to make explicit judgments about the palatability of the food, although the onset of this information emerged later, at 740 ms after stimulus onset. We recently used multivariate pattern analysis on EEG data to investigate whether more general food preferences across the population were represented and whether the brain represented the level of arousal in addition to the valence (palatability) of the food (Moerel, Psihoyos, et al., 2024). Our study found neither valence nor arousal information could be decoded from EEG recordings for visually presented food stimuli. The studies differed in terms of whether the valence and arousal ratings were personal or from another group of participants, and whether the participants had fasted. Therefore, in this study we obtained personal appeal models and studied the brain in both a fasted and fed state.

Finally, we investigated the effect of attention, manipulated through task relevance, on the coding of visual food information in the brain. As noted previously, it is possible that attention mediates the effect of hunger on visual information processing. For example, there is evidence for attentional capture of food over non-food words (Mogg et al., 1998; Placanica et al., 2002) and images (Piech et al., 2010) when participants are hungry. Further evidence for a role of attention in the modulation of hunger effects comes from EEG (Ilse et al., 2020; Nijs et al., 2010; Stockburger et al., 2008). Here, studies have shown evidence for an enhanced N2pc component (Ilse et al., 2020), linked to orienting of spatial attention, and P300 component (Nijs et al., 2010), linked to attention and working memory, for food compared to

non-food stimuli when participants were hungry. Previous work has also shown that attention can affect the representation of visual information generally, with stronger and/or more sustained representations of the information that is within the focus of attention compared to information that is not attended (Battistoni et al., 2020; Goddard et al., 2022; Grootswagers et al., 2021; Kaiser et al., 2016; Moerel et al., 2022; Moerel, Rich, et al., 2024). It is therefore possible that hunger state and attention modulate the processing of information about food.

The goal of this study was to investigate the effect of the state of the observer on the visual processing of information about food. Specifically, we investigated the effects of 1) hunger state, 2) current personal appeal, and 3) task relevance. Participants took part in two different EEG sessions, once after fasting overnight ('fasted': 13.00 ± 2.37 h since eaten), and once after eating as normal ('fed': 1.53 ± 0.78 h since eaten). We use two tasks to manipulate whether the stimuli were task relevant, and therefore within the focus of attention. Using multivariate decoding methods on EEG data, we studied the effect of hunger on the coding of edibility and food identity information in the brain. In addition, we used Representational Similarity Analysis to determine the time-course of information about food flavour, appeal and arousal, obtained through ratings at the start of the session. Our results showed that there was information about edibility and the identity of the food items. Critically, the hunger state of the participant did not affect the coding of this information, with no reliable difference between when the participant was in fasted or sated state. We found evidence for the representation of information about the flavour profile, regardless of whether the food was task relevant. Information about personal appeal and arousal emerged later in time but was only found when the food was task relevant. This information about personal appeal and arousal did not generalise across different sessions or participants, reconciling previous findings and suggesting this information is driven, at least in part, by the immediate state of the participant. Together, these results provide insight into the effects of personal food appeal and physiological states of the perceiver on the representation of food information in the brain.

2. MATERIALS AND METHODS

2.1. Participants

Twenty-four individuals participated in two experiment sessions. One participant was excluded from the analysis as they reported their last food intake as 13 hours prior to the session in the fed condition. Therefore, data were analysed for 23 participants (14 female/9 male) with a mean age of 22.83 years (SD = 8.45, range = 18-62) and a mean BMI of 21.87 points (SD = 2.65, range = 17.3-28). All participants reported normal or corrected-to-normal visual acuity, were able to fast, and reported no history of epilepsy, head injury, neurological disorder or

eating disorder. No participants were currently on a diet, however, three participants had dietary restrictions (gluten free, pescatarian, and intermittent fasting respectively). Participants completed additional surveys, that were unrelated to this study, about their self-reported health, knowledge about calories, and information about their culture and language. Participants received a total of $100 AUD for their participation in both sessions. The study was approved by the Human Ethics Committee of the University of Sydney, and participants provided written and verbal consent prior to participating in the experiment.

2.2. Hunger manipulation

We used a within-subjects design to test the effect of hunger on the representation of food in the brain. All participants attended two two-hour EEG sessions, one session for the fasted condition, and one session for the fed condition. The sessions were scheduled on two separate days, ranging from 3–14 days apart (mean = 7.13 days, SD = 3.17 days). The order of the sessions was counterbalanced to minimise order effects. Both sessions were scheduled at the same time of day to minimise variations in circadian rhythm (Poggiogalle et al., 2018; Stockburger et al., 2008). In the fasted condition, participants were instructed to fast for at least eight hours prior to participating in the experiment, based on previous work (Führer et al., 2008; LaBar et al., 2001; Piech et al., 2010), but they were allowed to drink water. Participants in the fed condition were instructed to eat as they normally would prior to the session. To ensure the effectiveness of the hunger manipulation, participants completed subjective physiological ratings at the beginning of each session. Before participating in the EEG experiment, participants were instructed to rate their hunger, thirst and fatigue levels on a one-hundred-point sliding scale, using an iPad, with 100 indicating very hungry/ thirsty/ tired. Participants also stated the time of their last full meal and the time of their last food intake (including snacks). Note that one participant provided the same subjective hunger rating across both the fasted and fed condition, although they complied with the food intake instructions. As removing this participant did not change the results, we have included this participant in the sample for completeness.

2.3. Stimuli and experimental design

The visual stimuli consisted of 280 segmented colour objects images (see Figure 1A). Half of these were food images (70 natural foods, 70 transformed foods) and half were non-food images (70 animate objects, 70 inanimate objects). All food images were selected from the Foodcast Research Image Database (FRIDa) (Foroni et al., 2013). The subset of images used was selected to be familiar for Australian participants, and rotten food items were excluded.

We controlled for the transformation of the foods by selecting 70 natural and 70 transformed foods. Natural foods are defined as being in their original state (e.g. banana), while transformed foods are foods whose form has been changed by humans in some way, for example, by cooking or preserving (e.g. donut). For the non-food images, 88 were selected from FRIDa (Foroni et al., 2013), and the remaining 52 images were obtained from PNG images.

Figure 1. Example stimuli and EEG task overview.

Figure 1. Example stimuli and EEG task overview. A) Examples of the stimuli that were used in the experiment. There were 140 food stimuli (examples displayed on the left) and 140 non-food objects (examples displayed on the right). The food stimuli were selected from the

Foodcast Research Image Database (FRIDa) (Foroni et al., 2013), and there were 70 natural foods and 70 prepared foods. The non-food objects were obtained from FRIDa (Foroni et al., 2013), and PNG images, and there were 70 animate objects and 70 inanimate objects. Note the stimuli depicting humans have been replaced with silhouettes in this figure to comply with bioRxiv guidelines. **B)** An overview of the letter task. Participants monitored a stream of letters on the screen and pressed a button when they saw a vowel. The images behind the letters are not task relevant. The letters and images were on the screen for 100 ms, with a 50 ms blank screen between presentations. **C)** An overview of the calorie categorisation task. Participant were presented with a food image for 100 ms and were asked to rate the calorie content as higher or lower compared to bread. There was a 1500 ms response time-out.

The stimuli were presented in the centre of the computer screen on a white background, using the Psychopy library (Peirce et al., 2019) in Python. The participants were seated approximately 60 cm from the computer screen, and the stimuli were approximately 6.25 degrees of visual angle in size (256 x 256 pixels). The fixation bullseye was displayed at the centre and was approximately 0.61 degrees of visual angle (24 x 24 pixels) in size. Participants were instructed to keep fixation on the central bullseye during the task. The EEG experiment consisted of a letter task and a calorie categorisation task (see Figure 1B). There were 6 blocks in total, and each block consisted of the letter task followed by the calorie categorisation task. The experiment took approximately 40 minutes to complete.

2.3.1. Letter task

In the letter task, participants saw a rapid stream of food and non-food images, with a simultaneously presented stream of letters, overlaid on top of the object images. Participants were instructed to ignore the objects and press a button whenever they saw a vowel (A, E, I, O, or U) in the stream of letters, as quickly and as accurately as possible. This means that the object images were not task relevant and attention was directed away from the stimuli and towards the letters.

Each of the 140 food images and 140 non-food images were presented three times per block and thus 18 times in total across the 6 blocks. Therefore, participants saw 5040 images in total. The order of images was randomised such that food and non-food images were intermixed and all items were presented before they were repeated a second and third time in each block. The object images with overlaid letters were presented at a rate of 6.67 Hz, where each stimulus was on the screen for 100ms, followed by a blank screen for 50 ms.

Each sequence contained 2-4 targets (vowels). Letters were presented in a randomised order, however, there were always at least 15 random letter presentations between two targets and the first 10 and last 10 letters within each sequence were never targets. Letters 'Q' and 'Y' were not used, as 'Q' looked too similar to 'O' at a rapid presentation

rate and 'Y' could be confused for a vowel. Task performance was indicated by mean accuracy and reaction time. Mean accuracy was calculated per session, as the number of correct responses made within 1.5 seconds post target presentation, divided by the total number of targets seen. Mean reaction time in seconds was calculated per session, as the mean time of correct responses.

2.3.2. Calorie categorisation task

In the calorie categorisation task, participants saw images of food items and were asked to classify the foods as higher or lower in caloric density compared to bread as quickly and accurately as possible (Figure 1C). Specifically, they were asked to compare 100g of the presented food to 100g of bread in terms of the caloric density. We chose bread as a reference point as it is familiar to Australian participants and was close to the median perceived calorie rating in FRIDa. Participants completed this classification by pressing the left or right outer buttons of a button box. The buttons for "higher calories" and "lower calories" changed between blocks to minimise the effect of motor preparation and execution signals that could contaminate the EEG signal.

The 140 food stimuli were shown once per block in a random order, meaning each stimulus was shown 6 times in total across the 6 blocks of the classification task. Participants took voluntary breaks after every 20 trials to minimise fatigue. During these breaks, the instructions reappeared on the screen to remind the subject which button mapped to high or low calories. In each trial, the fixation point was shown on the screen for 500-1000ms to prevent participants from anticipating the stimulus onset. The food was shown for 100ms, then replaced by the fixation point again for 1500ms or until the participant responded. If the participant responded within 1500ms, the circle would change to black to indicate that their response was registered and the next trial would start. If no response was registered, a message saying "Too late!" appeared on the screen to encourage the participant to respond faster on the next trial.

2.4. EEG Recording and Pre-processing

We continuously recorded the 64-channel EEG data at a sampling rate of 1000Hz using a BrainVision (GmbH, Herrsching, Germany) ActiChamp system. The electrode locations corresponded to the international 10-10 system of electrode placement (Oostenveld & Praamstra, 2001) with the reference at FCz. EEG data pre-processing was conducted offline using the EEGLAB toolbox (Delorme & Makeig, 2004) in MATLAB. We used minimal EEG pre-processing, following earlier work (Moerel, Psihoyos, et al., 2024). We used a high-pass

and low- pass filter to remove signals below 0.1Hz and above 100Hz, respectively, and down-sampled the data from 1000Hz to 250Hz. We then created epochs for each trial, starting 100 ms before stimulus onset and ending 1200 ms after stimulus onset.

2.5. Decoding analysis

To determine whether the hunger state of the individual affects how information about food is processed in the brain, we decoded information about edibility and food identity from the EEG data. To gain insight into the time-course of information about edibility, we decoded whether a stimulus item was a food or non-food object. To determine the time-course of food identity, we used pair-wise decoding of all food items. We used CoSMoMVPA toolbox for Matlab (Oosterhof et al., 2016) for both of these decoding analyses, using linear discriminant analysis classifiers on the pattern of activation across all electrodes to distinguish between the classes of interest. We did this separately for each time-point and participant. Critically, we also did this separately for the sessions where the participant was fasted and fed, and then compared decoding accuracies across the two different sessions.

For the decoding of edibility information, we trained a classifier to determine whether a stimulus was a food or a non-food object for each time-point. We only used the data from the letter task for this analysis, as the calorie categorisation task did not have non-food object items. We used an exemplar-by-block cross validation approach, following previous work (Carlson et al., 2013; Grootswagers et al., 2019; Moerel, Psihoyos, et al., 2024). We left out one pair of images, a randomly matched food and non-food object item, from one block as the test data. The 140 food items were randomly paired with the 140 non-food object items, and we iterated over the different test pairs. We iteratively trained the classifier on the remaining items from the remaining five blocks, leaving each block out as a test block once. We repeated this analysis 10 times, with different random image pairings, and averaged the decoding accuracies across the repeats.

To determine *if* and *when* there was information about the identity of the food, we decoded pairs of food stimuli. For this analysis, we used the data from both the letter task, including the only the food items, as well as the data from the calorie categorisation task. We repeated the decoding analysis for each possible combination of two food stimuli, and then averaged the decoding accuracy across the possible pairs. We repeated the decoding analysis for each time-point and participant. For the letter task we used a leave-one-block out cross-validation approach for each pair of stimuli, where all stimulus presentations from a single test block were left out. We trained the classifier on all remaining blocks in an iterative fashion. For the calorie categorisation task, participants used the button box to indicate

whether the stimulus was higher or lower in calories compared to bread. This means that the signal could be contaminated with the motor response or preparation. We therefore swapped the response button mapping between blocks, and tested the classifier on data from two blocks, one with each response mapping. We iteratively trained the classifier on the data from all remaining blocks, with balanced response mapping. We used all combinations of two blocks with opposite response mapping as the test blocks and averaged across the decoding accuracies.

2.6. Representational Similarity Analysis

We used Representational Similarity Analysis (RSA) (Kriegeskorte et al., 2008; Kriegeskorte & Kievit, 2013) to examine whether information about the flavour of food and current valence and arousal were represented in the brain. We calculated Representational Dissimilarity Matrices (RDMs) for the EEG data, representing the dissimilarity in the pattern of activation across the sensors evoked by two different food stimuli. We used the classification accuracy from the pair-wise decoding analysis described above as a measure of dissimilarity. We did this for all possible combinations of two food items, resulting in a 140 by 140 RDM (Figure 2A). We repeated this analysis for each time-point in the epoch, separately for each participant and session. We obtained three model RDMs, coding for 1) the flavour profile of the food, 2) the current appeal of the food, and 3) the current arousal of the food. Models for the current appeal and arousal were based on behavioural ratings obtained from the same EEG session. We then calculated partial correlations between the models and the EEG RDMs, for each time-point, participant, and session. To account for possible low-level visual differences between the stimuli, we partialled out three artificial neural network control models (V1, V2, and V4 layers of CORnet-S (Kubilius et al., 2019)), resulting in partial correlations between the EEG and the models.

The model of the flavour profile (Figure 2C) was based on a different sample of 84 participants (mean age = 20.48, SD age = 2.60, age range = 18 – 33, 74 right-handed / 10 left-handed, 57 female / 26 male / 1 non-binary). Participants rated the 140 food stimuli on four different flavour dimensions: sweet, salty, sour, and bitter. Participants used a sliding scale to rate each of the food items on a single taste dimension per block, covering all taste dimensions across four different blocks. We obtained the flavour profile RDM by calculating the Euclidean distance between two pairs of food stimuli, on the pattern of responses across the four flavour dimensions. We did this separately for each participant, and then averaged across participants to obtain a single flavour profile RDM.

Figure 2. Representational Similarity Analysis overview. We made 140 x 140 Representational Dissimilarity Matrices (RDMs) representing the dissimilarity between all pairs of food images. **A)** The average RDM for the EEG data from the letter task (left) and calorie categorisation task (right). The EEG RDMs show the dissimilarity in the pattern of activation across the sensors, for each pair of stimuli. We make a unique RDM for each participant, session, and time-point. The RDMs plotted here show the mean across all participants and sessions, for the time window between 100 ms and 400 ms relative to stimulus onset. **B)** The individual appeal (left) and arousal (right) models. We made a personal model for each participant and session, based on behavioural ratings obtained at the start of the experiment. Mean RDMs across all participants and sessions are shown here. **C)** The general flavour model. This model was based on behavioural ratings from a different group of

84 participants. **D)** The three CORnet-S control models. These RDMs model the low-level visual differences between the stimuli.

The appeal and arousal models are shown in Figure 2B. Unlike the flavour profile model, the appeal and arousal models were based on behavioural data from the same participants that participated in the EEG session. At the start of each EEG session, participants used a sliding scale from 0 to 100 to rate each food stimulus on how appealing the food was to them. We used these ratings to calculate an individual appeal and arousal RDM for each participant and session. The appeal RDM, coding for how appealing the food was to the participant, was obtained by calculating the absolute difference (dissimilarity) in the appeal rating between each combination of two food stimuli. The arousal RDM coded for the similarity in the emotional response that was elicited by the food, regardless of whether the food was considered appealing or unappealing. For example, a participant might rate cake as very appealing and brussels sprouts as very unappealing. This would mean that both the cake and brussels sprouts are very high arousal (i.e. they elicit a strong response), but the cake is high in appeal and the brussels sprouts low in appeal. According to the arousal model, the cake and brussels sprouts would therefore be very similar, whereas they would be very different according to the appeal model. To convert the appeal rating into an arousal rating, we recoded the ratings as the absolute distance to the middle possible rating (50). We then calculated the arousal RDM in the same way as the appeal RDM.

To control for possible low-level differences between stimuli, we used three artificial neural network control models (Figure 2D). We used the V1, V2, and V4 layers of CORnet-S (Kubilius et al., 2019), designed to reflect the response of visual areas V1, V2, and V4 respectively. We obtained the neural network responses to all stimuli and created the RDM by calculating the cosine distance in response between each pair of images. We then calculated the partial correlations between the EEG and model RDMs by partialling out the three control RDMs.

2.7. Exploratory analyses

We explored whether the information about personal appeal and arousal generalises 1) across time and hunger state, and 2) across different participants. To test whether information about appeal and/or arousal generalises across hunger states and time, we calculated the correlations between the EEG data, and the appeal and hunger models based on the behavioural data from the other session. If this information generalises across sessions, that suggests we are tapping into preferences that are stable across time. If not, this suggests that the appeal/arousal information is driven by the temporary state of the participant. We also

tested whether information about appeal and/or arousal generalises across participants. If so, this would suggest the information is driven by general preferences across the population, rather than personal preferences. For this analysis, we calculated the correlations between the EEG data, and the appeal and hunger models based on the behavioural data from a randomly matched participant in the same hunger state condition. This means the hunger level was matched between participants. We repeated this analysis 10 times, with different random matches between participants, and then averaged the correlation across the 10 repeats.

In addition, we also explored whether there was an effect of appeal on the coding of edibility information in the brain. We used the results from the edibility decoding analysis described above, training and testing on all 140 food stimuli and 140 non-food object items. As described above, this analysis was only performed on EEG data from the letter task. For this exploratory analysis, we split the decoding accuracies based on whether the presented item was high appeal (top 25%) or low appeal (bottom 25%). We collapsed across hunger state conditions. We then compared the edibility decoding for high appeal and low appeal food items.

2.8. Statistics

We used Bayesian statistics (Dienes, 2011; Kass & Raftery, 1995; Morey et al., 2016; Rouder et al., 2009; Wagenmakers, 2007), implemented through the Bayes Factor R package (Morey & Rouder, 2018), to determine whether there was evidence for above chance decoding or partial correlations above 0. We applied Bayesian t-tests at each time-point. All statistical analyses were applied at the group level, and we used the same analyses for the letter task and calorie categorisation task. For the decoding analysis, we tested whether there was evidence for the chance decoding (null hypothesis) or the above chance decoding (alternative hypothesis). For the RSA analysis, we tested whether there was evidence for a partial correlation of 0 (null hypothesis) or above 0 (alternative hypothesis). We used a point null for the null hypothesis and a half-Cauchy, directional, prior for the alternative. The prior was centred around chance ($d = 0$, chance decoding or a partial correlation of 0), and we used the default prior width of $d = 0.707$ (Jeffreys, 1998; Rouder et al., 2009; Wetzels et al., 2011). We excluded small effect sizes, that may occur by chance under the null hypothesis, by excluding the interval between $d = 0$ and $d = 0.5$ from the prior (Teichmann et al., 2022).

We also applied Bayesian t-tests for our behavioural ratings. We followed the same steps as described above with the following differences. To obtain Bayes Factors for the differences between the fasted and fed conditions, we subtracted the ratings for the fasted and fed conditions and performed Bayesian t-tests against 0. We used a full-Cauchy rather

than a half-Cauchy prior, to make the test 2-tailed. We excluded the interval between d = -0.5 and d = 0.5 from the prior for consistency with the t-tests described above.

3. RESULTS

3.1. Behavioural results

At the start of both EEG sessions, participants answered questions to determine their level of hunger, thirst, and fatigue, as well as the time since their last food intake and full meal. An overview of the results is shown in figure 3A. There was evidence for a difference in the reported hunger level between sessions, with higher hunger ratings for the fasted compared to fed session (BF = 6.75 x 10^5). There was evidence for no difference between sessions in the reported thirst level (BF = 0.05) or fatigue (BF = 0.01). There were differences between sessions in the reported time since the last full meal (BF = 9.35 x 10^8) and last food intake, including snacks (BF = 1.30 x 10^{14}), with longer reported times in the fasted compared to fed session. Together, these results suggest that the hunger manipulation was successful, resulting in higher hunger levels when participants were asked to fast.

Participants also rated the appeal of each food item at the start of both EEG sessions. We tested whether food appeal changed across the sessions. Previous work showed faster response times for high-fat (i.e. high-calorie) foods over low-fat foods in a visual search task when participants were hungry compared to when they were sated (Sawada et al., 2019). In addition, there is evidence participants choose high calorie over low calorie foods when hungry (Read & van Leeuwen, 1998; Tuorila et al., 2001). This suggests that hunger effects on appeal ratings could be stronger for high-calorie food items. We therefore split the appeal ratings based on the perceived caloric content, comparing the appeal of high calorie (top 25%) and low calorie (bottom 25%) foods across the fasted and fed conditions. For the low-calorie foods, we found no evidence for an effect of hunger state (BF = 0.08). However, there was an effect of hunger state for the high calorie foods (BF = 18.15), with an increased appeal of high calorie foods when the participants were hungry compared to sated.

A) State and time since last food intake

B) Food appeal ratings

Figure 3. Behavioural results. A) The responses to the questionnaire taken at the start of each EEG session, in the session where participants had eaten (purple) or were asked to fast (green). Dots show individual participants, and the thick grey line shows the mean across participants. The top panel shows the rating for the hunger level (left), thirst (middle), and fatigue (right). The bottom panel shows the reported time since the last full meal was consumed (left) and the time since any food, including snacks, was consumed (right). **B)** Participants also rated the appeal of each food items at the start of each EEG session, with food appeal ratings for the session where participants had eaten (purple) or were asked to fast (green) shown here. We split the data based on the perceived calorie content (Foroni et

al., 2013), showing high calorie foods (top 25%) in light colours and low calorie foods (bottom 25%) in dark colours. Plotting conventions are the same as Figure 3A.

3.2. No effect of hunger on edibility and food identity decoding

We investigated whether the hunger state of the participant influences the processing of food stimuli in the brain. First, we used the EEG data from the letter task to decode whether a stimulus item was a food or non-food object. This allowed us to investigate the time-course of edibility information when participants were asked to fast or had just eaten. The results are plotted in Figure 4. There was evidence for edibility information from 84 ms after stimulus onset for both the fasted condition and the fed condition. However, there was evidence for no difference between the fasted and fed condition for most of the time-course, and no time-points showed strong evidence for a difference.

Figure 4. The time-course of edibility decoding for the fasted and fed conditions. We used a linear classifier on the EEG data from the letter task to distinguish between food items and non-food objects. The decoding accuracy over time is shown for the fasted condition (green) and the fed condition (purple). Shaded areas around the plot line show the standard error of the mean. Theoretical chance is at 50% decoding accuracy. The Bayes factors (BFs) are shown on a logarithmic scale below the plot for the fasted condition (top), fed condition (middle) and the difference between the two conditions (bottom). BFs above 1, showing evidence for above chance decoding, are plotted in warm colours and BFs below 1, showing evidence for chance decoding, are plotted in cool colours.

We also investigated whether the hunger state of the participant could influence the coding of more fine-grained information about the food. To investigate this question, we decoded the identity of the food in a pair-wise way. Figure 5 shows the food identity decoding for the calorie categorisation task (Figure 5A) and the letter task (Figure 5B). For the calorie categorisation task, we found evidence for food identity decoding from 96 ms after stimulus onset in the fasted condition, and 100 ms after stimulus onset in the fed condition. There was evidence for no difference in food identity decoding between the fasted and fed conditions for most time-points, with a few time-points showing insufficient evidence for either hypothesis. The pattern of results for the letter task was similar, with evidence for food identity decoding from 88 ms onwards in the fasted condition, and from 96 ms after stimulus onset in the fed condition. Again, there was evidence for no effect of hunger on food identity decoding for most time-points, with insufficient evidence found for a few time-points. Together, the decoding results suggest that the brain processes whether an object is a food, as well as the identity of the food, but this information is not affected by the hunger state of the participant.

Figure 5. The time-course of food identity decoding for the fasted and fed conditions for the calorie categorisation task (A) and letter task (B). We used a linear classifier on the EEG data to distinguish between two food items in a pair-wise way. Plotting conventions are the same for both plots. The decoding accuracy is plotted over time for the fasted condition (green) and fed condition (purple). Theoretical chance is at 50% decoding accuracy, and shaded areas around the plot line show the standard error of the mean. The Bayes factors (BFs) are shown below the plot on a logarithmic scale. The top plot shows the BFs for the fasted condition, the middle plot for the fed condition, and the bottom plot for the difference between the two conditions. BFs above 1 are plotted in warm colours and BFs below 1 are plotted in cool colours.

3.3. The representation of flavour, appeal, and arousal information investigated through Representational Similarity Analysis

We used RSA to gain further insight into the coding information about the flavour profile, appeal, and arousal of food. The flavour profile was based on data from a different group of participants, i.e., one general flavour profile model. The appeal and arousal models were based on data from the EEG participants, obtained in the same session, with a unique appeal and arousal model made for each participant and for each session. We calculated partial correlations between the EEG data from the calorie categorisation task or the letter task and the models, controlling for three low-level visual CORnet models. Consistent with the findings from the decoding analysis, there was no evidence for an effect of hunger on information about the flavour profile, appeal, and arousal of food, for either the calorie categorisation task or the letter task. Most time-points in all tests showed evidence for no difference between the partial correlation from the fasted and fed conditions. The data, separated for the fasted and fed condition, can be found in Supplementary Figure 1.

To maximise the power, we collapsed across the fasted and fed conditions. Figure 6 shows the time-course of information about the flavour profile, appeal, and arousal of food. For the calorie categorisation task, there was evidence for information about the flavour profile from approximately 192 ms onwards. There was also information about the appeal and the arousal of the food item, but this information emerged later, around 520 ms and 464 ms after stimulus onset respectively. The letter task, where the food items are not task-relevant, shows a different pattern of results. For this task, we find information about the flavour profile from approximately 212 ms onwards. However, there is no consistent information about either appeal or arousal, with evidence for no partial correlation for most time-points. Together, these results show that there is information about the flavour profile of the food, regardless of the task. However, there is only information about personal appeal and arousal in the calorie categorisation task, when the task is task relevant, and the stimuli are presented at a slower rate.

Figure 6. The time-course of partial correlations between the EEG data and the flavour profile, appeal, and arousal models. We calculated partial correlations between the EEG data for the calorie categorisation task (left) and the letter task (right) and the models. The top panels show the partial correlation with the flavour profile model, the middle panels show the partial correlation with the individual appeal models, and the bottom panels with the individual arousal models. To account for possible low-level visual differences between the stimuli, we partialled out three CORnet control models from the correlations. The shaded areas around the plot lines show the standard error of the mean. The Bayes Factors (BFs) are shown below the plot. BFs above 1 are plotted in warm colours and BFs below 1 are plotted in cool colours.

3.4. Exploratory analyses of food appeal and arousal

Our results in the previous section show that there is information about food appeal and arousal in the brain, but only for the slow task where the stimuli were task relevant. We therefore only focus on the calorie categorisation task to further explore whether this information about appeal and arousal generalises to 1) appeal/arousal ratings from the other session for the same participant and 2) appeal/arousal ratings from other participants. If there is a correlation between the EEG and the personal appeal/arousal model within participants across sessions, this suggest that this information reflects long term food preferences of that individual, that are stable over time. If there is a correlation between the EEG and the personal appeal/arousal model across participants, this suggest that this information reflects universal food preferences that generalise across the population. Figure 7 shows the generalisation of appeal and arousal information. As described in the previous section, there was information about the appeal of the food item from approximately 520 ms after stimulus onset when we used the individual appeal model that was matched for the participant and session. When using the appeal model for the same participant, but the other session, we found no reliable evidence for information about appeal. Although visual inspection shows there was a trend for a correlation, showing a similar time-course as the EEG correlation with the model based on the same session, the BFs show the evidence for this correlation is anecdotal. We find a similar pattern of results for the correlation with a model from a randomly matched participant. Visual inspection shows a small correlation with a similar time-course as the matched appeal model, but this is not supported by the BF evidence. For the arousal model, we previously found a correlation between the EEG RDM and the individual arousal model that was matched for the participant and session from approximately 464 ms after stimulus onset. There was no evidence for a correlation between the EEG and the arousal model from the same participant but other session, with most time-points showing evidence for the null. In addition, the model based on a random matched participant from the same session also did not correlate with the EEG, most time-points showing evidence for the null. However, there were 2 consecutive timepoints with BF > 10 around 150 ms after stimulus onset. Neighbouring time-points did not show evidence for a correlation above 0, suggesting this is likely driven by noise. Together, these results show that only the matched appeal and arousal within the same session models reliably correlate with the EEG RDMs. However, there is a possibility that appeal could generalise to the other session and perhaps even across participants, but this evidence is merely anecdotal.

Figure 7. the generalisation of food appeal (A) and arousal (B) information across sessions and participants. We explored the correlation between the EEG for the calorie categorisation task and the appeal and arousal models, that were based on 1) the same session and participant (dark blue), the same participant but the other session (teal), a randomly matched participant but the same session (light green). The shaded areas show the standard error of the mean. The Bayes Factors (BF) are shown below the plot on a logarithmic scale. BFs above 1 are shown in warm colours and those below 1 are shown in cool colours. We also explored if the representation of edibility information, whether an item is a food or not, is affected by the appeal or arousal of the food. For this analysis, we used the same edibility

decoding data presented in section 3.2, collapsed the edibility decoding accuracy across the fasted and fed conditions, and split the results based on whether the food item had high appeal (top 25%) or low appeal (bottom 25%) or high or low arousal (top vs. bottom 25%). We did not include the results for the middle 50% of food items and the non-food items. The EEG data were obtained from the letter task, where the objects were not task relevant, and the food items were therefore likely not within the focus of attention. The results are shown in Figure 8. When splitting the results based on appeal, there was information about whether an item was edible from 92 ms onwards for the high appeal items and 84 ms onwards for the low appeal items. However, after approximately 176 ms, there was an effect of appeal, with stronger edibility decoding for the high appeal compared to low appeal items. When splitting the data by arousal, we found edibility decoding from 88 ms onwards for the high arousal items and after 92 ms for the low arousal items. We did not find an effect of arousal on edibility decoding, with most timepoints showing not enough evidence either way, or evidence for no difference between high and low arousal. Together, these results suggest that edibility information in the brain is affected by the appeal, but not arousal, of the food stimulus.

Figure 8. The time-course of edibility decoding, split by appeal (A) or arousal (B) level of the food items. A) We split the edibility decoding results, presented in Figure 4, based on whether the food item had high appeal (top 25%) or low appeal (bottom 25%). We excluded the results for the middle 50% of food items and non-food items. The decoding data for high appeal items is presented in red, and the data for the low appeal items is presented in yellow. Shaded areas around the plot lines show the standard error of the mean. The Bayes Factors (BFs) are presented below the plot, with BFs above 1 shown in warm colours and BFs below 1 in cool colours. **B)** shows the decoding data split by high arousal (top 25%) in red and low arousal (bottom 25%) in yellow. All plotting conventions are the same as in A.

4. DISCUSSION

In this study, we investigated 1) whether the hunger state of the participant influenced the representation of edibility information and object identity information, 2) whether information about personal appeal was reflected in brain activity, and 3) whether focused attention, manipulated through task-relevance, was needed for the coding of food information. Our behavioural results confirmed that there was a difference between the reported hunger levels in the fasted and fed conditions. In addition, we observed increased appeal ratings for the high-calorie items, but not the low-calorie items, when participants had fasted. These findings suggest that the hunger manipulation was successful. Our EEG data showed that there was information about edibility (food vs. non-food) from approximately 84 ms after stimulus onset, and about the identity of the food items (e.g., hamburger vs. pizza) from approximately 96-100 ms after stimulus onset. Critically, the hunger state of the participant did not affect the coding of this information, with no reliable difference between when the participant had fasted or eaten. We found evidence for the representation of information about the flavour profile from approximately 192 – 212 ms after stimulus onset. This information was present regardless of whether the food stimulus was task relevant or not. Information about the personal appeal and valence of the food emerged later, after approximately 520 ms and 464 ms after stimulus onset respectively. This information was only present for the calorie categorisation task, where the stimuli where task relevant and presented at a slow rate. There was no effect of hunger on the coding of flavour, appeal, or arousal information. In addition, the results showed that there was no reliable evidence that information about either appeal or arousal transferred to the other session or participants, although there was a trend. This suggests that appeal and arousal information is driven, at least in part, by the temporary state of the participant. Finally, our results showed that edibility decoding is mediated by appeal, with more information about edibility for food items that were highly appealing compared to those that had low appeal.

Our results show no effect of hunger state on the representation of information about edibility, food identity, flavour profile, personal appeal, or personal valence. Previous work has shown evidence that hunger mediates neural responses to food items, using fMRI (Charbonnier et al., 2018; Cheng et al., 2007; Führer et al., 2008; Holsen et al., 2005; LaBar et al., 2001; Mohanty et al., 2008) and EEG (Ilse et al., 2020; Nijs et al., 2010; Stockburger et al., 2008, 2009; Sultson et al., 2019). There are several possible explanations for this apparent difference in findings between these previous studies and our work. First, previous studies used univariate methods to assess the effect of hunger on the neural activation and are therefore not able to discern whether the coding of information is affected by hunger. It is

possible that the differences in activation, observed in previous work, are driven by other processes such as increased inhibitory control or differences in reward value when participants had fasted. For example, Charbonnier and colleagues (2018) used fMRI to show increased activation in right dorsolateral prefrontal cortex when participants viewed high calorie images in a hungry state, which was interpreted as evidence for increased inhibitory control when participants had fasted. If differences in activation are indeed driven by processes such as cognitive control or reward value, we would not expect to find differences in terms of the food features that are represented in the brain, which would be in line with our findings.

An alternative explanation for a lack of hunger effect is that our hunger manipulation was not successful. We relied on self-reported ratings of hunger rather than biochemical measures of hunger. However, it is important to note that all included participants reported complying with the instructions regarding food intake. In addition, there was evidence for a difference in reported hunger levels, hours since the last full meal, and hours since the last food intake (including snacks) between the fed and fasted conditions. There was also behavioural evidence showing an increase in the reported appeal for high-calorie food items when participants had fasted compared to when they had eaten. This difference was not observed for the low-calorie items, suggesting that our hunger manipulation was effective.

Another explanation for a lack of an effect of hunger is the stimulus set we used. There is some evidence that the stimuli that are chosen can influence whether hunger effects are observed. Using a visual mismatch negativity paradigm in EEG, Sultson and colleagues (2019) showed an effect of hunger between 100 ms and 220 ms after stimulus onset for high fat savoury stimuli. Importantly, there was no effect of hunger for the high fat sweet stimuli, highlighting the importance of stimulus choice. In line with this, other work has suggested hunger effects could be mediated by the motivational saliency of the stimuli. Piech and colleagues (2009) used fMRI to show enhanced responses to high compared to low appeal foods when the participant was hungry, but not when the participant had eaten. In addition, the calorie content of the food has been found to mediate hunger effects (Frank et al., 2010; Goldstone et al., 2009; Siep et al., 2009). Our study used a large stimulus set of 140 diverse food images, covering both high and low-calorie food of various appeal. It is therefore possible that the choice of stimuli can influence whether hunger effects are found.

Another possibility is that hunger effects were masked by our diverse sample of participants. For example, it is possible that collapsing results across men and women obscured hunger effects due to differences between groups. Previous fMRI work shows mixed findings, with some studies showing an interaction between gender and hunger (Frank et al.,

2010), and other studies showing no interaction (Uher et al., 2006). However, a range of studies have found effects of hunger on the neural response to food stimuli when using samples of both women and men, in fMRI (Charbonnier et al., 2018; Cheng et al., 2007; Holsen et al., 2005; LaBar et al., 2001; Mohanty et al., 2008) and EEG (Ilse et al., 2020; Stockburger et al., 2009). This shows that evidence for hunger effects can be found when including both male and female participants. Another factor that could potentially mediate hunger effects is BMI (Nijs et al., 2010). Most of our participants had a BMI in the healthy range (18.5 – 24.9), two participants were underweight (BMI = 17.31 and 18.19), and three participants were overweight (BMI ranging between 26.35 and 27.98). No obese participants (BMI of 30 or higher) were included in the sample. As most participants had a BMI within the healthy range, it is unlikely this can explain the lack of an effect of hunger. Finally, most participants (20 of the 23) reported not being on a diet, with one participant reporting a pescatarian diet, one reporting a gluten free diet, one reporting an intermittent fasting diet. It is therefore also unlikely that differences in diet could have caused the lack of a hunger effect.

Our results showed that there was information about personal appeal and arousal with regard to food stimuli during the categorisation task. This information emerged at approximately 464 – 520 ms after stimulus onset. This was found to be later than information about the flavour profile, which emerged at 192 – 212 ms after stimulus onset. There was no strong evidence for information about the appeal and arousal of food items generalising to a different session with the same participant, or across participants. The finding that the preference model from the same session explained variance in the EEG data, whereas there was only anecdotal evidence for the preference model from the other session, suggests that the information about appeal and valence was, at least partly, driven by the temporary state of the participant. Some preferences that are more stable over time might also have contributed to the information, driving the trend of shared appeal and arousal information across sessions. Although there was a trend suggesting some transfer between session and, to a lesser extent, participants, the current study does not have the power to statistically assess this. Future work could further disentangle the contribution of immediate appeal, potentially driven by temporary cravings, long-term food preferences, and general food preferences across the population. Overall, our study reconciles the different findings from previous work (Chae et al., 2025; Moerel, Psihoyos, et al., 2024; Schubert et al., 2021). Whereas several previous studies found that the brain carries information about the subjective pleasantness or 'tastiness' of food, using EEG (Chae et al., 2025; Schubert et al., 2021) and fMRI (Avery et al., 2025), our previous EEG work showed valence and arousal models based on a different group of participants did not explain the neural data (Moerel, Psihoyos, et al.,

2024). Our current findings suggest that personal appeal and temporary cravings, rather than general preferences across the population, are a better predictor of the brain's encoding of the food stimuli that did not include highly unappetising items. However, Chae and colleagues (2025) did find evidence that models of positive valence, negative valence and tastiness, based on one group of participants, could explain variance in the EEG data from another group of participants. A likely explanation, as suggested by Chae and colleagues (2025), is that their study included items that were specifically chosen to edible but highly unappetising. It is likely that agreement across the population about these unappetising items was high, thus enhancing between-subject model generalisation. In contrast, no highly unappetising items were included in our current and previous work, suggesting that any coding of appeal would be driven by more fine-grained, highly personal, and dynamic preferences in this case.

We further explored whether there was an effect of personal appeal on the coding of edibility information in the brain. Our results showed evidence for stronger edibility decoding in the brain for food items that were highly appealing compared to foods that were low in appeal. No such link was observed for arousal ratings. It is possible that the observed link between edibility decoding and subjective appeal is mediated by the personal experience with the food. It is reasonable to assume that the participants are more likely to consume the foods on a regular basis that are appealing to them (top 25%) compared to the foods that are unappealing (bottom 25%). This means participants might be more familiar with the appealing compared to unappealing items. In other words, the edibility decoding could be partly driven by "is it a food that I would eat" rather than "is it a food". We did not ask participants about how often they consumed each of the food items in the stimulus set, which means it is not possible to further investigate the role of personal experience in edibility decoding. Future work could investigate the factors that mediate edibility information in the brain, disentangling the roles of appeal and experience.

Finally, our results showed there was evidence for coding of both food identity (e.g., hamburger vs. pizza) and flavour profile (e.g. savory vs. sweet), regardless of whether the food items were task relevant. However, information about the current personal appeal and arousal was only present for the categorisation task, where the food stimuli were task relevant. It is thus possible that focussed attention is needed for information about personal appeal and arousal to be encoded by the brain, whereas information about the food identity and flavour profile is coded automatically. However, the stimuli in the calorie categorisation task were also presented at a slower rate compared to the letter task. It is therefore also possible that the faster speed of the letter task caused earlier masking (Robinson et al., 2019), limiting processing of the stimuli.

This EEG study revealed that hunger does not influence the brain's processing of food-related information, including flavour and personal appeal. Whereas food features such as edibility, food identity, and the flavour profile are encoded relatively early and regardless of whether the food item is relevant, subjective qualities, like current appeal and arousal, only emerge later and are influenced by task relevance. This study provides insight into the time-course of how the brain processes different aspects of food information, from basic edibility and identity to more complex attributes like flavour and personal appeal.

5. ACKNOWLEDGMENTS

This work was supported by an Australian Research Council discovery project awarded to TAC (DP200101787). We would like to acknowledge the university of Sydney HPC service for providing high performance computing resources.

6. DATA AND CODE AVAILABILITY STATEMENT

The analysis code and an example dataset can be found via the Open Science Framework (https://doi.org/10.17605/OSF.IO/ZFD7P). The full dataset will be made available upon acceptance.

Supplementary figure 1. No effect of hunger on the representation of information about the flavour profile, appeal or arousal. We calculated partial correlations between the EEG data for the calorie categorisation task (left) and the letter task (right) and the flavour profile (top), appeal (middle), and arousal (bottom) models. We did this separately for the fasted session (green) and fed session (purple). The shaded areas show the standard error of the mean, plotted in the same colour. We partialled out three CORnet control models from the correlations to control for possible low-level visual differences between the stimuli. The Bayes Factors (BFs) below the plot show evidence for a correlation above 0 in warm colours (BF > 1) and evidence for no correlation in cool colours (BF < 1).

7. REFERENCES

Avery, J. A., Carrington, M., Ingeholm, J. E., Darcey, V., Simmons, W. K., Hall, K. D., & Martin, A. (2025). Automatic engagement of limbic and prefrontal networks in response to food images reflects distinct information about food hedonics and inhibitory control. *Communications Biology*, *8*(1), 1–12. https://doi.org/10.1038/s42003-025-07704-w

Avery, J. A., Liu, A. G., Ingeholm, J. E., Gotts, S. J., & Martin, A. (2021). Viewing images of foods evokes taste quality-specific activity in gustatory insular cortex. *Proceedings of the National Academy of Sciences*, *118*(2), e2010932118. https://doi.org/10.1073/pnas.2010932118

Battistoni, E., Kaiser, D., Hickey, C., & Peelen, M. V. (2020). The time course of spatial attention during naturalistic visual search. *Cortex*, *122*, 225–234. https://doi.org/10.1016/j.cortex.2018.11.018

Carlson, T. A., Tovar, D. A., Alink, A., & Kriegeskorte, N. (2013). Representational dynamics of object vision: The first 1000 ms. *Journal of Vision*, *13*(10), 1. https://doi.org/10.1167/13.10.1

Chae, V. J., Grootswagers, T., Bode, S., & Feuerriegel, D. (2025). *Characterising the neural time-courses of food attribute representations* (p. 2025.05.13.653572). bioRxiv. https://doi.org/10.1101/2025.05.13.653572

Charbonnier, L., van Meer, F., Johnstone, A. M., Crabtree, D., Buosi, W., Manios, Y., Androutsos, O., Giannopoulou, A., Viergever, M. A., & Smeets, P. A. M. (2018). Effects of hunger state on the brain responses to food cues across the life span. *NeuroImage*, *171*, 246–255. https://doi.org/10.1016/j.neuroimage.2018.01.012

Chen, J., Papies, E. K., & Barsalou, L. W. (2016). A core eating network and its modulations underlie diverse eating phenomena. *Brain and Cognition*, *110*, 20–42. https://doi.org/10.1016/j.bandc.2016.04.004

Cheng, Y., Meltzoff, A. N., & Decety, J. (2007). Motivation Modulates the Activity of the Human Mirror-Neuron System. *Cerebral Cortex*, *17*(8), 1979–1986. https://doi.org/10.1093/cercor/bhl107

Coricelli, C., Toepel, U., Notter, M.-L., Murray, M. M., & Rumiati, R. I. (2019). Distinct brain representations of processed and unprocessed foods. *European Journal of Neuroscience*, *50*(8), 3389–3401. https://doi.org/10.1111/ejn.14498

Delorme, A., & Makeig, S. (2004). EEGLAB: An open source toolbox for analysis of single-trial EEG dynamics including independent component analysis. *Journal of Neuroscience Methods*, *134*(1), 9–21. https://doi.org/10.1016/j.jneumeth.2003.10.009

Dienes, Z. (2011). Bayesian Versus Orthodox Statistics: Which Side Are You On?
Perspectives on Psychological Science, *6*(3), 274–290.
https://doi.org/10.1177/1745691611406920

Foroni, F., Pergola, G., Argiris, G., & Rumiati, R. (2013). The FoodCast research image
database (FRIDa). *Frontiers in Human Neuroscience*, *7*.
https://www.frontiersin.org/articles/10.3389/fnhum.2013.00051

Frank, S., Laharnar, N., Kullmann, S., Veit, R., Canova, C., Hegner, Y. L., Fritsche, A., &
Preissl, H. (2010). Processing of food pictures: Influence of hunger, gender and
calorie content. *Brain Research*, *1350*, 159–166.
https://doi.org/10.1016/j.brainres.2010.04.030

Führer, D., Zysset, S., & Stumvoll, M. (2008). Brain Activity in Hunger and Satiety: An
Exploratory Visually Stimulated fMRI Study. *Obesity*, *16*(5), 945–950.
https://doi.org/10.1038/oby.2008.33

Goddard, E., Carlson, T. A., & Woolgar, A. (2022). Spatial and Feature-selective Attention
Have Distinct, Interacting Effects on Population-level Tuning. *Journal of Cognitive
Neuroscience*, *34*(2), 290–312. https://doi.org/10.1101/530352

Goldstone, A. P., Prechtl de Hernandez, C. G., Beaver, J. D., Muhammed, K., Croese, C.,
Bell, G., Durighel, G., Hughes, E., Waldman, A. D., Frost, G., & Bell, J. D. (2009).
Fasting biases brain reward systems towards high-calorie foods. *European Journal
of Neuroscience*, *30*(8), 1625–1635. https://doi.org/10.1111/j.1460-
9568.2009.06949.x

Grootswagers, T., Robinson, A. K., & Carlson, T. A. (2019). The representational dynamics
of visual objects in rapid serial visual processing streams. *NeuroImage*, *188*, 668–
679. https://doi.org/10.1016/j.neuroimage.2018.12.046

Grootswagers, T., Robinson, A. K., Shatek, S. M., & Carlson, T. A. (2021). The neural
dynamics underlying prioritisation of task-relevant information. *Neurons, Behavior,
Data Analysis, and Theory*, 2020.06.25.172643. https://doi.org/10.51628/001c.21174

Holsen, L. M., Zarcone, J. R., Thompson, T. I., Brooks, W. M., Anderson, M. F., Ahluwalia, J.
S., Nollen, N. L., & Savage, C. R. (2005). Neural mechanisms underlying food
motivation in children and adolescents. *NeuroImage*, *27*(3), 669–676.
https://doi.org/10.1016/j.neuroimage.2005.04.043

Ilse, A., Donohue, S. E., Schoenfeld, M. A., Hopf, J. M., Heinze, H.-J., & Harris, J. A. (2020).
Unseen food images capture the attention of hungry viewers: Evidence from event-
related potentials. *Appetite*, *155*, 104828.
https://doi.org/10.1016/j.appet.2020.104828

Jain, N., Wang, A., Henderson, M. M., Lin, R., Prince, J. S., Tarr, M. J., & Wehbe, L. (2022). *Food for thought: Selectivity for food in human ventral visual cortex* [Preprint]. Neuroscience. https://doi.org/10.1101/2022.05.22.492983

Jeffreys, H. (1998). *The Theory of Probability*. OUP Oxford.

Kaiser, D., Oosterhof, N., & Peelen, M. (2016). The Neural Dynamics of Attentional Selection in Natural Scenes. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *36*, 10522–10528. https://doi.org/10.1523/JNEUROSCI.1385-16.2016

Kass, R. E., & Raftery, A. E. (1995). Bayes Factors. *Journal of the American Statistical Association*, *90*(430), 773–795. https://doi.org/10.1080/01621459.1995.10476572

Khosla, M., Ratan Murty, N. A., & Kanwisher, N. (2022). A highly selective response to food in human visual cortex revealed by hypothesis-free voxel decomposition. *Current Biology*. https://doi.org/10.1016/j.cub.2022.08.009

Killgore, W. D. S., Young, A. D., Femia, L. A., Bogorodzki, P., Rogowska, J., & Yurgelun-Todd, D. A. (2003). Cortical and limbic activation during viewing of high- versus low-calorie foods. *NeuroImage*, *19*(4), 1381–1394. https://doi.org/10.1016/S1053-8119(03)00191-5

Kriegeskorte, N., & Kievit, R. A. (2013). Representational geometry: Integrating cognition, computation, and the brain. *Trends in Cognitive Sciences*, *17*(8), 401–412. https://doi.org/10.1016/j.tics.2013.06.007

Kriegeskorte, N., Mur, M., & Bandettini, P. A. (2008). Representational similarity analysis—Connecting the branches of systems neuroscience. *Frontiers in Systems Neuroscience*, *2*. https://doi.org/10.3389/neuro.06.004.2008

Kubilius, J., Schrimpf, M., Kar, K., Rajalingham, R., Hong, H., Majaj, N., Issa, E., Bashivan, P., Prescott-Roy, J., Schmidt, K., Nayebi, A., Bear, D., Yamins, D. L., & DiCarlo, J. J. (2019). Brain-Like Object Recognition with High-Performing Shallow Recurrent ANNs. *Advances in Neural Information Processing Systems*, *32*. https://proceedings.neurips.cc/paper_files/paper/2019/hash/7813d1590d28a7dd372ad54b5d29d033-Abstract.html

LaBar, K. S., Gitelman, D. R., Parrish, T. B., Kim, Y.-H., Nobre, A. C., & Mesulam, M.-M. (2001). Hunger selectively modulates corticolimbic activation to food stimuli in humans. *Behavioral Neuroscience*, *115*(2), 493–500. https://doi.org/10.1037/0735-7044.115.2.493

Mengotti, P., Foroni, F., & Rumiati, R. I. (2019). Neural correlates of the energetic value of food during visual processing and response inhibition. *NeuroImage*, *184*, 130–139. https://doi.org/10.1016/j.neuroimage.2018.09.017

Moerel, D., Grootswagers, T., Robinson, A. K., Shatek, S. M., Woolgar, A., Carlson, T. A., & Rich, A. N. (2022). The time-course of feature-based attention effects dissociated from temporal expectation and target-related processes. *Scientific Reports*, *12*(1), 6968. https://doi.org/10.1038/s41598-022-10687-x

Moerel, D., Psihoyos, J., & Carlson, T. A. (2024). The Time-Course of Food Representation in the Human Brain. *Journal of Neuroscience*. https://doi.org/10.1523/JNEUROSCI.1101-23.2024

Moerel, D., Rich, A. N., & Woolgar, A. (2024). Selective Attention and Decision-Making Have Separable Neural Bases in Space and Time. *Journal of Neuroscience*, *44*(38). https://doi.org/10.1523/JNEUROSCI.0224-24.2024

Mogg, K., Bradley, B. P., Hyare, H., & Lee, S. (1998). Selective attention to food-related stimuli in hunger: Are attentional biases specific to emotional and psychopathological states, or are they also found in normal drive states? *Behaviour Research and Therapy*, *36*(2), 227–237. https://doi.org/10.1016/S0005-7967(97)00062-4

Mohanty, A., Gitelman, D. R., Small, D. M., & Mesulam, M. M. (2008). The Spatial Attention Network Interacts with Limbic and Monoaminergic Systems to Modulate Motivation-Induced Attention Shifts. *Cerebral Cortex*, *18*(11), 2604–2613. https://doi.org/10.1093/cercor/bhn021

Morey, R. D., Romeijn, J.-W., & Rouder, J. N. (2016). The philosophy of Bayes factors and the quantification of statistical evidence. *Journal of Mathematical Psychology*, *72*, 6–18. https://doi.org/10.1016/j.jmp.2015.11.001

Morey, R. D., & Rouder, J. N. (2018). *BayesFactor: Computation of Bayes Factors for Common Designs*. https://CRAN.R-project.org/package=BayesFactor

Nijs, I. M. T., Muris, P., Euser, A. S., & Franken, I. H. A. (2010). Differences in attention to food and food intake between overweight/obese and normal-weight females under conditions of hunger and satiety. *Appetite*, *54*(2), 243–254. https://doi.org/10.1016/j.appet.2009.11.004

Oostenveld, R., & Praamstra, P. (2001). The five percent electrode system for high-resolution EEG and ERP measurements. *Clinical Neurophysiology*, *112*(4), 713–719. https://doi.org/10.1016/S1388-2457(00)00527-7

Oosterhof, N. N., Connolly, A. C., & Haxby, J. V. (2016). CoSMoMVPA: Multi-Modal Multivariate Pattern Analysis of Neuroimaging Data in Matlab/GNU Octave. *Frontiers in Neuroinformatics*, *10*. https://doi.org/10.3389/fninf.2016.00027

Peirce, J., Gray, J. R., Simpson, S., MacAskill, M., Höchenberger, R., Sogo, H., Kastman, E., & Lindeløv, J. K. (2019). PsychoPy2: Experiments in behavior made easy. *Behavior Research Methods*, *51*(1), 195–203. https://doi.org/10.3758/s13428-018-01193-y

Pergola, G., Foroni, F., Mengotti, P., Argiris, G., & Rumiati, R. I. (2017). A neural signature of food semantics is associated with body-mass index. *Biological Psychology*, *129*, 282–292. https://doi.org/10.1016/j.biopsycho.2017.09.001

Piech, R. M., Lewis, J., Parkinson, C. H., Owen, A. M., Roberts, A. C., Downing, P. E., & Parkinson, J. A. (2009). Neural Correlates of Appetite and Hunger-Related Evaluative Judgments. *PLOS ONE*, *4*(8), e6581. https://doi.org/10.1371/journal.pone.0006581

Piech, R. M., Pastorino, M. T., & Zald, D. H. (2010). All I saw was the cake. Hunger effects on attentional capture by visual food cues. *Appetite*, *54*(3), 579–582. https://doi.org/10.1016/j.appet.2009.11.003

Placanica, J. L., Faunce, G. J., & Soames Job, R. F. (2002). The effect of fasting on attentional biases for food and body shape/weight words in high and low Eating Disorder Inventory scorers. *International Journal of Eating Disorders*, *32*(1), 79–90. https://doi.org/10.1002/eat.10066

Poggiogalle, E., Jamshed, H., & Peterson, C. M. (2018). Circadian regulation of glucose, lipid, and energy metabolism in humans. *Metabolism*, *84*, 11–27. https://doi.org/10.1016/j.metabol.2017.11.017

Read, D., & van Leeuwen, B. (1998). Predicting Hunger: The Effects of Appetite and Delay on Choice. *Organizational Behavior and Human Decision Processes*, *76*(2), 189–205. https://doi.org/10.1006/obhd.1998.2803

Robinson, A. K., Grootswagers, T., & Carlson, T. A. (2019). The influence of image masking on object representations during rapid serial visual presentation. *NeuroImage*, *197*, 224–231. https://doi.org/10.1016/j.neuroimage.2019.04.050

Rothemund, Y., Preuschhof, C., Bohner, G., Bauknecht, H.-C., Klingebiel, R., Flor, H., & Klapp, B. F. (2007). Differential activation of the dorsal striatum by high-calorie visual food stimuli in obese individuals. *NeuroImage*, *37*(2), 410–421. https://doi.org/10.1016/j.neuroimage.2007.05.008

Rouder, J. N., Speckman, P. L., Sun, D., Morey, R. D., & Iverson, G. (2009). Bayesian t
tests for accepting and rejecting the null hypothesis. *Psychonomic Bulletin & Review*,
16(2), 225–237. https://doi.org/10.3758/PBR.16.2.225

Sawada, R., Sato, W., Minemoto, K., & Fushiki, T. (2019). Hunger promotes the detection of
high-fat food. *Appetite*, *142*, 104377. https://doi.org/10.1016/j.appet.2019.104377

Schubert, E., Rosenblatt, D., Eliby, D., Kashima, Y., Hogendoorn, H., & Bode, S. (2021).
Decoding explicit and implicit representations of health and taste attributes of foods
in the human brain. *Neuropsychologia*, *162*, 108045.
https://doi.org/10.1016/j.neuropsychologia.2021.108045

Siep, N., Roefs, A., Roebroeck, A., Havermans, R., Bonte, M. L., & Jansen, A. (2009).
Hunger is the best spice: An fMRI study of the effects of attention, hunger and calorie
content on food reward processing in the amygdala and orbitofrontal cortex.
Behavioural Brain Research, *198*(1), 149–158.
https://doi.org/10.1016/j.bbr.2008.10.035

Simmons, W. K., Martin, A., & Barsalou, L. W. (2005). Pictures of Appetizing Foods Activate
Gustatory Cortices for Taste and Reward. *Cerebral Cortex*, *15*(10), 1602–1608.
https://doi.org/10.1093/cercor/bhi038

Stingl, K. T., Rogić, M., Stingl, K., Canova, C., Tschritter, O., Braun, C., Fritsche, A., &
Preissl, H. (2010). The temporal sequence of magnetic brain activity for food
categorization and memorization—An exploratory study. *NeuroImage*, *52*(4), 1584–
1591. https://doi.org/10.1016/j.neuroimage.2010.04.278

Stockburger, J., Schmälzle, R., Flaisch, T., Bublatzky, F., & Schupp, H. T. (2009). The
impact of hunger on food cue processing: An event-related brain potential study.
NeuroImage, *47*(4), 1819–1829. https://doi.org/10.1016/j.neuroimage.2009.04.071

Stockburger, J., Weike, A. I., Hamm, A. O., & Schupp, H. T. (2008). Deprivation selectively
modulates brain potentials to food pictures. *Behavioral Neuroscience*, *122*(4), 936–
942. https://doi.org/10.1037/a0012517

Sultson, H., Vainik, U., & Kreegipuu, K. (2019). Hunger enhances automatic processing of
food and non-food stimuli: A visual mismatch negativity study. *Appetite*, *133*, 324–
336. https://doi.org/10.1016/j.appet.2018.11.031

Teichmann, L., Moerel, D., Baker, C. I., & Grootswagers, T. (2022). An empirically-driven
guide on using Bayes Factors for M/EEG decoding. *Aperture Neuro*, *1*(8), 1–10.
https://doi.org/10.1101/2021.06.23.449663

Toepel, U., Knebel, J.-F., Hudry, J., le Coutre, J., & Murray, M. M. (2009). The brain tracks
the energetic value in food images. *NeuroImage*, *44*(3), 967–974.
https://doi.org/10.1016/j.neuroimage.2008.10.005

Tsourides, K., Shariat, S., Nejati, H., Gandhi, T. K., Cardinaux, A., Simons, C. T., Cheung,
N.-M., Pavlovic, V., & Sinha, P. (2016). Neural correlates of the food/non-food visual
distinction. *Biological Psychology*, *115*, 35–42.
https://doi.org/10.1016/j.biopsycho.2015.12.013

Tuorila, H., Kramer, F. M., & Engell, D. (2001). The choice of fat-free vs. regular-fat fudge:
The effects of liking for the alternative and the restraint status. *Appetite*, *37*(1), 27–
32. https://doi.org/10.1006/appe.2001.0410

Uher, R., Treasure, J., Heining, M., Brammer, M. J., & Campbell, I. C. (2006). Cerebral
processing of food-related stimuli: Effects of fasting and gender. *Behavioural Brain
Research*, *169*(1), 111–119. https://doi.org/10.1016/j.bbr.2005.12.008

van der Laan, L. N., de Ridder, D. T. D., Viergever, M. A., & Smeets, P. A. M. (2011). The
first taste is always with the eyes: A meta-analysis on the neural correlates of
processing visual food cues. *NeuroImage*, *55*(1), 296–303.
https://doi.org/10.1016/j.neuroimage.2010.11.055

Vignando, M., Aiello, M., Rinaldi, A., Cattarruzza, T., Mazzon, G., Manganotti, P., Eleopra,
R., & Rumiati, R. I. (2019). Food knowledge depends upon the integrity of both
sensory and functional properties: A VBM, TBSS and DTI tractography study.
Scientific Reports, *9*(1), 7439. https://doi.org/10.1038/s41598-019-43919-8

Wagenmakers, E.-J. (2007). A practical solution to the pervasive problems ofp values.
Psychonomic Bulletin & Review, *14*(5), 779–804.
https://doi.org/10.3758/BF03194105

Wetzels, R., Matzke, D., Lee, M. D., Rouder, J. N., Iverson, G. J., & Wagenmakers, E.-J.
(2011). Statistical Evidence in Experimental Psychology: An Empirical Comparison
Using 855 t Tests. *Perspectives on Psychological Science*, *6*(3), 291–298.
https://doi.org/10.1177/1745691611406923