1 **Single capsid mutations modulating phage adsorption, persistence, and plaque**
2 **morphology shape evolutionary trajectories in ΦX174**
3

4 Short title:

5 **Life history evolution in ΦX174**

6
7 Key words:
8 **Adsorption, persistence, ΦX174, lytic bacteriophage, phage therapy, phage evolution**
9

10 Manuela Reuter[1], Michael Sieber[1], Octavio Reyes-Matte[1], Christina Vasileiou[1], Christopher
11 Böhmker[1], Jordan Romeyer Dherbey[2], Frederic Bertels[1,*] and Javier Lopez-Garrido[1,*]
12
13 [1]Max Planck Institute for Evolutionary Biology, Plön, Germany
14 [2]University of Cambridge, Cambridge, United Kingdom
15
16 *Correspondence: frederic.bertels.publications@gmail.com; lopezgarrido@evolbio.mpg.de

17
18

**Abstract**

The evolutionary success of lytic bacteriophages depends on key life-history traits, including

adsorption, lysis time, burst size and persistence in the environment. However, how these traits

evolve to allow adaptation to different environments remains poorly understood. Here, we

explored this question in phage $\Phi$X174, combining experimental evolution and mathematical

modelling. By investigating how serial transfer conditions shape evolutionary outcomes in

liquid culture, we found that the time between transfers imposes divergent selection on

adsorption and context-dependent directional selection on persistence. Longer transfer

intervals, which allow multiple infection cycles until host depletion, favoured fast-adsorbing,

highly persistent mutants that could rapidly initiate infections and remained viable in the

absence of the host. In contrast, shorter transfer intervals selected for slower adsorption without

substantially altering persistence. Mathematical modelling of phage infection dynamics

revealed that slow adsorption reduces the likelihood of irreversible phage attachment to already

infected cells, thereby increasing the number of free phages able to initiate new infections upon

dilution. These trade-offs resulted in distinct plaque phenotypes on solid media: fast adsorbers

formed small plaques, while slow adsorbers formed large ones. A single-point mutation in the

major capsid protein was sufficient to drive these changes in adsorption. In the case of fast-

adsorbing mutants, this mutation was pleiotropic and also enhanced environmental persistence.

Our findings show how simple changes in propagation conditions can steer phage phenotypes,

providing insights relevant to evolutionary biology and phage therapy.

## Introduction

The success of lytic bacteriophages relies on four central life-history traits, each of which may evolve under different selective pressures to maximise fitness [1–6]. These traits are: (I) adsorption constant, which determines the ease with which a phage irreversibly attaches to a host cell, (II) latent period, or time required to complete the infection cycle inside the host, (III) burst size, defined as the number of new phage particles produced per infected host cell, and (IV) decay rate, representing the loss of infectious phage particles in the absence of susceptible host cells [7–12].

Interest in these traits has grown due to the need to predict phage virulence for medical and biocontrol applications [13–16]. However, such predictions remain challenging because the fitness effects of individual traits are often highly context-dependent [17–19]. For example, high adsorption rates are typically considered beneficial in well-mixed liquid environments, where they facilitate rapid infection initiation [20, 21]. However, in spatially-structured environments such as biofilms, these same traits can hinder phage spread to uninfected neighbouring cells by promoting phage attachment to already infected cells at the centre of infection [17, 22]. Understanding how life-history traits influence phage fitness across environments is therefore essential for predicting and potentially optimising phage performance.

The model coliphage ΦX174 provides a powerful platform for dissecting how the relationship between trait and environment impacts fitness. [23–27]. ΦX174 is a strictly lytic, tailless phage of the *Microviridae* family, best known as the first DNA-based organism whose entire genome was sequenced [28]. Its 5386 bp circular single-stranded DNA genome encodes only 11 genes, most of which are necessary for phage DNA replication, morphogenesis and host lysis [28–31]. Only four genes (F, G, H, and J) encode for proteins that form the mature virion [32, 33]. The icosahedral capsid is made of 60 F, 60 J and 12 H protein monomers, decorated at each of its 12 vertices by 5 G spike monomers [32, 34]. G likely mediates host recognition and subsequent attachment to host lipopolysaccharide (LPS), after which it dissociates, allowing F to bind irreversibly to the host cell surface [35–37]. Irreversible binding triggers a conformational change that results in the injection of phage DNA into the host cell through the pilot protein H [37, 38].

Here, we have explored how life-history traits influence phage fitness across different
environments by evolving ΦX174 under contrasting serial transfer regimes. Through a
combination of life-history trait quantification and mathematical modelling, we have identified
individual mutations that alter adsorption constant and persistence, leading to fitness effects
that are strongly dependent on the environment. By disentangling these context-dependent
fitness effects, our results improve our understanding of phage ecology and evolution and also
provide insights for rational trait optimisation in medical and biocontrol applications.
80
81

## Materials and Methods

### Strains and culture conditions

The ancestral ΦX174 and its host *E. coli* C were provided by H.A. Wichman (Department of Biological Sciences, University of Idaho). All evolved phage strains were derived from the same ΦX174 genotype (GenBank: AF176034). Cultures were grown in Lysogeny Broth (LB, Miller) supplemented with 10 mM $MgCl_2$ and 5 mM $CaCl_2$ and incubated at 37°C, shaking at 250 rpm (orbital shaking diameter 2.5 cm). Mid-exponential *E. coli* C cultures (~$2*10^8$ colony-forming units [CFU] per ml) were prepared from overnight cultures and used for all infection experiments.

### Preparation of phage lysates and quantification

Phage lysates were generated as described in Romeyer Dherbey *et al*. [39] by infecting exponentially growing *E. coli* C with ΦX174 for 3 hours. After 3 hours, the remaining bacteria were removed with chloroform and subsequent centrifugation. Phage-containing supernatants were collected and stored for assays at 4°C or at -80°C for long-term storage.

Phage titers were quantified by plaque assays in which plaque-forming units (PFUs) were assessed by plating 100 µl of diluted phage lysate with 100 µl of *E. coli* C overnight culture pre-mixed in 4 ml semi-solid agar (SSA, 0.5% agar, supplemented with 10 mM $MgCl_2$ and 5 mM $CaCl_2$). The mixture was overlayed on a plate of LB-agar (1.5%) and incubated at 37°C for 18 hours. Phage lysates were diluted in phage buffer juice (PBJ: 2.03% Tris-HCl, 0.61% $MgCl_2$).

### Serial transfer regimes

For every transfer regime, five independent selection lines were initiated by infecting mid-exponential phage-sensitive *E. coli* C cultures with individual ancestral ΦX174 extracted from independent plaques. Consecutive transfers were inoculated from phage lysate at $MOI_{input}$ ~0.1. The 3-hour regime involved 3-hour infections followed by chloroform phage extraction. Serial transfers continued until heritable small-plaque mutants emerged, which were isolated and replated multiple times to confirm the phenotype. The 30-min regime involved four daily 30-minute transfers followed by one 3-hour amplification step, repeated for ten days. Individual plaques were isolated at the end for sequencing. An additional line was propagated

using only 30-minute transfers, without the amplification step. Phages were extracted with
chloroform and transferred after each round at $MOI_{input}$ ~0.1 for 40 passages. Population
sequencing was used to monitor mutation dynamics. Additional details can be found in **Fig. S1**
**and Supplementary Methods**.

**Sanger sequencing of evolved phages**

Whole-genome Sanger sequencing was conducted for individual phage isolates from
both transfer regimes according to Biggs *et al*. [40]. DNA was extracted from high-titer lysates
(QIAprep Spin Miniprep Kit® Qiagen, Hilden, Germany) and amplified via PCR in two
genome halves. PCR products were purified (QIAprep PCR Purification Kit® Qiagen, Hilden,
Germany) and sequenced at the Max Planck Institute for Evolutionary Biology. Primers and
settings used for amplification and sequencing are in **Tables S1, S2, and S3**. Sequences were
aligned against the reference ΦX174 genome using Geneious Prime 2023.0.1
(https://www.genious.com). Identified amino acid positions in which single-point mutations
were located were numbered by including the start codon as position one [39].

**Adsorption assays**

Adsorption constants were determined following a modified version of the Hyman and
Abedon assay [41]. Mid-exponential, phage-sensitive *E. coli* C cultures were infected with
phages at $MOI_{input}$ ~0.05 at 37°C in a shaking water bath at 200 rpm. Phages were sampled 2,
4, and 6 minutes post-infection and diluted 10-fold into non-supplemented LB. Samples were
cooled on ice to stop infection, treated with chloroform, and quantified. Adsorption constants
were calculated by fitting linear models to log-transformed phage titers and dividing slope
estimates by bacterial density. Differences between genotypes were tested using a linear mixed
model (replicate as random effect) and post hoc Tukey tests for multiple comparisons
(**Supplementary Methods**).

**Phage persistence assays**

Persistence was measured over a 5-hour infection period. An exponentially-growing *E.*
*coli* C culture was infected at $MOI_{input}$ ~0.1, and samples were taken every 30 minutes. Mature
phages were extracted with chloroform, and phage titers were determined by PFU counting.

Decay rates were estimated by fitting exponential decay models to the titer decline period after
peak productivity. Model fitting procedures are detailed in **the Supplementary Methods**.

**Simulations of phage evolution**

The following system of delay differential equations (DDEs) describes the dynamics of
the bacterial host $B$ (cells per ml), the infected bacterial population $B_I$ (cells per ml), and the
three phage types $P_A$ (ancestral, particles per ml), $P_L$ (large-plaque type, particles per ml), and
$P_S$ (small-plaque type, particles per ml). In addition, $B_\tau = B(t - \tau)$ denotes the bacterial
density $\tau$ minutes in the past, reflecting the delay introduced by the lysis time. We use a similar
notation for the phage densities, e.g. $P_{A,\tau}$ denotes the density of the ancestral phage $\tau$ minutes in
the past.

Bacterial growth throughout the transfer period was found not to play a significant role
in the simulations, so it is neglected in the model for simplicity. All parameter values and their
biological meaning are given in **Table 1** in the main text.

$$\frac{dB}{dt} = -(\phi_A P_A + \phi_L P_L + \phi_S P_S)B$$

$$\frac{dB_I}{dt} = \phi_A(BP_A - B_\tau P_{A,\tau}) + \phi_L(BP_L - B_\tau P_{L,\tau}) + \phi_S(BP_S - B_\tau P_{S,\tau})$$

152
$$\frac{dP_A}{dt} = \phi_A\left((1-m)\beta B_\tau P_{A,\tau} - (B + B_I)P_A\right) + \frac{1}{2}m\beta B_\tau(\phi_L P_{L,\tau} + \phi_S P_{S,\tau}) - \delta_A P_A$$

$$\frac{dP_L}{dt} = \phi_L\left((1-m)\beta B_\tau P_{L,\tau} - (B + B_I)P_L\right) + \frac{1}{2}m\beta B_\tau(\phi_A P_{A,\tau} + \phi_S P_{S,\tau}) - \delta_L P_L$$

$$\frac{dP_S}{dt} = \phi_S\left((1-m)\beta B_\tau P_{S,\tau} - (B + B_I)P_S\right) + \frac{1}{2}m\beta B_\tau(\phi_A P_{A,\tau} + \phi_L P_{L,\tau}) - \delta_S P_S$$

The DDE system was solved numerically in Python using the *scipy*-based *ddeInt* solver.
After 30 minutes or 3 hours of simulated time, depending on the transfer regime, the system
was reset to the initial density of fresh bacteria $B$ ($10^8$ CFU), and 2% of the final densities of
infected bacteria $B_I$ and all three phage types, respectively. This starts the next growth cycle
and corresponds to the transfer protocol of the experiment.

**Table 1: Parameters and corresponding parameter values used in the mathematical model.**

| Parameter | Value |
|---|---|
| Adsorption constant, ancestor | $4.15*10^{-9}$ (ml$^{-1}$ min-1) |
| Adsorption constant, large-plaque type | $9.87*10^{-10}$ (ml$^{-1}$ min-1) |
| Adsorption constant, small-plaque type | $6*10^{-9}$ (ml$^{-1}$ min-1) |
| Burst size, all types | 200 (per infected host cell) |
| Lysis time, all types | 17 min |
| Decay rate, ancestor | $1.7*10^{-2}$ (min$^{-1}$) |
| Decay rate, large-plaque type | $1.4*10^{-2}$ (min$^{-1}$) |
| Decay rate, small-plaque type | $6*10^{-3}$ (min$^{-1}$) |
| Mutation rate | $10^{-5}$ (per minute) |

## Image analysis of lysis plaques

Lysis plaque morphology was quantified using a custom image analysis pipeline.
Plaques were imaged under standardised conditions (ChemiDoc: Bio-Rad, California, USA;
0.1s exposure time) and processed using Ilastik [42] for pixel classification based on intensity,
texture, and edge features. Post-processing steps included erosion, whole-filling, and
watershed transformation using SciPy [43] and scikit-image [44]. Each plaque was assigned
an ID and analysed for area (cm²), distance to periphery (via distance transform) and grayscale
intensity (overlaying mask on original image) (details see **Supplementary Methods**).

## Data analysis and visualisation

Statistical analysis of adsorption constants, model fitting for adsorption and decay and
all data visualisations were performed in R (version 4.2.1) [45]. Details of the packages are
found in **Table S13**. The protein structure, resolved by McKenna *et al*. [32], shown in **Fig. 5A**,
was taken from RCSB PDB (reference 2BPA:
https://www.wwpdb.org/pdb?id=pdb_00002bpa), and the mutation visualisation was done in
UCSF Chimera (version 1.18). All cartoons were created using BioRender® as indicated in the
corresponding figure legends.

**Results**

**Differences in transfer intervals during serial passaging drive the evolution of contrasting lysis plaque phenotypes**

In the context of a previous study [39], we observed that small and turbid plaques frequently emerged when ΦX174 was serially transferred after 3-hour incubation periods with host cells in liquid culture. To confirm this observation, we evolved multiple independent selection lines of the ancestral ΦX174 under the same selection regime (hereafter, 3-h regime) and plated the evolving phage populations onto lawns of phage-sensitive host cells at every transfer (**Fig. 1; Fig. S1A**). Small, turbid plaques appeared in all lines after four to seven transfers (**Fig. 1; Fig. S2**).

**Fig. 1: 30-min and 3-h transfer regimes select for large- and small-plaque-forming phages, respectively.** Phage ΦX174 was propagated under 30-min (top) or 3-h (bottom) regimes, leading to the evolution of large- and small-plaque-forming phages, respectively. Ancestral plaques are shown on the left, and evolved large (top) and small (bottom) plaques on the right. Scale bars, 1 cm. All plaque images have the same magnification. Diagrams of the 30-min and 3-h regimes are in the middle. The approximate input multiplicity of infection (MOI$_{input}$) and incubation times are the same for each transfer. For illustration purposes, only four transfers are shown, but the experiment involved more transfers. See methods and **Fig. S1** for details. Figure was created in BioRender. Reuter, M. (2025) https://BioRender.com/20imrdd.

Since differences in plaque morphology often reflect underlying changes in life history

traits [2, 17, 22, 46–48], we hypothesised that our experimental regime was selecting for phages

with changes in one or more life history traits. The 3-h regime provides sufficient time for

phages to complete multiple infection cycles, potentially depleting host cells and altering

selective pressures. To test whether transfer length affects plaque phenotype evolution, we

performed a second evolution experiment in which phages were serially transferred every 30

minutes (hereafter, referred to as the 30-min regime) (**Fig. 1, Fig. S1B**), a timeframe during

which most phages can complete only a single infection cycle (**Fig. S3**). In contrast to the 3-h

transfer regime, no small plaques were detected under the 30-min regime, even after 40

transfers (**Fig. 1**). However, to our surprise, we began to observe plaques larger than those

formed by the ancestral phage after approximately 20 transfers (**Fig. 1; Fig. S2**).

**Large- and small-plaque phenotypes are caused by different mutations in the major**

**capsid (F) protein**

To identify the genetic modifications leading to the small- and large-plaque phenotypes,

we isolated five phages of each type, one per selection line, and sequenced their genomes. Each

evolved isolate differed from the ancestor by only one or two single-point mutations (**Fig. 2**).

All five large-plaque mutants isolated from the 30-min regime lines carried an A-to-G

transition at position 1301 in the phage genome, resulting in a threonine-to-alanine substitution

at position 101 of the F gene [A1301G/F(T101A)] (**Fig. 2A**). The mutant isolated from line 3

carried only this substitution. The other four isolates carried one additional mutation in either

the F or the H protein (**Fig. 2A**) acquired after the F(T101A) mutation (see Sanger Sequencing,

Zenodo: 10.5281/zenodo.17076221). Phenotypically, however, all large-plaque mutants

resembled each other (**Fig. 2A, Fig. S4**), suggesting that the T101A mutation in the F protein

is sufficient for the manifestation of the large-plaque phenotype.

None of the mutations observed in large-plaque mutants were found in the small-

plaque-forming phages isolated from the 3-h regime lines. However, all five small-plaque

phages also differed from the ancestor by a single-point mutation in the F gene (**Fig. 2B**), each

producing a distinct small-plaque phenotype (**Fig. 2B and S4**). The most drastic phenotype

was caused by the G1965A/F(G322D) mutation, resulting in the formation of extremely small

and turbid plaques. Phages carrying the C2280A/F(S427*) mutation, which truncates the major

capsid protein by the last amino acid, produced slightly larger and clearer plaques (**Fig. S4**).

The G1639A/F(M213I) mutation led to a heterogeneous phenotype, with both small and large plaques, and was excluded from further analysis.

**Fig. 2: Mutations linked to large- and small-plaque phenotypes occur in the major capsid protein.** (A) Large-plaque phenotypes (top) and corresponding genotypes (bottom) of mutants isolated from independent lines of the 30-min regime. All the mutants carried an A-to-G transition at position 1301 (A1301G) in the phage genome, resulting in a threonine-to-alanine substitution at position 101 of the F protein [F(T101A)]. Four isolates carried an additional mutation, either in the F (green) or the H (blue) gene. The ancestor is shown in grey (WT). (B) Small-plaque phenotypes (top) and genotypes (bottom) of mutants isolated from independent lines of the 3-h regime. All mutants differed from the ancestor by different mutation in the F gene (green), resulting in different amino acid changes: G322D in the clones from lines 1, 2 and 5; M213I in the clone from line 3; and the formation of a premature stop codon at the very last codon of the F gene (S427*) in the clone from line 4. The nucleotide change in the phage genome is shown at the top, and the corresponding amino acid change at the bottom. The ancestor is shown in grey (WT). Scale bars, 1 cm. Genome illustrations were generated using Geneious Prime 2023.0.1. Figure created in BioRender. Reuter, M. (2025) https://BioRender.com/du5tsj2.

251    Altogether, the sequencing results indicate that different modifications of the major
252 capsid protein are responsible for the contrasting plaque phenotypes emerging in our selection
253 regimes.

**Plaque size negatively correlates with the adsorption constant**

The major capsid protein is likely essential for the irreversible adsorption of ΦX174 to
the host cells [37], suggesting the large- and small-plaque-forming phages may have evolved
different adsorption constants compared to the ancestral ΦX174. To test this hypothesis, we
measured the adsorption constant of the ancestral phage, a large-plaque-forming mutant
carrying the single F(T101A) mutation, and the two small-plaque mutants, F(G322D)
and F(S427*). We focused on mutants with single substitutions to ensure that each mutation
could be directly linked to specific changes in life history traits.

The adsorption constants of the evolved phages differed significantly from that of the
ancestral ΦX174 and correlated inversely with plaque size (**Fig. 3**). The ancestral mean
adsorption constant was $4.15 \times 10^{-9}$ ml$^{-1}$ min$^{-1}$ (**Table S4**), in close agreement with previous
estimations [49, 50]. The adsorption constants of the small-plaque-forming mutants were
slightly but significantly higher than the ancestral. Conversely, the large-plaque mutant had an
absorption rate constant approximately five-fold lower than the ancestral (**Table S4**).

The relationship between adsorption constant and plaque size was not linear. The large-
plaque mutant with a five-fold reduction in adsorption constant produced plaques that were
1.5-fold larger than those of the ancestor, whereas the small-plaque mutants with less than two-
fold increase in adsorption constant formed plaques about 10 times smaller (**Table S4**). These
results are in line with previous predictions of a non-linear relationship between adsorption
constant and plaque size [48].

These results demonstrate that the adsorption constants evolved in opposite directions
under the 3-h and 30-min regimes, and suggest that the differences in plaque size may be caused
by variation in phage-host cell binding affinity.

**Fig. 3: The adsorption constant is inversely correlated with plaque size.** (A) Adsorption constant "k" of the

ancestor (grey), the large-plaque mutant F(T101A) (purple) and the two small-plaque mutants F(G322D) (orange)

and F(S427*) (green). At least three independent experiments were performed for each genotype. Each black dot

represents the adsorption constant of an individual experiment, the horizontal lines indicate the median, and the

boxes span the middle 50% of the data. The whiskers span the lower and upper 25% of the data. Different letters

on top of the boxes indicate significant adsorption constant differences between genotypes (significance level

$p<0.05$). (B) Lysis plaque area in $cm^2$ (y-axis) for the ancestor (grey, n=76), and the evolved phages F(T101A)

(purple, n=60), F(G322D) (green, n=303) and F(S427*) (orange, n=431). Violin plots show distribution shape;

boxplots display medians (black lines), interquartile range (boxes), and upper/lower 25% (whiskers). See

**Methods** and **Supplementary Methods** for details of adsorption and plaque phenotype analysis.

**Fast adsorption is detrimental in the 30-min regime**

We expected fast adsorption to be beneficial in the well-mixed liquid environment in

which our experimental evolution regimes took place, as it would allow rapid infection of host

cells [20, 21]. Hence, we were surprised that phages with dramatically reduced adsorption

constants were selected under the 30-min regime. To further explore the role of absorption rate

in phage fitness under different transfer intervals, we developed a mathematical model of phage

infection dynamics (**Fig. 4**).

The model captures the interaction between the bacterial host, the ancestral phage, the

slow-adsorbing mutant, and the fast-adsorbing mutant during the transfer experiments. Upon

adsorption of a phage to an uninfected host cell, the cell moves into the infected class. Phages

of all types keep adsorbing to already infected cells, but only phages corresponding to the type

that infected first are released upon lysis. We explicitly consider the lysis time, reflecting the

time delay between adsorption of a phage and lysis of the infected cell, triggering the release

of new viral particles. The full model is given in the **Supplementary Methods**.

**Fig. 4: Simulations of the 30-min regime.** (A) Simulated dynamics of free phage populations (y-axis) across

serial transfers (x-axis) in the 30-min regime, using genotypes with experimentally measured adsorption constants

(green: mutant F(G322D); purple: mutant F(T101A); grey: ancestor). The simulation predicts the rapid emergence

of the low-adsorption mutant, in line with experimental results. (B) Predicted optimal adsorption constants based

on simulated competitions between the ancestor and hypothetical mutants with lower adsorption constants (x-

axis), starting from an initial 1:1 ratio. The y-axis shows the predicted mutant-to-ancestor titer ratios after one

transfer. The evolved mutant (purple line) closely matches the predicted optimum. (C) Simulation of free phage

densities (y-axis) within four (T1-T4) 30-min transfer (x-axis) for the ancestor (grey) and large-plaque mutant

F(T101A) (purple). The free phage dynamics within one transfer predict a higher number of free phages of the

low-adsorption genotype F(T101A) at the end of a transfer. The high loss of comparably fast-adsorbing ancestral

phages to dead-end adsorption at the moment of transfer results in the selection of mutant F(T101A). (D) Bar

chart of experimentally measured free phages of mutant F(T101A) (purple bars) relative to ancestral free phages

(y-axis) after a 30-min transfer for a total of four transfers (x-axis). The number of free phages of mutant F(T101A)

was normalised for the free phages of the ancestor, here indicated by the dashed black line. The height of the bar

represents the mean of three independent biological replicates, with error bars indicating the standard error.

We started the *in silico* transfer experiments with phages with the ancestral adsorption

constant. We allowed for the appearance of mutations that increased or decreased the

adsorption constant to the values measured experimentally for the fast-adsorbing mutant

F(G322D) and the slow-adsorbing mutant F(T101A), respectively. All other life-history traits

were held constant (**Table 1**). The simulations predicted the evolution of the slow-adsorbing

mutant under the 30-min regime within a timescale compatible with our experimental

observations (**Fig. 4A; Fig. S2**). Note that the 30-min transfer regime was interrupted by a 3-h

incubation step after every fourth transfer to amplify the phage population (see Materials

&Methods, **Fig. S1**). Although this step slowed down the increase in the frequency of slow-

adsorbing mutants, it did not prevent their eventual fixation in the simulations.

To further investigate the benefit of low adsorption constants during 30-min transfers,

we simulated competition between the ancestral phage and hypothetical mutants with

adsorption constants ranging from the ancestor value to 1000-fold lower (**Fig. 4B**). Each

simulation began with a 1:1 ratio of ancestral and mutant phages at an input multiplicity of

infectiousness ($MOI_{input}$) of 0.1, and predicted the relative mutant titer after a single 30-min

transfer. The results revealed an optimal adsorption constant approximately 6-fold lower than

that of the ancestor, very close to the value observed in the F(T101A) mutant. Mutants with

very low adsorption constants were predicted to perform poorly due to limited infection

initiation, whereas mutants with very high adsorption constants were expected to also perform

poorly, as their rapid binding to host cells increased the risk of multiple attachment to one host

cell, resulting in a high number of phages being lost to "dead-end" infections.

A detailed analysis of the simulations revealed that, in the 30-min regime, the phage

adsorption constant influenced the number of free phages present at the moment of transfer

(**Fig. 4C**). For fast-adsorbing phages, the model predicted that a large fraction of phages would

be lost to dead-end adsorptions after the first infection cycle, due to a rapid MOI increase. In

contrast, slow-adsorbing phages remained largely unbound, resulting in a higher proportion of

free phages at the moment of transfer. This was beneficial, as free phages could readily infect

sensitive host cells in the subsequent cycle. Quantification of free phages using sterile filtration

after each 30-min transfer confirmed these predictions. The number of free phages was

consistently higher for slow-adsorbing mutants, whose adsorption constants were closer to the

simulated optimum, than for ancestral phages (**Fig. 4D**). These results indicate that reduced

adsorption constants can increase phage fitness under the 30-min regime by increasing the pool

of phages available to initiate new infections in the next cycle.

**3-h transfers select for high persistence outside the host**

In contrast to the 30-min regime, simulation under the 3-h regime predicted the eventual evolution of fast-adsorbing, small-plaque mutants, consistent with our experimental observations (**Fig. S5**). However, the predicted timescale was substantially longer than that observed experimentally (**Fig. S2**). This suggests that, although increased adsorption was beneficial in the 3-h regime, selection on adsorption alone was insufficient to explain the evolution of the observed mutants.

To get insight into which additional life history traits might promote the rapid selection of small-plaque mutants under the 3-h regime, we mapped the mutations underpinning the small- and large-plaque phenotypes onto the ΦX174 F-protein structure. The large-plaque-forming F(T101A) substitution was located in the centre of the F monomer. However, the two small-plaque-associated mutations, F(G322D) and F(S427*), were at the peripheral interfaces between F monomers in the assembled capsid (**Fig. 5A**). These peripheral mutations might influence capsid symmetry or stability by altering inter-subunit interactions, potentially contributing to the persistence of small-plaque-forming mutant phages during extended incubation periods.

To test this, we performed time-course infections using the ancestral ΦX174, the large-plaque mutant, and the two small-plaque mutants. Infections were initiated at an MOI$_{input}$ of ~0.1, and infectious phage titers were quantified at regular intervals over five hours following chloroform extraction. All populations exhibited an initial production phase, during which phage particles increased, followed by a decay phase after the depletion of susceptible host cells (**Fig. S6**). The production period differed slightly between the different phages. The ancestor and the large-plaque-forming mutant reached their maximum phage titer ~60 minutes post-infection, with a ~5,000-fold increase in population size. Small-plaque mutants peaked slightly later but achieved a 10,000-fold increase in population size. The higher population size peak of small-plaque mutants may be attributed to their higher adsorption constant, allowing for earlier infection initiation. The most striking differences, however, were observed during the decay phase (**Fig. 5B**). Both the ancestor and the large-plaque-forming mutant showed a drastic reduction in infective phage particles, with titers declining more than 15-fold by the end of the experiment. In contrast, the small-plaque mutants maintained a relatively constant number of infective particles throughout the remainder of the assay, showing only a minor decline over time.

**Fig. 5: Single-point mutations in the major capsid protein modulate adsorption and persistence.** (A)
Location of point mutations linked to plaque size differences in the F protein monomer (left) or multimer (right)
structure. Mutations F(G322D) and F(S427) leading to small plaque phenotypes are shown in green and orange,
respectively. Mutation F(T101) leading to a large plaque phenotype is shown in purple. Mutations associated with
the small plaque phenotype are localised at monomer-monomer interfaces, whereas mutation F(T101) is located
at the F protein centre. The protein structure, resolved by McKenna *et al*. [32], was taken from RCSB PDB
(reference 2BPA: https://www.wwpdb.org/pdb?id=pdb_00002bpa). The figure was prepared in UCSF Chimera
(version 1.18). (B) The y-axis shows the ratio of remaining particles relative to peak titer for the ancestor (grey),
large-plaque mutant T101A (purple), and small-plaque mutants G322D (green) and S427* (orange). One
experimental run, representative of multiple replicates, is shown. See **Supplementary Methods** for model
selection and parameter extraction. (C) Simulations of free phage population dynamics (y-axis) across transfers
(x-axis) for the 30-min (top) and 3-h (bottom) regimes, incorporating measured adsorption and decay for the
ancestor (grey), T101A (purple), and G322D (green). Including decay improved prediction accuracy, especially
under 3-h conditions, where it predicted the correct timing of small-plaque mutant evolution.

414    To test if differences in decay rates could explain the rapid evolution of the small-
415 plaque mutants in the 3-h regime, we incorporated the experimentally observed decay
416 parameters into our in silico serial transfer model (**Table 1**). In this case, the model predicted
417 the emergence of small-plaque mutants with enhanced persistence on a timescale consistent
418 with experimental observations (**Fig. 5C**). In contrast, adding persistence to simulations of the
419 30-min regime had minimal impact, slightly extending the predicted fixation time of the large-
420 plaque mutant and improving alignment with experimental observations (**Fig. 4A; Fig. 5C and**
421 **Fig. S2**).

**Discussion**

Our results reveal context-dependent fitness effects of persistence and adsorption in

ΦX174, modulated by the transfer time in serial passaging experiments. Lower adsorption

constants increase fitness under short transfer times. In contrast, longer transfer times select for

phages with higher adsorption constants and greater ability to persist outside the host.

These patterns can be explained by the ecological dynamics specific to each regime

(**Fig. 6**). In the 3-h regime, multiple infection cycles can occur per transfer. Fast-adsorbing

phages initiate these cycles more efficiently. Moreover, since phage populations increase by

more than two orders of magnitude per infection cycle (**Fig. S3**) and transfers are performed at

an $MOI_{input}$ of ~0.1 phages per bacterial cell, the vast majority of sensitive host cells are lysed

after two or three cycles, forcing phages to persist in the absence of host cells for extended

periods before being supplied with new susceptible hosts (**Fig. S6**). Thus, both enhanced

adsorption and environmental persistence are selected in this regime.

In contrast, under the 30-min regime, host cells are not exhausted between transfers.

Here, fast-adsorbing phages are more likely to adsorb to already infected cells (**Fig. 4**), leading

to unproductive infections due to superinfection inhibition or intracellular resource competition

[11, 40, 51]. Our simulations show that in these conditions, the early and frequent initiation of

infections by fast-adsorbing phages does not offset the losses incurred from unproductive

adsorption events (**Fig. 4C**). In contrast, slow-adsorbing phages are less likely to attach to

already infected cells and are more prone to being transferred as free phages. Upon transfer,

they encounter a fresh pool of uninfected hosts, allowing them to initiate productive infections

and outcompete their fast-adsorbing counterparts (**Fig. 6**). These findings challenge the

common assumption that high adsorption constants are universally advantageous in well-

mixed environments [20, 21].

Both adsorption and persistence are influenced by mutations in the major capsid protein

F, which likely plays a key role in irreversible adsorption to the host cell [32, 37]. The T101A

substitution, associated with a low adsorption constant, is located in an α-helix at the centre of

the F protein. Previous studies have identified this region as key to host specificity and binding

affinity [23, 25, 27, 39, 52, 53]. For example, Pepin *et al*. [53] identified a threonine-to-serine

substitution in the same residue that enabled phages to infect host cells with modified LPS.

Similarly, Crill *et al*. [25] described a glycine-to-aspartate substitution at position 102 that

drastically reduced adsorption to *E. coli* C but increased binding affinity to *Salmonella*, highlighting the role of this region in attachment to the host.

**Fig. 6: Schematic overview of phage population dynamics during 30-min and 3-h transfers.** In both panels, low-adsorption and -persistence phages are shown in orange and corresponding infected host cells in brown. High-persistence and -adsorption phages in blue and their infected hosts in light blue. Uninfected hosts are shown in yellow. (A) In the 30-min regime, mutants with low adsorption constants (orange) are selected, as they are transferred at high rates as free phage particles due to their delayed adsorption. Free phages can immediately initiate new infection cycles when they are transferred to fresh, phage-sensitive hosts. In contrast, phages with high adsorption constants have a lower chance of being transferred as free phage particles and engage in unproductive infections by attaching to already infected cells. (B) In the 3-h regime, phages are left without susceptible host cells (persistence period) after 2-3 infection cycles (growth period). High persistence phage mutants, which also have a high adsorption constant, are selected under these conditions due to both their faster adsorption and slower decay. In this case, high adsorption is beneficial as it allows phages to more effectively scavenge sensitive host cells. Figure created in BioRender. Reuter, M. (2025) https://BioRender.com/xvtmw9p.

Two additional F mutations, both located at the monomer-monomer interface in the assembled capsid (**Fig. 5A**), increased adsorption and dramatically increased persistence. Changes in this region can affect capsid symmetry and stability in icosahedral phages [54, 55]. The pleiotropic nature of these mutations makes it difficult to disentangle the relative fitness contributions of adsorption and persistence. However, the drastic increase in persistence compared to the modest elevation in the adsorption constant suggests that persistence is the primary trait under selection. This is further supported by our simulations, which show that

although both increased adsorption and persistence are favoured under the 3-h regime (**Fig.**
**5C**), selection on persistence alone leads to a more rapid evolution of small-plaque mutants
than selection on adsorption alone (**Fig. S5**).

Plaque phenotypes mirrored the contrasting selective pressures in our experiments, with
small- and large-plaque-forming phages evolving under the 3-h and 30-min regimes,
respectively. These differences in plaque size may reflect adsorption differences, as we found
an inverse correlation between adsorption constant and plaque size (**Fig. 3**). This pattern is
consistent with previous findings in phage λ, where mutations in the host recognition protein J
that increased the adsorption constant led to reduced plaque size [17, 22]. Similar observations
were also made in phage U136B, where fast adsorbers often formed smaller plaques [56]. A
likely explanation for this is that rapid adsorption promotes local reinfection, trapping phages
at the focus of infection and preventing them from reaching uninfected regions of the bacterial
lawn [22].

Our findings identify adsorption and persistence as key life-history traits mediating
ΦX174 adaptation to different environmental conditions. The contrasting effects of different
adsorption constants across environments suggest ecological trade-offs that may drive
divergent selection in natural populations. For instance, sewage water, a major source of
phages, contains both a liquid phase where bacteria grow pelagically and a semi-solid phase
where bacteria form biofilms [57, 58]. High adsorption may be favoured in the liquid phase to
exploit transient host availability. In contrast, slow adsorption may be advantageous in biofilms
or host-rich environments to avoid attachment to already infected cells [17]. Low adsorption
could also be beneficial in particle-rich environments, where phages risk irreversible binding
to inert material [20, 59, 60]. Recurrent transitions between these environments may have
shaped the evolutionary plasticity of adsorption in ΦX174, where single-point mutations in the
F protein can lead to changes in adsorption with profound fitness consequences in different
environments.

The rapid decay of the ancestral ΦX174 in our experimental conditions, losing more
than 90% of infectious particles in less than two hours, prompts a fundamental question: how
does ΦX174 persist in natural environments where host cells may be absent or scarce for
extended periods? One possibility, supported by our results, is that ΦX174 populations
diversify through mutation, generating subpopulations of highly persistent variants. However,

increased persistence may come at a cost [3, 18, 61, 62]. In our experiments, increased
persistence was consistently linked to enhanced adsorption, a combination that may be
maladaptive in host-rich and structured environments due to superinfection and local trapping.
Thus, phenotypic diversification may be an adaptive strategy allowing lytic phages to cope
with environmental heterogeneity.

These findings have potential practical relevance. Phages with different life-history
traits may be more or less suited for treating different types of infections, such as biofilm or
bloodstream infections. Our results illustrate how experimental evolution can uncover key
environmental trade-offs, which may inform the selection or engineering of phages for therapy.
In addition, understanding how phage life-history traits evolve under different selective
pressures is essential for anticipating phage evolutionary dynamics once they are administered
to patients. Thus, experimental evolution serves not only as a valuable research approach but
also as a practical tool to guide the development of safer, more effective, and more predictable
phage-based therapies.

**Acknowledgements**

We acknowledge the Sequencing Service Team from the Max Planck Institute for Evolutionary Biology in Plön for performing the Sanger Sequencing. We thank Daniel Martens and Anja Baade for their support regarding lab logistics as well as the entire Microbial Population Biology Department for their support. We also want to thank David Rogers for helping with PCR troubleshooting and Eva Lievens and Stephen T. Abedon for the insightful discussions on viral life history evolution. MR, CB, and ORM were supported by the International Max-Planck Research School for Evolutionary Biology (IMPRS EvolBio).

**Funding**

The funding for this study was received from the Max Planck Society. Work in JLG lab was also supported by ERC starting grant 853323.

**Data availability**

All raw sequencing, plaque imaging, and counting data, used to generate figures Fig. 1, Fig. 2, Fig. 3, Fig. 4B, Fig. 5B and Fig. S2 is available under **10.5281/zenodo.1707622**. The code used for simulations and analysis is available on GitHub under https://github.com/OReyesMatte/Reuter_2025.

**Supplementary material**

All supplementary material is found in file "Supplementary.pdf". The file contains supplementary figures **Fig. S1-S9**, supplementary tables **Table S1-S13**, and **Supplementary Methods**.

**Conflicts of interest**

The authors declare no conflict of interest.

**Author Orchids**

Manuela Reuter: https://orcid.org/0009-0000-5191-3414

Michael Sieber: https://orcid.org/0000-0003-2703-3665

Octavio Reyes-Matte: https://orcid.org/0009-0006-7031-6213

Christopher Böhmker: https://orcid.org/0009-0003-0098-9816

Jordan Romeyer Dherbey: https://orcid.org/0000-0002-6125-1722

Frederic Bertels: https://orcid.org/0000-0001-6222-4139

Javier Lopez-Garrido: https://orcid.org/0000-0002-8907-502X

## References

1. Abedon ST, Hyman P, Thomas C. Experimental examination of bacteriophage latent-period evolution as a response to bacterial availability. *Appl Environ Microbiol* 2003;**69**:7499–7506. https://doi.org/10.1128/AEM.69.12.7499-7506.2003

2. Eshelman CM *et al*. Unrestricted migration favours virulent pathogens in experimental metapopulations: Evolutionary genetics of a rapacious life history. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2010;**365**:2503–2513. https://doi.org/10.1098/rstb.2010.0066

3. Dessau M *et al*. Selective pressure causes an RNA virus to trade reproductive fitness for increased structural and thermal stability of a viral enzyme. *PLoS Genet* 2012;**8**:e1003102. https://doi.org/10.1371/journal.pgen.1003102

4. Heineman RH, Bull JJ. Testing optimality with experimental evolution: Lysis time in a bacteriophage. *Evolution (N Y)* 2007;**61**:1695–1709. https://doi.org/10.1111/j.1558-5646.2007.00132.x

5. DeLong JP *et al*. Towards an integrative view of virus phenotypes. *Nat Rev Microbiol* 2022;**20**:83–94. https://doi.org/10.1038/s41579-021-00612-w

6. Abedon ST. Impact of phage properties on bacterial survival. In: Adams HT (ed.), Contemporary Trends in Bacteriophage Research. Nova Science Publishers, Inc., 2009, 217–235.

7. Ellis EL, Delbrück M. The growth of bacteriophage. *J Gen Physiol* 1939;**22**:365–384. https://doi.org/10.1085/jgp.22.3.365

8. Delbrück M. Adsorption of bacteriophage under various physiological conditions of the host. *J Gen Physiol* 1940;**23**:631–642. https://doi.org/10.1085/jgp.23.5.631

9. Delbrück M. The growth of bacteriophage and lysis of the host. *J Gen Physiol* 1940;**23**:643–660. https://doi.org/10.1085/jgp.23.5.643

10. Suttle CA, Chen F. Mechanisms and rates of decay of marine viruses in seawater. *Appl Environ Microbiol* 1992;**58**:3721–3729. https://doi.org/10.1128/aem.58.11.3721-3729.1992

11. Abedon ST. Phage Ecology. In: Calendar R, Abedon ST (eds.), The bacteriophages. Oxford: Oxford University Press, 2006, 37–46.

12. Adams MH, Wassermann FE. Frequency distribution of phage release in the one-step growth experiment. *Virology* 1956;**2**:96–108. https://doi.org/10.1016/0042-6822(56)90079-4

13. Abedon ST. Kinetics of phage-mediated biocontrol of bacteria. *Foodborne Pathog Dis* 2009;**6**:807–815. https://doi.org/10.1089/fpd.2008.0242

599    14.    Lindberg HM, McKean KA, Wang I-N. Phage fitness may help predict phage therapy
600           efficacy. *Bacteriophage* 2014;**4**:e964081.
601           https://doi.org/10.4161/21597073.2014.964081

602    15.    Abedon S. Phage therapy pharmacology: Calculating phage dosing. In: Laskin A,
603           Gadd G, Sariaslani S (eds.), Advances in Applied Microbiology. Academic Press Inc.,
604           2011, 1–40.

605    16.    Kowalska JD *et al.* Growing trend of fighting infections in aquaculture environment—
606           opportunities and challenges of phage therapy. *Antibiotics* 2020;**9**:301.
607           https://doi.org/10.3390/antibiotics9060301

608    17.    Gallet R, Shao Y, Wang IN. High adsorption rate is detrimental to bacteriophage
609           fitness in a biofilm-like environment. *BMC Evol Biol* 2009;**9**:241.
610           https://doi.org/10.1186/1471-2148-9-241

611    18.    Heineman RH, Brown SP. Experimental evolution of a bacteriophage virus reveals the
612           trajectory of adaptation across a fecundity/longevity trade-off. *PLoS One*
613           2012;**7**:e46322. https://doi.org/10.1371/journal.pone.0046322

614    19.    Mäkelä K, Laanto E, Sundberg LR. Determinants in the phage life cycle: The dynamic
615           nature of ssDNA phage FLiP and host interactions under varying environmental
616           conditions and growth phases. *Environ Microbiol* 2024;**26**:e16670.
617           https://doi.org/10.1111/1462-2920.16670

618    20.    Daniels LL, Wais AC. Virulence in phage populations infecting *Halobacterium*
619           *cutirubrum*. *FEMS Microbiol Ecol* 1998;**25**:129–134. https://doi.org/10.1016/S0168-
620           6496(97)00089-5

621    21.    Shao Y, Wang IN. Bacteriophage adsorption rate and optimal lysis time. *Genetics*
622           2008;**180**:471–482. https://doi.org/10.1534/genetics.108.090100

623    22.    Gallet R, Kannoly S, Wang IN. Effects of bacteriophage traits on plaque formation.
624           *BMC Microbiol* 2011;**11**:181. https://doi.org/10.1186/1471-2180-11-181

625    23.    Bull JJ *et al.* Exceptional convergent evolution in a virus. *Genetics* 1997;**147**:1497–
626           1507. https://doi.org/https://doi.org/10.1093/genetics/147.4.1497

627    24.    Wichman HA *et al.* Different trajectories of parallel evolution during viral adaptation.
628           *Science (1979)* 1999;**285**:422–424. https://doi.org/10.1126/science.285.5426.422

629    25.    Crill WD, Wichman HA, Bull JJ. Evolutionary reversals during viral adaptation to
630           alternating hosts. *Genetics* 2000;**154**:27–37. https://doi.org/10.1093/genetics/154.1.27

631    26.    Wichman HA, Brown CJ. Experimental evolution of viruses: *Microviridae* as a model
632           system. *Philosophical Transactions of the Royal Society B: Biological Sciences*
633           2010;**365**:2495–2501. https://doi.org/10.1098/rstb.2010.0053

634    27.    Wichman HA, Millstein J, Bull JJ. Adaptive molecular evolution for 13,000 phage
635           generations: A possible arms race. *Genetics* 2005;**170**:19–31.
636           https://doi.org/10.1534/genetics.104.034488

637    28.    Sanger F *et al*. The nucleotide sequence of bacteriophage φX174. *J Mol Biol*
638           1978;**125**:225–246. https://doi.org/10.1016/0022-2836(78)90346-7

639    29.    Sinsheimer RL. A single-stranded deoxyribonucleic acid from bacteriophage φX174. *J*
640           *Mol Biol* 1959;**1**:43–53. https://doi.org/10.1016/S0022-2836(59)80006-1

641    30.    Fane B. ΦX174 *et al*. the *Microviridae*. In: Calendar R (ed.), The Bacteriophages.
642           Oxford University Press, Inc., 2006, 129–145.

643    31.    Baas PD. DNA replication of single-stranded *Escherichia coli* DNA phages. *Biochim*
644           *Biophys Acta* 1985;**825**:111–139. https://doi.org/10.1016/0167-4781(85)90096-X

645    32.    McKenna R *et al*. Atomic structure of single-stranded DNA bacteriophage ΦX174 and
646           its functional implications. *Nature* 1992;**355**:137–143.
647           https://doi.org/10.1038/355137a0

648    33.    Sun L *et al*. Icosahedral bacteriophage Φx174 forms a tail for DNA transport during
649           infection. *Nature* 2014;**505**:432–435. https://doi.org/10.1038/nature12816

650    34.    Burgess AB. Studies on the proteins of ΦX174 II. The protein composition of the ΦX
651           coat. *Proceedings of the National Academy of Sciences* 1969;**64**:613–617.
652           https://doi.org/10.1073/pnas.64.2.613

653    35.    Incardona NL, Selvidge L. Mechanism of adsorption and eclipse of bacteriophage
654           ΦX174 II. Attachment and eclipse with isolated *Escherichia coli* cell wall
655           lipopolysaccharide. *J Virol* 1973;**11**:775–782.

656    36.    Incardona NL, Tuech JK, Murti G. Irreversible binding of phage ΦX174 to cell-bound
657           lipopolysaccharide receptors and release of virus-receptor complexes. *Biochemistry*
658           1985;**24**:6439–6446.

659    37.    Sun Y *et al*. Structural changes of tailless bacteriophage ΦX174 during penetration of
660           bacterial cell walls. *Proceedings of the National Academy of Sciences*
661           2017;**114**:13708–13713. https://doi.org/10.1073/pnas.1716614114

662    38.    Ilag LL *et al*. Calcium ion-induced structural changes in bacteriophage ΦX174.
663           *Journal of Molecular Biologie* 1994;**244**:291–300.
664           https://doi.org/10.1006/jmbi.1994.1730

665    39.    Romeyer Dherbey J *et al*. Stepwise evolution of *E. coli* C and φX174 reveals
666           unexpected lipopolysaccharide (LPS) diversity. *Mol Biol Evol* 2023;**40**:msad154.
667           https://doi.org/10.1093/molbev/msad154

668    40.    Biggs KRH *et al*. Ecological approach to understanding superinfection inhibition in
669           bacteriophage. *Viruses* 2021;**13**:1389. https://doi.org/10.3390/v13071389

670    41.    Hyman P, Abedon ST. Practical methods for determining phage growth parameters.
671           In: Clokie MRJ, Kropinski AM (eds.), Methods in molecular biology, 1st ed. Totowa,
672           NJ: Humana Press, 2009, 175–202.

673    42.    Berg S *et al*. ilastik: interactive machine learning for (bio)image analysis. *Nat Methods*
674           2019;**16**:1226–1232. https://doi.org/10.1038/s41592-019-0582-9

43. Virtanen P *et al*. SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat Methods* 2020;**17**:261–272. https://doi.org/10.1038/s41592-019-0686-2

44. Van Der Walt S *et al*. scikit-image: Image processing in Python. *PeerJ* 2014;**2**:e453. https://doi.org/10.7717/peerj.453

45. R Core Team. A language and environment for statistical computing. R Foundation for Statistical Computing. 2022. Vienna, Austria, 2022.

46. Abedon ST. Detection of Bacteriophages: Phage Plaques. In: Harper DR et al. (eds.), Bacteriophages: biology, technology, therapy. Springer Cham, 2021, 507–538.

47. Lee Y, Yin J. Detection of evolving viruses. *Nat Biotechnol* 1996;**14**:491–493. https://doi.org/10.1038/nbt0496-491

48. Abedon ST, Yin J. Bacteriophage plaques: theory and analysis. In: Clokie MRJ, Kropinski AM (eds.), Bacteriophages: Methods and Protocols, Volume 1: Isolation, Characterization, and Interactions. Totowa, NJ: Humana Press, 2009, 161–174.

49. Fujimura R, Kaesberg P. The adsorption of bacteriophage $\phi$X174 to its host. *Biophys J* 1962;**2**:433–449. https://doi.org/10.1016/S0006-3495(62)86866-0

50. Bayer ME, Starkey TW. The adsorption of bacteriophage $\Phi$X174 and its interaction with *Escherichia coli*; a kinetic and morphological study. *Virology* 1972;**49**:236–256. https://doi.org/10.1016/S0042-6822(72)80026-6

51. Van Der Avoort HGAM *et al*. Regions of incompatibility in single-stranded DNA bacteriophages $\Phi$X174 and G4. *J Virol* 1984;**50**:533–540. https://doi.org/10.1128/jvi.50.2.533-540.1984

52. Wichman HA *et al*. Experimental evolution recapitulates natural evolution. *Philosophical Transactions of the Royal Society B: Biological Sciences* 2000;**355**:1677–1684. https://doi.org/10.1098/rstb.2000.0731

53. Pepin KM, Domsic J, McKenna R. Genomic evolution in a virus under specific selection for host recognition. *Infection, Genetics and Evolution* 2008;**8**:825–834. https://doi.org/10.1016/j.meegid.2008.08.008

54. Lee KH *et al*. First-step mutations for adaptation at elevated temperature increase capsid stability in a virus. *PLoS One* 2011;**6**:e25640. https://doi.org/10.1371/journal.pone.0025640

55. Bull JJ, Badgett MR, Wichman HA. Big-benefit mutations in a bacteriophage inhibited with heat. *Mol Biol Evol* 2000;**17**:942–950. https://doi.org/10.1093/oxfordjournals.molbev.a026375

56. Burmeister AR *et al*. Experimental evolution of the TolC-receptor phage U136B functionally identifies a tail fiber protein involved in adsorption through strong parallel adaptation. *Appl Environ Microbiol* 2023;**89**:e0007923. https://doi.org/10.1128/aem.00079-23

57.  Jensen H, Biggs CA, Karunakaran E. The importance of sewer biofilms. *Wiley*
*Interdisciplinary Reviews: Water* 2016;**3**:487–494. https://doi.org/10.1002/wat2.1144

58.  Balleste E *et al*. Bacteriophages in sewage: abundance, roles, and applications. *FEMS*
*microbes* 2022;**3**:xtac009. https://doi.org/10.1093/femsmc/xtac009

59.  Harvey RW, Ryan JN. Use of PRD1 bacteriophage in groundwater viral transport,
inactivation, and attachment studies. *FEMS Microbiol Ecol* 2004;**49**:3–16.
https://doi.org/10.1016/j.femsec.2003.09.015

60.  Richter Ł *et al*. Adsorption of bacteriophages on polypropylene labware affects the
reproducibility of phage research. *Sci Rep* 2021;**11**:7387.
https://doi.org/10.1038/s41598-021-86571-x

61.  De Paepe M, Taddei F. Viruses' life history: Towards a mechanistic basis of a trade-
off between survival and reproduction among phages. *PLoS Biol* 2006;**4**:e193.
https://doi.org/10.1371/journal.pbio.0040193

62.  García-Villada L, Drake JW. Experimental selection reveals a trade-off between
fecundity and lifespan in the coliphage Qß. *Open Biol* 2013;**3**:130043.
https://doi.org/10.1098/rsob.130043

730